

1 | **Uneven distribution of enamel in the tooth crown of a Plains Zebra *Equus quagga***

2 | Daniela E. Winkler^{1,3} and Thomas M. Kaiser^{2,3}

3 | ¹Corresponding Author, daniela.winkler@uni-hamburg.de; phone: +49 40 428386231

4 | ²thomas.kaiser@uni-hamburg.de

5 | ³Center of **Natural** History (CeNak), University of Hamburg, Martin-Luther-King-Platz 3, 20146
6 | Hamburg, Germany

7 |

8 | Introduction

9 | Hypsodonty is a common evolutionary strategy **of** herbivorous mammals to counter **high** abrasive
10 | loads in the **ingested diet**, which result in **a high degree of** dental wear. Hypsodonty can be easily
11 | achieved in all tooth positions by extending specific ontogenetic phases during tooth development
12 | (von Koenigswald, 2011). Newly erupted hypsodont cheek teeth share a feature between all taxa:
13 | they are not immediately functional. To disintegrate tough plant matter, the relatively rounded apex
14 | of the (pre)molar tooth crowns has to wear down slightly, exposing the enamel ridges which may
15 | then act as shearing blades during mastication. The rapid wear of the topmost tooth crown has been
16 | noted in selenodont molars (Osborn and Lumsden, 1978)₂ and several authors have hypothesised
17 | how this initial wear is facilitated. One theory is that empty chewing movements (thegosis; **i.e. tooth-**
18 | **on-tooth contacts or attrition**) sharpen teeth in adults and **initiate** wear in young **animals** (Every,
19 | 1972; Every et al. 1998). More often, however, such empty chewing is considered a behavioural
20 | anomaly (termed bruxism or pathological thegosis) which appears in livestock, other domestic or
21 | captive animals (eg. Murray et al., 1998; Troxler, 2007; 2012) and also in man.

22 | We propose that the top of the tooth crown should be less resistant to both attritional and
23 | abrasional contacts in order to promote early wear and hence expose functional enamel ridges
24 | quickly. This could be accomplished by either building the top of the tooth crown from less and/or
25 | thinner enamel or by building a less resistant enamel microstructure. Both hypotheses suggest that
26 | the top of the tooth crown is structurally different from the rest of the tooth. Analysis of enamel
27 | microstructure at different tooth crown heights is a destructive and time consuming method,
28 | therefore we chose to study enamel distribution within the tooth crown of a subadult Plains Zebra
29 | (*Equus quagga* sp.) using micro CT-scanning. The Plains Zebra is an ideal model organism for large,
30 | hypsodont herbivores, because it is adapted to grazing in both arid and savannah climates and
31 | therefore needs to have a high tolerance of abrasional tooth wear. Amongst extant large herbivore
32 | species, the Equidae exhibit the highest degree of hypsodonty, only equalled by a few Bovidae like
33 | *Bison bison* (compare hypsodonty indices in Janis, 1988).

34 |

35 | Material and Methods

36 | The selected individual is a loan from Museum für Naturkunde (Berlin). It shows very low or no wear
37 | on the premolar and molar teeth and is therefore in the optimal stage to investigate enamel
38 | distribution within all tooth positions of the same individual. The tooth eruption sequence for upper
39 | permanent teeth in *Equus burchelli* (which is synonymous to *E. quagga*) is M1, M2, I1, P2, P3, P4, I2,

marcus clauss 7.12.14 12:19

Kommentar [1]: this modification to the title is in my view the most elegant way to represent the fact that only one specimen was investigated; this MUST be represented somehow in the title (another possibility would be to use the words 'pilot study')

marcus clauss 7.12.14 12:18

Gelöscht: the hypsodont

marcus clauss 7.12.14 11:26

Kommentar [2]: reviewed by Marcus Claus, Zurich (does not wish to remain anonymous)

marcus clauss 7.12.14 11:26

Gelöscht: natural

marcus clauss 7.12.14 11:26

Kommentar [3]: large amounts or high loads, not large loads

marcus clauss 7.12.14 11:26

Gelöscht: by

marcus clauss 7.12.14 11:26

Gelöscht: large

marcus clauss 7.12.14 11:27

Gelöscht: food consumed

marcus clauss 7.12.14 11:29

Gelöscht: helps to bring the teeth

marcus clauss 7.12.14 11:29

Gelöscht: in

marcus clauss 7.12.14 11:29

Gelöscht: due to pure attritional contacts

marcus clauss 7.12.14 11:38

Kommentar [4]: I think it would be maybe appropriate to cite the Teaford paper on thgosis in utero in guinea pigs here?

48 C, M3, I3 (Erz, 1964). Hence we see small amounts of material loss in the earlier erupting teeth M1,
49 M2, P2, and P3 compared to the unworn P4 and M3. However, we chose not to use unworn
50 premolars and molars of several individuals in order to exclude inter-individual variation in enamel
51 distribution. We focus on the upper permanent dentition, because upper teeth are employed as the
52 standard in studying dental characteristics (Fortelius and Solounias, 2000; Solounias and Semperebon,
53 2002; Archer and Sanson, 2002) and functional traits should be more pronounced as compared to
54 lower teeth (Kaiser and Fortelius, 2003) due to the lack of gravity impact. High resolution computed
55 tomography (microCT) scans with an x-y-z resolution between 0.075 and 1.0 mm were obtained at
56 Steinmann-Institut für Geologie, Mineralogie und Paläontologie (Universität Bonn, Germany) on the
57 CT scanner v|tome|x s (GE phoenix|x-ray). The software VG StudioMax 2.1 (Volume Graphics,
58 Heidelberg) was used for reconstruction of virtual models and further processing. First, each tooth
59 was recreated with all dental tissues (enamel, dentin and cementum) as a voxel model using manual
60 and automatic segmentation tools. Next the mineralised enamel was selected and pure enamel voxel
61 models were created (Fig. 1). We then cut both, the enamel and the full tooth model at approx. 75%,
62 50% and 25% of the initial crown height and created individual models of four tooth sections: Section
63 1 from 100-75% crown height, Section 2 from 75-50% crown height, Section 3 from 50-25% crown
64 height and Section 4 from 25% down to the base of the crown (Fig. 2). Volumes of the enamel
65 sections and full tooth sections were taken directly from these models using VG StudioMax. We
66 further measured thickness of enamel ridges on virtual cross sections through txM1. Measurements
67 were taken at approximately the same position at the metacone for the outer enamel ridge and the
68 inner enamel ridge (Fig. 3A) at the apical and basal part of each section. The approximate height of
69 measurements is indicated in Fig. 3B.

71 Results

72 Data on enamel content are given in Fig. 4. Though distribution of enamel content per section was
73 variable between teeth, it was consistently smallest in Section 1 (the most apical section) for all tooth
74 positions. Section 2 contained 2.5-5.5% more enamel than Section 1, Section 3 even 4.5 -9.0% more
75 enamel. The largest ontogenetic increase in enamel content was found for txM1 and txM2, where
76 Section 4 contained more than 9.6% more enamel. The highest enamel contents were found either in
77 Section 3 (txP2, txP4, txM3) or Section 4 (txP3, txM1, txM2). It is notable that txM3 was composed of
78 more enamel than all other tooth positions and also showed the lowest differences in enamel
79 content between sections. Results for enamel thickness measurements are given in Table 1. The
80 thickness of the inner enamel ridge is largest in Section 2, but very similar in all other sections. The
81 outer enamel ridge is getting thinner from the apical part of Section 1 to the apical part of Section 2,
82 but then increases in thickness from the basal part of Section 2 down to the crown base (compare
83 Fig. 3A and 3B for location of measurements).

85 Discussion

86 The results of this study support our hypothesis that the top of the tooth crown is structurally
87 different from the rest of the tooth. We have shown that the overall enamel content is lowest at the
88 crown top and highest in the lower half of the crown. Our measurements of enamel thickness
89 indicate that both thickness and distribution of enamel vary along the tooth crown. The thinnest

marcus clauss 7.12.14 12:13

Kommentar [5]: this is not a really good excuse and I suggest to delete it. Just say this is a pilot study. Because, if the pattern of increasing proportion of enamel of overall dental tissue from section 1-4 was a common thing, then the difference between the sections (in %) should be rather independent of such inter-individual variation, and occur in all investigated specimens. Actually, accounting for inter-individual variation is the reason why one usually does higher n than 1.

marcus clauss 7.12.14 11:41

Kommentar [6]: see our comment on the 'gravity impact' in Müller J, Clauss M, Codron D, Schulz E, Hummel J, Fortelius M, Kircher P, Hatt J-M (2014) Growth and wear of incisor and cheek teeth in domestic rabbits (*Oryctolagus cuniculus*) fed diets of different abrasiveness. Journal of Experimental Zoology A 321:283-298 personally, I think the 'gravity' argument is not valid (because even if food rests on the lower teeth, this 'resting' does not induce wear. Which tooth (maxilla or mandibular) is more affected depends on other factors than gravity – but this need not be addressed or changed here – just a side-thought

marcus clauss 7.12.14 11:44

Gelöscht: .

marcus clauss 7.12.14 11:45

Gelöscht: 4.

marcus clauss 7.12.14 11:46

Gelöscht: .

marcus clauss 7.12.14 11:46

Gelöscht: .

marcus clauss 7.12.14 11:48

Gelöscht: remainder

95 enamel ridges were not found at the top of the crown; however the overall amount of enamel was
96 lowest at this level.

97 There are relatively more soft dental tissues (dentin and cementum) at the top of the crown and
98 therefore this part of the tooth is prone to fast wear. We further note that the base of each tooth
99 seems to be structurally “enhanced”, as the larger content of enamel should strengthen it and help
100 resist high pressure and stress loads. This interpretation is consistent with our enamel thickness
101 measurements at the base of the crown (Section 4 basal). There the greatest thickness of the outer
102 enamel ridge is recorded, but the inner enamel bands are no longer present, because the two
103 fossettes are worn out.

104 In *Equus quagga*, we find the third upper molar to be structurally different from all other upper
105 teeth, as it has the highest proportion of enamel and the least variation of enamel distribution along
106 the tooth crown. We relate this phenomenon to adaptive pressures related to generally two
107 phenomena:

- 108 1. Mechanical constraint: As the upper M3 is closest to the temporomandibular joint, the
109 highest masticatory forces can be generated here (Greaves, 2012). The high enamel
110 content will then prevent excessive wear and maintain chewing evenly distributed forces
111 induced.
- 112 2. Biogenetic constraints: The M3 is the last tooth to erupt in most mammals, and also in
113 the Zebra. Therefore it is also the tooth position maintaining function when anterior
114 teeth have already been worn out.

115 In general, by being more resistant to wear, txM3 can thus compensate for the functional loss of
116 anterior teeth. Because it comes in occlusion while shear-cutting functionality in anterior teeth is
117 well established, there is no need for a weakened crown top as in other cheek teeth.

118 Though *Equus quagga* is an appropriate model organism, these observations are still restricted to
119 one single specimen of this very taxon. They can, however, help us to refine hypotheses on how
120 mechanical and ontogenetic constraints of wear and resistance may be solved in a biological system,
121 by slight modifications of common structures. The findings suggest that the Zebra as a hypsodont
122 herbivore has undergone severe need of optimisation of its chewing system and that the acquisition
123 of hypsodonty does not mean that basic constraints are rendered insignificant in terms of functional
124 optimisation. As these constraints are universal for all mammals feeding on abrasive diets, we expect
125 to find similar adaptations in other herbivorous species, including bovids.

126

127 Acknowledgements

128 We thank our colleagues at Steinmann Institut for CT scanning of the specimen, the Museum für
129 Naturkunde Berlin for specimen loan and Lucy A. Taylor (University of Oxford) for her suggestions to
130 improve the language. This research is publication no. XX of the DFG Research Unit 771 “Function
131 and performance enhancement in the mammalian dentition—phylogenetic and ontogenetic impact
132 on the masticatory apparatus”.

133

134

marcus clauss 7.12.14 11:50

Kommentar [7]: please be consistent – sometimes you call it txM3, sometimes upper M3, sometimes M3 – please use one word throughout

marcus clauss 7.12.14 11:49

Gelöscht: so

marcus clauss 7.12.14 11:50

Gelöscht: .

marcus clauss 7.12.14 11:58

Gelöscht: singular and

marcus clauss 7.12.14 11:58

Gelöscht: understand

marcus clauss 7.12.14 11:51

Kommentar [8]: a very general statement: in English, no comma in front of a ‘that’ when it would be a ‘dass’ in German

marcus clauss 7.12.14 11:53

Gelöscht: also illustrate,

marcus clauss 7.12.14 11:53

Gelöscht: at least

marcus clauss 7.12.14 11:53

Kommentar [9]: in my view such words are too drastic. Also, you don’t ‘undergo’ a ‘need’.

marcus clauss 7.12.14 11:58

Gelöscht: ,

marcus clauss 7.12.14 12:03

Kommentar [10]: in my view, this sentence is very difficult to understand and, if I understand it correctly, makes a very broad statement that is, in my view, of little help in understanding the phenomenon you present here. I suggest a more sober, precise wording:

The findings suggest that different functional requirements at different tooth positions and tooth wear stages have shaped tooth morphology.

in the next sentence, I would then replace ‘constraint’ with requirement

142 **References**

- 143 Archer, D., Sanson, G., 2002. Form and function of the selenodont molar in southern African
144 ruminants in relation to their feeding habits. *Journal of Zoology* 257, 13–26.
- 145 Erz, W., 1964. Tooth eruption and replacement in Burchell's Zebra *Equus burchelli* Grau 1825.
146 *Arnolda* 22, 1-8.
- 147 Every, R.G., 1972. A New Terminology for Mammalian Teeth Founded on the Phenomenon of
148 Thegosis (Parts I and II) Christchurch Pegasus Press, 1-65.
- 149 Every, D., Tunncliffe, G. A., Every, R.G., 1998. Tooth-sharpening behaviour (thegosis) and other
150 causes of wear on sheep teeth in relation to mastication and grazing mechanisms. *Journal of the*
151 *Royal Society of New Zealand* 28(1), 169-184.
- 152
153
154 Fortelius, M., Solounias, N., 2000. Functional characterization of ungulate molars using the abrasion-
155 attrition wear gradient: A new method for reconstructing paleodiets. *American Museum Novitates*
156 3301, 1–36.
- 157
158 Greaves, W.S., 2012. *The Mammalian Jaw – A Mechanical Analysis*. Cambridge University Press,
159 Cambridge.
- 160
161 Janis, C.M., 1988. An estimation of tooth volume and hypsodonty indices in ungulate mammals, and
162 the correlation of these factors with dietary preference. In: Russell, D.E., Santoro, J.-P., Sigogneau-
163 Russell, D. (Eds.), *Teeth revisited: Proceedings of the VIIth International Symposium on dental*
164 *Morphology: Memoirs of the Museum National History Natural*, Paris, pp. 367-387.
- 165
166 Kaiser, T.M., Fortelius, M., 2003. Differential Mesowear in Occluding Upper and Lower Molars:
167 Opening Mesowear Analysis for Lower Molars and Premolars in Hypsodont Horses. *Journal of*
168 *Morphology* 258, 67–83.
- 169 Murray, C.G., Sanson, G.D., 1998. Thegosis - A critical review. *Australian Dental Journal* 43(3), 192–
170 198.
- 171 Osborn, J.W., Lumsden, A.G.S., 1978. An alternative to thegosis and a re-examination of the ways in
172 which mammalian molars work. *Neues Jahrbuch für Geologie und Paläontologie* 156(3), 371-392.
- 173 Solounias, N., Semprebon, G., 2002. Advances in the reconstruction of ungulate ecomorphology with
174 application to early fossil equids. *American Museum Novitates* 3366, 1–49.
- 175 Troxler, J., 2007. Verhaltensstörungen bei Haustieren. In: Bostedt, H. (Eds.), *Zur Rolle der*
176 *Veterinärmedizin in Forschung und Gesellschaft*. Nova Acta Leopoldina NF 95(353).
- 177 Troxler, J., 2012. Das Verhalten als Grundlage zur Beurteilung des Wohlbefindens von Tieren. Tagung
178 der Plattform Österreichische TierärztInnen für Tierschutz; MAY 10, 2012; Wien. In: Baumgartner, J.
179 (Eds.), *Tierschutz: Anspruch - Verantwortung – Realität*.
- 180 von Koenigswald, W., 2011. Diversity of hypsodont teeth in mammalian dentitions - construction and
181 classification. *Palaeontographica Abteilung A Band 294(1-3)*, 63-94.

marcus clauss 7.12.14 11:30

Kommentar [11]: please use a consistent reference style.

marcus clauss 7.12.14 11:33

Kommentar [12]: here, you use capitalisation of every word in the title – maybe because it is a book, but it looks strange

marcus clauss 7.12.14 11:34

Kommentar [13]: here, you use capitalisation although it is a journal paper

marcus clauss 7.12.14 11:34

Gelöscht:

marcus clauss 7.12.14 11:34

Kommentar [14]: here, you use an issue number – but you do not do that in all references, which makes it inconsistent. Giving issue numbers is unusual anyhow.

marcus clauss 7.12.14 11:35

Kommentar [15]: this is only one person, so it can't be 'Eds.' but must be 'Ed.'

185 Figures and Tables

186 | Fig. 1A. Cross section through a virtual tooth model with all dental tissues. B. As Fig. 1A., enamel only.
187 | Scale bar 50mm.

188 | Fig. 2. Virtual 3D-model of txP3 with all dental tissues. The tooth is separated in four sections, which
189 | are slightly separated from each other for better illustration.

190 | Fig. 3A. Cross section through the virtual model of txM1. Black bars indicate the approximate position
191 | where measurements of enamel thickness were taken on the outer and inner enamel ridge. Scale bar
192 | is 50mm. B. Dashed lines show approximate heights where thickness measurements were taken.
193 | Solid lines represent borders of each section. The tooth is in anatomically correct position with the
194 | top of the crown facing the bottom of the image. Hence, the lower dashed line within one section
195 | marks the positions referred to as “apical”, the upper dashed line as “basal” in Tab. 1.

196 | Fig. 4. Enamel content per section and tooth position. Each bar represents 100% enamel content per
197 | tooth position and shows relative enamel content per section. Percentages above bars give the
198 | relative enamel content per tooth position.

199 | Tab. 1. Measurements of enamel thickness for the outer and inner buccal enamel ridge at two
200 | positions (“apical”, “basal”) of each section. For the basal part of Section 4 no thickness could be
201 | measured for the inner enamel ridge as it has already ended at another height.

marcus clauss 7.12.14 12:04

Gelöscht: .

marcus clauss 7.12.14 12:04

Gelöscht: full

marcus clauss 7.12.14 12:04

Gelöscht: .

marcus clauss 7.12.14 12:04

Gelöscht: .