

# On the genus *Mesopontonia* Bruce, 1967 (Crustacea: Decapoda: Palaemonidae) in Korea, with the description of a new species

Jin-Ho Park<sup>1</sup>, Sammy De Grave<sup>2</sup>, Taeseo Park<sup>Corresp. 3</sup>

<sup>1</sup> College of Natural Sciences, Seoul National University, Seoul, Republic of Korea

<sup>2</sup> Oxford University Museum of Natural History, Oxford University, Oxford, United Kingdom

<sup>3</sup> Overseas Biological Resources Team, National Institute of Biological Resources, Incheon, Republic of Korea

Corresponding Author: Taeseo Park

Email address: polychaeta@gmail.com

*Mesopontonia verrucimanus* and *Mesopontonia kimwoni* sp. nov. are recorded from high-latitude temperate waters in Munseom Islet, Jeju Island, Republic of Korea, with both species collected on gorgonians and sponges by trimix diving between 50 and 75 m depth. *Mesopontonia kimwoni* sp. nov. is morphologically allied to *M. brevicarpus*, but can be distinguished by the cutting edges of the fingers of the first chela being entire. Significant morphological variation in the rostrum as well as the second pereopods is documented in *M. verrucimanus*, although cytochrome c oxidase subunit I (COI) barcode analysis proves this to be infra-specific variation. A key to species of the genus *Mesopontonia* is provided.

**On the genus *Mesopontonia* Bruce, 1967 (Crustacea:  
Decapoda: Palaemonidae) in Korea, with the  
description of a new species**

Jin-Ho Park<sup>1</sup>, Sammy De Grave<sup>2</sup>, Taeseo Park<sup>3</sup>

<sup>1</sup> College of Natural Sciences, Seoul National University, Seoul, Republic of Korea

<sup>2</sup> Oxford University Museum of Natural History, Oxford University, Oxford, United Kingdom

<sup>3</sup> Overseas Biological Resources Team, National Institute of Biological Resources, Incheon,  
Republic of Korea

Corresponding Author:

Taeseo Park<sup>3</sup>

Hwangyeong-ro 42, Seo-gu, Incheon, 22689, Republic of Korea

Email address: polychaeta@gmail.com

# On the genus *Mesopontonia* Bruce, 1967 (Crustacea: Decapoda: Palaemonidae) in Korea, with the description of a new species

Jin-Ho Park<sup>1</sup>, Sammy De Grave<sup>2</sup>, Taeseo Park<sup>3</sup>

<sup>1</sup> College of Natural Sciences, Seoul National University, Seoul, Republic of Korea

<sup>2</sup> Oxford University Museum of Natural History, Oxford University, Oxford, United Kingdom

<sup>3</sup> Overseas Biological Resources Team, National Institute of Biological Resources, Incheon, Republic of Korea

Corresponding Author:

Taeseo Park<sup>3</sup>

Hwangyeong-ro 42, Seo-gu, Incheon, 22689, Republic of Korea

Email address: polychaeta@gmail.com

## Abstract

*Mesopontonia verrucimanus* and *Mesopontonia kimwoni* sp. nov. are recorded from high-latitude temperate waters in Munseom Islet, Jeju Island, Republic of Korea, with both species collected on gorgonians and sponges by trimix diving between 50 and 75 m depth. *Mesopontonia kimwoni* sp. nov. is morphologically allied to *M. brevicarpus*, but can be distinguished by the cutting edges of the fingers of the first chela being entire. Significant morphological variation in the rostrum as well as the second pereopods is documented in *M. verrucimanus*, although cytochrome *c* oxidase subunit I (COI) barcode analysis proves this to be infra-specific variation. A key to species of the genus *Mesopontonia* is provided.

## Introduction

The deep-sea palaemonid shrimp fauna of the Indo-West Pacific is relatively well documented, with to date 23 genera and about 84 species recorded from depths of more than 100m by trawling and dredging (Bruce 1991; De Grave & Fransen 2011; Kou et al. 2016; Li et al. 2008; Marin & Chan 2014; Mitsuhashi & Chan 2006; Okuno 2017; Wang et al. 2015). Among them is the rarely recorded genus, *Mesopontonia* Bruce, 1967 which can be distinguished from related genera by the combination of the absence of both supraorbital and antennal teeth on the carapace, as well as the absence of an exopod on the third maxilliped (Bruce 1967; Bruce 1995; Chace & Bruce 1993).

The most recent classification of carideans by De Grave and Fransen (2011) listed six species in the genus, namely *M. gorgoniophila* Bruce, 1967 (type species), *M. gracilicarpus* Bruce, 1990, *M. brucei* Burukovsky, 1991, *M. monodactylus* Bruce, 1991, *M. verrucimanus* Bruce, 1996 and *M. brevicarpus* Li & Bruce, 2006. The type species was originally described from the

northern part of the South China Sea (Bruce, 1967), in association with two species of the gorgonian genus *Melithaea* Milne Edwards, 1857 (Cnidaria: Octocorallia: Melithaeidae) in depths of 117–183 m (Fig. 1A). Since then, it has been sparingly reported upon from New Caledonia, the Philippines, and eastern Australia within depths of 130 to 270 m (Bruce 1979; Bruce 1984; Bruce 1985; Bruce 1991). *Mesopontonia gracilicarpus* has been reported from New Caledonia and the Chesterfield Islands in depths of 226–600 m (Bruce 1990; Bruce 1991; Li & Bruce, 2006). The southwestern Indian Ocean species, *M. brucei* was recorded from depths of 415–460 m from about 850 km south of Madagascar (Burukovsky, 1991). *Mesopontonia monodactylus* was described from the Loyalty Islands (Bruce 1990), associated with species of the hexactinellid sponge genus *Pheronema* Leidy, 1868 (Porifera: Hexactinellida: Pheronematidae) from 460 m (Bruce 1990); *M. verrucimanus* was reported from the Tanimbar Islands, Indonesia in depths of 184–186 m (Bruce 1996). Finally, another western Indian Ocean species, *M. brevicarpus* was recorded from off La Réunion from 270 m (Li & Bruce, 2006).

Jejudo Island, the largest island in Korea, is located about 80 km off the southwestern coast of the mainland (Fig. 1B). In summer, the Tsushima current mixes with low-salinity, high-turbidity waters from the Yangtze River to influence the environment around Jejudo Island. In winter, the Yangtze River discharge reduces, resulting in a higher local salinity (Rebstock and Kang, 2003; Lim et al. 2019). Munseom Islet (Fig. 1C) is located off the south coast of the main island, and consists of volcanic rocks covered with rich invertebrate communities (Cho et al. 2014; Lutaenko et al. 2019; Lee et al. 2019), with a maximum depth of less than 75m. Thus far, only five symbiotic palaemonid shrimps have been reported from the Jejudo Island area (Koo & Kim 2003; Lee & Ko 2011; Lee & Ko 2014; Park et al. 2019a; Park et al. 2019b), although many more remain unrecorded and will be covered in future contributions.

During a faunal survey for deep-water invertebrate species from previously unexplored habitats around Jejudo Island in 2015–2020, numerous specimens of *Mesopontonia* were collected from gorgonians and sponges by trimix SCUBA diving between 50–75 m depth. Detailed examination of their morphology as well as a phylogenetic analysis including related genera indicated that these belong to *M. verrucimanus* (new record for Korea) and an undescribed species in the same genus, constituting the most northerly record for the genus, as well as the first for temperate waters in the Western Pacific.

In this study, we thus describe both *Mesopontonia* species and present an identification key for the genus *Mesopontonia*. To support systematic studies for deep-sea palaemonid shrimps, molecular analyses and ecological information through direct observations are provided.

## Materials & Methods

**Sample collection.** Specimens of *Mesopontonia* and other palaemonid shrimps were collected by trimix diving at depths between 50–75 m around Munseom Islet, Jejudo Island, organized by Seoul National University (SNU) and the National Institute of Biological Resources (NIBR) during 2015–2020. Comparative material used in the phylogenetic part was collected by conventional SCUBA diving at depths between 10–30 m in Korea (2018–2019), Palau (2019),

Philippines (2018–2019), and Taiwan (2016), organized by Academia Sinica, University of the Philippines Visayas (UPV), Korea Institute of Ocean Science & Technology (KIOST), Marine Biodiversity Institute of Korea (MABIK), and SNU. All specimens were collected together with their host invertebrates and preserved in 80% ethanol. Specimens are deposited in the Marine Arthropod depository Bank of Korea, Seoul National University, Seoul (MADBK), Seoul National University, Seoul (SNU), National Institute of Biological Resources, Incheon (NIBR) and the Zoological Collections of the Oxford University Museum of Natural History, Oxford (OUMNH.ZC).

**Morphological examination.** Shrimps were isolated from the host invertebrate using forceps, with their morphological characteristics observed using stereo microscopes (Leica M205C and M125, Germany) and a light microscope (Olympus BX51, Japan). Digital illustrations were done using a microscope digital camera (Leica MC170, Germany), Helicon focus software (Helicon focus 7.5.6, Ukraine) and drawing tablet (Wacom Intuos Pro PTH-660, China) with Adobe Illustrator software (Adobe Systems, USA), following Coleman (2006).

**Molecular data and phylogenetic analysis.** Molecular phylogenetic analyses were performed to understand the phylogenetic position of the new species, as well as the genus more broadly. Two species of *Mesopontonia* (*M. verrucimanus* and the new species) and 12 deep-sea species from the genera *Altopontonia* Bruce, 1990, *Anchistioides* Paulson, 1875, *Bathymenes* Kou, Li & Bruce, 2016, *Cuapetes* Clark, 1919, *Echinopericlimenes* Marin & Chan, 2014, *Lipkemenes* Bruce & Okuno, 2010, *Palaemonella* Dana, 1852, *Paraclimenes* Bruce, 1995, *Periclimenes* Costa, 1844 and *Thaumastocaris* Kemp, 1922 were selected as the ingroup. *Stenopus hispidus* (Olivier, 1811) was used as an outgroup (Table 1). Total genomic DNA was extracted from eggs or pleopod tissue using the QIAamp® DNA Micro Kit (QIAGEN, Hilden, Germany), following the manufacturer's instructions. Partial sequences of the COI (~658bp) and 16S (~538bp) markers were amplified via polymerase chain reaction (PCR) with the primers jgHCO2198/jgLCO1490 (Geller et al. 2013) and 16S-ar/16S-1472 (Crandall & Fitzpatrick 1996; Palumbi et al. 2002), respectively. PCR reactions and sequence data analysis were performed following Park et al. (2019a).

**Zoobank registration.** The electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Commission on Zoological Nomenclature (ICZN), and hence the new names contained in the electronic version are effectively published under that Code from the electronic edition alone. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix <http://zoobank.org/>. The LSID for this publication is: [urn:lsid:zoobank.org:pub:3CB43670-472F-49AE-80F2-EAE9597E12BD](http://zoobank.org/pub:3CB43670-472F-49AE-80F2-EAE9597E12BD). The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central and CLOCKSS.

# Results

## Taxonomy

Family Palaemonidae Rafinesque, 1815

Genus *Mesopontonia* Bruce, 1967

***Mesopontonia verrucimanus* Bruce, 1996**

Figs. 2–7

*Mesopontonia verrucimanus* Bruce, 1996: 198, 216–218, figs. 8, 29c (type locality: Tanimbar Islands, Indonesia, 7°59'S, 133°02'E, 184–186 m).

**Material examined.** 2 males, 2 females (pocl 2.0–2.5, R 1+7–8/0); Dec. 20, 2015; Munseom Islet, Jejudo Island, Korea (33°13'36"N 126°34'9"E), 55 m, on *Raspailia* (*Raspaxilla*) *hirsuta* Thiele, 1898, leg. JH Park (MADBK 120533\_001); 2 males, 1 female (pocl 2.5, R 1+6–8/0); Apr. 12, 2017; same location (33°13'39"N 126°34'7"E), 58 m, on *R. (R.) hirsuta*, leg. JH Park (MADBK 120533\_002); 1 male, 2 females (pocl 2.3–2.5, R 1+7–8/0–1); Apr. 12, 2017; same location (33°13'39"N 126°34'7"E), 58 m, on *R. (R.) hirsuta*, leg. JH Park (OUMNH.ZC.2018-03-04-06); 1 male, 4 females (pocl 2.5–3.5, R 1+7–8/0–1); May 30, 2017; same location (33°13'36"N 126°34'9"E), 54 m, on *Ellisella* cf. *limbaughi* Bayer & Deichmann, 1960, leg. JH Park (MADBK 120533\_003); 1 male, 1 female (pocl 2.5, R 1+7–8/1); Jan. 18, 2018; same location (33°13'36"N 126°34'9"E), 60m, on *E. cf. limbaughi*, leg. JH Park (MADBK 120533\_004); 2 males, 3 females (pocl 2.0–2.5, R 1+7–8/0–1); Jan. 18, 2018; same location (33°13'36"N 126°34'9"E), 60m, on *E. cf. limbaughi*, leg. JH Park (NIBRIV0000837760–0000837764); 1 female (pocl 3.0, R 1+8/1); Mar. 31, 2018; same location (33°13'36"N 126°34'9"E), 60m, on *E. cf. limbaughi*, leg. JH Park (MADBK 120533\_005); 1 male (pocl 2.6, R 1+8/0); Jun. 20, 2018; same location (33°13'31"N 126°34'11"E), 60m, on *E. cf. limbaughi*, leg. JH Park (MADBK 120533\_006); 1 male, 2 females (pocl 2.78–2.87, R 1+7–8/0–2); Jun. 20, 2018; same location (33°13'31"N 126°34'11"E), 60m, on *E. cf. limbaughi*, leg. JH Park (NIBRIV0000837778–NIBRIV0000837780); 1 male, 1 female, 1 ovigerous female (pocl 2.67–3.47, R 1+6–8/0x1); Jun. 21, 2018; same location (33°13'31"N 126°34'11"E), 55m, on *E. cf. limbaughi*, leg. JH Park (NIBRIV0000862982–NIBRIV0000837784); 1 male (pocl 3.13, R 1+7/0); Sep. 12, 2018; same location (33°13'29"N 126°33'52"E), 60m, on *Myriopathes lata* (Silberfeld, 1909), leg. JH Park (NIBRIV0000862987); 1 male (pocl 3.41, R 1+7/0); Sep. 13, 2018; same location (33°13'29"N 126°33'52"E), 60m, on *E. cf. limbaughi*, leg. JH Park (NIBRIV0000862988); 2 ovigerous females (pocl 3.7–3.8, R 1+8–9/0); Sep. 13, 2018; same location (33°13'29"N 126°33'52"E), 60m, on *Cirripathes* cf. *anguina* (Dana, 1846), leg. JH Park (NIBRIV0000862989–0000837790); 2 males (pocl 3.2, R 1+8–9/0); Jul. 26, 2019; same location (33°13'35"N 126°34'11"E), 60m, on *E. cf. limbaughi*, leg. JH Park (NIBRIV0000862991–0000837792); 1 ovigerous female (pocl 3.4, R 1+9/1); Jul. 27, 2019; same location (33°13'35"N 126°34'11"E), 55m, on *E. cf. limbaughi*, leg. JH Park (NIBRIV0000862993); 1 female (pocl 1.5, R 1+7/0); Aug. 16, 2019; same location (33°13'41"N 126°34'6"E), 53m, on *C. anguina*, leg. JH Park (NIBRIV0000862995); 1 male, 3 females, 2

ovigerous females (pocl 1.5–3.7, R 1+8–9/0–1); Oct. 22, 2019; same location (33°13'37"N 126°34'11"E), 67m, on *E. cf. limbaughi*, leg. JH Park (NIBRIV0000862996–NIBRIV0000863001); 2 males, 1 female (pocl 2.4–3.7, R 1+8–9/0–2); Jan. 15, 2020; same location (33°13'34"N 126°33'45"E), 75m, on *E. cf. limbaughi*, leg. JH Park (MADBK120533\_007); 3 females (pocl 2.5–2.8, R 1+7–8/0); Jan. 15, 2020; same location (33°13'34"N 126°33'45"E), 75m, on *C. anguina*, leg. JH Park (SNU KR JH1100–1102); 1 male (pocl 2.5, R 2–8/1); Jan. 15, 2020; same location (33°13'34"N 126°33'45"E), 75m, on *Raspailia* sp., leg. JH Park (SNU KR JH1095); 1 female (pocl 2.3, R 1–8/0); Jan. 15, 2020; same location (33°13'34"N 126°33'45"E), 75m, on *Raspailia* sp., leg. JH Park (SNU KR JH1096); 1 female (pocl 2.9, R 1–9/0); Jan. 15, 2020; same location (33°13'34"N 126°33'45"E), 75m, on *C. anguina*, leg. JH Park (SNU KR JH1104).

**Description of Korean specimens.** Body (Fig. 2) small-sized, subcylindrical form. Rostrum (Figs. 2 and 3A–D) straight, horizontal, almost as long as pocl, reaching or overreaching distal end of antennular peduncle, 6–9 dorsal teeth, spaced along entire length, 0–2 ventral teeth.

Carapace (Figs. 2 and 3A–D) smooth, glabrous, with epigastric tooth at anterior 0.3 of pocl; without supraorbital and antennal teeth; inferior orbital angle produced; hepatic tooth large, acute, extending to anterior margin of carapace; pterygostomial angle bluntly rounded.

Abdomen (Fig. 2) smooth; pleura of first five segments rounded; sixth pleura with pointed posterolateral angle, posteroventral angle subacute.

Telson (Figs. 2 and 3E, F) about 0.75 of pocl, 4.0 times as long as proximal width; two pairs of small dorsal spiniform setae at 0.4 and 0.65 of telson length respectively, with three pairs of posterior spiniform setae, lateral pair shortest, medial pair long and stout.

Eye (Figs. 2 and 4A) with hemispherical cornea, dorsolaterally with nebenaugae, diameter about 0.20 of pocl.

Antennule (Figs. 2 and 4B) with proximal peduncle bearing distolateral tooth, with small acute tooth at ventromedial margin; stylocerite narrow, bearing sharp point, reaching to 0.45 times of proximal segment; intermediate segment short, 0.4 times of proximal segment length, 0.8 of distal segment; upper flagellum biramous, proximal five segments fused, lower flagellum slender, filiform.

Antenna (Figs. 2 and 4C) basicerite with sharp pointed distodorsal margin; ischiocerite and merocerite unarmed; carpocerite reaching to about 0.4 of scaphocerite length; scaphocerite 4 times as long as maximal width, lateral margin rounded, medial margin convex, distolateral tooth large, at 0.9 of lamella length.

Mouthparts typical for genus. Third maxilliped (Fig. 4D) without exopod, reaching to middle of carpocerite; ultimate segment about 0.4 of antepenultimate segment length, tapering distally, with transverse rows of setae; penultimate segment about 0.6 of antepenultimate segment length, with sparsely row of long setae ventromedially; ischiomerus completely fused to basis, antepenultimate segment feebly compressed distally, with long setae ventromedially; coxa with rounded medial lobe, with rounded lateral plate.

First pereopod (Figs. 2, 5A and 5B) reaching to distal end of scaphocerite; fingers about 0.81 of palm length, tips hooked, cutting edge entire, with transverse row of setae and group of terminal setae; palm ventrolaterally with transverse row of serrulate setae; carpus 1.3 times length of chela with row of serrulate setae along distomesial margin; merus as long as carpus; ischium about 0.7 times length of merus; basis and coxa without special features.

Second pereopods (Figs. 2A, 5C–G) well developed, dissimilar in shape, unequal in size. Major second pereopod (Fig. 5C–E) overreaching distal end of rostrum by middle of propodus; chela about 0.63 of poel, with group of terminal setae; fingers about 0.5 of palm length; dactylus slender, about 3.6 times longer than proximal depth, distally curved with acute tip, proximally with two acute teeth at 0.3 and 0.40, distally entire without dorsolateral flange; fixed finger with acute tip, proximally with two small teeth at 0.2 and 0.3, distally entire; palm subcylindrical, about 4.0 times longer than distal width, covered with minutely blunt tubercles and simple setae; carpus about 0.45 of palm length, about 2 times longer than distal width; merus about 2.0 times as long as carpus, as long as palm length, about 6.2 times longer than distal width; ischium as long as carpus length, about 6.0 times longer than distal width; basis and coxa without special features.

Minor second pereopod (Fig. 5F, G) slightly overreaching distal end of scaphocerite; chela about 0.4 of poel, 0.7 of major chela length, with group of terminal setae; fingers about 0.7 of palm length, distally curved with acute tips, cutting edge entire; palm subcylindrical, about 3 times longer than distal width, smooth slightly tapering proximally; carpus about 1.4 of palm length, about 0.9 of chela length, about 5.6 times longer than distal width; merus about 1.1 times as long as carpus length, about 7.8 times longer than distal width; ischium about 1.1 times as long as merus length, about 7.8 times longer than distal width; basis and coxa without special features.

Ambulatory pereopods (Figs. 2 and 6) subequal in shape and size. Third pereopod (Fig. 6A, B) overreaching distal end of rostrum by distal end of propodus; dactylus about 0.26 of propodus length, about 4.2 times longer than proximal width, not biunguiculate; propodus about 10 times longer than distal width, ventral border with four distolateral spiniform setae including pair distoventral one, with long setae distally, with distodorsal plumose setae and distal setae; carpus about 0.54 times length of propodus, unarmed; merus as long as carpus length, unarmed; ischium about 0.5 length of propodus, unarmed; basis and coxa without special features. Fourth and fifth pereopods (Fig. 6C–F) similar to third pereopod.

Uropod (Figs. 2 and 3E) overreaching distal end of telson; exopod with distolateral tooth and movable acute tooth.

**Variation.** Bruce (1996) described *Mesopontonia verrucimanus* based on a single specimen (holotype), with nine dorsal teeth, unarmed ventrally and markedly unequal, dissimilar second pereopods. The Korean specimens exhibit morphological variation in rostral dentition and the major second pereopod. The number of dorsal and ventral rostral teeth (Figs. 2 and 3A–D) varies from 6–9 and 0–2 respectively. A single specimen (Fig. 3D) harbours two epigastric teeth on the carapace. Several specimens (Fig. 2B) bear two symmetrical second pereopods, which



are very similar to the minor second pereopods in the original description and in other Korean specimens (Fig 2A).

**Color in life.** Whole body and appendages almost transparent with scattered emerald green chromatophores (Fig. 7); longitudinal pale red band along the ventral surface of the body from the carapace to the fifth abdominal somite.

**Geographical distribution.** Presently known from the Tanimbar Islands (Indonesia) and Jejudo Island (Republic of Korea) (Fig. 1).

**Habitat and host.** The present specimens were obtained from gorgonian and sponge colonies below 50 m (Fig. 14A, B), with the deepest samples from 75 m depth. The present specimens demonstrate a lack of host specificity and the species cannot be considered as restricted to gorgonians, as previously postulated. Most specimens were collected on the orange colored sea whip, *Ellisella* cf. *limbaughi* (Bayer & Deichmann 1960), with further specimens obtained on the white colored gorgonian, *Cirrhipathes* cf. *anguina* (Dana, 1846), as well as the orange colored sponge, *Raspailia* (*Raspaxilla*) *hirsuta* Thiele, 1898. A single specimen was collected on the white colored antipatharian, *Myriopathes lata* (Silberfeld, 1909).

*Mesopontonia verrucimanus* can be immediately separated from most other species in the genus which have a biunguiculate dactylus of ambulatory pereopods, except *M. monodactylus* with which it shares a non-biunguiculate dactylus. *M. monodactylus* differs from *M. verrucimanus* primarily by having a distinct dorsolateral flange on the chela of the major second pereopod (Bruce 1991; Bruce 1996).

***Mesopontonia kimwoni* sp. nov.**

urn:lsid:zoobank.org:act:BBA317A3-7140-4D97-BCF4-DB6EEF6617F5

Figs. 8–12

# **Material examined.**

**Type material. Holotype.** 1 ovigerous female (pocl 2.5, R 1+8/2); Jun. 21, 2018; Munseom Islet, Jejudo Island (33°13'31"N 126°34'11"E), 55m, on *Myriopathes lata* (Silberfeld, 1909), leg. JH Park (NIBRIV0000862985). **Paratype.** 1 ovigerous female (pocl 2.8, R 1+8/2); Aug. 16, 2019; same location (33°13'41"N 126°34'6"E), 53m, on *M. lata*, leg. JH Park (NIBRIV0000862994).

**Description.** Body (Fig. 8) small-sized, subcylindrical form. Rostrum (Figs. 8 and 9A) straight, horizontal, almost as long as pocl, reaching or slightly beyond end of antennular peduncle, 8 dorsal teeth, along entire length, 2 subterminal ventral teeth.

Carapace (Fig. 8) smooth, glabrous, with epigastric tooth at anterior 0.3 of pocl; without supraorbital and antennal teeth; inferior orbital angle produced; hepatic tooth large, acute, extending to anterior margin of carapace; pterygostomial angle bluntly rounded.

Abdomen (Fig. 8) smooth; pleura of first five segments rounded; sixth pleura with pointed posterolateral angle, posteroventral angle subacute.

Telson (Figs. 8 and 9D) 0.8 of poel, 4.0 times as long as proximal width; two pairs of small dorsal spiniform setae at 0.45 and 0.7 of telson length respectively, with three pairs of posterior spiniform setae, lateral pair shortest, medial pair long and stout.

Eye (Figs. 8 and 9B–C) with hemispherical cornea, dorsolaterally with nebulose, diameter about 0.23 of poel.

Antennule (Fig. 9B) with proximal peduncle bearing distolateral tooth, with small acute tooth at ventromedial margin; stylocerite narrow, bearing sharp point, reaching to 0.45 times of proximal segment; intermediate segment short, 0.3 times of proximal segment length, subequal to distal segment; upper flagellum biramous, proximal four segments fused, lower flagellum slender, filiform.

Antenna (Figs. 8 and 9B) basicerite with sharp pointed distodorsal margin; ischiocerite and merocerite unarmed; carpocerite reaching to about 0.4 of scaphocerite length; scaphocerite 4 times as long as maximal width, lateral margin rounded, medial margin convex, distolateral tooth large, at 0.9 of lamella length.

Mouthparts not dissected. Third maxilliped (Fig. 9F) without exopod, reaching to 0.7 of carpocerite; ultimate segment about 0.35 of antepenultimate segment length, tapering distally, with transverse rows of setae; penultimate segment about 0.7 of antepenultimate segment length, with sparsely row of long setae ventromedially; ischiomerus completely fused to basis, antepenultimate segment feebly compressed distally, with long setae ventromedially; coxa with rounded medial lobe, with rounded lateral plate.

First pereopod (Figs. 8 and 9A, B) overreaching distal end of scaphocerite; fingers about 0.6 of palm length, tips hooked, cutting edge entire, with transverse row of setae and group of terminal setae; palm ventrolaterally with transverse row of serrulate setae; carpus 1.1 times length of chela with row of serrulate setae along distomesial margin; merus 1.1 times length of carpus; ischium about 0.5 times length of merus; basis and coxa without special features.

Second pereopods (Figs. 8 and 10C–G) well developed, similar in shape, unequal in size. Major second pereopod (Fig. 10C–E) overreaching distal end of rostrum by middle of propodus; chela about 1.3 times as long as poel, with group of terminal setae; fingers about 0.4 of palm length; dactylus slender, about 3.8 times longer than proximal depth, distally curved with acute tip, proximally with two blunt teeth at proximal 0.3 and 0.4, distally entire without dorsolateral flange; fixed finger with distally curved with acute tip, proximally with single acute tooth at 0.4, distally entire; palm subcylindrical, about 4.2 times longer than distal width, covered with minutely blunt tubercles and short simple setae; carpus about 0.4 of palm length, about 2.8 times longer than distal width; merus about 2.1 times as long as carpus, about 0.8 of palm length, 7.0 times longer than distal width; ischium subequal to carpus length, about 7.0 times longer than distal width; basis and coxa without special features.

Minor second pereopod (Fig. 10F, G) overreaching distal end of rostrum by end of carpus; chela about 0.7 of poel, 0.7 of major chela length, with group of terminal setae; fingers about 0.7 of palm length, with distally curved with acute tips, cutting edge entire; palm subcylindrical, about 3.75 times longer than distal width, smooth, slightly tapering proximally; carpus about 1.3

of palm length, about 0.75 of chela length, about 7.2 times longer than distal width; merus about 1.1 times as long as carpus length, about 9 times longer than distal width; ischium about 0.9 of merus length, about 10 times longer than distal width; basis and coxa without special features.

Ambulatory pereopods (Figs. 8 and 11) subequal in shape and size, only third pereopod with distodorsal plumose setae and distal serrulate setae on propodus. Third pereopod (Fig. 11A, B) overreaching distal end of rostrum by distal half of propodus; dactylus about 0.2 of propodus, about 4 times longer than proximal width, about 0.65 of corpus length, biunguiculate; propodus about 0.8 of poel, 16 times longer than distal width, ventral border with four distolateral spiniform setae including single distoventral one, with long setae distally; carpus about 0.4 times length of propodus, unarmed; merus about 0.9 times length of propodus, unarmed; ischium about 0.5 length of propodus, unarmed; basis and coxa without special features.

Fourth pereopod (Fig. 11C, D) with dactylus about 0.2 times length of propodus, about 4 times longer than proximal width, about 0.65 of corpus length, biunguiculate; propodus with four distolateral spiniform setae including single distoventral one, with long simple setae distally; carpus about 0.45 times length of propodus, unarmed; merus subequal to propodus length, unarmed; ischium about 0.46 length of propodus; basis and coxa without special feature. Fifth pereopod (Fig. 11E) similar to fourth pereopod.

Uropod (Figs. 11D) overreaching distal end of telson; exopod with distolateral tooth and movable acute tooth.

**Etymology.** The specific name “kimwoni” is in honor of Dr. Kim, Won, professor in the School of Biological Sciences, Seoul National University. He made a considerable contribution to the systematics of Korean crustaceans.

**Color in life.** Whole body and appendages almost transparent (Figs. 12 and 13); longitudinal red bands along the ventral surface of the body from antennular peduncle to the fifth abdominal somite; tiny white and red chromatophore scattered along the dorsal surface of cornea of eyes, proximal segment of antennular peduncle and abdomen.

**Type locality.** Munseom Islet, Jeju Island, Korea

**Geographical distribution.** Presently only known for the type locality.

**Habitat.** The two specimens of *M. kimwoni* sp. nov. were collected from the gorgonian antipatharian, *Myriopathes lata* (Silberfeld, 1909) in 53–55 m (Fig. 14C).

**Remarks.** Based on the presence of the biunguiculate dactyli of ambulatory pereopods, the new species is morphologically allied to four species: *M. brevicarpus* Li & Bruce, 2006, *M. brucei* Burukovsky, 1991, *M. gorgoniophila* Bruce, 1967, and *M. gracilicarpus* Bruce, 1990; and can be easily separated from the remaining two species in the genus (*M. monodactylus* Bruce, 1991, *M. verrucimanus* Bruce, 1996) which have simple dactyli. The new species differs from *M. gorgoniophila* by the straight distodorsal carina in the major second pereopod (vs. oblique distodorsal carina in *M. gorgoniophila*), with two teeth on the cutting edge of dactylus of major second pereopod (vs. with single tooth in *M. gorgoniophila*), dorsolateral dactylar flange of major second pereopod absent (vs. present in *M. gorgoniophila*), as well as the proportions of the carpus of the minor second pereopod (about 1.3 time palm length in *M. kimwoni* sp. nov. vs.

about 0.8 in *M. gorgoniophila*). The new species differs from *M. gracilicarpus* primarily by the relative short size of the carpus of the first pereopods (about 1.1 times the chela length in *M. kimwoni* **sp. nov.**, vs. about 1.4 in *M. gracilicarpus*), as well as the proportions of the carpus of the minor second pereopod (0.8 times chela length in *M. kimwoni* **sp. nov.** vs. about 1.5 in *M. gracilicarpus*). The new species can be distinguished from *M. brucei* by the presence of a minute tubercle on the chela of the major second pereopod (vs. absent in *M. brucei*), relatively long fingers of first pereopod (about 0.6 of palm length in *M. kimwoni* **sp. nov.** vs. about 0.5 in *M. brucei*), relatively short carpus of major second pereopod (about 0.4 of the palm length in *M. kimwoni* **sp. nov.** vs 0.6 in *M. brucei*) and the relatively long carpus of the minor second pereopod (about 1.3 times the palm length in *M. kimwoni* **sp. nov.** vs. about 0.9 in *M. brucei*).

*M. kimwoni* **sp. nov.**, appears most closely related to the west Indian species, *M. brevicarpus*, sharing a similar rostral formulation, a tuberculate major second pereopod chela, as well as the ratio of the ambulatory pereopods. Both species can be most easily distinguished on the basis of the combination of the following characters, 1) fingers of first chela with entire cutting edge (vs. fine pectinated serrations subapically on both fingers in *M. brevicarpus*); 2) hepatic tooth reaching to the anterior margin of the carapace (vs. reaching or extending to anterior margin of carapace in *M. brevicarpus*) and 3) the finger of minor chela being about 0.7 of the palm length (vs. fingers subequal to palm in *M. brevicarpus*).

Bruce (1996) suggested two further new species may be present in the genus, one collected from Indonesia (Bruce, 1996), as well as the juvenile specimen assigned to *M. gorgoniophila* in Bruce (1985), both however were left unnamed. Given their incomplete or juvenile status these taxa are not considered herein, but are unlikely to be the same species as *M. kimwoni*, as the details of the first and second pereopods are different.

### Molecular data analyses

Fragments of 658 and 462 bp were obtained for the COI and 16S markers, respectively. The multiple sequence alignment revealed that the K2P distance between the five specimens of *M. verrucimanus* which showed minor morphological variations in the dentition of the rostrum and proportions of the second pereopods fall within an intraspecific range, being 0–0.5% (Table 2). The intraspecific divergence between both specimens of *M. kimwoni* **sp. nov.** was higher at 1.1% (Table 2).

To elucidate the phylogenetic position of the genus, an analysis was performed on 22 specimens of 16 species of 12 genera (Table 1). The ML and BI analyses showed the same topology (Fig. 15), and the combined phylogenetic tree clearly demonstrated the monophyly of *Mesopontonia* with high support values (BP=100, PP=100). Furthermore, their distant relationship was supported by the K2P distance, which was 13.6% (Table 2).

From the concatenated tree in the present analysis, the genera *Mesopontonia* and *Paraclimenes* are postulated to be sister taxa with high support values (BP=100, PP=92), indicating that they are more genetically related to each other than the remaining analysed genera, supported by morphological similarities.

# Discussion

The present study explored the commensal palaemonid fauna of Jeju Island, recording *Mesopontonia kimwoni* sp. nov. and *M. verrucimanus* at higher latitude temperate waters than the genus was previously known from. While the six species previously known in the genus had been reported from between 117 and 600 m depth by trawling and dredging (Bruce 1967; Bruce 1979; Bruce 1984; Bruce 1985; Bruce, 1990a; Bruce 1991; Bruce, 1996; Burukovsky, 1991; Li & Bruce, 2006), the present specimens were collected from shallower depths of less than 75 m. As they were directly collected with technical SCUBA diving equipment, more details are available on their habitat and ecology, whilst color patterns are recorded for the first time for the genus as a whole.

*Mesopontonia kimwoni* sp. nov. can be distinguished from all other *Mesopontonia* species by the combination of the biunguiculate dactylus of the ambulatory pereopods, the lack of a dorsolateral dactylar flange on the major second chela, the relatively long carpus of the minor second pereopod and the entire cutting edge of fingers of first chela. Specimens of *M. verrucimanus* exhibited minor morphological variation in rostral dentition and proportions of the major second pereopod, but all specimens are clearly conspecific.

Due to its rarity, *Mesopontonia* had not been previously included in family level phylogenies (e.g. Kou et al. 2013; Gan et al. 2015; Horká et al. 2016), but is herein shown to be phylogenetically close to *Paraclimenes*. This is also supported by a relatively similar morphology with both antennal and supraorbital teeth being absent; and the epigastric and hepatic teeth being present. Nevertheless, *Paraclimenes* can be readily distinguished from *Mesopontonia* by the presence of a well-developed exopod on the third maxilliped (Bruce 1995).

## Key to species of *Mesopontonia* Bruce, 1967 (Adapted from Bruce, 1996; Li & Bruce, 2006)

1. Dactylus of ambulatory pereopods simple.....2
  - Dactylus of ambulatory pereopods biunguiculate.....3
2. Major second pereopod with dorsolateral dactylar flange .....*M. monodactylus* Bruce, 1991
  - Major second pereopod without dorsolateral dactylar flange .....*M. verrucimanus* Bruce, 1996
3. Major second pereopod with dorsolateral dactylar flange .....*M. gorgoniophila* Bruce, 1967
  - Major second pereopod without dorsolateral dactylar flange .....4
4. Carpus of minor second pereopod longer than chela .....*M. gracilicarpus* Bruce, 1990
  - Carpus of minor second pereopod shorter than chela .....5
5. Carpus of minor second pereopod shorter than palm .....*M. brucei* Burukovsky, 1991
  - Carpus of minor second pereopod longer than palm .....6
6. Fingers of first chela with pectinated serrations, subapically..*M. brevicarpus* Li & Bruce, 2006
  - Fingers of first chela with clear entire cutting edge; R. 1+8/2 .....*M. kimwoni* sp. nov.

## Abbreviations used in text, tables and figures

453 **BI** Bayesian Inference  
 454 **BP** Maximum likelihood bootstrap percentage  
 455 **K2P** Kimura 2-parameter  
 456 **ML** Maximum likelihood  
 457 **POCL** postorbital carapace length  
 458 **PP** Bayesian posterior probabilities  
 459 **R** Rostrum formula; formulation of epigastric tooth and teeth on rostrum.  
 460

## 461 **Acknowledgements**

462 We thank Dr. Hyi-Seung Lee (KISOT, Korea) and Dr. Wilfredo L. Campos (UPV, Philippines)  
 463 for their invitation and management in Philippines fieldwork. We are also grateful to Dr. Benny  
 464 K. K. Chan (Academia Sinica, Taiwan) for the invitation to participate in the exploration of  
 465 Green Island, Taiwan. Authors also wish to thank to Prof. Sung-Jin Hwang (Woosuk University,  
 466 Korea) and Dr. Hyung June Kim (MABIK, Korea) for helping identification of the host species,  
 467 and Dr. Sanghui Lee (MABIK, Korea) and Dr. Damin Lee (SNU, KOREA) for their assistance  
 468 in field survey, and to Ms. Eunseon Jun (Sahmyook University, Korea) for helping with  
 469 molecular experiments. TP would like to also thank Ms. Yun Kyoung Kim for her worthful  
 470 assistance with the color drawing in figure 13. We extend sincere thanks to Capt. Jong Moon  
 471 Choi (Blue Research, Korea) for his kind help of photographing the habitats and guidance in  
 472 Deep-Sea Research Diving.  
 473

# References

- Bayer FM, and Elisabeth D. 1960. The Ellisellidae (Octocorallia) and their bearing on the zoogeography of the eastern Pacific. *Proceedings of the Biological Society of Washington* 73:175–182.
- Bruce AJ. 1967. Notes on some Indo-Pacific Pontoniinae III–IX. Descriptions of some new genera and species from the western Indian Ocean and the South China Sea. *Zoologische Verhandelingen* 87:1–73.
- Bruce AJ. 1979. Records of some pontoniine shrimps from the South China Sea. *Cahiers de l'Indo-Pacifique* 1:215–248.
- Bruce AJ. 1984. The pontoniine shrimp fauna of Australia. *Australian Museum Memoir* 18:195–218 DOI:10.3853/j.0067-1967.18.1984.385.
- Bruce AJ. 1985. Decapod Crustacea: Pontoniinae (MUSORSTOM II). *Mémoires du Muséum national d'Histoire naturelle (A) Zoologie* 133:229–260.
- Bruce AJ. 1990a. Crustacea Decapoda: Deep-sea palaemonoid shrimps from New Caledonian Waters. In: Crosnier A, ed. *Résultats des Campagnes MUSORSTOM Vol 6*. Paris: Mémoires du Muséum national d'Histoire naturelle, A, Zoologie, 149–215.
- Bruce AJ. 1990b. *Periclimenes franklini* sp. nov., a new deep-sea shrimp from the Coral Sea (Crustacea: Decapoda: Palaemonidae). *Beagle: Records of the Museum and Art Galleries of the Northern Territory* 7:55–64.
- Bruce AJ. 1991. Crustacea Decapoda: Further deep-sea palaemonoid shrimps from New Caledonian waters. In: Crosnier A, ed. *Résultats des Campagnes MUSORSTOM Vol 9*. Paris: Mémoires du Muséum national d'Histoire naturelle, A, Zoologie, 299–411.
- Bruce AJ. 1995. *A synopsis of the Indo-West Pacific genera of the Pontoniinae (Crustacea: Decapoda: Palaemonidae)*. Königstein: Koeltz Scientific Books.
- Bruce AJ. 1996. Crustacea Decapoda: Palaemonoid shrimps from the Indo-West Pacific region mainly from New Caledonia. In: Crosnier A, ed. *Résultats des Campagnes MUSORSTOM, Volume 15*. Paris: Mémoires du Muséum national d'Histoire naturelle, 197–267.
- Bruce AJ, and Okuno J. 2010. Designation of a new genus *Lipkemenes*, with supplementary description and range extension of its type species, *L. lanipes* (Kemp, 1922) (Decapoda, Palaemonidae). In: Fransen CHJM, De Grave S, and Ng PKL, eds. *Studies on Malacostraca: Lipke Bijdeley Holthuis Memorial Volume*. Leiden: Brill, 159–171 DOI:10.1163/9789047427759\_008.
- Burukovsky RN. 1991. New and rare species of shrimps from the south-west part of the Indian Ocean [in Russian]. *Zoologicheskii Zhurnal* 70:36–41.
- Chace FAJ, and Bruce AJ. 1993. The Caridean Shrimps (Crustacea: Decapoda) of the Albatross Philippine Expedition, 1907–1910, Part 6: Superfamily Palaemonoidea. *Smithsonian Contributions to Zoology*:1–152.
- Cho I-Y, Kang D-W, Kang J, Hwang H, Won J-H, Paek WK and Seo S-Y. 2014. A study on the biodiversity of benthic invertebrates in the waters of Seogwipo, Jeju Island, Korea. *Journal of Asia-Pacific Biodiversity* 7:e11–e18 DOI:10.1016/j.japb.2014.03.003.
- Clark AH. 1919. Some necessary changes in crustacean nomenclature. *Proceedings of the Biological Society of Washington* 32:199.
- Coleman CO. 2006. Substituting time-consuming pencil drawings in arthropod taxonomy using stacks of digital photographs. *Zootaxa* 1360:61–68 DOI:10.11646/zootaxa.1360.1.4.
- Costa OG. 1844. Su due nuovi generi di Crostacei decapodi macrouri. *Annali delle Accademia de-gli Aspiranti Naturalisti, Napoli* 2:285–292.
- Crandall KA, and Fitzpatrick JF, Jr. 1996. Crayfish molecular systematics: using a combination of procedures to estimate phylogeny. *Systematic Biology* 45:1–26 DOI:10.1093/sysbio/45.1.1.
- Dana JD. 1846. United States Exploring Expedition during the years 1838–1842. *Zoophytes* 7:1–740 DOI:10.5962/bhl.title.69333.

- 522 Dana JD. 1852. Conspectus crustaceorum, &c. Conspectus of the Crustacea of the exploring expedition  
523 under Capt. C. Wilkes, U.S.N. Macroura. *Proceedings of the Academy of Natural Sciences of*  
524 *Philadelphia* 6:10–28.
- 525 De Grave S. 1999. Variation in rostral dentition and telson setation in a saltmarsh population of  
526 *Palaemonetes varians* (Leach) (Crustacea: Decapoda: Palaemonidae). *Hydrobiologia* 397:101–  
527 108 DOI:10.1023/A:1003698615660.
- 528 De Grave S, and Fransen CHJM. 2011. Carideorum Catalogus: The Recent Species of the  
529 Dendrobranchiate, Stenopodidean, Procarididean and Caridean Shrimps (Crustacea: Decapoda).  
530 *Zoologische Mededelingen* 85:195–588.
- 531 De Grave S, and Sakihara TS. 2011. Further records of the anchialine shrimp, *Periclimenes pholeter*  
532 Holthuis, 1973 (Crustacea, Decapoda, Palaemonidae). *Zootaxa* 2903:64–68  
533 DOI:10.11646/zootaxa.2903.1.7.
- 534 Ďuriš Z, and Bruce AJ. 1995. A revision of the ‘petitthouarsii’ species-group of the genus *Periclimenes*  
535 Costa, 1844 (Crustacea: Decapoda: Palaemonidae). *Journal of Natural History* 29:619–671  
536 DOI:10.1080/00222939500770221.
- 537 Ďuriš Z, and Šobánková A. 2020. Deep sea palaemonid shrimps of Papua New Guinea (Crustacea:  
538 Decapoda: Palaemonidae). In: Corbari L, Ahyong ST, and CHAN T-Y, eds. *Deep-Sea*  
539 *Crustaceans from Papua New Guinea. Tropical Deep-Sea Benthos 31*. Paris: Muséum national  
540 d’Histoire naturelle, 369–401.
- 541 Gan ZB, Li XZ, Chan T-Y, Chu KH and Kou Q. 2015. Phylogeny of Indo-West Pacific pontoniine  
542 shrimps (Crustacea: Decapoda: Caridea) based on multilocus analysis. *Journal of Zoological*  
543 *Systematics and Evolutionary Research* 53:282–290 DOI:10.1111/jzs.12108.
- 544 Geller J, Meyer C, Parker M, and Hawk H. 2013. Redesign of PCR primers for mitochondrial cytochrome  
545 c oxidase subunit I for marine invertebrates and application in all-taxa biotic surveys. *Molecular*  
546 *Ecology Resources* 13:851–861 DOI:10.1111/1755-0998.12138.
- 547 Horká I, De Grave S, Fransen CHJM, Petrusek A, and Ďuriš Z. 2016. Multiple host switching events  
548 shape the evolution of symbiotic palaemonid shrimps (Crustacea: Decapoda). *Scientific Reports*  
549 6:26486 DOI:10.1038/srep26486.
- 550 Kemp SW. 1922. Notes on Crustacea Decapoda in the Indian Museum, XV. Pontoniinae. *Records of the*  
551 *Indian Museum* 24:133–288.
- 552 Koo H, and Kim W. 2003. First report of palaemonid shrimp *Onyccaris callyspongiae* (Decapoda:  
553 Caridea: Palaemonidae) from Korea. *The Korean Journal of Systematic Zoology* 19:251–255.
- 554 Kou Q, Li XZ, Chan T-Y, Chu KH and Gan Z. 2013. Molecular phylogeny of the superfamily  
555 Palaemonoidea (Crustacea :Decapoda : Caridea) based on mitochondrial and nuclear DNA  
556 reveals discrepancies with the current classification. *Invertebrate Systematics* 27, 502–514  
557 DOI:10.1071/IS13005.
- 558 Kou Q, Li XZ, and Bruce AJ. 2016. Designation of a new genus *Bathymenes* for the deep-sea pontoniine  
559 shrimps of the ‘ *Periclimenes alcocki* speciesgroup’ (Decapoda, Caridea, Palaemonidae), with a  
560 checklist of the species assigned to the genus. *Chinese Journal of Oceanology and Limnology*  
561 34:170–176 DOI:10.1007/s00343-015-4359-4.
- 562 Kwak M-T, Seo G-H, Cho Y-K, Kim B-G, You SH, and Seo J-W. 2015. Long-term comparison of  
563 satellite and in-situ sea surface temperatures around the Korean Peninsula. *Ocean Science*  
564 *Journal* 50:109–117 DOI:10.1007/s12601-015-0009-1.
- 565 Lee KH, and Ko HS. 2011. First records of tow pontoniid shrimps (Crustacea: Decapoda: Caridea:  
566 Palaemonidae) from Korea. *The Korean Journal of Systematic Zoology* 27:239–245  
567 DOI:10.5635/KJSZ.2011.27.3.239.
- 568 Lee KH, and Ko HS. 2014. Occurrence of pontoniine shrimp, *Periclimenes brevicarpalis* (Decapoda:  
569 Caridea: Palaemonidae) in Korean waters. *Animal Systematics, Evolution and Diversity* 30:319–  
570 322 DOI:10.5635/ASED.2014.30.4.319.



- 571 Lee SH, and Ko HS. 2013. The first record of *Conchodytes nipponensis* (De Haan, 1844) (Crustacea:  
572 Decapoda: Palaemonidae) associated with pen shell *Atrina pectinata* (Linnaeus, 1767) (Mollusca:  
573 Bivalvia) from Korea. *Journal of Species Research* 2:1–6 DOI:10.12651/JSR.2013.2.1.001.
- 574 Lee T, Stöhr S, Bae YJ and Shin S. 2019. A new fissiparous brittle star, *Ophiacantha scissionis* sp. nov.  
575 (Echinodermata, Ophiuroidea, Ophiacanthida), from Jeju Island, Korea. *Zoological Studies* 58:e8  
576 DOI:10.6620/ZS.2019.58-08.
- 577 Li XZ, and Bruce AJ. 2006. Further Indo-Pacific palaemonoid shrimps (Crustacea: Decapoda:  
578 Palaemonoidea), principally from the New Caledonian region. *Journal of Natural History*  
579 40:611–738 DOI:10.1080/00222930600763627.
- 580 Li XZ, Mitsuhashi M, and Chan T-Y. 2008. Deep-sea pontoniines (Decapoda: Palaemonidae) from the  
581 Philippine “panglao 2005” expedition, with descriptions of four new species. *Journal of*  
582 *Crustacean Biology* 28:385–411 DOI:10.1163/20021975-99990384.
- 583 Lim N-L, Lee H-M, Jeung H-D, Noseworthy RG, Jung S, Choi K-S. 2019. Early larval development and  
584 annual gametogenesis of the brooding oyster *Ostrea cicumpicta* (Pilsbry, 1904) in the shallow  
585 subtidal benthic ecosystem in Jeju Island, off the south coast of Korea. *Zoological studies* 58:e29  
586 DOI:10.6620/ZS.2019.58-29.
- 587 Lutaenko KA, Noseworthy RG and Choi K-S. 2019. Marine bivalve mollusks of Jeju Island (Korea). Part  
588 1. *The Korean Journal of Malacology* 35:149–238 DOI:10.9710/kjm.2019.35.2.149.
- 589 Marin I, and Chan T-Y. 2014. Deep water echinoid-associated pontonine shrimp “*Periclimenes hertwigi*  
590 Balss, 1913” species group (Crustacea: Decapoda: Caridea: Palaemonidae): species review,  
591 description of a new genus and species from Philippines. *Zootaxa* 3835:301–324  
592 DOI:10.11646/zootaxa.3835.3.1.
- 593 Milne-Edwards H. 1857. *Histoire Naturelle des Coralliaires ou Polypes Proprement Dits. Vol. 1.*  
594 *Librairie*. Paris: Librairie Encyclopédique de Roret DOI:10.5962/bhl.title.11574.
- 595 Mitsuhashi M, and Chan T-Y. 2006. A new genus and species of deep-water pontonine shrimp  
596 (Decapoda, Caridea, Palaemonidae) from Taiwan. *Zoosystema* 28:389–398.
- 597 Okuno J. 2017. Designation of a new genus *Michaelimenes* (Decapoda: Caridea: Palaemonidae), with  
598 new host record and range extension of its type species, *M. perlucidus* (Bruce, 1969). *Nauplius*  
599 25:e2017013 DOI:10.1590/2358-2936e2017013.
- 600 Olivier AG. 1811. Suite de l’introduction à l’histoire naturelle des insectes. Paléon. In: Olivier AG, ed.  
601 *Encyclopédie méthodique Histoire naturelle Insectes*. Paris: H. Agasse, Imprimeur-Libraire, 656–  
602 670.
- 603 Palumbi S, Martin A, Romano S, Mcmillan WO, Stice L, and Grabowski G. 2002. *The simple fool’s*  
604 *guide to PCR. version 2.0*. Honolulu: Department of Zoology and Kewalo Marine Laboratory,  
605 University of Hawaii.
- 606 Park J-H, De Grave S, and Kim W. 2019a. A new species of the genus *Cristimenes* Đuriš & Horká, 2017  
607 (Decapoda, Caridea, Palaemonidae). *Zookeys* 852:53–71 DOI:10.3897/zookeys.852.34959.
- 608 Park J-H, De Grave S, and Kim W. 2019b. On the systematic status of *Isopericlimenaeus* Marin, 2012  
609 and its type species, *Periclimenaeus gorgonidarum* (Balss, 1913) (Crustacea: Decapoda:  
610 Palaemonidae). *Zootaxa* 4614:353–367 DOI:10.11646/zootaxa.4614.2.5.
- 611 Paul’son O. 1875. *Studies on Crustacea of the Red Sea with notes regarding other seas. Part 1*  
612 *Podophthalmata and Edriophthalmata (Cumacea) [in Russian]*. Kiev: S. V. Kul’zhenko.
- 613 Rafinesque CS. 1815. *Analyse de la nature ou tableau de l’univers et des corps organisés*. Palermo.
- 614 Rebstock GA, Kang YS. 2003. A comparison of three marine ecosystems surrounding the Korean  
615 peninsula: Responses to climate change. *Progress in Oceanography* 59:357–379  
616 DOI:10.1016/j.pocean.2003.10.002.
- 617 Silberfeld E. 1909. Diagnosen neuer japanischer Antipatharien aus der Sammlung von Herrn Prof Doflein  
618 (München). *Zoologischer Anzeiger* 34:760–763.
- 619 Thiele J. 1898. Studien über pazifische Spongien. I. Japanische Demospongien. *Zoologica: Original-*  
620 *Abhandlungen aus dem Gesamtgebiete der Zoologie. Stuttgart* 24:1–72.

- 621 Wang Y-R, Chan T-Y, and Sha Z-L. 2015. A new deep-sea species of the genus *Urocaridella* (Crustacea:  
622 Decapoda: Caridea: Palaemonidea) from Yap Seamount in the Western Pacific. *Zootaxa*  
623 4012:191–197 DOI:10.11646/zootaxa.4012.1.11.
- 624 Wicksten MK. 1995. Within-species variation in *Periclimenes yucatanicus* (Ives), with taxonomic  
625 remarks on *P. pedersoni* Chace (Crustacea: Decapoda: Caridea: Palaemonidae). *Proceedings of*  
626 *the Biological Society of Washington* 108:458–464.

**Figure 1 Map of Indo West Pacific Ocean (IWP).** Map showing (A) distribution of *Mesopontonia* species, (B) location of Jejudo Island, and (C) the type locality of *M. kimwoni* sp. nov.

**Figure 2 *Mesopontonia verrucimanus* Bruce, 1996, habitus, lateral view.** (A) Ovigerous female, pocl 3.7 mm (NIBRIV0000862989); (B) male, pocl 3.2 mm (NIBRIV0000862991).

**Figure 3 *Mesopontonia verrucimanus* Bruce, 1996.** (A) Frontal margin and rostrum, lateral view; (B, C) same, lateral view; (D) same, lateral view; (E) telson and uropods, dorso-lateral view; (F) distal end of telson, dorsal view. (A, E, F) Ovigerous female, pocl 3.8 mm (NIBRIV0000862990); (B) Male, pocl 2.7 mm (NIBRIV0000862982); (C) Female, 2.78 mm (NIBRIV0000862978); (D) Male, pocl 2.5 mm (SNU KR JH1095).

**Figure 4 *Mesopontonia verrucimanus* Bruce, 1996.** (A) Left eye, dorsal view; (B) left antennule, dorsal view; (C) left antenna, dorsal view; (D) left third maxilliped. (A–C) Ovigerous female, pocl 3.8 mm (NIBRIV0000862990); (D) Ovigerous female, pocl 3.7 mm (NIBRIV0000862989).

**Figure 5 *Mesopontonia verrucimanus* Bruce, 1996, ovigerous female, pocl 3.7 mm (NIBRIV0000862989).** (A) Left first pereopod. (B) same, chela; (C) major right second pereopod; (D) same, chela and carpus; (E) same, fingers; (F) minor left second pereopod; (G) same, chela.

**Figure 6 *Mesopontonia verrucimanus* Bruce, 1996, ovigerous female, pocl 3.7 mm (NIBRIV0000862989).** (A) Left third pereopod; (B) same, dactylus and distal propodus; (C) left fourth pereopod; (D) same, dactylus and distal propodus; (E) left fifth pereopod; (F) same, dactylus and distal propodus.

**Figure 7 *Mesopontonia verrucimanus* Bruce, 1996 on *Ellisella limbaughii* Bayer & Deichmann, 1960 from Munseom Islet, Jejudo Island.** (A) Female specimen, pocl 2.5 mm (MADBK 120533\_004); (B) male specimens, pocl 3.0 mm (MADBK 120533\_003). Photographic Credits: (A) Jin-Ho Park, (B) Jong Moon Choi.

**Figure 8 *Mesopontonia kimwoni* sp. nov., habitus, lateral view.** Holotype, ovigerous female, pocl 2.5 mm (NIBRIV0000862985).

**Figure 9 *Mesopontonia kimwoni* sp. nov., ovigerous female pocl 2.5 mm (NIBRIV0000862985), holotype.** (A) frontal margin and rostrum, lateral view; (B) frontal region, dorsal view; (C) left eye, dorsal view; (D) left antenna, ventral view; (E) telson and uropod, dorsal view; (F) distal end of telson, dorsal view; (G) left third maxilliped, mesial view.

**Figure 10 *Mesopontonia kimwoni* sp. nov., ovigerous female pocl 2.5 mm (NIBRIV0000862985), holotype.** (A) left first pereopod; (B) same, chela; (C) major right second pereopod; (D) same, chela and carpus; (E) same, fingers; (F) minor left second pereopod; (G) same, chela.

**Figure 11 *Mesopontonia kimwoni* sp. nov., ovigerous female pocl 2.5 mm (NIBRIV0000862985), holotype.** (A) left third pereopod; (B) same, dactylus and distal propodus; (C) left fourth pereopod; (D) same, dactylus and distal propodus; (E) left fifth pereopod.

**Figure 12 *Mesopontonia kimwoni* sp. nov. from Munseom Islet, Jejudo Island.** Holotype, ovigerous female, pocl 2.5 mm (NIBRIV0000862985). Photographic credit: Jin-Ho Park.

**Figure 13 Artistic interpretation of *Mesopontonia kimwoni* sp. nov.** Holotype, ovigerous female, pool 2.5 mm (NIBRIV0000862985). Painting by Kyoung Kim.

**Figure 14 Habitat and host of *Mesopontonia* from Munseom Islet, Jeju Island, Korea.** Habitat and host specimens of *M. verrucimanus* (A, B) and *M. kimwoni* sp. nov. (C). (A) host sea whip, *Ellisella* cf. *limbaughii* and habitat in depth of 57 m; (B) host sponge *Raspailia* (*Raspaxilla*) *hirsuta* and habitat in depth of 55 m; (C) host black coral *Myriopathes lata* and habitat in depth of 55m. Photographic Credits: Jong Moon Choi.

**Figure 15 Phylogenetic tree obtained by the Maximum likelihood (ML) analysis based on the combined dataset for COI and 16S sequences.** Numbers at nodes represent Maximum Likelihood bootstrap percentage (above) and Bayesian posterior probabilities (below), numbers less than 75% are not shown.

**Table 1:** Species used in the phylogenetic analysis with GenBank accession numbers and source. Used symbols: N/A - not available.

**Table 2:** Pairwise distances of COI sequences and selected morphological characteristics for specimens used in the analysis. (P2 - second pereopods).

**Table 1**(on next page)

Species used in the phylogenetic analysis with GenBank accession numbers and source.  
Used symbols: N/A - not available.

**Table 1:**

Species used in the phylogenetic analysis with GenBank accession numbers and source. Used symbols: N/A - not available.

Species	Voucher ID	Location	GenBank accession number		Sources
			COI	16S	
<i>Mesopontonia kimwoni</i> sp. nov. (1)	NIBRIV0000862985	Korea	MT311866	MT311974	Present study
<i>M. kimwoni</i> sp. nov. (2)	NIBRIV0000862994	Korea	MT311865	MT311973	Present study
<i>M. verrucimanus</i> (1)	NIBRIV0000862999	Korea	MT311867	MT311975	Present study
<i>M. verrucimanus</i> (2)	NIBRIV0000862991	Korea	MT311868	MT311976	Present study
<i>M. verrucimanus</i> (3)	NIBRIV0000862990	Korea	MT311869	MT311977	Present study
<i>M. verrucimanus</i> (4)	SNU KR JH1095	Korea	MT311870	MT311978	Present study
<i>M. verrucimanus</i> (5)	MADBK 120533_007	Korea	MT311871	MT311979	Present study
<i>Paraclimenes</i> sp. (1)	SNU-KR_JH357	Korea	MT311872	MT311980	Present study
<i>Paraclimenes</i> sp. (2)	SNU-KR-JH466	Korea	MT311873	MT311981	Present study
<i>Altopontonia disparostris</i>	RMNH.CRUS.D.51028	Norfolk Ridge	KM921671	KU064797	Horká et al. 2016
<i>Cuapetes tenuipes</i>	UO V08-48	Vietnam	KU064965	KU064814	Horká et al. 2016
<i>Palaemonella rotumana</i>	MTQ W-33176	Australia	KR088755	KU064830	Horká et al. 2016
<i>Anchistioides willeyi</i>	SNU-Pal_Pal12	Palau	MT311864	MT311972	Present study
<i>Lipkemenes lanipes</i>	SNU-PH_PH99	Philippines	MT311863	MT311971	Present study
<i>Thaumastocaris streptopus</i>	SNU-TW_TW62	Taiwan	MT311862	MK602867	Present, Park et al. 2019
<i>Bathymenes aleator</i>	UO-948	Papua New Guinea	MK940921	MK940946	Đuriš & Šobáňová 2020
<i>Echinopericlimenes aurorae</i>	UO-1057	Philippines	MK940923	MK940948	Đuriš & Šobáňová 2020
<i>Echinopericlimenes dentidactylus</i>	UO-1059	Philippines	MK940930	MK940954	Đuriš & Šobáňová 2020
<i>Echinopericlimenes calcaratus</i>	UO-960	Papua New Guinea	MK940925	MK940949	Đuriš & Šobáňová 2020
<i>Periclimenes laccadivensis</i>	UO-881	Papua New Guinea	MK940931	MK940955	Đuriš & Šobáňová 2020
<i>Periclimenes uniungiculatus</i>	UO-586	Papua New Guinea	MK940932	MK940956	Đuriš & Šobáňová 2020
<i>Stenopus hispidus</i> (outgroup)	UO V10-17	Vietnam	KJ690260	KU064859	Horká et al. 2016

## **Table 2**(on next page)

Pairwise distances of COI sequences and selected morphological characteristics for specimens used in the analysis. (P2 - second pereopods).

**Table 2:**

**Pairwise distances of COI sequences and selected morphological characteristics for specimens used in the analysis. (P2 - second pereopods).**


	Species	Rostral formula	Shape and size of P2	1	2	3	4	5	6
1	<i>M. kimwoni</i> sp. nov. (1)	1-8/2	Unequal and dissimilar						
2	<i>M. kimwoni</i> sp. nov. (2)	1-8/2	Unequal and dissimilar	0.011					
3	<i>M. verrucimanus</i> (1)	1-8/1	Unequal and dissimilar	0.142	0.1360				
4	<i>M. verrucimanus</i> (2)	1-8/0	Equal and similar	0.1380	0.1320	0.0030			
5	<i>M. verrucimanus</i> (3)	1-9/0	Unequal and dissimilar	0.1380	0.1320	0.0030	0.0000		
6	<i>M. verrucimanus</i> (4)	2-8/1	Unequal and dissimilar	0.1400	0.1340	0.0050	0.0020	0.0020	
7	<i>M. verrucimanus</i> (5)	1-9/2	Equal and similar	0.1380	0.1320	0.0030	0.0000	0.0000	0.0020



# Figure 1

Map of Indo West Pacific Ocean (IWP).


Map showing (A) distribution of *Mesopontonia* species, (B) location of Jeju Island, and (C) the type locality of *M. kimwoni* sp. nov.



# Figure 2

*Mesopontonia verrucimanus* Bruce, 1996, habitus, lateral view.


(A) Ovigerous female, pocl 3.7 mm (NIBRIV0000862989); (B) male, pocl 3.2 mm (NIBRIV0000862991).



# Figure 3

*Mesopontonia verrucimanus* Bruce, 1996.


(A) Frontal margin and rostrum, lateral view; (B, C) same, lateral view; (D) same, lateral view; (E) telson and uropods, dorso-lateral view; (F) distal end of telson, dorsal view. (A, E, F) Ovigerous female, pochl 3.8 mm (NIBRIV0000862990); (B) Male, pochl 2.7 mm (NIBRIV0000862982); (C) Female, 2.78 mm (NIBRIV0000862978); (D) Male, pochl 2.5 mm (SNU KR JH1095).



# Figure 4

*Mesopontonia verrucimanus* Bruce, 1996.

(A) Left eye, dorsal view; (B) left antennule, dorsal view; (C) left antenna, dorsal view; (D) left third maxilliped. (A–C) Ovigerous female, pocl 3.8 mm (NIBRIV0000862990); (D) Ovigerous female, pocl 3.7 mm (NIBRIV0000862989).




# Figure 5

*Mesopontonia verrucimanus* Bruce, 1996, ovigerous female, pocl 3.7 mm (NIBRIV0000862989).

(A) Left first pereopod. (B) same, chela; (C) major right second pereopod; (D) same, chela and carpus; (E) same, fingers; (F) minor left second pereopod; (G) same, chela.






# Figure 6

*Mesopontonia verrucimanus* Bruce, 1996, ovigerous female, pocl 3.7 mm (NIBRIV0000862989).


(A) Left third pereopod; (B) same, dactylus and distal propodus; (C) left fourth pereopod; (D) same, dactylus and distal propodus; (E) left fifth pereopod; (F) same, dactylus and distal propodus.



# Figure 7

*Mesopontonia verrucimanus* Bruce, 1996 on *Ellisella limbaughi* Bayer & Deichmann, 1960 from Munseom Islet, Jejudo Island.


(A) Female specimen, pocl 2.5 mm (MADBK 120533\_004); (B) male specimens, pocl 3.0 mm (MADBK 120533\_003). Photographic Credits: (A) JH Park, (B) JM Choi.



# Figure 8

*Mesopontonia kimwoni* sp. nov., habitus, lateral view.

Holotype, ovigerous female, pochl 2.5 mm (NIBRIV0000862985).




# Figure 9

*Mesopontonia kimwoni* sp. nov., ovigerous female pochl 2.5 mm (NIBRIV0000862985), holotype.

(A) frontal margin and rostrum, lateral view; (B) frontal region, dorsal view; (C) left eye, dorsal view; (D) left antenna, ventral view; (E) telson and uropod, dorsal view; (F) distal end of telson, dorsal view; (G) left third maxilliped, mesial view.








# Figure 10

*Mesopontonia kimwoni* sp. nov., ovigerous female pochl 2.5 mm (NIBRIV0000862985), holotype.


(A) left first pereopod; (B) same, chela; (C) major right second pereopod; (D) same, chela and carpus; (E) same, fingers; (F) minor left second pereopod; (G) same, chela.



# Figure 11

*Mesopontonia kimwoni* sp. nov., ovigerous female pochl 2.5 mm (NIBRIV0000862985), holotype.

(A) left third pereopod; (B) same, dactylus and distal propodus; (C) left fourth pereopod; (D) same, dactylus and distal propodus; (E) left fifth pereopod.



# Figure 12

*Mesopontonia kimwoni* sp. nov. from Munseom Islet, Jeju Island.


Holotype, ovigerous female, pochl 2.5 mm (NIBRIV0000862985). Photographic credit: JH Park.



# Figure 13

Artistic interpretation of *Mesopontonia kimwoni* sp. nov.

Holotype, ovigerous female, pochl 2.5 mm (NIBRIV0000862985). Painting by YK Kim.




# Figure 14

Habitat and host of *Mesopontonia* from Munseom Islet, Jeju Island, Korea.

Habitat and host specimens of *M. verrucimanus* (A, B) and *M. kimwoni* **sp. nov.** (C). (A) host sea whip, *Ellisella* cf. *limbaughii* and habitat in depth of 57 m; (B) host sponge *Raspailia* (*Raspaxilla*) *hirsuta* and habitat in depth of 55 m; (C) host black coral *Myriopathes lata* and habitat in depth of 55m. Photographic Credits: JM Choi.







# Figure 15

Phylogenetic tree obtained by the Maximum likelihood (ML) analysis based on the combined dataset for COI and 16S sequences.

Numbers at nodes represent Maximum Likelihood bootstrap percentage (above) and Bayesian posterior probabilities (below), numbers less than 75% are not shown.

