

Relative availability of natural prey versus livestock predicts landscape suitability for cheetahs *Acinonyx jubatus* in Botswana

Hanlie EK Winterbach, Christiaan W Winterbach, Lorraine Boast, Rebecca Klein, Michael MJ Somers

Prey availability and human-carnivore conflict are strong determinants that govern the spatial distribution and abundance of large carnivore species and determine the suitability of areas for their conservation. For wide-ranging large carnivores such as cheetahs (*Acinonyx jubatus*) suitable conservation areas beyond protected area boundaries are crucial to effectively conserve them both inside and outside protected areas. Although cheetahs prefer preying on wild prey, they also cause conflict with people by predating on especially small livestock. We investigated whether the distribution of cheetahs' preferred prey and small livestock biomass can be used to explore the ~~current~~ potential suitability of agricultural areas in Botswana for the long-term persistence of its cheetah population. We found it gave a good point of departure for identifying priority areas for land management, the threat to connectivity between cheetah populations and areas where the reduction and mitigation of human-cheetah conflict is critical. Our analysis showed the existence of a wide prey base for cheetahs across large parts of Botswana's agricultural areas which provide additional large areas with high conservation potential. Twenty percent wild prey biomass proved to be possibly the critical point to distinguish between high and low predicted levels of human-cheetah conflict. We identified focal areas in the agricultural zones where restoring wild prey numbers in concurrence with effective human-cheetah conflict mitigation efforts are the most immediate conservation strategies needed to maintain Botswana's still large and contiguous cheetah population.

Hanlie E. K. Winterbach Tau Consultants (Pty) Ltd, Maun, Botswana and Centre for Wildlife Management, University of Pretoria, Pretoria, South Africa.

Christiaan W. Winterbach, Tau Consultants (Pty) Ltd, Maun, Botswana and Centre for Wildlife Management, University of Pretoria, Pretoria, South Africa.

Lorraine Boast, Cheetah Conservation Botswana, Private Bag 0457, Gaborone, Botswana.

R. Klein, Cheetah Conservation Botswana, Private Bag 0457, Gaborone, Botswana.

Michael J. Somers Centre for Wildlife Management, University of Pretoria, Pretoria, South Africa and Centre for Invasion Biology, University of Pretoria, Pretoria, South Africa,

Corresponding author: Hanlie E. K Winterbach, Private Bag 83, Maun, tel: +26772782522, tauconsultants@gmail.com

2 The strong linear relationships that exist between the density of African large carnivores and the
 3 biomass of their natural prey (Hayward, O'Brien, & Kerley, 2007) point to prey availability as the
 4 primary natural determinant that governs the spatial distribution and abundance of large carnivore
 5 species and determines the suitability of an area for their conservation (Broekhuis et al., 2013; Fuller &
 6 Sievert, 2001; Gittleman & Harvey, 1982; Hayward, O'Brien, & Kerley, 2007). For competitively
 7 inferior species, such as cheetahs and African wild dogs (*Lycaon pictus*), interspecific competition,
 8 especially from lions (*Panthera leo*) and spotted hyaenas (*Crocuta crocuta*), can also exert a strong
 9 influence on their movements, behaviour and density (Durant, 2000; Van der Meer et al., 2013). This
 10 occurs predominantly inside protected areas where densities of lions and spotted hyaenas tend to be
 11 high (Creel, Spong & Creel, 2001). In human-dominated landscapes human activities and their conflict
 12 with predators are often as strong a determinant factor as prey availability in the occurrence and
 13 survival of large carnivores (Gusset et al., 2009; Marker et al., 2003; Schuette, Creel & Christianson,
 14 2013; Woodroffe & Ginsberg, 1998) With few protected areas large enough to contain the wide-
 15 ranging behaviour of large carnivores, the management of suitable conservation areas beyond protected
 16 area boundaries are necessary to effectively conserve large carnivore species both inside and outside
 17 protected areas (Woodroffe & Ginsberg, 1998). This requires the assessment of both the distribution
 18 and abundance of suitable prey and the potential levels of human-carnivore conflict, often on a large
 19 geographic scale.

20 The cheetah is Africa's most endangered felid (Marker et al., 2007) and is listed as Vulnerable with
 21 a declining population trend by the IUCN Red List of Threatened Species (Durant et al., 2008). They
 22 are one of the most wide-ranging terrestrial carnivores and need extensive areas to sustain viable
 23 populations (IUCN/SCC, 2007). Cheetahs feed on a diverse wild prey base ranging from animals as
 24 small as scrub hares (*Lepus saxatilis*) to as large as zebras (*Equus quagga*), but generally select the
 25 most abundant locally available prey up to 135 kg with a strong preference for those with a body mass

26 between 14 kg – 40 kg (Clements et al., 2014). At least 75% of the cheetah's resident range in southern
 27 Africa falls outside protected areas (IUCN/SSC, 2007). In southern Africa this falls mostly on
 28 farmlands where competition with other large carnivores is low and a sufficient small to medium-sized
 29 wild prey base still occurs (Lindsey & Davies-Mostert, 2009; Klein, 2007; Marker & Dickman, 2004).
 30 Consequently, conservation efforts in human-dominated landscapes are critical for this species long-
 31 term survival. Although cheetahs prefer ~~preying on~~ wild prey (Marker et al., 2003), they also cause
 32 conflict with people by predating on livestock which generally involves small stock (sheep and goats)
 33 and occasionally calves and foals (Marker et al., 2003; Ogada et al., 2003; Selebatso, Moe & Swenson,
 34 2008; Woodroffe et al., 2007).

35 Botswana is important for the regional and global long-term survival of cheetahs. It has a large and
 36 still contiguous cheetah population and hosts the second largest national population with $\pm 1\,786$
 37 animals (Klein, 2007) after Namibia with $\pm 3\,138 - 5\,775$ animals (Hanssen & Stander, 2004; Marker
 38 et al., 2007). It also forms the major connecting range for the southern African cheetah population
 39 which is largest known free-ranging resident cheetah population comprising $\pm 6\,500$ animals
 40 (IUCN/SCC, 2007). Around half of the Botswana cheetah population occurs outside conservation
 41 areas on rangeland (Winterbach & Winterbach, 2003). In 2009, a Draft National Conservation Action
 42 Plan for Cheetahs and African Wild Dogs in Botswana was prepared by the Department of Wildlife
 43 and National Parks (DWNP). One of the primary targets set out in the national plan is obtaining
 44 quantitative knowledge regarding the main threats to securing a viable cheetah population across its
 45 range in Botswana. However, scientific information on cheetah distribution and density on a country-
 46 wide scale is nearly impossible to obtain, but the urgent conservation status of threatened species such
 47 as cheetahs can ill-afford to wait for detailed scientific information before policy decisions are made.
 48 Therefore, an objective, clear system of evaluation, based on the best available and reliably correlated
 49 information, is needed that can be used as a basis to support policy setting (Theobald et al., 2000) 

50 We used the distribution of cheetahs' preferred prey and small stock biomass as the most essential
 51 components to explore the **current potential suitability** of agricultural areas for the long-term
 52 persistence of cheetahs in Botswana. The percentage prey biomass of small stock and prey biomass
 53 combined ~~was~~ used as the primary indicator of probable levels of human-cheetah conflict. **Our analyses**
 54 **demonstrated that wild prey combined with livestock can provide a country-wide overview of the**
 55 **suitability of the Botswana landscape for cheetahs without the use of complex modelling. It allows for**
 56 **prudent conclusions as a point of departure for identifying specifically priority areas for land**
 57 **management, the threat to connectivity between cheetah populations and areas where the reduction and**
 58 **mitigation of human-cheetah conflict is critical.**

59

60 Study Area

61

62 The Republic of Botswana is ca 582 000 km² in size and is landlocked with Namibia, South Africa,
 63 Zimbabwe, and Zambia as its neighbours. Roughly 50% of its 2 million people (3.5 people / km²) live
 64 in rural villages and small settlements (Central Statistics Office, 2014).

65 The mean altitude above sea level is 1 000 m (515 – 1 491 m a.s.l.). The climate is arid to semi-arid
 66 with highly variable rainfall and periodic~~al~~ severe droughts. Mean annual rainfall varies from 650 mm
 67 in the north-east to 250 mm in the south-west. Average maximum daily temperatures range from 22° C
 68 in July to 33° C in January and average minimum temperatures from 5° C to 19° C respectively
 69 (Department of Surveys and Mapping, 2001). Only two perennial rivers occur; the Okavango River
 70 which fans out into the Okavango Delta and the Kwando/ Linyanti/Chobe river system which forms the
 71 boundary with Namibia and Zambia. The Makgadikgadi Pans is a seasonal wetland with natural
 72 perennial water holes in the Boteti River providing critical dry season water sources for wildlife in
 73 Makgadikgadi National Park (MNP). Across the rest of the country, scattered pans and ancient

74 riverbeds periodically hold water during the wet season. Considerable seasonal variations in the density
 75 and distribution of ungulate species occur and the blocking of migration routes by veterinary fences has
 76 led to ungulate die-offs during drought years (Bergström & Skarpe, 1999; Verlinden, 1998). Seasonal
 77 migrations of Burchell's zebra and blue wildebeest (*Connochaetes taurinus*) still occur inside MNP
 78 (Brooks, 2005), and zebra migrate between MNP and the Okavango Delta (Bartlam-Brooks, Bonyongo
 79 & Harris, 2011).

80 Approximately 38% of the land use in Botswana is designated for wildlife utilization; 17% as
 81 protected areas (national parks and game reserves) and 21% as Wildlife Management Areas (WMAs)
 82 (Figure 1). WMAs are primarily designed for wildlife conservation, utilisation, and management
 83 (Hachileka, 2003), however, unlike protected areas; people are permitted to reside within WMAs and
 84 to own and graze livestock there. Protected areas and WMAs do not have 'predator-proof' fences, with
 85 the exception of the western and southern boundary of MNP which provides only a partial barrier due
 86 to its poor upkeep.

87 Five percent of the country is residential areas and 57% consists of rangeland (of which roughly
 88 70% is tribal / communal grazing land), 25% is state land, 5% freehold land leased for large-scale
 89 commercial ranching (Department of Surveys and Mapping, 2001). In the Draft National Predator
 90 Strategy (Winterbach & Winterbach, 2003), the country was sub-divided into two main predator
 91 management zones; Conservation Zones comprising of national parks, forest reserves, sanctuaries and
 92 WMAs, and the Agricultural Zones consisting of rangelands, residential and mining areas (Figure 1).

93 Livestock (mainly cattle) rearing is the primary economic activity over large parts of Botswana and
 94 constitutes 70 - 80% of the agricultural GDP (Botswana Ministry of Agriculture 2011). In the 2012
 95 household survey, the livestock population in Botswana was estimated as 2.6 million cattle, 1.8 million
 96 goats and 300,000 sheep, most of which were located on the more fertile eastern side of the country
 97 (Botswana Ministry of Agriculture 2011). Approximately 92% of this livestock are in the traditional

98 cattle post system on communal grazing land (Botswana Ministry of Agriculture 2011). Botswana's
99 key environmental issues include water scarcity and pollution, rangeland degradation and
100 desertification, loss of biodiversity, deforestation, and an increased frequency of periodic droughts
101 (Wingqvist & Dahlberg, 2008).

102  Botswana, cheetahs are a protected species. Before 2000, cheetahs could be hunted or captured
103 under and in accordance with the terms and conditions of a Director's permit (Wildlife Conservation
104 and National Parks Act (Act No. 28 of 1992)). However, since 2000, a statutory instrument disallowed
105 the killing of cheetahs for any reason (Botswana Government Gazette, 2000), and in 2005, a US\$ 113
106 fine or term in prison was added (Botswana Government Gazette, 2005), although the latter is rarely
107 enforced. Botswana has an unutilized CITES quota of five cheetah per annum (Klein, 2007).

108

109 **Methods**

110

111  The first parameter we used to determine landscape suitability was the biomass of wild prey species
112 which occur in Botswana **that are preferred prey** or **in the preferred weight range** (body mass 14 kg –
113 40 kg) for cheetahs (Clements et al., 2014) (Table 1), hereafter termed 'wild prey'. We included Red
114 lechwe (*Kobus lechwe*)  as one of the preferred prey of cheetah in the seasonal floodplains of the
115 Okavango Delta (pers. com. ). We did not include calves of the larger wild prey since population
116 numbers were collected during the dry season when calves were not prevalent. The second parameter
117 we used was the biomass of goats and sheep (herein referred to as 'small stock') as the main livestock
118 whose depredation is a significant predictor of human-cheetah conflict levels  Supplementary material
119 Appendix 1). The third parameter was the percentage that wild prey biomass contributed to the total
120 biomass of wild prey and small stock combined to indicate probable levels of conflict  (herein referred
121 to as **'percentage wild prey'**).

122 We calculated wild prey and small stock biomass in Large Stock Units (LSU = body weight^{0.75})
123 per 1 grid cells from the combined data of six country-wide annual dry season aerial surveys
124 conducted by DWNP between 2001 and 2005 and in 2007. Aerial surveys during the drought years
125 before 2001 and very wet years after 2007 were excluded. Cheetah biomass strongly correlates with
126 lean season prey biomass (Fuller & Sievert, 2001) and we felt the six aerial surveys used in this study
127 best represented the general distribution of prey and small stock biomass on a country-wide scale.
128 Although aerial surveys tend to undercount small mammal species, such as steenbok (*Raphicerus*
129 *campestris*) and duiker (*Sylvicapra grimmia*), it is the only feasible method for wildlife monitoring on
130 a country-wide scale. We assumed that using a combined data set from six aerial surveys was sufficient
131 to determine the general distribution of wild prey and small stock biomass across Botswana.

132 We utilized the broad landscape suitability stratification for large carnivores in Botswana from
133 (Winterbach et al., 2014) and refined it in the agricultural zones based on the distribution of wild prey
134 biomass, small stock biomass and the percentage wild prey biomass to identify homogeneous strata. To
135 determine if there is a critical percentage wild prey that can be used to differentiate between high and
136 low probable levels of conflict, we used data on livestock attacks by cheetahs between 1995 and 2006
137 consisting of problem animal conflict reports and farms questionnaire surveys ($N = 188$) conducted
138 during 2004 and 2005. We calculated the number of aerial survey grids with livestock attacks in
139 different categories of percentage wild prey biomass in the Kgalagadi, Ghanzi Agricultural Zones and
140 the western strata of the Central Agricultural Zone. From this we calculated the frequency of livestock
141 attacks per grid cell for each category (Supplementary material Appendix 2) and used a chi-square test
142 to test whether the observed frequency of conflict reports has the same frequency as the grids per
143 category of percentage wild prey. Observations were classified into categories independently and all
144 categories had expected frequencies $> 5\%$. We used Bonferroni intervals (Byers, Steinhorst &

145 Krausman, et al. 1984) to test for categories with observed frequencies that differed significantly from
146 the expected.

147 To test our classification of suitable and unsuitable areas for cheetahs we investigated the reported
148 presence, transience and absence of cheetahs using questionnaire surveys ($N = 89$) conducted during
149 2012 and 2013 that targeted primarily game ranchers and commercial livestock farmers in the game
150 ranching regions of the Central, Ghanzi, Ngamiland and North East regional districts. Farmers were
151 asked to record the status of cheetahs on their property as present (visual sightings or tracks seen at
152 least quarterly), transient (visual sightings or tracks seen less frequently than quarterly) or absent
153 (never seen cheetahs or its tracks). We used a chi-square test with Bonferroni simultaneous confidence
154 intervals (Byers, Steinhorst & Krausman, et al. 1984) to test the hypothesis that farmers reported
155 cheetahs as present, transient or absent in similar proportions on grid cells with different percentage
156 wild prey categories. We used the Natural Breaks (Jenks) function in ArcMap 9.3.1 that best grouped
157 similar values and maximize the differences between groups to identify categories of landscape
158 suitability for the long-term persistence of cheetahs. We identified five categories of suitability, based
159 on the proportion of grid cells in each sub-stratum that had $\leq 20\%$ wild prey.

160

161 Result

162

163 The distribution of cheetah wild prey biomass and the percentages of wild prey biomass in the different
164 categories are shown in Figure 2 and Figure 3, respectively. Although cheetah wild prey occurred
165 widely in the agricultural zones, it contributed only 0 – 5% of the total biomass (wild prey plus small
166 stock) available to cheetahs in the eastern parts of both the Central Agricultural Zone and the
167 Kgalagadi Agricultural Zone 

168 The percentage wild prey biomass in the Kgalagadi and Ghanzi Agricultural Zones and the western
169 strata of the Central Agricultural Zone combined was $\leq 20\%$ in 235 of the 403 grids (58.3%) and $>$
170 80% in 136 grids (33.7%) with the remaining 32 grids (8%) between $> 20\%$ and $\leq 80\%$. The number of
171 conflict reports recorded per grid for the percentage wild prey biomass intervals ranged from 0.23 to
172 0.81 reports per grid cell with a mean of 0.49 and standard error of 0.25 ($N = 8$) (Table 2). The number
173 of conflict reports was consistently below the mean when the percentage wild prey biomass exceeded
174 20% (Figure 3).

175 We subsequently selected 0% , > 0 to $\leq 20\%$, > 20 to $\leq 80\%$, and > 80 to $\leq 100\%$ as the main
176 categories of percentage wild prey biomass. The observed frequency of conflict reports in grids from
177 the separate categories differed significantly from the expected ($\chi^2 = 52.42$, $df = 1$, $P < 0.001$). Conflict
178 reports were more frequently than expected ($P = 0.05$) in grids with 0% wild prey biomass, and
179 significantly lower than expected in areas with $> 20\%$ wild prey biomass ($\alpha = 0.05$, $Z = 2.4977$) (Table
180 3). We therefore took grids with $\leq 20\%$ wild prey biomass as representing areas with high predicted
181 levels of conflict, and $> 20\%$ wild prey biomass as areas with low predicted levels of conflict.

182 We rated the five categories of landscape suitability identified in ArcMap as very high ($0 - 6.7\%$
183 grid cells with $\leq 20\%$ wild prey), high ($6.8 - 25\%$), medium ($25.1 - 50$), low ($50.1 - 75\%$) or
184 unsuitable ($75.1 - 100\%$) and provide a country-wide landscape suitability map for the long-term
185 persistence of cheetahs in Botswana (Figure 4) (Supplementary material Appendix 3). The
186 conservation zones were the most suitable for the long-term persistence of cheetahs, while the
187 agricultural zones consisted of a mosaic of medium suitability to unsuitable. The classification of some
188 strata as unsuitable for cheetahs was supported by the questionnaire data where the proportion of
189 farmers that reported cheetahs present or absent differed significantly between suitable and unsuitable
190 cheetah areas ($\chi^2 = 129.11$, $df = 3$, $P < 0.001$). A hundred per-cent of farmers reported cheetahs absent

191 in the unsuitable areas ($n = 10$) significantly more than would be expected by chance and only 13.9 %
 192 of farmers reported cheetahs absent within the suitable areas ($N = 79$) ($\alpha = 0.05$, $Z = 2.4977$) (Table 4).

193

194 Discussion

195

196 Our results show that the distribution of cheetah wild prey and small stock biomass can provide a good
 197 information basis to evaluate the landscape suitability for cheetahs on a country-wide scale and
 198 indicate priority areas for conservation actions. In Botswana, 20% wild prey biomass showed to be a
 199 potentially critical point to distinguish between high and low predicted levels of human-cheetah
 200 conflict. The distribution of the categories of percentage wild prey biomass (Figure 3) clearly
 201 highlights areas where locally-adapted conflict mitigation strategies are a priority, for example along
 202 the western and eastern boundaries of the Okavango Delta. In addition, the landscape suitability map
 203 (Figure 4) shows strata where currently the long-term persistence of cheetahs is highly unlikely, such
 204 as in the eastern part of the Central Agricultural Zone, and where connectivity within the cheetah
 205 population is threatened, such as between the Northern and Southern Conservation Zones.

206 The distribution of cheetah wild prey biomass provides a wide prey base across large parts of the
 207 agricultural areas in Botswana. In fact, the greater resource availability in Botswana may be causal to
 208 the considerably higher density, smaller home range sizes and generally larger body size of cheetahs in
 209 Botswana compared to Namibia (Boast et al., 2013). The significant reduction in conflict reports in
 210 areas with $> 20\%$ wild prey biomass confirms adopting an integrated livestock-wildlife management
 211 approach in the communal rangelands of Botswana as an effective conflict mitigation strategy to
 212 maintain key areas in the agricultural zones for cheetah conservation.

213 Almost 50% of Botswana's cheetah population occurs in agricultural areas (Klein, 2007). Support
 214 from both livestock and game farmers for cheetah conservation outside protected areas is low in

215 Botswana (Selebatso, Moe & Swenson, 2008) and retaliatory killing of cheetahs is considered to be
216 widespread (Klein, 2007). Livestock farmers view cheetahs as the fourth most problematic predator,
217 following leopards (*Panthera pardus*), jackals (*Canis mesomelas*) and wild dogs, while sixty percent
218 (60%) of private game farmers rate them as the second top stock predators (Selebatso, Moe &
219 Swenson, 2008). As in Namibia (Marker et al., 2003), cheetahs in Botswana prey predominantly on
220 local native game (Cheetah Conservation Botswana unpubl data). This preference for wild prey has
221 also been shown for other large carnivores even in areas where livestock is predominant (Hemson,
222 2003; Ogara et al., 2010; Woodroffe et al., 2005), and maintaining wild prey populations within
223 livestock areas is viewed as a feasible way to decrease livestock depredation (de Azevedo & Murray,
224 2007; Mizutani, 1999).

225 Steenbok and duiker are two of the generally most common prey for cheetahs in Botswana (Klein,
226 2007). The high density of steenbok and duiker (range 0.261 - 4.319 animals / 100 km²) calculated
227 from the six aerial surveys in the Ghanzi Community Stratum (stratum 7.2) suggests that livestock do
228 not necessarily displace small ungulates to the extent that large ungulates are displaced (Riginos et al.,
229 2012). However, conflict with people seems to have a stronger influence on cheetah numbers than wild
230 prey biomass when livestock numbers are high. In the Ghanzi Farms Stratum (stratum 7.1) there was a
231 sample point where cheetah density was too low to calculate despite having more wild prey than
232 neighbouring areas with 0.7 cheetahs / 100 km² (Boast & Houser, 2012; Kent, 2010). This shows the
233 potential negative impact of human conflict on the cheetah population even where there is a good wild
234 prey base but it formed a small percentage to the total biomass due to the high small stock biomass.

235 The Okavango Delta is nearly surrounded by two agricultural strata (strata 6.2 and 6.4 in Figure 4)
236 that are currently unsuitable for the long-term persistence of cheetahs. Wild prey biomass is low and
237 PAC reports show conflict levels between farmers and cheetahs are high. Together with the low
238 suitability of agricultural stratum 6.3, the free movement of cheetahs from the Northern Conservation

239 Zone towards the Central Kalahari Game Reserve (CKGR) is impeded and we therefore assumed that
 240 cheetahs found in these strata are most likely transients. A second linkage between the Northern and
 241 Southern Conservation Zones extends from MNP to the south-west across the most northern part of the
 242 Central Agricultural Zone. PAC reports indicate cheetah presence in this corridor (Figure 4). Concerted
 243 efforts in conflict mitigation to ensure functional corridors are essential to maintain connectivity
 244 between the smaller northern cheetah population and that of the south, especially in the light that
 245 (Dalton et al., 2013) found that the northern cheetah population showed some degree of genetic
 246 isolation. However, the small sample size ($N = 4$) of cheetahs from the north and its isolation by
 247 distance from the rest of the samples may have contributed to their findings. Our study emphasises the
 248 urgent need for intensive genetic studies to accurately determine gene flow across the country.

249 In the Southern Conservation Zone, the WMAs in the Western Kgalagadi Conservation Corridor
 250 (WKCC) connect the CKGR to the Kgalagadi Transfrontier Park (KTP), Botswana's two largest
 251 nationally protected areas (Conservation International Botswana, 2010). The Schwelle, which lies
 252 south within the WKCC, provides crucial wet season forage to ungulates and is one of the most
 253 important wildlife areas in Botswana, preserving the Kalahari Ecosystem (Anonymous, 2008). The
 254 Schwelle also holds almost half of the cheetahs in the Southern Conservation Zone (Klein, 2007). The
 255 increasing demand for livestock grazing areas and already extensive use of parts of the WKCC for
 256 cattle production is a concern in maintaining this corridor for large predators and wildlife in general.
 257 For example, the proposed changing of the land use from wildlife to cattle in the eastern section of the
 258 WKCC (Anonymous, 2008) has the potential to enlarge the unsuitable areas for cheetahs across the
 259 Kgalagadi Agricultural Zones 1 and 2, and PAC reports show human-cheetah conflict is already
 260 widespread here. The implementation of effective conflict mitigation strategies will be essential to
 261 prevent the formation of a wide barrier to the free movement of cheetahs both between the CKGR and
 262 KTP, and further south connecting with the South African cheetah population.

263 The Dry North (stratum 3.1) provides a linkage with Hwange National Park in Zimbabwe. The
 264 importance of this linkage lies in that it enlarges the northern Botswana cheetah population. The aerial
 265 surveys showed wild prey biomass in the central part of the stratum overall to be low, to zero in some
 266 grid cells. However, large parts of this area are dominated by Miombo and mopane (*Colophospermum*
 267 *mopane*) woodland and close-tree *Acacia* savannah (Department of Surveys and Mapping, 2001). The
 268 low wild prey biomass is probably a function of the limitation of aerial surveys to detect small, cryptic
 269 species in these dense habitats (Jachmann, 2002). A high density of small wild prey, especially duiker
 270 and steenbok, was recorded in the western part of this area during a ground survey done in 2011
 271 (Winterbach, unpubl. data). Limiting the uncontrolled development of artificial water points in this
 272 stratum is an important conservation strategy for cheetahs as the wide-spread availability of water may
 273 increase large ungulate numbers leading to a corresponding increase in lions and spotted hyaenas that
 274 are dominant competitors of cheetahs (Creel, Spong & Creel, 2001; Durant, 2000; Mills & Gorman,
 275 1997).

276 Namibia and Botswana protect approximately 77% of the southern African cheetah population
 277 (IUCN/SCC, 2007). The most important linkage between the Namibian and Botswana population lies
 278 in the Ghanzi Agricultural Zone, where landscape suitability for cheetahs ranges between medium on
 279 the commercial farms (stratum 7.1) to low on the community farms (stratum 7.2). The proposed re-
 280 alignment of the western boundary of the WKCC to enlarge the communal grazing area for the cattle
 281 industry (Anonymous, 2008) will not only further threaten the functionality of the WKCC for wildlife,
 282 but also potentially threaten the connectivity with the Namibian cheetah population.

283 Conservation of free-ranging cheetah populations is multi-faceted and needs to be addressed from
 284 an ecological, biological and socio-economic management perspective. Despite the threats, Botswana
 285 has a large and still contiguous cheetah population with wide-spread natural movements allowing
 286 substantial gene flow (Dalton et al., 2013). However, its contiguous nature is threatened and may cease

if corridors are not maintained. Cheetahs have vast tracts of intact habitat in the conservation areas for persistence, and the wide-spread availability of wild prey across the agricultural zones provides additional large areas with high conservation potential. On a micro-scale, some studies found habitat structure, such as dense woodland and open savannah, have an even stronger effect on the areas the different social groups of cheetahs prefer to utilize than absolute wild prey density (Bissett & Bernard, 2007; Muntifering et al., 2006). Spatial data indicating land degradation were not available to include in the landscape suitability map. However, the wide-spread rangeland degradation and desertification in the agricultural areas of Botswana (Moleele et al., 2002) reduces the availability of suitable wild prey, as well as sufficient grass cover for cheetahs for stalking, concealment from other predators, and movement between areas (Broomhall, Mills & du Toit, 2003; Marker, 2003; Mills, Broomhall & du Toit, 2004; Purchase & du Toit, 2000). This study showed that restoring wild prey numbers in focal areas in concurrence with effective human-cheetah conflict mitigation efforts are the most immediate conservation strategies needed to ensure the long-term survival of cheetahs in Botswana. With this is the restoration of degraded rangeland which will not only recover habitat for cheetahs and their wild prey but also benefit farmers by increasing the carrying capacity for livestock, which in itself may increase their tolerance for cheetahs on their land (Marker, 2003).

Acknowledgements

Thanks to Laurie Marker, J. W. McNutt, E. Fabiano and R. Kotze for valuable comments on the manuscript. This manuscript was made possible by the Department of Wildlife and National Parks, Ministry of Environment, Wildlife and Tourism, Gaborone, Botswana.

References

311 **Anonymous. 2008.** Review of the national land use map. Final Report. LANDflow Solutions (Pty)
 312 Ltd, Gaborone.

313 **Conservation International Botswana. 2010.** Consultancy to identify important habitats for key
 314 wildlife in the Western Kgalagadi Conservation Corridor (WKCC). Conservation International
 315 Botswana, Gaborone.

316 **Bartlam-Brooks HLA, Bonyongo MC, Harris S. 2011.** Will reconnecting ecosystems allow long-
 317 distance mammal migrations to resume? A case study of a zebra *Equus burchelli* migration in
 318 Botswana. *Oryx* **45**:210-216.

319 **Bergström R, Skarpe C. (1999).** The abundance of large wild herbivores in a semi-arid savanna in
 320 relation to seasons, pans and livestock. *African Journal of Ecology* **37**:12-26.

321 **Bissett C, Bernard TF. 2007.** Habitat selection and feeding ecology of the cheetah (*Acinonyx jubatus*)
 322 in thicket vegetation: is the cheetah a savanna specialist? *Journal of Zoology* **271**:310-317.

323 **Boast LK, Houser A. 2012.** Density of large predators on commercial farmland in Ghanzi, Botswana.
 324 *South African Journal of Wildlife Research* **42(2)**:138-143..

325 **Boast LK, Houser AM, Good K, Gusset M. 2013.** Regional variation in body size of the cheetah
 326 (*Acinonyx jubatus*). *Journal of Mammalogy* **94(6)**:1293-1297.

327 **Botswana Government Gazette. 2000.** Statutory Instrument No. 69 of 2000 Restricting the Killing of
 328 Lions and Cheetahs by both Public and Wildlife Staff. Government of Botswana, Gaborone.

329 **Botswana Government Gazette. 2005.** Statutory Instrument No 27 of 2005 killing of cheetahs
 330 suspension order. Ministry of Wildlife , Environment and Tourism. Government of Botswana,
 331 Gaborone.

332 **Botswana Ministry of Agriculture. 2011.** Livestock sector. *Available at*
 333 *http://www.moa.gov.bw/downloads_agricultural_marketing_strategy.pdf.*

334 **Broekhuis F, Cozzi G, Valeix M, McNutt JW, Macdonald DW. 2013.** Risk avoidance in sympatric
 335 large carnivores: reactive or predictive? *Journal of Animal Ecology* **82**:1098-1105.

336 **Brooks CJ. 2005.** The foraging behaviour of Burchell's zebra (*Equus burchelli antiquorum*). D. Phil.
 337 Thesis, University of Bristol.

338 **Broomhall LS, Mills MGL, du Toit JT. 2003.** Home range and habitat use by cheetahs (*Acinonyx*
 339 *jubatus*) in the Kruger National Park. *Journal of Zoology* **261**:119-128.

340 **Byers R, Steinhorst RK, Krausman PR. 1984.** Clarification of a technique for analysis of utilization-
 341 availability data. *Journal of Wildlife Management* **48**:1050-1053.

342 **Central Statistics Office. 2014.** Population and housing census 2011 analytical report. Available at
 343 http://www.cso.gov.bw/images/analytical_report.pdf.

344 **Clements HS, Tambling CJ, Hayward MW, Kerley GIH. (2014).** An objective approach to
 345 determining the weight ranges of prey preferred by and accessible to the five large African carnivores.
 346 *PLOS ONE* **9(7)**: e101054. Available at <http://www.plosone.org>.

347 **Creel S, Spong G, Creel NM. 2001.** Interspecific competition and the population biology of
 348 extinction-prone carnivores. In: Gittleman JL, Funk SM, Macdonald DW, Wayne RK eds. *Carnivore*
 349 *Conservation*. Cambridge University Press, 35-60.

350 **Dalton DL, Charruau P, Boast L, Kotzé A. 2013.** Social and genetic population structure of free-
 351 ranging cheetah in Botswana: implications for conservation. *European Journal of Wildlife Research*
 352 **59**:281-285.

353 **de Azevedo FCC Murray DI. 2007.** Evaluation of potential factors predisposing livestock to
 354 predation by jaguars. *Journal of Wildlife Management* **71**:2379-2386.

355 **Department of Surveys and Mapping. 2001.** Botswana National Atlas. Government of Botswana.
 356 Available at <http://www.atlas.gov.bw>

- 357 **Durant S, Marker L, Purchase N, Belbachir F, Hunter L, Packer C, Breitenmoser-Würsten C,**
358 **Sogbohossou E, Bauer H. 2008.** *Acinonyx jubatus*. Available at <http://www.iucnredlist.org> .
- 359 **Durant SM. 2000.** Living with the enemy: avoidance of hyenas and lions by cheetahs in the Serengeti.
360 *Behavioral Ecology* **11**:624-632.
- 361 **Fuller TK, Sievert PR. 2001.** Carnivore demography and the consequences of changes in prey
362 availability. In: Gittleman JL, Funk SM, Macdonald DW, Wayne RK eds. *Carnivore Conservation*.
363 Cambridge University Press, 163-178.
- 364 **Gittleman JL Harvey P H. 1982.** Carnivore home range size, metabolic needs and ecology.
365 *Behavioural Ecology and Sociobiology* **10**:57-63.
- 366 **Gusset M, Swarner MJ, Mponwane L, Keletile K, McNutt JW. 2009.** Human-wildlife conflict in
367 northern Botswana: livestock predation by endangered African wild dog *Lycaon pictus* and other
368 carnivores. *Oryx* **43**:67-72.
- 369 **Hachileka E. 2003.** Sustainability of wildlife utilization in the Chobe District, Botswana. *South*
370 *African Geographic Journal* **85**:50-57.
- 371 **Hanssen L, Stander P. 2004.** Namibia Large Carnivore Atlas. Available at
372 http://www.desertlion.info/reports/atlas_July2004.pdf.
- 373 **Hayward MW, O'Brien J, Kerley GIH. 2007.** Carrying capacity of large African predators:
374 predictions and tests. *Biological Conservation* **139**:219-229.
- 375 **Hemson G. 2003.** The ecology and conservation of lions: human-wildlife conflict in semi-arid
376 Botswana. D. Phil. Thesis, University of Oxford.
- 377 **Jachmann H. 2002.** Comparison of aerial counts with ground counts for large African herbivores.
378 *Journal of Applied Ecology* **39**:841-852.
- 379 **Kent VT. 2010.** The status and conservation potential of carnivores in semi-arid rangelands, Botswana.
380 D. Phil. Thesis, Durham University.

381 **Klein R. 2007.** Status report for cheetah in Botswana. In: Breitenmoser C, Breitenmoser U, Durant S
 382 eds. Status and conservation needs of cheetahs in southern Africa. IUCN/Cat Specialist Group, 14-21.

383 **Lindsey PA, Davies-Mostert HT. 2009.** South African action plan for the conservation of cheetahs
 384 and African wild dogs. Report from a national conservation action planning workshop. IUCN Cat
 385 Specialist Group, 17-19.

386 **Marker L. 2003.** Aspects of cheetah (*Acinonyx jubatus*) biology, ecology and conservation strategies
 387 on Namibian farmlands. D. Phil. Thesis, University of Oxford.

388 **Marker L, Dickman A. (2004).** Human aspects of cheetah conservation: lessons learned from the
 389 Namibian farmlands. *Human Dimensions of Wildlife* **9**:297-305.

390 **Marker L, Dickman A, Wilkinson C, Schumann B, Fabiano E. 2007.** The Namibian cheetah: status
 391 report. In: Breitenmoser C, Breitenmoser U, Durant S eds. Status and conservation needs of cheetahs in
 392 southern Africa. IUCN Cat Specialist Group, 4-13.

393 **Marker LL, Muntifering JR, Dickman AJ, Mills MGL, Macdonald DW. 2003.** Quantifying prey
 394 preferences of free-ranging Namibian cheetahs. *South African Journal of Wildlife Research* **33**:43-53.

395 **Mills MGL, Broomhall LS, du Toit JT. 2004.** Cheetah *Acinonyx jubatus* feeding ecology in the
 396 Kruger National Park and a comparison across African savanna habitats: is the cheetah only a
 397 successful hunter on open grassland plains? *Wildlife Biology* **10**:177-186.

398 **Mills MGL, Gorman ML. 1997.** Factors affecting the density and distribution of wild dogs in the
 399 Kruger National Park. *Conservation Biology* **11**:1397-1406.

400 **Mizutani F. 1999.** Impact of leopards on a working ranch in Laikipia, Kenya. *African Journal of*
 401 *Ecology* **37**:211-225.

402 **Moleele NM, Ringrose S, Matheson W, Vanderpost C. 2002.** More woody plants? The status of
 403 bush encroachment in Botswana's grazing areas. *Journal of Environmental Management* **64**:3-11.

- 404 **Muntifering JR, Dickman AJ, Perlow LM, Hruska T, Ryan PG, Marker LL, Jeo RM. 2006.**
 405 Managing the matrix for large carnivores: a novel approach and perspective from cheetah (*Acinonyx*
 406 *jubatus*) habitat suitability modelling. *Animal Conservation* **9**:103-112.
- 407 **Ogada MO, Woodroffe R, Oguge NO, Frank LG. 2003.** Limiting depredation by African
 408 carnivores: the role of livestock husbandry. *Conservation Biology* **17**:1521-1530.
- 409 **Ogara WO, Gitahi NJ, Andanje SA, Oguge N, Nduati DW, Mainga AO. 2010.** Determination of
 410 carnivores prey base by scat analysis in Samburu community group ranches in Kenya. *African Journal*
 411 *of Environmental Science and Technology* **4**:540-546.
- 412 **Purchase GK, du Toit JT. 2000.** The use of space and prey by cheetahs in Matusadona National Park,
 413 Zimbabwe. *South African Journal of Wildlife Research* **30**:139-144.
- 414 **Riginos C, Porensky LM, Veblen KE, Odadi WO, Sensenig R, Kimuyu D, Keesing F, Wilkerson**
 415 **ML, Young TP. 2012.** Lessons on the relationship between livestock husbandry and biodiversity from
 416 the Kenya Long-term Exclosure Experiment (KLEE). *Pastoralism* **22**:1-22.
- 417 **Schuetz P, Creel C, Christianson D. 2013.** Coexistence of African lions, livestock, and people in a
 418 landscape with variable human land use and seasonal movements. *Biological Conservation* **157**:148-
 419 154.
- 420 **Selebatso M, Moe SR, Swenson JE. 2008.** Do farmers support cheetah *Acinonyx jubatus* conservation
 421 in Botswana despite livestock depredation? *Oryx* **42**:430-436.
- 422 **Theobald DM, Hobbs NT, Bearly T, Zack JA, Shenk T, Riebsame WE. 2000.** Incorporating
 423 biological information in local land-use decision making: designing a system for conservation
 424 planning. *Landscape Ecology* **15**:35-45.
- 425 **Van der Meer E, Mpofu J, Rasmussen GSA, Fritz H. 2013.** Characteristics of African wild dog
 426 natal dens selected under different interspecific predation pressures. *Mammalian Biology* **78**:336-343.

- 427 **Verlinden A. 1998.** Seasonal movement patterns of some ungulates in the Kalahari ecosystem of
428 Botswana between 1990 and 1995. *African Journal of Ecology* **36**:117-128.
- 429 **Wingqvist GÖ, Dahlberg E 2008.** Botswana Environmental and Climate Change Analysis. University
430 of Gothenburg, Stockholm.
- 431 **Winterbach CW Winterbach HEK. 2003.** Draft predator management strategy. Department of
432 Wildlife and National Parks, Gaborone, Botswana.
- 433 **Winterbach HEK, Winterbach CW, Somers MJ. 2014.** Landscape suitability in Botswana for the
434 conservation of its six large African carnivores. *PLOS ONE* **9(6)**:e100202 . Available at
435 <http://www.plosone.org> .
- 436 **Woodroffe R, Frank LG, Lindsey PA, ole Ranah MK, Romañach S. (2007).** Livestock husbandry
437 as a tool for carnivore conservation in Africa's community rangelands: a case-control study.
438 *Biodiversity Conservation* **16**:1245-1260.
- 439 **Woodroffe R, Ginsberg JR. 1998.** Edge effects and the extinction of populations inside protected
440 areas. *Science* **280**:2126-2128.
- 441 **Woodroffe R, Lindsey P, Romañach S, Stein A, ole Ranah SMK. 2005.** Livestock predation by
442 endangered African wild dogs (*Lycaon pictus*) in northern Kenya. *Biological Conservation* **124**:225-
443 234.
- 444




Figure 1. Land use categories present in Botswana






Figure 2. The distribution of cheetah prey biomass (LSU) across Botswana.




Figure 3. The distribution of the percentage cheetah prey biomass (LSU) of small stock and prey biomass combined across Botswana.




Figure 4. Map of the landscape suitability of Botswana for the long-term persistence of cheetah



Table 1. Wild prey species occurring in Botswana and identified as within the preferred weight range of cheetah's prey (body mass 14 – 135 kg, Clements et al, 2014).

<i>Species</i>	<i>Scientific name</i>	<i>Weight (kg)</i>	<i>LSU conversion</i>
Red lechwe	<i>Kobus leche</i>	72	0.25298
Ostrich	<i>Struthio camelus</i>	68	0.24237
Warthog	<i>Phacochoerus africanus</i>	45	0.17783
Impala	<i>Aepyceros melampus</i>	45	0.17783
Reedbuck	<i>Redunca arundinum</i>	40	0.16279
Springbok	<i>Antidorcas marsupialis</i>	26	0.11785
Duiker	<i>Sylvicapra grimmia</i>	15	0.07801
Steenbok	<i>Raphicerus campestris</i>	10	0.05756

Table 2. The frequency distributions of 12' grids in the Kgalagadi and Ghanzi Agricultural Zones and the western strata of the Central Agricultural Zone in Botswana, and livestock attacks by cheetahs in these areas from 1995 to 2006 categorised by the percentage that cheetah's wild prey biomass contributed to the total biomass of wild prey and small stock combined.

<i>Percentage wild prey</i>	<i>Number of grids</i>	<i>Number of livestock attacks</i>	<i>Attacks per grid</i>
0	43	34	0.79
>0 to ≤ 1	48	39	0.81
>1 to ≤ 2	33	9	0.27
>2 to ≤ 5	48	27	0.56
>5 to ≤ 10	40	9	0.23
>10 to ≤ 20	23	15	0.65
>20 to ≤ 80	32	8	0.25
>80 to ≤ 100	136	47	0.35
TOTAL	403	188	

Table 3. Simultaneous confidence intervals for cheetah livestock attacks ($N = 188$) recorded (observed values) with the number of grid cells ($N = 403$) in the categories of the percentage that cheetah's wild prey biomass contributed to the total biomass of wild prey and small stock combined in the Kgalagadi and Ghanzi Agricultural Zones and the western strata of the Central Agricultural Zone in Botswana ($k = 4$, $\alpha = 0.05$, $Z = 2.4977$).

<i>Percentage</i>	<i>Expected</i>	<i>Observed</i>	<i>Bonferonni intervals</i>	<i>Use index</i>	<i>Significant</i>
<i>wild prey</i>	<i>Proportion</i>	<i>Proportion</i>	<i>for P_i</i>	<i>P_i / P_{io}</i>	
	<i>P_{io}</i>	<i>P_i</i>			
0	0.106700	0.180851	$0.1107 \leq P_i \leq 0.2510$	1.69	+
>0 to ≤ 20	0.476427	0.526596	$0.4357 \leq P_i \leq 0.6175$	1.11	0
>20 to ≤ 80	0.079404	0.042553	$0.0058 \leq P_i \leq 0.0793$	0.54	-
>80 to ≤ 100	0.337469	0.250000	$0.1711 \leq P_i \leq 0.3289$	0.74	-

Table 4. Simultaneous confidence intervals for the presence, transience and absence of cheetahs based on 89 questionnaire completed by farmers in areas deemed suitable and unsuitable for cheetahs in Botswana ($k = 4$, $\alpha = 0.05$, $Z = 2.4977$).

Observation Type	<i>Expected</i>	<i>Observed</i>	<i>Use</i>		
	<i>Proportion</i>	<i>Proportion</i>	Bonferonni	<i>index Pi/</i>	
	<i>Pio</i>	<i>Pi</i>	<i>intervals for Pi</i>	<i>Pio</i>	<i>Significant</i>
Absent in unsuitable area	0.026462	0.112360	$0.0287 \leq P_i \leq 0.1960$	4.25	*+
Present in unsuitable area	0.085687	0.000000	$0.0000 \leq P_i \leq 0.0000$	0.00	*-
Absent in suitable area	0.209493	0.123596	$0.0365 \leq P_i \leq 0.2107$	0.59	0
Present in suitable area	0.678358	0.764045	$0.6516 \leq P_i \leq 0.8765$	1.13	0