

Assessing Anxiety, Depression and Quality of Life in Patients with Peripheral Facial Palsy: A Systematic Review

Ferran Cuenca-Martínez^{1, 2}, Eva Zapardiel-Sánchez¹, Enrique Carrasco-González¹, Roy La Touche^{Corresp., 1, 2, 3}, Luis Suso-Martí^{2, 4}

¹ Departamento de Fisioterapia. Centro Superior de Estudios Universitarios La Salle, Universidad Autónoma de Madrid, Madrid, Spain

² Motion in Brains Research Group, Institute of Neuroscience and Sciences of the Movement (INCIMOV), Centro Superior de Estudios Universitarios La Salle, Universidad Autónoma de Madrid, Spain., Madrid, Spain

³ Instituto de Neurociencia y Dolor Craneofacial (INDCRAN), Madrid, Spain

⁴ Universidad CEU Cardenal Herrera, Valencia, Spain

Corresponding Author: Roy La Touche
Email address: roylatouche@yahoo.es

Objective: Peripheral facial palsy (PFP) is predominantly a unilateral disorder of the facial nerve , which can lead to psychological disorders that can result in decreased quality of life. The aim of this systematic review was to assess anxiety, depression and quality of life symptoms associated with PFP.

Data sources: We searched the Medline, PEDro, CINAHL and Google Scholar databases to conduct this systematic review while following Preferred Reporting Items for Systematic Reviews and Meta-Analyses standards. The search was performed by two independent reviewers, and differences between the two reviewers were resolved by consensus.

Study Selection: The search terms used were derived from the combination of the following MeSH terms: "facial paralysis", "bell palsy", "anxiety", "anxiety disorders", "depression", "depressive disorders", "quality of life" and not MeSH: "facial palsy", "hemifacial paralysis", "facial paresis", "Peripheral Facial Paralysis", using the combination of different Boolean operators (AND/OR).

Data Extraction: On November 1st (2019).

Data Synthesis: In total, 18 cross-sectional articles and two case-control studies were selected.

Conclusions: The cross-sectional articles showed low methodological quality, while the case-control studies showed acceptable methodological quality. Limited evidence suggests that patients with PFP might have increased levels of anxiety and depressive symptoms. A qualitative analysis also showed limited evidence that quality of life might be diminished in patients with PFP.

PROSPERO: CRD42020159843

Assessing Anxiety, Depression and Quality of Life in Patients with Peripheral Facial Palsy: A Systematic Review

Psychological factors in PFP

Ferran Cuenca-Martínez^{a,b}, Eva Zapardiel-Sánchez^a, Enrique Carrasco-González^a, Roy La
Touche^{a,b,c,*} and Luis Suso-Martí^{b,d}

*a. Departamento de Fisioterapia, Centro Superior de Estudios Universitarios La Salle,
Universidad Autónoma de Madrid, Spain.*

*b. Motion in Brains Research Group, Instituto de Neurociencias y Ciencias del Movimiento
(INCIMOV), Centro Superior de Estudios Universitarios La Salle, Universidad Autónoma
de Madrid, Spain*

c. Instituto de Neurociencia y Dolor Craneofacial (INDCRAN), Madrid, Spain

*d. Department of Physiotherapy, University CEU Cardenal Herrera, CEU Universities,
Valencia, Spain*

***Author for Correspondence: Roy La Touche**

Facultad de Ciencias de la Salud, Centro Superior de Estudios Universitarios La Salle, Calle La
Salle, nº 10, 28023 Madrid, Spain.

Teléfono: (+34) 91 740 19 80 Fax: (+34) 91 357 17 30

E-Mail: roylatouche@yahoo.es

Abstract

Objective: Peripheral facial palsy (PFP) is predominantly a unilateral disorder of the facial nerve, which can lead to psychological disorders that can result in decreased quality of life. The aim of this systematic review was to assess anxiety, depression and quality of life symptoms associated with PFP. **Data sources:** We searched the Medline, PEDro, CINAHL and Google Scholar databases to conduct this systematic review while following Preferred Reporting Items for Systematic Reviews and Meta-Analyses standards. The search was performed by two independent reviewers, and differences between the two reviewers were resolved by consensus.

Study Selection: The search terms used were derived from the combination of the following MeSH terms: "facial paralysis", "bell palsy", "anxiety", "anxiety disorders", "depression", "depressive disorders", "quality of life" and not MeSH: "facial palsy", "hemifacial paralysis", "facial paresis", "Peripheral Facial Paralysis", using the combination of different Boolean operators (AND/OR). **Data Extraction:** On November 1st (2019). **Data Synthesis:** In total, 18 cross-sectional articles and two case-control studies were selected. **Conclusions:** The cross-sectional articles showed low methodological quality, while the case-control studies showed acceptable methodological quality. Limited evidence suggests that patients with PFP might have increased levels of anxiety and depressive symptoms. A qualitative analysis also showed limited evidence that quality of life might be diminished in patients with PFP.

PROSPERO: CRD42020159843

Key words: Peripheral facial palsy; Psychological factors; Anxiety; Depressive symptoms; Quality of life.

Introduction

Peripheral facial palsy (PFP) is predominantly a unilateral disorder of the facial nerve (VII cranial nerve), a mixed nerve with both motor and sensory fibers, whose main function is to control the facial expression muscles. In addition, specifically the *chorda tympani* is a uniquely sensitive branch that innervates the anterior two-thirds of the tongue (Alonso Navarro et al., 2005; De Diego-Sastre, Prim-Espada & Fernández-García, 2005). However, despite being predominantly a unilateral clinical entity, it has been found that PFP can also be a bilateral disorder (Adour et al., 1978; Stahl & Ferit, 1989).

PFP has been shown to have several etiologies, which can be classified as a) idiopathic, commonly referred to as Bell's palsy; b) infectious, such as Lyme disease, otitis media or even related to herpes virus like the Ramsay Hunt syndrome, which occurs by reactivation of the varicella-zoster virus at the geniculate ganglion; c) genetic such as Melkersson-Rosenthal syndrome or Albers-Schönberg disease; d) tumorous, e) traumatic and f) post-surgical (Herrero Velázquez et al., 2009; Williams, 2010; Lorch & Teach, 2010).

At the motor level, PFP is related to a loss of essential facial functions, such as blinking, nasal breathing, lip sealing, smiling or speaking (Roob, Fazekas & Hartung, 1999; Lorch & Teach, 2010). Involvement of the musculature around the eyes is a key clinical finding for differentiating PFP from central facial palsy although at the beginning of the course of an incomplete PFP, the closing of the eyes may not be affected (Beurskens & Heymans, 2004). At the sensory level, PFP is characterized by a diminished sense of taste (Roob, Fazekas & Hartung, 1999; Lorch & Teach, 2010).

PFP can lead to psychosocial disorders such as depressive symptoms, high anxiety levels and reduced quality of life. Changes in facial symmetry can lead to depressed mood, which has

previously been associated with maladaptive behavior and depressive symptoms (Macgregor, 1990; Valente, 2004). The facial expressions of patients with PFP are often perceived negatively by observers, even when the patients are smiling (Ishii et al., 2012). Norris et al. (2019) showed that patients with PFP reported a decline in psychological wellbeing due to their perceived appearance and healthcare experience. Further research is therefore needed to relate the functional disorders of PFP to the possible psychosocial effects that aesthetic changes can have on this type of patient. Greater knowledge in this regard could increase the therapeutic efficacy of treatments and the quality of life of patients with PFP.

The main objective of this systematic review was to evaluate the involvement of psychological variables such as anxiety, depressive symptoms and quality of life in patients with PFP.

Material and Methods

The present systematic review study was conducted with the defined protocol and subdivided into four phases based on the standards of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement (**Fig. 1**) (Moher et al., 2009). The protocol of this systematic review was registered in an international register prior to starting the article (PROSPERO, CRD42020159843).

Inclusion criteria

The studies included in this systematic review had to meet the following criteria: a) cross-sectional, cohort or case-control study with a methodological design; b) patients with PFP; and c) evaluating only the following variables: anxiety, depression and quality of life.

Search strategy

The systematic search of the included articles was conducted on the following databases: MEDLINE (from 1950 to November 2019), CINAHL (from 1982 to November 2019), PEDro

(from 1950 to November 2019) and Google Scholar carried out in the latter on November 1st, 2019. The search terms used were derived from the combination of the following MeSH terms: "facial paralysis", "bell palsy", "anxiety", "anxiety disorders", "depression", "depressive disorders", "quality of life", and not MeSH: "facial palsy", "hemifacial paralysis", "facial paresis", "Peripheral Facial Paralysis", using the combination of different Boolean operators (AND/OR). We used the study design type (observational studies) as a filter. Two independent reviewers (RLT and FCM) performed the search using the same methodology. Differences between the two reviewers were resolved by consensus.

Selection criteria and data extraction

Two independent reviewers (EVS and ECG) conducted the first evaluation of the studies to assess the relevance (or lack thereof) of the systematic review. The reviewers used the articles' title, abstract and keywords for this initial screening process. If there was no initial consensus or the information provided by the abstract contained insufficient information, the full text was reviewed.

The full text was then reviewed to ensure that the studies met the inclusion criteria. A third reviewer (LSM) measured when there were differences between the evaluators until a consensus was reached. The data were extracted using a structured protocol that ensured that the most relevant information was obtained from each study.

Methodological quality assessment

The methodological quality of the articles included in this review was assessed using the Newcastle-Ottawa modified quality assessment scale (NOS) (Deeks et al., 2003). NOS is an appropriate tool for assessing the quality of case control, cohort and cross-sectional studies and has moderate interexaminer reliability (Hootman et al., 2011). NOS consists of 3 criteria with a

minimum of 0 stars and a maximum of 4: participant selection, exposure evaluation, results and comparability. In the total star count, each study is rated as one of the following categories: 1) poor, 0-3 stars; 2) acceptable, 4-5 stars; 3) good, 6-7 stars; and 4) excellent, 8-9 stars.(Wells et al., 2015) For the cross-sectional studies, we employed the modified version proposed by Fingleton et al. (2015) in which 1) 0-1 out of a possible 3 stars is considered poor quality, 2) 2 out of 3 is considered acceptable quality and 3) 3 out of 3 is considered good quality. Two independent evaluators (EVS and ECG) reviewed and examined the methodological quality of the studies included in the present study, and disagreements were resolved by consensus through the mediation of a third evaluator (FCM). Interevaluator reliability was determined using the kappa (κ) coefficient, where 1) $\kappa > 0.7$ indicated a high level of interevaluator agreement, 2) $\kappa = 0.5-0.7$ indicated moderate interevaluator agreement, and 3) $\kappa < 0.5$ indicated low interevaluator agreement (Cohen, 1960).

Qualitative analysis

For the qualitative analysis of the selected studies, we used an adaptation of the classification criteria proposed by van Tulder et al. for randomized clinical trials adapted for cross-sectional observational, cohort and case-control studies (van Tulder et al., 2003). The results were divided into five levels depending on the methodological quality: 1) strong evidence, representing results from multiple studies (at least three studies); 2) moderate evidence, representing results from multiple low-quality studies and/or one high-quality study; 3) limited evidence, one case-control and/or low-quality cohort study and/or at least two cross-sectional studies; and 4) conflicting evidence, inconsistent findings across multiple studies (at least three studies), and 5) No evidence, where there are no studies of any kind for an outcome category.

Results

We selected 18 cross-sectional studies (Weir et al., 1995; VanSwearingen et al., 1998; VanSwearingen, Cohn & Bajaj-Luthra, 1999; Coulson et al., 2004; Bradbury, Simons & Sanders, 2006; Lee et al., 2007; Fu, Bundy & Sadiq, 2011; Walker et al., 2012; Pouwels et al., 2014; Lindsay, Bhama & Hadlock, 2014; Sun et al., 2015; Kleiss et al., 2015; Togni et al., 2016; Volk et al., 2017; Nellis et al., 2017; Worrack et al., 2018; Díaz-Aristizabal et al., 2019; Muhammad Kassim, Abdullahi & Sammani Usman, 2019) and 2 case-control studies (Silva et al., 2011; Goines et al., 2016) in the first phase of the analysis. For all articles, we analyzed the variables depression, anxiety and quality of life. **Table 1** presents the study sample's characteristics, the symptom duration, the inclusion criteria, the measures of the variables and their assessment, and the conclusions of the analyzed psychological variables.

Characteristics of the study population

All studies were conducted with populations with PFP. A total of 2362 patients were evaluated, with a mean age of 45.7 years (range, 14–81 years). In 9 studies, the sample consisted of more women than men (Weir et al., 1995; VanSwearingen, Cohn & Bajaj-Luthra, 1999; Bradbury, Simons & Sanders, 2006; Walker et al., 2012; Pouwels et al., 2014; Lindsay, Bhama & Hadlock, 2014; Togni et al., 2016; Worrack et al., 2018; Díaz-Aristizabal et al., 2019), while two studies had more men than women (Kleiss et al., 2015; Muhammad Kassim, Abdullahi & Sammani Usman, 2019). Two studies (Pouwels et al., 2014; Díaz-Aristizabal et al., 2019) differentiated between patients with left or right PFP. Another study (Silva et al., 2011) observed a division between different phases of PFP, including a flaccid phase, a recovery phase and a sequel phase. Finally Coulson et al., 2004 and Muhammad Kassim, Abdullahi & Sammani Usman (2019) included participants with unilateral lower motor neuron facial nerve paralysis.

Results of the methodological quality assessment

Of the 20 articles included in this systematic review, 18 were cross-sectional observational studies, and 2 were case-control studies. The 18 cross-sectional studies showed low methodological quality (with scores of 1 out of 3 stars), while the 2 case-control studies showed acceptable methodological quality (with scores of 6 out of 9 stars). The third evaluator was needed to reach consensus on 2 of the articles. According to the kappa coefficient ($\kappa=0.736$), agreement among the assessors was high. The modified NOS results on the quality of the cross-sectional studies are presented in **Table 2**, while the results of the quality of the case-control studies are presented in **Table 3**.

Qualitative analysis

There was limited evidence that anxiety levels are higher in patients with PFP (Weir et al., 1995; Bradbury, Simons & Sanders, 2006; Fu, Bundy & Sadiq, 2011; Silva et al., 2011; Walker et al., 2012; Pouwels et al., 2014; Sun et al., 2015; Worrack et al., 2018; Díaz-Aristizabal et al., 2019; Muhammad Kassim, Abdullahi & Sammani Usman, 2019). This variable was assessed with the Hospital Anxiety and Depression Scale in 6 studies (Bradbury, Simons & Sanders, 2006; Fu, Bundy & Sadiq, 2011; Walker et al., 2012; Pouwels et al., 2014; Díaz-Aristizabal et al., 2019; Muhammad Kassim, Abdullahi & Sammani Usman, 2019), with the Self-reported Anxiety Scale in 1 study (Sun et al., 2015), with the General Health Questionnaire in 1 study (Weir et al., 1995), by open interview in 1 study (Silva et al., 2011) and with the Liebowitz Social Anxiety Scale in 1 study (Worrack et al., 2018).

There was limited evidence that depressive symptoms are present in patients with PFP (Weir et al., 1995; VanSwearingen et al., 1998; VanSwearingen, Cohn & Bajaj-Luthra, 1999; Bradbury, Simons & Sanders, 2006; Fu, Bundy & Sadiq, 2011; Silva et al., 2011; Walker et al., 2012;

Pouwels et al., 2014; Sun et al., 2015; Worrack et al., 2018; Díaz-Aristizabal et al., 2019; Muhammad Kassim, Abdullahi & Sammani Usman, 2019). One study found that impaired smiling and physical disability might be the main predictors of symptoms compatible with depression. Depressive symptoms was assessed with the Hospital Anxiety and Depression Scale in 6 studies (Bradbury, Simons & Sanders, 2006; Fu, Bundy & Sadiq, 2011; Walker et al., 2012; Pouwels et al., 2014; Díaz-Aristizabal et al., 2019; Muhammad Kassim, Abdullahi & Sammani Usman, 2019), with the Patient Health Questionnaire-9 in 1 study (Worrack et al., 2018), with the Self-reported Depression Scale in 1 study (Sun et al., 2015), with the Beck Depression Inventory in 2 studies (VanSwearingen et al., 1998; VanSwearingen, Cohn & Bajaj-Luthra, 1999) with the General Health Questionnaire in 1 study (Weir et al., 1995) and by open interview in 1 study (Silva et al., 2011).

There was limited evidence that quality of life might be reduced in patients with PFP (Lee et al., 2007; Lindsay, Bhama & Hadlock, 2014; Kleiss et al., 2015; Goines et al., 2016; Nellis et al., 2017). Quality of life was assessed with the Facial Clinical Evaluation Scale in 2 studies (Lee et al., 2007; Lindsay, Bhama & Hadlock, 2014) with a 100-point analog scale in 1 study (Goines et al., 2016), with a visual analog scale in 1 study (Nellis et al., 2017) and with the House-Brackmann scale in 1 study (Kleiss et al., 2015).

Discussion

The main objective of this review was to analyze the involvement of psychological variables in PFP. The results showed that anxiety and symptoms compatible with depression can be present in patients with PFP. Quality of life can also be diminished in these patients and is related to the presence of psychological factors.

Psychological factors can play a role in patients with PFP for several reasons, one of which is the possible involvement of facial palsy in an individual's self-concept. The functional and motor disorders of the facial region present in these patients significantly affect their perception of their self-image (Cross et al., 2000). This perception of physical attributes and self-image (or self-concept) significantly influences mood and self-esteem (Campbell, Assanand & Di Paula, 2003). The self-concept of patients with PFP is disrupted, in other words, there is a dissonance between their perceived image (the image altered by the PFP) and their learned self-concept, which has previously been associated with a decrease in self-esteem, the presence of anxiety and the appearance of depressive symptoms (Campbell, Assanand & Di Paula, 2003). Previous studies have shown that patients with facial palsy have an altered visualization of the affected side and that their self-concept is jeopardized, which could be related to the results of this study (Ishii et al., 2011).

Patients with PFP routinely report psychological distress due to the negative implications of PFP in social interactions. Previous studies have indicated that these patients' physical appearance reduces their self-esteem and attractiveness, which, due to social stereotypes, makes them appear less intelligent or confident, leading many people to avoid social relationships with patients with PFP (Ishii et al., 2012). Many patients with PFP are unable to express themselves or recognize facial expressions, which can also have a negative effect on their ability to communicate in their social setting, accentuating their difficulties interacting with others (Korb et al., 2016).

These patients might therefore view social interactions negatively, leading them to limit or avoid completely such interactions, resulting in social isolation, a factor highly related to poorer moods and depressive symptoms (Macgregor, 1990; Ishii et al., 2016). A number of studies have shown

an association between greater severity of facial palsy and greater depressive symptoms related to these social aspects (Nellis et al., 2017).

Some types of facial palsy such as Bell's palsy have been linked to immune system responses in nerve tissue. Previous research studies have shown increased proinflammatory cytokine levels in patients with this disease, which has been linked as a pathogenetic factor to the origin and maintenance of the disease (Yilmaz et al., 2002). Elevated anxiety and stress levels can also increase inflammatory levels, which in turn can exacerbate symptoms. For example, one study showed a two-way relationship between anxiety levels and Bell's palsy (Tseng et al., 2017).

Lastly, quality of life is related to the subjective perception of health, which includes the relationship between physical and mental health, as well as functionality. Quality of life therefore includes individual satisfaction with the state of health and the emotional response associated with it. Therefore, the aspects mentioned above can have a significant influence on the quality of life of patients with PFP (Taşkapılıoğlu & Karli, 2013). Our results show that the quality of life can be diminished in these patients and that their perception is closely linked to the psychological factors mentioned above. These results are similar to those obtained for other clinical entities such as temporomandibular disorders (Trize et al., 2018) and chronic migraine (Fernández-Concepción & Canuet-Delis, 2003).

From a clinical point of view, the results of the present review emphasize the need to thoroughly assess psychological aspects such as anxiety and depressive symptoms in patients with PFP. The correct management of these factors from a biopsychosocial paradigm can have a major impact on therapeutic success and an increase in the patients' quality of life.

Limitations

There are several limitations to be considered when interpreting the results of this systematic review. Firstly, the main limitation to be considered is the heterogeneity of the sample included in this article. It is very complicated to draw solid conclusions when you include patients with different states, moments and durations of PFP. For example, both Bradbury et al. (2006) and Coulson et al. (2004) include patients with PFP lasting at least one year. The work carried out by Diaz-Arastizabal et al. (2019) includes patients with PFP between 6 and 752 months. Fu et al. older than 6 months. Lee et al. (2007) deals with patients with PFP secondary to surgery, same as Sun et al. (2015), and with at least 24 months of duration, Lindsay et al. (2014) includes flaccid PFP or Silva et al. (2011) had a sample containing patients with PFP in the flaccid phase, another sample in the recovery phase and another in the sequelae phase. This limitation is very important to consider because the conclusions are established using the whole of this very heterogeneous sample. The results must therefore be considered with great caution. This fact also makes it impossible to carry out a statistical aggregation. Secondly, probably a greater number of psychological variables would have provided more information and could be considered a limitation. Thirdly, a number of the studies presented a high risk of bias due to their methodological quality or limited sample sizes, which could affect the results. Finally, the low number of articles could also be considered a limitation due to the impossibility of performing a meta-analysis or only because of the difficulty in drawing solid conclusions.

Conclusions

The results of this systematic review showed that patients with PFP can have increased anxiety levels and depressive symptoms, as well as having a decrease of quality of life. The quality of evidence of all outcome measures was limited. It is therefore that the presence of these psychological variables in patients with PFP should be considered clinically. However, the

272 authors stress the heterogeneity of the sample and therefore the findings should be taken with
273 caution.

274 **Author Disclosure Statement**

275 No competing financial interests exist.

276 **Funding Information**

277 No funding was received for this article.

278 **Acknowledgements**

279 The authors would like to thank Centro Superior de Estudios Universitarios CSEU La Salle for
280 its services in editing this manuscript.

281

References

- Adour KK, Byl FM, Hilsinger RL, Kahn ZM, Sheldon MI. 1978. The true nature of Bell’s palsy: analysis of 1,000 consecutive patients. *The Laryngoscope* 88:787–801. DOI: 10.1002/lary.1978.88.5.787.
- Alonso Navarro H, Zurdo Hernández JM, Ortí Pareja M, Jiménez Jiménez FJ. 2005. Parálisis facial periférica recurrente familiar. *Revista de Neurología* 40:61. DOI: 10.33588/rn.4001.2004565.
- Beurskens CHG, Heymans PG. 2004. Physiotherapy in patients with facial nerve paresis: Description of outcomes. *American Journal of Otolaryngology - Head and Neck Medicine and Surgery* 25:394–400. DOI: 10.1016/j.amjoto.2004.04.010.
- Bradbury ET, Simons W, Sanders R. 2006. Psychological and social factors in reconstructive surgery for hemi-facial palsy. *Journal of Plastic, Reconstructive and Aesthetic Surgery* 59:272–278. DOI: 10.1016/j.bjps.2005.09.003.
- Campbell JD, Assanand S, Di Paula A. 2003. The structure of the self-concept and its relation to psychological adjustment. *Journal of personality* 71:115–40. DOI: 10.1111/1467-6494.t01-1-00002.
- Cohen J. 1960. A Coefficient of Agreement for Nominal Scales. *Educational and Psychological Measurement* 20:37–46. DOI: 10.1177/001316446002000104.
- Coulson SE, O’Dwyer NJ, Adams RD, Croxson GR. 2004. Expression of emotion and quality of life after facial nerve paralysis. *Otology and Neurotology* 25:1014–1019. DOI: 10.1097/00129492-200411000-00026.
- Cross T, Sheard CE, Garrud P, Nikolopoulos TP, O’Donoghue GM. 2000. Impact of facial paralysis on patients with acoustic neuroma. *The Laryngoscope* 110:1539–42. DOI:

10.1097/00005537-200009000-00024.

Deeks JJ, Dinnes J, D'Amico R, Sowden AJ, Sakarovich C, Song F, Petticrew M, Altman DG.

2003. Evaluating non-randomised intervention studies. *Health Technology Assessment* 7.

DOI: 10.3310/hta7270.

Díaz-Aristizabal U, Valdés-Vilches M, Fernández-Ferreras TR, Calero-Muñoz E, Bienzobas-

Allué E, Moracén-Naranjo T. 2019. Correlations between impairment, psychological

distress, disability, and quality of life in peripheral facial palsy. *Neurologia* 34:423–428.

DOI: 10.1016/j.nrl.2017.03.004.

De Diego-Sastre JI, Prim-Espada MP, Fernández-García F. 2005. Epidemiología de la parálisis

facial del Bell. *Revista de neurologia* 41:287–90. DOI: 10.33588/rn.4105.2004593.

Fernández-Concepción O, Canuet-Delis L. 2003. [Disability and quality of life in patients with

migraine: determining factors]. *Revista de neurologia* 36:1105–12.

Fingleton C, Smart K, Moloney N, Fullen BM, Doody C. 2015. Pain sensitization in people with

knee osteoarthritis: a systematic review and meta-analysis. *Osteoarthritis and Cartilage*

23:1043–1056. DOI: 10.1016/j.joca.2015.02.163.

Fu L, Bundy C, Sadiq SA. 2011. Psychological distress in people with disfigurement from facial

palsy. *Eye* 25:1322–1326. DOI: 10.1038/eye.2011.158.

Goines JB, Ishii LE, Dey JK, Phillis M, Byrne PJ, Boahene KDO, Ishii M. 2016. Association of

facial paralysis-related disability with patient- and observer-perceived quality of life. *JAMA*

Facial Plastic Surgery 18:363–369. DOI: 10.1001/jamafacial.2016.0483.

Herrero Velázquez S, Guerrero Peral ÁL, Fuertes JJ, Rojo Martínez E, Peñas Martínez ML,

Cortijo García E, Fernández Herranz R. 2009. Neuralgia del nervio intermediario y parálisis

facial con o sin herpes. *Revista de Neurología* 49:670. DOI: 10.33588/rn.4912.2009233.

329 Hootman JM, Driban JB, Sitler MR, Harris KP, Cattano NM. 2011. Reliability and validity of
 330 three quality rating instruments for systematic reviews of observational studies. *Research*
 331 *Synthesis Methods* 2:110–118. DOI: 10.1002/jrsm.41.

332 Ishii L, Dey J, Boahene KDO, Byrne PJ, Ishii M. 2016. The social distraction of facial paralysis:
 333 Objective measurement of social attention using eye-tracking. *The Laryngoscope* 126:334–
 334 9. DOI: 10.1002/lary.25324.

335 Ishii LE, Godoy A, Encarnacion CO, Byrne PJ, Boahene KDO, Ishii M. 2011. What faces reveal:
 336 impaired affect display in facial paralysis. *The Laryngoscope* 121:1138–43. DOI:
 337 10.1002/lary.21764.

338 Ishii L, Godoy A, Encarnacion CO, Byrne PJ, Boahene KDO, Ishii M. 2012. Not just another
 339 face in the crowd: Society’s perceptions of facial paralysis. *Laryngoscope* 122:533–538.
 340 DOI: 10.1002/lary.22481.

341 Kleiss IJ, Hohman MH, Susarla SM, Marres HAM, Hadlock TA. 2015. Health-related quality of
 342 life in 794 patients with a peripheral facial palsy using the FaCE Scale: A retrospective
 343 cohort study. *Clinical Otolaryngology* 40:651–656. DOI: 10.1111/coa.12434.

344 Korb S, Wood A, Banks CA, AgoulNIK D, Hadlock TA, Niedenthal PM. 2016. Asymmetry of
 345 Facial Mimicry and Emotion Perception in Patients With Unilateral Facial Paralysis. *JAMA*
 346 *facial plastic surgery* 18:222–7. DOI: 10.1001/jamafacial.2015.2347.

347 Lee J, Fung K, Lownie SP, Parnes LS. 2007. Assessing impairment and disability of facial
 348 paralysis in patients with vestibular schwannoma. *Archives of Otolaryngology - Head and*
 349 *Neck Surgery* 133:56–60. DOI: 10.1001/archotol.133.1.56.

350 Lindsay RW, Bhama P, Hadlock TA. 2014. Quality-of-life improvement after free gracilis
 351 muscle transfer for smile restoration in patients with facial paralysis. *JAMA Facial Plastic*

Surgery 16:419–424. DOI: 10.1001/jamafacial.2014.679.

Lorch M, Teach SJ. 2010. Facial nerve palsy: Etiology and approach to diagnosis and treatment. *Pediatric Emergency Care* 26:763–769. DOI: 10.1097/PEC.0b013e3181f3bd4a.

Macgregor FC. 1990. Facial disfigurement: problems and management of social interaction and implications for mental health. *Aesthetic plastic surgery* 14:249–57. DOI: 10.1007/bf01578358.

Moher D, Liberati A, Tetzlaff J, Altman DG. 2009. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg* 8:6. DOI: 10.1016/j.ijssu.2010.02.007.

Muhammad Kassim N, Abdullahi A, Sammani Usman J. 2019. Relationship Between Psychological Symptoms, Age, Severity of Symptoms, and Societal Integration in Patients With Facial Nerve Paralysis. *Iranian Rehabilitation Journal*:235–240. DOI: 10.32598/irj.17.3.235.

Nellis JC, Ishii M, Byrne PJ, Boahene KDO, Dey JK, Ishii LE. 2017. Association among facial paralysis, depression, and quality of life in facial plastic surgery patients. *JAMA Facial Plastic Surgery* 19:190–196. DOI: 10.1001/jamafacial.2016.1462.

Norris JH, Longmire NM, Kilcoyne S, Johnson D, Fitzpatrick R, Klassen AF. 2019. Exploring Patient Experience of Facial Nerve Palsy to Inform the Development of a PROM. *Plastic and reconstructive surgery. Global open* 7:e2072. DOI: 10.1097/GOX.0000000000002072.

Pouwels S, Ingels K, Van Heerbeek N, Beurskens C. 2014. Cosmetic appreciation of lateralization of peripheral facial palsy: “Preference for left or right, true or mirror image?” *European Archives of Oto-Rhino-Laryngology* 271:2517–2521. DOI: 10.1007/s00405-013-2790-8.

375 Roob G, Fazekas F, Hartung H-P. 1999. Peripheral Facial Palsy: Etiology, Diagnosis and
376 Treatment. *European Neurology* 41:3–9. DOI: 10.1159/000007990.

377 Silva MFF, Cunha MC, Lazarini PR, Fouquet ML. 2011. Conteúdos psíquicos e efeitos sociais
378 associados à paralisia facial periférica: abordagem fonoaudiológica. *International Archives*
379 *of Otorhinolaryngology* 15:450–460. DOI: 10.1590/s1809-48722011000400008.

380 Stahl N, Ferit T. 1989. Recurrent bilateral peripheral facial palsy. *Journal of Laryngology and*
381 *Otology* 103:117–119. DOI: 10.1017/S0022215100108199.

382 Sun D, Shi Z, Li P, Shi S, Cai Y. 2015. Psychological status and quality of life in acoustic
383 neuroma patients with facial palsy after microsurgery: a 1-year postoperative follow-up
384 study. *Acta Neurologica Belgica* 115:311–316. DOI: 10.1007/s13760-014-0382-z.

385 Taşkapılıoğlu Ö, Karlı N. 2013. Assessment of Quality of Life in Migraine. *Noro psikiyatri*
386 *arsivi* 50:S60–S64. DOI: 10.4274/Npa.y7310.

387 Togni R, Abbamonte M, Comelli M, Mandrini S, Dall’angelo A, Pavese C, De Bernardi E,
388 Mariani F, Sala V, Caverzasi E, Giorgi I, Carlisi E, Dalla Toffola E. 2016. Personality
389 Traits and Perception of Disability after Facial Palsy. *J Psychol Psychother* 6:276. DOI:
390 10.4172/2161-0487.1000276.

391 Trize D de M, Calabria MP, Franzolin S de OB, Cunha CO, Marta SN. 2018. Is quality of life
392 affected by temporomandibular disorders? *Einstein (Sao Paulo, Brazil)* 16:eAO4339. DOI:
393 10.31744/einstein_journal/2018AO4339.

394 Tseng C-C, Hu L-Y, Liu M-E, Yang AC, Shen C-C, Tsai S-J. 2017. Bidirectional association
395 between Bell’s palsy and anxiety disorders: A nationwide population-based retrospective
396 cohort study. *Journal of Affective Disorders* 215:269–273. DOI:
397 10.1016/J.JAD.2017.03.051.

398 van Tulder M, Furlan A, Bombardier C, Bouter L. 2003. Updated Method Guidelines for
399 Systematic Reviews in the Cochrane Collaboration Back Review Group. *Spine* 28:1290–
400 1299. DOI: 10.1097/01.brs.0000065484.95996.af.

401 Valente SM. 2004. Visual disfigurement and depression. *Plastic surgical nursing : official*
402 *journal of the American Society of Plastic and Reconstructive Surgical Nurses* 24:140–6;
403 quiz 147–8. DOI: 10.1097/00006527-200410000-00003.

404 VanSwearingen JM, Cohn JF, Bajaj-Luthra A. 1999. Specific impairment of smiling increases
405 the severity of depressive symptoms in patients with facial neuromuscular disorders.
406 *Aesthetic Plastic Surgery* 23:416–423. DOI: 10.1007/s002669900312.

407 VanSwearingen JM, Cohn JF, Turnbull J, Mrzai T, Johnson P. 1998. Psychological distress:
408 Linking impairment with disability in facial neuromotor disorders. *Otolaryngology - Head*
409 *and Neck Surgery* 118:790–796. DOI: 10.1016/S0194-5998(98)70270-0.

410 Volk GF, Granitzka T, Kreysa H, Klingner CM, Guntinas-Lichius O. 2017. Initial severity of
411 motor and non-motor disabilities in patients with facial palsy: an assessment using patient-
412 reported outcome measures. *European Archives of Oto-Rhino-Laryngology* 274:45–52.
413 DOI: 10.1007/s00405-016-4018-1.

414 Walker DT, Hallam MJ, Ni Mhurchadha S, McCabe P, Nduka C. 2012. The psychosocial impact
415 of facial palsy: Our experience in one hundred and twenty six patients. *Clinical*
416 *Otolaryngology* 37:474–477. DOI: 10.1111/coa.12026.

417 Weir A, Pentland B, Crosswaite A, Murray J, Mountain R. 1995. Bell’s palsy: the effect on self-
418 image, mood state and social activity. *Clinical Rehabilitation* 9:121–125. DOI:
419 10.1177/026921559500900206.

420 Wells G, Tugwell P, O’Connell D, Welch V, Peterson J, Shea B, Losos M. 2015. The Newcastle-

421 Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in meta-analyses.
 422 Williams DS. 2010. Ramsay Hunt syndrome. *Journal of insurance medicine (New York, N.Y.)*
 423 42:41–43. DOI: 10.17245/jdapm.2018.18.6.333.
 424 Worrack S, Guntinas-Lichius O, Volk GF, Kaczmarek MC, Mühleck J, Brenk-Franz K, Strau
 425 BM, Altmann U. 2018. Restricted sleep as a secondary psychosocial consequence of facial
 426 palsy. *Laryngo- Rhino- Otologie* 97:398–404. DOI: 10.1055/a-0573-2119.
 427 Yilmaz M, Tarakcioglu M, Bayazit N, Bayazit YA, Namiduru M, Kanlikama M. 2002. Serum
 428 cytokine levels in Bell’s palsy. *Journal of the neurological sciences* 197:69–72. DOI:
 429 10.1016/s0022-510x(02)00049-7.
 430

Figure 1

Prisma flow chart.

Figura 1. PRISMA Flowchart




Table 1(on next page)

Characteristics of the studies included.

PFP: peripheral facial paralysis; M: man; W: woman; SD: standard deviation; FGS: Sunnybrook Facial Grading System; HADS: Hospital Anxiety and Depression Scale; FDI: Facial Disability Index; FaCE: Facial Clinimetric Evaluation; FPEM: Facial Paralysis Evaluation Profile; GBI: Glasgow Benefit Inventory; IPQ-R: Illness Perception Questionnaire-Revised; HBS: House-Brackmann Scale; BDI: Beck Depression Inventory; FAIR: Facial Nerve Palsy Integration Register; VAS = Visual Analogue Scale; TCI: Temperament and Character Inventory; SAS: Self-assessment Anxiety Scale, SDS: Self-assessed Depression Scale; ABS: Affect Balance Scale; SDS: Zung Self-Rating Depression Scale; FSI: Facial Self-Image Scale; FDQ: Funcional Disability Questionnaire; GHQ: General Health Questionnaire; MSRA: Measerement of Facial Motion; LSAS: Liebowitz Social Anxiety Scale; PHQ9: Patient Health Questionnaire.

1 **Table 1.** Characteristics of the studies included.

Article	Design	Characteristics of the sample of PFP group	Characteristics of the sample of control group	Average duration of the symptoms in PFP group	Inclusion criteria in PFP group	Outcome measures	Measurements instruments	General conclusions on psychological variables
Díaz-Aristizabal et al. (2019)	Cross-sectional study	N = 30 7 (M); 23 (W) Age: 51,1 (SD = 16,02) Affectation: 12 left; 18 right	-	-	Over 18 years old, minimum evaluation time of 6 month since the injury, incomplete resolution of the PFP and collaboration in the completion of the questionnaires.	Degree of PFP deficiency: FGS. Paralysis disability: FDI. Quality of life for patients with PFP.	Anxiety and depressive symptoms: HADS.	Patients with PFP with higher deficits had greater physical and overall disability and a worse quality of life. However, they did not show greater social disability, nor greater psychological impairment in the form of anxiety and depressive symptoms. Patients with greater physical, social and global disability had greater psychological impairment (anxiety and depressive symptoms) and a worse quality of life.
Bradbury et al. (2006)	Cross-sectional study	N = 106 Age: 44,7 34 (M); 72 (W) Form of PFP: 71 acquired paralysis; 35 congenital paralysis	-	At least 12 months after the operation	Patients undergoing vascularized muscle graft reconstruction to correct PFP. The operations had been performed by the same surgeon and the patients were taken from a database at the same hospital.	-	Anxiety and depressive symptoms: HADS. Satisfaction degree after PFP: FPEM Other psychosocial variables: semi-structured interview	There is no relationship between dissatisfaction with surgery and anxiety; however, this relationship does occur with depression. Most reported feeling social pressure and distress about their condition. Participants with consistent patterns of avoidance and social isolation, both before and after surgery, were more likely to be depressed than the rest of the study group and more likely to be dissatisfied with the surgery.
Coulson et al.	Cross-	N = 24	N = 24	1 year	-	Degree of PFP	Difficulties	A movement deficit associated

(2004)	sectional study	10 (M); 14 (H) Age: 46,1	17 couples, 5 family members, 2 friends	Mean age: 38,7		deficiency: Sunnybrook Facial Grading System, Sydney Facial Grading System, HBS. Quality of life: SF-36 PFP disability: FDI. Quality of life after surgery: GBI.	in facial expression of emotions: questionnaire	with the expression of specific emotions and a decrease in quality of life was identified in patients with long-term PFP.
Fu et al. (2011)	Cross-sectional study	N = 103 Age: 59 (SD = 17)	-	6 months - 50 years since PFP diagnosis	Patients with a diagnosis of PFP, adequate level of English, PFP longer than 6 months, over 18 years old	Degree of PFP deficiency: HBS Disease beliefs: IPQ-R.	Anxiety and depressive symptoms: HADS.	Participants have a significant level of psychological distress, which in turn is significantly related to perceptions about their FP. Significant differences were found between the levels of anxiety among women and men.
Muhammad Kassim et al. (2019)	Cross-sectional study	N = 37 16 (M); 21 (W) Age: 14-70	-	-	Patients with unilateral PFP over 18 years old with good cognitive ability (more than 24 points in Mini-Mental State Examination).	Degree of PFP deficiency: HBS Community integration of patients with PFP: FAIR	Anxiety and depressive symptoms: HADS.	People with PFP may suffer from emotional problems such as the presence of depressive symptoms, social isolation and self-awareness. Women and older people with PFP may be more vulnerable to social isolation. Women may be less depressed than men.
Nellis et al. (2017)	Cross-sectional study	N= 263 Age: 48,8	-	PFP developed post-surgery.	Patients without FP that are performed a facial plastic surgery. Older than 18 years old and speak English. Are excluded patients with a previous facial	Affectation level of PFPa PFP: HBS Quality of life: EVA (0 = bad, 100 = very good).	Psychometric data: Validated formulary: depression: BDI. Self-reported attractiveness	Depression significantly greater in patients that developed PFP after surgery. The also saw themselves less good looking, were in a worse mood and presented a decrease in quality of life. There were more women than men depressed.

					palsy surgery, if they have malignant neoplasm in head or neck, if had head or neck surgery and if couldn't fill the formulary' data on their own.		s: EVA (0 =I see myself very bad, 100 = I see myself very good). General mood: EVA (0 = very bad; 100 = very good).	
Kleiss et al. (2015)	Cross-sectional study	N = 794 40,1 (M); 59,9% (W) Age: 47 (SD = 16)	-	-	Patients with PFP. Children under 14 were excluded	Affectation level of PFP: HBS.	Quality of life in patients with PFP: FaCE.	Quality of life is more affected in patients with chronic facial paralysis in a psychosocial level. It is also related to increase of the age.
Volk et al. (2017)	Cross-sectional study	N = 256 103 (M); 153 (W) Age: 52 (SD= 18)	-	-	Patients with unilateral PFP. Patients with bilateral PFP were excluded as well as children under 14.	Affectation level of PFP: HBS. Disability induced by PFP (physical and social): FDI. Quality of life: SF-36.	Quality of life in patients with PFP: FaCE.	Patients with PFP suffer from a severe social and psychological disability. Patients with chronic PFP showed a greater disability than patients with acute PFP. Female an older patients showed more social and psychological disability
Togni et al. (2016)	Cross-sectional study	N = 61 41(W); 20(M) Age: 49,7 (SD = 13,92)	-	-	-	Affectation level of PFP: FGS. Facial disability level: FDI. Temper and character: TCI.	Depression: BDI.	Personality features modulate facial paralysis disability. Self-management shows up to mediate between the facial paralysis and the disability. Understanding how personality features influence in disability perception can be useful to improve the relationship between the doctor and the patient and to allow personalized and effective therapeutic interventions
Pouwels et al. (2014)	Cross-sectional	N = 59 Age=18-75	N=59 Randomly	-	PFP diagnostic, good level of german.	-	Anxiety and depressive	Significant differences have been found in anxiety and

	study	Affectation: 28 PF left 30 PF right	chosen, without previous PF symptoms, psychiatric or psychologi cal				symptoms: HADS.	depression between FP patients and healthy controls. Clinically significant differences have not been found in the presence of anxiety or depressive symptoms between right or left PFP.
Worrack et al. (2018)	Cross- sectional study	N=81	-	-	-	PFP symptoms and sleep evaluation: PSQI Quality of life: SF-36.	Depression: PHQ9. Social Anxiety: LSAS.	Facial changes related to PFP causes psychological problems that lead to reduction in sleep quality.
Sun et al. (2015)	Cross- sectional study	N= 21 11(M); 10(W) Age: 23-67	-	-	All the patients were diagnosed with acoustic neuroma and the surgery was made by the same doctor in all cases.	Level of Affectation of PF: HBS and Burres- Fisch. Quality of life: SF-36. Questionary of post-surgery perceptions.	Anxiety: SAS. Autoreporte d depression: SDS.	The PFP caused by the treatment of microsurgery can provoke psychological symptoms.
VanSwearing en et al. (1998)	Cross- sectional study	N = 48 Age: 49 (18-48) (SD = 16,3)	-	-	Older than 18. Had to be part of the facial center of Pittsburgh University.	Affectation level of PFP: FGS. Facial disability level: FDI. Subscale of physical and social wellness.	Depresión: BDI.	In this study of relations between clinical variables in facial neuromotor disorders, the facial deterioration was related with physical and social disability, with psychological disorders that stress the impact in physical and psychological disability and are connected with the social disability.
VanSwearing en et al. (1999)	Cross- sectional study	N= 29 Age: 50,2 (18- 81)	-	-	-	Facial disability level : FDI.	Depression: BDI.	The smile and physical disability were the main predictors of depression in

		(SD: 17,0)				Measure of facial movement: MSRA. Highest static response, a quantitative measurement of deterioration,a auto-notification rate, especific from the región of social and physical disability, related to facial deterioration.		patients of neuromuscular facial conditions.Global deterioration of facial movement was not a main predictor.
Lee et al. (2007)	Cross-sectional study	N = 56 30(M); 26(W) Age: 54,9 (SD: 1,7)	-	-	All the patients spoke english. Nobody with facial pathology previous to surgery.	Affectation PFP level: HBS.	Quality of life in patients with PFP: FaCE.	This impact magnitude in quailty of life can be not predicted by the severity of facial paralysis, the age or the gender of the patient, the time passed since the surgery or the size of the tumour.
Lindsay et al. (2014)	Cross-sectional study	N = 148	-	-	Patients with PFP.	Facial evaluation by Software: FACE-gram.	Quality of life in patients with PFP: FaCE.	The treatment of gracilis muscle produces and increase of the quality of life in patients with PF.
Walker et al. (2012)	Cross-sectional study	N= 126 (2 women for each man) Age: 50,1 (17-93)	-	-	Patients with PFP older than 16.	-	Anxiety and depression: HADS.	The severity of the paralysis is what affects to the psychological wellness of the patient and not the presents of the pathology.
Weir et al. (1995)	Cross-sectional	N = 20 12(W); 8(M)	-	6 days from PFP – 7 years	Patients with idiopathic PFP.	Perception of the self-image:	Anxiety and depression:	Increase in anxiety an depression levels.

study	Age: 41 (15-78)	(average of 65 days).			FSI. General health state: GHQ. Functional disability: FDQ.	part of GHQ.		
Silva et al. (2011)	Case-control study	N = 16 5 (M); 11 (W) Age: 43-48 PFP phase: 4 flaccid, 6 recovery y 6 sequel	-	-	Patients with idiopathic PFP older than 18. Patients with traumatic PFP are excluded.	PFP phase: otorhinolaryngologic and speech-language exam. Severity level of PFP: HBS.	Psychosocial variables: open interview.	Patients with PFP in an aftermath phase presented a greater affectation in psychosocial ways followed by those who were in a flaccid and recovery phase.
Goines et al. (2016)	Case-control study	N = 16	N = 4 N = 84 observers 59(W); 25(M)	-	Older than 18, without autism nor schizophrenia.	Facial clinic rating scale, quality of life, the paralysis severity and the disability were evaluated by a visual analogical 100 points scale.	Effective balance: ABS (happy, content, vigorous, affectionate, anxious, depressed, guilty and hostile).	The observers rated with a less quality of life in patients that presented facial paralysis in comparison to the self-reported disability by the patients.

- 2 PFP: peripheral facial paralysis; M: man; W: woman; SD: standard deviation; FGS: Sunnybrook Facial Grading System; HADS:
- 3 Hospital Anxiety and Depression Scale; FDI: Facial Disability Index; FaCE: Facial Clinimetric Evaluation; FPED: Facial Paralysis
- 4 Evaluation Profile; GBI: Glasgow Benefit Inventory; IPQ-R: Illness Perception Questionnaire-Revised; HBS: House-Brackmann Scale;
- 5 BDI: Beck Depression Inventory; FAIR: Facial Nerve Palsy Integration Register; VAS = Visual Analogue Scale; TCI:
- 6 Temperament and Character Inventory; SAS: Self-assessment Anxiety Scale, SDS: Self-assessed Depression Scale; ABS: Affect
- 7 Balance Scale; SDS: Zung Self-Rating Depression Scale; FSI: Facial Self-Image Scale; FDQ: Funcional Disability Questionnaire; GHQ:

8 General Health Questionnaire; MSRA: Measerement of Facial Motion; LSAS: Liebowitz Social Anxiety Scale; PHQ9: Patient Health
 9 Questionnaire.

10

11

12

13

14

15

Table 2(on next page)

Quality assessment of cross-sectional studies.

S = selection; C = comparability; O = outcome

1 **Table 2.** Quality assesment of cross-sectional studies.

Cross-sectional studies	S1: Representativeness of the exposed cohort	S2: Selection of the non- exposed cohort	S3: Ascertainment of exposure	S4: Demonstration that outcome of interest was not present at the start of the study	C1a: Study controls for previous injury	C1b: Study controld for age	O1: Assessment of outcome	O2: Follow- up long enought for outcomes to occur	O3: Adecuacy of follow up of cohorts	Total	%
Díaz-Aristizabal et al. (2019)		-	★	-	-	-		-	-	1/3	33
Bradbury et al. (2006)		-	★	-	-	-		-	-	1/3	33
Coulson et al. (2004)		-	★	-	-	-		-	-	1/3	33
Fu et al. (2011)		-	★	-	-	-		-	-	1/3	33
Muhammad Kassim et al. (2019)		-	★	-	-	-		-	-	1/3	33
Nellis et al. (2017)		-	★	-	-	-		-	-	1/3	33
Kleiss et al. (2015)		-	★	-	-	-		-	-	1/3	33
Volk et al. (2017)		-	★	-	-	-		-	-	1/3	33
Togni et al. (2016)		-	★	-	-	-		-	-	1/3	33
Pouwels et al. (2014)		-	★	-	-	-		-	-	1/3	33
Worrack et al. (2018)		-	★	-	-	-		-	-	1/3	33
Sun et al. (2015)		-	★	-	-	-		-	-	1/3	33
VanSwearingen et al. (1998)		-	★	-	-	-		-	-	1/3	33
		-	★	-	-	-		-	-	1/3	33

VanSwearingen et al. (1999)										
Lee et al. (2007)	-	★	-	-	-	-	-	1/3	33	
Lindsay et al. (2014)	-	★	-	-	-	-	-	1/3	33	
Walker et al. (2012)	-	★	-	-	-	-	-	1/3	33	
Weir et al. (1995)	-	★	-	-	-	-	-	1/3	33	

2

3

4 S = selection; C = comparability; O = outcome

5

Table 3(on next page)

Quality assessment of case studies and controls.

S = selection; C = comparability; E = Exposure

1 **Table 3.** Quality assesment of case studies and controls.

Case- Control Studies	S1: Definitio n of the cases	S2: Representativine ss of the cases	S3: Selectio n of controls	S4: Definitio n of controls	C1a: Study controls for previou s injury	C1b: Study control s for age	E1: Ascertainme t of exposure	E2: Same method of ascertainmen t for cases and controls	E3: Non- respons e rate	Tota l	%
Silva et al. (2011)	★		★	★	★	★	★			6/9	6 7
Goines et al. (2016)	★		★	★	★	★	★			6/9	6 7

2

3 S = selection; C = comparability; E = Exposure

