

The p53 signaling pathway of the large yellow croaker (*Larimichthys crocea*) responds to acute cold stress: Evidence via spatiotemporal expression analysis of *p53*, *p21*, *MDM2*, *IGF-1*, *Gadd45*, *Fas*, and *Akt*

Baoying Qian^{Corresp., 1}, Xin Qi¹, Yi Bai¹, Yubo Wu¹

¹ School of Life Science, Taizhou University, Taizhou, Zhejiang, China

Corresponding Author: Baoying Qian
Email address: wutongye1979@126.com

The *p53* activation is induced by stressors, such as DNA damage, oxidative stress, and activated oncogenes, and can promote cell cycle arrest, cellular senescence, and apoptosis. The large yellow croaker (*Larimichthys crocea*) is an important warm temperate marine fish in the Chinese aquaculture industry. However, few studies have investigated the role of *p53* in the response of *L. crocea* to environmental stressors. Therefore, the aim of the present study was to assess the spatiotemporal mRNA expression levels of genes involved in the *p53* signaling pathway of the large yellow croaker in response to cold stress. The results showed significant changes in the expression levels of *p53*, *p21*, *MDM2*, *IGF-1*, *Gadd45*, *Fas*, and *Akt* in various tissues of the large yellow croaker in response to cold stress for different times. As compared to the control group, *p53* mRNA expression was upregulated in most of the examined tissues at 24 h with the exception of the gill. In the liver, the expression levels of *p53* and *Fas* were significantly decreased at 12 h, while those of *p21*, *MDM2*, *IGF-1*, *Gadd45* were dramatically increased. *Akt* expression was notably changed in response to cold in several tissues. These results suggested that *p53* was potentially a key gene in the large yellow croaker response to cold and possibly other environmental stressors.

The p53 signaling pathway of the large yellow croaker (*Larimichthys crocea*) responds to acute cold stress: Evidence via spatiotemporal expression analysis of *p53*, *p21*, *MDM2*, *IGF-1*, *Gadd45*, *Fas*, and *Akt*

Baoying Qian¹, Xin Qi¹, Yi Bai¹, Yubo Wu¹

¹ School of Life Science, Taizhou University, Taizhou, Zhejiang 318000, P. R. China;

Corresponding author:

Baoying Qian

1139 Shifu Road, Taizhou, Zhejiang, 318000, P. R. China

Email address: wutongye1979@126.com

Abstract: The *p53* activation is induced by stressors, such as DNA damage, oxidative stress, and activated oncogenes, and can promote cell cycle arrest, cellular senescence, and apoptosis. The large yellow croaker (*Larimichthys crocea*) is an important warm temperate marine fish in the Chinese aquaculture industry. However, few studies have investigated the role of *p53* in the response of *L. crocea* to environmental stressors. Therefore, the aim of the present study was to assess the spatiotemporal mRNA expression levels of genes involved in the *p53* signaling pathway of the large yellow croaker in response to cold stress. The results showed significant changes in the expression levels of *p53*, *p21*, *MDM2*, *IGF-1*, *Gadd45*, *Fas*, and *Akt* in various tissues of the large yellow croaker in response to cold stress for different times. As compared to the control group, *p53* mRNA expression was upregulated in most of the examined tissues at 24 h with the exception of the gill. In the liver, the expression levels of *p53* and *Fas* were significantly decreased at 12 h, while those of *p21*, *MDM2*, *IGF-1*, *Gadd45* were dramatically increased. *Akt* expression was notably changed in response to cold in several tissues. These results suggested that *p53* was potentially a key gene in the large yellow croaker response to cold and possibly other environmental stressors.

Keywords: large yellow croaker; cold stress; *p53* signaling pathway; mRNA expression; gene network

1. Introduction

Net cage culturing of the large yellow croaker (*Larimichthys crocea*) is economically important to the marine aquaculture industry in China (Liu and Han, 2011). Over the past decade, low temperature resistance of *L. crocea* has gained considerable attention because of extensive economic losses caused by cold stress in winter, especially in the East China Sea (Xu et al, 2012). In order to breed large yellow croaker with stronger resistance to low temperatures and a lower death rate during the winter season, previous studies have investigated serum expression levels of physiological and biochemical markers, antioxidant production, enzymatic activities, and the proteome in response to cold stress, which showed that the serum biochemical indicators glutamate pyruvate transaminase, glutamic oxaloacetic transaminase, and alkaline phosphatase were significantly affected (Ji et al, 2009; Li et al, 2010; Zhang et al, 2013). More recent studies have focused on genes in response to cold stress. Liver transcriptome analysis indicated that the expression levels of numerous genes were either up- or down-regulated after 12 h of cold stress (Qian and Xue, 2016). Moreover, the expression profile of cold-inducible RNA-binding protein was significantly changed in different tissues of the large yellow croaker during acute cold stress (Miao et al, 2017). As the molecular mechanisms underlying the activation of these genes are complicated, further studies are needed to fully understand the genetic responses of the large yellow croaker to cold stress.

The tumor suppressor *p53* not only plays key roles in the inhibition of cell carcinogenesis and tumor development, but also promotes cell cycle arrest and apoptosis (Levine and Oren, 2009;

Kastenhuber and Lowe, 2017), and is involved in autophagy modulation, homeostatic regulation of metabolism, pluripotency, and repression of cellular plasticity (Aylon and Oren, 2016). On account of the negative regulation of *MDM2* (induced by *p53*), basal levels of *p53* are low. The *p53* gene acts as a “guardian of the genome” under physiological conditions (Momand et al, 1992; Haupt et al, 1997; Honda et al, 1997; Kubbutat et al, 1997) and is activated by stress signals, such as DNA damage, oncogene activation, and environmental stress. However, the response of *p53* is dependent on the intensity of the stress signal, the cell type, and the stage of cellular differentiation (Horn and Vousden, 2007; Kastenhuber and Lowe, 2017). Notably, the *p53* response is exceedingly flexible, as even a very low basal level of *p53* can protect the cell from the accumulation of DNA damage and subsequent carcinogenesis, which under different stress signals occurs through two typical mechanisms: (1) the promotion of cell senescence and apoptosis in response to severe or constant genotoxic and cellular stressors, and (2) the promotion of temporary cell cycle arrest in order to maintain cell survival prior to DNA repair (Jones et al, 2005; Maddocks et al, 2013; Chen et al, 2013; Pappas et al, 2017).

Various target genes of the *p53* signaling pathway involved in the arrest of cellular growth have been investigated, which include growth arrest and DNA damage-inducible protein (*Gadd45*), cyclin-dependent kinase inhibitor 1A (*p21*), and tumor necrosis factor receptor superfamily member 6 (*Fas*) (reviewed in Levine and Oren, 2009). The results of our previous study showed that the *p53* signaling pathway was significantly enriched in the liver of the large yellow croaker in response to cold stress for 12 h, while numerous genes related to cell cycle arrest,

apoptosis, and DNA repair and damage prevention were remarkably affected (Qian and Xue, 2016).
Reportedly, *p53* promotes apoptosis in the gills of the Nile tilapia (*Oreochromis niloticus*) and zebrafish (*Danio rerio*) in response to cold stress (Wang, 2016). Similarly, *p53* mRNA expression was significantly upregulated in the muscle tissue of *D. rerio* under low temperature stress (Li et al, 2018).

Therefore, the aim of the present study was to investigate the role of *p53* in response to cold-induced stress via analysis of the spatiotemporal expression of various genes (i.e., *p53*, *p21*, *MDM2*, *IGF-1*, *Gadd45*, *Fas*, and *Akt*) in different tissues of the large yellow croaker.

2. Materials and methods

2.1 Ethics statement

The study protocol was approved by the Medical Ethics Committee of Taizhou University Medical College (TZYXY2019-211) and conducted in accordance with the guidelines of the Institutional Animal Care and Use Committee. To minimize suffering, all experiment fish in this study were anesthetized by Tricaine-S (TMS, MS-222)(50 mg /L) firstly. And then we collected the tissues of large yellow croaker after the fish lost consciousness. Lastly, the fish which has been taken a surgery would be sprayed with anesthetic (200 mg/L), and let it death with euthanasia.

2.2 Fish and induction of acute cold stress

91 A total of 180 large yellow croakers (mean weight, 80 ± 0.7 g) were purchased from a
 92 mariculture farm located in Xiangshan Bay (Zhejiang Province, China), randomly assigned to
 93 one of six groups (30 fish/group), and then cultured in 500-L plastic aerated tanks in the
 94 laboratory of the Ningbo Marine and Fishery Science and Technology Innovation Base (Zhejiang
 95 Province, China) for 7 days. During the adaption period, the fish were fed granulated feed twice
 96 per day (04:30 and 18:30 h). A total of 90 fish cultured in three tanks were exposed to acute cold
 97 stress with the use of ice wrapped in thick plastic bags until the water temperature decreased to
 98 14°C within 2 h (cold stress group), while the other 90 fish in three tanks were cultured at
 99 environmental temperature and received no treatment (control group). After 1, 3, 6, 12, 24, 48,
 100 and 72 h of acute cold stress, the liver, muscle, gill, heart, spleen, intestine, brain, and kidney
 101 were collected from three fish in the cold stress group and control group, respectively, and
 102 immediately snap-frozen in liquid nitrogen, then stored at -80°C .

103 *2.3 Total RNA extraction and cDNA synthesis*

104 Total RNA was extracted from each sample using the E.Z.N.A.[®] Total RNA Kit I (Omega Bio-
 105 Tek, Inc., Norcross, GA, USA) in accordance with the manufacturer's instructions. Total RNA
 106 was quantified with a NanoDrop[™] 1000 Spectrophotometer (NanoDrop Technologies, LLC,
 107 Wilmington, DE, USA) and RNA integrity was assessed with the use of an Agilent 2100
 108 Bioanalyzer (Agilent Technologies, Inc., Santa Clara, CA, USA). The RNA integrity value of all
 109 samples was ≥ 8 . The extracted RNA was stored at -80°C until analysis.

First stand cDNA was synthesized from total RNA using the PrimeScript™ RT reagent Kit with gDNA Eraser (Takara Bio, Inc., Kusatsu, Shiga Prefecture, Japan) in accordance with the manufacturer's instructions and stored at -20°C until quantitative real-time polymerase chain reaction (qRT-PCR) analysis.

2.4 Spatiotemporal expression analysis

The qRT-PCR analyses of the spatiotemporal expression profiles of *p53*, *p21*, *MDM2*, *IGF-1*, *Gadd45*, *Fas*, and *Akt* were conducted with primers designed using Primer Premier 5 software (Premier Biosoft, Palo Alto, CA, USA) (Table 1). β -actin was used as a housekeeping gene. Before qRT-PCR, the amplification efficiency of the primers was evaluated with five 10-fold serial dilutions of cDNA of all tissues.

The qRT-PCR analysis was performed using a CFX96 Real-Time PCR System (Bio-Rad Laboratories, Hercules, CA, USA) with a total reaction volume of 20 μ L, consisting of 1 μ L of cDNA diluted to 1:5 with sterile DNase/ RNase-free distilled water, 0.6 μ L of the forward primer, 0.6 μ L of the reverse primer, 9 μ L of FastStart Universal SYBR Green Master mix (Sigma-Aldrich Corporation, St. Louis, MO, USA), and 8.8 μ L of sterile DNase/RNase-free distilled water. The following thermal cycling conditions were used: 95°C for 10 min followed by 40 cycles at 95°C for 15 s, 58°C for 20 s, and 72°C for 20 s. A melting curve was generated. Each sample was amplified in triplicate and the relative expression levels of *p53*, *p21*, *MDM2*, *IGF-1*, *Gadd45*, *Fas*, and *Akt* were normalized to that of β -actin with the $2^{-\Delta\Delta CT}$ method (Thomas and

Livak, 2008). Statistical significance was determined using one-way analysis of variance. All statistical analyses were performed using IBM SPSS Statistics for Windows, version 21.0. (IBM Corporation, Armonk, NY, USA). A probability (p) value of < 0.05 was considered statistically significant and < 0.01 as highly significant. All qPCR data could be obtained in “Supplemental File” which named with raw data.

3. Results

3.1 Analysis of the $p53$ signaling pathway

The results of our previous study showed that the expression profiles of genes involved in the $p53$ signaling pathway were significantly affected and this pathway was remarkably enriched in the liver transcriptome of the large yellow croaker in response to acute cold stress for 12 h (Qian and Xue, 2016). Gene networks associated with cell cycle arrest, apoptosis, $p53$ negative feedback, and DNA repair and damage prevention were evaluated in the present study based on the transcriptome data of previous studies (Fig. 1).

3.2 The mRNA profiles of *Akt*, *MDM2*, *p53*, *p21*, *Gadd45*, *Fas*, and *IGF-1* in liver tissue

The mRNA expression profiles of genes and gene networks in the liver of the large yellow croaker in response to acute cold stress were investigated (Fig. 2a₁₋₇, b). Liver mRNA expression levels of $p53$ were significantly decreased in response to cold stress at 3, 6 and 72 h, while

significantly increased at 24 and 48 h. In addition, the mRNA levels of *Akt*, *MDM2*, *p21*, and *Gadd45* were significantly increased at 1 and 3 h, while the expression levels of *p21* and *Gadd45* were significantly upregulated with the exception of 48 h. There was a good agreement between the qRT-PCR findings and previous liver transcriptome data of the large yellow croaker in response to acute cold stress at 12 h. The changes in mRNA expression levels of the selected genes were comparable between the two methods, although *IGF-1* expression was increased by qRT-PCR, while decreased according to the transcriptome data. The qRT-PCR results showed that the mRNA expression levels of *Akt*, *MDM2*, *p21*, and *Gadd45* were significantly increased, while that of *p53* was increased after 12 h of acute cold stress, as compared to the control group, although this difference was not statistically significant.

3.3 Spatiotemporal expression patterns of *Akt*, *MDM2*, *p53*, *p21*, *Gadd45*, *Fas*, and *IGF-1*

The spatiotemporal expression profiles of *Akt*, *MDM2*, *p53*, *p21*, *Gadd45*, *Fas*, and *IGF-1* in sampled tissues of large yellow croaker were determined by qRT-PCR analysis. The results of qRT-PCR analysis are shown in Figs. 2a₁₋₇ and 2b (liver), 3a₁₋₇ and 3b (muscle), 4a₁₋₇ and 4b (brain), 5a₁₋₇ and 5b (spleen), 6a₁₋₇ and 6b (gill), 7a₁₋₇ and 7b (kidney), 8a₁₋₇ and 8b (intestine), and 9a₁₋₇ and 9b (heart). *p53* mRNA expression levels in muscle were obviously increased after 1, 6, 12, 24, and 72 h of acute cold stress, but were significantly decreased at 3 h. mRNA levels of *Gadd45* and *p21* were increased in muscle throughout the cold stress period, although *Gadd45* expression was downregulated at 3 h. The mRNA expression levels of *Akt* and *Fas* were

downregulated, while those of *MDM2*, *p53*, *p21*, *Gadd45*, and *IGF-1* were upregulated in muscle tissue at 12 h (Fig. 3a₁₋₇, b).

Brain mRNA expression levels of all selected genes were significantly affected after 1 and 48 h of acute cold treatment. Notably, *p53* expression was significantly increased at 1, 12, 24, and 48 h, but not at 3, 6, and 72 h. Also, the expression levels of *MDM2* and *Gadd45* were dramatically upregulated in the brain tissue of the large yellow croaker throughout most of the treatment period (Fig. 4a₁₋₇, b).

In contrast to the brain tissue, *p53* mRNA expression in the spleen tissue was obviously increased at 24 and 48 h, but not at all times. Spleen mRNA expression levels of *Gadd45* were increased after cold stress, although there was no statistical significance at 72 h. In addition, the expression levels of *Akt*, *MDM2*, *p53*, *p21*, *Gadd45*, and *IGF-1* were significantly upregulated at 24 h, while that of *Fas* was significantly downregulated (Fig. 5a₁₋₇, b).

In the gill tissues, *p53* expression was significantly downregulated at 1, 3, 48, and 72 h, and significantly upregulated at 6 and 12 h. *MDM2* and *Akt* mRNA expression levels were obviously increased at 1, 24, 48, and 72 h, and at 3, 6, and 12 h of cold stress, respectively. *MDM2* mRNA expression was not significantly affected, while *Akt* expression was significantly decreased. Gill mRNA expression levels of *p21* were significantly increased throughout the cold stress period, as was that of *Gadd45* with the exception at 72 h (Fig. 6a₁₋₇, b).

Kidney mRNA expression levels of *p21* and *Gadd45* were significantly increased throughout the acute cold stress period, while that of *Fas* was obviously decreased with the exception of 1 and 24 h. The mRNA expression levels of *p53* in the kidney tissues were significantly upregulated at 1, 6, 12, 24, and 48 h. *MDM2* mRNA expression was upregulated throughout the cold stress period, although there was no significant change at 6 and 12 h (Fig. 7a₁₋₇, b).

The mRNA expression of *p53* in the intestinal tissue was significantly upregulated at 1, 24, and 48 h, but downregulated at 3, 6, 12, and 72 h. At 24 and 48 h after acute cold stress, the mRNA expression levels of *Akt*, *MDM2*, *p53*, *p21*, *Gadd45*, *Fas*, and *IGF-1* were all significantly increased. At 72 h, the mRNA expression levels of *Akt*, *MDM2*, *p21*, *Gadd45*, and *IGF-1* were significantly upregulated, while those of *p53* and *Fas* were downregulated (Fig. 8a₁₋₇, b).

The qRT-PCR results indicated that in the heart tissues, the mRNA expression levels of *Akt*, *MDM2*, *p53*, and *Gadd45* were significantly increased, while *p53* was significantly downregulated at 3 h with no significant difference at 1 and 12 h (Fig. 9a₁₋₇, b).

4. Discussion

Low temperatures that exceed the tolerance range of fish are known to disrupt energy metabolism, biochemical composition, immune function, and gene expression (Lu et al, 2019; Song et al, 2019). However, few studies have investigated the specific molecular mechanisms underlying the physiological changes in large yellow croaker in response to acute cold stress.

The results of our previous study of the liver transcriptome of *L. crocea* in response to 12 h of acute cold stress, identified a large number of differentially expressed genes that were enriched in the *p53* signaling pathway. Specifically, the mRNA expression levels of *MDM2*, *p21*, *Gadd45*, *CytC*, and *Apaf-1* were significantly increased, while those of *Fas*, *IGF*, and *CDK4/6* were significantly decreased. In addition, there was no significant change in the mRNA expression levels *p53* or other genes related to cell cycle arrest, apoptosis, inhibition of angiogenesis and metastasis, DNA repair and damage prevention, and *p53* negative feedback (Qian and Xue, 2016). In the present study, acute cold stress altered the expression profiles of genes related to cell cycle arrest and apoptosis in the liver, muscle, brain, spleen, gill, kidney, intestine, and heart. Of interest, the expression levels of *p21* and *Gadd45*, which are related to cell cycle arrest, were significantly changed in the liver, muscle, and kidney tissues throughout the cold stress period, while those of *Fas* and *IGF-1*, which are related to apoptosis, were also significantly altered in the heart tissue.

As a guardian of the genome, *p53* is remarkably sensitive to environmental factors and is readily activated by multiple stress signals, especially in aquatic organisms in response to temperature change. Li et al. (2018) demonstrated that upregulation of *p53* expression in response to low temperature stress can cause tail malformation of the zebrafish. The molecular mechanism of the *p53* pathway in response to cold stress also involves *MDM2* (Wang, 2016; Sun et al, 2019). In this study, the mRNA expression levels of *p53* and *MDM2* were significantly increased in the brain tissue of the large yellow croaker after 1 h of acute cold stress. Although there was no

change in the *p53* expression profile, *MDM2* expression was increased at 3 and 6 h. *p53* induces the expression of *MDM2* and *MDM2* inhibits the activity and promotes the degradation of *p53* in a negative feedback loop (Barak et al, 1993; Wu et al, 1993, Picksley and Lane, 1993). In the present study, the expression levels of *p53* and *MDM2* in response to the same cold stress period differed among tissues. In contrast to that observed in the brain, muscle, kidney, and intestine, there was no significant change in *p53* mRNA expression in the liver after 1 h of cold stress, while *MDM2* mRNA expression was significantly upregulated in the liver, heart, and spleen. These results may be due to the increased expression levels of *p53* in the liver (1.29-fold), heart (1.06-fold), and spleen (1.77-fold), which may have also impacted the expression of *MDM2* after 1 h of cold stress. In addition, the upregulation of *MDM2* inhibited *p53* expression in the liver at 3 and 6 h.

Akt plays key roles in glucose metabolism, apoptosis, cell proliferation, transcription, and cell migration. Activated *Akt* inhibits apoptosis through *MDM2* phosphorylation, which then inactivates *p53* (Song et al, 2005; Farrell et al, 2009). It has been reported that decreased mRNA expression of *Akt* and increased expression of *p53* can cause apoptosis of hepatocytes, suggesting an inverse correlation between these two genes (Wu et al, 2016). A delicate relationship between *Akt* and *p53* also occurred in the liver of the large yellow croaker after 1, 3, and 12 h of cold stress. During the cold stress period, the expression levels of *Akt* and *MDM2* were significantly increased, whereas *p53* expression was unchanged or decreased (Fig. 2b). In the present study, the relationships among *Akt*, *MDM2*, and *p53* seemed to be more intricate in different tissues in response to acute cold stress. *Akt* mRNA expression in the brain was significantly increased at 1

h, returned to normal levels from 3 to 6 h, decreased at 12 h, and increased again from 24 to 72 h, while *MDM2* expression was upregulated during most of the cold stress period, but not at 12 h (Fig. 4b). These results indicated that *MDM2* was regulated by *Akt* as well as *p53*.

Fas and *p21* are target genes of the *p53* pathway. The activation of *p21* usually predicts the beginning of cell cycle arrest, whereas *Fas* promotes apoptosis. Previous studies have reported that short pulses of *p53* activity usually lead to cell cycle arrest, as the *p21* promoter is more sensitive to this signaling output, while sustained *p53* signaling usually leads to changes in *Fas* expression, resulting in apoptosis. However, the mRNA signal of *p53*, either short or sustained, had no impact on the maximal level of the translated *p53* protein (Espinosa et al, 2003; Gomes et al, 2010; Morachis et al, 2010; Kastenhuber and Lowe, 2017). In this study, hepatic mRNA levels of *p53* increased by 1.29-fold in response to cold stress at 1 h. Although this change was not statistically significant, this slight increase in *p53* expression could be sufficient to result in a change in *p21* expression. In addition, mRNA expression of *p21* was remarkably increased from 1 to 24 h, even at 72 h, although *p53* expression was decreased, while *Fas* expression was decreased or remained comparatively unchanged. It is not clear why there were such differences in the expression patterns of *p21* and *Fas*, as it seems that targeting of *p53* by these genes had no impact on the expression profiles of *p53* at 3, 6, 12, 48, and 72 h in the liver tissues. Similar changes in expression levels occurred in the other tissues at certain times. One possibility could be complex regulatory signals from other genes that may be unknown in the *p53* signaling pathway in response to acute cold stress, which could affect the expression levels of *p21* and *Fas*.

Hence, further studies are warranted to fully understand the molecular mechanism of all of genes involved in the *p53* signaling pathway.

Gadd45 is as an important carcinogenic stress response factor that is sensitive to physiological and environmental stressors, and usually induced by cell cycle stagnation, DNA damage, and apoptosis (Liebermann and Hoffman, 2008; Salvador et al, 2013; Peng et al, 2015). It has been observed that *Gadd45* can interact with *p21*, which activates *p53* via *p38* to maintain *p53* signaling (Smish et al, 1994; Vairapandi et al, 1996; Azam et al, 2001; Liebermann and Hoffman, 2008). In the present study, *Gadd45* was sensitive to both long- and short-term cold stress, as indicated by the consistently high expression levels in the liver, intestine, kidney, and muscle at all time points. Even in the other tissues (spleen, heart, and gill), *Gadd45* mRNA expression was significantly increased at most time points. *Gadd45* is known to prevent DNA damage and promote DNA repair (Peng et al, 2015). It was possible that DNA damage was induced by acute cold stress, which resulted in significant increases in *Gadd45* expression. In addition, increased *Gadd45* expression may impact *p21* expression as mentioned previously.

The results of the present study indicated that genes involved in the *p53* signaling pathway were largely affected by acute cold stress. There were significant changes in the mRNA expression levels of *Akt*, *MDM2*, *p53*, *p21*, *Gadd45*, *Fas*, and *IGF-1* in the liver, brain, muscle, gill, kidney, intestine, heart, and spleen in response to acute cold stress. *p53* target *p21* and *Gadd45*, which are involved with cell cycle arrest and were more sensitive to cold stress than *Fas*. mRNA expression of *Gadd45*, which is involved in DNA repair, was significantly increased

283 in most of the studied tissues (liver, muscle, kidney, and intestine) in response to cold stress. The
 284 results of this study are in agreement with those of prior studies, which reported that genes
 285 involved in the *p53* signaling pathway could be affected by acute cold stress. However, further
 286 studies are needed to elucidate the molecular mechanisms of genes in the *p53* signaling pathway
 287 that are activated by low temperature stress.

288 **Acknowledgements**

289 This project was funded by Zhejiang Provincial Natural Science Foundation of China (Grant No.
 290 LGN18C190007) and Taizhou science and technology project (Grant No. 1901ny09). We thank
 291 International Science Editing for language editing this manuscript.

292 **Conflict of interest**

293 We declare that we have no conflicts of interest.

References:

- Liu, J.F., Han, K.H., 2011.** Current development situation and counter measure of large yellow croaker industry in China. *Journal of Fujian Fisheries*. 33(5): 4-8.
- Xu, Y.Q., Chen, Z., Ding, Z.K., 2012.** Effects of low temperature on fish and its prevention. *Journal of Southern Agriculture*. 39(15): 132-136.
- Ji, D.W., Li, M.Y., Wang, T.Z., Zhang, C.N., Xu, Z., Xu, W.S., 2009.** Effects of low temperature stress periods on serum biochemical indexed in large yellow croaker *Pseudosciaena crocea*. *Fisheries Science*. 28(1): 1-4.
- Li, M.Y., Ji D.W., Wu, H.Q., Chen, J., Shi, Y.H., Xu, Z., Xu, W.S., 2010.** 2-DE analysis in liver of *pseudosciaena crocea* during low temperature stress. *Journal of Agricultural Biotechnology*. 18(2): 323-328.
- Zhang, X.L., Hu, Y.Z., Li, M.Y., Miao, L., Chen, J., Shi, Y.H., Lu, X.J., 2013.** Changes in antioxidant level and serum enzyme activity of farmed large yellow croaker during natural water cooling in winter. *Marine Sciences*. 37(11): 27-34.
- Qian, B.Y., Xue, L.Y., 2016.** Liver transcriptome sequencing and *de novo* annotation of the large yellow croaker (*Larimichthys crocea*) under heat and cold stress. *Marine Genomics*. 25(1): 95-102.
- Miao, L., Li, M.Y., Chen, Y.Y., Hu, M., Chen, J., 2017.** Cloning of cold inducible RNA-binding protein (CIRP) gene in *Larimichthys crocea* and its expression analysis under cold treatments. *Journal of Fisheries of China*. 41(4): 481-489.
- Levine, A.J., Oren, M., 2009.** The first 30 years of p53: growing ever more complex. *Nature Reviews Cancer*. 9(10): 749-758.
- Kastenhuber, E.R., Lowe, S.W., 2017.** Putting p53 in Context. *Cell*. 170(6):1062-1078.
- Aylon, Y., Oren, M., 2016.** The paradox of p53: what, how, and why? *Cold Spring Harbor Perspectives in Medicine*. 6(10): a026328.
- Momand, J., Zambetti, G.P., Olson, D.C., George, D., Levine, A.J., 1992.** The mdm-2 oncogene product forms a complex with the p53 protein and inhibits p53-mediated transactivation. *Cell*. 69(7): 1237-1245.
- Haupt, Y., Maya, R., Kazaz, A., Oren, M., 1997.** Mdm2 promotes the rapid degradation of p53. *Nature*. 387(6630): 296-299.
- Honda, R., Tanaka, H., Yasuda, H., 1997.** Oncoprotein MDM2 is a ubiquitin ligase E3 for tumor suppressor p53. *FEBS Letters*. 420(1): 25-27.
- Kubbutat, M.H., Jones, S.N., Vousden, K.H., 1997.** Regulation of p53 stability by Mdm2. *Nature*. 387(6630): 299-303.
- Horn, H.F., Vousden, K.H., 2007.** Coping with stress: multiple ways to activate p53. *Oncogene*. 26(9): 1306-1316.
- Jones, R.G., Plas, D.R., Kubek, S., Buzzai, M., Mu, J., Xu, Y., Birnbaum, M.j., Thompson, C.B., 2005.** AMP-activated protein kinase induces a p53-dependent metabolic checkpoint. *Molecular Cell*. 18(3): 283-293.
- Moddocks, O.D., Berkers, C.R., Mason, S.M., Zheng, L., Blyth, K., Gottlieb, E., Vousden, K.H., 2013.** Serine starvation induces stress and p53-dependent metabolic remodeling in cancer cells. *Nature*. 493(7433): 542-546.
- Chen, K., Ding, Y.P., Wang, J.L., Shao, B.P., 2016.** Research progress on p53-involved metabolic regulation. *Biotechnology Bulletin*. 32(11): 52-58.
- Pappas, K., Xu, j., Zairis, S., Resnick-Silverman, L., Abate, F., Steinbach, N., Ozturk, S., Saal, L.H., Su, T., Cheung, P., Schmidt, H., Aaronson, S., Hibshoosh, H., Manfredi, J., Rabadan, R. Parsons, R., 2017.** p53 maintains baseline expression of multiple tumor suppressor genes. *Molecular Cancer Research*. 15(8): 1051-1062.
- Wang, J.F., 2016.** The molecular mechanisms underlying differential cold-induced apoptosis in gills of tilapia and zebrafish. Shanghai: Shanghai Ocean University.
- Li, W.H., Ran, H.Y., Chen, L.B., 2018.** Molecular mechanism of zebrafish tai developmental malformation caused by low temperature stress. *Genomics and Applied Biology*. 37(6): 2383-2388.
- Thomas, D.S., Livak, K.J., 2008.** Analyzing real-time PCR data by the comparative CT method. *Nature Protocols*. 3:1101-1108.
- Stoner, A.W., Ottmar, M.L., Copeman, L.A., 2010.** Temperature effects on the molting, growth, and lipid composition of

newly-settled red king crab. *Journal of Experimental Marine Biology and Ecology*. 393 (1-2): 138-147.

Lu, D.L., Ma, Q., Sun, S.X., Zhang, H., Chen, L.Q., Zhang, M.L., Du, Z.Y., 2019. Reduced oxidative stress increases acute cold stress tolerance in zebrafish. *Comparative Biochemistry and Physiology, Part A*. 235: 166-173.

Song, H.B., Xu, D.D., Tian, L., Chen, R.Y., Wang, L.G., Tan, P., You, Q.C., 2019. Overwinter mortality in yellow drum (*Nibea albiflora*): Insights from growth and immune responses to clod and starvation stress. *Fish and Shellfish Immunology*. 92: 341-347.

Barak, Y., Juven, T., Haffner, R., Oren, M., 1993. Mdm2 expression is induced by wild type p53 activity. *Embo Journal*. 12(2): 461-468.

Wu, X., Bayle, J.H., Olson, D., Levine, A.J., 1993. The p53-mdm-2 autoregulatory feedback loop. *Genes & Development*. 7(7a): 1126-1132.

Picksley, S.M., Lane, D.P., 1993. The p53-mdm2 autoregulatory feedback loop: a paradigm for the regulation of growth control by p53? *Bioessays*. 15(10): 689-690.

Sun, Z.Z., Tan, X.H., Xu, M.L., Liu, Q.Y., Ye, H.Q., Zou, C.Y., Ye, C.X., 2019. Liver transcriptome analysis and de novo annotation of the orange-spotted groupers (*Epinephelus coioides*) under cold stress. *Comparative Biochemistry and Physiology: Part D, Genomics & Proteomics*. 29: 264-273.

Song, G., Ouyang, G.L., Bao, S.D., 2005. The activation of Akt/PKB signaling pathway and cell survival. *Journal of Cellular and molecular medicine*. 9(1): 59-71.

Farrel, G.C., Larter, C.Z., Hou, J.Y., Zhang, R.H., Yeh, M.M., Williams, J., Dela peña, A., Francisco, R., Osvath, S.R., Brooling, J., Teoh, N., Sedger, L.M., 2009. Apoptosis in experimental NASH is associated with p53 activation and TRAIL receptor expression. *Journal of Gastroenterology and Hepatology*. 24(3): 443-452.

Wu, J.L., Chen, C., Hu, X., Cai, X.B., Guan, Y.H., Hu, H., Wang, Q.J., Chen, X.F., Cai, B.Z., Jing, X.B., 2016. Suppressing cyclooxygenase-2 prevents nonalcoholic and inhibits apoptosis of hepatocytes that are involved in the Akt/p53 signal pathway. *Biochemical and Biophysical Research Communications*. 469(4): 1034-1040.

Espinosa, J.M., Verdun, R.E., Emerson, B.M., 2003. p53 functions through stress- and promoter-specific recruitment of transcription initiation components before and after DNA damage. *Molecular Cell*. 12(4): 1015-1027.

Gomes, N.P., Espinosa, J.M., 2010. Disparate chromatin landscapes and kinetics of inactivation impact differential regulation of p53 target genes. *Cell Cycle (Georgetown, Tex.)*. 9(17): 3428-3437.

Morachis, J.M., Murawsky, C.M., Emerson, B.M., 2010. Regulation of the p53 transcriptional response by structurally diverse core promoters. *Genes & Development*. 24(2): 135-147.

Liebermann, D.A., Hoffman, B., 2008. Gadd45 in stress signaling. *Journal of Molecular signaling*. 3(1): 15.

Salvador, J.M., Brown-Clay, J.D., Fornace, A.J., 2013. Gadd45 in stress signaling, cell cycle control, and apoptosis. *Advances in Experimental Medicine & Biology*. 793(10):1-19.

Peng, Q.Q., Li, S.H., Zhang, W.L., Hu, H., Zhu, L.F., Yang, X.G., 2015. Influence of *Gadd45* gene expression regulation on tumor genesis. *Immunology Studies*. 3(2): 19-24.

Smith, M.L., Chen, I.T., Zhan, Q., Bae, I., Chen, C.Y., Gilmer, T.M., Kastan, M.B., O'Connor, P.M., Fornace, A.J., 1994. Protein interaction of the p53-regulated protein Gadd45 with proliferating cell nuclear antigen. *Science*. 266(5189): 1376-1380.

Vairapandi, M., Balliet, A.G., Hoffman, B., Liebermann, D.A., 1996. The differentiation primary response gene MyD118, related to GADD45, encodes for a nuclear protein which interacts with PCNA and p21WAF1/CIP1. *Oncogene*. 12(12): 2579-2594.

Azam, N., Vairapandi, M., Zhang, W., Hoffman, B., Liebermann, D.A., 2001. Interaction of CR6 (GADD45 gamma) with

376 proliferating cell nuclear antigen impedes negative growth control. Journal of Biological Chemistry. 276: 2766-2774.

377 Legends:

378 **Fig. 1.** Putative gene networks in large yellow croaker stressed by 12 h acute cold based on the transcriptome
379 data of previous studies. Enriched gene networks associated with cell cycle arrest, apoptosis, p53 negative
380 feedback, and DNA repair and damage prevention. Red indicates up-regulated, blue down-regulated, white
381 indicates no changes. Full names of abbreviated genes are listed in Abbreviation.

382
383 **Fig. 2.** qPCR analysis of genes in the liver of large yellow croaker under acute cold stress at different time
384 point (1, 3, 6, 12, 24, 72 h) (Fig. 2a₁₋₇); putative gene networks based on qPCR datas (Fig. 2b). Namely, tumor
385 protein p53 (*p53*), protein kinase B (*Akt*), insulin-like growth factor 1 (*IGF-I*), tumor necrosis factor receptor
386 superfamily member 6 (*Fas*), cyclin-dependent kinase inhibitor 1A (*p21*), growth arrest and DNA damage-
387 inducible protein (*Gadd45*), E3 ubiquitin-protein ligase mdm2 (*MDM2*). The results are expressed as mean fold
388 change \pm SD (n = 3 fish per treatment). Significant differences were considered at $*0.01 < P \leq 0.01$ and $**P <$
389 0.01.

390
391 **Fig. 3.** qPCR analysis of genes in the muscle of large yellow croaker under acute cold stress at different time
392 point (1, 3, 6, 12, 24, 72 h) (Fig. 3a₁₋₇); putative gene networks based on qPCR datas (Fig. 3b).

393
394 **Fig. 4.** qPCR analysis of genes in the brain of large yellow croaker under acute cold stress at different time
395 point (1, 3, 6, 12, 24, 72 h) (Fig. 4a₁₋₇); putative gene networks based on qPCR datas (Fig. 4b).

396
397 **Fig. 5.** qPCR analysis of genes in the spleen of large yellow croaker under acute cold stress at different time
398 point (1, 3, 6, 12, 24, 72 h) (Fig. 5a₁₋₇); putative gene networks based on qPCR datas (Fig. 5b).

399
400 **Fig. 6.** qPCR analysis of genes in the gill of large yellow croaker under acute cold stress at different time point
401 (1, 3, 6, 12, 24, 72 h) (Fig. 6a₁₋₇); putative gene networks based on qPCR datas (Fig. 6b).

402
403 **Fig. 7.** qPCR analysis of genes in the kidney of large yellow croaker under acute cold stress at different time
404 point (1, 3, 6, 12, 24, 72 h) (Fig. 7a₁₋₇); putative gene networks based on qPCR datas (Fig. 7b).

405
406 **Fig. 8.** qPCR analysis of genes in the intestine of large yellow croaker under acute cold stress at different time
407 point (1, 3, 6, 12, 24, 72 h) (Fig. 8a₁₋₇); putative gene networks based on qPCR datas (Fig. 8b).

408
409 **Fig. 9.** qPCR analysis of genes in the heart of large yellow croaker under acute cold stress at different time
410 point (1, 3, 6, 12, 24, 72 h) (Fig. 9a₁₋₇); putative gene networks based on qPCR datas (Fig. 9b).

412 Table 1

413 Primers for quantitative real time PCR.

414

415 **Abbreviations:**

416 p53: tumor protein p53

417 Akt: protein kinase B

418 IGF-1: insulin-like growth factor 1

419 CytC: cytochrome c

420 Apaf-1: apoptotic protease-activating factor

421 CASP8: caspase 8

422 CHK1: serine/threonine-protein kinase

423 Siah: E3 ubiquitin-protein ligase SIAH1

424 Bax: apoptosis regulator BAX

425 Fas: tumor necrosis factor receptor superfamily member 6

426 PIDD: leucine-rich repeats and death domain-containing protein

427 p21: cyclin-dependent kinase inhibitor 1A

428 CDK4/6: cyclin-dependent kinase 4/6

429 Gadd45: growth arrest and DNA damage-inducible protein

430 P53R2: ribonucleoside-diphosphate reductase subunit M2

431 Sestrins: sestrin 1/3

432 PAI: plasminogen activator inhibitor 1

433 BAI-1: adhesion G protein-coupled receptor B1

434 PTEN: phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase and dual-specificity protein phosphatase PTEN

435 TSC2: tuberous sclerosis 2

436 MDM2: E3 ubiquitin-protein ligase mdm2

437 Siah-1: E3 ubiquitin-protein ligase SIAH1

438 Cyclin G: cyclin G1

439 PIRH-2: RING finger and CHY zinc finger domain-containing protein 1

440 Cop-1: E3 ubiquitin-protein ligase RFWD2

441

Figure 1

Putative gene networks in large yellow croaker stressed by 12 h acute cold based on the transcriptome data of previous studies.

Putative gene networks in large yellow croaker stressed by 12 h acute cold based on the transcriptome data of previous studies. Enriched gene networks associated with cell cycle arrest, apoptosis, p53 negative feedback, and DNA repair and damage prevention. Red indicates up-regulated, blue down-regulated, white indicates no changes. Full names of abbreviated genes are listed in Abbreviation.




Figure 2

qPCR analysis of *Akt* in the liver of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 3

qPCR analysis of *p53* in the liver of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 4

qPCR analysis of *p21* in the liver of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 5

qPCR analysis of *MDM2* in the liver of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 6

qPCR analysis of *IGF-1* in the liver of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 7

qPCR analysis of *Gadd45* in the liver of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 8

qPCR analysis of *Fas* in the liver of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 9

putative gene networks based on qPCR datas in liver




Figure 10

qPCR analysis of *Akt* in the muscle of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 11

qPCR analysis of *p53* in the muscle of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 12

qPCR analysis of *MDM2* in the muscle of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 13

qPCR analysis of *p21* in the muscle of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 14

qPCR analysis of *IGF-1* in the muscle of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 15

qPCR analysis of *Gadd45* in the muscle of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 16

qPCR analysis of *Fas* in the muscle of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 17

putative gene networks based on qPCR datas in muscle




Figure 18

qPCR analysis of *Akt* in the brain of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 19

qPCR analysis of *p53* in the brain of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 20

qPCR analysis of *p21* in the brain of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 21

qPCR analysis of *MDM2* in the brain of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 22

qPCR analysis of *IGF-1* in the brain of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 23

qPCR analysis of *Gadd45* in the brain of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 24

qPCR analysis of *Fas* in the brain of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 25

putative gene networks based on qPCR datas in brain




Figure 26

qPCR analysis of *Akt* in the spleen of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 27

qPCR analysis of *p53* in the spleen of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 28

qPCR analysis of *p21* in the spleen of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 29

qPCR analysis of *MDM2* in the spleen of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 30

qPCR analysis of *IGF-1* in the spleen of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 31

qPCR analysis of *Gadd45* in the spleen of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)





Figure 32

qPCR analysis of *Fas* in the spleen of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)



putative gene networks based on qPCR datas in spleen




Figure 34

qPCR analysis of *Akt* in the gill of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 35

qPCR analysis of *p53* in the gill of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 36

qPCR analysis of *p21* in the gill of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 37

qPCR analysis of *MDM2* in the gill of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 38

qPCR analysis of *IGF-1* in the gill of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 39

qPCR analysis of *Gadd45* in the gill of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)





Figure 40

qPCR analysis of *Fas* in the gill of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)



putative gene networks based on qPCR datas in gill




Figure 42

qPCR analysis of *Akt* in the kidney of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 43

qPCR analysis of *p53* in the kidney of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 44

qPCR analysis of *p21* in the kidney of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 45

qPCR analysis of *MDM2* in the kidney of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 46

qPCR analysis of *IGF-1* in the kidney of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 47

qPCR analysis of *Gadd45* in the kidney of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 48

qPCR analysis of *Fas* in the kidney of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 49

putative gene networks based on qPCR datas in kidney




Figure 50

qPCR analysis of *Akt* in the intestine of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 51

qPCR analysis of *p53* in the intestine of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 52

qPCR analysis of *p21* in the intestine of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 53

qPCR analysis of *MDM2* in the intestine of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 54

qPCR analysis of *IGF-1* in the intestine of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 55

Gadd45




Figure 56

qPCR analysis of *Fas* in the intestine of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 57

putative gene networks based on qPCR datas in intestine




Figure 58

qPCR analysis of *Akt* in the heart of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 59

qPCR analysis of *p53* in the heart of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 60

qPCR analysis of *p21* in the heart of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 61

qPCR analysis of *MDM2* in the heart of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 62

qPCR analysis of *IGF-1* in the heart of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 63

qPCR analysis of *Gadd45* in the heart of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 48, 72 h)




Figure 64

qPCR analysis of *Fas* in the heart of large yellow croaker under acute cold stress at different time point (1, 3, 6, 12, 24, 72 h)




Figure 65

putative gene networks based on qPCR datas in heart




Table 1 (on next page)

Primers for quantitative real time PCR

Table 1 Primers for quantitative real time PCR.

Gene	Primer sequence (5'—3')	Gene	Primer sequence (5'—3')
<i>β-actin</i>	F: TCGGTATGGAATCTTGCG R: GTATTTACGCTCAGGTGGG	<i>Fas</i>	F: CACTCCAGCAGGGAAATGGA R: GCCATTTTGCTACGTCTCGC
<i>P53</i>	F: ACTACTGCCGGCCTAATGTG R: GCAAACCTGCATGGTTGGAGG	<i>Akt</i>	F: TGCCCCAGCATGAATGAAGT R: GTTGTGGTCACTGGACACCT
<i>MDM2</i>	F: TAGACGCCGTGCATGGATTT R: CCAGTTTGTGTGCATCGGCG	<i>Gadd45</i>	F: ATCAACGTGGTGCAGTCAA R: CATTGCAGTAGCGTGTGCAG
<i>P21</i>	F: GGGAAATGGCACCAATGTGCG R: GACGAAGAAGATGTCCGCCT	<i>IGF-1</i>	F: GTTCATTTTCGCCGGGCTTT R: ACAGCACATCGCACTCTTGA