

Vision of conspecifics decreases the effectiveness of ethanol on zebrafish behaviour

Rachel Dean ¹, Nicole Hurst-Radke ¹, Nirudika Velupillai ², Brian C Franczak ², Trevor J Hamilton ^{Corresp. 1, 3}

¹ Department of Psychology, MacEwan University, Edmonton, Alberta, Canada

² Department of Mathematics and Statistics, MacEwan University, Edmonton, Alberta, Canada

³ Neuroscience and Mental Health Institute, University of Alberta, Edmonton, Alberta, Canada

Corresponding Author: Trevor J Hamilton
Email address: trevorjameshamilton@gmail.com

Aquatic organisms in pharmacology and toxicology research are often exposed to compounds in isolation prior to physiological or behavioural testing. Recent evidence suggests that the presence of conspecifics during a stressful event can modulate behavioural outcomes (called 'social buffering') when testing occurs within the same context. It is unknown, however, whether the social environment during exposure interacts with the efficacy of anxiety-altering substances when subsequently tested in the absence of conspecifics. In this study, zebrafish were individually exposed to habitat water or ethanol (1.0% vol/vol) while untreated conspecifics were visually present or absent during dosing. Using the novel object approach test, a validated test of boldness and anxiety-like behaviour, we observed significantly greater effects of ethanol in isolated fish, compared to fish with a view of conspecifics during dosing. These results were not explained by altered locomotion during exposure, which might otherwise increase drug uptake. This highlights the need to consider the social environment during exposure when conducting and interpreting behavioural research involving drug or toxicant exposure.

Vision of conspecifics decreases the effectiveness of ethanol on zebrafish behaviour

Rachel Dean¹, Nicole Hurst-Radke,¹Nirudika Velupillai², Brian C. Franczak², & Trevor James
Hamilton.^{1,3}

¹Department of Psychology, MacEwan University, Edmonton, AB, Canada.

²Department of Mathematics and Statistics, MacEwan University, Edmonton, AB, Canada.

³Neuroscience and Mental Health Institute, University of Alberta, Edmonton, AB, Canada.

Corresponding Author: Trevor James Hamilton
10700 104 Ave NW, Edmonton, Alberta, T5J 4S2, Canada
Email address: trevorjameshamilton@gmail.com

ABSTRACT

Aquatic organisms in pharmacology and toxicology research are often exposed to compounds in isolation prior to physiological or behavioural testing. Recent evidence suggests that the presence of conspecifics during a stressful event can modulate behavioural outcomes (called ‘social buffering’) when testing occurs within the same context. It is unknown, however, whether the social environment during exposure interacts with the efficacy of anxiety-altering substances when subsequently tested in the absence of conspecifics. In this study, zebrafish were individually exposed to habitat water or ethanol (1.0% vol/vol) while untreated conspecifics were visually present or absent during dosing. Using the novel object approach test, a validated test of boldness and anxiety-like behaviour, we observed significantly greater effects of ethanol in isolated fish, compared to fish with a view of conspecifics during dosing. These results were not explained by altered locomotion during exposure which might otherwise increase drug uptake. This highlights the need to consider the social environment during exposure when conducting and interpreting behavioural research involving drug or toxicant exposure.

INTRODUCTION

Living in a social environment offers many evolutionary advantages. Belonging to a group can facilitate reproduction, enable earlier detection and evasion of predators, and improve resource efficiency (Rubinstein, 1978). Social cues are commonly the mechanism that conveys these messages between conspecifics and can guide responses in uncertain situations (Suboski et al., 1990). Another benefit of conspecifics is a decrease in stress level that minimizes the impact of stressful situations (Kikusui, Winslow & Mori, 2006). This phenomenon, known as ‘social buffering’, has been experimentally demonstrated in many species including cats (Masserman, 1943), goats (Liddell, 1949), rats (Davitz & Donald, 1955; Latané, 1969), humans (Hostinar, Johnson & Gunnar, 2015), and, recently, zebrafish (Oliveira & Faustino, 2017; Faustino, Tação-Monteiro & Oliveira, 2017).

The zebrafish has become a popular model organism for use in a variety of scientific disciplines including pharmacology and toxicology. Behavioural neuroscience tests can be used to analyze a wide variety of cognitive processes in zebrafish including episodic-like memory (Hamilton et al., 2016), object recognition memory (May et al., 2016), classically conditioned memory (Sison & Gerlai, 2010), fear (Speedie & Gerlai, 2008), boldness (Dean et al., 2020), and anxiety-like behaviour (Maximino, de Brito & da Silva Batista, 2010). To test anxiety-like behaviour there are a variety of paradigms available, with the most common being the light/dark preference and novel tank diving tests (for a review see (Maximino, de Brito & da Silva Batista, 2010). Due to the reliability of these tests and the practical simplicity in which psychopharmacological substances can be administered to zebrafish (Gerald, Lee & Blaser, 2006), adaptive behavioural responses can be easily manipulated with anxiolytic (anxiety-reducing) and anxiogenic (anxiety-enhancing) compounds (Collier & Echevarria, 2013).

Ethanol is a classic anxiolytic compound that has been shown to reduce zebrafish anxiety-like behaviour in a variety of paradigms. For instance, 0.25%, 0.5% and 1.0% ethanol increase the amount of time zebrafish spend exploring the light zone of the light/dark test (Gebauer et al., 2011; Fontana et al., 2020) and 0.3% and 0.5% ethanol increase time spent in the top zone of a novel tank dive test (Egan et al., 2009; Müller et al., 2020), both indications of the anxiolytic nature of ethanol. In the novel object approach test, used to test boldness and anxiety-like behaviour, 1.0% ethanol reduces the amount of time zebrafish spend in the outer ‘thigmotaxis zone’ adjacent to the wall of the arena (Johnson & Hamilton, 2017) further exemplifying the anxiolytic nature of ethanol. Using the same test, 1.5% ethanol also increases the time zebrafish spend in the inner zone exploring a novel object (Hamilton et al., 2017b), thus increasing boldness. Taken together, these findings illustrate zebrafish display less anxiety and more boldness following an acute administration of ethanol. Recent evidence, however, suggests that the social environment in which anxiety-altering compounds are administered and/or tested in may influence the behavioural effects of these substances in zebrafish (Faustino, Tacão-Monteiro & Oliveira, 2017), which may complicate conclusions.

Visual and olfactory conspecific cues can decrease the response of zebrafish to an anxiogenic compound when exposure and testing occurs within the same environment (Faustino, Tacão-Monteiro & Oliveira, 2017). Specifically, conspecific water and alarm substance along with a visual of untreated conspecifics induced significantly less freezing and erratic movements than when the adjacent tank was empty and no conspecific water was added (Faustino, Tacão-Monteiro & Oliveira, 2017). When the effectiveness of each type of cue was tested, visual cues were superior to olfactory cues in reducing aversive behaviours and promoting ‘social buffering’ (Faustino, Tacão-Monteiro & Oliveira, 2017). However, it is unknown whether the presence of

consppecifics can alter behavioural response when the fish are subsequently removed from the compound and tested in a separate arena. In other words, does the effect of social buffering persist beyond the exposure to the cues?

In the majority of acute pharmacological experiments that test individual fish behaviour, substances are administered while fish are physically isolated, and the exposed fish is then transferred to a behavioural arena for testing (Stewart et al., 2012). Moreover, almost all of the studies published that have examined acute ethanol exposure on behaviour of individual fish have not specified whether conspecifics were within view during dosing (Blaser and Peñalosa, 2011, Echevarria et al., 2008; Egan et al., 2009, Fontana et al., 2020, Gerlai et al., 2008). Moreover, only a few other studies have stated that fish were isolated during ethanol dosing (Hamilton et al., 2017b, Johnson and Hamilton 2010). To the best of our knowledge, no study has examined whether the view of conspecifics during dosing may influence the anxiety level of a fish subsequently tested in an isolated testing arena. It is also unknown whether social buffering may also act to alter the effects of anxiety *reducing*, ‘anxiolytic’ substances. To test these questions, we exposed individual zebrafish to either habitat water or ethanol (1.0% vol/vol) while untreated conspecifics were visually present or absent for the entire exposure period. Following exposure, the fish were transferred to the novel object approach test for quantification of anxiety-like behaviour and boldness (Dean et al., 2020; Krook et al., 2019; Leighton et al., 2018). Finally, we tested whether fish move at different rates and remain closer to conspecifics during the dosing period itself, in order to determine whether the social condition (*Isolated* vs. *In-view*) influences behaviour during drug exposure.

METHODS

Subjects and housing

Short-fin wild-type zebrafish (n = 90) were acquired from Aquatic Imports (Calgary, AB) at a minimum age of 9-months. Fish were experimentally naïve and comprised of mixed males and females (~50/50 ratio). Following a month-long quarantine period, the fish were held in either 3 or 10L polypropylene tanks within a three-shelf bench top system (Aquatic Habitats, Aquatic Ecosystems, Inc. Apopka, FL, USA) which was controlled for filtration and aeration. No fish was ever housed in isolation and tank capacities never exceeded five fish per liter. Temperature and pH remained between 26 - 30°C and 6.0 – 8.0, respectively. Lights were kept on a 12-hour light/dark cycle with lights on at 8AM and off at 8PM. Fish were fed dry brine shrimp (Omega One Freeze Dried Mysis Shrimp nutri-treat, OmegaSea Ltd., Germany) once per day, and after experimentation on test days. All experiments were approved by the MacEwan University Animal Research Ethics Board (AREB) under protocol number 05-12-13 in compliance with the Canadian Council for Animal Care (CCAC) guidelines for the care and use of experimental animals.

Experimental design

Experiment 1: This experiment used a 2 x 2 factorial design. The between-subject experimental variables included visual access to conspecifics (Fig. 1A,B, *Isolated or In-view*) and the type of substance the fish were exposed to (habitat water (*CTL*) or *Ethanol*) while in the dosing containers. Prior to experimentation, fish were randomly assigned to one of four groups: *Isolated-CTL*, *Isolated-Ethanol*, *In-view-CTL*, and *In-view-Ethanol*. Following exposure, behaviour was tested in the novel object approach test (Fig. 1C,D) to examine whether the social environment during exposure influences the efficacy of ethanol.

Experiment 2: A follow-up experiment was performed in which we tested the movement of individual zebrafish within the dosing containers in the *Isolated* or *In-view* conditions while exposed to control water (Fig. 2A,B).

Experiment 1: Experimental conditions

Ethanol exposure: On the day of experimentation, fish were carried in their habitat tanks into the experimental room prior to feeding and were given at least 10 minutes to acclimatize to this new environment. A white corrugated plastic barrier was set up surrounding habitat tanks to limit external stimuli. Following the habituation period, fish were individually netted from their habitat tanks and placed into one of two experimental dosing containers (600 mL). Each dosing container contained 500 mL of solution and was also surrounded by white corrugated plastic barriers (Fig. 1A). Two dosing containers were used rather than one to increase testing efficiency by allowing two fish to be dosed with a staggered schedule. Once in the dosing container, a square piece of the same plastic was placed on top to prevent evaporation of the solution and to ensure fish remained inside (Cachat et al., 2010; Holcombe et al., 2013). Fish exposed to control water (*Isolated-CTL* (n = 15) or *In-view-CTL* (n = 15)), were placed into dosing containers that only contained habitat water (500 mL). Fish in the ethanol groups (*Isolated-Ethanol* (n = 15) or *In-view-Ethanol* (n = 15)) were placed into dosing containers with 1.0% ethanol. Solutions for each compound were made fresh each day by mixing 5.26 mL of non-denatured, 95% ethanol into 495 mL of habitat water in the respective dosing containers. The selected concentration and duration of ethanol exposure was based on previous experiments in zebrafish (Johnson & Hamilton, 2017).

View of conspecifics: Fish were assigned to one of the two *Isolated* conditions (*Isolated-CTL* (n = 15), or *Isolated-Ethanol* (n = 15)) with no view of conspecifics while in the dosing container, or fish were assigned to the *In-view* conditions (*In-view-CTL* (n = 15) or *In-view-Ethanol* (n = 15)), with a view of a tank containing 12 untreated conspecifics during dosing. The dosing containers used in the *In-view* conditions were positioned in front of each other to ensure fish in both dosing containers had equal view of their conspecifics. The same group of conspecifics were used for each *In-view* condition. A white corrugated plastic barrier covered the remaining two sides of the conspecific tank (Fig. 1B) and water temperatures were maintained between 26 and 30°C by seedling heat mats (Hydrofarm Horticultural Products, Petaluma CA). Fish in the *Isolated* condition were surrounded fully by a white plastic barrier which was also placed on the heat mat. Fish in all conditions remained in the dosing containers for 30 minutes, after which the solution (including the fish) was carefully poured into a net, with a second dosing container collecting the solution. Once in the net, the fish was placed into the adjacent behavioural arena for testing. No data was acquired during dosing in experiment 1.

Experiment 1: Behavioural testing

Fish were individually tested in the novel object approach test following the 30-minute exposure period. The behavioural arena used in this experiment was circular and made from white opaque plastic ($\varnothing = 34$ cm; depth = 15 cm; Fig. 1C). The arena was placed on top of a heat mat to maintain habitat water temperatures and was surrounded by a three-sided white corrugated plastic enclosure to limit external stimuli during testing. Habitat water was added to the arena up to a maximum height of 5 cm and was replaced with fresh habitat water every four hours. An equal amount of heated habitat water was also exchanged whenever temperatures fell below

26°C. The object used in this study was a 2 cm x 4.25 cm Lego figurine which was multi-coloured to rule out possible colour preferences (Fig. 1D; Dean et al., 2020; Hamilton et al., 2017b; Johnson & Hamilton, 2017) and was adhered using velcro to the bottom of the arena's center. Prior to testing, three virtual zones representing the thigmotaxis (the outermost zone, 23-34 cm in diameter), inner (center zone of 12 cm in diameter) and transition zones (in between the other two zones, 12-23 cm in diameter) were defined using EthoVision XT motion tracking software (Fig. 1E; version 11.0, Noldus, VA, USA). All experimental procedures occurred between 9AM and 6PM prior to feeding. The time, in seconds, fish spent in each zone (thigmotaxis, transition, inner) was recorded and used as a proxy for anxiety-like behaviour (i.e. increased time in the thigmotaxis zone related to increased anxiety) and boldness behaviour (i.e. increased time in the inner zone is related to increased boldness) (Ou et al., 2015). Locomotion was also assessed by tracking the distance moved(cm) and immobility(s). Fish were tested individually for a period of 10 minutes following dosing and recording began as soon as the fish was placed into the transition zone facing the object.

Experiment 2: Experimental conditions and behavioural testing

After the first round of experimentation we sought to determine if the social context during exposure affected the distance fish moved while in the dosing container; possibly this could account for any differences in behavioural outcomes observed? In a second experiment, we tested the activity of a new group of fish in the dosing containers while in either the *Isolated-CTL* (n=15) or *In-view-CTL* (n=15) condition. We also quantified whether fish preferred the half of the dosing container close to conspecifics (conspecific side). Following a 10-minute habituation period, one fish was individually netted from their habitat tank and placed into a 600

mL dosing container with habitat water (500 mL), identical to procedures in our first experiment. A rectangular piece of white corrugated plastic was placed beneath the dosing container to assist with motion tracking. As in the novel object approach test, a three-sided enclosure was set up during behavioural tracking and seedling heat mats maintained water temperatures. To ensure these fish received the same treatment as fish in the *Isolated-CTL* and *In-view-CTL* conditions in the first experiment, a white piece of corrugated plastic was also set up across the front of the three-sided enclosure (Fig. 2A). For fish in both the *Isolated-Dosing* and *In-view-Dosing* conditions, EthoVision was set up to record the distance (cm) each fish moved throughout the 30-minute exposure period while in the dosing container. For fish in the *In-view-Dosing* condition, a habitat tank containing the same conspecifics ($n = 12$) that were used in the other *In-view* conditions was positioned to the right of the beaker. Using EthoVision, the beaker was then vertically split into two equal-sized virtual sections to compare the amount of time, in seconds, fish explored the side of the beaker closest to conspecifics (conspecific side) and the side farthest from conspecifics (empty side; Fig.2B). To rule out external variables potentially contributing to a side preference, the habitat tank was placed to the left of the beaker for the final three of fifteen trials per condition. No differences were observed in the time spent exploring either side of the beaker regardless of whether the habitat tank was on the right or left side of the beaker (Mann-Whitney; conspecific side, $U = 10$, $p = 0.2549$; opposite side, $U = 10$, $p = 0.2945$) so these were combined for analysis.

Statistical analysis

R (version 4.0.2) was used to analyze the data. The majority of the utilized functions are a part of the *stats* library. The exceptions are the *brunnermunzel.test* and *leveneTest* functions, which are a

part of the *brunnermunzel* and *car* libraries, respectively (Fox, 2016; Fox and Weisberg, 2019; Munzel and Brunner, 2000). Prior to the model fitting process, an exploratory data analysis was performed on all variables. In particular, we used the *cor.test* function to determine if there was evidence of statistically significant correlations among the measurements in the thigmotaxis, transitions, and inner zones and among the measurements in the distance moved and immobility variables. To analyze the effect of social condition and/or ethanol on anxiety levels, two-way ANOVAs were fitted to each variable using the *aov* function. The normality and constant variance assumptions were checked using Shapiro-Wilks' and Levene's tests, respectively, via the *shapiro.test* and *leveneTest* functions. If either assumption was violated, a Gamma Generalized Linear Model (GLM) with log link was fitted to the data using the *glm* function. If there was evidence that the Gamma GLM did not sufficiently fit the data, then Wilcoxon rank-sum (WRS) tests were fitted to compare both main effects and the treatment combinations of interest via the *wilcox.test* and *pairwise.wilcox.test* functions. In cases where the Gamma GLM fit sufficiently, the Brunner-Munzel test (Munzel and Brunner, 2000) was used to analyze differences between the treatment combinations of interest via the *brunnermunzel.test* function. Significance across all tests was determined using a 5% significance level. In what follows, we use $t(df)$ to designate a Student's t test statistic with df degrees of freedom, $F(df1,df2)$ to represent a F-test statistic with $df1$ and $df2$ degrees of freedom, W to represent the test statistic from a Wilcoxon Rank Sum test, and W_{BM} to represent the test statistic from the Brunner-Munzel test.

RESULTS

Analysis of Correlation

Time spent in the thigmotaxis zone was negatively correlated with time spent in the transition zone ($r = -0.816$, $t(58) = -10.74$, $p < 0.001$) and time spent in the transition zone ($r = -0.754$, $t(58) = -8.74$, $p < 0.001$). On the contrary, time spent in the inner zone was positively correlated, but the result was not statistically significant ($r = 0.235$, $t(58) = 1.84$, $p = 0.071$). Distanced moved was significantly negatively correlated with time spent immobile ($r = -0.643$, $t(58) = -6.400$, $p < 0.001$). While the results show that there are significant correlations between the times spent in each zone and between distance moved and immobility we follow Hamilton et al., (2017) and Thomson et al., (2020) and study individual behavioural measures rather than consider a principal component analysis as it has been shown that these behavioural measures have a large variance across reports from different research groups.

Effect of social context

Social context was found to have a significant effect on behaviour when comparing the time fish in the *In-view* and *Isolate* conditions spent in the thigmotaxis, transition, and inner zones. Specifically, fish spent more time in the thigmotaxis zone in the *In-view* condition than they did in the *Isolate* condition ($t(56) = 4.26$, $p < 0.0001$; Fig. 3A), and less time in the transition zone ($t(56) = 3.98$, $p < 0.001$; Fig. 3B) and the inner zone ($t(56) = 3.40$, $p = 0.001$; Fig. 3C), on average. No significant differences were found between the mean distance fish in the *In-view* and *Isolate* conditions moved ($F(1,56) = 2.44$, $p = 0.12$; Fig. 3D) nor between the median time fish in these groups spent immobile ($W = 470$, $p = 0.77$; Fig. 3E).

Investigating social context interactions

Among the control (CTL) groups, the Brunner-Munzel test provided sufficient evidence to conclude that fish in the *Isolate-CTL* group spent more time in the thigmotaxis zone than fish in the *In-View-CTL* group ($W_{BM} = -4.34, p < 0.001$, Fig. 3A). Whereas, fish in the *In-View-CTL* group spent more time in both the inner and transition zones than fish in the *Isolate-CTL* group ($W_{BM} = 3.99, p < 0.001$, Fig. 3C and $W_{BM} = 4.92, p < 0.0001$, Fig. 3B respectively).

Among the ethanol groups, the Brunner-Munzel test provided sufficient evidence to conclude that fish in the *In-view-Ethanol* group spent more time in the thigmotaxis zone than fish in the *Isolate-Ethanol* group ($W_{BM} = 2.07, p = 0.05$, Fig. 3A). Whereas, fish in the *Isolate-Ethanol* group spent more time in the inner zone compared to fish in the *In-View-Ethanol* group ($W_{BM} = -2.34, p = 0.03$, Fig. 3C). No differences were found when comparing the time fish in the *Isolate-Ethanol* and *In-View-Ethanol* groups spent in the transition zone ($W_{BM} = -0.93, p = 0.361$, Fig. 3B).

Effect of type of solution

The type of solution was found to have a significant effect on fish behaviour when comparing the time fish spent in their habitat water (CTL) to the time fish spent in ethanol. In particular, we found that fish dosed in the ethanol solution spent less time in the in the thigmotaxis zone ($t(56) = 8.78, p < 0.0001$; Fig. 3A), and more time in the transition zone ($t(56) = 7.20, p < 0.0001$; Fig. 3B) and the inner zone ($t(56) = 9.31, p < 0.0001$; Fig. 3C), on average, compared to fish dosed in control water. In addition, a significant difference was also detected between the median time spent immobile when comparing fish in their habitat water to those in ethanol ($W = 169.5, p < 0.0001$). No significant difference was found when comparing the mean distance moved for

these two groups ($F(1,56) = 3.75, p = 0.058$; Fig. 3D).

Investigating solution interactions

Investigating the treatment combinations revealed that time spent in the thigmotaxis zone was statistically greater for fish in the *In-view-CTL* group compared to fish in the *In-view-Ethanol* group ($W_{BM} = -3.24, p = 0.003$, Fig. 3A). Whereas, time spent in the transition zone was statistically greater for fish in the *In-view-Ethanol* group compared to fish in the *In-view-CTL* group ($W_{BM} = 3.52, p = 0.002$, Fig. 3B). A significant difference was not found when comparing the time fish in these two groups spent in the inner zone ($W_{BM} = 1.82, p = 0.079$, Fig. 3C). In addition, the Wilcoxon rank-sum test concluded that the median time spent immobile for fish in the *In-view-Ethanol* group was not significantly different compared to fish in *In-view-CTL* group ($W = 69, p = 0.074$, Fig. 3E).

Post hoc comparison of the isolated treatment groups using the Brunner-Munzel test indicated that fish in the *Isolate-CTL* group spent more time in the thigmotaxis zone than fish in the *Isolate-Ethanol* group ($W_{BM} = -32.19, p < 0.0001$, Fig. 3A). Whereas, fish in the *Isolate-Ethanol* group spent more time in both the transition zone ($W_{BM} = 24.80, p < 0.0001$, Fig. 3B) and the inner zone ($W_{BM} = 13.16, p < 0.0001$, Fig. 3C) compared to fish in *Isolate-CTL* group. The Wilcoxon rank-sum test indicated that the median time spent immobile by fish in *Isolate-CTL* group was significantly greater than median time spent in the in *Isolate-Ethanol* group ($W = 16.5, p < 0.001$, Fig. 3E).

Distance moved and side-preference during exposure

The distance fish moved while in the dosing container did not significantly differ between *Isolated-Dosing* and *In-view-Dosing* groups ($t_{28} = 1.255$, $p = 0.2198$; Fig. 4A). A highly significant preference for the conspecific side of the dosing container was found in fish from the *In-view-Dosing* group ($t_{28} = 10.21$, $p < 0.0001$; Fig. 4B).

DISCUSSION

To examine whether the sight of conspecifics during dosing impacts effectiveness of anxiety-like behaviour measurements and anxiety-altering substances, we exposed zebrafish to habitat water or ethanol (1.0%) while fish were isolated or able to observe conspecifics. Following dosing, behaviour was tested in the novel object approach test. Fish that were able to view conspecifics during dosing with control water had significantly less anxiety-like behaviour and increased boldness compared to fish dosed in isolation. The behavioural effects of ethanol also varied depending on the social condition in which it was administered. Ethanol had a significantly greater effect on anxiety-like behaviour and boldness in isolated fish compared to fish that were able to view conspecifics during dosing.

Both the sight of conspecifics and ethanol exposure significantly increased the time fish spent in the zones closest to the novel object (transition and inner; Fig. 3B-C), consistent with previous research demonstrating that ethanol increases boldness (Hamilton et al., 2017b; Johnson & Hamilton, 2017). Ethanol and sight of conspecifics also both decreased time spent in the thigmotaxis zone; an indication that anxiety-like behaviour was decreased. The most pronounced change in behaviour occurred with the combination of ethanol exposure and isolation during dosing compared to ethanol exposure and view of conspecifics; anxiety-like behaviour was significantly reduced (Fig. 3A, $P < 0.0001$) and boldness was increased (Fig. 3C, $P < 0.0001$). In

fish with a view of conspecifics, ethanol still did decrease anxiety-like behaviour, however to much less of an extent than in isolated fish, and had little effect on approach to the novel object. A similar pattern emerged in ethanol's influence on locomotion. Ethanol did not impact the distance *Isolated* or *In-view* fish moved (Fig. 3D), or the time *In-view* fish spent immobile; it only increased immobility in *Isolated* fish (Fig. 3E). This suggests that social isolation either increases sensitivity to ethanol's anxiolytic and depressant effects, or the presence of conspecifics suppresses these effects.

In an attempt to understand how the social context contributes to differences observed in behavioural outcomes, we analyzed the behaviour of a second group of fish while in the dosing container during the 30-minute dosing period. Because mobility may affect the rate of intake when fish are dosed via immersion, with greater physiological demands resulting in more ventilation and therefore increased uptake of the drug through the gills (Blaser & Vira, 2014), we sought to determine whether the heightened effect of ethanol observed in *Isolated* fish could be explained by greater movement during dosing. We analyzed the behaviours of fish exposed to habitat water while isolated or within view of conspecifics while they were in the dosing container. Interestingly, no differences were observed in the distance fish moved (Fig. 4A), indicating differences in locomotion during dosing could not explain the behaviours we observed. Not surprisingly, zebrafish spent significantly more time on the side of the dosing container closest to conspecifics when in view (Fig. 4B), demonstrating their preference to remain near other zebrafish.

An explanation for the increased anxiolytic effect of ethanol in *Isolated* relative to *In-view* groups may be related to 'social buffering.' Previous research in zebrafish has shown that the presence of conspecifics helps to suppress anxiety evoked by a fearful stimulus (Faustino,

378 Tacão-Monteiro & Oliveira, 2017). Faustino, Tacão-Monteiro and Oliveira (2017) first
 379 demonstrated this in zebrafish by exposing fish to a conspecific alarm substance with or without
 380 the presence of conspecific cues. They found that the anxiogenic effects were dampened by the
 381 presence of olfactory and/or visual cues. In other words, fish that could observe or smell their
 382 conspecifics showed less anxiety in response to the alarm substance (Faustino, Tacão-Monteiro
 383 & Oliveira, 2017). The mechanisms of social buffering have not been well explored in zebrafish;
 384 however, it is possible that zebrafish use the behaviours of their conspecifics as a source of
 385 information to guide their own responses in unfamiliar or fearful environments. This would
 386 explain why there was less of an effect of ethanol in the *In-view* condition in our experiment (ie.
 387 higher time in the thigmotaxis zone and less time near the object) compared to the *Isolated*
 388 condition. Presumably, fish in the dosing container were observing their conspecifics behaving
 389 normally and this may have minimized the effect of ethanol. However, social buffering has only
 390 been shown to decrease stress responses and in our study anxiolysis was reduced. The effect of
 391 ethanol may have been ‘buffered’ by the presence of conspecifics. The sight of conspecifics did
 392 significantly lower time spent in the thigmotaxis zone independent of the presence of ethanol,
 393 which suggests that the sight of conspecifics may lower anxiety overall. However, if this were
 394 the only factor at play then the effect of ethanol should be greater in the *In-view* condition
 395 compared to the *Isolated* condition, because of the combined anxiolytic effect of conspecifics
 396 and the anxiolytic effect of ethanol, but this is the opposite of what we observed. Somehow the
 397 visual cues from conspecifics decreased the effectiveness of ethanol, and the candidate
 398 mechanism is social mimicry, which has been demonstrated in zebrafish (Dresoti et al., 2015).
 399 To further investigate this future, studies could manipulate the emotional state of conspecifics
 400 with stress or pharmacology then examine how these *In-view* fish affect the behavioural

outcomes of the individual fish observing them. In this study, we did not examine sex differences and we did not manipulate the number of fish used as conspecifics nor the distance of the conspecific tank from the focal fish. These factors may also impact the strength of the social buffering response. It would also be valuable to explore the neurochemical basis of social buffering with analysis of brain chemistry after dosing in these social conditions.

CONCLUSIONS

Taken together, the sight of conspecifics decreases anxiety, and more so, buffers the anxiolytic effect of ethanol. These findings have important implications in the fields of pharmacology, toxicology and behavioural neuroscience as isolated drug administration seems to be more effective in eliciting a behavioural response. Additionally, zebrafish behavioural research findings can be inconsistent across laboratories, and this may be due a lack of detailed methodological reporting, including whether conspecifics are within visual range during drug exposure. Social buffering may offer a potential explanation for at least some of these discrepancies and necessitates detailed descriptions of dosing conditions in future experiments. Overall, this study provides the first evidence that the social condition during dosing effects the efficacy of anxiolytic substances when subsequently tested in isolation. This highlights the need to consider the social environment during exposure when conducting or interpreting behavioural research.

421 Figure legends:

422 **FIGURE 1.** Experimental dosing set-up. (A) *Isolated* and (B) *In-view* dosing. During dosing an
 423 individual fish was netted from the holding tank and placed into one of the two dosing
 424 containers. *In-view* fish had visual access to 12 conspecifics held in the conspecific tank but were
 425 not able to see the other fish being dosed. Focal fish remained in the dosing containers for 30-
 426 minutes prior to behavioural testing. C) The circular arena used was 34 cm in diameter and 16
 427 cm in height. D) The novel object used was a multi-coloured LEGO® figurine. E) The
 428 thigmotaxis, transition and inner zones were calibrated to 34, 23 and 12 cm in diameter
 429 respectively.

430

431 **FIGURE 2.** Experimental set-up for motion-tracking during exposure. (A) *Isolated* dosing, or
 432 (B) *In-view* dosing. The circle in the bottom left of figure (B) represents the virtual zones created
 433 in EthoVision to test whether fish spend more time on the side of the beaker closest to
 434 conspecifics when in view.

435

436 **FIGURE 3.** Effects of social context and ethanol on zone preference. The time, in seconds, fish
 437 spent in the thigmotaxis (A), transition (B) and inner zones (C). (D) and (E) represent the effect
 438 of social context and substance on the distance fish moved (D) and the time fish spent immobile
 439 (E). Graphs show individual data points (n = 15 per group) with horizontal lines that represent
 440 mean values \pm SEM. The number of * symbols identifies significant differences between groups
 441 at varying levels of significance: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$.

442

FIGURE 4. Distance and side preferences during the dosing procedure. (A) The distance, *Isolated-CTL* and *In-view-CTL* fish moved and the (B) amount of time, in seconds, *In-view-CTLs* spent on either side of the dosing container during dosing. The social condition did not have a significant effect on the distance fish moved, however when in-view, fish had a significant preference for the side of the dosing container closest to conspecifics. Data was analyzed using independent *t*-tests. Graphs show individual data points (*n* = 15 per group) with horizontal lines that represent mean values \pm SEM. The number of * symbols identifies significant differences between groups at varying levels of significance: *****P* < 0.0001.

Acknowledgements

We would like to thank Jasmin Bajwa, Shayna Chaput, Dr. Melike Schalomon, and Aleah McCory (Animal Care Technician) for their help with daily husbandry and aquarium maintenance.

References

- Blaser RE, Pénalosa YM. 2011. Stimuli affecting zebrafish (*Danio rerio*) behavior in the light/dark preference test. *Physiology & Behavior*, 104: 831-838. DOI: 10.1016/j.physbeh.2011.07.029
- Blaser RE, Vira DG. 2014. Experiments on learning in zebrafish (*Danio rerio*): a promising model of neurocognitive function. *Neuroscience & Biobehavioural Reviews*, 42: 224-231. DOI: <https://doi.org/10.1016/j.neubiorev.2014.03.003>.
- Cachat J, Canavello P, Elegante M, Bartels B, Hart P, Bergner C, Egan R, Duncan A, Tien D, Chung A, Wong K, Goodspeed J, Tan J, Grimes C, Elkhayat S, Suciu C, Rosenberg M, Chung KM, Kadri F, Roy S, Gaikwad S, Stewart A, Zapolsky I, Gilder T, Mohnot S, Beeson E, Amri H, Zukowska Z, Soignier RD, Kaleuff AV. 2010. Modeling withdrawal syndrome in zebrafish. *Behavior Brain Research*, 208: 371-376. DOI: <https://doi.org/10.1016/j.bbr.2009.12.004>.
- Collier AD, Echevarria DJ. 2013. The utility of the zebrafish model in conditioned place preference to assess the rewarding effects of drugs. *Behaviour Pharmacology*, 24: 375-383. DOI: <https://doi.org/10.1097/FBP.0b013e328363d14a>.
- Davitz JR, Donald JM. 1955. Socially facilitated reduction of a fear response in rats. *Journal of Comparative Psychology*, 48: 149-151. DOI: <https://doi.org/10.1037/h0046411>.
- Dean R, Duperrault E, Newton D, Krook J, Ingraham E, Gallup J, Franczak BC, Hamilton TJ. 2020. Opposing effects of acute and repeated nicotine exposure on boldness in zebrafish. *Scientific Reports*, 10: 8570. DOI: <https://doi.org/10.1038/s41598-020-65382-6>
- Dresoti E, Lopes G, Kampff AR, Wilson WW. 2015. Development of social behavior in young zebrafish. *Frontiers in Neural Circuits*, 9:39. <https://doi.org/10.3389/fncir.2015.00039>
- Echevarria DJ, Hammock CM, Pratt DW, Hosemann, JD. 2008. A novel behavioural test battery to assess global drug effects using the zebrafish. *International Journal of Comparative Psychology*, 21: 19-35.
- Egan RJ, Bergner CL, Hart PC, Cachat JM, Canavello PR, Elegante MF, Elkhayat SI, Bartels BK, Tien AK, Tien DH, Mohnot S, Beeson E, Glasgow E, Amri H, Zukowska Z, Kalueff AV. 2009. Understanding behavioral and physiological phenotypes of stress and anxiety in zebrafish. *Behavioural Brain Research*, 205: 38-44. Doi: 10.1016/j.bbr.2009.06.022
- Faustino AI, Tacão-Monteiro A, Oliveira RF. 2017. Mechanisms of social buffering of fear in zebrafish. *Scientific Reports*, 7: 44329. DOI: <https://doi.org/10.1038/srep44329>.
- Fontana BD, Duarte T, Muller TE, Canzian J, Ziana PR, Mezzomo NJ, Parker MO, Rosemberg DB. 2020. Concomitant taurine exposure counteracts ethanol-induced changes in locomotor and

- anxiety-like responses in zebrafish. *Psychopharmacology*, 237: 735-743. DOI:
<https://doi.org/10.1007/s00213-019-05410-0>
- Fox J. 2016. *Applied Regression Analysis and Generalized Linear Models*, Third Edition. Sage.
- Fox J. and Weisberg, S. 2019. *An R Companion to Applied Regression*, Third Edition, Sage
- Gebauer DL, Pagnussat N, Piato AL, Schaefer IC, Bonn CD, Lara DR. 2011. Effects of
anxiolytics in zebrafish: Similarities and differences between benzodiazepines, buspirone and
ethanol. *Pharmacology Biochemistry and Behavior*, 99: 480- 487. DOI:
10.1016/j.pbb.2011.04.021
- Gerald R, Lee V, Blaser R. 2006. Effects of acute and chronic ethanol exposure on the behavior
of adult zebrafish (*Danio rerio*). *Pharmacology Biochemistry and Behavior*, 85: 752-761. DOI:
<https://doi.org/10.1016/j.pbb.2006.11.010>.
- Gerlai R, Ahmad F, Prajapati S. 2008. Differences in acute alcohol-induced behavioral responses
among zebrafish populations. *Alcoholism, Clinical and Experimental Research*, 32: 1763-177s.
DOI: 10.1111/j.1530-0277.2008.00761.x
- Hamilton TJ, Myggland A, Duperreault E, May Z, Gallup J, Powell RA, Schalomon M, Digweed
SM. 2016. Episodic-like memory in zebrafish. *Animal Cognition*. 19(6):1071–1079. doi:
10.1007/s10071-016-1014-1.
- Hamilton TJ, Morrill A, Lucas K, Gallup J, Harris M, Healey M, Pitman T, Schalomon M,
Digweed S, Tresguerres M. 2017. Establishing zebrafish as a model to study the anxiolytic
effects of scopolamine. *Scientific Reports*, 7: 1-9. DOI: <https://doi.org/10.1038/s41598-017-15374-w>.
- Holcombe A, Howorko A, Powell RA, Schalomon M, Hamilton TJ. 2013. Reversed scototaxis
during withdrawal after daily-moderate, but not weekly-binge, administration of ethanol in
zebrafish. *PloS one*, 8. DOI: <https://doi.org/10.1371/journal.pone.0063319>.
- Hostinar CE, Johnson AE, Gunnar MR. 2015. Early social deprivation and the social buffering of
cortisol stress responses in late childhood: An experimental study. *Developmental Psychology*,
51: 1597-1608. DOI: <https://doi.org/10.1037/dev0000029>.
- Johnson A, Hamilton TJ. 2017. Modafinil decreases anxiety-like behaviour in zebrafish. *PeerJ*,
5. DOI: <https://doi.org/10.7717/peerj.2994>.
- Kikusui T, Winslow JT, Mori Y. 2006. Social buffering: relief from stress and
anxiety. *Philosophical Transactions: Biological Sciences*, 1476: 2215-2228. DOI:
<http://doi.org/10.1098/rstb.2006.1941>.

- Krook JT, Duperreault E, Newton D, Ross MS, Hamilton TJ. 2019. Repeated ethanol exposure increases anxiety-like behaviour in zebrafish during withdrawal. *PeerJ*, 7. DOI: <https://doi.org/10.7717/peerj.6551>.
- Latané B. Gregariousness and fear in laboratory rats. 1969. *Journal of Experimental Social Psychology*, 5: 61-69. DOI: [https://doi.org/10.1016/0022-1031\(69\)90006-7](https://doi.org/10.1016/0022-1031(69)90006-7).
- Leighton PLA, Nadolski NJ, Morrill A, Hamilton TJ, Allison WT. 2018. An ancient conserved role for prion protein in learning and memory. *Biology Open*, 7. DOI: <https://doi.org/10.1242/bio.025734>.
- Liddell H. 1949. Some specific factors that modify tolerance for environmental stress. *Research publications-Association for Research in Nervous and Mental Disease*, 29: 155-171.
- Masserman JH. 1943. Behavior and neurosis: An experimental psychoanalytic approach to psychobiologic principles. Chicago, IL: University of Chicago Press.
- Maximino C, de Brito TM, da Silva Batista AW, Herculano AM, Morato S, Gouveia Jr A. 2010. Measuring anxiety in zebrafish: a critical review. *Behavior Brain Research*, 214: 157-171. DOI: <https://doi.org/10.1016/j.bbr.2010.05.031>.
- May Z, Morrill A, Holcombe A, Johnston T, Gallup J, Fouad K, Schalomon M, Hamilton TJ. 2016. Object recognition memory in zebrafish. *Behavior Brain Research*, 296: 199-210. DOI: <https://doi.org/10.1016/j.bbr.2015.09.016>.
- Munzel U, Brunner E. 2000. Nonparametric methods in multivariate factorial planning. *Journal of Statistical Planning and Inference*, 88(1): 117-132. DOI: [https://doi.org/10.1016/S0378-3758\(99\)00212-8](https://doi.org/10.1016/S0378-3758(99)00212-8).
- Ou M, Hamilton TJ, Eom J, Lyall EM, Gallup J, Jiang A, Lee J, Close DA, Yun S-S, Brauner CJ. 2015. Responses of pink salmon to CO₂-induced aquatic acidification. *Nature Climate Change*. 5(10):950–955. doi: 10.1038/nclimate2694.
- Oliveira RF, Faustino AI. 2017. Social information use in threat perception: Social buffering, contagion and facilitation of alarm responses. *Communicative & Integrative Biology*, 10: 44329. DOI: <https://doi.org/10.1080/19420889.2017.1325049>.
- Rubinstein DI. 1978. On predation, competition, and the advantages of group living. *Social Behavior*, 205-231. DOI: https://doi.org/10.1007/978-1-4684-2901-5_9.
- Sison M, Gerlai R. 2010. Associative learning in zebrafish (*Danio rerio*) in the plus maze. *Behavior Brain Research*, 207: 99-104. DOI: <https://doi.org/10.1016/j.bbr.2009.09.043>.
- Speedie N, Gerlai R. 2008. Alarm substance induced behavioral responses in zebrafish (*Danio rerio*). *Behavior Brain Research*, 188: 168-177. DOI: <https://doi.org/10.1016/j.bbr.2007.10.031>.

594 Stewart A, Gaikwad S, Kyzar E, Green J, Roth A, Kalueff AV. 2012. Modeling anxiety using
595 adult zebrafish: a conceptual review. *Neuropharmacology*, 62: 135-143. DOI:
596 <https://doi.org/10.1016/j.neuropharm.2011.07.037>.
597
598 Suboski MD, Bain S, Carty AE, McQuoid LM, Seelen MI, Seifert M. 1990. Alarm reaction in
599 acquisition and social transmission of simulated predator recognition by zebra danio fish
600 (*Brachydanio rerio*). *Journal of Comparative Psychology*, 104: 101-112. DOI: 10.1037/0735-
601 7036.104.1.101
602
603 Thomson HR, Lamb SD, Besson AA, Johnson SL. 2020. Long-term repeatability of behaviours
604 in zebrafish (*Danio rerio*). *Ethology*, 126: 803-811. DOI: [10.1111/eth.13038](https://doi.org/10.1111/eth.13038)

Figure 1

Experimental dosing set-up

(A) *Isolated* and (B) *In-view* dosing. An Individual fish was netted from the holding tank and placed into one of the two dosing containers. *In-view* fish had visual access to 12 conspecifics held in the conspecific tank but were not able to see the other fish being dosed. Fish remained in the dosing containers for 30-minutes prior to behavioural testing. C) The circular arena used was 34 cm in diameter and 16 cm in height. D) The novel object used was a multi-coloured LEGO figurine. E) The thigmotaxis, transition and inner zones were calibrated to 34, 23 and 12 cm in diameter respectively.


A




B



C



D



E




Figure 2

Experimental set up for tracking behaviour during testing

(A) *Isolated* dosing, or (B) *In-view* dosing. The circle in the bottom left of figure (B) represents the virtual zones created in Ethovision to test whether fish spend more time on the side of the beaker closest to conspecifics when in view.

A



B




Figure 3

Effects of social context and ethanol on zone preference

The time, in seconds, fish spent in the thigmotaxis (A), transition (B) and inner zones (C). (D) and (E) represent the effect of social context and substance on the distance fish moved (D) and the time fish spent immobile (E). Individual data points represent mean values (n = 15 per group). Error bars represent the SE and * identifies significant differences between group means using 95% C.I. $**P < 0.01$; $***P < 0.001$; $****P < 0.0001$





Figure 4

Distance and side preferences during the dosing procedure

(A) The distance, *Isolated* and *In-view-CTL* fish moved and the (B) amount of time, in seconds, *In-view-CTLs* spent on either side of the dosing container during dosing. The social condition did not have a significant effect on the distance fish moved, however when in-view, fish had a significant preference for the side of the dosing container closest to conspecifics. Data was analyzed using independent *t*-tests. Individual data points (*n* = 15 per group) represent mean values \pm SEM. Error bars represent SE and * identifies significant differences between group means using 95% C.I. *****P* < 0.0001.

A



B

