

Refining the marine reptile turnover at the Early-Middle Jurassic transition (#53076)

1

First submission

Guidance from your Editor

Please submit by **19 Oct 2020** for the benefit of the authors (and your \$200 publishing discount) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

12 Figure file(s)

2 Table file(s)

4 Raw data file(s)



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Refining the marine reptile turnover at the Early-Middle Jurassic transition

Valentin Fischer^{Corresp., 1}, Robert Weis², Ben Thuy²

¹ UR Geology, Université de Liège, Liège, Belgium

² Département des sciences de la Terre, Musée national d'Histoire Naturelle, Luxembourg-Ville, Luxembourg

Corresponding Author: Valentin Fischer

Email address: v.fischer@uliege.be

Even though a handful of long-lived reptilian clades dominated Mesozoic marine ecosystems, several biotic turnovers drastically changed the taxonomic composition of these communities. A seemingly slow paced, within-System turnover took place across the Early-Middle Jurassic transition, seeing the demise of early neoichthyosaurians, rhomaleosaurid plesiosaurians and early plesiosauroids in favour of ophthalmosaurid ichthyosaurians and cryptoclidid and pliosaurid plesiosaurians, clades that will dominate the Late Jurassic and, for two of them, the entire Early Cretaceous as well. The fossil record of this turnover is however extremely poor, and this change of dominance appears to be spread across the entire middle Toarcian-Bathonian interval. We describe a series of ichthyosaurian and plesiosaurian specimens from successive geological formations in Luxembourg and Belgium that detail the evolution of marine reptile assemblages across the Early-Middle Jurassic, within a single sub-basin. These fossils reveal the continuing dominance of large rhomaleosaurid plesiosaurians, microcleidid plesiosaurians, and *Temnodontosaurus*-like ichthyosaurians up to the latest Toarcian, indicating that the structuration of the upper tier of marine ecosystems remained essentially constant up to the very end of the Early Jurassic. These fossils also suddenly record ophthalmosaurid ichthyosaurians and cryptoclidid plesiosaurians by the early Bajocian. As a result, the Early-Middle Jurassic marine reptile turnover appears restricted to the sole Aalenian stage, reducing the uncertainty of its duration, at least for ichthyosaurians and plesiosaurians, to 4 instead of 14 million years.

Refining the marine reptile turnover at the Early-Middle Jurassic transition

Valentin Fischer¹, Robert Weis², Ben Thuy²

¹ Evolution & Diversity Dynamics Lab, Université de Liège, Liège, Belgium.
v.fischer@uliege.be

² Muséum National d'Histoire Naturelle du Luxembourg, Luxembourg-Ville, Luxembourg.

Corresponding Author:

Valentin Fischer

14 Allée du 6 Août, Liège, 4000, Belgium

Email address: v.fischer@uliege.be

Abstract

Even though a handful of long-lived reptilian clades dominated Mesozoic marine ecosystems, several biotic turnovers drastically changed the taxonomic composition of these communities. A seemingly slow paced, within-System turnover took place across the Early-Middle Jurassic transition, seeing the demise of early neoichthyosaurians, rhomaleosaurid plesiosaurians and early plesiosauroids in favour of ophthalmosaurid ichthyosaurians and cryptoclidid and pliosaurid plesiosaurians, clades that will dominate the Late Jurassic and, for two of them, the entire Early Cretaceous as well. The fossil record of this turnover is however extremely poor, and this change of dominance appears to be spread across the entire middle Toarcian-Bathonian interval.

We describe a series of ichthyosaurian and plesiosaurian specimens from successive geological formations in Luxembourg and Belgium that detail the evolution of marine reptile assemblages across the Early-Middle Jurassic, within a single sub-basin. These fossils reveal the continuing dominance of large rhomaleosaurid plesiosaurians, microcleidid plesiosaurians, and *Temnodontosaurus*-like ichthyosaurians up to the latest Toarcian, indicating that the structuration of the upper tier of marine ecosystems remained essentially constant up to the very end of the Early Jurassic. These fossils also suddenly record ophthalmosaurid ichthyosaurians and cryptoclidid plesiosaurians by the early Bajocian. As a result, the Early-Middle Jurassic marine reptile turnover appears restricted to the sole Aalenian stage, reducing the uncertainty of its duration, at least for ichthyosaurians and plesiosaurians, to 4 instead of 14 million years.

Introduction

A series of diapsid clades dominated marine ecosystems during the entire Mesozoic (e.g. Motani, 2009; Benson, 2013; Pyenson, Kelley & Parham, 2014; Kelley & Pyenson, 2015). This long-term dominance was, however, pulsed by a series of extinctions and turnovers (Massare, 1987; Callaway & Massare, 1989; Bardet, 1994, 1995; Benson et al., 2010; Benson & Druckenmiller, 2014), the most-studied of which being those happening during the latest Triassic (Thorne, Ruta & Benton, 2011; Benson, Evans & Druckenmiller, 2012; Fischer et al., 2014c; Dick & Maxwell, 2015; Moon & Stubbs, 2020) and at the Jurassic-Cretaceous boundary (Tennant et al.; Fischer et al., 2012, 2013; Benson & Druckenmiller, 2014; Young et al., 2014a; Tennant, Mannion & Upchurch, 2016; Zverkov et al., 2018).

The dynamics of within-System marine reptile turnovers are less well known, despite evident modifications of assemblages outside System boundaries, such as during the early Late Cretaceous (Bakker, 1993; Bardet et al., 2008; Fischer et al., 2016, 2018). The composition of marine reptile communities clearly changed across the Early-Middle Jurassic transition (Maxwell, Fernández & Schoch, 2012; Vincent et al., 2013). Indeed, early neoichthyosaurian ichthyosaurs and microcleidid plesiosaurs seemingly met their demise and rhomaleosaurid plesiosaurs went near extinction (Maxwell, Fernández & Schoch, 2012; Fischer et al., 2013; Benson, Zverkov & Arkhangelsky, 2015), while ophthalmosaurid ichthyosaurs, thalattosuchian crocodylomorphs, cryptocleidid plesiosauroids, and pliosaurs radiated (Motani, 1999a; Ketchum & Benson, 2011a; Fischer et al., 2013; Benson & Druckenmiller, 2014; Moon, 2017; Foffa, Young & Brusatte, 2018). These newly diversified clades will then dominate the Late Jurassic and beyond.

However, the fossil record of marine reptiles across the middle Toarcian - Allovian interval is extremely poor (Benson et al., 2010; Maxwell, Fernández & Schoch, 2012) (Figure 1) and no region preserves fossiliferous successions spanning this transition. These biases blur both the tempo and the severity of this turnover. We tackle this issue by describing and analysing the composition of the late Toarcian-Bajocian marine reptile assemblages of Luxembourg and Belgium. We show that typical Early Jurassic forms such as rhomaleosaurid and microcleidid plesiosaurs and non-baracromian ichthyosaurs persisted in mid latitudes up to the latest Toarcian and that ophthalmosaurid ichthyosaurs and cryptocleidid plesiosaurs, the dominant Late Jurassic clades, suddenly appear during the early Bajocian, providing a refined timeline in a single sub-basin for what appears as an abrupt, Aalenian-restricted turnover.

Material & Methods

Upper Toarcian of Luxembourg: Thouarsense, Pseudoradosa, and Aalensis zones

Towards the end of the Early Jurassic, the area of present-day Luxembourg was located in the northeast Paris Basin within the northwestern Peri-Tethys Ocean (e.g. Pieńkowski et al., 2008; Schintgen & Förster, 2013). In a shallow, near-coastal sea between the former landmasses of the Rhenish Massif in the north and the Vosges and Black Forest in the south, a thick siliclastic succession was deposited during the late Toarcian, with clayey to silty sediments in the lower parts and silty to sandy, iron-grain rich sediments in the upper part following a general coarsening upwards trend (Siehl & Thein, 1989). The material described herein includes a single, semi-articulated post-cranial ichthyosaur skeleton from the so-called “Couches à *Astarte voltzi*” (Dittrich, 1993), a clay-rich siltstone corresponding to level 11 of Delsate and Weis (2010), dated to the late Toarcian Thouarsense Ammonite Zone of the Grandcourt Formation.

All the other late Toarcian marine reptile remains from Luxembourg described herein were found in the overlying Minette ironstone Formation (Lucius, 1945). Deposition of the Minette successions took place under the influence of strong tidal currents (Teyssen, 1984) during the upper Toarcian Pseudoradosa and Aalensis ammonite Zones (Di Cencio & Weis, 2020). The Aalenian Opalinum ammonite zone has not yet been detected in the area and probably forms a local hiatus (Maubeuge, 1972; Guérin-Franiatte & Weis, 2010).

Upper Toarcian of Belgium: Dispansum-Pseudoradosa-Aalensis zones

The Mont-Saint-Martin Formation crops out in southern Belgium and includes iron-rich beds forming a slightly diachronous equivalent of the Minette ironstones in Luxembourg. These ironstones contain all the marine reptiles and were deposited in a shallow marine setting during the late Toarcian (Dispansum to Aalensis ammonite zones; Boulvain et al., 2001). One fossil (IRSNB Vert-06455-0001) was found just outside the Belgian border, in the Meurthe-et-Moselle department, north-eastern France (Figure 2).

Middle-upper Aalenian of Luxembourg: Murchisonae and Concavum zones

The Toarcian part of the Minette ironstones in Luxembourg is locally overlain by a few meters of iron-rich marly sandstones and thin beds with calcareous nodules deposited during the Aalenian Murchisonae to Concavum ammonite zones (Guérin-Franiatte & Weis, 2010; Sadki, Weis & Braun, 2020). These deposits form the onset of a transgressive succession with increasingly finer sedimentation leading to the so-called “marnes micacées”, a succession of poorly lithified silty claystones with thin layers of phosphatic nodules, dated to the lowermost Bajocian Discites ammonite zone (Guérin-Franiatte & Weis, 2010).

Lower Bajocian of Luxembourg: Humphriesianum Zone

The youngest marine strata in Luxembourg correspond to a succession of calcarenites, marl-limestone alternations and coral limestones deposited on the northeast margin of the Burgundy carbonate platform at the end of the lower Bajocian. The marine reptile remains described herein were all recovered from the so-called “Marnes sableuses d’Audun-le-Tiche”, dated to the lower Bajocian Humphriesianum ammonite zone (Delsate et al., 2018; Popov, Delsate & Felten, 2019).

Fossilised diversity data

We extracted the number of fossil collections of Jurassic (Hettangian-Tithonian) marine reptiles (Ichthyosauria, Plesiosauria, Thalattosuchia, Pleurosauria, Angolachelonia) from the paleobiology database (<https://paleobiodb.org>) on the 26th March 2020. We binned these data per geological stage and generate a plot of the number of collections over time in R v3.6.2 (R Core Team, 2016), using the geoscale v2.0 package (Bell, 2015).

Centra proportions

We gathered a series of measurements on centra, recording their position within the vertebral column (Table 1). We treated and plotted these data in R v3.6.2 (R Core Team, 2016) to visualize changes through time, if any.

Institutional abbreviations

IRSNB: Royal Belgium Institute of Natural Sciences, Brussels, Belgium. **MNHNL:** Muséum national d’histoire naturelle du Luxembourg, Luxembourg-ville, Luxembourg. **ULg-PA:** Collections de paléontologie animale de l’Université de Liège, Liège, Belgium.

Comparative Descriptions

Thouarsense zone fauna, upper Toarcian, Luxembourg

ICHTHYOSAURIA De Blainville, 1835

PARVIPELVIA Motani, 1999

THUNNOSAURIA Motani, 1999

BARACROMIA Fischer et al., 2013

Baracromia indet.

(Figure 3)

A single specimen (MNHNL TM212) is recorded from the Thouarsense zone: it is a partial articulated ichthyosaur comprising fifty-two centra, several ribs and gastralia, and two partial forefemora from the Dudelange locality. All the preserved pre-apical centra are markedly rounded and anteroposteriorly short, bearing close similarities with Jurassic

thunnosaurians such as *Stenopterygius* and *Ophthalmosaurus* (Buchholtz, 2001; Massare et al., 2006). Posterior dorsal centra maintain clear bicapital rib articulations and all ribs have a 8-shaped cross-section, which also suggest that this specimen is a thunnosaurian (Sander, 2000; V.F. pers. obs. on NHMUK 2003, R5465, R498). The posterior apical centra are strongly compressed laterally, suggesting the presence of a lunate tailfin; although the angle of the tailfin is unknown. The humerus is incompletely-preserved, lacking the capitulum. The dorsal trochanter is proximodistally short, as it does not extend to mid-shaft, while the deltopectoral crest is elongated and parallel to the long axis of the humerus. The small dorsal trochanter suggest that this taxon is not an ophthalmosaurid (Motani, 1999a). The humerus forms a prominent anterior process that possesses a conspicuous, anterodorsally facing facet. This facet exhibits a pitted bone texture which suggests the presence of a cartilaginous cap in vivo. While many Early Jurassic ichthyosaurians possess a mesiodistally-thick anterodistal expansion of the humerus (*Temnodontosaurus*, *Leptonectidae*, *Hauffiopteryx*, *Stenopterygius* (Johnson, 1979; McGowan & Milner, 1999; McGowan, 2003; McGowan & Motani, 2003; Maisch, 2008; Caine & Benton, 2011; Martin et al., 2012), this expansion here forms a flat facet but not for articulation with an anterior accessory epipodial element (the rounded and notched anterior surface of the radius confirms the absence of such an element). This combination of features rules out earliest neoichthyosaurians and ophthalmosaurids (e.g. Maisch & Matzke, 2000; McGowan & Motani, 2003) and has only been reported in the poorly known, ?late Toarcian-lower Bajocian taxon *Dearcmhara shawcrossi* (Brusatte et al., 2015), although the facet is smaller and deeper in *D. shawcrossi*. Epipodial and proximal elements closely resemble those of *Stenopterygius* spp. and *Chacaicosaurus cayi* (Johnson, 1979; Fernández, 1994; Maxwell, Fernández & Schoch, 2012), being polygonal and dorsoventrally thick, with no spatium interosseum. The radius and the radiale are notched as in many early neoichthyosaurians (e.g. Huene, 1922; McGowan, 1974; Maxwell, 2012; Lomax & Massare, 2016). The intermedium forms two distal facets unequal in size: a large, distally facing facet for articulation with distal carpal 3 and a smaller, posterodistally facing facet for articulation with distal carpal 4 (the ‘latipinnate’ condition, although the latipinnate-longipinnate dichotomy fails to capture what is essentially a trait with continuous spectrum; see Mazin (1982) and Motani (1999b) for discussions). The specimen MNHNL TM212 is regarded as a non-ophthalmosaurid baracromian, resembling *Stenopterygius* spp., *Chacaicosaurus cayi*, and *Dearcmhara shawcrossi*.

Pseudoradiosa zone fauna, upper Toarcian, Luxembourg

PLESIOSAURIA De Blainville, 1835
Rhomaleosauridae Nopcsa, 1928

Rhomaleosauridae indet.

(Figure 4)

Two large plesiosaurian propodials have been recovered from **Pseudoradosa** zone: one (MNHNL DOU307) from the “Couche grise” of the Rodange locality, and one (MNHNL KA109) from the “Couche noire” of the Esch-sur-Alzette locality.

MNHNL DOU307 is a complete left propodial (proximal-distal length: 390mm; distal width: 210mm) with a weak posterior and dorsal curvature. The shaft is as wide as the capitulum, and there is no anterior expansion indicating that this propodial is an early plesiosaurian humerus (Storrs, 1997; Bardet, Godefroit & Sciau, 1999; O’Keefe, 2004; Smith & Vincent, 2010; Benson, Evans & Druckenmiller, 2012; Vincent & Storrs, 2019). The dorsal tuberosity is as anteroposteriorly wide as the capitulum and is weakly demarcated. A bulge with muscle scars is present on the ventral surface of the shaft, 30 mm distally to the capitulum. A very short preaxial flange is present distally, forming a small, anterodistally facing triangular facet. The radial/tibial facet is flat and marks an angle of ca. 45° with the ulnar/fibular facet. This facet is convex and rugose, indicating the presence of an extensive cartilage layer. A larger postaxial flange is present, forming distally a semioval facet that is hardly discernible from the ulnar/fibular facet. There is no distal ridge, unlike in the cryptoclidid *Colymbosaurus* (Benson & Bowdler, 2014). The absence of a marked curvature (the anterior surface is straight in dorsal and ventral views) and the presence of a conspicuous postaxial flange suggest rhomaleosaurid affinities (Smith & Vincent, 2010; Smith & Benson, 2014; Smith, 2015) (with the exception of *Lindwurmia thiuda* Vincent & Storrs, 2019), which are reinforced by the large size of the propodial.

MNHNL KA109 is the distal end of a plesiosaurian propodial; the anterior-posterior asymmetry suggests it is also a humerus. It resembles MNHNL DOU307 in lacking a preaxial flange and the postaxial flange is smaller than in MNHNL DOU307. The distal surface is markedly rounded in dorsoventral view. A prominent, strongly pitted anteroposterior ridge textures of the distal surface, obliterating radial/tibial and ulnar/fibular facets. The postaxial flange forms a triangular and concave facet posterior to the distal ridge. A distal ridge is regarded as an autapomorphic feature of *Colymbosaurus* (Benson & Bowdler, 2014; Arkhangelsky et al., 2019) but the lack of an extensive postaxial flange **precludes such a referral**; the size of the humerus and its similarities with MNHNL DOU307 suggest rhomaleosaurid affinities.

Dispansum-Pseudoradosa-Aalensis zones, upper Toarcian, Belgium

ICHTHYOSAURIA De Blainville, 1835

Ichthyosauria indet.

(Figure 5)

IRSNB Vert-06455-0001, IRSNB Vert-00000-00801, and IRSNB Vert-06659-0001 are ichthyosaurian centra, heavily eroded and damaged because their original shapes cannot be asserted, we refrained from calculating their shape ratios and assign them to Ichthyosauria indet.

IRSNB Vert-06462-0003 is a distal tooth that preserves the crown and the acellular cementum ring. The crown is recurved and robust (height: 14.83 mm, basal diameter: 8.14 mm, giving a crown shape ratio of 1.82), as is usually the case for distal ichthyosaurian teeth (Motani, 1997; McGowan & Motani, 2003). Apicobasal ridges texture the crown, unlike in leptonektids (Maisch, 1998; Maisch & Matzke, 2003; McGowan & Motani, 2003; Lomax, 2016) and the acellular cementum is itself ridged. This combination of features match that of *Ichthyosaurus*, *Protoichthyosaurus*, and *Temnodontosaurus* (McGowan, 1973; Godefroit, 1993a; Vincent et al., 2014; Brusatte et al., 2015; Lomax, Porro & Larkin, 2019). But it is also found in more primitive taxa such as *Contectopalatus* and *Cymbospondylus* (Maisch & Matzke, 2001; Klein et al., 2020). As a result, this specimen is referred to as Ichthyosauria indet.

IRSNB Vert-00000-00802 is a partial ichthyosaurian right quadrate, preserving the condyle, the stapedial facet and the posteroventral part of the anterior lamella. The condyle is massive, oval, but appears moderately short anteroposteriorly. The medial and lateral surfaces are pitted by numerous small foramina; the stapedial facet is semi-circular thickened ventral and posterior edges. This quadrate has the same size (maximal condylar width: 71 mm) and morphology as that of *Temnodontosaurus platyodon* (Godefroit, 1993a). In the absence of additional evidence, this quadrate is referred to Ichthyosauria indet. but indicates the presence of large ichthyosaurs by the latest Toarcian.

PARVPELVIA Motani, 1999

Parvipelvia indet.

(Figure 5)

ULg-PA 35.961 is an ichthyosaurian anterior dorsal centrum with a damaged dorsal surface. Its height-length ratio is 78 mm/35 mm = 2.23. The centrum is rounded as in parvipelvians (Merriam, 1908; Maisch & Matzke, 2000) and the diapophyses and parapophyses strongly protrude from the body of the centrum, as in MNHNL TM 212.

IRSNB Vert-00000-00800 is a partial left ichthyosaurian surangular, probably associated with the angular IRSNB Vert-06462-0002. It consists of posterior third, where the angular in a mesiolaterally-flattened bone with a slightly convex lateral surface and a slightly concave medial surface for accommodation of the Meckelian canal. The dorsal surface of the surangular is saddle-shaped, forming a long and shallow concavity anterior to the coronoid process. The preserved morphology of IRSNB Vert-00000-00800 is not diagnostic, but we assign it to Parvipelvina indet. because of its probable association with the angular IRSNB Vert-06462-0002 (see below).

IRSNB Vert-06462-0002 is a partial left ichthyosaurian angular bearing the typical double-grooved dorsal surface. The lateral wall of the angular is dorsoventrally short anteriorly, and only increases in size posterior to the level of the coronoid process. This narrow lateral exposure of the angular suggests that this specimen does not belong to Ophthalmosauridae, as this clade is notably characterized by large exposure of the angular in lateral view (Motani, 1999a).

IRSNB Vert-00000-00803 and IRSNB Vert-00000-00804 are large ichthyosaurian centra. IRSNB Vert-00000-00803 is 118 mm high and 48 mm long, giving a height-length ratio of 2.45; IRSNB Vert-00000-00804 is damaged on the edges and will not be measured but is of comparable size. The lateral surfaces of these centra are rounded in anteroposterior view, as in parvipelvians (Merriam, 1908; Maisch & Matzke, 2000), but their size departs from that of Early Jurassic thunnosaurians, even particularly large individuals (e.g. Maxwell, 2012; Lomax & Sachs, 2017). These centra rather match the size and shape of the dorsal-caudal centra of some large non-thunnosaurian neoichthyosaurians such as *Temnodontosaurus* spp. (Godefroit, 1993a; Martin et al., 2012) and *Excalibosaurus costini* (McGowan, 2003). IRSNB Vert-00000-00803 possesses a single apophysis (the parapophysis) placed laterally, suggesting it is middle preflexural caudal (Huene, 1922; McGowan & Motani, 2003). IRSNB Vert-00000-00804 possesses a diapophysis and a parapophysis placed ventrolaterally and is regarded as a posterior dorsal centrum. The size and shape of IRSNB Vert-00000-00803 and IRSNB Vert-00000-00804 match those of large early neoichthyosaurians and but are regarded as Parvipelvina indet. in the absence of unambiguous synapomorphies.

IRSNB Vert-00000-00808 is an ichthyosaurian anterior caudal centrum. Its height-length ratio is 65 mm / 22 mm = 2.94; this high shape ratio close to 3 might suggest thunnosaurian affinities (Buchholtz, 2001). The lateral surface of the centrum is rounded as in parvipelvians (Merriam, 1908; Maisch & Matzke, 2000) and slightly oval, as the ventral surface is flattened. The parapophyses are located lateroventrally.

IRSNB Vert-06462-0005 is an ichthyosaurian cervical centrum's height-length ratio is 63 mm / 24 mm = 2.62. The lateral surface of the centrum is rounded as in parvipelvians (Merriam, 1908; Maisch & Matzke, 2000). A faint ventral keel is present, giving the centrum a pentagonal shape. Such a shape is often seen in the cervical/anterior dorsals of parvipelvians (e.g. Huene, 1922; Fischer et al., 2014b; Lomax, Porro & Larkin, 2019) and is not of diagnostic value at a lower taxonomic level at the current state of our knowledge.

PLESIOSAURIA De Blainville, 1835

Plesiosauria indet.

IRSNB Vert-11312-00007 is a plesiosaurian dorsal centrum, as evidenced by the absence of transverse processes. Its height-length ratio is 67 mm / 44 mm = 1.52. The centrum is poorly preserved and we assign it to Plesiosauria indet.

Aalensis zone fauna, upper Toarcian, Luxembourg

The material from the Aalensis zone of the Minette ironstones is more abundant. We identified 12 ichthyosaurs and 8 plesiosaurians in the MNHL collections. Most of the material consist of isolated centra, complemented by three plesiosaurian propodials, one ichthyosaur coracoid, and one plesiosaurian tooth crown.

ICHTHYOSAURIA De Blainville, 1835

PARVIPELVIA Motani, 1999

Parvipelvia indet.

(Figure 6)

MNHL DOU94 is an ichthyosaurian anterior dorsal centrum from an unknown locality. A central keel is not present but the centrum forms a ventral expansion, giving the centrum a slight pentagonal shape. The diapophysis is small and rounded, as in parvipelvians and unlike in more primitive forms (Merriam, 1908; Huene, 1922) and is strongly protruding and has a rounded distal surface. Shallow dorsoventral ridges connect to the base of the diapophysis, anterior and posteriorly. The parapophysis is flatter and less prominent and its anterior margin merges with the anterior edge of the centrum. Such a pentagonal shape is often seen in parvipelvians (e.g. Huene, 1922; Fischer et al., 2014b; Lomax, Porro & Larkin, 2019) and is not of diagnostic value at a lower taxonomic level at the current state of our knowledge.

MNHNL DOU998 is an anterior caudal centrum from an unknown locality. The centrum has a rounded outline, as in usually parvipelvian ichthyosaurians (Merriam, 1908; e.g. Maisch & Matzke, 2000; McGowan & Motani, 2003). Its height/length ratio is $90/31=2.9$. This high ratio close to 3 might suggest thunnosaurian affinities (Buchholtz, 2001), but in the absence of additional data, this specimen is regarded as Parvipelvia indet.

MNHNL DOU369 is a series of disarticulated ichthyosaur centra. One originates from the mid-thoracic region and two others from the anterior caudal region, their respective height/length ratios are as follows: $68/32=2.1$, $61/35=1.74$, and $63/32=1.96$. The mid-thoracic resembles MNHNL DOU944. The rather elongated anterior caudal suggest these centra do not belong to a thunnosaurian, but rather to an early parvipelvian (Buchholtz, 2001).

MNHNL DOU351 is a fragmentary left coracoid from an unknown locality. It preserves the lateral part and a fragment of the intercoracoid facet. The ventral surface is moderately saddle-shaped (intercoracoid facet is only 56mm thick while the mesiolateral length is 180mm). The glenoid facet is large (anteroposterior length=75mm) and oval, while the scapular facet forms a small triangle (anteroposterior length= 39mm) and forms a ca. 60° angle with the glenoid facet. The posterolateral emargination is wide and deep, the coracoid surface posterior to the glenoid faces posteriorly and not posterolaterally, similar to the condition seen in *Ichthyosaurus communis*, *Ichthyosaurus anningae*, *Protoichthyosaurus postaxialis*, and many but not all specimens of *Temnodontosaurus platyodon* (Home, 1819; Godefroit, 1993a; Lomax & Massare, 2015; Lomax, Porro & Larkin, 2019). The Triassic shastasaurid *Shatasaurus neoscapularis* also possesses a wide posterior notch, but the coracoid of that taxon is different, having a very wide anterior notch (McGowan, 1994). We regard this coracoid as belonging to a non-baracromian parvipelvian.

MNHNL DOU378 is a very large centrum from an unknown locality. It originates from the posterior dorsal – sacral – anterior caudal region (length: 65mm and diameter: 175mm, estimated height: 180-185mm, H/L ratio: 2.79-2.84). Its size departs from the Early Jurassic thunnosaurians (*Ichthyosaurus* and *Stenopterygius*, which have much smaller centra even particularly large individuals (e.g. Maxwell, 2012; Lomax & Sachs, 2017)). It rather matches the size and shape of the anteriormost caudal centra of some large non-thunnosaurian neoichthyosaurians such as *Temnodontosaurus* spp. (Godefroit, 1993a; Martin et al., 2012) and *Excalibosaurus costini* (McGowan, 2003) (the anterior caudals are much longer in *Eurhinosaurus longirostris* [V.F., pers. obs. on MNHN 1946-20]).

PLESIOSAURIA De Blainville, 1835

Plesiosauria indet.

~~(Figures 7, 8)~~

MNHNL DOU906 is a plesiosaurian tooth crown from the Esch-sur-Alzette locality. It is slender (length/basal diameter= $23/8=2.875$), labiolingually flattened and recurved indicating it does not belong to derived rhomaleosaurids and thalassophoneans (Owen, 1865; Smith & Vincent, 2010; Ketchum & Benson, 2011b; Smith & Benson, 2014). The crown bears fine striations along the entire surface.

MNHNL DOU722 is a plesiosaurian sacral vertebra probably originating from the Esch-sur-Alzette locality. The articular surface is strongly dorsoventral depressed and the centrum is as wide as long. As in MNHNL DOU722, MNHNL DOU723, and MNHNL DOU954, the subcentral foramina are widely spaced and are separated by a well-rounded/convex surface. Similar to the other centra, the dorsal edge of the articular surface is notched by the neural canal. The transverse process is elongated, stout, and points posterolaterally. The rib facet is 8-shaped (waisted), this time with a larger, ear-shaped dorsal part and a smaller, semioval ventral part. The neural canal is also 8-shaped, resembling a 'plesiosauroid' pectoral vertebra from the Aalenian-Bajocian of Australia (Kear, 2012). Despite having similar size to the vertebrae we referred to as *Rhomaleosauridae* indet., none of the feature we highlight above are recorded in *Rhomaleosaurus* (Smith, 2013; Smith & Benson, 2014) and their representation in other derived rhomaleosaurids is unclear. Accordingly, we regard this specimen as *Plesiosauria* indet.

Rhomaleosauridae Nopcsa, 1928

Rhomaleosauridae indet.

~~(Figures 7, 8)~~

MNHNL DOU722 and MNHNL DOU723 are two plesiosaurian pectoral vertebrae from the Esch-sur-Alzette locality. The articular surface of the centrum is oval, with a dorsal notch corresponding to the neural canal. The subcentral foramina are displaced laterally, being located on the ventrolateral surface. There is no ventral facet, but the ventral surface in between the subcentral foramina is clearly transversely convex. The rib facets are oval, connect to the neural arch, and their long axis is vertical; they extend ventrally below the level of the notochordal pit. The semi-circular prezygaphophyses face dorsomedially and are clearly separated from one another; their ventral margin is set below the centre of the

neural canal, even though the neural canal is large. Anteriorly, the neural spine forms a basal triangular cavity containing a paired ridge. The height of this trough is about 1/3 of the total height of the neural spine. On the posterior surface, the basal region of the neural spines forms a deep tear-shaped concavity that extends for about one half of the total height of the neural spine, and appears similar to the condition of Microcleididae (Bardet, Godefroit & Sciau, 1999; Schwermann & Sander, 2011) and *Rhomaleosaurus* (Smith & Benson, 2014). However, the neural spines are much shorter than in microclidids (e.g. Owen, 1865; Bardet, Godefroit & Sciau, 1999). The neural spine possesses a posterior ridge while its anterior surface is concave, as in *Rhomaleosaurus thortoni* (Smith & Benson, 2014). The neural spine thickens dorsally and its dorsal half is posteriorly inclined, unlike in *Westphaliasaurus simonensis* (Schwermann & Sander, 2011). We assign these vertebrae to Rhomaleosauridae indet.

MNHNL DOU954  posterior pectoral/anterior dorsal vertebra from an unknown locality. It is essentially similar to MNHNL DOU722 and MNHNL DOU723. The main difference is the existence of ventral expansion of the centrum, even if a clear ventral keel remains absent. This ventral expansion gives the articular surface of the centrum a heart shape. The neural spine is concave anteriorly and bears a median ridges similar to the condition of *Rhomaleosaurus thortoni* (Smith & Benson, 2014). We assign this vertebra to Rhomaleosauridae indet.

MNHNL DOU558  from the Belvaux locality and MNHNL DOU324A and MNHNL DOU324B (broken, juvenile) from the Esch-sur-Alzette locality are three plesiosaurian humeri. These strongly resemble MNHNL DOU307: the shaft is long and slightly curved posteriorly and dorsally, the dorsal tuberosity is wide, a boss with muscle scars is present ventrally to the capitulum, the preaxial lamella is very short and forms a small triangular distal facet, the postaxial lamella is longer and forms a semioval posterodistal facet, radial/tibial and ulnar/fibular facets form a 45-55° angle and no distal ridge is present. The only difference is the position of the dorsal tuberosity, which is offset ventrally, creating a flattened, dorsomedially-facing surface between the dorsal part of the capitulum and the ventral part of the dorsal tuberosity. The larger propodial (MNHNL DOU324A) measures 331mm in proximal distal length and 168mm is maximal distal width. The smaller 'adult' propodial (MNHNL DOU558) measures 309mm in proximodistal length and 150mm in distal width.

Microcleididae Benson et al. 2012

Microcleidus Watson, 1909

cf. *Microcleidus*

~~(Figure 7)~~

MNHNL DOU978 is a plesiosaurian cervical centrum from an unknown locality. The centrum is slightly wider than high and is markedly elongated (height=42mm, width=53mm, length=55mm) unlike in derived rhomaleosaurids and thalassophoneans, which have anteroposteriorly short centra (Owen, 1865; Smith & Vincent, 2010; Ketchum & Benson, 2011b; Smith & Benson, 2014). The ventral surface is flattened as in *Microcleidus* (Owen, 1865; Vincent et al., 2017) and bears two subcentral foramina. There is no ventral keel. The lateral surface bears a (very) faint anteroposterior ridge; a more conspicuous lateral ridge is present in *Microcleidus* spp. (Owen, 1865; Bardet, Godefroit & Sciau, 1999; Vincent et al., 2017). The neural arch is fused with the centrum. The orientation of bone fibers on the right side suggests that the suture is V-shaped, which is mainly seen in rhomaleosaurids (Benson, Evans & Druckenmiller, 2012) and in the early pliosaurid *Hauffiosaurus* (Benson et al., 2011). The neural canal is very small in diameter, resulting in zygapophyses that are located close to the centrum, as in *Microcleidus* (Owen, 1865; Bardet, Godefroit & Sciau, 1999) and unlike in *Plesiosaurus dolichodeirus* (Owen, 1865), *Westphaliasaurus simonensis* (Schwermann & Sander, 2011), *Hauffiosaurus* spp. (White, 1940; Benson et al., 2011) The prezygapophyses are clearly separated, medially-inclined and their ventral surface is located more ventrally than the centre of the neural canal. The rib facets is 8-shaped, being formed of two semi-ovals separated by an anterior and a posterior furrows, as in *Plesiosaurus dolichodeirus* (Owen, 1865; Storrs, 1997) and unlike in many microcleidids (Owen, 1865; Schwermann & Sander, 2011), *Hauffiosaurus* (White, 1940), and cryptoclidids (Andrews, 1910a; Brown, 1981; Knutsen, Druckenmiller & Hurum, 2012a,b). The rib facet in posterior cervical centra is separated by a groove *Microcleidus melusinae* (Vincent et al., 2017). This centrum thus bears strong similarities with *Microcleidus* and is regarded as cf. *Microcleidus*.

Upper Aalenian fauna

ICHTHYOSAURIA De Blainville, 1835

Ichthyosauria indet.

(Figure 9)

MNHNL HU242 (Concavum zone) is a small partial ichthyosaur rostrum from the Rumelange locality. It contains the anterior part of the left dentary (or right premaxilla), showing the labial and lingual walls. Two teeth are preserved in situ. The root and the acellular cementum ring have a circular cross-section and are perfectly smooth externally, unlike in *Ichthyosaurus*, *Protoichthyosaurus*, and *Temnodontosaurus* (McGowan, 1973; Godefroit, 1993a; Vincent et al., 2014; Brusatte et al., 2015; Lomax, Porro & Larkin, 2019) and there is no evidence for plicidentine (unlike other Jurassic ichthyosaurs for

which this feature is known (Maxwell, Caldwell & Lamoureux, 2012)), but this latter feature cannot be used for a taxonomic purpose at the present state of our knowledge. The teeth are of a maximal diameter of 8mm and estimated 20mm apicobasal length; the size and shape of the tooth fits within the range of early baracromians (Godefroit, 1993b, 1994a; Fernández, 1994; Maisch, 2008; Maxwell, Fernández & Schoch, 2012) but in the lack of additional evidence, this material is here referred to Ichthyosauria indet.

PLESIOSAURIA De Blainville, 1835

Plesiosauria indet.

~~(Figure 9)~~



MNHNL HU384 (Murchisonae zone) is a small caudal centrum of a plesiosaurian of a juvenile individual, originating from the Tétange locality. Its height/width ratio is $32/36\text{mm} = 0.88$. Two small triangular chevron facets are present ventrally, merging with the edge of the articular surface. The ventral surface is flat. A rounded rib facet is present on the lateral surface. The neural arc is disconnected from the centrum and absent; the facets appear diamond shaped. Sacral vertebrae usually have less diagnostic features than other vertebrae in plesiosaurians, as could be inferred from their respective number of phylogenetic characters (Benson & Druckenmiller, 2014); this, coupled to the immaturity of the specimen would make an assignment ambiguous and we refer this material to Plesiosauria indet.

PLESIOSAUROIDEA Gray, 1825

Plesiosauroidea indet.

(Figure 9)



MNHNL BU157 (Concavum or Murchisonae zone) is a moderately elongated plesiosaurian cervical vertebra from the Rumelange locality. The shape of the centrum suggests it does not belong to derived rhomaleosaurids and thalassophoneans, which have anteroposteriorly short centra (Owen, 1865; Smith & Vincent, 2010; Ketchum & Benson, 2011b; Smith & Benson, 2014). The ventral surface is flattened and bears two subcentral foramina. No ventral keel is present. The edge of the centrum is rugose, as sometimes seen in various plesiosaurian clades (Owen, 1865; Seeley, 1874a; Fischer et al., 2020). An anteroposteriorly elongated bulge (rather than a ridge) is present on the dorsolateral surface. This bulge is separated from the rib facet by a median concave area. The rib facet is roughly oval but is very strongly waisted by an anterior and a posterior notch, giving the facet a marked 8 shape; this structure recalls – but is more marked than in – the late Toarcian centra MNHNL DOU978. The neural arch is fully fused to the centrum and no suture is discernible. A pair of supracentral foramina is present on the floor of foramen magnum. While MNHNL DOU978 could clearly be attributed to

Microcleidus, this vertebra – although superficially similar – is less elongated, possesses a lateral bulge rather than a ridge, and does not preserve the zygapophyses. As a result, a referral to Microclididae is too ambiguous. Nevertheless, the presence of an 8-shaped rib facet precludes a referral to cryptoclidids (Andrews, 1910a; Brown, 1981; Knutsen, Druckenmiller & Hurum, 2012a,b). Accordingly, this specimen is referred to as a non-cryptoclidid plesiosauroid.

Lower Bajocian fauna

ICHTHYOSAURIA De Blainville, 1835

Ichthyosauria indet.

~~(Figure 10)~~



MNHNL BM360 from the Rumelange locality contains one partial neural spine and two proximal parts of bicipital ribs. The neural spine bears a dorsoventrally-oriented median ridge on its anterior surface. The taxonomic values of these features is low and we assign this material to Ichthyosauria indet.

PARVPELVIA Motani, 1999

Parvipelvian indet.

~~(Figure 10)~~



MNHNL BM758 is an ichthyosaurian caudal centrum from the Rumelange locality. Its shape is rounded, with a slightly flattened ventral surface, as in Parvipelvian (Merriam, 1908; Maisch & Matzke, 2000). Its height/length ratio is $100\text{mm}/38\text{mm}=2.63$.



MNHNL BM725 from the Rumelange locality contains a cervical and a caudal ichthyosaurian centra. The cervical centrum has a height/length ratio of $83\text{mm}/38\text{mm}=2.18$ while the same ratio for the caudal centrum is $56\text{mm}/26\text{mm}=2.15$. The centra are rounded in shape and exhibit small and rounded apophyses, as in parvipelvians and unlike in more primitive forms (Merriam, 1908; Huene, 1922).



MNHNL BM461 is an ichthyosaurian cervical centrum from the Rumelange locality. Its height/length ratio is $72\text{mm}/30\text{mm}=2.4$. The centrum is rounded in shape and exhibits small and rounded apophyses, as in parvipelvians and unlike in more primitive forms (Merriam, 1908; Huene, 1922).



MNHNL BM392 from the Rumelange locality contains five caudal centra and one sclerotic plate of a single ichthyosaurian specimen. The height/length ratios of the first four caudal centra are as follows: $72\text{mm}/30\text{mm}=2.4$, $77\text{mm}/31\text{mm}=2.48$,

72mm/29mm=2.48, 75mm/32mm=2.34. Their rounded shape indicates they belong to Parvipelvina (Merriam, 1908; Huene, 1922; Maisch & Matzke, 2000). The sclerotic plate bears radiating striations on its lateral surface and its internal and external edges are crenulated, as is usually the case in ichthyosaurians (Andrews, 1910a; McGowan, 1973; Fischer et al., 2014a). The external third of the plate is deflected, facing dorsally.

OPHTHALMOSAURIDAE Baur, 1897

Ophthalmosauridae indet.

~~(Figures 10, 11)~~



MNHNL BM780–BM781 is a nearly complete right ichthyosaurian surangular (MNHNL BM780), associated with seven teeth (bearing the collection number MNHNL BM781), originating from the Rumelange locality. The surangular is straight in all planes (only its posterior quarter is slightly deflected medially) and bears a thickened dorsal margin, giving it a tear-shaped cross-section. A lateral, anteroposteriorly elongated and posteriorly-deepening concavity is present on the lateral surface ('surangular fossa'). This depressed area terminates anteriorly to the level of the coronoid process. The coronoid process is prominent and bears a rugose texture. A small, anteroposteriorly oriented ridge is present directly posteromedially to the coronoid process and is likely part of the muscle attachment. The dorsal margin of the surangular forms a concave plateau posteriorly to the coronoid process. A prominent M.a.m.e process is present directly posteromedially to the plateau; this process points dorsomedially. The angular facet extends anteriorly up to the level of the coronoid process and covers the ventral half of the surangular posteriorly. This indicates the presence of an extensive angular, which is an ophthalmosaurid synapomorphy (Motani, 1999a). The teeth are peculiar in forming externally-visible plicidentine, texturing the acellular cementum ring and the root by very deep apicobasal grooves, as in *Ichthyosaurus*, *Protoichthyosaurus*, and *Temnodontosaurus* (McGowan, 1973; Godefroit, 1993a; Vincent et al., 2014; Brusatte et al., 2015; Lomax, Porro & Larkin, 2019). The root cross-section is oval as in all parvipelvians bar most platypterygiines (Fischer et al., 2012; Fischer, 2016) and the base of the enamel is easy to discern, which is usually the case in ophthalmosaurids (Fischer et al., 2016). Accordingly, we refer this specimen as Ophthalmosauridae indet.



MNHNL BM779 is a fragmentary partially articulated ichthyosaurian, containing parts of the skull, axial skeleton, and scapular girdle, originating from the Rumelange locality. It belongs to Ophthalmosauridae, having a humerus with plate-like dorsal trochanter, a massive deltopectoral crest, a humeral facet for an anterior accessory element, a posterodistally deflected ulnar facet and a conspicuous acromial process on the scapula (Motani, 1999a; Fischer et al., 2012, 2013; Moon, 2017; Zverkov & Efimov, 2019). It resembles *Arthropterygius* in having an anteroposteriorly-short parietal symphysis, a

deep ulnar facet on the humerus, a small humeral facet for an anterior accessory epipodial elements (Maxwell, 2010; Fernández & Maxwell, 2012; Zverkov & Prilepskaya, 2019). However, the presence of several autapomorphies on the cranial and appendicular elements indicates that this specimen constitutes a novel taxon, for which a dedicated manuscript is in preparation.

PLESIOSAURIA De Blainville, 1835

CRYPTOCLIDIDAE Williston, 1925

Cryptoclididae indet.

(Figure 10)

MNHNL BM782 is a left plesiosaurian propodial from the Rumelange locality. The propodial is straight in the proximodistal direction and slightly curves dorsally. The shaft is fairly elongated. The dorsal tuberosity is wide and weakly demarcated from the capitulum; they form together an evenly rounded proximal surface. A median dorsal boss surrounded by muscle scars is present close to the distal end of the dorsal tuberosity. This boss is median (along the axis of the shaft) and thus differs from the anteriorly placed boss seen in the humeri of rhomaleosaurids and microcleidids (Owen, 1865; Smith & Benson, 2014). The propodial markedly expands distally, forming a preaxial flange and a longer postaxial flange. The preaxial flange forms a dorsoventrally-narrow triangular facet distally. The postaxial flange forms a semioval posterior facet that forms a marked angle (ca. 45°) with the ulnar/fibular. Both the radial/tibial and the anteroposteriorly- and dorsoventrally-shorter ulnar/fibular facets are slightly convex. However, a distal ridge is absent, unlike in *Colymbosaurus* (Benson & Bowdler, 2014; Arkhangelsky et al., 2019). The shape of the propodial with its fairly slender shaft and large preaxial and postaxial flanges indicate cryptocleidid affinities (Andrews, 1910a; Mehl, 1912; Brown, 1981; O’Keefe & Wahl, 2003; Knutsen, Druckenmiller & Hurum, 2012c; Roberts et al., 2017, 2020); it resembles the humerus of humerus of *Muraenosaurus leedsi* (Seeley, 1874b; Andrews, 1910a) and the femur of *Tricleidus seeleyi* (Owen, 1865).

Results

We visualized the size and shape of Toarcian and Bajocian ichthyosaur centra (Figure 12). These data reveal a decrease in overall size and a much smaller range of shapes from the late Toarcian to the Bajocian. However, a difference in sampling intensity between these two assemblages is evident. We randomly sampled our late Toarcian dataset to select 3 cervicals and 6 caudals 10000 times, yielding comparable datasets. We then compared the range of height-length ratios between these datasets. The Bajocian height-

length ratios range is smaller than Toarcian height-length ratios range about 99.85% of the times and two times smaller about 98.56% of the times. The data at hand thus ~~unambiguously~~ suggest a reduction of the disparity of ichthyosaurian centrum shapes across the Early–Middle Jurassic transition, i.e. going from an early neoichthyosaurian-dominated to an ophthalmosaurid-dominated assemblage.

Discussion

The Early–Middle Jurassic transition in marine reptiles, a state of the art

The very well-sampled (Benson et al., 2010) Lower Jurassic marine ecosystems of western Europe housed a vast menagerie of neoichthyosaurians, thalattosuchians, and plesiosaurians (Owen, 1860; Huene, 1922, 1931; McGowan, 1974, 1979; Benton & Taylor, 1984; Maisch & Matzke, 2000; O’Keefe, 2004; Großmann, 2007; Maisch, 2008, 2010; Benson, Evans & Druckenmiller, 2012; Martin et al., 2012; Bardet et al., 2014; Lomax & Massare, 2016; Johnson, Young & Brusatte, 2020). Evidence from body size, craniodental shape, and swimming capabilities suggests that these taxa occupied several niches within shallow marine ecosystems (Hauff, 1953; Massare, 1987  97; Böttcher, 1989; Godefroit, 1994a; McGowan, 1996; Buchholtz, 2001; Buchy, 2010; Fischer, Guimar & Godefroit, 2011; Martin et al., 2012; Dick, Schweigert & Maxwell, 2016; Maxwell & Cortés, 2020). Such a diversity of forms made the Early Jurassic marine ecosystems of western Europe an iconic representation of Mesozoic marine life (Taylor, 1997).

Even though a drastic reduction of apparent diversity is expected following the lagerstätten effect of the early Toarcian localities (Benson et al., 2010), the Middle Jurassic assemblages, when well sampled (i.e. not before the Callovian), are markedly distinct from their Toarcian counterparts. For ichthyosaurians, the dominant and diversified early neoichthyosaurians (leptonectids, *Suevoleviathan*, *Hauffiopteryx*, and *Temnodontosaurus*) are gone; the disappearance of non-thunnosaurian ichthyosaurians marks the end of large (>6m) ichthyosaurian top predators, at least up until the Aptian, when some derived platypterygiines will presumably fill similar roles (Fischer et al.,  2014b, 2016; Bardet, Fischer & Machalski, 2016; Fischer, 2016). Ichthyosaurians will never re-evolve carinated teeth like those seen in *Temnodontosaurus platyodon* and *Temnodontosaurus trigonodon* (Conybeare, 1822; Godefroit, 1993a), nor the sawfish-like morphology of Leptonectidae (Swinton, 1930; Huene, 1951; McGowan, 1986, 2003; Lomax, 2016). From the Callovian onwards, ichthyosaur assemblages overwhelmingly consist of ophthalmosaurids (but see Fischer et al. (2013)), up to the final extinction of ichthyosaurians (Fischer et al., 2016). While diverse ophthalmosaurid assemblages are known by the Kimmeridgian (Fernández, 1997; Druckenmiller et al., 2012; Arkhangel'sky

& Zverkov, 2014; Zverkov et al., 2015; Paparella et al., 2017; Moon & Kirton, 2018a; Delsett et al., 2019; Zverkov & Efimov, 2019; Campos, Fernández & Herrera, 2020; Zverkov & Jacobs, 2020), only the closely related *Ophthalmosaurus* and *Baptanodon* are known in the Callovian of Europe and North America, respectively (Seeley, 1874c; Marsh, 1895; Gilmore, 1902, 1906; Knight, 1903; Andrews, 1910b; Massare et al., 2006; Moon & Kirton, 2018b).

This leaves a fairly long interval (late Toarcian-Callovian) for which the ichthyosaur fossil record is very scarce and geographically dispersed: Sander & Bucher (1993) reported a large cf. *Stenopterygius* from the late Toarcian of southern France; Maisch & Matzke (2000: p72) mention the presence of *Temnodontosaurus* in the late Toarcian of southern Germany; Vincent et al. (2013) reported *Temnodontosaurus* sp., and *Stenopterygius*-like forms in the late Toarcian of southern France and a coracoid bearing many similarities with MNHN DOU353 from the early Aalenian of southern France; finally, Brusatte et al. (2015) described *Dearcmahara shawcrossi* from the late Toarcian-Bajocian of Scotland. The only diagnosable ichthyosaur remains unambiguously known from Aalenian deposits is the holotype of *Stenopterygius aalensis* from Germany (Maxwell, Fernández & Schoch, 2012) (see also Arnaud et al. (1976) and Maxwell et al. (2012) for additional indeterminate specimens). Fragmentary ophthalmosaurid specimens have been reported in the Aalenian-Bajocian boundary of Argentina (Fernández, 2003) and the early Bajocian of Argentina (Fernández, 1999; Gasparini et al., 2007) and Canada (Druckenmiller & Maxwell, 2014). It can be inferred from these data that a replacement of early neoichthyosaurians by ophthalmosaurids took place across the Early–Middle Jurassic transition, but the generalized absence of Aalenian-Bathonian ichthyosaur fossils worldwide and Early Jurassic fossils in South America make it difficult to identify a precise turnover.

While less abundant than ichthyosaurians in early Toarcian Lagerstätten (Hauff, 1953), plesiosaurians already had also evolved a vast array of morphotypes by the Toarcian (O’Keefe, 2001a,b; Benson, Evans & Druckenmiller, 2012; Smith & Araújo, 2017), presumably filling as many ecological niches: long-necked and small headed microclidids (Owen, 1865; Bardet, Godefroit & Sciau, 1999), gigantic, apex predatory rhomaeosaurids (Taylor, 1992; Cruickshank, 1994; Smith & Dyke, 2008; Smith & Vincent, 2010; Smith & Benson, 2014), and small-sized, moderately long-necked, and long-snout early pliosaurids (Benson et al., 2011; Ketchum & Benson, 2011a; Vincent, 2011; Fischer et al., 2017). Microcleidids supposedly go extinct after the Toarcian; their last definite record in the latest Toarcian of France (Sciau, Crochet & Mattei, 1990; Bardet, Godefroit & Sciau, 1999). However, an Aalenian specimen from France regarded as an indeterminate elasmosaurid by Vincent et al. (2007) possesses microcleidid features (8-shaped cervical rib facet, lateral ridge on cervical centra) and might possibly be

regarded as a member of that clade. Nevertheless, the long-necked morphotype is then colonised by cryptoclydids by the Callovian-Oxfordian (Brown, 1981).

Only a couple of rhomaleosaurid lineages will survive up to the Middle/Late Jurassic, in high latitudes (Gasparini, 1997; Sato & Wu, 2008; Benson, Zverkov & Arkhangelsky, 2015). Thalassophonean pliosaurids concomitantly radiated and became the main marine apex predators from the Middle Jurassic to the early Late Cretaceous (Noe, 1999, 2001; Ketchum & Benson, 2011a; Benson et al., 2013; Benson & Druckenmiller, 2014; Fischer et al., 2017; Zverkov et al., 2018). A narrative where thalassophoneans filled (either competitively or passively) the niches previously occupied by *Temnodontosaurus* and derived rhomaleosaurids is sensible, but these animals have clearly distinct body plans and probably hunted differently (Massare, 1988; Buchholtz, 2001; O'Keefe, 2001b, 2002). In any case, there is evidence for the continuous presence of gigantic apex predatory plesiosaurians in Europe and elsewhere: *Simolestes keileni* from the upper Bajocian of eastern France (described by Godefroit (1994b), and currently under revision by S. Sachs), an indeterminate fragmentary jaw from the lower Bajocian of Switzerland (Sachs, Klug & Kear, 2019) and 'pliosauroid' tooth from the Aalenian-Bajocian of Australia (Long & Cruickshank, 1998; Kear, 2012). Whether these specimens are rhomaleosaurids or thalassophoneans obviously yields different outcomes as they might push the origin of apex predatory pliosaurids to the Bajocian, but these fossils suggest that the niche of very large apex predators was not vacated for a long period, if at all.

Recent evidence revealed a series of basal metriorhynchoid crocodyliforms in the late Toarcian–Aalenian interval (Wilberg, 2015; Ösi et al., 2018; Aiglstorfer, Havlik & Herrera, 2020) suggesting an intense diversification across the Early-Middle Jurassic transition (Aiglstorfer, Havlik & Herrera, 2020). The effect of this radiation is evident by the late Middle and Late Jurassic, where abundant thalattosuchian taxa are known, occupying several ecological niches (Gasparini, Pol & Spalletti, 2006; Young & de Andrade, 2009; Young et al., 2012, 2014b; Herrera, Gasparini & Fernández, 2013; Foffa et al., 2017, 2018). Teleosauroids do not appear much affected by this transition, but this might be due to the problematic taxonomy of the group (Johnson, Young & Brusatte, 2020).

An abrupt turnover

In light of the short review above, it appears clear that fossiliferous successions in a geographically-restricted areas are crucial to decipher both the tempo and the severity of the marine reptile turnover at the early–Middle Jurassic transition. The late Toarcian–Bajocian successions of Luxembourg and Belgium we analysed above yields novel biostratigraphic information that precise the turnover dynamics of ichthyosaurians and

plesiosaurs during the Early–Middle Jurassic transition (the crocodyliform record recently reviewed by Johnson et al. (2018), is scarce and does not provide relevant information regarding this turnover). In a global, broad-brush view of marine reptile macroevolution, the presence of rhomaleosaurids, microcleidids, and non-thunnosaurian ichthyosaurs in the late Toarcian of Luxembourg (Table 2) was expected, as these clades are already known to survive after the early Toarcian Lagerstätten (Bardet, Godefroit & Sciau, 1999; Vincent et al., 2013; Benson, Zverkov & Arkhangelsky, 2015). With the exception of possible remains from the Aalenian of France (Vincent, Bardet & Morel, 2007; see discussion above), MNHNL DOU978 and the holotype of *Microcleidus tournemirensis* (Sciau, Crochet & Mattei, 1990; Bardet, Godefroit & Sciau, 1999) are the youngest-known microcleidid to date and indicate that the clade extended at least up to the very end of the Early Jurassic. Recording microcleidids, large rhomaleosaurids, *Temnodontosaurus*-like forms, and more derived ichthyosaurs within the same basin that formed the early Toarcian Lagerstätten yields important palaeobiogeographic implications. These fossils indicate that the main marine reptile clades of the Early Jurassic remained abundant and dominant in mid-latitude, epicontinental seas up to the end of the Toarcian at least. Although much less prolific than the underlying strata, the upper Toarcian fossil record suggest that no major turnover took place within the Toarcian as the structuration of the upper tier of marine ecosystems remained intact. The taxonomic diversity crash that follows the early Toarcian appears to be mainly a Lagerstätten effect, confirming previous suspicions (Benson et al., 2010; Maxwell & Vincent, 2015).

In stark contrast with the sense of continuity displayed by the upper Toarcian occurrences, the Bajocian fossil record of Luxembourg suddenly records the dominant clades of the Late Jurassic: cryptocleidids, ophthalmosaurids, and probable pliosaurids, with no evidence – so far – for the presence of more ancient clades. The specimen MNHNL BM782 is oldest-known cryptocleidid and MNHNL BM770 and MNHNL BM780–BM781 are the oldest ophthalmosaurids known, after a single partial forefin from the Aalenian–Bajocian boundary of Argentina (Fernández, 2003) and fragmentary basicranium from an equivalent of the Sauzei Zone in Canada (Druckenmiller & Maxwell, 2014), which directly underlies the Humphresianum Zone. The new occurrences we report here indicate that ophthalmosaurids rapidly dispersed, being almost simultaneously recorded in Canada, Argentina, and Luxembourg by the Early Bajocian.

The Belgo-Luxembourgian marine reptile record provides a clearer picture of the marine reptile turnover occurring at the Early–Middle Jurassic transition, indicating that this replacement is restricted to the sole Aalenian stage instead of possibly spanning the entire middle Toarcian–Bathonian interval. This reduces the uncertainty on the timing and

duration of this turnover to 4 instead of 14 million years and packs a series of extinctions and diversifications within a short period of abrupt climate cooling and changes in oceanic currents (Korte et al., 2015). This Aalenian shift from a Toarcian Warm Mode to Aalenian-Bajocian Cool Mode (Korte et al., 2015) also appears associated with a marked faunal disruption in belemnites, with random (i.e. non-morphologically-selective) extinctions leading to a distinct drop in belemnite biodiversity at least in the northwestern Peri-Tethys Ocean (Dera, Toumoulin & de Baets, 2016; Neige, Weis & Fara, 2020). This major disruption in the evolutionary history of Jurassic belemnites ended at the Aalenian-Bajocian boundary and resulted in a radiation of the suborder Belemnopseina that partially replaced the previously dominant Belemnitina in the Western Tethys (Weis, Mariotti & Riegraf, 2012; Weis, Sadki & Mariotti, 2017), furthermore entailing a distinct Boreal vs. Tethyan belemnite provincialism (Doyle, 1987; Mariotti, Santantonio & Weis, 2007; Weis & Mariotti, 2007; Mariotti et al., 2012; Dzyuba et al., 2019). At the present state of knowledge, we can only speculate about whether faunal changes in belemnites and marine reptiles were independently impacted by the same factors, or even causally connected, since belemnites are an essential component of the food spectrum in some marine reptiles (Massare, 1987; Böttcher, 1989; Dick, Schweigert & Maxwell, 2016), their faunal change could have triggered a disruption of the trophic chain up to the giant top predators.



Conclusions

A generalised turnover affected marine reptile communities across the Early–Middle Jurassic transition. However, the extremely poor fossil record of the middle Toarcian–Bathonian interval leaves a ca. 14 million years window of the uncertainty for this important event. Our thorough analysis of the marine reptile record of the late Toarcian to Bajocian successions in a confined palaeogeographic setting (the Belgo-Luxembourgian sub-basin) indicates that:

- (i) the structuration of the upper tier of marine ecosystems remained unchanged up to the very end of the Early Jurassic, with the presence of large rhomaleosaurid plesiosaurians, microcroclidid plesiosaurians, as well as *Temnodontosaurus*-like and baracromian ichthyosaurians in the late Toarcian.
- (ii) the dominant clades of the Late Jurassic, cryptoclidid plesiosaurians and ophthalmosaurid ichthyosaurians, arose and dispersed earlier than expected, being recorded in the early Bajocian of Luxembourg.
- (iii) the within-System marine reptile turnover of the Early–Middle Jurassic transition is more abrupt than previously supposed, being restricted to the sole Aalenian stage, i.e. 4 million years.



Acknowledgements

We thank the curatorial staff of the IRSNB for their help. We also thank the main contributors of the Paleobiology Database data we used: Prof. R.B.J. Benson, Dr. M.T. Carrano, Dr. J.P. Tennant, Dr. H.P. Street, Prof. R.J. Butler, Prof. M. Clapham and Prof. M. Uhen. Finally, we warmly thank Dr. M. Trotta for her help in generating geographic maps. Reviewers+Editor.

References

- Aiglstorfer M, Havlik P, Herrera Y. 2020. The first metriorhynchoid crocodyliform from the Aalenian (Middle Jurassic) of Germany , with implications for the evolution of Metriorhynchoidea. *Zoological Journal of the Linnean Society* 188:522–551.
- Andrews CW. 1910a. *A descriptive catalogue of the marine reptiles of the Oxford clay. Based on the Leeds Collection in the British Museum (Natural History), London. Part I.* London,; Order of the Trustees of the British Museum.
- Andrews CW. 1910b. Note on the osteology of *Ophthalmosaurus icenicus* Seeley an ichthyosaurian Reptile from the Oxford Clay of Peterborough. *Geological Magazine* 4:202–208.
- Arkhangelsky MS, Zverkov NG. 2014. On a new ichthyosaur of the genus *Undorosaurus*. *Proceedings of the Zoological Institute RAS* 318:187–196.
- Arkhangelsky MS, Zverkov NG, Rogov MA, Stenshin IM, Baykina EM. 2019. Colymbosaurines from the Upper Jurassic of European Russia and their implication for palaeobiogeography of marine reptiles. *Palaeobiodiversity and Palaeoenvironments* 100:197–218. DOI: 10.1007/s12549-019-00397-0.
- Arnaud M, Monleau C, Wenz S. 1976. Découverte de restes d'ichthyosaure dans l'Aalénien du Massif de la Loube (Var). *Bulletin du Musée d'Histoire Naturelle de Marseille* 36:3.
- Bakker RT. 1993. Plesiosaur Extinction Cycles — Events that Mark the Beginning, Middle and End of the Cretaceous. In: Caldwell WGE, Kauffman EG eds. *Evolution of the Western Interior Basin: Geological Association of Canada, Special Paper.* Stittsville, Ontario, Canada, 641–664.
- Bardet N. 1994. Extinction events among Mesozoic marine reptiles. *Historical Biology* 7:313–324.
- Bardet N. 1995. Evolution et extinction des reptiles marins au cours du Mésozoïque. *Palaeovertebrata* 24:177–283.
- Bardet N, Falconnet J, Fischer V, Houssaye A, Jouve S, Pereda Suberbiola X, Pérez-García A, Rage J-CJ-C, Vincent P. 2014. Mesozoic marine reptile palaeobiogeography in response to drifting plates. *Gondwana Research* 26:869–887. DOI: 10.1016/j.gr.2014.05.005.
- Bardet N, Fischer V, Machalski M. 2016. Large predatory marine reptiles from the Albain–Cenomanian of Annopol, Poland. *Geological Magazine* 153:1–16. DOI:

- 10.1017/S0016756815000254.
- Bardet N, Godefroit P, Sciau J. 1999. A new elasmosaurid Plesiosaur from the Lower Jurassic of Southern France. *Palaeontology* 42:927–952.
- Bardet N, Houssaye A, Rage J-C, Suberbiola XP. 2008. The Cenomanian-Turonian (Late Cretaceous) radiation of marine squamates (Reptilia): the role of the Mediterranean Tethys. *Bulletin de la Société géologique de France* 179:605–622.
- Bell MA. 2015. geoscale: Geological Time Scale Plotting.
- Benson RBJ. 2013. Marine Reptiles. In: MacLeod N, Archibald JD, Levin P eds. *Grzimek's Animal Life Encyclopedia, Extinction*. Farmington Hills, Michigan: Gale Cengage Learning, 267–279.
- Benson RBJ, Bowdler T. 2014. Anatomy of *Colymbosaurus megadeirus* (Reptilia, Plesiosauria) from the Kimmeridge Clay Formation of the U.K., and high diversity among Late Jurassic plesiosauroids. *Journal of Vertebrate Paleontology* 34:1053–1071. DOI: 10.1080/02724634.2014.850087.
- Benson RBJ, Butler RJ, Lindgren J, Smith AS. 2010. Mesozoic marine tetrapod diversity: mass extinctions and temporal heterogeneity in geological megabiases affecting the vertebrates. *Proceedings of the Royal Society B: Biological Sciences* 277:829–834.
- Benson RBJ, Druckenmiller PS. 2014. Faunal turnover of marine tetrapods during the Jurassic–Cretaceous transition. *Biological Reviews* 89:1–23. DOI: 10.1111/brv.12038.
- Benson RBJ, Evans M, Druckenmiller PS. 2012. High diversity, low disparity and small body size in plesiosaurs (Reptilia, Sauropterygia) from the Triassic–Jurassic boundary. *PLoS ONE* 7:e31838.
- Benson RBJ, Evans M, Smith AS, Sassoon J, Moore-Faye S, Ketchum HF, Forrest R. 2013. A giant pliosaurid skull from the Late Jurassic of England. *PLoS ONE* 8:e65989. DOI: 10.1371/journal.pone.0065989.
- Benson RBJ, Ketchum HF, Noè LF, Gómez-Pérez M. 2011. New information on *Hauffiosaurus* (Reptilia, Plesiosauria) based on a new species from the Alum Shale member (Lower Toarcian: Lower Jurassic) of Yorkshire, UK. *Palaeontology* 54:547–571.
- Benson RBJ, Zverkov NG, Arkhangel'sky MS. 2015. Youngest occurrences of rhomaleosaurid plesiosaurs indicate survival of an archaic marine reptile clade at high palaeolatitudes. *Acta Palaeontologica Polonica* 60:769–780.
- Benton MJ, Taylor MA. 1984. Marine reptiles from the Upper Lias (Lower Toarcian, Lower Jurassic) of the Yorkshire coast. *Proceedings of the Yorkshire Geological Society* 44:399–429.
- Böttcher VR. 1989. Über die Nahrung eines *Leptopterygius* (Ichthyosauria, Reptilia) aus dem süddeutschen Posidonienschiefer (Unterer Jura) mit Bemerkungen über den Magen der Ichthyosaurier. *Stuttgarter Beiträge zur Naturkunde Serie B (Geologie und paläontologie)* 155:1–19.
- Boulvain F, Belanger I, Delsate D, Ghysel P, Godefroit P, Laloux M, Roche M. 2001. Triassic and Jurassic lithostratigraphic units (Belgian Lorraine). *Geologica Belgica* 4:113–119.
- Brown D. 1981. The English Upper Jurassic Plesiosauroidea (Reptilia) and a review of the phylogeny and classification of the Plesiosauria. *Bulletin of the British Museum (Natural History) Geology* 35:253–347.

- 968 Brusatte SL, Young MT, Challands TJ, Clark NDL, Fischer V, Fraser NC, Liston JJ,
969 MacFadyen CCJ, Ross DA, Walsh SA, Wilkinson M. 2015. Ichthyosaurs from the
970 Jurassic of Skye, Scotland. *Scottish Journal of Geology* 51:43–55. DOI:
971 10.1144/sjg2014-018.
- 972 Buchholtz EA. 2001. Swimming styles in Jurassic ichthyosaurs. *Journal of Vertebrate*
973 *Paleontology* 21:61–73.
- 974 Buchy M-C. 2010. Morphologie dentaire et régime alimentaire des reptiles marins du
975 Mésozoïque : revue critique et réévaluation. *Oryctos* 9:49–82.
- 976 Caine H, Benton MJ. 2011. Ichthyosauria from the upper Lias of Strawberry Bank,
977 England. *Palaeontology* 54:1069–1093.
- 978 Callaway JM, Massare J. 1989. Geographic and stratigraphic distribution of the Triassic
979 Ichthyosauria (Reptilia; Diapsida). *Neues Jahrbuch für Geologie und Paläontologie,*
980 *Abhandlungen* 187:37–58.
- 981 Campos L, Fernández MS, Herrera Y. 2020. A new ichthyosaur from the Late Jurassic of
982 north-west Patagonia (Argentina) and its significance for the evolution of the narial
983 complex of the ophthalmosaurids. *Zoological Journal of the Linnean Society*
984 188:180–201. DOI: 10.1093/zoolinnean/zlz095.
- 985 Di Cencio A, Weis R. 2020. Revision of upper Toarcian ammonites (Lytocerotidae,
986 Graphoceratidae and Hammatoceratidae) from the Minette ironstones, southern
987 Luxembourg. In: Di Cencio A, Sadki D, Weis R eds. *Paléontologie au Luxembourg*
988 (2) *Les ammonites de la Minette*. Ferrantia, in press.
- 989 Conybeare WD. 1822. Additional notes on the fossil genera *Ichthyosaurus* and
990 *Plesiosaurus*. *Transactions of the Geological Society of London* 2:103–123.
- 991 Cruickshank a. RI. 1994. Cranial Anatomy of the Lower Jurassic Pliosaur
992 *Rhomaleosaurus megacephalus* (Stutchbury) (Reptilia: Plesiosauria). *Philosophical*
993 *Transactions of the Royal Society B: Biological Sciences* 343:247–260. DOI:
994 10.1098/rstb.1994.0024.
- 995 Delsate D, Suberbiola XP, Felten R, Felten G. 2018. First thyreophoran dinosaur from the
996 Middle Jurassic (Bajocian) of Luxembourg. *Geologica Belgica* 21:19–26.
- 997 Delsate D, Weis R. 2010. La Couche à Crassum (Toarcien moyen) au Luxembourg:
998 stratigraphie et faunes de la coupe de Dudelange-Zoufftgen. *Ferrantia* 62:35–60.
- 999 Delsett LL, Roberts AJ, Druckenmiller PS, Hurum JH. 2019. Osteology and phylogeny
1000 of Late Jurassic ichthyosaurs from the Slotsmøya Member Lagerstätte (Spitsbergen,
1001 Svalbard). *Acta Palaeontologica Polonica*:717–743. DOI: 10.4202/app.00571.2018.
- 1002 Dera G, Toumoulin A, de Baets K. 2016. Diversity and morphological evolution of
1003 Jurassic belemnites from South Germany. *Palaeogeography, Palaeoclimatology,*
1004 *Palaeoecology* 457:80–97. DOI: 10.1016/j.palaeo.2016.05.029.
- 1005 Dick DG, Maxwell EE. 2015. The evolution and extinction of the ichthyosaurs from the
1006 perspective of quantitative ecospace modelling. *Biology letters* 11:20150339-. DOI:
1007 10.1098/rsbl.2015.0339.
- 1008 Dick DG, Schweigert G, Maxwell EE. 2016. Trophic niche ontogeny and palaeoecology
1009 of early Toarcian *Stenopterygius* (Reptilia: Ichthyosauria). *Palaeontology*:n/a-n/a.
1010 DOI: 10.1111/pala.12232.
- 1011 Dittrich D. 1993. *Erläuterungen zur geologischen Karte von Luxemburg, 1:25000: Blatt*
1012 *11 Grevenmacher und Blatt 13 Remic*.
- 1013 Doyle P. 1987. Lower Jurassic-Lower Cretaceous belemnite biogeography and the

- 1014 development of the Mesozoic Boreal Realm. *Palaeogeography, Palaeoclimatology,*
1015 *Palaeoecology* 61:237–254. DOI: [https://doi.org/10.1016/0031-0182\(87\)90052-6](https://doi.org/10.1016/0031-0182(87)90052-6).
- 1016 Druckenmiller PS, Hurum J, Knutsen EM, Nakrem HA. 2012. Two new
1017 ophthalmosaurids (Reptilia: Ichthyosauria) from the Agardhfjellet Formation (Upper
1018 Jurassic: Volgian/Tithonian), Svalbard, Norway. *Norwegian Journal of Geology*
1019 92:311–339.
- 1020 Druckenmiller PS, Maxwell EE. 2014. A Middle Jurassic (Bajocian) ophthalmosaurid
1021 (Reptilia, Ichthyosauria) from the Tuxedni Formation, Alaska and the early
1022 diversification of the clade. *Geological Magazine* 151:41–48. DOI:
1023 [doi:10.1017/S0016756813000125](https://doi.org/10.1017/S0016756813000125).
- 1024 Dzyuba OS, Schraer CD, Hults CP, Blodgett RB, Schraer DJ. 2019. Early Bajocian
1025 belemnites of Southcentral Alaska: new data and new perspectives on mid-Middle
1026 Jurassic Megateuthididae and Belemnopseidae biogeography. *Journal of Systematic*
1027 *Palaeontology* 17:911–935. DOI: [10.1080/14772019.2018.1486335](https://doi.org/10.1080/14772019.2018.1486335).
- 1028 Fernández M. 1994. A new long-snouted ichthyosaur from the early Bajocian of Neuquén
1029 basin (Argentina). *Ameghiniana* 31:291–297.
- 1030 Fernández M. 1997. A new ichthyosaur from the Tithonian (Late Jurassic) of the
1031 Neuquén Basin (Argentina). *Journal of Paleontology* 71:479–484.
- 1032 Fernández M. 1999. A new ichthyosaur from the Los Molles Formation (Early Bajocian),
1033 Neuquén basin, Argentina. *Journal of Paleontology* 73:677–681.
- 1034 Fernández M. 2003. Ophthalmosauria (Ichthyosauria) forefin from the Aalenian-Bajocian
1035 boundary of Mendoza Province, Argentina. *Journal of Vertebrate Paleontology*
1036 23:691–694.
- 1037 Fernández MS, Maxwell EE. 2012. The genus *Arthropterygius* Maxwell (Ichthyosauria:
1038 Ophthalmosauridae) in the Late Jurassic of the Neuquén Basin, Argentina. *Geobios*
1039 45:535–540.
- 1040 Fischer V. 2016. Taxonomy of *Platypterygius campylodon* and the diversity of the last
1041 ichthyosaurs. *PeerJ* 4:1–21. DOI: [10.7717/peerj.2604](https://doi.org/10.7717/peerj.2604).
- 1042 Fischer V, Appleby RM, Naish D, Liston J, Riding JB, Brindley S, Godefroit P. 2013. A
1043 basal thunnosaurian from Iraq reveals disparate phylogenetic origins for Cretaceous
1044 ichthyosaurs. *Biology Letters* 9:1–6. DOI: [10.1098/rsbl.2013.0021](https://doi.org/10.1098/rsbl.2013.0021).
- 1045 Fischer V, Arkhangelsky MS, Uspensky GN, Stenshin IM, Godefroit P. 2014a. A new
1046 Lower Cretaceous ichthyosaur from Russia reveals skull shape conservatism within
1047 Ophthalmosaurinae. *Geological Magazine* 151:60–70. DOI:
1048 [10.1017/S0016756812000994](https://doi.org/10.1017/S0016756812000994).
- 1049 Fischer V, Bardet N, Benson RBJ, Arkhangelsky MS, Friedman M. 2016. Extinction of
1050 fish-shaped marine reptiles associated with reduced evolutionary rates and global
1051 environmental volatility. *Nature communications* 7:1–11. DOI:
1052 [10.1038/ncomms10825](https://doi.org/10.1038/ncomms10825).
- 1053 Fischer V, Bardet N, Guimaraes M, Godefroit P. 2014b. High Diversity in Cretaceous
1054 Ichthyosaurs from Europe Prior to Their Extinction. *PLoS ONE* 9:e84709. DOI:
1055 [10.1371/journal.pone.0084709](https://doi.org/10.1371/journal.pone.0084709).
- 1056 Fischer V, Benson RBJ, Druckenmiller PS, Ketchum HF, Bardet N. 2018. The
1057 evolutionary history of polycotyloid plesiosaurians. *Royal Society Open Science* 5.
1058 DOI: [10.1098/rsos.172177](https://doi.org/10.1098/rsos.172177).
- 1059 Fischer V, Benson RBJ, Zverkov NG, Soul LC, Arkhangelsky MS, Lambert O, Stenshin

- IM, Uspensky GN, Druckenmiller PS. 2017. Plasticity and convergence in the evolution of short-necked plesiosaurs. *Current Biology* 27:1667–1676. DOI: 10.1016/j.cub.2017.04.052.
- Fischer V, Cappetta H, Vincent P, Garcia G, Goolaerts S, Martin JE, Roggero D, Valentin X. 2014c. Ichthyosaurs from the French Rhaetian indicate a severe turnover across the Triassic–Jurassic boundary. *Naturwissenschaften* 101:1027–1040. DOI: 10.1007/s00114-014-1242-7.
- Fischer V, Guimomar M, Godefroit P. 2011. New data on the palaeobiogeography of Early Jurassic marine reptiles: the Toarcian ichthyosaur fauna of the Vocontian Basin (SE France). *Neues Jahrbuch für Geologie und Paläontologie* 261:111–127. DOI: 10.1127/0077-7749/2011/0155.
- Fischer V, Maisch MW, Naish D, Kosma R, Liston J, Joger U, Krüger FJ, Pardo Pérez J, Tainsh J, Appleby RM. 2012. New ophthalmosaurid ichthyosaurs from the European Lower Cretaceous demonstrate extensive ichthyosaur survival across the Jurassic–Cretaceous boundary. *PLoS ONE* 7:e29234. DOI: 10.1371/journal.pone.0029234.
- Fischer V, Zverkov NG, Arkhangelsky MS, Stenshin IM, Blagoveschensky IV, Uspensky GN. 2020. A new elasmosaurid plesiosaurian from the Early Cretaceous of Russia marks an early attempt at neck elongation. *Zoological Journal of the Linnean Society*:in press. DOI: 10.1093/zoolinnean/zlaa103.
- Foffa D, Young MT, Brusatte SL. 2018. Filling the Corallian gap : New information on Late Jurassic marine reptile faunas from England. *Acta Palaeontologica Polonica* 63:287–313.
- Foffa D, Young MT, Brusatte SL, Graham MR, Steel L. 2017. A new metriorhynchid crocodylomorph from the Oxford Clay Formation (Middle Jurassic) of England, with implications for the origin and diversification of Geosaurini. *Journal of Systematic Palaeontology* 2019:1–21. DOI: 10.1080/14772019.2017.1367730.
- Foffa D, Young MT, Stubbs TL, Dexter KG, Brusatte SL. 2018. The long-term ecology and evolution of marine reptiles in a Jurassic seaway. *Nature Ecology and Evolution* 2:1548–1555. DOI: 10.1038/s41559-018-0656-6.
- Gasparini Z. 1997. A new pliosaur from the Bajocian of the Neuquen Basin, Argentina. *Palaeontology* 40:135–147.
- Gasparini Z, Fernández MS, de la Fuente MS, Salgado L. 2007. Reptiles marinos Jurásicos y Cretácicos de la Patagonia Argentina: su aporte al conocimiento de la herpetofauna mesozoica. *Ameghiniana* APA Public:125–136.
- Gasparini Z, Pol D, Spalletti LA. 2006. An unusual marine crocodyliform from the Jurassic–Cretaceous boundary of Patagonia. *Science* 311:70–73.
- Gilmore CW. 1902. Discovery of teeth in *Baptanodon*, an ichthyosaurian from the Jurassic of Wyoming. *Science* 16:913–914.
- Gilmore CW. 1906. Notes on osteology of *Baptanodon*. *Memoirs of the Carnegie Museum* II:325–337.
- Godefroit P. 1993a. Les grands ichthyosaures sinémuriens d’Arlon. *Bulletin de l’Institut Royal des Sciences Naturelles de Belgique Sciences de la Terre* 63:25–71.
- Godefroit P. 1993b. The skull of *Stenopterygius longifrons* (Owen, 1881). *Revue de Paléobiologie de Genève volume spécial* 7:67–84.
- Godefroit P. 1994a. Les reptiles marins du Toarcien (Jurassique inférieur) belgo-

- 1106 luxembourgeois. *Mémoires pour servir à l'Explication des Cartes Géologiques et*
1107 *Minières de la Belgique* 39:98.
- 1108 Godefroit P. 1994b. *Simolestes keileni* sp. nov., un Pliosaure (Plesiosauria, Reptilia) du
1109 Bajocien supérieur de Lorraine (France). *Bulletin des Académie et Société Lorraines*
1110 *des sciences* 33:77–95.
- 1111 Großmann F. 2007. The taxonomic and phylogenetic position of the Plesiosauroidea from
1112 the Lower Jurassic Posidonia shale of south-west Germany. *Palaeontology* 50:545–
1113 564.
- 1114 Guérin-Franiatte S, Weis R. 2010. Le passage Aalénien-Bajocien près de Rumelange: la
1115 série biostratigraphique dans le Bassin d'Esch-sur-Alzette (Grand-Duché de
1116 Luxembourg). In: Weis R, Guérin-Franiatte S eds. *Le Jurassique inférieur et moyen*
1117 *au Luxembourg*. Luxembourg ville, Luxembourg: Ferrantia, 73–96.
- 1118 Hauff B. 1953. *Das Holzmadenbuch*. Öhringen.
- 1119 Herrera Y, Gasparini Z, Fernández MS. 2013. A new Patagonian species of Cricosaurus
1120 (Crocodyliformes, Thalattosuchia): First evidence of Cricosaurus in Middle-Upper
1121 Tithonian lithographic limestone from Gondwana. *Palaeontology* 56:663–678. DOI:
1122 10.1111/pala.12010.
- 1123 Home E. 1819. Reasons for giving the name Proteo-Saurus to the fossil skeleton which
1124 has been described. *Philosophical transactions of the Royal Society of London*
1125 109:212–216.
- 1126 Huene F von. 1922. *Die Ichthyosaurier des Lias und ihre Zusammenhänge*. Berlin:
1127 Verlag von Gebrüder Borntraeger.
- 1128 Huene F von. 1931. Neue Studien über Ichthyosaurier aus Holzmaden. *Abhandlungen der*
1129 *Senckenbergischen Naturforschenden Gesellschaft* 42:345–382.
- 1130 Huene F von. 1951. Ein neuer Fund von *Eurhinosaurus longirostris*. *Neues Jahrbuch für*
1131 *Geologie und Paläontologie, Abhandlungen* 93:277–283.
- 1132 Johnson R. 1979. The osteology of the pectoral complex of *Stenopterygius* Jaekel
1133 (Reptilia: Ichthyosauria). *Neues Jahrbuch für Geologie und Paläontologie,*
1134 *Abhandlungen* 159:41–86.
- 1135 Johnson MM, Young MT, Brusatte SL. 2020. Emptying the wastebasket: a historical and
1136 taxonomic revision of the Jurassic crocodylomorph Steneosaurus. *Zoological*
1137 *Journal of the Linnean Society* 189:428–448. DOI: 10.1093/zoolinnean/zlaa027.
- 1138 Johnson MM, Young MT, Brusatte SL, Thuy B, Weis R. 2018. A catalogue of
1139 teleosauroids (Crocodylomorpha: Thalattosuchia) from the Toarcian and Bajocian
1140 (Jurassic) of southern Luxembourg. *Historical Biology* 2963:1–16. DOI:
1141 10.1080/08912963.2018.1427090.
- 1142 Kear BP. 2012. A revision of Australia's Jurassic plesiosaurs. *Palaeontology* 55:1125–
1143 1138. DOI: 10.1111/j.1475-4983.2012.01183.x.
- 1144 Kelley NP, Pyenson ND. 2015. Evolutionary innovation and ecology in marine tetrapods
1145 from the Triassic to the Anthropocene. *Science* 348:aaa3716. DOI:
1146 10.1126/science.aaa3716.
- 1147 Ketchum HF, Benson RBJ. 2011a. A new pliosaurid (Sauropterygia, Plesiosauria) from
1148 the Oxford Clay Formation (Middle Jurassic, Callovian) of England: evidence for a
1149 gracile, longirostrine grade of Early–Middle Jurassic pliosaurids. *Special Papers in*
1150 *Palaeontology* 86:109–129.
- 1151 Ketchum HF, Benson RBJ. 2011b. The cranial anatomy and taxonomy of *Peloneustes*

- 1152 *philarchus* (Sauropterygia, Pliosauridae) from the Peterborough Member (Callovian,
1153 Middle Jurassic) of the United Kingdom. *Palaeontology* 54:639–665. DOI:
1154 10.1111/j.1475-4983.2011.01050.x.
- 1155 Klein N, Schmitz L, Wintrich T, Sander PM. 2020. A new cymbospondylid ichthyosaur
1156 (Ichthyosauria) from the Middle Triassic (Anisian) of the Augusta Mountains,
1157 Nevada, USA. *Journal of Systematic Palaeontology* 18:1167–1191. DOI:
1158 10.1080/14772019.2020.1748132.
- 1159 Knight WC. 1903. Notes on genus *Baptanodon*. *American Journal of Science, Fourth*
1160 *Series* 16:78–81.
- 1161 Knutsen EM, Druckenmiller PS, Hurum JH. 2012a. A new plesiosauroid (Reptilia:
1162 Sauropterygia) from the Agardhfjellet Formation (Middle Volgian) of central
1163 Spitsbergen, Norway. *Norwegian Journal of Geology* 92:213–234.
- 1164 Knutsen EM, Druckenmiller PS, Hurum JH. 2012b. Two new species of long-necked
1165 plesiosaurs (Reptilia: Sauropterygia) from the Upper Jurassic (Middle Volgian)
1166 Agardhfjellet Formation of central Spitsbergen. *Norwegian Journal of Geology*
1167 92:187–212.
- 1168 Knutsen EM, Druckenmiller PS, Hurum JH. 2012c. Redescription and taxonomic
1169 clarification of “*Tricleidus*” svalbardensis based on new material from the
1170 Agardhfjellet Formation (Middle Volgian). *Norwegian Journal of Geology* 92:175–
1171 186.
- 1172 Korte C, Hesselbo SP, Ullmann C V., Dietl G, Ruhl M, Schweigert G, Thibault N. 2015.
1173 Jurassic climate mode governed by ocean gateway. *Nature Communications*
1174 6:10015. DOI: 10.1038/ncomms10015.
- 1175 Lomax DR. 2016. A new leptonektid ichthyosaur from the Lower Jurassic (Hettangian)
1176 of Nottinghamshire, England, UK, and the taxonomic usefulness of the
1177 ichthyosaurian coracoid. *Journal of Systematic Palaeontology*:1–15. DOI:
1178 10.1080/14772019.2016.1183149.
- 1179 Lomax DR, Massare JA. 2015. A new species of *Ichthyosaurus* from the Lower Jurassic
1180 of West Dorset, England, U.K. *Journal of Vertebrate Paleontology*:903260. DOI:
1181 10.1080/02724634.2014.903260.
- 1182 Lomax DR, Massare JA. 2016. Two new species of *Ichthyosaurus* from the lowermost
1183 Jurassic (Hettangian) of Somerset, England. *Papers in Palaeontology*. DOI:
1184 10.1002/SPP2.1065.
- 1185 Lomax DR, Porro LB, Larkin NR. 2019. Descriptive anatomy of the largest known
1186 specimen of *Protoichthyosaurus prostaxalis* (Reptilia: Ichthyosauria) including
1187 computed tomography and digital reconstruction of a three-dimensional skull. *PeerJ*
1188 7:e6112. DOI: 10.7717/peerj.6112.
- 1189 Lomax DR, Sachs S. 2017. On the largest *Ichthyosaurus* : A new specimen of
1190 *Ichthyosaurus somersetensis* containing an embryo. *Acta Palaeontologica Polonica*
1191 62:575–584.
- 1192 Long JA, Cruickshank ARI. 1998. Further Records of Plesiosaurian Reptiles of Jurassic
1193 and Cretaceous Age From Western Australia. *Records of the Western Australian*
1194 *Museum* 19:47.
- 1195 Lucius M. 1945. Die Luxemburger Minetteformation und die jüngeren Eisenerzbildungen
1196 unseres Landes. Beiträge zur Geologie von Luxemburg. *Service Carte géologique*
1197 *du Luxembourg* 4:1–347.

- 1198 Maisch MW. 1998. A new ichthyosaur genus from the Posidonia Shale (Lower Toarcian,
1199 Jurassic) of Holzmaden, SW-Germany with comments on the phylogeny of post-
1200 Triassic ichthyosaurs. *Neues Jahrbuch für Geologie und Paläontologie*,
1201 *Abhandlungen* 209:47–78.
- 1202 Maisch MW. 2008. Revision der Gattung *Stenopterygius* Jaekel, 1904 emend. von
1203 Huene, 1922 (Reptilia: Ichthyosauria) aus dem unteren Jura Westeuropas.
1204 *Palaeodiversity* 1:227–271.
- 1205 Maisch MW. 2010. Phylogeny, systematics, and origin of the Ichthyosauria - the state of
1206 the art. *Palaeodiversity* 3:151–214.
- 1207 Maisch MW, Matzke AT. 2000. The Ichthyosauria. *Stuttgarter Beiträge zur Naturkunde*
1208 *Serie B (Geologie und Paläontologie)* 298:1–159.
- 1209 Maisch MW, Matzke AT. 2001. The cranial osteology of the Middle Triassic ichthyosaur
1210 *Contectopalatus* from Germany. *Palaeontology* 44:1127–1156.
- 1211 Maisch MW, Matzke AT. 2003. The cranial osteology of the ichthyosaur *Leptonectes* cf.
1212 *tenuirostris* from the Lower Jurassic of England. *Journal of Vertebrate Paleontology*
1213 23:116–127. DOI: 10.1671/0272-4634(2003)23[116:TCOOTI]2.0.CO;2.
- 1214 Mariotti N, Santantonio M, Weis R. 2007. Aalenian-Early Bajocian Belemnite
1215 Assemblage From Peri-Mediterranean Tethyan Sediments (Calabria , Southern Italy
1216). *Geologica Romana* 40:1–19.
- 1217 Mariotti N, Weis R, Di Cencio A, Clément A, De Baets K. 2012. New records of early
1218 Middle Jurassic belemnites in the French Subalpine Basin and their
1219 paleobiogeographic significance. *Geobios* 45:99–108. DOI:
1220 <https://doi.org/10.1016/j.geobios.2011.11.012>.
- 1221 Marsh OC. 1895. The Reptilia of the *Baptanodon* Beds. *American Journal of Science*
1222 50:405–406.
- 1223 Martin JE, Fischer V, Vincent P, Suan G. 2012. A longirostrine *Temnodontosaurus*
1224 (Ichthyosauria) with comments on Early Jurassic ichthyosaur niche partitioning and
1225 disparity. *Palaeontology* 55:995–1005. DOI: 10.1111/j.1475-4983.2012.01159.x.
- 1226 Massare JA. 1987. Tooth morphology and prey preference of Mesozoic marine reptiles.
1227 *Journal of Vertebrate Paleontology* 7:121–137.
- 1228 Massare JA. 1988. Swimming capabilities of Mesozoic marine reptiles: implications for
1229 the methods of predation. *Palaeobiology* 14:187–205.
- 1230 Massare JA. 1997. Faunas, behavior, and evolution. In: Callaway JM, Nicholls EL eds.
1231 *Ancient Marine Reptiles*. San Diego: Academic Press, 401–421.
- 1232 Massare JA, Buchholtz EA, Kenney J, Chomat A-M. 2006. Vertebral morphology of
1233 *Ophthalmosaurus natans* (Reptilia: Ichthyosauria) from the Jurassic Sundance
1234 Formation of Wyoming. *Paludicola* 5:242–254.
- 1235 Maubeuge PL. 1972. *Etudes stratigraphiques sur la formation ferrifère de Lorraine et ses*
1236 *morts-terrains*. Metz: Self-edited.
- 1237 Maxwell EE. 2010. Generic reassignment of an ichthyosaur from the Queen Elizabeth
1238 Islands, Northwest Territories, Canada. *Journal of Vertebrate Paleontology* 30:403–
1239 415.
- 1240 Maxwell EE. 2012. New metrics to differentiate species of *Stenopterygius* (Reptilia:
1241 Ichthyosauria) from the Lower Jurassic of southwestern Germany. *Journal of*
1242 *Paleontology* 86:105–115. DOI: 10.1666/11-038.1.
- 1243 Maxwell EE, Caldwell MW, Lamoureux DO. 2012. Tooth histology, attachment, and

- 1244 replacement in the Ichthyopterygia reviewed in an evolutionary context.
- 1245 *Paläontologische Zeitschrift* 86:1–14.
- 1246 Maxwell EE, Cortés D. 2020. A revision of the Early Jurassic ichthyosaur *Hauffiopteryx*
- 1247 (Reptilia: Ichthyosauria), and description of a new species from Southwestern
- 1248 Germany. *Palaeontologia Electronica* 23:1–43. DOI: 10.26879/937.
- 1249 Maxwell EE, Fernández MS, Schoch RR. 2012. First diagnostic marine reptile remains
- 1250 from the Aalenian (Middle Jurassic): a new ichthyosaur from southwestern
- 1251 Germany. *PLoS ONE* 7:e41692. DOI: 10.1371/journal.pone.0041692.
- 1252 Maxwell EE, Vincent P. 2015. Effects of the early Toarcian Oceanic Anoxic Event on
- 1253 ichthyosaur body size and faunal composition in the Southwest German Basin.
- 1254 *Paleobiology*:1–10. DOI: 10.1017/pab.2015.34.
- 1255 Mazin J-M. 1982. Affinités et phylogénie des Ichthyopterygia. *Geobios, mémoire spécial*
- 1256 6:85–98.
- 1257 McGowan C. 1973. The cranial morphology of the Lower Liassic latipinnate
- 1258 ichthyosaurs of England. *Bulletin of the British Museum (Natural History) Geology*
- 1259 24:1–109.
- 1260 McGowan C. 1974. A revision of the longipinnate ichthyosaurs of the Lower Jurassic of
- 1261 England, with description of the new species (Reptilia, Ichthyosauria). *Life Science*
- 1262 *Contributions, Royal Ontario Museum* 97:1–37.
- 1263 McGowan C. 1979. A revision of the Lower Jurassic ichthyosaurs of Germany with
- 1264 descriptions of two new species. *Palaeontographica. Abteilung A. Paläozoologie,*
- 1265 *Stratigraphie* 166:93–135.
- 1266 McGowan C. 1986. A putative ancestor for the swordfish-like ichthyosaur
- 1267 *Eurhinosaurus*. *Nature* 322:454–456.
- 1268 McGowan C. 1994. A new species of *Shastasaurus* (Reptilia: Ichthyosauria) from the
- 1269 Triassic of British Columbia; the most complete exemplar of the genus. *Journal of*
- 1270 *Vertebrate Paleontology* 14:168–179.
- 1271 McGowan C. 1996. Giant ichthyosaurs of the Early Jurassic. *Canadian Journal of Earth*
- 1272 *Sciences* 33:1011–1021.
- 1273 McGowan C. 2003. A new specimen of *Excalibosaurus* from the English Lower Jurassic.
- 1274 *Journal of Vertebrate Paleontology* 23:950–956.
- 1275 McGowan C, Milner AC. 1999. A new Pliensbachian ichthyosaur from Dorset, England.
- 1276 *Palaeontology* 42:761–768.
- 1277 McGowan C, Motani R. 2003. *Part 8. Ichthyopterygia*. München: Verlag Dr. Friedrich
- 1278 Pfeil.
- 1279 Mehl MG. 1912. *Muraenosaurus? Reedii*, Sp. Nov and *Tricleidus? Laramiensis* Knight,
- 1280 American Jurassic plesiosaurs. *The Journal of Geology* 20:344–352.
- 1281 Merriam JC. 1908. Triassic Ichthyosauria with special reference to the American forms.
- 1282 *Memoirs of the University of California* 1:1–154.
- 1283 Moon BC. 2017. A new phylogeny of ichthyosaurs (Reptilia: Diapsida). *Journal of*
- 1284 *Systematic Palaeontology* 17:129–155. DOI: 10.1080/14772019.2017.1394922.
- 1285 Moon BC, Kirton AM. 2018a. Ichthyosaurs of the British Middle and Upper Jurassic Part
- 1286 2 *Brachypterygius, Nannopterygius, Macropterygius* and Taxa invalida.
- 1287 *Monographs of the Palaeontographical Society* 172:85–197. DOI:
- 1288 10.1080/02693445.2018.1468139.
- 1289 Moon BC, Kirton AM. 2018b. Ichthyosaurs of the British Middle and Upper Jurassic Part

- 1290 1, *Ophthalmosaurus*. *Monographs of the Palaeontographical Society* 170:1–84.
- 1291 DOI: 10.1080/02693445.2016.11963958.
- 1292 Moon BC, Stubbs TL. 2020. Early high rates and disparity in the evolution of
- 1293 ichthyosaurs. *Communications Biology* 3:1–8. DOI: 10.1038/s42003-020-0779-6.
- 1294 Motani R. 1997. Temporal and Spatial Distribution of Tooth Implantation in
- 1295 Ichthyosaurs. In: Callaway JM, Nicholls EL eds. *Ancient Marine Reptiles*. San
- 1296 Diego, California: Academic Press, 81–103.
- 1297 Motani R. 1999a. Phylogeny of the Ichthyopterygia. *Journal of Vertebrate Paleontology*
- 1298 19:473–496.
- 1299 Motani R. 1999b. On the evolution and homologies of ichthyosaurian forefins. *Journal of*
- 1300 *Vertebrate Paleontology* 19:28–41.
- 1301 Motani R. 2009. The Evolution of Marine Reptiles. *Evo Edu Outreach* 2:224–235.
- 1302 Neige P, Weis R, Fara E. 2020. Ups and downs of belemnite diversity in the Early
- 1303 Jurassic of Western Tethys. *Palaeontology*:In Press.
- 1304 Noe LF. 1999. The Callovian pliosaurs of the Oxford Clay—evidence and implications for
- 1305 the consumption of marine invertebrates. In: *Secondary Adaptation to Life in*
- 1306 *Water/La réadaptation au milieu aquatique*. 13–17.
- 1307 Noe LF. 2001. A taxonomic and functional study of the Callovian (Middle Jurassic)
- 1308 Pliosauroida (Reptilia, Sauropterygia). University of Derby.
- 1309 O’Keefe FR. 2001a. Ecomorphology of plesiosaur flipper geometry. *Journal of*
- 1310 *Evolutionary Biology* 14:987–991.
- 1311 O’Keefe FR. 2001b. A cladistic analysis and taxonomic revision of the Plesiosauria
- 1312 (Reptilia:Sauropterygia). *Acta Zoologica Fennica* 213:1–63.
- 1313 O’Keefe FR. 2002. The evolution of plesiosaur and pliosaur morphotypes in the
- 1314 Plesiosauria (Reptilia: Sauropterygia). *Palaeobiology* 28:101–112.
- 1315 O’Keefe RF. 2004. Preliminary description and phylogenetic position of a new Plesiosaur
- 1316 (Reptilia : Sauropterygia) from the Toarcian of Holzmaden, Germany. *Journal of*
- 1317 *Paleontology* 78:973–988. DOI: 10.1666/0022-
- 1318 3360(2004)078<0973:pdappo>2.0.co;2.
- 1319 O’Keefe FR, Wahl WJ. 2003. Preliminary report on the osteology and relationships of a
- 1320 new aberrant cryptocleidoid plesiosaur from the Sundance Formation, Wyoming.
- 1321 *Paludicola* 4:48–68.
- 1322 Ősi A, Young MT, Galácz A, Rabi M. 2018. A new large-bodied thalattosuchian
- 1323 crocodyliform from the Lower Jurassic (Toarcian) of Hungary, with further
- 1324 evidence of the mosaic acquisition of marine adaptations in Metriorhynchoidea.
- 1325 *PeerJ* 6:e4668. DOI: 10.7717/peerj.4668.
- 1326 Owen R. 1860. On the orders of fossil and recent Reptilia and their distribution in time.
- 1327 *Report of the British Association for the Advancement of Science* 29:153–166.
- 1328 Owen R. 1865. A monograph of the Fossil Reptilia of the Liassic Formations, part third:
- 1329 *Plesiosaurus, Dimorphodon, and Ichthyosaurus*. *Monograph of the*
- 1330 *Palaeontographical society*:12–130.
- 1331 Paparella I, Maxwell EE, Cipriani A, Roncace S, Caldwell MW. 2017. The first
- 1332 ophthalmosaurid ichthyosaur from the Upper Jurassic of the Umbrian-Marchean
- 1333 Apennines (Marche, Central Italy). *Geological Magazine* 154:837–858. DOI:
- 1334 10.1017/S0016756816000455.
- 1335 Pieńkowski G, Schudack ME, Bosák P, Enay R, Feldman-Olszewska A, Golonka J,

- 1336 Gutowski J, Herngreen GFW, Jordan P, Krobicki M, Lathuiliere B, Leinfelder RR,
1337 Michalík J, Mönnig E, Noe-Nygaard N, Pálffy J, Pint A, Rasser MW, Reisdorf AG,
1338 Schmid DU, Schweigert G, Surlyk F, Wetzel A, Wong TE. 2008. Jurassic. *The*
1339 *Geology of Central Europe Volume 2: Mesozoic and Cenozoic* 2:0. DOI:
1340 10.1144/CEV2P.2.
- 1341 Popov E V, Delsate D, Felten R. 2019. A New Callorhynchid Genus (Holocephali,
1342 Chimaeroidei) from the Early Bajocian of Ottange-Rumelange, on the Luxembourg-
1343 French Border. *Paleontological Research* 23:220–230. DOI: 10.2517/2018PR021.
- 1344 Pyenson ND, Kelley NP, Parham JF. 2014. Marine tetrapod macroevolution: Physical
1345 and biological drivers on 250 million years of invasions and evolution in ocean
1346 ecosystems. *Palaeogeography, Palaeoclimatology, Palaeoecology* 400:1–8. DOI:
1347 10.1016/j.palaeo.2014.02.018.
- 1348 R Core Team. 2016. R: A language and environment for statistical computing.
- 1349 Roberts AJ, Druckenmiller PS, Cordonnier B, Delsett LL, Hurum JH. 2020. A new
1350 plesiosaurian from the Jurassic-Cretaceous transitional interval of the Slottsmøya
1351 Member (Volgian), with insights into the cranial anatomy of cryptoclidids using
1352 computed tomography. *PeerJ* 2020. DOI: 10.7717/peerj.8652.
- 1353 Roberts AJ, Druckenmiller PS, Delsett LL, Hurum JH. 2017. Osteology and relationships
1354 of *Colymbosaurus* Seeley, 1874, based on new material of *C. svalbardensis* from the
1355 Slottsmøya Member, Agardhfjellet Formation of central Spitsbergen. *Journal of*
1356 *Vertebrate Paleontology* 4634:e1278381. DOI: 10.1080/02724634.2017.1278381.
- 1357 Sachs S, Klug C, Kear BP. 2019. Rare evidence of a giant pliosaurid-like plesiosaur from
1358 the Middle Jurassic (lower Bajocian) of Switzerland. *Swiss Journal of*
1359 *Palaeontology* 138:337–342. DOI: 10.1007/s13358-019-00200-9.
- 1360 Sadki D, Weis R, Braun P. 2020. Graphoceratidés (Ammonitina) de l’Aalénien moyen
1361 (Jurassique) de Rumelange-Hutberg (Grand-Duché de Luxembourg). In: Di Cencio
1362 A, Sadki D, Weis R eds. *Paléontologie au Luxembourg (2). Les ammonites de la*
1363 *Minette*. Ferrantia, in press.
- 1364 Sander PM. 2000. Ichthyosauria: their diversity, distribution, and phylogeny.
1365 *Paläontologische Zeitschrift* 74:1–35.
- 1366 Sander PM, Bucher H. 1993. An ichthyosaur from the uppermost Toarcian of southern
1367 France. *Neues Jahrbuch für Geologie und Paläontologie. Monatshefte* 1993:631–
1368 640.
- 1369 Sato T, Wu XC. 2008. A new Jurassic pliosaur from Melville Island, Canadian Arctic
1370 Archipelago. *Canadian Journal of Earth Sciences* 45:303–320. DOI: 10.1139/E08-
1371 003.
- 1372 Schintgen T, Förster A. 2013. Geology and basin structure of the Trier-Luxembourg
1373 Basin – implications for the existence of a buried Rotliegend graben. *Zeitschrift der*
1374 *Deutschen Gesellschaft für Geowissenschaften* 164:615–637. DOI: 10.1127/1860-
1375 1804/2013/0025.
- 1376 Schwermann L, Sander PM. 2011. Osteologie und Phylogenie von *Westphaliasaurus*
1377 *simonsensii*: Ein neuer Plesiosauride (Sauropterygia) aus dem Unteren Jura
1378 (Pliensbachium) von Sommersell (Kreis Höxter), Nordrhein-Westfalen,
1379 Deutschland. *Geologie und Paläontologie in Westfalen* 79:5–56.
- 1380 Sciau J, Crochet J-YY, Mattei J. 1990. Le premier squelette de plésiosaure de France sur
1381 le Causse du Larzac (Toarcien, Jurassique inférieur). *Geobios* 23:111–116. DOI:

- 1382 10.1016/0016-6995(90)80021-7.
- 1383 Seeley HG. 1874a. Note on some of the generic modifications of the plesiosaurian
- 1384 pectoral arch. *Quarterly Journal of the Geological Society* 30:436–449. DOI:
- 1385 10.1144/GSL.JGS.1874.030.01-04.48.
- 1386 Seeley HG. 1874b. On *Muraenosaurus Leedsii*, a Plesiosaurian from the Oxford Clay.
- 1387 Part I. *Quarterly Journal of the Geological Society of London* 30:197–208. DOI:
- 1388 10.1144/GSL.JGS.1874.030.01-04.35.
- 1389 Seeley HG. 1874c. On the pectoral arch and fore limb of *Ophthalmosaurus*, a new
- 1390 ichthyosaurian genus from the Oxford Clay. *Quarterly Journal of the Geological*
- 1391 *Society of London* 30:696–707.
- 1392 Siehl A, Thein J. 1989. Minette-type ironstones. In: *Phanerozoic ironstones*. Geological
- 1393 Society of London, Special Publications, 175–193.
- 1394 Smith AS. 2013. Morphology of the caudal vertebrae in *Rhomaleosaurus zetlandicus* and
- 1395 a review of the evidence for a tail fin in Plesiosauria. *Paludicola* 9:144–158.
- 1396 Smith AS. 2015. Reassessment of “*Plesiosaurus*” *megacephalus* (Sauropterygia:
- 1397 Plesiosauria) from the Triassic-Jurassic boundary, UK. *Palaeontologia Electronica*
- 1398 1820:1–19.
- 1399 Smith AS, Araújo R. 2017. *Thaumatodracon wiedenrothi*, a morphometrically and
- 1400 stratigraphically intermediate new rhomaleosaurid plesiosaurian from the Lower
- 1401 Jurassic (Sinemurian) of Lyme Regis. *Palaeontographica Abteilung A* 308:89–125.
- 1402 DOI: 10.1127/pala/308/2017/89.
- 1403 Smith AS, Benson RBJ. 2014. Osteology of *Rhomaleosaurus thorntoni* (Sauropterygia:
- 1404 Rhomaleosauridae) from the Lower Jurassic (Toarcian) of Northamptonshire,
- 1405 England. *Monographs of the Palaeontographical Society* 168:1–40. DOI:
- 1406 10.1080/02693445.2014.11963953.
- 1407 Smith AS, Dyke GJ. 2008. The skull of the giant predatory pliosaur *Rhomaleosaurus*
- 1408 *cramptoni*: implications for plesiosaur phylogenetics. *Naturwissenschaften* 95:975–
- 1409 980. DOI: 10.1007/s00114-008-0402-z.
- 1410 Smith AS, Vincent P. 2010. A new genus of pliosaur (Reptilia: Sauropterygia) from the
- 1411 Lower Jurassic of Holzmaden, Germany. *Palaeontology* 53:1049–1063. DOI:
- 1412 10.1111/j.1475-4983.2010.00975.x.
- 1413 Storrs GW. 1997. Morphological and taxonomic clarification of the genus *Plesiosaurus*.
- 1414 In: *Ancient marine reptiles*. 145–190.
- 1415 Swinton WE. 1930. On a new species of *Eurhinosaurus* in the British Museum.
- 1416 *Geological Magazine* 67:275–277.
- 1417 Taylor M. 1992. Functional anatomy of the head of the large aquatic predator
- 1418 *Rhomaleosaurus zetlandicus* (Plesiosauria, Reptilia) from the Toarcian (Lower
- 1419 Jurassic) of Yorkshire, England. *Philosophical Transactions of the Royal Society B:*
- 1420 *Biological Sciences* 335:247–280. DOI: 10.1098/rstb.1992.0022.
- 1421 Taylor MA. 1997. Before the Dinosaur: The historical significance of the fossil marine
- 1422 reptiles. In: Callaway JM, Nicholls EL eds. *Ancient Marine Reptiles*. San Diego
- 1423 California: Academic Press, xix–xlvi.
- 1424 Tennant JP, Mannion PD, Upchurch P. 2016. Sea level regulated tetrapod diversity
- 1425 dynamics through the Jurassic/Cretaceous interval. *Nature Communications*
- 1426 7:12737. DOI: 10.1038/ncomms12737.
- 1427 Tennant JP, Mannion PD, Upchurch P, Sutton MD, Price GD. Biotic and environmental

- 1428 dynamics through the Late Jurassic – Early Cretaceous transition : evidence for
- 1429 protracted faunal and ecological turnover. *Biological Reviews*. DOI:
- 1430 10.1111/brv.12255.
- 1431 Teyssen TAL. 1984. Sedimentology of the Minette oolitic ironstones of Luxembourg
- 1432 and Lorraine: a Jurassic subtidal sandwave complex. *Sedimentology* 31:195–2011.
- 1433 Thorne PM, Ruta M, Benton MJ. 2011. Resetting the evolution of marine reptiles at the
- 1434 Triassic-Jurassic boundary. *Proceedings of the National Academy of Sciences of the*
- 1435 *United States of America* 108:8339–8344. DOI: 10.1073/pnas.1018959108.
- 1436 Vincent P. 2011. A re-examination of *Hauffiosaurus zannoni*, a pliosauroid from the
- 1437 Toarcian (Early Jurassic) of Germany. *Journal of Vertebrate Paleontology* 31:340–
- 1438 351. DOI: 10.1080/02724634.2011.550352.
- 1439 Vincent P, Bardet N, Morel N. 2007. An elasmosaurid plesiosaur from the Aalenian
- 1440 (Middle Jurassic) of Western France. *Neues Jahrbuch für Geologie und*
- 1441 *Paläontologie, Abhandlungen* 243:363–370.
- 1442 Vincent P, Martin JE, Fischer V, Suan G, Khalloufi B, Sucheras-Marx B, Lena A,
- 1443 Janneau K, Rousselle B, Rulleau L, Suchéras-Marx B, Léna A, Janneau K,
- 1444 Rousselle B, Rulleau L. 2013. A marine vertebrate fauna from the Toarcian-
- 1445 Aalenian succession of southern Beaujolais, Rhône, France. *Geological Magazine*
- 1446 150:822–834. DOI: doi:10.1017/S0016756812000982.
- 1447 Vincent P, Storrs GW. 2019. *Lindwurmia*, a new genus of Plesiosauria (Reptilia:
- 1448 Sauropterygia) from the earliest Jurassic of Halberstadt, northwest Germany.
- 1449 *Science of Nature* 106. DOI: 10.1007/s00114-018-1600-y.
- 1450 Vincent P, Taquet P, Fischer V, Bardet N, Falconnet J, Godefroit P. 2014. Mary
- 1451 Anning’s legacy to French vertebrate palaeontology. *Geological Magazine* 151:7–
- 1452 20. DOI: 10.1017/S0016756813000861.
- 1453 Vincent P, Weis R, Kronz G, Delsate D. 2017. *Microcleidus melusinae*, a new
- 1454 plesiosaurian (Reptilia, Plesiosauria) from the Toarcian of Luxembourg. *Geological*
- 1455 *Magazine*:In Press. DOI: 10.1017/S0016756817000814.
- 1456 Weis R, Mariotti N. 2007. A belemnite fauna from the Aalenian-Bajocian boundary beds
- 1457 of the Grand Duchy of Luxembourg (NE Paris Basin). *Bollettino della Societa*
- 1458 *Paleontologica Italiana* 46:149–174.
- 1459 Weis R, Mariotti N, Riegraf W. 2012. The belemnite family Holcobelidae (Coleoidea) in
- 1460 the European Jurassic: systematics, biostratigraphy, palaeobiogeography and
- 1461 evolutionary trends. *Palaeodiversity* 5:13–49.
- 1462 Weis R, Sadki D, Mariotti N. 2017. Aalenian-Bajocian belemnites from the Middle and
- 1463 High Atlas, Morocco: taxonomy, biostratigraphy and palaeobiogeographical
- 1464 affinities. *Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen*
- 1465 284:215–240. DOI: 10.1127/njgpa/2017/0659.
- 1466 White TE. 1940. Holotype of *Plesiosaurus longirostris* Blake and classification of the
- 1467 plesiosaurs. *Journal of Paleontology* 14:451–467.
- 1468 Wilberg EW. 2015. A new metriorhynchoid (Crocodylomorpha, Thalattosuchia) from the
- 1469 Middle Jurassic of Oregon and the evolutionary timing of marine adaptations in
- 1470 thalattosuchian crocodylomorphs. *Journal of Vertebrate Paleontology* 902846. DOI:
- 1471 10.1080/02724634.2014.902846.
- 1472 Young MT, de Andrade MB. 2009. What is *Geosaurus*? Redescription of *Geosaurus*
- 1473 *giganteus* (Thalattosuchia: Metriorhynchidae) from the Upper Jurassic of Bayern,

- Germany. *Zoological Journal of the Linnean Society* 157:551–585.
- Young MT, Brandalise de Andrade M, Cornée J-J, Steel L, Foffa D. 2014a. Re-description of a putative Early Cretaceous “teleosaurid” from France, with implications for the survival of metriorhynchids and teleosaurids across the Jurassic-Cretaceous Boundary. *Annales de Paléontologie* 100:165–174. DOI: 10.1016/j.annpal.2014.01.002.
- Young MT, Brusatte SL, de Andrade MB, Desojo JB, Beatty BL, Steel L, Fernández MS, Sakamoto M, Ruiz-Omeñaca JI, Schöch RR. 2012. The cranial osteology and feeding ecology of the metriorhynchid crocodylomorph genera *Dakosaurus* and *Plesiosuchus* from the Late Jurassic of Europe. *PLoS ONE* 7:e44985. DOI: 10.1371/journal.pone.0044985.
- Young MT, Steel L, Brusatte SL, Foffa D, Lepage Y. 2014b. Tooth serration morphologies in the genus *Machimosaurus* (Crocodylomorpha, Thalattosuchia) from the Late Jurassic of Europe. *Royal Society Open Science* 1:1–14. DOI: 10.1098/rsos.140269.
- Zverkov NG, Arkhangel'sky MS, Pardo-Pérez J, Beznosov PA. 2015. On the Upper Jurassic ichthyosaur remains from the Russian North. *Proceedings of the Zoological Institute RAS* 319:81–97.
- Zverkov NG, Efimov VM. 2019. Revision of *Undorosaurus*, a mysterious Late Jurassic ichthyosaur of the Boreal Realm. *Journal of Systematic Palaeontology*:in press. DOI: 10.1080/14772019.2018.1515793.
- Zverkov NG, Fischer V, Madzia D, Benson RBJ. 2018. Increased Pliosaurid Dental Disparity Across the Jurassic – Cretaceous Transition. *Palaeontology* 61:825–846. DOI: 10.1111/pala.12367.
- Zverkov NG, Jacobs ML. 2020. Revision of Nannopterygius (Ichthyosauria: Ophthalmosauridae): reappraisal of the ‘inaccessible’ holotype resolves a taxonomic tangle and reveals an obscure ophthalmosaurid lineage with a wide distribution. *Zoological Journal of the Linnean Society*:1–48. DOI: 10.1093/zoolinnean/zlaa028.
- Zverkov NG, Prilepskaya NE. 2019. A prevalence of *Arthropterygius* (Ichthyosauria: Ophthalmosauridae) in the Late Jurassic—earliest Cretaceous of the Boreal Realm. *PeerJ* 7:e6799. DOI: 10.7717/peerj.6799.

Figure Captions

Figure 1. Number of collections of marine reptiles (Ichthyosauria, Plesiosauria, Thalattosuchia, Pleurosauria, Angolachelonia) per stage. Data extracted from the paleobiology database on the 26th March 2020 (see Acknowledgements for the main contributors of these data). Silhouettes originate from phylopic: *Dakosaurus* by Dmitry Bogdanov & T Michael Keesey; *Meyerasaurus*, *Eurhinosaurus*, *Temnodontosaurus*, *Plesiopterys*, and *Ophthalmosaurus* by Gareth Monger; *Rhomaleosaurus* and *Stenopterygius* by Scott Hartmann; *Peloneustes* by Nobu Tamura & T Michael Keesey; *Albertonectes* by Frank Denota.

Figure 2. Map of the fossiliferous localities investigated. Produced using data from Esri, HERE, Garmin, and OpenStreetMap.

Figure 3. Thoursense Zone fauna, late Toarcian, Luxembourg. Selected anatomy of specimen MNHNL TM212. (A–B) left coracoid in ventral (A) and anteroventral (B) views. (C, E, F) left humerus in anterodorsal (C), distal (E), and proximal (F) views. (D) left forefin in ventral view.

Figure 4. Pseudoradosa Zone fauna, late Toarcian, Luxembourg. (A–E) rhomaleosaurid left humerus MNHNL DOU307 in ventral (A), anterior (B), dorsal (C), proximal (D), and distal (E) views. (F–H) rhomaleosaurid humerus MNHNL KA109 in dorsal/ventral (F), proximal (H), and distal (H) views.

Figure 5. Ichthyosaurians of the late Toarcian of Belgium and Luxembourg. (A–B) large, *Temnodontosaurus*-like caudal centrum IRSNB Vert-00000-00803 in anterior (A) and lateral (B) views. (C–D) probable thunnosaurian caudal centrum IRSNB Vert-00000-00808 in anterior (C) and dorsal (D) views. (E–F) ichthyosaurian distal tooth IRSNB Vert-06462-0003 in labial (E) and mesial (F) views. (G–H) probable non-ophthalmosaurid left angular IRSNB Vert-06462-0002 in lateral (G) and medial (H) views. (I–K) ichthyosaurian right quadrate IRSNB Vert-00000-00802 in lateral (I), medial (J), and posterior (K) views.

Figure 6. Ichthyosaurians of the Aalensis Zone, late Toarcian, Luxembourg. (A–C) left coracoid of a non-baracromian parvipelvian MNHNL DOU353 in ventral (A), lateral (B), and posterior (C) views. (D–E) large parvipelvian caudal centrum MNHNL DOU378 in dorsal (D), anterior (E), and cross-sectional (F) views. (G) small parvipelvian caudal centrum MNHNL DOU998 in anteroventral view. (H–I) parvipelvian dorsal centrum

MNHNL DOU944 in anterior (H), and lateral (I) views. (J) parvipelvian dorsal centrum
MNHNL DOU352 in anterior view.

Figure 7. Plesiosaurian centra of the Aalensis Zone, late Toarcian, Luxembourg. (A–
B) rhomaleosaurid pectoral vertebra MNHNL DOU954 in anterior (A) and ventral (B)
views. (C–G) cf. *Microcleidus* cervical vertebra in anterior (C), ventral (D), dorsal (E),
lateral (F), and oblique (G) views. (H–J) rhomaleosaurid pectoral vertebra MNHNL
DOU723 in anterior (H), lateral (I), and posterior (J) views. (K–N) plesiosaurian sacral
vertebra MNHNL DOU724 in dorsal (K), anterior (L), posterolateral (M), and ventral (N)
views. (O–Q) rhomaleosaurid pectoral vertebra MNHNL DOU722 in anterior (H), lateral
(I), and posterior (J) views.

Figure 8. Plesiosaurians of the Aalensis Zone, late Toarcian, Luxembourg. (A)
plesiosaurian tooth crown MNHNL DOU906 in ?mesial view. (B) rhomaleosaurid right
humerus MNHNL DOU558 in dorsal view. (C–F) rhomaleosaurid right humerus
MNHNL DOU324a in proximal (C), dorsal (D), posterior (E), and distal (F) views. (G)
juvenile rhomaleosaurid propodial MNHNL DOU324b.

Figure 9. Late Aalenian fauna, Luxembourg. (A–D) non-cryptoclidid plesiosauroid
cervical centrum MNHNL BU157 in anterior (A), dorsal (B), ventral (C), and oblique (D)
views. (E–G) plesiosaur juvenile caudal centrum in lateral (E), anterior (F), and
anteroventral (G) views. (H–I) fragmentary ichthyosaurian rostrum.

Figure 10. Humphresianum Zone fauna, early Bajocian, Luxembourg. (A–F)
cryptoclidid propodial MNHNL BM782 in anterior (A), proximal (B), dorsal (C), ventral
(D), posterior (E), and distal (F) views. (G–H) parvipelvian MNHNL BM392 centra (G)
and sclerotic element in lateral view (H). (I–O) ophthalmosaurid surangular and teeth
MNHNL BM780_781: anterior tooth in labial (I) and basal (J) views; mid-rostrum tooth
in labial (K) and basal (L) views; posterior tooth in labial view (M); right angular in
lateral (N) and medial (O) views.

**Figure 11. The new Humphresianum Zone ophthalmosaurid, early Bajocian,
Luxembourg.** Selected anatomy of specimen MNHNL BM779. (A–C) right exoccipital
in anterolateral (A), posteromedial (B), and posterior (C) views. (D–E) suraoccipital in
posterior (D) and ventral (E) views. (F–H) left scapula in anterior (F), medial (G), and
lateral (H) views. (I–K) right quadrate in lateral (I), anterior (J), and condylar (K) views.
(L) right parietal in dorsal view. (M–N) right humerus in posterior (M) and dorsal (N)
views. (O–P) right ulna in posterior (O) and dorsal (P) views.

1586 **Figure 12. Ichthyosaurian centra proportions.** (A) Height over length; the grey lines
 1587 indicate a ratio of 2 and 3. (B) Height-length ratio, emphasizing the reduction of shape
 1588 disparity from the late Toarcian to the Bajocian. The silhouette originate from phylopic:
 1589 *Stenopterygius* by Scott Hartmann.

Figure 1

Number of collections of marine reptiles (Ichthyosauria, Plesiosauria, Thalattosuchia, Pleurosauria, Angolachelonia) per stage.




Figure 2

Map of the fossiliferous localities investigated





Figure 3


Thoursense Zone fauna, late Toarcian, Luxembourg.

Thouarsense Zone, late Toarcian


A Left coracoid, ventral view




B Left coracoid, anteroventral view




C Left humerus, anterodorsal view




E Left humerus, distal view



F Left humerus, proximal view



D Left forefin, ventral view



50 mm

Figure 4

Pseudoradosa Zone fauna, late Toarcian, Luxembourg.



Figure 5

Ichthyosaurians of the Aalensis Zone, late Toarcian of Luxembourg and Belgium.

Dispansum-Pseudoradosa-Aalensis Zone, late Toarcian




Figure 6

Ichthyosaurians of the Aalensis Zone, late Toarcian, Luxembourg.




Figure 7

Plesiosaurian centra of the Aalensis Zone, late Toarcian, Luxembourg.

Aalensis Zone, late Toarcian




Figure 8

Plesiosauroians of the Aalensis Zone, late Toarcian, Luxembourg.




Figure 9

Late Aalenian fauna, Luxembourg.





Figure 10

Humphresianum Zone fauna, early Bajocian, Luxembourg.

Humphriesianum Zone, early Bajocian

Criptoclididae



Parvipelvian




Figure 11

The new Humphresianum Zone ophthalmosaurid, early Bajocian, Luxembourg.

Humphriesianum Zone, early Bajocian



Figure 12

Ichthyosaurian centra proportions.




Table 1(on next page)

Centra dimensions (in millimeters).

1 **Table 1.** Centra measurements in millimeters.

Specimen	Clade	Age	Axial_zone	Height	Length	Width
ULgPA35961	Ichthyosauria	Toarcian	dorsal	78	35	86
IRSNB_Vert-00000-00803	Ichthyosauria	Toarcian	caudal	118	48	118
IRSNB_Vert-00000-00808	Ichthyosauria	Toarcian	caudal	62	22	60
IRSNB_Vert-06462-0005	Ichthyosauria	Toarcian	cervical	63	24	64
IRSNB_Vert-11312-00007	Plesiosauria	Toarcian	dorsal	67	44	81
MNHNL_DOU378	Ichthyosauria	Toarcian	caudal	190	65	NA
MNHNL_DOU978	Plesiosauria	Toarcian	cervical	41	55	53
MNHNL_DOU944	Ichthyosauria	Toarcian	dorsal	82	36	NA
MNHNL_DOU998	Ichthyosauria	Toarcian	sacral	90	31	NA
MNHNL_DOU722	Plesiosauria	Toarcian	dorsal	50	37	60
MNHNL_DOU723	Plesiosauria	Toarcian	dorsal	55	30	63
MNHNL_DOU954	Plesiosauria	Toarcian	pectoral	63	42	73
MNHNL_DOU724	Plesiosauria	Toarcian	sacral	39	40	57
MNHNL_DOU369a	Ichthyosauria	Toarcian	dorsal	68	32	NA
MNHNL_DOU369b	Ichthyosauria	Toarcian	caudal	61	35	NA
MNHNL_DOU369c	Ichthyosauria	Toarcian	caudal	63	32	NA
MNHNL_HU384	Plesiosauria	Aalenian	caudal	32	NA	NA
MNHNL_BU157	Plesiosauria	Aalenian	cervical	67	62	71
TMNHNL_MP212	Ichthyosauria	Toarcian	cervical	62	32	56
MNHNL_BM779	Ichthyosauria	Bajocian	caudal	91	39	NA
MNHNL_BM758	Ichthyosauria	Bajocian	caudal	100	39	NA
MNHNL_BM725a	Ichthyosauria	Bajocian	cervical	83	38	NA
MNHNL_BM725b	Ichthyosauria	Bajocian	caudal	56	26	NA
MNHNL_BM461	Ichthyosauria	Bajocian	cervical	72	30	NA
MNHNL_BM392a	Ichthyosauria	Bajocian	caudal	72	30	NA
MNHNL_BM392b	Ichthyosauria	Bajocian	caudal	77	31	NA
MNHNL_BM392c	Ichthyosauria	Bajocian	caudal	72	29	NA
MNHNL_BM392d	Ichthyosauria	Bajocian	caudal	75	32	NA
MNHNL_BM779	Ichthyosauria	Bajocian	cervical	87	35	90

2

Table 2(on next page)

List of the material and taxonomic identifications.

1 Table 2. List of the material and taxonomic identifications.

Specimen	Locality	Stage	Zone	Element	Clade	Identification	Note
ULgPA359 61	Arlon region	Toarcia n	Dispansum to Aalensis	Dorsal centrum	Ichthyosau ria	Parvipelvia indet.	
IRSNB Vert- 06455- 0001	Mont-Saint- Martin (France)	Toarcia n	Dispansum to Aalensis	Centrum	Ichthyosau ria	Ichthyosauria indet.	
IRSNB Vert- 00000- 00801	Halanzy (Belgiu m)	Toarcia n	Dispansum to Aalensis	Centrum	Ichthyosau ria	Ichthyosauria indet.	
IRSNB Vert- 06659- 0001	NA (Luxem bourg)	Toarcia n	Dispansum to Aalensis	Centrum	Ichthyosau ria	Ichthyosauria indet.	
IRSNB Vert- 06462- 0003	Halanzy (Belgiu m)	Toarcia n	Dispansum to Aalensis	Tooth	Ichthyosau ria	Ichthyosauria indet.	
IRSNB Vert- 00000- 00802	NA	Toarcia n	Dispansum to Aalensis	Right quadrate	Ichthyosau ria	Ichthyosauria indet.	
IRSNB Vert- 00000- 00800	Halanzy (Belgiu m)	Toarcia n	Dispansum to Aalensis	Left surangular	Ichthyosau ria	Parvipelvia indet.	Probable non- ophthalmosaurid
IRSNB Vert- 06462- 0002	Halanzy (Belgiu m)	Toarcia n	Dispansum to Aalensis	Left angular	Ichthyosau ria	Parvipelvia indet.	Probable non- ophthalmosaurid
IRSNB Vert- 00000- 00803	NA (Luxem bourg)	Toarcia n	Dispansum to Aalensis	Caudal centrum	Ichthyosau ria	Parvipelvia indet.	<i>Temnodontosaur us-like</i>
IRSNB Vert- 00000- 00804	NA (Luxem bourg)	Toarcia n	Dispansum to Aalensis	Dorsal centrum	Ichthyosau ria	Parvipelvia indet.	<i>Temnodontosaur us-like</i>
IRSNB Vert- 00000- 00808	NA	Toarcia n	Dispansum to Aalensis	Caudal centrum	Ichthyosau ria	Parvipelvia indet.	Possible thunnosaurian
IRSNB Vert- 06462- 0005	Halanzy (Belgiu m)	Toarcia n	Dispansum to Aalensis	Cervical centrum	Ichthyosau ria	Parvipelvia indet.	

IRSNB Vert- 11312- 00007	Halanz (Belgiu m)	Toarcia n	Dispansum to Aalensis	Dorsal centrum	Plesiosauri a	Plesiosauria indet.	
MNHN TM212	Dudelage	Toarcia n	Thouarsense	Partial skeleton	Ichthyosau ria	Baracromia indet.	Non- ophthalmosaurid
MNHN DOU9 44	NA (Luxem bourg)	Toarcia n	Aalensis	Dorsal centrum	Ichthyosau ria	Parvipelvia indet.	
MNHN DOU9 98	NA (Luxem bourg)	Toarcia n	Aalensis	Caudal centrum	Ichthyosau ria	Parvipelvia indet.	
MNHN DOU3 69	NA (Luxem bourg)	Toarcia n	Aalensis	Centra	Ichthyosau ria	Parvipelvia indet.	Probable non- thunnosaurian
MNHN DOU3 53	NA (Luxem bourg)	Toarcia n	Aalensis	Coracoid	Ichthyosau ria	Parvipelvia indet.	Non- baracromian
MNHN DOU3 78	NA (Luxem bourg)	Toarcia n	Aalensis	Posterior dorsal to anterior caudal centrum	Ichthyosau ria	Parvipelvia indet.	Probable non- thunnosaurian
MNHN HU242	Rumelage	Aalenia n	Concavum	Partial rostrum	Ichthyosau ria	Ichthyosauria indet.	
MNHN BM360	Rumelage	Bajocia n	Humphriesian um	Partial neural spine and ribs	Ichthyosau ria	Ichthyosauria indet.	
MNHN BM758	Rumelage	Bajocia n	Humphriesian um	Caudal centrum	Ichthyosau ria	Parvipelvia indet.	
MNHN BM725	Rumelage	Bajocia n	Humphriesian um	Cervical and caudal centra	Ichthyosau ria	Parvipelvia indet.	
MNHN BM461	Rumelage	Bajocia n	Humphriesian um	Cervical centrum	Ichthyosau ria	Parvipelvia indet.	
MNHN BM392	Rumelage	Bajocia n	Humphriesian um	Five caudal centrum and one sclerotic plate	Ichthyosau ria	Parvipelvia indet.	
MNHN BM780 _781	Rumelage	Bajocia n	Humphriesian um	Surangular, teeth	Ichthyosau ria	Ophthalmosauri dae indet.	
MNHN BM779	Rumelage	Bajocia n	Humphriesian um	Partial skeleton	Ichthyosau ria	Ophthalmosauri dae indet.	
MNHN DOU3 07	Rodange	Toarcia n	Pseudoradosa	Humerus	Plesiosauri a	Rhomaleosaurid ae indet.	

MNHNL KA109	Esch-sur-Alzette	Toarcia n	Pseudoradosa	Humerus	Plesiosauria	Rhomaleosauridae indet.	
MNHNL DOU9 06	Esch-sur-Alzette	Toarcia n	Aalensis	Tooth crown	Plesiosauria	Plesiosauria indet.	
MNHNL DOU7 24	Probably Esch-sur-Alzette	Toarcia n	Aalensis	Sacral vertebra	Plesiosauria	Plesiosauria indet.	
MNHNL DOU7 22	Esch-sur-Alzette	Toarcia n	Aalensis	Pectoral vertebra	Plesiosauria	Rhomaleosauridae indet.	
MNHNL DOU7 23	Esch-sur-Alzette	Toarcia n	Aalensis	Pectoral vertebra	Plesiosauria	Rhomaleosauridae indet.	
MNHNL DOU9 54	NA (Luxembourg)	Toarcia n	Aalensis	Posterior pectoral/anterior dorsal	Plesiosauria	Rhomaleosauridae indet.	
MNHNL DOU5 58	Belvaux	Toarcia n	Aalensis	Humerus	Plesiosauria	Rhomaleosauridae indet.	
MNHNL DOU3 24A	Esch-sur-Alzette	Toarcia n	Aalensis	Humerus	Plesiosauria	Rhomaleosauridae indet.	
MNHNL DOU3 24B	Esch-sur-Alzette	Toarcia n	Aalensis	Humerus	Plesiosauria	Rhomaleosauridae indet.	Juvenile
MNHNL DOU9 78	NA (Luxembourg)	Toarcia n	Aalensis	Cervical centrum	Plesiosauria	cf. Microcleidus	
MNHNL HU384	Esch-sur-Alzette	Aalenia n	Murchisonae	Caudal centrum	Plesiosauria	Plesiosauria indet.	Juvenile
MNHNL BU157	Rumelange	Aalenia n	Concavum or Murchisonae	Cervical vertebra	Plesiosauria	Plesiosauroidea indet.	Non-cryptoclidid
MNHNL BM782	Rumelange	Bajocian	Humphriesianum	Propodial	Plesiosauria	Cryptoclididae indet.	