

# Refining the marine reptile turnover at the Early-Middle Jurassic transition

Valentin Fischer<sup>Corresp., 1</sup>, Robert Weis<sup>2</sup>, Ben Thuy<sup>2</sup>

<sup>1</sup> Evolution & Diversity Dynamics Lab, UR Geology, Université de Liège, Liège, Belgium

<sup>2</sup> Département des sciences de la Terre, Musée national d'Histoire Naturelle, Luxembourg-Ville, Luxembourg

Corresponding Author: Valentin Fischer

Email address: v.fischer@uliege.be

Even though a handful of long-lived reptilian clades dominated Mesozoic marine ecosystems, several biotic turnovers drastically changed the taxonomic composition of these communities. A seemingly slow paced, within-geological period turnover took place across the Early-Middle Jurassic transition. This turnover saw the demise of early neoichthyosaurians, rhomaleosaurid plesiosaurians and early plesiosauroids in favour of ophthalmosaurid ichthyosaurians and cryptoclidid and pliosaurid plesiosaurians, clades that will dominate the Late Jurassic and, for two of them, the entire Early Cretaceous as well. The fossil record of this turnover is however extremely poor, and this change of dominance appears to be spread across the entire middle Toarcian-Bathonian interval. We describe a series of ichthyosaurian and plesiosaurian specimens from successive geological formations in Luxembourg and Belgium that detail the evolution of marine reptile assemblages across the Early-Middle Jurassic transition within a single area, the Belgo-Luxembourgian sub-basin. These fossils reveal the continuing dominance of large rhomaleosaurid plesiosaurians, microcleidid plesiosaurians, and *Temnodontosaurus*-like ichthyosaurians up to the latest Toarcian, indicating that the structuration of the upper tier of Western Europe marine ecosystems remained essentially constant up to the very end of the Early Jurassic. These fossils also suddenly record ophthalmosaurid ichthyosaurians and cryptoclidid plesiosaurians by the early Bajocian. These results from a geographically-restricted area provide a clearer picture of the shape of the marine reptile turnover occurring at the early-Middle Jurassic transition. This event appears restricted to the sole Aalenian stage, reducing the uncertainty of its duration, at least for ichthyosaurians and plesiosaurians, to 4 instead of 14 million years.

# Refining the marine reptile turnover at the Early-Middle Jurassic transition

Valentin Fischer<sup>1</sup>, Robert Weis<sup>2</sup>, Ben Thuy<sup>2</sup>

<sup>1</sup> Evolution & Diversity Dynamics Lab, UR Geology, Université de Liège, Liège, Belgium.  
v.fischer@uliege.be

<sup>2</sup> Département des sciences de la Terre, Musée national d'Histoire Naturelle, Luxembourg-Ville, Luxembourg.

Corresponding Author:

Valentin Fischer

14 Allée du 6 Août, Liège, 4000, Belgium

Email address: [v.fischer@uliege.be](mailto:v.fischer@uliege.be)

## Abstract

Even though a handful of long-lived reptilian clades dominated Mesozoic marine ecosystems, several biotic turnovers drastically changed the taxonomic composition of these communities. A seemingly slow paced, within-geological period turnover took place across the Early-Middle Jurassic transition. This turnover saw the demise of early neoichthyosaurians, rhomaleosaurid plesiosaurians and early plesiosauroids in favour of ophthalmosaurid ichthyosaurians and cryptoclidid and pliosaurid plesiosaurians, clades that will dominate the Late Jurassic and, for two of them, the entire Early Cretaceous as well. The fossil record of this turnover is however extremely poor, and this change of dominance appears to be spread across the entire middle Toarcian-Bathonian interval.

We describe a series of ichthyosaurian and plesiosaurian specimens from successive geological formations in Luxembourg and Belgium that detail the evolution of marine reptile assemblages across the Early-Middle Jurassic transition within a single area, the Belgo-Luxembourgian sub-basin. These fossils reveal the continuing dominance of large rhomaleosaurid plesiosaurians, microcleidid plesiosaurians, and *Temnodontosaurus*-like ichthyosaurians up to the latest Toarcian, indicating that the structuration of the upper tier of Western Europe marine ecosystems remained essentially constant up to the very end of the Early Jurassic. These fossils also suddenly record ophthalmosaurid ichthyosaurians and cryptoclidid plesiosaurians by the early Bajocian. These results from a geographically-restricted area provide a clearer picture of the shape of the marine reptile turnover occurring at the early–Middle Jurassic transition. This event appears restricted to the sole Aalenian stage, reducing the uncertainty of its duration, at least for ichthyosaurians and plesiosaurians, to 4 instead of 14 million years.

# Introduction

A series of diapsid clades dominated marine ecosystems during the entire Mesozoic (e.g. Motani, 2009; Benson, 2013; Pyenson, Kelley & Parham, 2014; Kelley & Pyenson, 2015). This long-term dominance was, however, pulsed by a series of extinctions and turnovers (Massare, 1987; Callaway & Massare, 1989; Bardet, 1994, 1995; Benson et al., 2010; Benson & Druckenmiller, 2014), the most-studied of which being those happening during the latest Triassic (Thorne, Ruta & Benton, 2011; Benson, Evans & Druckenmiller, 2012; Fischer et al., 2014c; Dick & Maxwell, 2015; Moon & Stubbs, 2020) and at the Jurassic-Cretaceous boundary (Tennant et al.; Fischer et al., 2012, 2013; Benson & Druckenmiller, 2014; Young et al., 2014a; Tennant, Mannion & Upchurch, 2016; Zverkov et al., 2018).

The dynamics of marine reptile turnovers that took place outside of geological period boundaries are less well known, despite evident modifications of assemblages, such as during the early Late Cretaceous (Bakker, 1993; Bardet et al., 2008; Fischer et al., 2016, 2018). The composition of marine reptile communities clearly changed across the Early-Middle Jurassic transition (Massare, 1987; Maxwell, Fernández & Schoch, 2012; Vincent et al., 2013). On one hand, early neoichthyosaurian ichthyosaurs and microcleidid plesiosaurs seemingly met their demise and rhomaleosaurid plesiosaurs nearly went extinct (Maxwell, Fernández & Schoch, 2012; Fischer et al., 2013; Benson, Zverkov & Arkhangelsky, 2015). On the other hand, ophthalmosaurid ichthyosaurs, thalattosuchian crocodylomorphs, cryptocleidid plesiosauroids, and pliosaurids radiated, dominating the Late Jurassic and beyond (Motani, 1999a; Cau & Fanti, 2011; Ketchum & Benson, 2011a; Fischer et al., 2013; Benson & Druckenmiller, 2014; Cau, 2014; Moon, 2017; Foffa, Young & Brusatte, 2018; Campos, Fernández & Herrera, 2020).

However, the fossil record of marine reptiles across the middle Toarcian – Callovian interval is extremely poor (Bardet, 1994; Benson et al., 2010; Benson & Butler, 2011; Cau & Fanti, 2011; Maxwell, Fernández & Schoch, 2012) (Figure 1) and no region preserves relevant fossiliferous successions spanning this transition, as a result of a generalized regression reducing the volume of marine epicontinental ecosystems (Bardet, 1994). These biases blur both the tempo and the severity of this turnover. We tackle this issue by describing and analysing the composition of the late Toarcian-Bajocian ichthyosaurs and plesiosaurs of Luxembourg and Belgium (northeastern part of Paris Basin; Figure 2; see Johnson et al. (2018) for an account on thalattosuchians), extending the spatiotemporal extent of several Jurassic clades and precisising the severity of the biotic turnover at the Early–Middle Jurassic transition.

# Material & Methods

## Upper Toarcian of Luxembourg: Thouarsense, Pseudoradosa, and Aalensis zones

Towards the end of the Early Jurassic, the area of present-day Luxembourg was located in the northeast Paris Basin within the northwestern Peri-Tethys Ocean (e.g. Pieńkowski et al., 2008; Schintgen & Förster, 2013). In a shallow, near-coastal sea between the former landmasses of the Rhenish Massif in the north and the Vosges and Black Forest in the south, a thick siliclastic succession was deposited during the late Toarcian, with clayey to silty sediments in the lower parts and silty to sandy, iron-grain rich sediments in the upper part following a general coarsening upwards trend (Siehl & Thein, 1989). The material described herein includes a single, semi-articulated post-cranial ichthyosaur skeleton from the so-called “Couches à *Astarte voltzi*” (Dittrich, 1993), a clay-rich siltstone corresponding to level 11 of Delsate and Weis (2010), dated to the late Toarcian Thouarsense Ammonite Zone of the Grandcourt Formation.

All the other late Toarcian marine reptile remains from Luxembourg described herein were found in the overlying Minette ironstone Formation (Lucius, 1945). Deposition of the Minette successions took place under the influence of strong tidal currents (Teyssen, 1984) during the upper Toarcian Pseudoradosa and Aalensis ammonite Zones (Di Cencio & Weis, 2020). The Aalenian Opalinum ammonite zone has not yet been detected in the area and probably forms a local hiatus (Maubeuge, 1972; Guérin-Franiatte & Weis, 2010).

#### **Upper Toarcian of Belgium: Dispansum-Pseudoradosa-Aalensis zones**

The Mont-Saint-Martin Formation crops out in southern Belgium and includes iron-rich beds forming a slightly diachronous equivalent of the Minette ironstones in Luxembourg. These ironstones contain all the marine reptiles and were deposited in a shallow marine setting during the late Toarcian (Dispansum to Aalensis ammonite zones; Boulvain et al., 2001). One fossil (IRSNB Vert-06455-0001) was found just outside the Belgian border, in the Meurthe-et-Moselle department, north-eastern France (Figure 2).

#### **Middle-upper Aalenian of Luxembourg: Murchisonae and Concavum zones**

The Toarcian part of the Minette ironstones in Luxembourg is locally overlain by a few meters of iron-rich marly sandstones and thin beds with calcareous nodules deposited during the Aalenian Murchisonae to Concavum ammonite zones (Guérin-Franiatte & Weis, 2010; Sadki, Weis & Braun, 2020). These deposits form the onset of a transgressive succession with increasingly finer sedimentation leading to the so-called “marnes micacées”, a succession of poorly lithified silty claystones with thin layers of phosphatic nodules, dated to the lowermost Bajocian Discites ammonite zone (Guérin-Franiatte & Weis, 2010).

#### **Lower Bajocian of Luxembourg: Humphriesianum Zone**

The youngest marine strata in Luxembourg correspond to a succession of calcarenites, marl-limestone alternations and coral limestones deposited on the northeast margin of the Burgundy carbonate platform at the end of the lower Bajocian. The marine reptile remains described herein were all recovered from the so-called “Marnes sableuses d’Audun-le-Tiche”, dated to the lower

Bajocian Humphriesianum ammonite zone (Delsate et al., 2018; Popov, Delsate & Felten, 2019). An indeterminate thalattosuchian has been reported from these strata in Luxembourg (Johnson et al., 2018).

### Marine reptile collections data

We extracted all the fossil collections of Jurassic (Hettangian-Tithonian) marine reptiles (selected clades: Ichthyosauria, Plesiosauria, Thalattosuchia, Pleuroosauria, and Angolachelonia) from the paleobiology database (<https://paleobiodb.org>) on the 26th March 2020. While the taxonomic record of Jurassic marine reptiles on the paleobiology database is not complete, the record we extracted samples the vast majority of the fossil localities of the Jurassic. This record is used as a proxy for sampling intensity for visualization purposes only; we binned these data per geological stage and generate a plot of the number of collections over time in R v3.6.2 (R Core Team, 2016), using the geoscale v2.0 package (Bell, 2015) (Figure 1; see supplementary files for the dataset and the R script).

### Institutional abbreviations

**IRSNB**: Royal Belgium Institute of Natural Sciences, Brussels, Belgium. **MNHNL**: Muséum national d'histoire naturelle du Luxembourg, Luxembourg-ville, Luxembourg. **ULg-PA**: Collections de paléontologie animale de l'Université de Liège, Liège, Belgium.

## Comparative Descriptions

### Thouarsense zone fauna, upper Toarcian, Luxembourg

ICHTHYOSAURIA De Blainville, 1835  
PARVIPELVIA Motani, 1999  
THUNNOSAURIA Motani, 1999  
BARACROMIA Fischer et al., 2013  
Baracromia indet.

A single specimen (MNHNL TM212) is recorded from the Thouarsense zone: it is a partial articulated ichthyosaur comprising fifty-two centra, several ribs and gastralia, a partial coracoid, and two partial forefins (Figure 3; Table 1; Table 2), from the Dudelage locality. All the preserved pre-apical centra are markedly rounded and anteroposteriorly short, bearing close similarities with Jurassic thunnosaurians such as *Stenopterygius* and *Ophthalmosaurus* (Buchholtz, 2001; Massare et al., 2006). Posterior dorsal centra maintain clear bicipital rib articulations and all ribs have a 8-shaped cross-section, which also suggest that this specimen is a thunnosaurian (Sander, 2000; V.F. pers. obs. on NHMUK 2003, R5465, R498), even though this feature is likely an ambiguous synapomorphy, being recorded in Early–Middle Triassic specimens as well (Schmitz, 2005; A.J. Roberts, pers. comm. October 2020). The posterior

apical centra are strongly compressed laterally, suggesting the presence of a lunate tailfin; although the angle of the tail is unknown. The coracoid (Figure 3 A-B) possesses a wide anterior notch, unlike in leptonektids (Huene, 1951; Maisch & Matzke, 2000; Lomax, 2016; Lomax, Evans & Carpenter, 2018). The presence of a posterolateral emargination, typical of *Ichthyosaurus*, *Protoichthyosaurus*, and *Temnodontosaurus* (Home, 1819; Godefroit, 1993a; Lomax & Massare, 2015; Lomax, Porro & Larkin, 2019) cannot be assessed because the coracoid is incomplete (Figure 3A). The humerus (Figure 3C–F) is incompletely-preserved, lacking the capitulum. The dorsal trochanter is proximodistally short, as it does not extend to mid-shaft, while the deltopectoral crest is elongated and parallel to the long axis of the humerus. The small dorsal trochanter suggests that this taxon is not an ophthalmosaurid (Motani, 1999a). The humerus forms a prominent anterior process that possesses a conspicuous, anterodorsally facing facet. This facet exhibits a pitted bone texture which suggests the presence of a cartilaginous cap in vivo. While many Early Jurassic ichthyosaurs possess a mesiodistally-thick anterodistal expansion of the humerus (*Temnodontosaurus*, Leptonektidae, *Hauffiopteryx*, *Stenopterygius* (Johnson, 1979; McGowan & Milner, 1999; McGowan, 2003; McGowan & Motani, 2003; Maisch, 2008; Caine & Benton, 2011; Martin et al., 2012), this expansion forms a flat facet but not for articulation with an anterior accessory epipodial element. The rounded and notched anterior surface of the radius also confirms the absence of such an element. This combination of features rules out earliest neoichthyosaurs and ophthalmosaurids (e.g. Maisch & Matzke, 2000; McGowan & Motani, 2003) and has only been reported in the poorly known, late Toarcian-lower Bajocian taxon *Dearcmhara shawcrossi* (Brusatte et al., 2015), although the facet is smaller and deeper in *D. shawcrossi*. Epipodial and proximal elements closely resemble those of *Stenopterygius* spp. and *Chacaicosaurus cayi* (Johnson, 1979; Fernández, 1994; Maxwell, Fernández & Schoch, 2012), being polygonal and dorsoventrally thick, with no spatium interosseum. The radius and the radiale are notched as in many early neoichthyosaurs (e.g. Huene, 1922; McGowan, 1974; Maxwell, 2012; Lomax & Massare, 2016). The intermedium forms two distal facets unequal in size (Figure 3D): a large, distally facing facet for articulation with distal carpal 3 and a smaller, posterodistally facing facet for articulation with distal carpal 4 (the ‘latipinnate’ condition, although the latipinnate-longipinnate dichotomy fails to capture what is essentially a trait with continuous spectrum; see Mazin (1982) and Motani (1999b) for discussions). The specimen MNHNL TM212 is regarded as a non-ophthalmosaurid baracromian, resembling *Stenopterygius* spp., *Chacaicosaurus cayi*, and *Dearcmhara shawcrossi*.

# **Pseudoradosa zone fauna, upper Toarcian, Luxembourg**

PLESIOSAURIA De Blainville, 1835

Rhomaleosauridae Nopcsa, 1928

Rhomaleosauridae indet.

Two large plesiosaurian propodials have been recovered from Pseudoradosa zone: one (MNHNL DOU307) from the “Couche grise” of the Rodange locality, and one (MNHNL KA109) from the “Couche noire” of the Esch-sur-Alzette locality (Table 2).

MNHNL DOU307 is a complete left propodial (proximal-distal length: 390mm; distal width: 210mm) with a weak posterior and dorsal curvature (Figure 4A–E). The shaft is as wide as the capitulum, and there is no anterior expansion indicating that this propodial is an early plesiosaurian humerus (Storrs, 1997; Bardet, Godefroit & Sciau, 1999; O’Keefe, 2004; Smith & Vincent, 2010; Benson, Evans & Druckenmiller, 2012; Vincent & Storrs, 2019). The dorsal tuberosity is as anteroposteriorly wide as the capitulum and is weakly demarcated. A bulge with muscle scars is present on the ventral surface of the shaft, 30 mm distally to the capitulum. A very short preaxial flange is present distally, forming a small, anterodistally facing triangular facet. The radial/tibial facet is flat and marks an angle of ca. 45° with the ulnar/fibular facet. This facet is convex and rugose, indicating the presence of an extensive cartilage layer. A larger postaxial flange is present, forming distally a semioval facet that is hardly discernible from the ulnar/fibular facet. There is no distal ridge, unlike in the cryptoclidid *Colymbosaurus megadeirus* (Benson & Bowdler, 2014; Roberts et al., 2017). The absence of a marked curvature (the anterior surface is straight in dorsal and ventral views) and the presence of a conspicuous postaxial flange suggest rhomaleosaurid affinities (Smith & Vincent, 2010; Smith & Benson, 2014; Smith, 2015) (with the exception of *Lindwurmia thiuda* Vincent & Storrs, 2019), which are reinforced by the large size of the propodial.

MNHNL KA109 is the distal end of a plesiosaurian propodial; the anterior-posterior asymmetry suggests it is also a humerus (Figure 4F–H). It resembles MNHNL DOU307 in lacking a preaxial flange and the postaxial flange is smaller than in MNHNL DOU307. The distal surface is markedly rounded in dorsoventral view. A prominent, strongly pitted anteroposterior ridge textures of the distal surface, obliterating radial/tibial and ulnar/fibular facets. The postaxial flange forms a triangular and concave facet posterior to the distal ridge. An anteroposteriorly elongated distal ridge is regarded as an autapomorphic feature of *Colymbosaurus megadeirus* (Benson & Bowdler, 2014; Roberts et al., 2017; Arkhangelsky et al., 2019) but the lack of an extensive postaxial flange precludes such a referral; the size of the humerus and its similarities with MNHNL DOU307 suggest rhomaleosaurid affinities.

# **Dispansum-Pseudoradosa-Aalensis zones, upper Toarcian, Belgium**

ICHTHYOSAURIA De Blainville, 1835

Ichthyosauria indet.

IRSNB Vert-06455-0001, IRSNB Vert-00000-00801, and IRSNB Vert-06659-0001 are ichthyosaurian centra, heavily eroded and damaged. Because their original shapes cannot be asserted, we refrained from calculating their shape ratios and assign them to Ichthyosauria indet (Table 2)..

IRSNB R 438 (Figure 5E–F) is a distal tooth that preserves the crown and the acellular cementum ring. The crown is recurved and robust (height: 14.83 mm, basal diameter: 8.14 mm, giving a crown shape ratio of 1.82), as is usually the case for distal ichthyosaurian teeth (Motani, 1997; McGowan & Motani, 2003). Apicobasal ridges texture the crown, unlike in leptoneotids (Maisch, 1998; Maisch & Matzke, 2003; McGowan & Motani, 2003; Lomax, 2016) and the acellular cementum is itself ridged. This combination of features match that of *Ichthyosaurus*, *Protoichthyosaurus*, and *Temnodontosaurus* (McGowan, 1973; Godefroit, 1993a; Vincent et al., 2014; Brusatte et al., 2015; Lomax, Porro & Larkin, 2019). But it is also found in more primitive taxa such as *Contectopalatus* and *Cymbospondylus* (Maisch & Matzke, 2001; Klein et al., 2020). As a result, this specimen is referred to as Ichthyosauria indet. (Table 2).

IRSNB R 440 (Figure 5I–K) is a partial ichthyosaurian right quadrate, preserving the condyle, the stapedial facet and the posteroventral part of the anterior lamella. The condyle is massive, oval, but appears moderately short anteroposteriorly. The medial and lateral surfaces are pitted by numerous small foramina; the stapedial facet is semi-circular thickened ventral and posterior edges. This quadrate has the same size (maximal condylar width: 71 mm) and morphology as that of *Temnodontosaurus platyodon* (Godefroit, 1993a). In the absence of additional evidence, this quadrate is referred to Ichthyosauria indet. (Table 2) but indicates the presence of large ichthyosaurs by the latest Toarcian.

# PARVIPELVIA Motani, 1999

## Parvipelvia indet.

ULg-PA 35.961 is an ichthyosaurian anterior dorsal centrum with a damaged dorsal surface. Its height-length ratio is 78 mm/35 mm = 2.23 (Table 1). The centrum is rounded as in parvipelvians (Merriam, 1908; Maisch & Matzke, 2000) and the diapophyses and parapophyses strongly protrude from the body of the centrum, as often seen in neoichthyosaurs (e.g. Owen, 1865; Huene, 1922).

IRSNB Vert-00000-00800 is a partial left ichthyosaurian surangular, probably associated with the angular IRSNB R 439. It consists of posterior third, where the angular in a mesiolaterally-flattened bone with a slightly convex lateral surface and a slightly concave medial surface for accommodation of the Meckelian canal. The dorsal surface of the surangular is saddle-shaped, forming a long and shallow concavity anterior to the coronoid process. The preserved



morphology of IRSNB Vert-00000-00800 is not diagnostic, but we assign it to *Parvipelvina* indet. because of its probable association with the angular IRSNB R 439 (see below).

IRSNB R 439 (Figure 5G–H) is a partial left ichthyosaurian angular, bearing the typical double-grooved dorsal surface. The lateral wall of the angular is dorsoventrally short anteriorly, and only increases in size posterior to the level of the coronoid process. This narrow lateral exposure of the angular suggests that this specimen does not belong to *Ophthalmosauridae*, as this clade is notably characterized by large exposure of the angular in lateral view (Motani, 1999a).

IRSNB R 436 (Figure 5A–B) and IRSNB Vert-00000-00804 are large ichthyosaurian centra. IRSNB R 436 is 118 mm high and 48 mm long, giving a height-length ratio of 2.45 (Table 1).; IRSNB Vert-00000-00804 is damaged on the edges and will not be measured but is of comparable size. The lateral surfaces of these centra are rounded in anteroposterior view, as in parvipelvians (Merriam, 1908; Maisch & Matzke, 2000), but their size departs from that of Early Jurassic thunnosaurians, even particularly large individuals (e.g. Maxwell, 2012; Lomax & Sachs, 2017). These centra rather match the size and shape of the dorsal-caudal centra of some large non-thunnosaurian neoichthyosaurians such as *Temnodontosaurus* spp. (Godefroit, 1993a; Martin et al., 2012) and *Excalibosaurus costini* (McGowan, 2003). IRSNB R 436 possesses a single apophysis (the parapophysis) placed laterally, suggesting it is middle preflexural caudal (Huene, 1922; McGowan & Motani, 2003). IRSNB Vert-00000-00804 possesses a diapophysis and a parapophysis placed ventrolaterally and is regarded as a posterior dorsal centrum. The size and shape of IRSNB R 436 and IRSNB Vert-00000-00804 match those of large early neoichthyosaurians and but are regarded as *Parvipelvina* indet. in the absence of unambiguous synapomorphies (Table 2).

IRSNB R 437 (Figure 5C–D) is an ichthyosaurian anterior caudal centrum. Its height-length ratio is  $65 \text{ mm} / 22 \text{ mm} = 2.94$  (Table 1).; this high shape ratio close to 3 might suggest thunnosaurian affinities (Buchholtz, 2001). The lateral surface of the centrum is rounded as in parvipelvians (Merriam, 1908; Maisch & Matzke, 2000) and slightly oval, as the ventral surface is flattened. The parapophyses are located lateroventrally.

IRSNB Vert-06462-0005 is an ichthyosaurian cervical centrum. Its height-length ratio is  $63 \text{ mm} / 24 \text{ mm} = 2.62$ . The lateral surface of the centrum is rounded as in parvipelvians (Merriam, 1908; Maisch & Matzke, 2000). A faint ventral keel is present, giving the centrum a pentagonal shape. Such a shape is often seen in the cervical/anterior dorsals of parvipelvians (e.g. Huene, 1922; Fischer et al., 2014b; Lomax, Porro & Larkin, 2019) and is not of diagnostic value at a lower taxonomic level at the current state of our knowledge.

PLESIOSAURIA De Blainville, 1835

Plesiosauria indet.

IRSNB Vert-11312-00007 is a plesiosaurian dorsal centrum, as evidenced by the absence of transverse processes. Its height-length ratio is 67 mm / 44 mm = 1.52. The centrum is poorly preserved and we assign it to Plesiosauria indet. (Table 2).

# **Aalensis zone fauna, upper Toarcian, Luxembourg**

The material from the Aalensis zone of the Minette ironstones is more abundant. We identified 12 ichthyosaurians and 8 plesiosaurians in the MNHL collections. Most of the material consist of isolated centra, complemented by three plesiosaurian propodials, one ichthyosaur coracoid, and one plesiosaurian tooth crown.

ICHTHYOSAURIA De Blainville, 1835

PARVIPELVIA Motani, 1999

Parvipelvia indet.

MNHNL DOU944 (Figure 6H–I) is an ichthyosaurian anterior dorsal centrum from an unknown locality. A central keel is not present but the centrum forms a ventral expansion, giving the centrum a slight pentagonal shape. The diapophysis is small and rounded, as in parvipelvians and unlike in more primitive forms (Merriam, 1908; Huene, 1922) and is strongly protruding and has a rounded distal surface. Shallow dorsoventral ridges connect to the base of the diapophysis, anterior and posteriorly. The parapophysis is flatter and less prominent and its anterior margin merges with the anterior edge of the centrum. Such a pentagonal shape is often seen in parvipelvians (e.g. Huene, 1922; Fischer et al., 2014b; Lomax, Porro & Larkin, 2019) and is not of diagnostic value at a lower taxonomic level at the current state of our knowledge.

MNHNL DOU998 (Figure 6G) is an anterior caudal centrum from an unknown locality. The centrum has a rounded outline, as in usually parvipelvic ichthyosaurians (Merriam, 1908; e.g. Maisch & Matzke, 2000; McGowan & Motani, 2003). Its height/length ratio is 90/31=2.9. This high ratio close to 3 might suggest thunnosaurian affinities (Buchholtz, 2001), but in the absence of additional data, this specimen is regarded as Parvipelvia indet. (Tables 1–2).

MNHNL DOU369 is a series of disarticulated ichthyosaur centra. One originates from the mid-thoracic region and two others from the anterior caudal region, their respective height/length ratios are as follows: 68/32=2.1, 61/35=1.74, and 63/32=1.96. The mid-thoracic resembles MNHNL DOU944. The rather elongated anterior caudal suggest these centra do not belong to a thunnosaurian, but rather to an early parvipelvic (Buchholtz, 2001).

MNHNL DOU353 (Figure 6A–C) is a fragmentary left coracoid from an unknown locality. It preserves the lateral part and a fragment of the intercoracoid facet. The ventral surface is moderately saddle-shaped (intercoracoid facet is only 56mm thick while the mesiolateral length is 180mm). The glenoid facet is large (anteroposterior length=75mm) and oval, while the scapular facet forms a small triangle (anteroposterior length= 39mm) and forms a ca. 60° angle with the glenoid facet. The posterolateral emargination is wide and deep: the coracoid surface posterior to the glenoid faces posteriorly and not posterolaterally, similar to the condition seen in *Ichthyosaurus communis*, *Ichthyosaurus anningae*, *Protoichthyosaurus postaxialis*, and many but not all specimens of *Temnodontosaurus platyodon* (Home, 1819; Godefroit, 1993a; Lomax & Massare, 2015; Lomax, Porro & Larkin, 2019). The Triassic shastasaurid *Shatasaurus neoscapularis* also possesses a wide posterior notch, but the coracoid of that taxon is different, having a very wide anterior notch (McGowan, 1994). We regard this coracoid as belonging to a non-baracromian parvipelvian.

MNHNL DOU378 (Figure 6D–F) is a very large centrum from an unknown locality. It originates from the posterior dorsal – sacral – anterior caudal region (length: 65mm and diameter: 175mm, estimated height: 180-185mm, H/L ratio: 2.79-2.84; Table 1). Its size departs from the Early Jurassic thunnosaurians (*Ichthyosaurus* and *Stenopterygius*, which have much smaller centra even particularly large individuals (e.g. Maxwell, 2012; Lomax & Sachs, 2017)). It rather matches the size and shape of the anteriormost caudal centra of some large non-thunnosaurian neoichthyosaurians such as *Temnodontosaurus* spp. (Godefroit, 1993a; Martin et al., 2012) and *Excalibosaurus costini* (McGowan, 2003) (the anterior caudals are much longer in *Eurhinosaurus longirostris* [V.F., pers. obs. on MNHN 1946-20]).

# PLESIOSAURIA De Blainville, 1835

## Plesiosauria indet.

MNHNL DOU906 (Figure 7A) is a plesiosaurian tooth crown from the Esch-sur-Alzette locality. It is slender (length/basal diameter= 23/8= 2.875), labiolingually flattened and recurved indicating it does not belong to derived rhomaleosaurids and thalassophoneans (Owen, 1865; Smith & Vincent, 2010; Ketchum & Benson, 2011b; Smith & Benson, 2014). The crown bears fine striations along the entire surface.

MNHNL DOU724 (Figure 8K–N) is a plesiosaurian sacral vertebra probably originating from the Esch-sur-Alzette locality. The articular surface is strongly dorsoventral depressed and the centrum is as wide as long. As in MNHNL DOU722, MNHNL DOU723, and MNHNL DOU954, the subcentral foramina are widely spaced and are separated by a well-rounded/convex

surface. Similar to the other centra, the dorsal edge of the articular surface is notched by the neural canal. The transverse process is elongated, stout, and points posterolaterally. The rib facet is 8-shaped (waisted), this time with a larger, ear-shaped dorsal part and a smaller, semioval ventral part. The neural canal is also 8-shaped, resembling a ‘plesiosauroid’ pectoral vertebra from the Aalenian-Bajocian of Australia (Kear, 2012). Despite having similar size to the vertebrae we referred to as Rhomaleosauridae indet., none of the feature we highlight above are recorded in *Rhomaleosaurus* (Smith, 2013; Smith & Benson, 2014) and their representation in other derived rhomaleosaurids is unclear. Accordingly, we regard this specimen as Plesiosauria indet.

#### Rhomaleosauridae Nopcsa, 1928

##### Rhomaleosauridae indet.

MNHNL DOU722 (Figure 8O–Q) and MNHNL DOU723 (Figure 8H–J) are two plesiosaurian pectoral vertebrae from the Esch-sur-Alzette locality. The articular surface of the centrum is oval, with a dorsal notch corresponding to the neural canal. The subcentral foramina are displaced laterally, being located on the ventrolateral surface. The ventral surface in between the subcentral foramina is clearly transversely convex. The rib facets are oval, connect to the neural arch, and their long axis is vertical; they extend ventrally below the level of the notochordal pit. The semi-circular prezygophyses face dorsomedially and are clearly separated from one another; their ventral margin is set below the centre of the neural canal, even though the neural canal is large. Anteriorly, the neural spine forms a basal triangular cavity containing a paired ridge. The height of this trough is about 1/3 of the total height of the neural spine. On the posterior surface, the basal region of the neural spines forms a deep tear-shaped concavity that extends for about one half of the total height of the neural spine, and appears similar to the condition of Microcleididae (Bardet, Godefroit & Sciau, 1999; Schwermann & Sander, 2011) and *Rhomaleosaurus* (Smith & Benson, 2014). However, the neural spines are much shorter than in microclidids (e.g. Owen, 1865; Bardet, Godefroit & Sciau, 1999). The neural spine possesses a posterior ridge while its anterior surface is concave, as in *Rhomaleosaurus thortoni* (Smith & Benson, 2014). The neural spine thickens dorsally and its dorsal half is posteriorly inclined, unlike in *Westphaliasaurus simonensis* (Schwermann & Sander, 2011). We assign these vertebrae to Rhomaleosauridae indet. (Table 2).

MNHNL DOU954 (Figure 8A–B) is a posterior pectoral/anterior dorsal vertebra from an unknown locality. It is essentially similar to MNHNL DOU722 and MNHNL DOU723. The main difference is the existence of ventral expansion of the centrum, even if a clear ventral keel remains absent. This ventral expansion gives the articular surface of the centrum a heart shape. The neural spine is concave anteriorly and bears a median ridges similar to the condition of *Rhomaleosaurus thortoni* (Smith & Benson, 2014). We assign this vertebra to Rhomaleosauridae indet. (Table 2).

MNHNL DOU558 (Figure 7B) from the Belvaux locality and MNHNL DOU324a (Figure 7C–F) and MNHNL DOU324b (broken, juvenile; Figure 7G) from the Esch-sur-Alzette locality are three plesiosaurian humeri. These strongly resemble MNHNL DOU307: the shaft is long and slightly curved posteriorly and dorsally, the dorsal tuberosity is wide, a boss with muscle scars is present ventrally to the capitulum, the preaxial lamella is very short and forms a small triangular distal facet, the postaxial lamella is longer and forms a semioval posterodistal facet, radial/tibial and ulnar/fibular facets form a 45–55° angle and no distal ridge is present. The only difference is the position of the dorsal tuberosity, which is offset ventrally, creating a flattened, dorsomedially-facing surface between the dorsal part of the capitulum and the ventral part of the dorsal tuberosity. The larger propodial (MNHNL DOU324A) measures 331mm in proximal distal length and 168mm in maximal distal width. The smaller ‘adult’ propodial (MNHNL DOU558) measures 309mm in proximodistal length and 150mm in distal width.

Microcleididae Benson et al. 2012

*Microcleidus* Watson, 1909

cf. *Microcleidus*

MNHNL DOU978 (Figure 8C–G) is a plesiosaurian cervical centrum from an unknown locality. The centrum is slightly wider than high and is markedly elongated (height=42mm, width=53mm, length=55mm) unlike in derived rhomaleosaurids and thalassophoneans, which have anteroposteriorly short cervical centra (Owen, 1865; Smith & Vincent, 2010; Ketchum & Benson, 2011b; Smith & Benson, 2014). The ventral surface is flattened as in *Microcleidus* (Owen, 1865; Vincent et al., 2017) and bears two subcentral foramina. There is no ventral keel, unlike in rhomaleosaurids (O’Keefe, 2001a; Smith & Dyke, 2008; Benson & Druckenmiller, 2014). The lateral surface bears a (very) faint anteroposterior ridge; a more conspicuous lateral ridge is present in *Microcleidus* spp. (Owen, 1865; Bardet, Godefroit & Sciau, 1999; Vincent et al., 2017). The neural arch is fused with the centrum. The orientation of bone fibers on the right side suggests that the suture is V-shaped, which is mainly seen in rhomaleosaurids (Benson, Evans & Druckenmiller, 2012) and in the early pliosaurid *Hauffiosaurus* (Benson et al., 2011). The neural canal is very small in diameter, resulting in zygapophyses that are located close to the centrum, as in *Microcleidus* (Owen, 1865; Bardet, Godefroit & Sciau, 1999) and unlike in *Plesiosaurus dolichodeirus* (Owen, 1865), *Westphaliasaurus simonensis* (Schwermann & Sander, 2011), *Hauffiosaurus* spp. (White, 1940; Benson et al., 2011) The prezygapophyses are clearly separated, medially-inclined and their ventral surface is located more ventrally than the centre of the neural canal. The rib facets is 8-shaped, being formed of two semi-ovals separated by an anterior and a posterior furrows, as in *Plesiosaurus dolichodeirus* (Owen, 1865; Storrs, 1997) and unlike in many microcleidids (Owen, 1865; Schwermann & Sander, 2011), *Hauffiosaurus* (White, 1940), and cryptoclidids (Andrews, 1910a; Brown, 1981; Knutsen, Druckenmiller & Hurum, 2012a,b). The rib facet in posterior cervical centra is separated by a

groove *Microcleidus melusinae* (Vincent et al., 2017). This centrum thus bears strong similarities with *Microcleidus* and is regarded as cf. *Microcleidus* (Table 2).

# **Upper Aalenian fauna**

## ICHTHYOSAURIA De Blainville, 1835

### Ichthyosauria indet.

MNHNL HU242 (Concavum zone; Figure 9H-I) is a small partial ichthyosaur rostrum from the Rumelange locality. It contains the anterior part of the left dentary (or right premaxilla), showing the labial and lingual walls. Two teeth are preserved in situ. The root and the acellular cementum ring have a circular cross-section and are perfectly smooth externally, unlike in *Ichthyosaurus*, *Protoichthyosaurus*, and *Temnodontosaurus* (McGowan, 1973; Godefroit, 1993a; Vincent et al., 2014; Brusatte et al., 2015; Lomax, Porro & Larkin, 2019) and there is no evidence for plicidentine (unlike other Jurassic ichthyosaurians for which this feature is known (Maxwell, Caldwell & Lamoureux, 2012)), but this latter feature cannot be used for a taxonomic purpose at the present state of our knowledge. The teeth are of a maximal diameter of 8mm and estimated 20mm apicobasal length; the size and shape of the tooth fits within the range of early baracromians (Godefroit, 1993b, 1994a; Fernández, 1994; Maisch, 2008; Maxwell, Fernández & Schoch, 2012) but in the lack of additional evidence, this material is here referred to Ichthyosauria indet. (Table 2).

## PLESIOSAURIA De Blainville, 1835

### Plesiosauria indet.

MNHNL HU384 (Murchisonae zone; Figure 9E–G) is a small caudal centrum of a plesiosaurian of a juvenile individual, originating from the Tétange locality. Its height/width ratio is 32/36mm = 0.88. Two small triangular chevron facets are present ventrally, contacting the edge of the articular surface. The ventral surface is flat. A rounded rib facet is present on the lateral surface. The neural arc is disconnected from the centrum and absent; the facets appear diamond shaped. Caudal vertebrae usually have less diagnostic features than other vertebrae in plesiosaurians, as could be inferred from their respective number of phylogenetic characters (Benson & Druckenmiller, 2014); this, coupled to the immaturity of the specimen would make an assignment ambiguous and we refer this material to Plesiosauria indet. (Table 2).

## PLESIOSAUROIDEA Gray, 1825

### Plesiosauroidea indet.

MNHNL BU157 (Concavum or Murchisonae zone; Figure 9A–D) is a moderately elongated plesiosaurian cervical vertebra from the Rumelange locality. The shape of the centrum suggests it does not belong to derived rhomaleosaurids and thalassophoneans, which have anteroposteriorly short centra (Owen, 1865; Smith & Vincent, 2010; Ketchum & Benson, 2011b; Smith & Benson, 2014). The ventral surface is flattened and bears two subcentral foramina. No ventral keel is present. The edge of the centrum is rugose, as sometimes seen in various plesiosaurian clades (Owen, 1865; Seeley, 1874a; Fischer et al., 2020). An anteroposteriorly elongated bulge (rather than a ridge) is present on the dorsolateral surface. This bulge is separated from the rib facet by a median concave area. The rib facet is roughly oval but is very strongly waisted by an anterior and a posterior notch, giving the facet a marked 8 shape; this structure recalls – but is more marked than in – the late Toarcian centra MNHNL DOU978. The neural arch is fully fused to the centrum and no suture is discernible. A pair of supracentral foramina is present on the floor of foramen magnum. While MNHNL DOU978 could clearly be attributed to *Microcleidus*, this vertebra – although superficially similar – is less elongated, possesses a lateral bulge rather than a ridge, and does not preserve the zygapophyses. As a result, a referral to Microclididae is too ambiguous. Nevertheless, the presence of an 8-shaped rib facet precludes a referral to cryptoclidids (Andrews, 1910a; Brown, 1981; Knutsen, Druckenmiller & Hurum, 2012a,b). Accordingly, this specimen is referred to as a non-cryptoclidid plesiosauroid (Table 2).

# **Lower Bajocian fauna**

## ICHTHYOSAURIA De Blainville, 1835

### Ichthyosauria indet.

MNHNL BM360 from the Rumelange locality contains one partial neural spine and two proximal parts of bicipital ribs. The neural spine bears a dorsoventrally-oriented median ridge on its anterior surface. The taxonomic values of these features is low and we assign this material to Ichthyosauria indet. (Table 2).

## PARVPELVIA Motani, 1999

### Parvipelvia indet.

MNHNL BM758 is an ichthyosaurian caudal centrum from the Rumelange locality. Its shape is rounded, with a slightly flattened ventral surface, as in Parvipelvia (Merriam, 1908; Maisch & Matzke, 2000). Its height/length ratio is  $100\text{mm}/38\text{mm}=2.63$ .

MNHNL BM725 from the Rumelange locality contains a cervical and a caudal ichthyosaurian centra. The cervical centrum has a height/length ratio of  $83\text{mm}/38\text{mm}=2.18$  while the same ratio for the caudal centrum is  $56\text{mm}/26\text{mm}=2.15$ . The centra are rounded in shape and exhibit small

and rounded apophyses, as in parvipelvians and unlike in more primitive forms (Merriam, 1908; Huene, 1922).

MNHNL BM461 is an ichthyosaurian cervical centrum from the Rumelange locality. Its height/length ratio is  $72\text{mm}/30\text{mm}=2.4$ . The centrum is rounded in shape and exhibits small and rounded apophyses, as in parvipelvians and unlike in more primitive forms (Merriam, 1908; Huene, 1922).

MNHNL BM392 (Figure 10G–H) from the Rumelange locality contains five caudal centra and one sclerotic plate of a single ichthyosaurian specimen. The height/length ratios of the first four caudal centra are as follows:  $72\text{mm}/30\text{mm}=2.4$ ,  $77\text{mm}/31\text{mm}=2.48$ ,  $72\text{mm}/29\text{mm}=2.48$ ,  $75\text{mm}/32\text{mm}=2.34$  (Table 1). Their rounded shape indicates they belong to Parvipelvia (Merriam, 1908; Huene, 1922; Maisch & Matzke, 2000) (Table 2). The sclerotic plate bears radiating striations on its lateral surface and its internal and external edges are crenulated, as is usually the case in ichthyosaurians (Andrews, 1910a; McGowan, 1973; Fischer et al., 2014a). The external third of the plate is deflected, facing dorsally.

#### OPHTHALMOSAURIDAE Baur, 1897

##### Ophthalmosauridae indet.

MNHNL BM780–BM781 (Figure 10I–O) is a nearly complete right ichthyosaurian surangular (MNHNL BM780), associated with seven teeth (bearing the collection number MNHNL BM781), originating from the Rumelange locality. The surangular is straight in all planes (only its posterior quarter is slightly deflected medially) and bears a thickened dorsal margin, giving it a tear-shaped cross-section. A lateral, anteroposteriorly elongated and posteriorly-deepening concavity is present on the lateral surface ('surangular fossa'). This depressed area terminates anteriorly to the level of the coronoid process. The coronoid process is prominent and bears a rugose texture. A small, anteroposteriorly oriented ridge is present directly posteromedially to the coronoid process and is likely part of the muscle attachment. The dorsal margin of the surangular forms a concave plateau posteriorly to the coronoid process. A prominent M.a.m.e process is present directly posteromedially to the plateau; this process points dorsomedially. The angular facet extends anteriorly up to the level of the coronoid process and covers the ventral half of the surangular posteriorly. This indicates the presence of an extensive angular, which is an ophthalmosaurid synapomorphy (int. al. Motani, 1999a; Druckenmiller et al., 2012; Fischer et al., 2012, 2014a; Zverkov & Prilepskaya, 2019). The teeth are peculiar in forming externally-visible plicidentine, texturing the acellular cementum ring and the root by very deep apicobasal grooves, as in *Ichthyosaurus*, *Protoichthyosaurus*, and *Temnodontosaurus* (McGowan, 1973; Godefroit, 1993a; Vincent et al., 2014; Brusatte et al., 2015; Lomax, Porro & Larkin, 2019). The root cross-section is oval as in all parvipelvians bar most platypterygiines (Fischer et al., 2012; Fischer, 2016) and the base of the enamel is easy to discern, which is usually the case in



ophthalmosaurids (Fischer et al., 2016). Accordingly, we refer this specimen as Ophthalmosauridae indet. (Table 2).

MNHNL BM779 (Figure 11) is a fragmentary partially articulated ichthyosaurian, containing parts of the skull, axial skeleton, and scapular girdle, originating from the Rumelange locality. It belongs to Ophthalmosauridae, having a humerus with plate-like dorsal trochanter, a massive deltopectoral crest, a humeral facet for an anterior accessory element, a posterodistally deflected ulnar facet and a conspicuous acromial process on the scapula (Motani, 1999a; Fischer et al., 2012, 2013; Moon, 2017; Zverkov & Efimov, 2019) (Table 2). It resembles *Arthropterygius* in having an anteroposteriorly-short parietal symphysis, a deep ulnar facet on the humerus, a small humeral facet for an anterior accessory epipodial elements (Maxwell, 2010; Fernández & Maxwell, 2012; Zverkov & Prilepskaya, 2019). However, the presence of several autapomorphies on the cranial and appendicular elements indicates that this specimen constitutes a novel taxon, for which a dedicated manuscript is in preparation.

PLESIOSAURIA De Blainville, 1835  
CRYPTOCLIDIDAE Williston, 1925  
Cryptocleididae indet.

MNHNL BM782 (Figure 10A–F) is a left plesiosaurian propodial from the Rumelange locality. The propodial is straight in the proximodistal direction and slightly curves dorsally. The shaft is fairly elongated. The dorsal tuberosity is wide and weakly demarcated from the capitulum; they form together an evenly rounded proximal surface. A median dorsal boss surrounded by muscle scars is present close to the distal end of the dorsal tuberosity. This boss is median (along the axis of the shaft) and thus differs from the anteriorly placed boss seen in the humeri of rhomaleosaurids and microcleidids (Owen, 1865; Smith & Benson, 2014). The propodial markedly expands distally, forming a preaxial flange and a longer postaxial flange. The preaxial flange forms a dorsoventrally-narrow triangular facet distally. The postaxial flange forms a semioval posterior facet that forms a marked angle (ca. 45°) with the ulnar/fibular. Both the radial/tibial and the anteroposteriorly- and dorsoventrally-shorter ulnar/fibular facets are slightly convex. The slightly longer radial facet forms a ca. 120° angle with the ulnar facet. However, a distal ridge is absent, unlike in *Colymbosaurus* (Benson & Bowdler, 2014; Arkhangelsky et al., 2019). The shape of the propodial with its fairly slender shaft and large preaxial and postaxial flanges indicate cryptocleidid affinities (Andrews, 1910a; Mehl, 1912; Brown, 1981; O’Keefe & Wahl, 2003; Knutsen, Druckenmiller & Hurum, 2012c; Roberts et al., 2017, 2020) (Table 2), notably resembling the humerus of *Muraenosaurus leedsi* (Seeley, 1874b; Andrews, 1910a) and the femur of *Tricleidus seeleyi* (Owen, 1865).

# Discussion

## The Early–Middle Jurassic transition in marine reptiles, a state of the art

The very well-sampled (Benson et al., 2010) Lower Jurassic marine ecosystems of western Europe housed a vast menagerie of neoichthyosaurians, thalattosuchians, and plesiosaurians (Owen, 1860; Huene, 1922, 1931; McGowan, 1974, 1979; Benton & Taylor, 1984; Maisch & Matzke, 2000; O’Keefe, 2004; Großmann, 2007; Maisch, 2008, 2010; Benson, Evans & Druckenmiller, 2012; Martin et al., 2012; Bardet et al., 2014; Lomax & Massare, 2016; Johnson, Young & Brusatte, 2020). Evidence from body size, craniodental shape, and swimming capabilities suggests that these taxa occupied several niches within shallow marine ecosystems (Hauff, 1953; Massare, 1997, 1987, 1988; Böttcher, 1989; Godefroit, 1994a; McGowan, 1996; Buchholtz, 2001; Buchy, 2010; Fischer, Guimar & Godefroit, 2011; Martin et al., 2012; Dick, Schweigert & Maxwell, 2016; Maxwell & Cortés, 2020). Such a diversity of forms made the Early Jurassic marine ecosystems of western Europe an iconic representation of Mesozoic marine life (Taylor, 1997).

Even though a drastic reduction of apparent diversity is expected following the lagerstätten effect of the early Toarcian localities (Benson et al., 2010), the Middle Jurassic assemblages, when well sampled (i.e. not before the Callovian), are markedly distinct from their Toarcian counterparts. For ichthyosaurians, the dominant and diversified early neoichthyosaurians (leptonectids, *Suevoleiathan*, *Hauffiopteryx*, and *Temnodontosaurus*) are gone; the disappearance of non-thunnosaurian ichthyosaurians marks the end of large (>6m) ichthyosaurian top predators (Massare, 1987), at least up until the Aptian, when some derived platypterygiines will presumably fill similar roles (Fischer et al., 2014b, 2016; Bardet, Fischer & Machalski, 2016; Fischer, 2016). Ichthyosaurians will never re-evolve carinated teeth like those seen in *Temnodontosaurus platyodon* and *Temnodontosaurus trigonodon* (Conybeare, 1822; Massare, 1987; Godefroit, 1993a), nor the sawfish-like morphology of Leptonectidae (Sinton, 1930; Huene, 1951; McGowan, 1986, 2003; Lomax, 2016). From the Callovian onwards, ichthyosaur assemblages overwhelmingly consist of ophthalmosaurids (but see Fischer et al. (2013)), up to the final extinction of ichthyosaurians (Fischer et al., 2016). While diverse ophthalmosaurid assemblages are known by the Kimmeridgian (Valenciennes, 1861; Druckenmiller et al., 2012; Arkhangelsky & Zverkov, 2014; Zverkov et al., 2015; Paparella et al., 2017; Moon & Kirton, 2018a; Delsett et al., 2019; Zverkov & Efimov, 2019; Barrientos-Lara & Alvarado-Ortega, 2020; Campos, Fernández & Herrera, 2020; Zverkov & Jacobs, 2020), only the closely related *Ophthalmosaurus* and *Baptanodon* as well as indeterminate but compatible forms are known in the Callovian-Oxfordian (Seeley, 1874c; Marsh, 1895; Gilmore, 1902, 1906; Knight, 1903; Andrews, 1910b; Arkhangelsky, 1999; Fernández & Iturralde-Vinent, 2000; Massare et al., 2006, 2014; Moon & Kirton, 2018b; Arkhangelsky et al., 2018; Otero et al., 2020).

This leaves a fairly long interval (late Toarcian-latest Callovian) for which the ichthyosaur fossil record is very scarce and geographically dispersed: Sander & Bucher (1993) reported a large cf. *Stenopterygius* from the late Toarcian of southern France; Maisch & Matzke (2000: p72) mention the presence of *Temnodontosaurus* in the late Toarcian of southern Germany; Vincent et al. (2013) reported *Temnodontosaurus* sp., and *Stenopterygius*-like forms in the late Toarcian of southern France and a coracoid bearing many similarities with MNHN DOU353 from the early Aalenian of southern France; finally, Brusatte et al. (2015) described *Dearcmhara shawcrossi* from the late Toarcian-Bajocian of Scotland. The only diagnosable ichthyosaur remains unambiguously known from Aalenian deposits is the holotype of *Stenopterygius aalensis* from Germany (Maxwell, Fernández & Schoch, 2012) (see also Arnaud et al. (1976) and Maxwell et al. (2012) for additional indeterminate specimens). Fragmentary ophthalmosaurid specimens have been reported in the Aalenian-Bajocian boundary of Argentina (Fernández, 2003) and in the early Bajocian of Argentina (Cabrera, 1939; Fernández, 1999; Gasparini et al., 2007; Fernández & Talevi, 2014) and Canada (Druckenmiller & Maxwell, 2014). A fragmentary rostrum from the early Bajocian of Argentina has been regarded as a representative of *Stenopterygius* (Cabrera, 1939) or as a *Temnodontosaurus/Suevoleviathan*-like form because of relatively long dental roots (Fernández & Talevi, 2014). However, this material lacks diagnostic features and is currently referred to as Ichthyosauria indet. (Fernández & Talevi, 2014). It can be inferred from these occurrences that a replacement of early neoichthyosaurians by ophthalmosaurids took place across the Early–Middle Jurassic transition, but the generalized absence of Aalenian-Bathonian ichthyosaur fossils worldwide and Early Jurassic fossils in South America make it difficult to identify a precise turnover.

While less abundant than ichthyosaurians in early Toarcian Lagerstätten (Hauff, 1953), plesiosaurians already had also evolved a vast array of morphotypes by the Toarcian (O’Keefe, 2001b,a; Benson, Evans & Druckenmiller, 2012; Smith & Araújo, 2017), presumably filling as many ecological niches: long-necked and small headed microclidids (Owen, 1865; Bardet, Godefroit & Sciau, 1999), gigantic, apex predatory rhomaleosaurids (Taylor, 1992; Cruickshank, 1994; Smith & Dyke, 2008; Smith & Vincent, 2010; Smith & Benson, 2014), and small-sized, moderately long-necked, and long-snout early pliosaurids (Benson et al., 2011; Ketchum & Benson, 2011a; Vincent, 2011; Fischer et al., 2017). Microcleidids supposedly go extinct after the Toarcian; their last definite record in the latest Toarcian of France (Sciau, Crochet & Mattei, 1990; Bardet, Godefroit & Sciau, 1999). However, an Aalenian specimen from France regarded as an indeterminate elasmosaurid by Vincent et al. (2007) possesses microcleidid features (8-shaped cervical rib facet, lateral ridge on cervical centra) and might possibly be regarded as a member of that clade. Nevertheless, the long-necked morphotype is then colonised by cryptoclydids by the Callovian-Oxfordian (Andrews, 1909; Brown, 1981).

Only a couple of rhomaleosaurid lineages will survive up to the Middle/Late Jurassic, in high latitudes (Gasparini, 1997; Sato & Wu, 2008; Benson, Zverkov & Arkhangelsky, 2015).

Thalassophonean pliosaurids concomitantly radiated and became the main marine apex predators from the Middle Jurassic to the early Late Cretaceous (Noe, 1999, 2001; Ketchum & Benson, 2011a; Benson et al., 2013; Benson & Druckenmiller, 2014; Fischer et al., 2017; Zverkov et al., 2018). A narrative where thalassophoneans filled (either competitively or passively) the niches previously occupied by *Temnodontosaurus* and derived rhomaleosaurids is sensible, but these animals have clearly distinct body plans and probably hunted differently (Massare, 1988; Buchholtz, 2001; O’Keefe, 2001a, 2002). In any case, there is evidence for the continuous presence of gigantic apex predatory plesiosaurians in Europe and elsewhere: *Simolestes keileni* from the upper Bajocian of eastern France (described by Godefroit (1994b), and currently under revision by S. Sachs), an indeterminate fragmentary jaw from the lower Bajocian of Switzerland (Sachs, Klug & Kear, 2019) and ‘pliosauroid’ tooth from the Aalenian-Bajocian of Australia (Long & Cruickshank, 1998; Kear, 2012). Whether these specimens are rhomaleosaurids or thalassophoneans obviously yields different outcomes as they might push the origin of apex predatory pliosaurids to the Bajocian, but these fossils nevertheless suggest that the niche of very large apex predators was not vacated for a long period, if at all.

Recent evidence revealed a series of basal metriorhynchoid crocodyliforms in the late Toarcian–Aalenian interval (Wilberg, 2015; Ösi et al., 2018; Aiglstorfer, Havlik & Herrera, 2020) suggesting an intense diversification across the Early-Middle Jurassic transition (Aiglstorfer, Havlik & Herrera, 2020), associated with a radiation into the pelagic realm (Cau & Fanti, 2011; Cau, 2014; Wilberg, Turner & Brochu, 2019). The effect of this radiation is evident by the late Middle and Late Jurassic, where abundant pelagic thalattosuchian taxa are known, occupying several ecological niches (Gasparini, Pol & Spalletti, 2006; Young & de Andrade, 2009; Young et al., 2012, 2014b; Herrera, Gasparini & Fernández, 2013; Foffa et al., 2017, 2018). Teleosauroids do not appear much affected by this transition however (Wilberg, Turner & Brochu, 2019), as many lineages and ecomorphotypes crossed this interval unscathed (Johnson, Young & Brusatte, 2020).

## **An abrupt turnover**

In light of the short review above, it appears clear that fossiliferous successions in a geographically-restricted area are crucial to decipher both the tempo and the severity of the marine reptile turnover at the early–Middle Jurassic transition. The late Toarcian–Bajocian successions of Luxembourg and Belgium we analysed above yields novel biostratigraphic information that precise the turnover dynamics of ichthyosaurians and plesiosaurians during the Early–Middle Jurassic transition. In a global, broad-brush view of marine reptile macroevolution, the presence of rhomaleosaurids, microcleidids, and non-thunnosaurian ichthyosaurians in the late Toarcian of Luxembourg (Table 2) was expected, as these clades are already known to survive after the early Toarcian Lagerstätten (Bardet, Godefroit & Sciau, 1999; Vincent et al., 2013; Benson, Zverkov & Arkhangelsky, 2015). With the exception of possible remains from the

Aalenian of France (Vincent, Bardet & Morel, 2007; see discussion above), MNHNL DOU978 and the holotype of *Microcleidus tournemirensis* (Sciau, Crochet & Mattei, 1990; Bardet, Godefroit & Sciau, 1999) are the youngest-known microcleidid to date and indicate that the clade extended at least up to the very end of the Early Jurassic. However, recording microcleidids, large rhomaleosaurids, *Temnodontosaurus*-like forms, and more derived ichthyosaurs within the same basin that formed the early Toarcian Lagerstätten yields more important palaeobiogeographic implications. These fossils indicate that the main marine reptile clades of the Early Jurassic remained abundant and dominant in mid-latitude, epicontinental seas up to the end of the Toarcian at least. Although much less prolific than the underlying strata, the upper Toarcian fossil record suggest that no major turnover took place within the Toarcian as the structuration of the upper tier of marine ecosystems remained largely unchanged. The taxonomic diversity crash that follows the early Toarcian appears to be mainly a Lagerstätten effect, confirming previous suspicions (Benson et al., 2010; Maxwell & Vincent, 2015).

In stark contrast with the sense of continuity displayed by the upper Toarcian occurrences, the Bajocian fossil record of the Luxembourg area suddenly records the dominant clades of the Late Jurassic: cryptoclidids, ophthalmosaurids, thalattosuchians (Johnson et al., 2018), and probable pliosaurids (Godefroit, 1994b), with no evidence – so far – for the presence of more ancient clades. The specimen MNHNL BM782 is oldest-known cryptoclidid and MNHNL BM770 and MNHNL BM780–BM781 are two of the oldest ophthalmosaurids known, after a single partial forefin from the Aalenian-Bajocian boundary of Argentina (Fernández, 2003) and fragmentary basicranium from an equivalent of the Sauzei Zone in Canada (Druckenmiller & Maxwell, 2014), which directly underlies the Humphresianum Zone. The new occurrences we report here indicate that ophthalmosaurids rapidly dispersed, being almost simultaneously recorded in Canada, Argentina, and Luxembourg by the early Bajocian.

The Belgo-Luxembourgian marine reptile record provides a clearer picture of the marine reptile turnover occurring at the Early–Middle Jurassic transition, indicating that this replacement is restricted to the sole Aalenian stage instead of possibly spanning the entire middle Toarcian–Bathonian interval. This reduces the uncertainty on the timing and duration of this turnover to 4 instead of 14 million years and packs a series of extinctions and diversifications within a short period of abrupt climate cooling and changes in oceanic currents (Korte et al., 2015). This Aalenian shift from a Toarcian Warm Mode to Aalenian-Bajocian Cool Mode (Korte et al., 2015) also appears associated with a marked faunal disruption in belemnites, with random (i.e. non-morphologically-selective) extinctions leading to a distinct drop in belemnite biodiversity at least in the northwestern Peri-Tethys Ocean (Dera, Toumoulin & de Baets, 2016; Neige, Weis & Fara, 2020). This major disruption in the evolutionary history of Jurassic belemnites ended at the Aalenian-Bajocian boundary and resulted in a radiation of the suborder Belemnopseina that partially replaced the previously dominant Belemnitina in the Western Tethys (Weis, Mariotti & Riegraf, 2012; Weis, Sadki & Mariotti, 2017), furthermore entailing a distinct Boreal vs. Tethyan

belemnite provincialism (Doyle, 1987; Mariotti, Santantonio & Weis, 2007; Weis & Mariotti, 2007; Mariotti et al., 2012; Dzyuba et al., 2019). At the present state of knowledge, we can only speculate about whether faunal changes in belemnites and marine reptiles were independently impacted by the same factors, or even causally connected, since belemnites are an essential component of the food spectrum in some marine reptiles (Massare, 1987; Böttcher, 1989; Dick, Schweigert & Maxwell, 2016), their faunal change could have triggered a disruption of the trophic chain up to the giant top predators.

## Conclusions

A generalised turnover affected marine reptile communities across the Early–Middle Jurassic transition. However, the extremely poor fossil record of the middle Toarcian-Bathonian interval leaves a ca. 14 million years window of the uncertainty for this important event. Our thorough analysis of the ichthyosaurian and plesiosaurian record of the late Toarcian to Bajocian successions in a confined palaeogeographic setting (the Belgo-Luxembourgian sub-basin) indicates that:

- (i) the structuration of the upper tier of marine ecosystems of Western Europe remained unchanged up to the very end of the Early Jurassic, with the presence of large rhomaleosaurid plesiosaurians, microcroclidid plesiosaurians, as well as *Temnodontosaurus*-like and baracromian ichthyosaurians in the late Toarcian.
- (ii) the dominant clades of the Late Jurassic, cryptoclidid plesiosaurians and ophthalmosaurid ichthyosaurians, arose and dispersed earlier than expected, being now recorded in the early Bajocian of Luxembourg.
- (iii) the marine reptile turnover of the Early–Middle Jurassic transition is more abrupt than previously supposed, being restricted to the sole Aalenian stage, i.e. 4 million years in the Belgo-Luxembourgian sub-basin.

## Acknowledgements

We thank the curatorial staff of the IRSNB for their help. We also thank the main contributors of the Paleobiology Database data we used: Prof. R.B.J. Benson, Dr. M.T. Carrano, Dr. J.P. Tennant, Dr. H.P. Street, Prof. R.J. Butler, Prof. M. Clapham and Prof. M. Uhen. We warmly thank We warmly thank Dr. M. Trotta for her help in creating the geographic map, reviewers Prof. M. Fernández, Dr. A. Roberts, and Dr. D. Foffa, and editor Dr. M.T. Young for their insightful comments and constructive suggestions.

## References

- 838 Aiglstorfer M, Havlik P, Herrera Y. 2020. The first metriorhynchoid crocodyliform from the  
839 Aalenian ( Middle Jurassic ) of Germany , with implications for the evolution of  
840 Metriorhynchoidea. *Zoological Journal of the Linnean Society* 188:522–551.
- 841 Andrews CW. 1909. On some new Plesiosauria from the Oxford Clay of Peterborough. *The*  
842 *Annals and magazine of natural history; zoology, botany, and geology* 4:418–429. DOI:  
843 10.1080/00222930908692691.
- 844 Andrews CW. 1910a. *A descriptive catalogue of the marine reptiles of the Oxford clay. Based on*  
845 *the Leeds Collection in the British Museum (Natural History), London. Part I.* London, :  
846 Order of the Trustees of the British Museum.
- 847 Andrews CW. 1910b. Note on the osteology of *Ophthalmosaurus icenicus* Seeley an  
848 ichthyosaurian Reptile from the Oxford Clay of Peterborough. *Geological Magazine* 4:202–  
849 208.
- 850 Arkhangel'sky MS. 1999. On a new ichthyosaur from the Callovian stage of the Volga region  
851 near Saratov. *Paleontological Journal* 33:88–90.
- 852 Arkhangel'sky MS, Zverkov NG. 2014. On a new ichthyosaur of the genus *Undorosaurus*.  
853 *Proceedings of the Zoological Institute RAS* 318:187–196.
- 854 Arkhangel'sky MS, Zverkov NG, Rogov MA, Stenshin IM, Baykina EM. 2019. Colymbosaurines  
855 from the Upper Jurassic of European Russia and their implication for palaeobiogeography  
856 of marine reptiles. *Palaeobiodiversity and Palaeoenvironments* 100:197–218. DOI:  
857 10.1007/s12549-019-00397-0.
- 858 Arkhangel'sky MS, Zverkov NG, Spasskaya OS, Evgrafov A V. 2018. On the First Reliable  
859 Record of the Ichthyosaur *Ophthalmosaurus icenicus* Seeley in the Oxfordian –  
860 Kimmeridgian Beds of European Russia. *Paleontological Journal* 52:49–57. DOI:  
861 10.1134/S0031030118010033.
- 862 Arnaud M, Monleau C, Wenz S. 1976. Découverte de restes d'ichthyosaure dans l'Aalénien du  
863 Massif de la Loube (Var). *Bulletin du Musée d'Histoire Naturelle de Marseille* 36:3.
- 864 Bakker RT. 1993. Plesiosaur Extinction Cycles — Events that Mark the Beginning, Middle and  
865 End of the Cretaceous. In: Caldwell WGE, Kauffman EG eds. *Evolution of the Western*  
866 *Interior Basin: Geological Association of Canada, Special Paper.* Stittsville, Ontario,  
867 Canada, 641–664.
- 868 Bardet N. 1994. Extinction events among Mesozoic marine reptiles. *Historical Biology* 7:313–  
869 324.
- 870 Bardet N. 1995. Evolution et extinction des reptiles marins au cours du Mésozoïque.  
871 *Palaeovertebrata* 24:177–283.
- 872 Bardet N, Falconnet J, Fischer V, Houssaye A, Jouve S, Pereda Suberbiola X, Pérez-García A,  
873 Rage J-CJ-C, Vincent P. 2014. Mesozoic marine reptile palaeobiogeography in response to  
874 drifting plates. *Gondwana Research* 26:869–887. DOI: 10.1016/j.gr.2014.05.005.
- 875 Bardet N, Fischer V, Machalski M. 2016. Large predatory marine reptiles from the Albian–  
876 Cenomanian of Annopol, Poland. *Geological Magazine* 153:1–16. DOI:  
877 10.1017/S0016756815000254.
- 878 Bardet N, Godefroit P, Sciau J. 1999. A new elasmosaurid Plesiosaur from the Lower Jurassic of  
879 Southern France. *Palaeontology* 42:927–952.
- 880 Bardet N, Houssaye A, Rage J-C, Suberbiola XP. 2008. The Cenomanian-Turonian (Late  
881 Cretaceous) radiation of marine squamates (Reptilia): the role of the Mediterranean Tethys.  
882 *Bulletin de la Société géologique de France* 179:605–622.
- 883 Barrientos-Lara JJ, Alvarado-Ortega J. 2020. *Acuetzpalin carranzai* gen et sp. nov. A new

- ophthalmosauridae (Ichthyosauria) from the Upper Jurassic of Durango, North Mexico. *Journal of South American Earth Sciences* 98:102456. DOI: 10.1016/j.jsames.2019.102456.
- Bell MA. 2015. geoscale: Geological Time Scale Plotting.
- Benson RBJ. 2013. Marine Reptiles. In: MacLeod N, Archibald JD, Levin P eds. *Grzimek's Animal Life Encyclopedia, Extinction*. Farmington Hills, Michigan: Gale Cengage Learning, 267–279.
- Benson RBJ, Bowdler T. 2014. Anatomy of *Colymbosaurus megadeirus* (Reptilia, Plesiosauria) from the Kimmeridge Clay Formation of the U.K., and high diversity among Late Jurassic plesiosauroids. *Journal of Vertebrate Paleontology* 34:1053–1071. DOI: 10.1080/02724634.2014.850087.
- Benson RBJ, Butler RJ. 2011. Uncovering the diversification history of marine tetrapods: ecology influences the effect of geological sampling biases. In: McGowan AJ, Smith AB eds. *Comparing the geological and fossil records: implications for biodiversity studies*. London: Geological Society, Special Publications, 191–208.
- Benson RBJ, Butler RJ, Lindgren J, Smith AS. 2010. Mesozoic marine tetrapod diversity: mass extinctions and temporal heterogeneity in geological megabiases affecting the vertebrates. *Proceedings of the Royal Society B: Biological Sciences* 277:829–834.
- Benson RBJ, Druckenmiller PS. 2014. Faunal turnover of marine tetrapods during the Jurassic–Cretaceous transition. *Biological Reviews* 89:1–23. DOI: 10.1111/brv.12038.
- Benson RBJ, Evans M, Druckenmiller PS. 2012. High diversity, low disparity and small body size in plesiosaurs (Reptilia, Sauropterygia) from the Triassic–Jurassic boundary. *PLoS ONE* 7:e31838.
- Benson RBJ, Evans M, Smith AS, Sassoon J, Moore-Faye S, Ketchum HF, Forrest R. 2013. A giant pliosaurid skull from the Late Jurassic of England. *PLoS ONE* 8:e65989. DOI: 10.1371/journal.pone.0065989.
- Benson RBJ, Ketchum HF, Noè LF, Gómez-Pérez M. 2011. New information on *Hauffiosaurus* (Reptilia, Plesiosauria) based on a new species from the Alum Shale member (Lower Toarcian: Lower Jurassic) of Yorkshire, UK. *Palaeontology* 54:547–571.
- Benson RBJ, Zverkov NG, Arkhangelsky MS. 2015. Youngest occurrences of rhomaleosaurid plesiosaurs indicate survival of an archaic marine reptile clade at high palaeolatitudes. *Acta Palaeontologica Polonica* 60:769–780.
- Benton MJ, Taylor MA. 1984. Marine reptiles from the Upper Lias (Lower Toarcian, Lower Jurassic) of the Yorkshire coast. *Proceedings of the Yorkshire Geological Society* 44:399–429.
- Böttcher VR. 1989. Über die Nahrung eines *Leptopterygius* (Ichthyosauria, Reptilia) aus dem süddeutschen Posidonienschiefer (Unterer Jura) mit Bemerkungen über den Magen der Ichthyosaurier. *Stuttgarter Beiträge zur Naturkunde Serie B (Geologie und paläontologie)* 155:1–19.
- Boulvain F, Belanger I, Delsate D, Ghysel P, Godefroit P, Laloux M, Roche M. 2001. Triassic and Jurassic lithostratigraphic units (Belgian Lorraine). *Geologica Belgica* 4:113–119.
- Brown D. 1981. The English Upper Jurassic Plesiosauroidea (Reptilia) and a review of the phylogeny and classification of the Plesiosauria. *Bulletin of the British Museum (Natural History) Geology* 35:253–347.
- Brusatte SL, Young MT, Challands TJ, Clark NDL, Fischer V, Fraser NC, Liston JJ, MacFadyen CCJ, Ross DA, Walsh SA, Wilkinson M. 2015. Ichthyosaurs from the Jurassic of Skye, Scotland. *Scottish Journal of Geology* 51:43–55. DOI: 10.1144/sjg2014-018.



- 930 Buchholtz EA. 2001. Swimming styles in Jurassic ichthyosaurs. *Journal of Vertebrate*  
931 *Paleontology* 21:61–73.
- 932 Buchy M-C. 2010. Morphologie dentaire et régime alimentaire des reptiles marins du  
933 Mésozoïque : revue critique et réévaluation. *Oryctos* 9:49–82.
- 934 Cabrera A. 1939. Sobre un nuevo ictiosaurio del Neuquén. *Notas del Museo de La Plata -*  
935 *Paleontología* 4:485–491.
- 936 Caine H, Benton MJ. 2011. Ichthyosauria from the upper Lias of Strawberry Bank, England.  
937 *Palaeontology* 54:1069–1093.
- 938 Callaway JM, Massare J. 1989. Geographic and stratigraphic distribution of the Triassic  
939 Ichthyosauria (Reptilia; Diapsida). *Neues Jahrbuch für Geologie und Paläontologie,*  
940 *Abhandlungen* 187:37–58.
- 941 Campos L, Fernández MS, Herrera Y. 2020. A new ichthyosaur from the Late Jurassic of north-  
942 west Patagonia (Argentina) and its significance for the evolution of the narial complex of  
943 the ophthalmosaurids. *Zoological Journal of the Linnean Society* 188:180–201. DOI:  
944 10.1093/zoolinnean/zlz095.
- 945 Cau A. 2014. The affinities of ‘*Steneosaurus barettoni*’ (Crocodylomorpha, Thalattosuchia),  
946 from the Jurassic of Northern Italy, and implications for cranial evolution among geosaurine  
947 metriorhynchids. *Historical Biology* 26:433–440. DOI: 10.1080/08912963.2013.784906.
- 948 Cau A, Fanti F. 2011. The oldest known metriorhynchid crocodylian from the Middle Jurassic of  
949 North-eastern Italy: *Neptunidraco ammoniticus* gen. et sp. nov. *Gondwana Research*  
950 19:550–565.
- 951 Di Cencio A, Weis R. 2020. Revision of upper Toarcian ammonites (Lytoceratidae,  
952 Graphoceratidae and Hammatoceratidae) from the Minette ironstones, southern  
953 Luxembourg. In: Di Cencio A, Sadki D, Weis R eds. *Paléontologie au Luxembourg (2) Les*  
954 *ammonites de la Minette*. Ferrantia, in press.
- 955 Conybeare WD. 1822. Additional notes on the fossil genera *Ichthyosaurus* and *Plesiosaurus*.  
956 *Transactions of the Geological Society of London* 2:103–123.
- 957 Cruickshank a. RI. 1994. Cranial Anatomy of the Lower Jurassic Pliosaur *Rhomaleosaurus*  
958 *megacephalus* (Stutchbury) (Reptilia: Plesiosauria). *Philosophical Transactions of the*  
959 *Royal Society B: Biological Sciences* 343:247–260. DOI: 10.1098/rstb.1994.0024.
- 960 Delsate D, Suberbiola XP, Felten R, Felten G. 2018. First thyreophoran dinosaur from the  
961 Middle Jurassic (Bajocian) of Luxembourg. *Geologica Belgica* 21:19–26.
- 962 Delsate D, Weis R. 2010. La Couche à Crassum (Toarcien moyen) au Luxembourg: stratigraphie  
963 et faunes de la coupe de Dudelange-Zoufftgen. *Ferrantia* 62:35–60.
- 964 Delsett LL, Roberts AJ, Druckenmiller PS, Hurum JH. 2019. Osteology and phylogeny of Late  
965 Jurassic ichthyosaurs from the Slottsmøya Member Lagerstätte (Spitsbergen, Svalbard).  
966 *Acta Palaeontologica Polonica*:717–743. DOI: 10.4202/app.00571.2018.
- 967 Dera G, Toumoulin A, de Baets K. 2016. Diversity and morphological evolution of Jurassic  
968 belemnites from South Germany. *Palaeogeography, Palaeoclimatology, Palaeoecology*  
969 457:80–97. DOI: 10.1016/j.palaeo.2016.05.029.
- 970 Dick DG, Maxwell EE. 2015. The evolution and extinction of the ichthyosaurs from the  
971 perspective of quantitative ecospace modelling. *Biology letters* 11:20150339-. DOI:  
972 10.1098/rsbl.2015.0339.
- 973 Dick DG, Schweigert G, Maxwell EE. 2016. Trophic niche ontogeny and palaeoecology of early  
974 Toarcian *Stenopterygius* (Reptilia: Ichthyosauria). *Palaeontology*:n/a-n/a. DOI:  
975 10.1111/pala.12232.

- 976 Dittrich D. 1993. *Erläuterungen zur geologischen Karte von Luxemburg, 1:25000: Blatt 11*  
977 *Grevenmacher und Blatt 13 Remic.*
- 978 Doyle P. 1987. Lower Jurassic-Lower Cretaceous belemnite biogeography and the development  
979 of the Mesozoic Boreal Realm. *Palaeogeography, Palaeoclimatology, Palaeoecology*  
980 61:237–254. DOI: [https://doi.org/10.1016/0031-0182\(87\)90052-6](https://doi.org/10.1016/0031-0182(87)90052-6).
- 981 Druckenmiller PS, Hurum J, Knutsen EM, Nakrem HA. 2012. Two new ophthalmosaurids  
982 (Reptilia: Ichthyosauria) from the Agardhfjellet Formation (Upper Jurassic:  
983 Volgian/Tithonian), Svalbard, Norway. *Norwegian Journal of Geology* 92:311–339.
- 984 Druckenmiller PS, Maxwell EE. 2014. A Middle Jurassic (Bajocian) ophthalmosaurid (Reptilia,  
985 Ichthyosauria) from the Tuxedni Formation, Alaska and the early diversification of the  
986 clade. *Geological Magazine* 151:41–48. DOI: [doi:10.1017/S0016756813000125](https://doi.org/10.1017/S0016756813000125).
- 987 Dzyuba OS, Schraer CD, Hults CP, Blodgett RB, Schraer DJ. 2019. Early Bajocian belemnites  
988 of Southcentral Alaska: new data and new perspectives on mid-Middle Jurassic  
989 Megateuthididae and Belemnopseidae biogeography. *Journal of Systematic Palaeontology*  
990 17:911–935. DOI: [10.1080/14772019.2018.1486335](https://doi.org/10.1080/14772019.2018.1486335).
- 991 Fernández M. 1994. A new long-snouted ichthyosaur from the early Bajocian of Neuquén basin  
992 (Argentina). *Ameghiniana* 31:291–297.
- 993 Fernández M. 1999. A new ichthyosaur from the Los Molles Formation (Early Bajocian),  
994 Neuquén basin, Argentina. *Journal of Paleontology* 73:677–681.
- 995 Fernández M. 2003. Ophthalmosauria (Ichthyosauria) forefin from the Aalenian-Bajocian  
996 boundary of Mendoza Province, Argentina. *Journal of Vertebrate Paleontology* 23:691–  
997 694.
- 998 Fernández M, Iturralde-Vinent M. 2000. An oxfordian Ichthyosauria (Reptilia) from Viñales,  
999 Western Cuba: paleobiogeographic significance. *Journal of Vertebrate Paleontology*  
1000 20:191–193.
- 1001 Fernández MS, Maxwell EE. 2012. The genus *Arthropterygius* Maxwell (Ichthyosauria:  
1002 Ophthalmosauridae) in the Late Jurassic of the Neuquén Basin, Argentina. *Geobios* 45:535–  
1003 540.
- 1004 Fernández M, Talevi M. 2014. Ophthalmosaurian (Ichthyosauria) records from the Aalenian–  
1005 Bajocian of Patagonia (Argentina): an overview. *Geological Magazine* 151:49–59. DOI:  
1006 [10.1017/S0016756813000058](https://doi.org/10.1017/S0016756813000058).
- 1007 Fischer V. 2016. Taxonomy of *Platypterygius campylodon* and the diversity of the last  
1008 ichthyosaurs. *PeerJ* 4:1–21. DOI: [10.7717/peerj.2604](https://doi.org/10.7717/peerj.2604).
- 1009 Fischer V, Appleby RM, Naish D, Liston J, Riding JB, Brindley S, Godefroit P. 2013. A basal  
1010 thunnosaurian from Iraq reveals disparate phylogenetic origins for Cretaceous ichthyosaurs.  
1011 *Biology Letters* 9:1–6. DOI: [10.1098/rsbl.2013.0021](https://doi.org/10.1098/rsbl.2013.0021).
- 1012 Fischer V, Arkhangelsky MS, Uspensky GN, Stenshin IM, Godefroit P. 2014a. A new Lower  
1013 Cretaceous ichthyosaur from Russia reveals skull shape conservatism within  
1014 Ophthalmosaurinae. *Geological Magazine* 151:60–70. DOI: [10.1017/S0016756812000994](https://doi.org/10.1017/S0016756812000994).
- 1015 Fischer V, Bardet N, Benson RBJ, Arkhangelsky MS, Friedman M. 2016. Extinction of fish-  
1016 shaped marine reptiles associated with reduced evolutionary rates and global environmental  
1017 volatility. *Nature communications* 7:1–11. DOI: [10.1038/ncomms10825](https://doi.org/10.1038/ncomms10825).
- 1018 Fischer V, Bardet N, Guimaraes M, Godefroit P. 2014b. High Diversity in Cretaceous Ichthyosaurs  
1019 from Europe Prior to Their Extinction. *PLoS ONE* 9:e84709. DOI:  
1020 [10.1371/journal.pone.0084709](https://doi.org/10.1371/journal.pone.0084709).
- 1021 Fischer V, Benson RBJ, Druckenmiller PS, Ketchum HF, Bardet N. 2018. The evolutionary

- 1022 history of polycotyloid plesiosaurians. *Royal Society Open Science* 5. DOI:  
1023 10.1098/rsos.172177.
- 1024 Fischer V, Benson RBJ, Zverkov NG, Soul LC, Arkhangelsky MS, Lambert O, Stenshin IM,  
1025 Uspensky GN, Druckenmiller PS. 2017. Plasticity and convergence in the evolution of  
1026 short-necked plesiosaurs. *Current Biology* 27:1667–1676. DOI: 10.1016/j.cub.2017.04.052.
- 1027 Fischer V, Cappetta H, Vincent P, Garcia G, Goolaerts S, Martin JE, Roggero D, Valentin X.  
1028 2014c. Ichthyosaurs from the French Rhaetian indicate a severe turnover across the  
1029 Triassic–Jurassic boundary. *Naturwissenschaften* 101:1027–1040. DOI: 10.1007/s00114-  
1030 014-1242-7.
- 1031 Fischer V, Guimar M, Godefroit P. 2011. New data on the palaeobiogeography of Early  
1032 Jurassic marine reptiles: the Toarcian ichthyosaur fauna of the Vocontian Basin (SE  
1033 France). *Neues Jahrbuch für Geologie und Paläontologie* 261:111–127. DOI:  
1034 10.1127/0077-7749/2011/0155.
- 1035 Fischer V, Maisch MW, Naish D, Kosma R, Liston J, Joger U, Krüger FJ, Pardo Pérez J, Tainsh  
1036 J, Appleby RM. 2012. New ophthalmosaurid ichthyosaurs from the European Lower  
1037 Cretaceous demonstrate extensive ichthyosaur survival across the Jurassic-Cretaceous  
1038 boundary. *PLoS ONE* 7:e29234. DOI: 10.1371/journal.pone.0029234.
- 1039 Fischer V, Zverkov NG, Arkhangelsky MS, Stenshin IM, Blagoveschensky IV, Uspensky GN.  
1040 2020. A new elasmosaurid plesiosaurian from the Early Cretaceous of Russia marks an  
1041 early attempt at neck elongation. *Zoological Journal of the Linnean Society*:in press. DOI:  
1042 10.1093/zoolinnean/zlaa103.
- 1043 Foffa D, Young MT, Brusatte SL. 2018. Filling the Corallian gap : New information on Late  
1044 Jurassic marine reptile faunas from England. *Acta Palaeontologica Polonica* 63:287–313.
- 1045 Foffa D, Young MT, Brusatte SL, Graham MR, Steel L. 2017. A new metriorhynchid  
1046 crocodylomorph from the Oxford Clay Formation (Middle Jurassic) of England, with  
1047 implications for the origin and diversification of Geosaurini. *Journal of Systematic  
1048 Palaeontology* 2019:1–21. DOI: 10.1080/14772019.2017.1367730.
- 1049 Foffa D, Young MT, Stubbs TL, Dexter KG, Brusatte SL. 2018. The long-term ecology and  
1050 evolution of marine reptiles in a Jurassic seaway. *Nature Ecology and Evolution* 2:1548–  
1051 1555. DOI: 10.1038/s41559-018-0656-6.
- 1052 Gasparini Z. 1997. A new pliosaur from the Bajocian of the Neuquen Basin, Argentina.  
1053 *Palaeontology* 40:135–147.
- 1054 Gasparini Z, Fernández MS, de la Fuente MS, Salgado L. 2007. Reptiles marinos Jurásicos y  
1055 Cretácicos de la Patagonia Argentina: su aporte al conocimiento de la herpetofauna  
1056 mesozoica. *Ameghiniana* APA Public:125–136.
- 1057 Gasparini Z, Pol D, Spalletti LA. 2006. An unusual marine crocodyliiform from the Jurassic-  
1058 Cretaceous boundary of Patagonia. *Science* 311:70–73.
- 1059 Gilmore CW. 1902. Discovery of teeth in *Baptanodon*, an ichthyosaurian from the Jurassic of  
1060 Wyoming. *Science* 16:913–914.
- 1061 Gilmore CW. 1906. Notes on osteology of *Baptanodon*. *Memoirs of the Carnegie Museum*  
1062 II:325–337.
- 1063 Godefroit P. 1993a. Les grands ichthyosaures sinémuriens d’Arlon. *Bulletin de l’Institut Royal  
1064 des Sciences Naturelles de Belgique Sciences de la Terre* 63:25–71.
- 1065 Godefroit P. 1993b. The skull of *Stenopterygius longifrons* (Owen, 1881). *Revue de  
1066 Paléobiologie de Genève volume spécial* 7:67–84.
- 1067 Godefroit P. 1994a. Les reptiles marins du Toarcien (Jurassique inférieur) belgo-

- 1068 luxembourgeois. *Mémoires pour servir à l'Explication des Cartes Géologiques et Minières*  
1069 *de la Belgique* 39:98.
- 1070 Godefroit P. 1994b. *Simolestes keileni* sp. nov., un Pliosau (Plesiosauria, Reptilia) du Bajocien  
1071 supérieur de Lorraine (France). *Bulletin des Académie et Société Lorraines des sciences*  
1072 33:77–95.
- 1073 Großmann F. 2007. The taxonomic and phylogenetic position of the Plesiosauroidea from the  
1074 Lower Jurassic Posidonia shale of south-west Germany. *Palaeontology* 50:545–564.
- 1075 Guérin-Franiatte S, Weis R. 2010. Le passage Aalénien-Bajocien près de Rumelange: la série  
1076 biostratigraphique dans le Bassin d'Esch-sur-Alzette (Grand-Duché de Luxembourg). In:  
1077 Weis R, Guérin-Franiatte S eds. *Le Jurassique inférieur et moyen au Luxembourg*.  
1078 Luxembourg ville, Luxembourg: Ferrantia, 73–96.
- 1079 Hauff B. 1953. *Das Holzmadenbuch*. Öhringen.
- 1080 Herrera Y, Gasparini Z, Fernández MS. 2013. A new Patagonian species of Cricosaurus  
1081 (Crocodyliformes, Thalattosuchia): First evidence of Cricosaurus in Middle-Upper  
1082 Tithonian lithographic limestone from Gondwana. *Palaeontology* 56:663–678. DOI:  
1083 10.1111/pala.12010.
- 1084 Home E. 1819. Reasons for giving the name Proteo-Saurus to the fossil skeleton which has been  
1085 described. *Philosophical transactions of the Royal Society of London* 109:212–216.
- 1086 Huene F von. 1922. *Die Ichthyosaurier des Lias und ihre Zusammenhänge*. Berlin: Verlag von  
1087 Gebrüder Borntraeger.
- 1088 Huene F von. 1931. Neue Studien über Ichthyosaurier aus Holzmaden. *Abhandlungen der*  
1089 *Senckenbergischen Naturforschenden Gesellschaft* 42:345–382.
- 1090 Huene F von. 1951. Ein neuer Fund von *Eurhinosaurus longirostris*. *Neues Jahrbuch für*  
1091 *Geologie und Paläontologie, Abhandlungen* 93:277–283.
- 1092 Johnson R. 1979. The osteology of the pectoral complex of *Stenopterygius* Jaekel (Reptilia:  
1093 Ichthyosauria). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 159:41–86.
- 1094 Johnson MM, Young MT, Brusatte SL. 2020. Emptying the wastebasket: a historical and  
1095 taxonomic revision of the Jurassic crocodylomorph Steneosaurus. *Zoological Journal of the*  
1096 *Linnean Society* 189:428–448. DOI: 10.1093/zoolinnean/zlaa027.
- 1097 Johnson MM, Young MT, Brusatte SL, Thuy B, Weis R. 2018. A catalogue of teleosauroids  
1098 (Crocodylomorpha: Thalattosuchia) from the Toarcian and Bajocian (Jurassic) of southern  
1099 Luxembourg. *Historical Biology* 2963:1–16. DOI: 10.1080/08912963.2018.1427090.
- 1100 Kear BP. 2012. A revision of Australia's Jurassic plesiosaurs. *Palaeontology* 55:1125–1138.  
1101 DOI: 10.1111/j.1475-4983.2012.01183.x.
- 1102 Kelley NP, Pyenson ND. 2015. Evolutionary innovation and ecology in marine tetrapods from  
1103 the Triassic to the Anthropocene. *Science* 348:aaa3716. DOI: 10.1126/science.aaa3716.
- 1104 Ketchum HF, Benson RBJ. 2011a. A new pliosaurid (Sauropterygia, Plesiosauria) from the  
1105 Oxford Clay Formation (Middle Jurassic, Callovian) of England: evidence for a gracile,  
1106 longirostrine grade of Early–Middle Jurassic pliosaurids. *Special Papers in Palaeontology*  
1107 86:109–129.
- 1108 Ketchum HF, Benson RBJ. 2011b. The cranial anatomy and taxonomy of *Peloneustes philarchus*  
1109 (Sauropterygia, Pliosauridae) from the Peterborough Member (Callovian, Middle Jurassic)  
1110 of the United Kingdom. *Palaeontology* 54:639–665. DOI: 10.1111/j.1475-  
1111 4983.2011.01050.x.
- 1112 Klein N, Schmitz L, Wintrich T, Sander PM. 2020. A new cymbospondylid ichthyosaur  
1113 (Ichthyosauria) from the Middle Triassic (Anisian) of the Augusta Mountains, Nevada,

- 1114 USA. *Journal of Systematic Palaeontology* 18:1167–1191. DOI:  
1115 10.1080/14772019.2020.1748132.
- 1116 Knight WC. 1903. Notes on genus *Baptanodon*. *American Journal of Science, Fourth Series*  
1117 16:78–81.
- 1118 Knutsen EM, Druckenmiller PS, Hurum JH. 2012a. A new plesiosauroid (Reptilia:  
1119 Sauropterygia) from the Agardhfjellet Formation (Middle Volgian) of central Spitsbergen,  
1120 Norway. *Norwegian Journal of Geology* 92:213–234.
- 1121 Knutsen EM, Druckenmiller PS, Hurum JH. 2012b. Two new species of long-necked  
1122 plesiosaurs (Reptilia: Sauropterygia) from the Upper Jurassic (Middle Volgian)  
1123 Agardhfjellet Formation of central Spitsbergen. *Norwegian Journal of Geology* 92:187–  
1124 212.
- 1125 Knutsen EM, Druckenmiller PS, Hurum JH. 2012c. Redescription and taxonomic clarification of  
1126 “*Tricleidus*” svalbardensis based on new material from the Agardhfjellet Formation (Middle  
1127 Volgian). *Norwegian Journal of Geology* 92:175–186.
- 1128 Korte C, Hesselbo SP, Ullmann C V., Dietl G, Ruhl M, Schweigert G, Thibault N. 2015. Jurassic  
1129 climate mode governed by ocean gateway. *Nature Communications* 6:10015. DOI:  
1130 10.1038/ncomms10015.
- 1131 Lomax DR. 2016. A new leptonectid ichthyosaur from the Lower Jurassic (Hettangian) of  
1132 Nottinghamshire, England, UK, and the taxonomic usefulness of the ichthyosaurian  
1133 coracoid. *Journal of Systematic Palaeontology*:1–15. DOI:  
1134 10.1080/14772019.2016.1183149.
- 1135 Lomax DR, Evans M, Carpenter S. 2018. An ichthyosaur from the UK Triassic – Jurassic  
1136 boundary : A second specimen of the leptonectid ichthyosaur *Wahlisaurus massarae* Lomax  
1137 2016. *Geological Journal*:1–8. DOI: 10.1002/gj.3155.
- 1138 Lomax DR, Massare JA. 2015. A new species of *Ichthyosaurus* from the Lower Jurassic of  
1139 West Dorset, England, U.K. *Journal of Vertebrate Paleontology*:903260. DOI:  
1140 10.1080/02724634.2014.903260.
- 1141 Lomax DR, Massare JA. 2016. Two new species of *Ichthyosaurus* from the lowermost Jurassic  
1142 (Hettangian) of Somerset, England. *Papers in Palaeontology*. DOI: 10.1002/SPP2.1065.
- 1143 Lomax DR, Porro LB, Larkin NR. 2019. Descriptive anatomy of the largest known specimen of  
1144 *Protoichthyosaurus prostaurodon* (Reptilia: Ichthyosauria) including computed tomography  
1145 and digital reconstruction of a three-dimensional skull. *PeerJ* 7:e6112. DOI:  
1146 10.7717/peerj.6112.
- 1147 Lomax DR, Sachs S. 2017. On the largest *Ichthyosaurus* : A new specimen of *Ichthyosaurus*  
1148 *somersetensis* containing an embryo. *Acta Palaeontologica Polonica* 62:575–584.
- 1149 Long JA, Cruickshank ARI. 1998. Further Records of Plesiosaurian Reptiles of Jurassic and  
1150 Cretaceous Age From Western Australia. *Records of the Western Australian Museum* 19:47.
- 1151 Lucius M. 1945. Die Luxemburger Minetteformation und die jüngeren Eisenerzbildungen  
1152 unseres Landes. Beiträge zur Geologie von Luxemburg. *Service Carte géologique du*  
1153 *Luxembourg* 4:1–347.
- 1154 Maisch MW. 1998. A new ichthyosaur genus from the Posidonia Shale (Lower Toarcian,  
1155 Jurassic) of Holzmaden, SW-Germany with comments on the phylogeny of post-Triassic  
1156 ichthyosaurs. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 209:47–78.
- 1157 Maisch MW. 2008. Revision der Gattung *Stenopterygius* Jaekel, 1904 emend. von Huene, 1922  
1158 (Reptilia: Ichthyosauria) aus dem unteren Jura Westeuropas. *Palaeodiversity* 1:227–271.
- 1159 Maisch MW. 2010. Phylogeny, systematics, and origin of the Ichthyosauria - the state of the art.

- 1160 *Palaeodiversity* 3:151–214.
- 1161 Maisch MW, Matzke AT. 2000. The Ichthyosauria. *Stuttgarter Beiträge zur Naturkunde Serie B*
- 1162 *(Geologie und Paläontologie)* 298:1–159.
- 1163 Maisch MW, Matzke AT. 2001. The cranial osteology of the Middle Triassic ichthyosaur
- 1164 *Contectopalatus* from Germany. *Palaeontology* 44:1127–1156.
- 1165 Maisch MW, Matzke AT. 2003. The cranial osteology of the ichthyosaur *Leptonectes* cf.
- 1166 *tenuirostris* from the Lower Jurassic of England. *Journal of Vertebrate Paleontology*
- 1167 23:116–127. DOI: 10.1671/0272-4634(2003)23[116:TCOOTI]2.0.CO;2.
- 1168 Mariotti N, Santantonio M, Weis R. 2007. Aalenian-Early Bajocian Belemnite Assemblage From
- 1169 Peri-Mediterranean Tethyan Sediments ( Calabria , Southern Italy ). *Geologica Romana*
- 1170 40:1–19.
- 1171 Mariotti N, Weis R, Di Cencio A, Clément A, De Baets K. 2012. New records of early Middle
- 1172 Jurassic belemnites in the French Subalpine Basin and their paleobiogeographic
- 1173 significance. *Geobios* 45:99–108. DOI: <https://doi.org/10.1016/j.geobios.2011.11.012>.
- 1174 Marsh OC. 1895. The Reptilia of the *Baptanodon* Beds. *American Journal of Science* 50:405–
- 1175 406.
- 1176 Martin JE, Fischer V, Vincent P, Suan G. 2012. A longirostrine *Temnodontosaurus*
- 1177 (Ichthyosauria) with comments on Early Jurassic ichthyosaur niche partitioning and
- 1178 disparity. *Palaeontology* 55:995–1005. DOI: 10.1111/j.1475-4983.2012.01159.x.
- 1179 Massare JA. 1987. Tooth morphology and prey preference of Mesozoic marine reptiles. *Journal*
- 1180 *of Vertebrate Paleontology* 7:121–137.
- 1181 Massare JA. 1988. Swimming capabilities of Mesozoic marine reptiles: implications for the
- 1182 methods of predation. *Palaeobiology* 14:187–205.
- 1183 Massare JA. 1997. Faunas, behavior, and evolution. In: Callaway JM, Nicholls EL eds. *Ancient*
- 1184 *Marine Reptiles*. San Diego: Academic Press, 401–421.
- 1185 Massare JA, Buchholtz EA, Kenney J, Chomat A-M. 2006. Vertebral morphology of
- 1186 *Ophthalmosaurus natans* (Reptilia: Ichthyosauria) from the Jurassic Sundance Formation of
- 1187 Wyoming. *Paludicola* 5:242–254.
- 1188 Massare JA, Wahl WR, Ross M, Connely M V. 2014. Palaeoecology of the marine reptiles of the
- 1189 Redwater Shale Member of the Sundance Formation (Jurassic) of central Wyoming, USA.
- 1190 *Geological Magazine* 151:167–182. DOI: [doi:10.1017/S0016756813000472](https://doi.org/10.1017/S0016756813000472).
- 1191 Maubeuge PL. 1972. *Etudes stratigraphiques sur la formation ferrifère de Lorraine et ses morts-*
- 1192 *terrains*. Metz: Self-edited.
- 1193 Maxwell EE. 2010. Generic reassignment of an ichthyosaur from the Queen Elizabeth Islands,
- 1194 Northwest Territories, Canada. *Journal of Vertebrate Paleontology* 30:403–415.
- 1195 Maxwell EE. 2012. New metrics to differentiate species of *Stenopterygius* (Reptilia:
- 1196 Ichthyosauria) from the Lower Jurassic of southwestern Germany. *Journal of Paleontology*
- 1197 86:105–115. DOI: 10.1666/11-038.1.
- 1198 Maxwell EE, Caldwell MW, Lamoureux DO. 2012. Tooth histology, attachment, and
- 1199 replacement in the Ichthyopterygia reviewed in an evolutionary context. *Paläontologische*
- 1200 *Zeitschrift* 86:1–14.
- 1201 Maxwell EE, Cortés D. 2020. A revision of the Early Jurassic ichthyosaur *Hauffiopteryx*
- 1202 (Reptilia: Ichthyosauria), and description of a new species from Southwestern Germany.
- 1203 *Palaeontologia Electronica* 23:1–43. DOI: 10.26879/937.
- 1204 Maxwell EE, Fernández MS, Schoch RR. 2012. First diagnostic marine reptile remains from the
- 1205 Aalenian (Middle Jurassic): a new ichthyosaur from southwestern Germany. *PLoS ONE*

- 1206 7:e41692. DOI: 10.1371/journal.pone.0041692.
- 1207 Maxwell EE, Vincent P. 2015. Effects of the early Toarcian Oceanic Anoxic Event on
- 1208 ichthyosaur body size and faunal composition in the Southwest German Basin.
- 1209 *Paleobiology*:1–10. DOI: 10.1017/pab.2015.34.
- 1210 Mazin J-M. 1982. Affinités et phylogénie des Ichthyopterygia. *Geobios, mémoire spécial* 6:85–
- 1211 98.
- 1212 McGowan C. 1973. The cranial morphology of the Lower Liassic latipinnate ichthyosaurs of
- 1213 England. *Bulletin of the British Museum (Natural History) Geology* 24:1–109.
- 1214 McGowan C. 1974. A revision of the longipinnate ichthyosaurs of the Lower Jurassic of
- 1215 England, with description of the new species (Reptilia, Ichthyosauria). *Life Science*
- 1216 *Contributions, Royal Ontario Museum* 97:1–37.
- 1217 McGowan C. 1979. A revision of the Lower Jurassic ichthyosaurs of Germany with descriptions
- 1218 of two new species. *Palaeontographica. Abteilung A. Paläozoologie, Stratigraphie* 166:93–
- 1219 135.
- 1220 McGowan C. 1986. A putative ancestor for the swordfish-like ichthyosaur *Eurhinosaurus*.
- 1221 *Nature* 322:454–456.
- 1222 McGowan C. 1994. A new species of *Shastasaurus* (Reptilia: Ichthyosauria) from the Triassic of
- 1223 British Columbia; the most complete exemplar of the genus. *Journal of Vertebrate*
- 1224 *Paleontology* 14:168–179.
- 1225 McGowan C. 1996. Giant ichthyosaurs of the Early Jurassic. *Canadian Journal of Earth*
- 1226 *Sciences* 33:1011–1021.
- 1227 McGowan C. 2003. A new specimen of *Excalibosaurus* from the English Lower Jurassic.
- 1228 *Journal of Vertebrate Paleontology* 23:950–956.
- 1229 McGowan C, Milner AC. 1999. A new Pliensbachian ichthyosaur from Dorset, England.
- 1230 *Palaeontology* 42:761–768.
- 1231 McGowan C, Motani R. 2003. *Part 8. Ichthyopterygia*. München: Verlag Dr. Friedrich Pfeil.
- 1232 Mehl MG. 1912. *Muraenosaurus? Reedii*, Sp. Nov and *Tricleidus? Laramiensis* Knight,
- 1233 American Jurassic plesiosaurs. *The Journal of Geology* 20:344–352.
- 1234 Merriam JC. 1908. Triassic Ichthyosauria with special reference to the American forms.
- 1235 *Memoirs of the University of California* 1:1–154.
- 1236 Moon BC. 2017. A new phylogeny of ichthyosaurs (Reptilia: Diapsida). *Journal of Systematic*
- 1237 *Palaeontology* 17:129–155. DOI: 10.1080/14772019.2017.1394922.
- 1238 Moon BC, Kirton AM. 2018a. Ichthyosaurs of the British Middle and Upper Jurassic Part 2
- 1239 *Brachypterygius, Nannopterygius, Macropterygius* and Taxa invalida. *Monographs of the*
- 1240 *Palaeontographical Society* 172:85–197. DOI: 10.1080/02693445.2018.1468139.
- 1241 Moon BC, Kirton AM. 2018b. Ichthyosaurs of the British Middle and Upper Jurassic Part 1,
- 1242 *Ophthalmosaurus*. *Monographs of the Palaeontographical Society* 170:1–84. DOI:
- 1243 10.1080/02693445.2016.11963958.
- 1244 Moon BC, Stubbs TL. 2020. Early high rates and disparity in the evolution of ichthyosaurs.
- 1245 *Communications Biology* 3:1–8. DOI: 10.1038/s42003-020-0779-6.
- 1246 Motani R. 1997. Temporal and Spatial Distribution of Tooth Implantation in Ichthyosaurs. In:
- 1247 Callaway JM, Nicholls EL eds. *Ancient Marine Reptiles*. San Diego, California: Academic
- 1248 Press, 81–103.
- 1249 Motani R. 1999a. Phylogeny of the Ichthyopterygia. *Journal of Vertebrate Paleontology* 19:473–
- 1250 496.
- 1251 Motani R. 1999b. On the evolution and homologies of ichthyosaurian forefins. *Journal of*

- 1252 *Vertebrate Paleontology* 19:28–41.
- 1253 Motani R. 2009. The Evolution of Marine Reptiles. *Evo Edu Outreach* 2:224–235.
- 1254 Neige P, Weis R, Fara E. 2020. Ups and downs of belemnite diversity in the Early Jurassic of
- 1255 Western Tethys. *Palaeontology*:In Press.
- 1256 Noe LF. 1999. The Callovian pliosaurs of the Oxford Clay—evidence and implications for the
- 1257 consumption of marine invertebrates. In: *Secondary Adaptation to Life in Water/La*
- 1258 *réadaptation au milieu aquatique*. 13–17.
- 1259 Noe LF. 2001. A taxonomic and functional study of the Callovian (Middle Jurassic)
- 1260 Pliosauroida (Reptilia, Sauropterygia). University of Derby.
- 1261 O’Keefe FR. 2001a. A cladistic analysis and taxonomic revision of the Plesiosauria
- 1262 (Reptilia:Sauropterygia). *Acta Zoologica Fennica* 213:1–63.
- 1263 O’Keefe FR. 2001b. Ecomorphology of plesiosaur flipper geometry. *Journal of Evolutionary*
- 1264 *Biology* 14:987–991.
- 1265 O’Keefe FR. 2002. The evolution of plesiosaur and pliosaur morphotypes in the Plesiosauria
- 1266 (Reptilia: Sauropterygia). *Palaeobiology* 28:101–112.
- 1267 O’Keefe RF. 2004. Preliminary description and phylogenetic position of a new Plesiosaur
- 1268 (Reptilia : Sauropterygia) from the Toarcian of Holzmaden, Germany. *Journal of*
- 1269 *Paleontology* 78:973–988. DOI: 10.1666/0022-3360(2004)078<0973:pdappo>2.0.co;2.
- 1270 O’Keefe FR, Wahl WJ. 2003. Preliminary report on the osteology and relationships of a new
- 1271 aberrant cryptocleidoid plesiosaur from the Sundance Formation, Wyoming. *Paludicola*
- 1272 4:48–68.
- 1273 Ösi A, Young MT, Galácz A, Rabi M. 2018. A new large-bodied thalattosuchian crocodyliform
- 1274 from the Lower Jurassic (Toarcian) of Hungary, with further evidence of the mosaic
- 1275 acquisition of marine adaptations in Metriorhynchoidea. *PeerJ* 6:e4668. DOI:
- 1276 10.7717/peerj.4668.
- 1277 Otero RA, Alarcón-Muñoz J, Soto-Acuña S, Rojas J, Rojas O, Ortíz H. 2020. Cryptoclidid
- 1278 plesiosaurs (Sauropterygia, Plesiosauria) from the Upper Jurassic of the Atacama Desert.
- 1279 *Journal of Vertebrate Paleontology*:In Press. DOI: 10.1080/02724634.2020.1764573.
- 1280 Owen R. 1860. On the orders of fossil and recent Reptilia and their distribution in time. *Report of*
- 1281 *the British Association for the Advancement of Science* 29:153–166.
- 1282 Owen R. 1865. A monograph of the Fossil Reptilia of the Liassic Formations, part third:
- 1283 *Plesiosaurus, Dimorphodon, and Ichthyosaurus. Monograph of the Palaeontographical*
- 1284 *society*:12–130.
- 1285 Paparella I, Maxwell EE, Cipriani A, Roncace S, Caldwell MW. 2017. The first ophthalmosaurid
- 1286 ichthyosaur from the Upper Jurassic of the Umbrian-Marchean Apennines (Marche, Central
- 1287 Italy). *Geological Magazine* 154:837–858. DOI: 10.1017/S0016756816000455.
- 1288 Pieńkowski G, Schudack ME, Bosák P, Enay R, Feldman-Olszewska A, Golonka J, Gutowski J,
- 1289 Herngreen GFW, Jordan P, Krobicki M, Lathuiliere B, Leinfelder RR, Michalik J, Mönnig
- 1290 E, Noe-Nygaard N, Pálffy J, Pint A, Rasser MW, Reisdorf AG, Schmid DU, Schweigert G,
- 1291 Surlyk F, Wetzel A, Wong TE. 2008. Jurassic. *The Geology of Central Europe Volume 2:*
- 1292 *Mesozoic and Cenozoic* 2:0. DOI: 10.1144/CEV2P.2.
- 1293 Popov E V, Delsate D, Felten R. 2019. A New Callorhynchid Genus (Holocephali, Chimaeroidei)
- 1294 from the Early Bajocian of Ottange-Rumelange, on the Luxembourg-French Border.
- 1295 *Paleontological Research* 23:220–230. DOI: 10.2517/2018PR021.
- 1296 Pyenson ND, Kelley NP, Parham JF. 2014. Marine tetrapod macroevolution: Physical and
- 1297 biological drivers on 250 million years of invasions and evolution in ocean ecosystems.



- 1298 *Palaeogeography, Palaeoclimatology, Palaeoecology* 400:1–8. DOI:  
1299 10.1016/j.palaeo.2014.02.018.
- 1300 R Core Team. 2016. R: A language and environment for statistical computing.
- 1301 Roberts AJ, Druckenmiller PS, Cordonnier B, Delsett LL, Hurum JH. 2020. A new plesiosaurian  
1302 from the Jurassic-Cretaceous transitional interval of the Slottsmøya Member (Volgian),  
1303 with insights into the cranial anatomy of cryptoclidids using computed tomography. *PeerJ*  
1304 2020. DOI: 10.7717/peerj.8652.
- 1305 Roberts AJ, Druckenmiller PS, Delsett LL, Hurum JH. 2017. Osteology and relationships of  
1306 *Colymbosaurus* Seeley, 1874, based on new material of *C. svalbardensis* from the  
1307 Slottsmøya Member, Agardhfjellet Formation of central Spitsbergen. *Journal of Vertebrate*  
1308 *Paleontology* 4634:e1278381. DOI: 10.1080/02724634.2017.1278381.
- 1309 Sachs S, Klug C, Kear BP. 2019. Rare evidence of a giant pliosaurid-like plesiosaur from the  
1310 Middle Jurassic (lower Bajocian) of Switzerland. *Swiss Journal of Palaeontology* 138:337–  
1311 342. DOI: 10.1007/s13358-019-00200-9.
- 1312 Sadki D, Weis R, Braun P. 2020. Graphoceratidés (Ammonitina) de l’Aalénien moyen  
1313 (Jurassique) de Rumelange-Hutberg (Grand-Duché de Luxembourg). In: Di Cencio A,  
1314 Sadki D, Weis R eds. *Paléontologie au Luxembourg (2). Les ammonites de la Minette*.  
1315 Ferrantia, in press.
- 1316 Sander PM. 2000. Ichthyosauria: their diversity, distribution, and phylogeny. *Paläontologische*  
1317 *Zeitschrift* 74:1–35.
- 1318 Sander PM, Bucher H. 1993. An ichthyosaur from the uppermost Toarcian of southern France.  
1319 *Neues Jahrbuch für Geologie und Paläontologie. Monatshefte* 1993:631–640.
- 1320 Sato T, Wu XC. 2008. A new Jurassic pliosaur from Melville Island, Canadian Arctic  
1321 Archipelago. *Canadian Journal of Earth Sciences* 45:303–320. DOI: 10.1139/E08-003.
- 1322 Schintgen T, Förster A. 2013. Geology and basin structure of the Trier-Luxembourg Basin –  
1323 implications for the existence of a buried Rotliegend graben. *Zeitschrift der Deutschen*  
1324 *Gesellschaft für Geowissenschaften* 164:615–637. DOI: 10.1127/1860-1804/2013/0025.
- 1325 Schmitz L. 2005. The taxonomic status of *Mixosaurus nordenskioeldii* (Ichthyosauria). *Journal*  
1326 *of Vertebrate Paleontology* 25:983–985.
- 1327 Schwermann L, Sander PM. 2011. Osteologie und Phylogenie von *Westphaliasaurus*  
1328 *simonsensii*: Ein neuer Plesiosauride (Sauropterygia) aus dem Unteren Jura (Pliensbachium)  
1329 von Sommersell (Kreis Höxter), Nordrhein-Westfalen, Deutschland. *Geologie und*  
1330 *Paläontologie in Westfalen* 79:5–56.
- 1331 Sciau J, Crochet J-YY, Mattei J. 1990. Le premier squelette de plésiosaure de France sur le  
1332 Causse du Larzac (Toarcien, Jurassique inférieur). *Geobios* 23:111–116. DOI:  
1333 10.1016/0016-6995(90)80021-7.
- 1334 Seeley HG. 1874a. Note on some of the generic modifications of the plesiosaurian pectoral arch.  
1335 *Quarterly Journal of the Geological Society* 30:436–449. DOI:  
1336 10.1144/GSL.JGS.1874.030.01-04.48.
- 1337 Seeley HG. 1874b. On *Murænosaurus Leedsii*, a Plesiosaurian from the Oxford Clay. Part I.  
1338 *Quarterly Journal of the Geological Society of London* 30:197–208. DOI:  
1339 10.1144/GSL.JGS.1874.030.01-04.35.
- 1340 Seeley HG. 1874c. On the pectoral arch and fore limb of *Ophthalmosaurus*, a new  
1341 ichthyosaurian genus from the Oxford Clay. *Quarterly Journal of the Geological Society of*  
1342 *London* 30:696–707.
- 1343 Siehl A, Thein J. 1989. Minette-type ironstones. In: *Phanerozoic ironstones*. Geological Society

- of London, Special Publications, 175–193.
- Smith AS. 2013. Morphology of the caudal vertebrae in *Rhomaleosaurus zetlandicus* and a review of the evidence for a tail fin in Plesiosauria. *Paludicola* 9:144–158.
- Smith AS. 2015. Reassessment of “*Plesiosaurus*” *megacephalus* (Sauropterygia: Plesiosauria) from the Triassic-Jurassic boundary, UK. *Palaeontologia Electronica* 1820:1–19.
- Smith AS, Araújo R. 2017. *Thaumatodracon wiedenrothi*, a morphometrically and stratigraphically intermediate new rhomaleosaurid plesiosaurian from the Lower Jurassic (Sinemurian) of Lyme Regis. *Palaeontographica Abteilung A* 308:89–125. DOI: 10.1127/pala/308/2017/89.
- Smith AS, Benson RBJ. 2014. Osteology of *Rhomaleosaurus thorntoni* (Sauropterygia: Rhomaleosauridae) from the Lower Jurassic (Toarcian) of Northamptonshire, England. *Monographs of the Palaeontographical Society* 168:1–40. DOI: 10.1080/02693445.2014.11963953.
- Smith AS, Dyke GJ. 2008. The skull of the giant predatory pliosaur *Rhomaleosaurus cramptoni*: implications for plesiosaur phylogenetics. *Naturwissenschaften* 95:975–980. DOI: 10.1007/s00114-008-0402-z.
- Smith AS, Vincent P. 2010. A new genus of pliosaur (Reptilia: Sauropterygia) from the Lower Jurassic of Holzmaden, Germany. *Palaeontology* 53:1049–1063. DOI: 10.1111/j.1475-4983.2010.00975.x.
- Storrs GW. 1997. Morphological and taxonomic clarification of the genus *Plesiosaurus*. In: *Ancient marine reptiles*. 145–190.
- Swinton WE. 1930. On a new species of *Eurhinosaurus* in the British Museum. *Geological Magazine* 67:275–277.
- Taylor M. 1992. Functional anatomy of the head of the large aquatic predator *Rhomaleosaurus zetlandicus* (Plesiosauria, Reptilia) from the Toarcian (Lower Jurassic) of Yorkshire, England. *Philosophical Transactions of the Royal Society B: Biological Sciences* 335:247–280. DOI: 10.1098/rstb.1992.0022.
- Taylor MA. 1997. Before the Dinosaur: The historical significance of the fossil marine reptiles. In: Callaway JM, Nicholls EL eds. *Ancient Marine Reptiles*. San Diego California: Academic Press, xix–xlvi.
- Tennant JP, Mannion PD, Upchurch P. 2016. Sea level regulated tetrapod diversity dynamics through the Jurassic/Cretaceous interval. *Nature Communications* 7:12737. DOI: 10.1038/ncomms12737.
- Tennant JP, Mannion PD, Upchurch P, Sutton MD, Price GD. Biotic and environmental dynamics through the Late Jurassic – Early Cretaceous transition : evidence for protracted faunal and ecological turnover. *Biological Reviews*. DOI: 10.1111/brv.12255.
- Teyssen TAL. 1984. Sedimentology of the Minette oolitic ironstones of Luxembourg and Lorraine: a Jurassic subtidal sandwave complex. *Sedimentology* 31:195–2011.
- Thorne PM, Ruta M, Benton MJ. 2011. Resetting the evolution of marine reptiles at the Triassic-Jurassic boundary. *Proceedings of the National Academy of Sciences of the United States of America* 108:8339–8344. DOI: 10.1073/pnas.1018959108.
- Valenciennes MA. 1861. D’un nouveau reptile très voisin du genre Ichthyosaure, trouvé dans l’argile du Kimméridge de Bléville, au nord du cap de la Hève du Havre. *Comptes rendus hebdomadaires des Séances de l’Académie des Sciences* 53:999–1001.
- Vincent P. 2011. A re-examination of *Hauffiosaurus zanoni*, a pliosauroid from the Toarcian (Early Jurassic) of Germany. *Journal of Vertebrate Paleontology* 31:340–351. DOI:

- 10.1080/02724634.2011.550352.
- Vincent P, Bardet N, Morel N. 2007. An elasmosaurid plesiosaur from the Aalenian (Middle Jurassic) of Western France. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 243:363–370.
- Vincent P, Martin JE, Fischer V, Suan G, Khalloufi B, Sucheras-Marx B, Lena A, Janneau K, Rousselle B, Rulleau L, Suchéras-Marx B, Léna A, Janneau K, Rousselle B, Rulleau L. 2013. A marine vertebrate fauna from the Toarcian-Aalenian succession of southern Beaujolais, Rhône, France. *Geological Magazine* 150:822–834. DOI: doi:10.1017/S0016756812000982.
- Vincent P, Storrs GW. 2019. *Lindwurmia*, a new genus of Plesiosauria (Reptilia: Sauropterygia) from the earliest Jurassic of Halberstadt, northwest Germany. *Science of Nature* 106. DOI: 10.1007/s00114-018-1600-y.
- Vincent P, Taquet P, Fischer V, Bardet N, Falconnet J, Godefroit P. 2014. Mary Anning’s legacy to French vertebrate palaeontology. *Geological Magazine* 151:7–20. DOI: 10.1017/S0016756813000861.
- Vincent P, Weis R, Kronz G, Delsate D. 2017. *Microcleidus melusinae*, a new plesiosaurian (Reptilia, Plesiosauria) from the Toarcian of Luxembourg. *Geological Magazine*:In Press. DOI: 10.1017/S0016756817000814.
- Weis R, Mariotti N. 2007. A belemnite fauna from the Aalenian-Bajocian boundary beds of the Grand Duchy of Luxembourg (NE Paris Basin). *Bollettino della Societa Paleontologica Italiana* 46:149–174.
- Weis R, Mariotti N, Riegraf W. 2012. The belemnite family Holcobelidae (Coleoidea) in the European Jurassic: systematics, biostratigraphy, palaeobiogeography and evolutionary trends. *Palaeodiversity* 5:13–49.
- Weis R, Sadki D, Mariotti N. 2017. Aalenian-Bajocian belemnites from the Middle and High Atlas, Morocco: taxonomy, biostratigraphy and palaeobiogeographical affinities. *Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen* 284:215–240. DOI: 10.1127/njgpa/2017/0659.
- White TE. 1940. Holotype of *Plesiosaurus longirostris* Blake and classification of the plesiosaurs. *Journal of Paleontology* 14:451–467.
- Wilberg EW. 2015. A new metriorhynchoid (Crocodylomorpha, Thalattosuchia) from the Middle Jurassic of Oregon and the evolutionary timing of marine adaptations in thalattosuchian crocodylomorphs. *Journal of Vertebrate Paleontology* 902846. DOI: 10.1080/02724634.2014.902846.
- Wilberg EW, Turner AH, Brochu CA. 2019. Evolutionary structure and timing of major habitat shifts in Crocodylomorpha. *Scientific Reports* 9:1–10. DOI: 10.1038/s41598-018-36795-1.
- Young MT, de Andrade MB. 2009. What is *Geosaurus*? Redescription of *Geosaurus giganteus* (Thalattosuchia: Metriorhynchidae) from the Upper Jurassic of Bayern, Germany. *Zoological Journal of the Linnean Society* 157:551–585.
- Young MT, Brandalise de Andrade M, Cornée J-J, Steel L, Foffa D. 2014a. Re-description of a putative Early Cretaceous “teleosaurid” from France, with implications for the survival of metriorhynchids and teleosaurids across the Jurassic-Cretaceous Boundary. *Annales de Paléontologie* 100:165–174. DOI: 10.1016/j.annpal.2014.01.002.
- Young MT, Brusatte SL, de Andrade MB, Desojo JB, Beatty BL, Steel L, Fernández MS, Sakamoto M, Ruiz-Omeñaca JI, Schoch RR. 2012. The cranial osteology and feeding ecology of the metriorhynchid crocodylomorph genera *Dakosaurus* and *Plesiosuchus* from

- the Late Jurassic of Europe. *PLoS ONE* 7:e44985. DOI: 10.1371/journal.pone.0044985.
- Young MT, Steel L, Brusatte SL, Foffa D, Lepage Y. 2014b. Tooth serration morphologies in the genus *Machimosaurus* (Crocodylomorpha, Thalattosuchia) from the Late Jurassic of Europe. *Royal Society Open Science* 1:1–14. DOI: 10.1098/rsos.140269.
- Zverkov NG, Arkhangelsky MS, Pardo-Pérez J, Beznosov PA. 2015. On the Upper Jurassic ichthyosaur remains from the Russian North. *Proceedings of the Zoological Institute RAS* 319:81–97.
- Zverkov NG, Efimov VM. 2019. Revision of *Undorosaurus*, a mysterious Late Jurassic ichthyosaur of the Boreal Realm. *Journal of Systematic Palaeontology*:in press. DOI: 10.1080/14772019.2018.1515793.
- Zverkov NG, Fischer V, Madzia D, Benson RBJ. 2018. Increased Pliosaurid Dental Disparity Across the Jurassic – Cretaceous Transition. *Palaeontology* 61:825–846. DOI: 10.1111/pala.12367.
- Zverkov NG, Jacobs ML. 2020. Revision of *Nannopterygius* (Ichthyosauria: Ophthalmosauridae): reappraisal of the ‘inaccessible’ holotype resolves a taxonomic tangle and reveals an obscure ophthalmosaurid lineage with a wide distribution. *Zoological Journal of the Linnean Society*:1–48. DOI: 10.1093/zoolinnean/zlaa028.
- Zverkov NG, Prilepskaya NE. 2019. A prevalence of *Arthropterygius* (Ichthyosauria: Ophthalmosauridae) in the Late Jurassic—earliest Cretaceous of the Boreal Realm. *PeerJ* 7:e6799. DOI: 10.7717/peerj.6799.

# Figure Captions

**Figure 1. Number of collections of marine reptiles (Ichthyosauria, Plesiosauria, Thalattosuchia, Pleuroosauria, Angolachelonia) per stage.** Data extracted from the paleobiology database on the 26th March 2020 (see Acknowledgements for the main contributors of these data). Silhouettes originate from phylopic: *Dakosaurus* by Dmitry Bogdanov & T Michael Keesey; *Meyerasaurus*, *Eurhinosaurus*, *Temnodontosaurus*, *Plesiopterys*, and *Ophthalmosaurus* by Gareth Monger; *Rhomaleosaurus* and *Stenopterygius* by Scott Hartmann; *Peloneustes* by Nobu Tamura & T Michael Keesey; *Albertonectes* by Frank Denota.

**Figure 2. Map of the fossiliferous localities investigated.**

**Figure 3. Thoursense Zone fauna, late Toarcian, Luxembourg.** Selected anatomy of specimen MNHNL TM212. (A–B) left coracoid in ventral (A) and anteroventral (B) views. (C, E, F) left humerus in anterodorsal (C), distal (E), and proximal (F) views. (D) left forefin in ventral view.

**Figure 4. Pseudoradosa Zone fauna, late Toarcian, Luxembourg.** (A–E) rhomaleosaurid left humerus MNHNL DOU307 in ventral (A), anterior (B), dorsal (C), proximal (D), and distal (E) views. (F–H) rhomaleosaurid humerus MNHNL KA109 in dorsal/ventral (F), proximal (H), and distal (H) views.

**Figure 5. Ichthyosaurians of the late Toarcian of Belgium and Luxembourg.** (A–B) large, *Temnodontosaurus*-like (Parvipelvia indet.) caudal centrum IRSNB R 436 in anterior (A) and lateral (B) views. (C–D) probable thunnosaurian caudal centrum IRSNB R 437 in anterior (C) and dorsal (D) views. (E–F) ichthyosaurian distal tooth IRSNB R 438 in labial (E) and mesial (F) views. (G–H) probable non-ophthalmosaurid (Parvipelvia indet.) left angular IRSNB R 439 in lateral (G) and medial (H) views. (I–K) ichthyosaurian right quadrate IRSNB R 440 in lateral (I), medial (J), and posterior (K) views.

**Figure 6. Ichthyosaurians of the Aalensis Zone, late Toarcian, Luxembourg.** (A–C) left coracoid of a non-baracromian parvipelvia MNHNL DOU353 in ventral (A), lateral (B), and posterior (C) views. (D–F) large parvipelvia caudal centrum MNHNL DOU378 in dorsal (D), anterior (E), and cross-sectional (F) views. (G) small parvipelvia caudal centrum MNHNL DOU998 in anteroventral view. (H–I) parvipelvia dorsal centrum MNHNL DOU944 in anterior (H), and lateral (I) views.

**Figure 7. Plesiosaurians of the Aalensis Zone, late Toarcian, Luxembourg.** (A) plesiosaurian tooth crown MNHNL DOU906 in ?mesial view. (B) rhomaleosaurid right humerus MNHNL

DOU558 in dorsal view. (C–F) rhomaleosaurid right humerus MNHNL DOU324a in proximal (C), dorsal (D), posterior (E), and distal (F) views. (G) juvenile rhomaleosaurid propodial MNHNL DOU324b.

**Figure 8. Plesiosaurian centra of the Aalensis Zone, late Toarcian, Luxembourg.** (A–B) rhomaleosaurid pectoral vertebra MNHNL DOU954 in anterior (A) and ventral (B) views. (C–G) cf. *Microcleidus* cervical vertebra MNHNL DOU978 in anterior (C), ventral (D), dorsal (E), lateral (F), and oblique (G) views. (H–J) rhomaleosaurid pectoral vertebra MNHNL DOU723 in anterior (H), lateral (I), and posterior (J) views. (K–N) plesiosaurian sacral vertebra MNHNL DOU724 in dorsal (K), anterior (L), posterolateral (M), and ventral (N) views. (O–Q) rhomaleosaurid pectoral vertebra MNHNL DOU722 in anterior (O), lateral (P), and posterior (Q) views.

**Figure 9. Late Aalenian fauna, Luxembourg.** (A–D) non-cryptoclidid plesiosauroid cervical centrum MNHNL BU157 in anterior (A), dorsal (B), ventral (C), and oblique (D) views. (E–G) plesiosaur juvenile caudal centrum MNHNL HU384 in lateral (E), anterior (F), and anteroventral (G) views. (H–I) fragmentary ichthyosaurian rostrum MNHN HU242.


**Figure 10. Humphresianum Zone fauna, early Bajocian, Luxembourg.** (A–F) cryptoclidid propodial MNHNL BM782 in anterior (A), proximal (B), dorsal (C), ventral (D), posterior (E), and distal (F) views. (G–H) parvipelvian MNHNL BM392 centra (G) and sclerotic element in lateral view (H). (I–O) ophthalmosaurid surangular and teeth MNHNL BM780\_781: anterior tooth in labial (I) and basal (J) views; mid-rostrum tooth in labial (K) and basal (L) views; posterior tooth in labial view (M); right angular in lateral (N) and medial (O) views. Abbreviation: PAE: posterior accessory epipodial element.

**Figure 11. The new Humphresianum Zone ophthalmosaurid, early Bajocian, Luxembourg.** Selected anatomy of specimen MNHNL BM779. (A–C) right exoccipital in anterolateral (A), posteromedial (B), and posterior (C) views. (D–E) suraoccipital in posterior (D) and ventral (E) views. (F–H) left scapula in anterior (F), medial (G), and lateral (H) views. (I–K) right quadrate in lateral (I), anterior (J), and condylar (K) views. (L) right parietal in dorsal view. (M–N) right humerus in posterior (M) and dorsal (N) views. (O–P) right ulna in posterior (O) and dorsal (P) views.

# Figure 1


Number of collections of marine reptiles (Ichthyosauria, Plesiosauria, Thalattosuchia, Pleurosauria, Angolachelonia) per stage.

Data extracted from the paleobiology database on the 26th March 2020 (see Acknowledgements for the main contributors of these data). Silhouettes originate from phylopic.org , Licensed under CC BY 3.0 SA: Dakosaurus by Dmitry Bogdanov & T Michael Keesey; Meyerasaurus, Eurhinosaurus, Temnodontosaurus, Plesiopterys, and Ophthalmosaurus by Gareth Monger; Rhomaleosaurus and Stenopterygius by Scott Hartmann; Peloneustes by Nobu Tamura & T Michael Keesey; Albertonectes by Frank Denota (CC 1.0).



# Figure 2

Map of the fossiliferous localities investigated.






# Figure 3

Thoursense Zone fauna, late Toarcian, Luxembourg.


Selected anatomy of specimen MNHNL TM212. (A-B) left coracoid in ventral (A) and anteroventral (B) views. (C, E, F) left humerus in anterodorsal (C), distal (E), and proximal (F) views. (D) left forefin in ventral view.

Thouarsense Zone, late Toarcian


A Left coracoid, ventral view




B Left coracoid, anteroventral view





C Left humerus, anterodorsal view




D Left forefin, ventral view



E Left humerus, distal view



F Left humerus, proximal view




50 mm

# Figure 4

Pseudoradosa Zone fauna, late Toarcian, Luxembourg.

(A-E) rhomaleosaurid left humerus MNHNL DOU307 in ventral (A), anterior (B), dorsal (C), proximal (D), and distal (E) views. (F-H) rhomaleosaurid humerus MNHNL KA109 in dorsal/ventral (F), proximal (H), and distal (H) views.

Pseudoradosa Zone, late Toarcian




# Figure 5

Ichthyosaurians of the Aalensis Zone, late Toarcian of Luxembourg and Belgium.

(A-B) large, *Temnodontosaurus*-like (*Parvipelvia* indet.) caudal centrum IRSNB R 436 in anterior (A) and lateral (B) views. (C-D) probable thunnosaurian caudal centrum IRSNB R 437 in anterior (C) and dorsal (D) views. (E-F) ichthyosaurian distal tooth IRSNB R 438 in labial (E) and mesial (F) views. (G-H) probable non-opthalmosaurid (*Parvipelvia* indet.) left angular IRSNB R 439 in lateral (G) and medial (H) views. (I-K) ichthyosaurian right quadrate IRSNB R 440 in lateral (I), medial (J), and posterior (K) views.



Dispansum-Pseudoradosa-Aalensis Zone, late Toarcian




# Figure 6

Ichthyosaurians of the Aalensis Zone, late Toarcian, Luxembourg.

(A–C) left coracoid of a non-baracromian parvipelvian MNHNL DOU353 in ventral (A), lateral (B), and posterior (C) views. (D–F) large parvipelvian caudal centrum MNHNL DOU378 in dorsal (D), anterior (E), and cross-sectional (F) views. (G) small parvipelvian caudal centrum MNHNL DOU998 in anteroventral view. (H–I) parvipelvian dorsal centrum MNHNL DOU944 in anterior (H), and lateral (I) views.

Aalensis Zone, late Toarcian






# Figure 7

Plesiosaurs of the Aalensis Zone, late Toarcian, Luxembourg.

(A) plesiosaurian tooth crown MNHNL DOU906 in ?mesial view. (B) rhomaleosaurid right humerus MNHNL DOU558 in dorsal view. (C-F) rhomaleosaurid right humerus MNHNL DOU324a in proximal (C), dorsal (D), posterior (E), and distal (F) views. (G) juvenile rhomaleosaurid propodial MNHNL DOU324b.


Aalensis Zone, late Toarcian



# Figure 8

Plesiosaurian centra of the Aalensis Zone, late Toarcian, Luxembourg.

(A-B) rhomaleosaurid pectoral vertebra MNHNL DOU954 in anterior (A) and ventral (B) views. (C-G) cf. *Microcleidus* cervical vertebra MNHNL DOU978 in anterior (C), ventral (D), dorsal (E), lateral (F), and oblique (G) views. (H-J) rhomaleosaurid pectoral vertebra MNHNL DOU723 in anterior (H), lateral (I), and posterior (J) views. (K-N) plesiosaurian sacral vertebra MNHNL DOU724 in dorsal (K), anterior (L), posterolateral (M), and ventral (N) views. (O-Q) rhomaleosaurid pectoral vertebra MNHNL DOU722 in anterior (O), lateral (P), and posterior (Q) views.






# Figure 9

Late Aalenian fauna, Luxembourg.

(A-D) non-cryptoclidid plesiosauroid cervical centrum MNHNL BU157 in anterior (A), dorsal (B), ventral (C), and oblique (D) views. (E-G) plesiosaur juvenile caudal centrum MNHNL HU384 in lateral (E), anterior (F), and anteroventral (G) views. (H-I) fragmentary ichthyosaurian rostrum MNHN HU242.




# Figure 10

Humphresianum Zone fauna, early Bajocian, Luxembourg.

(A-F) cryptoclidid propodial MNHNL BM782 in anterior (A), proximal (B), dorsal (C), ventral (D), posterior (E), and distal (F) views. (G-H) parvipelvian MNHNL BM392 centra (G) and sclerotic element in lateral view (H). (I-O) ophthalmosaurid surangular and teeth MNHNL BM780\_781: anterior tooth in labial (I) and basal (J) views; mid-rostrum tooth in labial (K) and basal (L) views; posterior tooth in labial view (M); right angular in lateral (N) and medial (O) views. Abbreviation: PAE: posterior accessory epipodial element.

Humphriesianum Zone, early Bajocian



# Figure 11

The new Humphresianum Zone ophthalmosaurid, early Bajocian, Luxembourg.

Selected anatomy of specimen MNHNL BM779. (A-C) right exoccipital in anterolateral (A), posteromedial (B), and posterior (C) views. (D-E) suraoccipital in posterior (D) and ventral (E) views. (F-H) left scapula in anterior (F), medial (G), and lateral (H) views. (I-K) right quadrate in lateral (I), anterior (J), and condylar (K) views. (L) right parietal in dorsal view. (M-N) right humerus in posterior (M) and dorsal (N) views. (O-P) right ulna in posterior (O) and dorsal (P) views.



Humphriesianum Zone, early Bajocian



**Table 1** (on next page)

Centra dimensions (in millimeters).

1 **Table 1. Centra dimensions (in millimeters).**

Specimen	Clade	Age	Axial_zone	Height	Length	Width
ULgPA35961	Ichthyosauria	Toarcian	dorsal	78	35	86
IRSNB_R_436	Ichthyosauria	Toarcian	caudal	118	48	118
IRSNB_R_437	Ichthyosauria	Toarcian	caudal	62	22	60
IRSNB_Vert-06462-0005	Ichthyosauria	Toarcian	cervical	63	24	64
IRSNB_Vert-11312-00007	Plesiosauria	Toarcian	dorsal	67	44	81
MNHNL_DOU378	Ichthyosauria	Toarcian	caudal	190	65	NA
MNHNL_DOU978	Plesiosauria	Toarcian	cervical	41	55	53
MNHNL_DOU944	Ichthyosauria	Toarcian	dorsal	82	36	NA
MNHNL_DOU998	Ichthyosauria	Toarcian	sacral	90	31	NA
MNHNL_DOU722	Plesiosauria	Toarcian	dorsal	50	37	60
MNHNL_DOU723	Plesiosauria	Toarcian	dorsal	55	30	63
MNHNL_DOU954	Plesiosauria	Toarcian	pectoral	63	42	73
MNHNL_DOU724	Plesiosauria	Toarcian	sacral	39	40	57
MNHNL_DOU369a	Ichthyosauria	Toarcian	dorsal	68	32	NA
MNHNL_DOU369b	Ichthyosauria	Toarcian	caudal	61	35	NA
MNHNL_DOU369c	Ichthyosauria	Toarcian	caudal	63	32	NA
MNHNL_HU384	Plesiosauria	Aalenian	caudal	32	NA	NA
MNHNL_BU157	Plesiosauria	Aalenian	cervical	67	62	71
MNHNL_TM212	Ichthyosauria	Toarcian	cervical	62	32	56
MNHNL_BM779	Ichthyosauria	Bajocian	caudal	91	39	NA
MNHNL_BM758	Ichthyosauria	Bajocian	caudal	100	39	NA
MNHNL_BM725a	Ichthyosauria	Bajocian	cervical	83	38	NA
MNHNL_BM725b	Ichthyosauria	Bajocian	caudal	56	26	NA
MNHNL_BM461	Ichthyosauria	Bajocian	cervical	72	30	NA
MNHNL_BM392a	Ichthyosauria	Bajocian	caudal	72	30	NA
MNHNL_BM392b	Ichthyosauria	Bajocian	caudal	77	31	NA
MNHNL_BM392c	Ichthyosauria	Bajocian	caudal	72	29	NA
MNHNL_BM392d	Ichthyosauria	Bajocian	caudal	75	32	NA
MNHNL_BM779	Ichthyosauria	Bajocian	cervical	87	35	90

2

## **Table 2**(on next page)

List of the material and taxonomic identifications.

1 Table 2. List of the material and taxonomic identifications.

Specimen	Clade	Stage	Zone	Element	Identification	Note	Figured
MNHNL TMP212	Ichthyosauria	Toarcian	Thouarsense	Partial skeleton	Baracromia indet.	Non-opthalmosaurid	Figure 3
MNHNL DOU307	Plesiosauria	Toarcian	Pseudoradosa	Humerus	Rhomaleosauridae indet.		Figure 4 A-E
MNHNL KA109	Plesiosauria	Toarcian	Pseudoradosa	Humerus	Rhomaleosauridae indet.		Figure 4 F-H
ULgPA35961	Ichthyosauria	Toarcian	Dispansum to Aalensis	Dorsal centrum	Parvipelvia indet.		No
IRSNB Vert-06455-0001	Ichthyosauria	Toarcian	Dispansum to Aalensis	Centrum	Ichthyosauria indet.		No
IRSNB Vert-00000-00801	Ichthyosauria	Toarcian	Dispansum to Aalensis	Centrum	Ichthyosauria indet.		No
IRSNB Vert-06659-0001	Ichthyosauria	Toarcian	Dispansum to Aalensis	Centrum	Ichthyosauria indet.		No
IRSNB Vert-00000-00800	Ichthyosauria	Toarcian	Dispansum to Aalensis	Left surangular	Parvipelvia indet.		No
IRSNB Vert-00000-00804	Ichthyosauria	Toarcian	Dispansum to Aalensis	Dorsal centrum	Parvipelvia indet.	<i>Temnodontosaurus</i> -like	No
IRSNB Vert-06462-0005	Ichthyosauria	Toarcian	Dispansum to Aalensis	Cervical centrum	Parvipelvia indet.		No
IRSNB R 436	Ichthyosauria	Toarcian	Dispansum to Aalensis	Caudal centrum	Parvipelvia indet.	<i>Temnodontosaurus</i> -like	Figure 5 A-B
IRSNB R 437	Ichthyosauria	Toarcian	Dispansum to Aalensis	Caudal centrum	Parvipelvia indet.	Possible thunnosaurian	Figure 5 C-D
IRSNB R 438	Ichthyosauria	Toarcian	Dispansum to Aalensis	Tooth	Ichthyosauria indet.		Figure 5 E-F
IRSNB R 439	Ichthyosauria	Toarcian	Dispansum to Aalensis	Left angular	Parvipelvia indet.		Figure 5 G-H
IRSNB R 440	Ichthyosauria	Toarcian	Dispansum to Aalensis	Right quadrate	Ichthyosauria indet.		Figure 5 I-K
IRSNB Vert-11312-00007	Plesiosauria	Toarcian	Dispansum to Aalensis	Dorsal centrum	Plesiosauria indet.		No
MNHNL DOU369	Ichthyosauria	Toarcian	Aalensis	Centra	Parvipelvia indet.	Probable non-thunnosaurian	No
MNHNL DOU353	Ichthyosauria	Toarcian	Aalensis	Coracoid	Parvipelvia indet.	Non-baracromian	Figure 6 A-C
MNHNL DOU378	Ichthyosauria	Toarcian	Aalensis	Posterior dorsal to anterior caudal centrum	Parvipelvia indet.	Probable non-thunnosaurian	Figure 6 D-F
MNHNL DOU998	Ichthyosauria	Toarcian	Aalensis	Caudal centrum	Parvipelvia indet.		Figure 6 G
MNHNL DOU944	Ichthyosauria	Toarcian	Aalensis	Dorsal centrum	Parvipelvia indet.		Figure 6 H-I
MNHNL DOU906	Plesiosauria	Toarcian	Aalensis	Tooth crown	Plesiosauria indet.		Figure 7 A
MNHNL DOU558	Plesiosauria	Toarcian	Aalensis	Humerus	Rhomaleosauridae indet.		Figure 7 B

<b>MNHNL DOU324 A</b>	Plesiosauria	Toarcian	Aalensis	Humerus	Rhomaleosauridae indet.		Figure 7 C- F
<b>MNHNL DOU324 B</b>	Plesiosauria	Toarcian	Aalensis	Humerus	Rhomaleosauridae indet.	Juvenile	Figure 7 G
<b>MNHNL DOU954</b>	Plesiosauria	Toarcian	Aalensis	Posterior pectoral/anterior dorsal	Rhomaleosauridae indet.		Figure 8 A- B
<b>MNHNL DOU978</b>	Plesiosauria	Toarcian	Aalensis	Cervical centrum	cf. Microcleidus		Figure 8 C- G
<b>MNHNL DOU723</b>	Plesiosauria	Toarcian	Aalensis	Pectoral vertebra	Rhomaleosauridae indet.		Figure 8 H- J
<b>MNHNL DOU724</b>	Plesiosauria	Toarcian	Aalensis	Sacral vertebra	Plesiosauria indet.		Figure 8 K- N
<b>MNHNL DOU722</b>	Plesiosauria	Toarcian	Aalensis	Pectoral vertebra	Rhomaleosauridae indet.		Figure 8 O- Q
<b>MNHNL BU157</b>	Plesiosauria	Aalenian	Concavum or Murchisonae	Cervical vertebra	Plesiosauroidea indet.	Non-cryptoclidid	Figure 9 A- D
<b>MNHNL HU384</b>	Plesiosauria	Aalenian	Murchisonae	Caudal centrum	Plesiosauria indet.	Juvenile	Figure 9 E- G
<b>MNHNL HU242</b>	Ichthyosauria	Aalenian	Concavum	Partial rostrum	Ichthyosauria indet.		Figure 9 H- I
<b>MNHNL BM360</b>	Ichthyosauria	Bajocian	Humphriesianum	Partial neural spine and ribs	Ichthyosauria indet.		No
<b>MNHNL BM758</b>	Ichthyosauria	Bajocian	Humphriesianum	Caudal centrum	Parvipelvina indet.		No
<b>MNHNL BM725</b>	Ichthyosauria	Bajocian	Humphriesianum	Cervical and caudal centra	Parvipelvina indet.		No
<b>MNHNL BM461</b>	Ichthyosauria	Bajocian	Humphriesianum	Cervical centrum	Parvipelvina indet.		No
<b>MNHNL BM782</b>	Plesiosauria	Bajocian	Humphriesianum	Propodial	Cryptoclididae indet.		Figure 10 A-F
<b>MNHNL BM392</b>	Ichthyosauria	Bajocian	Humphriesianum	Five caudal centrum and one sclerotic plate	Parvipelvina indet.		Figure 10 G-H
<b>MNHNL BM780_7 81</b>	Ichthyosauria	Bajocian	Humphriesianum	Surangular, teeth	Ophthalmosauridae indet.		Figure 10 I- O
<b>MNHNL BM779</b>	Ichthyosauria	Bajocian	Humphriesianum	Partial skeleton	Ophthalmosauridae indet.		Figure 11