

High expression of MAPK-14 promoting the death of chondrocytes is an important signal of osteoarthritis process

Zhiqin Deng^{Equal first author, 1}, Xiaotian Hu^{Equal first author, 1, 2}, Murad Alahdal¹, Jianquan Liu¹, Zhe Zhao¹, Xiaoqiang Chen¹, Junxiong Xie¹, Li Duan¹, Daping Wang^{Corresp., 1, 2}, Wencui Li^{Corresp. 1}

¹ Hand and Foot Surgery Department, Shenzhen Second People's Hospital (The First Hospital Affiliated to Shenzhen University), Shenzhen, Guangdong, China

² Anhui Medical University, Hefei, China

Corresponding Authors: Daping Wang, Wencui Li
Email address: wangdp@mail.sustech.edu.cn, 13923750767@163.com

Background Osteoarthritis (OA) is one of the most common degenerative diseases worldwide. Many researchers are studying the pathogenesis of OA, however, it is still unclear. **Methods** Screening and validation of OA relevant hub genes are an important part of exploring their potential molecular mechanism. Therefore, this study aims to explore and verify the mechanisms of hub genes in the OA by bioinformatics, qPCR, fluorescence and propidium iodide staining. **Results** Microarray datasets GSE43923, GSE55457 and GSE12021 were collected in the Gene Expression Omnibus (GEO), including 45 samples, which divided into 23 osteoarthritis knee joint samples and 22 samples of normal knee joint. Thereafter, 265 differentially expressed genes (DEGs) were identified in all, which divided into 199 upregulated genes and 66 downregulated genes. The hub genes MAPK-14, PTPRC, PTPN12 were upregulated, while B9D1 was downregulated. In order to further confirm the expression of screening differential genes in human chondrocytes, the human chondrocytes were extracted from a joint replacement surgery and stained with toluidine blue for identification. Compared with normal chondrocytes, OA chondrocytes had high expression of COL I protein and low expression of COL II protein. The expression levels of MAPK-14, PTPRC and PTPN12 in OA chondrocytes were significantly higher than the expression levels of B9D1 in normal chondrocytes. Moreover, the inflammatory necrosis of OA chondrocytes was increased compared with the normal chondrocytes by propidium iodide staining. **Conclusions** The high expression of MAPK-14 works as a promoter of chondrocytes death and an important signal of the osteoarthritis process.

High expression of MAPK-14 promoting the death of chondrocytes is an important signal of osteoarthritis process

Zhiqing, Deng ^{1, a}, Xiaotian Hu ^{1, 2, a}, Murad Alahdal ¹, Jianquan Liu¹, Zhe Zhao¹, Xiaoqiang Chen¹, Junxiong Xie¹, Li Duan ¹, Daping Wang ^{1, 2, *}, Wencui Li ^{1, *}

¹ Hand and Foot Surgery Department, Shenzhen Second People's Hospital (The First Hospital Affiliated to Shenzhen University), Shenzhen, China.

² Anhui Medical University, Hefei, Anhui province, China.

Corresponding Author:

* Daping Wang^{1, 2, *}, Wencui Li^{b, *}

Daping Wang, Shenzhen Second People's Hospital, 3002 Sungang West Road, Shenzhen 518000, Tel.: +86-0755-83791866, Fax: +86-0755-83366388, E-mail: dapingwang1963@qq.com.

Wencui Li, Shenzhen Second People's Hospital, 3002 Sungang West Road, Shenzhen 518000, Tel.: +86-0755-83366388, Fax: +86-0755-83366388, E-mail: 13923750767@163.com. ORCID ID: 0000-0003-2787-5360.

^a Zhiqin Deng, and Xiaotian Hu contributed equally to this work.

Disclosure of potential conflicts of interest: There are no potential conflicts of interest.

Abstract

Background Osteoarthritis (OA) is one of the most common degenerative diseases worldwide. Many researchers are studying the pathogenesis of OA, however, it is still unclear.

Methods Screening and validation of OA relevant hub genes are an important part of exploring their potential molecular mechanism. Therefore, this study aims to explore and verify the mechanisms of hub genes in the OA by bioinformatics, qPCR, fluorescence and propidium iodide staining.

Results Microarray datasets GSE43923, GSE55457 and GSE12021 were collected in the Gene Expression Omnibus (GEO), including 45 samples, which divided into 23 osteoarthritis knee joint samples and 22 samples of normal knee joint. Thereafter, 265 differentially expressed genes (DEGs) were identified in all, which divided into 199 upregulated genes and 66 downregulated genes. The hub genes MAPK-14, PTPRC, PTPN12 were upregulated, while B9D1 was downregulated. In order to further confirm the expression of screening differential genes in human chondrocytes, the human chondrocytes were extracted from a joint replacement surgery and stained with toluidine blue for identification. Compared with normal chondrocytes, OA chondrocytes had high expression of COL I protein and low expression of COL II protein. The expression levels of MAPK-14, PTPRC and PTPN12 in OA chondrocytes were significantly higher than the expression levels of B9D1 in normal chondrocytes. Moreover, the inflammatory necrosis of OA chondrocytes was increased compared with the normal chondrocytes by propidium iodide staining.

Conclusions The high expression of MAPK-14 works as a promoter of chondrocytes death and an important signal of the osteoarthritis process.

Key words: Osteoarthritis; MAPK-14, chondrocytes, differentially expressed genes; protein-protein interaction; Gene Ontology;

1 Introduction

Osteoarthritis is the most serious chronic bone disease and can cause severe pain and even disability in the elderly(Song et al. 2006). OA not only brings suffering to patients but also aggravates their economic burden. Nearly half of the elderly patients are receiving treatment, and most of them are still suffering or disabled(McKenzie & Torkington 2010). About one-third of elderly over 60 years old are supposed to be diagnosed with OA by 2030(Croft 2005). One of the most important risks in OA development is the pathogenic changes of hub genes (Hochberg et al. 2013). Heritability of osteoarthritis reached approximately 50% (Cibrian Uhalte et al. 2017). However, the molecular mechanism genetic changes in OA remains unknown.

However, current researches showed some disadvantages such as using small amount of genetic samples and the lack of relevant experimental verification. Therefore, the expanding of the number of screening sample size on the basis of the predecessors, optimizing the relevant screening conditions and carrying out the relevant verification through cell lines experiments could contribute to further understanding and enhances further study of the internal mechanism regarding the occurrence and development of osteoarthritis. Recently, microarray technology and bioinformatics have made significant advances which contribute to understand differentially expressed genes (DEGs) in OA and enable us to do further studies. Here, we got 3 datasets from Gene Expression Omnibus (GEO) for analysis and obtain DEGs between OA knee joint samples and normal knee joint samples. Based on this, we haven't only analyzed Gene Ontology (GO), Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment of DEGs, but also produced protein-protein interaction (PPI) network to help us unlock some of the unstudied parts of the molecular mechanisms of OA. The purpose of this research was to explore regarding underlying mechanisms of OA, so as to promote the further development of OA treatment.

2 Materials & Methods

2.1. Microarray Data Information

The Gene Expression Omnibus (GEO, <http://www.ncbi.nlm.nih.gov/geo>) is a genomic database for all users, from which users can get all kinds of data free of charge(Barrett et al. 2013).The expression profiles of GSE43923, GSE55457 and GSE12021 were all searched and got from the GEO. GSE43923 include 3 OA knee joint samples and 3 normal knee joint samples. GSE12021 include 10 OA knee joint samples and 9 normal knee joint samples, and GSE55457 include 10 OA knee joint samples and 10 normal knee joint samples.

2.2. Identification of DEGs

GEO2R (<http://www.ncbi.nlm.nih.gov/geo/geo2r>), a free public network platform, was used to process the DEGs between OA knee joint samples and normal knee joint samples. Using GEO2R, we have integrated and analyzed multiple datasets for identifying DEGs. $|\log FC| > 1$ and P-value < 0.05 were considered the cutoff criterion and have statistically significant.

2.3. GO and KEGG enrichment analyses of DEGs

DAVID(<http://david.ncifcrf.gov>), a free public online biological information database was used to provide biological function analysis for a large number of genes and can visualize the results(Huang et al. 2007). GO is a biological information analysis tool that can induce and analyze a large number of genes and visually export corresponding results, including BP, CC, and MF(Shannon et al. 2003). KEGG is a utility database resource for understanding advanced functions and biological systems (such as cells, organisms, and ecosystems), generating genomic sequencing and other high-throughput experimental technologies from molecular level information, especially large molecular datasets(Szklarczyk et al. 2011). Deep analyses were

implemented by using the DAVID online database. $P < 0.05$ was considered statistically significant.

2.4. PPI network construction and module analysis.

STRING(<https://string-db.org/cgi/input.pl>), a free public online search platform was used to elucidate the relationship between interacting genes (Franceschini et al. 2013). DEGs previously screened in this research were imported into STRING to product the PPI network, and interaction with a confidence score > 0.4 was considered significant. Cytoscape, a free and public source bioinformatics application that we imported the data into it. The built-in APP Molecular Complex Detection (MCODE) of Cytoscape is convenient to select the most significant modules in all DEGs and identify the most prominent gene in each module (Bader & Hogue 2003).

2.5 Isolation and culture of chondrocytes

Cartilage specimens of the fracture patients' femoral head and the knee joint of patients with OA in Shenzhen second people's hospital were collected to isolate and cultivate chondrocytes. The adherent cells were cultured in the chondrocyte medium, which contains the DMEM-LG (Gibco), 10% FBS (Gibco), 1% penicillin-streptomycin (Gibco), 1% HEPES, 1% vitamin solution, 1% proline solution and 1% non-essential amino acid NEAA. These adherent cells are what we call chondrocytes. chondrocytes were cultured in at 37°C and 5% CO_2 . The study was reviewed and approved by the First Affiliated Hospital of Shenzhen University Health Science Center Research Ethics Committee. All patients signed a written informed consent before participating in this study.

2.6 Toluidine blue staining of chondrocytes

First, we washed the Petri dishes with PBS for 2 times and fixed the cells with 4% paraformaldehyde directly for about 0.5 h (4°C). Then wash with tap water for 15 minutes, then with distilled water for 5 minutes, then add 1% toluidine blue for 2h. Finally, remove excess dye and filter impurities. The dyeing condition was observed under the microscope, dried after satisfaction, and sealed with neutral gum.

2.7 Immunofluorescence staining of chondrocytes

PBS washed the cells twice. The cells were fixed with paraformaldehyde for 15 min and washed with PBS for 6 times / 5min. 0.5% Triton-X100 was incubated and drilled for 15 min. Wash PBS 6 times / 5min. The cells were sealed with 5% BSA at room temperature for 2 h and cleaned with 0.5% BSA for 6 times / 5min. The primary antibody (Rabbit Anti-Collagen II/FITC Conjugated antibody (bs-10589R-FITC) and Rabbit Anti-Collagen I/Cy3 Conjugated antibody (bs-10423R-Cy3), Bioss, Beijing, China) were added and incubated at room temperature for 2h and washed with PBS for 6 times / 5min. DAPI staining was added for 5min. Photos were taken using a confocal microscope ZEISS LSM800, Germany.

2.8 Western Blot

OA chondrocytes and normal control were cultured for three days, and then it was taken out of the CO₂ incubator, cells were washed twice with PBS. Total protein was obtained by lysis buffer (Beyotime, China), and then placed on ice for 30 minutes. Centrifuged the sample at 12000g for 25 min at 4°C. The supernatants were transferred to a new EP tube. The protein concentration was measured by BCA quantitative method. Protein samples were electrophoresed on 10% SDS-PAGE gel. After that, protein samples were transferred on PVDF membranes

according to the three-character sandwich method. Use 5% skimmed milk powder to block for 2h. Use the primary antibody containing COL I (1:1000, boss, China), COL II (1:1000, Thermo Scientific, USA) and GAPDH (1:1000, abcam, USA) to co-incubate with the sample overnight at 4 °C. TBST cleaning, 5 times/10 min. Then use the second antibody containing goat anti-rabbit incubation bath for two hours at room temperature. After washing the unbound antibody, use chemiluminescence to take pictures in the two-color laser and chemiluminescence imaging system (Odyssey FC, Gene Company Limited, China).

2.9 Apoptosis assay of chondrocytes

The chondrocytes were isolated from the cartilage tissue, washed and then resuspended, and the precipitate was taken. About 50uL cell suspension was added into each flap-throwing tank and centrifuged in a flap-throwing centrifuge (1500 g, 10 min). The chondrocytes can be attached to a slide after the surrounding water is absorbed by the filter paper. Then wash with PBS for 3 times and fix with 4% paraformaldehyde for 15min. Wash again with PBS for 3 times, add PI immunofluorescence staining and incubate in a wet tank for 30 min. After washing with PBS again for 3 times, fluorescence microscope observation and image collection were performed.

2.10 qRT-PCR Validation and Statistical Analysis

After previous research, we've found four hub genes, including MAPK14, PTPRC, B9D1, PTPN12. We searched the primers of the four hub genes through the PrimerBank (<https://pga.mgh.harvard.edu/primerbank/>) and ordered the relevant primers through the Sangon Biotech. Quantitative real-time polymerase chain reaction primers are detailed in Table 1. Total

RNA was extracted from chondrocytes using the RNeasy mini kit (Takara), according to the manufacturer's instructions. The RNA was reversely transcribed into cDNA by iScript™ cDNA Synthesis Kit (Takara) and qRT-PCR was performed with a Real-time quantitative PCR instrument (Life Technologies).

2.11 Statistical analysis

Use SigmaPlot 10.0 to draw. Use IBM SPSS Statistics 19 to conduct statistical analysis, data are expressed as mean \pm std. A significant statistical difference was $P < 0.05$.

3 Results

3.1. Identification of DEGs in Osteoarthritis

A total of 23 OA knee joint samples and 22 matched normal knee joint samples were analyzed; taking $P < 0.05$ and $|\log FC| > 1$ as a threshold, DEGs (2555 in GSE43923, 1347 in GSE55457 and 1398 in GSE11201) were identified. First, we made a pairwise comparison and got three Venn graphs (Figure 1A, B, C). Then we comprehensively compared and plotted the final Venn diagram of the differentially expressed gene (Figure 1 D). A total of 265 common differential genes were screened, including 66 down-regulated genes and 199 up-regulated genes.

3.2 KEGG and GO enrichment analyses of DEGs

By using DAVID, the DEGs functions were classified into 3 main categories, including BP, MF, and CC (Figure 2). We can see that otherness in BP were significantly enriched in anatomical structure formation involved in morphogenesis, cell adhesion, vasculature development, blood vessel development (Figure 3A). Otherness in MF was mainly enriched in

calcium channel inhibitor activity, signaling receptor binding, growth factor activity, cytoskeletal protein binding and adenylate cyclase activator activity (Figure 3B). Otherness in CC was mainly enriched in the cytoplasmic part, cytoplasm and extracellular region (Figure 3C).

3.3 PPI network construction and module analysis

We use the data obtained through STRING and then build the PPI network through Cytoscape, containing 140 nodes and 356 edges (Figure 4). Next, we identified 4 hub genes, including MAPK14, PTPN12, PTPRC and B9D1. Then we analyzed the 4 modules using MCODE (the plug-in for Cytoscape), the four modules were shown in (Figure 5). Functional enrichment analyses were also carried out for these modules. Pathway enrichment analysis showed that Module 1 is the signaling pathways regulating pluripotency of stem cells and mainly associated with leukocyte transendothelial migration. Module 2 is mainly associated with the regulation of actin polymerization or depolymerization. Module 3 is mainly associated with the positive regulation of response to external stimulus and leukocyte migration. It also can response to lipopolysaccharide. Module 4 is mainly associated with the docking of the ciliary basal body-plasma membrane (Table 2).

3.4 Morphological observation and authenticate of normal chondrocytes and OA chondrocytes

When chondrocytes are completely adherent to the dish, they are mainly flat and polygonal. The nuclei are located in the center of the cell, with 2-3 nucleoli. The morphology of OA chondrocytes was more irregular, with rare cell community aggregation (Figure 6A).

The cytoplasm of chondrocytes stained with toluidine blue was blue, and 1-3 purplish blue nucleoli were observed (Figure 6B). The results showed that all the cells we cultured were chondrocytes.

3.5 Biological identification of OA and control chondrocytes

COL I, COL II and DAPI immunohistochemical tested were carried out on the Chondrocytes. And then we looked at it through a fluorescence microscope, the nuclei are dyed blue, collagen type I dyed red, collagen type II dyed green. Results showed that OA chondrocytes compared with normal chondrocytes, the COL I expression increased (Figure 6 C-H and O), and the expression of COL II reduced (Figure 6 I-N and O) detected by immunofluorescence and westernbolt. This indicates that OA chondrocytes mainly express collagen type I, which means that fibrochondrocytes are newly formed on the surface of cartilage, and the spontaneous depolymerization of fibrochondrocytes further accelerates the death of chondrocytes.

3.6 Validation of Hub Gene

4 hub genes have been found. Their details are shown in Table 3. We validated our results by measuring the relative expression of our hub genes using qRT-PCR. As can be seen, the relative expression of MAPK-14, PTPN12 and PTPRC in OA chondrocytes were significantly increased, while the expression of B9D1 were significantly decreased ($P < 0.05$) (Figure 7A-D), which was consistent with our expected results.

3.7 Apoptosis test of chondrocytes

The chondrocytes were stained with iodide propidium (PI) and examined by fluorescence microscope. Results showed that compared with normal chondrocytes, a large number of OA

chondrocytes were surrounded by pathological vesicles and showed intracellular red staining, which was the PI entering the cells (Figure 7E-J). It suggested that the cell membrane was incomplete and the cells death of OA chondrocytes was increased.

4 Discussion

The incidence of OA is a second after heart diseases in the statistical reports of some countries. OA, cardiovascular, cerebrovascular diseases and cancer are also called "three major killers" threatening human health. The most immediate symptom of OA is joint pain (Torres et al. 2006). A large number of epidemiological studies have confirmed that osteoarthritis is a complex disease involving multiple factors (Neogi & Zhang 2013). Therefore, the molecular mechanisms of OA are the most important direction we can study. It can help us to make early diagnosis and treatment of OA.

In previous researches on bioinformatic analysis, many of them are only limited in two cohort studies or two genetic events. This study makes up for the deficiency in this aspect, we used 3 datasets to obtain DEGs between OA knee joint samples and normal knee joint samples. This can reduce the contingency of analysis. 265 DEGs were identified in all, which divided into 199 upregulated genes and 66 downregulated genes.

GO and KEGG enrichment analyses were performed found that the upregulated genes were mainly enriched in calcium channel inhibitor activity, cytoskeletal protein binding, adenylate cyclase activator activity, while the downregulated genes were mainly enriched in vasculature development, and blood vessel development. According to previous studies, calcium channels have been shown to mediate the OA pain pathway (Rahman et al. 2015). In addition, some studies have confirmed that the cytoskeletal structure is incomplete and diseased in cartilage chondrocytes of OA(Blain 2009). Other researches had indicated a protective role for pituitary

adenylate cyclase-activating polypeptide (PACAP) in chondrocyte differentiation and metabolism (Strange-Vognsen et al. 1997), (Juhasz et al. 2014), (Juhasz et al. 2015), (Juhasz et al. 2015). In addition, some studies have pointed out that one of the most important members of osteogenesis is the skeletal vasculature and distinct blood vessel subtypes play different roles in osteogenesis (Sivan et al. 2019), (Cui et al. 2016). Moreover, the enriched KEGG pathways of DEGs and modules analysis included NF-kappa B signaling pathway, TNF signaling pathway, PI3K-Akt signaling pathway and MAPK signaling pathway. Previous studies showed that PI3K-Akt, NF-kappa B and TNF signaling pathways are involved in synovial hyperplasia and osteoarthritis cartilage degeneration (Fu et al. 2016), (Lai et al. 2014) and changes in the expression of cytokines, biological factors and enzymes on chondrocytes in human osteoarthritis may be caused by the p38-MAPK signaling pathway (Feng et al. 2017). These theories are also consistent with our results.

In this study, normal chondrocytes and OA chondrocytes were extracted from the collected cartilage specimens of the knee joint of patients with femoral head and OA. The cytoplasm of chondrocytes stained with toluidine blue was purplish blue, and 1-3 purplish blue nucleoli were observed. We found that the expression levels of MAPK-14, PTPN12 and PTPRC in OA chondrocytes were significantly increased, while the expression levels of B9D1 were significantly decreased by qPCR test. This indicates that the changes in the expression level of the potential key genes in the development of OA are consistent with the changes in the expression level of OA chondrocytes, which is consistent with our conjecture.

However, does the high expression of MAPK-14 predict chondrocyte death? The chondrocyte apoptosis experiment showed that the cell membrane of OA chondrocytes was more incomplete than that of normal chondrocytes, and pathological vesicles appeared, indicating that the amount

of necrosis of OA chondrocytes was significantly increased. It has been proved that chondrocytes are doomed to die in the development of OA, while high expression of MAPK-14 plays a role in inducing the autophagy of cells.

MAPK14 is one of the representatives of the MAPK family. Furthermore, the MAPK signaling pathway is the most important signaling system to mediate osteoarthritic cartilage damage (Chowdhury et al. 2008). Studies have shown that the major pathological changes of osteoarthritis are degradation of the extracellular matrix, leading to progressive loss of cartilage components and destruction of chondrocyte structure and function(Stoddart et al. 2009). Inflammatory factors and growth factors specifically bind to receptors on cell membranes to activate the intracellular MAPK signaling pathway, causing an increase in matrix metalloproteinases (MMPs) expression, chondrocyte apoptosis and cartilage destruction. MMPs can degrade almost all the extracellular matrix of chondrocytes and has been shown involved in the pathogenesis of OA(Murphy et al. 2002). In addition, MAPK signaling pathway also participates in and regulates chondrocyte enlargement, calcification and proliferation in the process of OA pathology (Huang et al. 2008). Therefore, the MAPK signaling pathway can be used as an effective target for the study of new drugs.

Other DEGs PRPRC and PTPN12 are members of the PTP family. On the basis of previous studies, an important part of the pathogenesis of OA is the immune system(Goldring & Otero 2011). The proliferation of T lymphocytes can regulate the immune response, and the biochemical activities of human articular cartilage can directly affect the proliferation of T lymphocytes(Pereira et al. 2016). T lymphocytes are lymphoid stem cells derived from bone marrow and distributed to immune systems throughout the body through lymph and blood circulation (Moro-Garcia et al. 2018). On the other hand, dendritic cells(DCs) as a crucial

component in triggering an immune response, as well as in inflammatory and autoimmune diseases(Ganguly et al. 2013). Furthermore, the PTP family is important in the activation of DCs ability to induce T cell activation and function (Rhee et al. 2014). Therefore, the PTP family and its related pathways may provide a new idea to treat OA.

The involvement of B9D1 in OA has not been well studied, but in recent research, mutations of B9D1 result in common skeleton defects(Yuan & Yang 2015). RB9D1 and its related pathways can be used as a supplement to the current traditional treatment of OA.

In brief, four potential key genes in the pathogenesis of OA were identified through bioinformatics screening, including MAPK-14, PTPRC, PTPN12 and B9D1. The imbalance of matrix synthesis and catabolism of OA chondrocytes and the increase of cell membrane destruction may be related to the high expression of MAPK-14 promoting the death of OA chondrocytes. The regulation of the expression of these key genes can provide a potential reference target for the treatment of OA.

Conclusions

The hub genes MAPK-14, PTPRC, PTPN12 were upregulated while B9D1 was downregulated after screening from the 265 differentially expressed genes. OA chondrocytes had high expression of COL I protein and low expression of COL II protein. The inflammatory necrosis of OA chondrocytes was increased compared with the normal chondrocytes. The expression levels of MAPK-14, PTPRC and PTPN12 in OA chondrocytes were significantly higher, while the expression levels of B9D1 were significantly lower than that in normal chondrocytes. The MAPK-14 among the four potential key genes has been proved to play an important role in the treatment of OA.

Acknowledgements

We thank Dr. Qian Yi, Dr. Weichao Sun, and Dr. Desheng Sun for their assistance in bioinformatics and cell culture methods.

Funding

This study was supported by the National Natural Science Foundation of China (81800785, 81972085, 81772394), the Natural Science Foundation of Guangdong Province (2018A0303100027), the Sanming Project of Shenzhen Health and Family Planning Commission (SZSM201612086), Shenzhen Science and Technology Planning (JCYJ20180228163401333), and the Doctor Innovation Project of Shenzhen Health System (SZBC2018015), and Shenzhen Peacock Project (KQTD20170331100838136).

Data Availability

The following information was supplied regarding data availability:
Raw data are available in the Supplemental Files.

Author Contributions

Review conception and design: WL and DW.
Drafting of the article: ZD, XH and MA.
Drawing the schematic: ZD, XH, and MA.
Revising the manuscript: JL, MA, ZZ, XC, GW, JX, and DL.
Final approval of the article: All authors.

Ethics approval and consent to participate

The study was reviewed and approved by the First Affiliated Hospital of Shenzhen University Health Science Center Research Ethics Committee.

Patient consent for publication

All patients signed informed consent forms for publication.

Competing interests

The authors declare that they have no competing interests.

References

- Bader GD, and Hogue CW. 2003. An automated method for finding molecular complexes in large protein interaction networks. *BMC Bioinformatics* 4:2. 10.1186/1471-2105-4-2
- Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, Marshall KA, Phillippy KH, Sherman PM, Holko M, Yefanov A, Lee H, Zhang N, Robertson CL, Serova N, Davis S, and Soboleva A. 2013. NCBI GEO: archive for functional genomics data sets--update. *Nucleic Acids Res* 41:D991-995. 10.1093/nar/gks1193
- Blain EJ. 2009. Involvement of the cytoskeletal elements in articular cartilage homeostasis and pathology. *Int J Exp Pathol* 90:1-15. 10.1111/j.1365-2613.2008.00625.x
- Chowdhury TT, Salter DM, Bader DL, and Lee DA. 2008. Signal transduction pathways involving p38 MAPK, JNK, NFkappaB and AP-1 influences the response of chondrocytes cultured in agarose constructs to IL-1beta and dynamic compression. *Inflamm Res* 57:306-313. 10.1007/s00011-007-7126-y
- Cibrian Uhalte E, Wilkinson JM, Southam L, and Zeggini E. 2017. Pathways to understanding the genomic aetiology of osteoarthritis. *Hum Mol Genet* 26:R193-R201. 10.1093/hmg/ddx302
- Croft P. 2005. The epidemiology of osteoarthritis: Manchester and beyond. *Rheumatology (Oxford)* 44 Suppl 4:iv27-iv32. 10.1093/rheumatology/kei058
- Cui Z, Crane J, Xie H, Jin X, Zhen G, Li C, Xie L, Wang L, Bian Q, Qiu T, Wan M, Xie M, Ding S, Yu B, and Cao X. 2016. Halofuginone attenuates osteoarthritis by inhibition of TGF-beta activity and H-type vessel formation in subchondral bone. *Ann Rheum Dis* 75:1714-1721. 10.1136/annrheumdis-2015-207923
- Feng Z, Zheng W, Li X, Lin J, Xie C, Li H, Cheng L, Wu A, and Ni W. 2017. Cryptotanshinone protects against IL-1beta-induced inflammation in human osteoarthritis chondrocytes and ameliorates the progression of osteoarthritis in mice. *Int Immunopharmacol* 50:161-167. 10.1016/j.intimp.2017.06.017
- Franceschini A, Szklarczyk D, Frankild S, Kuhn M, Simonovic M, Roth A, Lin J, Minguez P, Bork P, von Mering C, and Jensen LJ. 2013. STRING v9.1: protein-protein interaction networks, with increased coverage and integration. *Nucleic Acids Res* 41:D808-815. 10.1093/nar/gks1094

- 394 Fu D, Shang X, Ni Z, and Shi G. 2016. Shikonin inhibits inflammation and chondrocyte
395 apoptosis by regulation of the PI3K/Akt signaling pathway in a rat model of osteoarthritis.
396 *Exp Ther Med* 12:2735-2740. 10.3892/etm.2016.3642
- 397 Ganguly D, Haak S, Sisirak V, and Reizis B. 2013. The role of dendritic cells in autoimmunity.
398 *Nat Rev Immunol* 13:566-577. 10.1038/nri3477
- 399 Goldring MB, and Otero M. 2011. Inflammation in osteoarthritis. *Curr Opin Rheumatol* 23:471-
400 478. 10.1097/BOR.0b013e328349c2b1
- 401 Hochberg MC, Yerges-Armstrong L, Yau M, and Mitchell BD. 2013. Genetic epidemiology of
402 osteoarthritis: recent developments and future directions. *Curr Opin Rheumatol* 25:192-
403 197. 10.1097/BOR.0b013e32835cfb8e
- 404 Huang D, Ding Y, Luo WM, Bender S, Qian CN, Kort E, Zhang ZF, VandenBeldt K, Duesbery
405 NS, Resau JH, and Teh BT. 2008. Inhibition of MAPK kinase signaling pathways
406 suppressed renal cell carcinoma growth and angiogenesis in vivo. *Cancer Res* 68:81-88.
407 10.1158/0008-5472.CAN-07-5311
- 408 Huang DW, Sherman BT, Tan Q, Collins JR, Alvord WG, Roayaei J, Stephens R, Baseler MW,
409 Lane HC, and Lempicki RA. 2007. The DAVID Gene Functional Classification Tool: a
410 novel biological module-centric algorithm to functionally analyze large gene lists.
411 *Genome Biol* 8:R183. 10.1186/gb-2007-8-9-r183
- 412 Juhasz T, Matta C, Katona E, Somogyi C, Takacs R, Gergely P, Csernoch L, Panyi G, Toth G,
413 Reglodi D, Tamas A, and Zakany R. 2014. Pituitary adenylate cyclase activating
414 polypeptide (PACAP) signalling exerts chondrogenesis promoting and protecting effects:
415 implication of calcineurin as a downstream target. *PLoS One* 9:e91541.
416 10.1371/journal.pone.0091541
- 417 Juhasz T, Szentleleky E, Somogyi CS, Takacs R, Dobrosi N, Engler M, Tamas A, Reglodi D,
418 and Zakany R. 2015. Pituitary Adenylate Cyclase Activating Polypeptide (PACAP)
419 Pathway Is Induced by Mechanical Load and Reduces the Activity of Hedgehog
420 Signaling in Chondrogenic Micromass Cell Cultures. *Int J Mol Sci* 16:17344-17367.
421 10.3390/ijms160817344
- 422 Lai Y, Bai X, Zhao Y, Tian Q, Liu B, Lin EA, Chen Y, Lee B, Appleton CT, Beier F, Yu XP,
423 and Liu CJ. 2014. ADAMTS-7 forms a positive feedback loop with TNF-alpha in the
424 pathogenesis of osteoarthritis. *Ann Rheum Dis* 73:1575-1584. 10.1136/annrheumdis-

2013-203561

McKenzie S, and Torkington A. 2010. Osteoarthritis - management options in general practice. *Aust Fam Physician* 39:622-625.

Moro-Garcia MA, Mayo JC, Sainz RM, and Alonso-Arias R. 2018. Influence of Inflammation in the Process of T Lymphocyte Differentiation: Proliferative, Metabolic, and Oxidative Changes. *Front Immunol* 9:339. 10.3389/fimmu.2018.00339

Murphy G, Knauper V, Atkinson S, Butler G, English W, Hutton M, Stracke J, and Clark I. 2002. Matrix metalloproteinases in arthritic disease. *Arthritis Res* 4 Suppl 3:S39-49. 10.1186/ar572

Neogi T, and Zhang Y. 2013. Epidemiology of osteoarthritis. *Rheum Dis Clin North Am* 39:1-19. 10.1016/j.rdc.2012.10.004

Pereira RC, Martinelli D, Cancedda R, Gentili C, and Poggi A. 2016. Human Articular Chondrocytes Regulate Immune Response by Affecting Directly T Cell Proliferation and Indirectly Inhibiting Monocyte Differentiation to Professional Antigen-Presenting Cells. *Front Immunol* 7:415. 10.3389/fimmu.2016.00415

Rahman W, Patel R, and Dickenson AH. 2015. Electrophysiological evidence for voltage-gated calcium channel 2 (Cav2) modulation of mechano- and thermosensitive spinal neuronal responses in a rat model of osteoarthritis. *Neuroscience* 305:76-85. 10.1016/j.neuroscience.2015.07.073

Rhee I, Zhong MC, Reizis B, Cheong C, and Veillette A. 2014. Control of dendritic cell migration, T cell-dependent immunity, and autoimmunity by protein tyrosine phosphatase PTPN12 expressed in dendritic cells. *Mol Cell Biol* 34:888-899. 10.1128/MCB.01369-13

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, and Ideker T. 2003. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 13:2498-2504. 10.1101/gr.1239303

Sivan U, De Angelis J, and Kusumbe AP. 2019. Role of angiocrine signals in bone development, homeostasis and disease. *Open Biol* 9:190144. 10.1098/rsob.190144

Song J, Chang RW, and Dunlop DD. 2006. Population impact of arthritis on disability in older adults. *Arthritis Rheum* 55:248-255. 10.1002/art.21842

Stoddart MJ, Grad S, Eglin D, and Alini M. 2009. Cells and biomaterials in cartilage tissue

456 engineering. *Regen Med* 4:81-98. 10.2217/17460751.4.1.81

457 Strange-Vognsen HH, Arnbjerg J, and Hannibal J. 1997. Immunocytochemical demonstration of

458 pituitary adenylate cyclase activating polypeptide (PACAP) in the porcine epiphyseal

459 cartilage canals. *Neuropeptides* 31:137-141. 10.1016/s0143-4179(97)90082-2

460 Szklarczyk D, Franceschini A, Kuhn M, Simonovic M, Roth A, Minguéz P, Doerks T, Stark M,

461 Müller J, Bork P, Jensen LJ, and von Mering C. 2011. The STRING database in 2011:

462 functional interaction networks of proteins, globally integrated and scored. *Nucleic Acids*

463 *Res* 39:D561-568. 10.1093/nar/gkq973

464 Torres L, Dunlop DD, Peterfy C, Guermazi A, Prasad P, Hayes KW, Song J, Cahue S, Chang A,

465 Marshall M, and Sharma L. 2006. The relationship between specific tissue lesions and

466 pain severity in persons with knee osteoarthritis. *Osteoarthritis Cartilage* 14:1033-1040.

467 10.1016/j.joca.2006.03.015

468 Yuan X, and Yang S. 2015. Cilia/Ift protein and motor -related bone diseases and mouse models.

469 *Front Biosci (Landmark Ed)* 20:515-555. 10.2741/4323

470

Figure 1

Figure 1. Venn diagram.

A. GSE43923 and GSE55457 B. GSE11201 and GSE55457 C. GSE43923 and GSE11201 D. GSE43923, GSE55457 and GSE11201.




Figure 2

Figure 2. Gene Ontology (GO) analysis classified the DEGs into 3 groups: molecular function, biological process, and cellular component.




Figure 3

Figure 3. Gene Ontology (GO) enrichment analysis.

A Enrichment analysis of biological processes (BP). **B** Enrichment analysis of molecular function (MF). **C** Enrichment analysis of cellular component (CC). The gradual color represents the P value; the size of the black spots represents the gene number.




Figure 4

Fig 4. The PPI network of DEGs.

(Upregulated genes are marked in red; downregulated genes are marked in blue).





Figure 5


Figure 5. The 4 significant modules were obtained from PPI network.

Upregulated genes are marked in red; downregulated genes are marked in blue. **A.** MAPK14 module. **B.** PTPN12 module. **C.** PTPRC module. **D.** B9D1 module.


A



B



C



D




Figure 6

Figure 6. Cell biology detection.

A. Morphological observation of chondrocytes. **B.** Microscopic observation of toluidine blue staining of chondrocytes. **C.** - **N.** are the expression of COL I and COL II by immunofluorescence staining in human chondrocytes. **O.** is the expression of COL I and COL II by fluorescence western blot in human chondrocytes.




Figure 7

Figure 7. Comparison of expression of potential key genes and Apoptosis test in chondrocytes.

(**A.** MAPK-14; **B.** PTPRC; **C.** PTPN12; **D.** B9D1). All samples were normalized to the expression of GAPDH, and the relative expression levels of each gene were analyzed using the $2^{-\Delta\Delta C_t}$ method. **E.-J.** Apoptosis test of chondrocytes (DIC: bright field imaging; PI: piridine iodide staining imaging; Merged: composite imaging). The red arrow refers to cells that are stained or not by PI.




Table 1(on next page)

Table 1: The primers of 4 hub genes

1

Table 1: The primers of 4 hub genes

Gene		Primer sequence	Annealing Temperature (°C)	Length	PrimerBank ID
MAPK14		Fwd 5'- CCCGAGCGTTACCAGAACC -3'	62.4	19	194578904c1
		Rev 5'- TCGCATGAATGATGGACTGAAAT-3'	60.4	23	
PTPRC		Fwd 5'- ACCACAAGTTTACTAACGCAAGT-3'	60.4	23	18641362a1
		Rev 5'- TTTGAGGGGGATTCCAGGTAAT-3	60.5	22	
B9D1		Fwd 5'- TGGAGGAGGGGATCTCACAG-3'	61.9	20	343478278c1
		Rev 5'- CCGTAGGGGTTGGTGCTTTT-3	62.4	20	
PTPN12		Fwd 5'- AGTTGCCTTGTTGAAGGGGAT-3'	61.6	21	196114950c1
		Rev 5' - AGAAGGTGTCAAGATGGGTGG-3	61.4	21	

2

Table 2(on next page)

Table 2 Signaling pathway enrichment analysis of hub DEGs function in OA

1

Table 2 Signaling pathway enrichment analysis of hub DEGs function in OA

Expression	Term	Description	Gene Count	LogP
UP-DEGs	hsa04550	Signaling pathways regulating pluripotency of stem cells	3	-5.43
	hsa04670	Leukocyte transendothelial migration	3	-5.69
	GO:0008064	regulation of actin polymerization or depolymerization	3	-5.4
	GO:0032103	positive regulation of response to external stimulus	3	-6.33
	GO:0097529	myeloid leukocyte migration	3	-4.88
	GO:0032496	response to lipopolysaccharide	3	-4.31
DOWN-DEGs	GO:0097711	ciliary basal body-plasma membrane docking	3	-7.23

2

3

4

Table 3(on next page)

Table 3. Functional roles of 4 hub genes

1

Table 3. Functional roles of 4 hub genes

No.	Gene symbol	Full name	Function
1	MAPK-14	mitogen-activated protein kinase 14	Play an important role in the cascades of cellular responses evoked and inhibit the lysosomal degradation pathway of autophagy
2	PTPRC	protein tyrosine phosphatase receptor type C	High expression of PTPRC is associated with progression of PTP family
3	PTPN12	protein tyrosine phosphatase non-receptor type 12	High expression of PTPN12 is associated with progression of PTP family
4	B9D1	B9 domain containing 1	Component of the tectonic-like complex and acting as a barrier that prevents diffusion of transmembrane proteins between the cilia and plasma membranes

2

3