

Differential relieving effects of shikonin and its derivatives on inflammation and mucosal barrier damage caused by ulcerative colitis

Hongwei Han^{Equal first author, 1, 2}, Wenxue Sun^{Equal first author, 1, 2}, Lu Feng^{1, 2}, Zhongling Wen^{1, 2}, Minkai Yang^{1, 2}, Yingying Ma^{1, 2}, Jiangyan Fu^{1, 2}, Xiaopeng Ma^{1, 2}, Xinhong Xu^{1, 2}, Zhaoyue Wang^{1, 2}, Tongming Yin², Xiao-Ming Wang^{1, 2}, Gui-Hua Lu^{1, 3}, Jin-Liang Qi^{1, 2}, Hongyan Lin^{Corresp., 1, 2}, Yonghua Yang^{Corresp. 1, 2}

¹ State Key Laboratory of Pharmaceutical Biotechnology, Institute of Plant Molecular Biology, School of Life Sciences, Nanjing University, Nanjing, China

² Co-Innovation Center for Sustainable Forestry in Southern China, MOE Key Laboratory of Forest Genetics and Biotechnology, Nanjing Forestry University, Nanjing, China

³ School of Life Sciences, Huaiyin Normal University, Huaian, China

Corresponding Authors: Hongyan Lin, Yonghua Yang
Email address: linhy@nju.edu.cn, yangyh@nju.edu.cn

Background: Ulcerative colitis (UC) is one of the most challenging human diseases. Natural shikonin (SK) and its derivatives (with have higher accumulation) isolated from the root of *Lithospermum erythrorhizon* have numerous beneficial effects, such as wound healing and anti-inflammatory activities. Some researchers have reported that hydroxynaphthoquinone mixture (HM) and SK attenuate the acute UC induced by dextran sulfate sodium (DSS). However, no existing study has systemically investigated the effectiveness of SK and other hydroxynaphthoquinone natural derivative monomers on UC.

Methods: In this study, mice were treated with SK and its derivatives (25 mg/kg) and mesalazine (200 mg/kg) after DSS administration daily for one week. Disease progression was monitored daily by observing the changes in clinical signs and body weight.

Results: Intragastric administration natural single naphthoquinone attenuated the malignant symptoms induced by DSS. SK or its derivatives remarkably suppressed the serum levels of pro-inflammatory cytokines while increasing the inflammatory cytokine interleukin (IL)-10. Additionally, both SK and alkanin restrained the activities of cyclooxygenase-2 (COX-2), myeloperoxidase (MPO) and inducible nitric oxide synthase (iNOS) in serum and colonic tissues. SK and its derivatives inhibited the activation of nucleotide binding oligomerization domain-like receptors (NLRP3) inflammasome and NF-κB signaling pathway, thereby relieving the DSS-induced disruption of epithelial tight junction (TJ) in colonic tissues.

Conclusions: Our findings shed more lights on the pharmacological efficacy of SK and its derivatives in UC against inflammation and mucosal barrier damage.

Differential Relieving effects of shikonin and its derivatives on inflammation and mucosal barrier damage caused by ulcerative colitis

Hongwei Han^{1,2†}, Wenxue Sun^{1,2†}, Lu Feng^{1,2}, Zhongling Wen^{1,2}, Minkai Yang^{1,2}, Yingying Ma^{1,2}, Jiangyan Fu^{1,2}, Xiaopeng Ma^{1,2}, Xinhong Xu^{1,2}, Zhaoyue Wang^{1,2}, Tongming Yin², Xiaoming Wang^{1,2}, Guihua Lu^{1,3}, Jinliang Qi^{1,2}, Hongyan Lin^{1,2*}, Yonghua Yang^{1,2*}

¹State Key Laboratory of Pharmaceutical Biotechnology, Institute of Plant Molecular Biology, School of Life Sciences, Nanjing University, Nanjing, PR China

²Co-Innovation Center for Sustainable Forestry in Southern China, MOE Key Laboratory of Forest Genetics and Biotechnology, Nanjing Forestry University, Nanjing, PR China

³School of Life Sciences, Huaiyin Normal University, Huaian, PR China

[†]These authors contributed equally to this work

*Corresponding authors

Contact information of main corresponding author:

Dr. Yonghua Yang,

State Key Laboratory of Pharmaceutical Biotechnology,

Nanjing University, Nanjing, PR China.

Tel/Fax: 86-25-89686305,

Email: yangyh@nju.edu.cn

Abstract

Background: Ulcerative colitis (UC) is one of the most challenging human diseases. Natural shikonin (SK) and its derivatives (with have higher accumulation) isolated from the root of *Lithospermum erythrorhizon* have numerous beneficial effects, such as wound healing and anti-inflammatory activities. Some researchers have reported that hydroxynaphthoquinone mixture (HM) and SK attenuate the acute UC induced by dextran sulfate sodium (DSS). However, no existing study has systemically investigated the effectiveness of SK and other hydroxynaphthoquinone natural derivative monomers on UC.

Methods: In this study, mice were treated with SK and its derivatives (25 mg/kg) and mesalazine (200 mg/kg) after DSS administration daily for one week. Disease progression was monitored daily by observing the changes in clinical signs and body weight.

Results: Intragastric administration natural single naphthoquinone attenuated the malignant symptoms induced by DSS. SK or its derivatives remarkably suppressed the serum levels of pro-inflammatory cytokines while increasing the anti-inflammatory cytokine interleukin (IL)-10. Additionally, both SK and alkanin restrained the activities of cyclooxygenase-2 (COX-2), myeloperoxidase (MPO) and inducible nitric oxide synthase (iNOS) in serum and colonic tissues. SK and its derivatives inhibited the activation of nucleotide binding oligomerization domain-like receptors (NLRP3) inflammasome and NF- κ B signaling pathway, thereby relieving the DSS-induced disruption of epithelial tight junction (TJ) in colonic tissues.

Conclusions: Our findings shed more lights on the pharmacological efficacy of SK and its derivatives in UC against inflammation and mucosal barrier damage.

59

60 Introduction

61 Inflammatory bowel disease (IBD), which encompasses Crohn’s disease (CD) and ulcerative
 62 colitis (UC), is a kind of refractory and relapsing gastrointestinal inflammatory disease. IBD has
 63 not only affected the living quality of patients but also increased the risk of colon cancer if not
 64 treated in a timely manner (*Ungaro et al., 2017; Stone et al., 2003*). An increasing amount of
 65 evidence indicates that the incidence of colorectal cancer in IBD patients is generally higher than
 66 that in the general population (*Terzic et al., 2010*). Currently, traditional treatment strategies for
 67 IBD can be classified as three categories, namely, aminosalicyclic acid agents, glucocorticoids,
 68 and immunosuppressive agents (*Huang et al., 2009*). Typically, 5-aminosalicylic acid remains
 69 the preferred drug in treating IBD, which can directly treat enteritis through inhibiting
 70 inflammatory reaction in the intestinal tract and exerting antibacterial function. Nonetheless, this
 71 drug will be quickly absorbed by the small intestine if taken orally, leading to acute and chronic
 72 renal injury. Moreover, the drug cannot reach the inflammatory site, which can hardly meet the
 73 treatment goal (*Munk et al., 2004*). In addition, glucocorticoid is likely to develop dependence
 74 after long-term application, and disease recurrence is common after drug withdrawal (*Katz et al.,*
 75 *2004*). By contrast, immunosuppressant is suitable for patients with chronic enteritis, which can
 76 inhibit lymphocyte proliferation, to achieve the therapeutic effect; nonetheless, it is associated
 77 with strong toxic side effects (*Schroder et al., 2003*). The emergence of anti-tumor necrosis
 78 factor therapy has made a major advance in the treatment of patients with IBD for the past two
 79 decades (*Khan et al., 2019*). However, despite the fact that this therapy is useful to over half of
 80 IBD patients, a substantial proportion of patients are primary nonresponders or lose response
 81 with long-term use (*Hanauer et al., 2002; Ma et al., 2014*). Moreover, this therapy causes some
 82 disturbing safety issues encompassing the increased risk of infections and malignant tumor in

some groups (*Lichtenstein et al., 2012*). These factors are the reason for searching other safe, mild, durable, and effective drugs to cure IBD.

The disease characteristics of IBD and shortcomings of current treatments forced us to turn to the traditional Chinese medicine (TCM). *Lithospermum erythrorhizon* (*L. erythrorhizon*) is a commonly used TCM in the Chinese pharmacopoeia of the People's Republic of China and is typically abundant in shikonin (SK) derivatives, which have anti-tumor, anti-virus, and anti-inflammatory activities (*Chen et al., 2019; Yoshida et al., 2017*). For example, Fan et al. reported that the crude extract of *Arnebia euchroma* is effective on rats with experimental colitis (*Fan et al., 2013*). Dozens of species of SK are available in the Boraginaceae plants, among which the representative ones are SK, β , β -dimethylacryl-SK (β , β -dimethylacryl-SK), acetyl-SK (acetyl-SK), 5,8-dihydroxy-1, 4-naphthoquinone (naphthoquinone), and alkanin (AK, the enantiomeric of SK) with high contents (*Papageorgiou et al., 2006*). They belong to the naphthoquinones that are collectively known as SKs, which typically possess the advantages of bright color, long-lasting effect, and nontoxic side effects. Currently, the efficacy of each individual component of the crude extract has not been studied in detail. The pathological changes in the DSS-induced IBD model are similar to human UC. Thus, it is an ideal model that has been widely used to study the mechanism of UC and for screening potential drugs (*Farooq et al., 2018*).

In the present work, we successfully established a murine IBD model by treating C57BL/6 mice with DSS and the effectiveness of some natural SKs with high contents in *L. erythrorhizon* were further studied and compared for their efficacy in IBD treatment to explore the potential mechanisms.

Materials & Methods

Chemicals, Regents and antibodies

Chemicals and Regents were described in Supplementary File 1. All of compounds were characterized by ¹³C NMR, ¹H NMR in Supplementary File 2. All antibodies are described in Table S1. ECL Kit (#34077) was purchased from Thermo Scientific (USA). Haematoxylin-Eosin/HE Staining Kit (#20170804) was purchased from Solarbio Science & Technology Co. Ltd (Beijing, China).

Animals

A total of 66 health male C57BL/6 mice weighted 18-20 g (6-8 weeks old) were selected for in vivo experiment. All of them were purchased from Model Animal Research Center of Nanjing University (Nanjing, China), All animal experiments and welfare were treated in strict accordance with the relevant Guidelines for Care and Use of Laboratory Animals of Nanjing University and approved by the Laboratory Animal Ethics Committee of School of Life Sciences, Nanjing University (IACUC-1909002). The mice were group-housed in a laboratory with controlled conditions (22±2°C and 60%±5% humidity) under a 12 h light-dark cycle throughout the experiment. Animals were supplied with standard diet and sterile water ad libitum. All efforts were aimed at reducing the number and suffering of experimental animals while meeting the needs of the experiment.

DSS-induced colitis and design of drug treatment

The mice were randomly divided into eight groups (n = 6 per group) after one week of acclimatization, including the model (the dextran sulfate sodium; 3.5% (w/v) DSS), SK (6.25, 12.5, 25 mg/kg), AK (25 mg/kg), naphthoquinone (25 mg/kg), acetyl-SK (25 mg/kg), β, β-

dimethylacryl-SK (25 mg/kg), mesalazine (positive drug; 100, 200 mg/kg) and control group, then all the groups mice except for control group were exposed to 3.5% DSS, which was dissolved in drinking water. All drugs were dissolved in olive oil and were perfused 200 μ L per mice once daily at a predetermined dose by gavage which started at the same time as the DSS treatment and continued one week. In the control and DSS-induced groups, the mice fed orally with the same amount of olive oil. There was no significant difference in water intake among all groups. Body weight and the disease activity index (DAI) were measured and recorded everyday (*Farooq et al., 2018*). The DAI was calculated according to the standard listed in Table S2. After one week of treatment, the entire mice monocular eyeball in each mouse was removed under lightly anesthetized (carbon dioxide euthanasia) for blood collection. Then, all mice were euthanized by cervical dislocation in the unconscious state. The colonic tissues were quickly and safely removed and flushed with ice-cold PBS for pathological analysis. The same protocol was carried out at least three times in independent experiments.

Measurement of Cytokines, COX-2, MPO and iNOS in serum

After drug treatment, mice were sacrificed at the end of the experiment. The whole bloods samples were collected from peripheral blood after reperfusion. Afterwards, the blood was allowed to clot by standing at room temperature for 30 min, and later the clot was removed by centrifuging at 2000 g for 10 min in a refrigerated centrifuge. The resultant supernatant (serum) was then immediately transferred into a clean polypropylene tube. Kits were used to assay the level of various cytokines (IL-6, IL-1 β , TNF- α and IL-10), COX-2, MPO and iNOS in the serum of mice according to the product's guidelines (*Yang et al., 2020*).

Haematoxylin & Eosin (H&E) staining

Small sections of colonic tissue were fixed in 10% buffered formalin and embedded in paraffin. Thereafter, the sections were stained with haematoxylin and eosin (H&E) and the colonic mucosa was histologically evaluated by a pathologist in a blinded fashion, which was widely used as evaluation criterion (*Fan et al., 2013*).

Immunohistochemical analysis

Immunohistochemistry was performed for Cyclooxygenase-2 (COX-2), NF- κ B, NLRP3, the assay was conducted as described in (*Xu et al., 2019*). Briefly, the tissues were incubated with primary antibodies: anti-NF- κ B p65 (1:100), anti-COX-2 (1:200) and anti-NLRP3 (1:100) at 4°C for at least 12 h. Then, the sections were wash with PBS thrice and added with Alexa Fluor 488 labeled anti-mouse secondary antibody (Invitrogen, USA) at 25°C for 60 min. Signals were developed with Hematoxylin and DAB (Dako, Agilent Technologies, USA). Sections were examined using light microscopy and image analysis software (Image-pro plus 6.0; Media Cybernetics, Inc., Rockville, MD, USA) was used. The positive cumulative optical density (IOD) and the tissue pixel AREA (AREA) of each photo were measured, and thus average optical value (AO) was calculated by using $AO=IOD/AREA$.

Western blot analysis

Colonic samples were lysed with RIPA lysis buffer which containing 1% PMSF at 1:100 dilution on ice for 30 min. The insoluble components of cell lysates were removed by centrifugation (10000 rpm at 4°C for 10 min) and the harvested protein concentrations were assessed by using the BCA Protein Assay kit. Then the following immunoblot analysis was referred to the study by Farooq et al (*Farooq et al., 2018*). Finally, the immunoblots were quantified by densitometry with Image J Software (National Institutes of Health, Bethesda, Maryland, USA).

Data and Statistical analysis

Results were presented as mean±S.E.M. of three independent experiments, with six mice (n=6) per group to ensure their reliability. Statistical comparisons between the treated and untreated groups were performed using Student's *t*-test (two-tailed) using GraphPad PRISM5. Differences with a *P*-value < 0.05 were considered statistically significant.

Results

SK and its derivatives protected the colon from dextran sulfate sodium (DSS)-induced damage in an acute ulcerative colitis mouse model

An acute DSS-induced ulcerative colitis (UC) mouse model was established to investigate the effects of SK and its derivatives on colitis. Mice were given either sterile water or 3.5% DSS dissolved in sterile water in the model. Six other groups received 3.5% DSS and were administered SK (25 mg/kg), naphthoquinone (25 mg/kg), β , β -dimethylacryl-SK (25 mg/kg), acetyl-SK (25 mg/kg), AK (25 mg/kg), and mesalazine (200 mg/kg) orally both at the beginning of the experiment (n=6 mice per group). SK and its derivatives could significantly attenuate the severity level of inflammation at the end of this experiment. DSS group mice suffered a shortening of about 30% compared with normal mice in the case of the colorectum length. SK and its derivatives significantly prevented this shortening (Fig. 1A-1B). Moreover, SK and its derivatives-treated mice showed improvements in body weight and disease activity index, as well as the alleviation of splenomegaly (Fig. 1C–1F). Therefore, SK and its derivatives ameliorated the colon syndromes of DSS-induced damage in mice.

SK and its derivatives ameliorated the DSS-induced histological damage

The colon tissues of mice in each group were stained with H&E staining to observe the pathological changes. The results (Fig. 2A) indicated that compared with other derivatives group, inflammatory cells infiltration significantly decreased, the number of goblet cells decreased,

and crypt abscesses were distinctly reduced in SK and AK-treated group. Grading was performed in a blinded manner by a pathologist and the inflammation score result was shown in Fig. 2B. In brief, the administration of SK and AK markedly relieved the symptoms of DSS-induced IBD in mice.

SK and its derivatives relieved DSS-induced colonic epithelial TJ disruption in mice

Zonula occludens 1 (ZO-1), vascular cell adhesion molecule-1 (VCAM-1), Occludin and Claudin-1 proteins play an important role in epithelial TJ, which can protect mucosa epithelial cells from breaching by the harmful substances, maintain cellular permeability and integrity, and ensure homeostasis of internal environment. Therefore, the effect of SK and its derivatives on epithelial TJ proteins were further investigated. These results suggested that the expression of ZO-1, VCAM-1, Occludin, and Claudin-1 were remarkably reduced in the DSS group compared with the control group. However, SK and its derivatives at 25 mg/kg can restore the expression of these epithelial TJ proteins (Fig. 3A-3B).

SK and its derivatives inhibit the activation of NLRP3 inflammasome in colonic tissues

Many studies have revealed that NLRP3 affects several inflammatory disorders to a large extent including colitis (Lesuis *et al.*, 2012). Fig. 4A-4B shows the significantly up-regulated expression of NLRP3, apoptosis-associated speck-like protein containing a CARD (ASC), caspase-1, and IL-1 β in DSS group compared with control group. However, changes of caspase-1 and IL-1 β were significantly reduced by SK and naphthoquinone. SK- and β,β -Dimethylacryl-SK-treatment groups evidently reduced the level of ASC protein.

SK and its derivatives limited the expression of pro-inflammatory cytokines

When the circulating leukocytes are recruited into the colon, pro-inflammatory mediators are released, including pro-inflammatory cytokines such as IL-6, IL-1 β and TNF- α . The levels of IL-

6, IL-1 β , TNF- α and IL-10 were detected in the serum from 3.5% DSS-exposed mice given either control diet or SK and its derivatives to determine whether SK and its derivatives attenuated the release of pro-inflammatory cytokines, thereby protecting the intestinal mucosa. Fig. 5A–5D showed that treatment with SK and its derivatives at 25 mg/kg significantly reduced the levels of pro-inflammatory cytokines (IL-6, IL-1 β and TNF- α) and increased the level of anti-inflammatory cytokine (IL-10) in the serum. SK showed the best anti-inflammation effect that was similar to mesalazine group at 200 mg/kg. Fig. 5E–5H showed that SK and mesalazine (the positive control) reduced the levels of pro-inflammatory cytokines (IL-6, IL-1 β and TNF- α) and increased the level of anti-inflammatory cytokine (IL-10) in a dose-dependent manner, and SK treatment resulted in improved anti-inflammatory effect at a dose of 25 mg/kg. SK and its derivatives may improve the severity of colon damage induced by DSS by inhibiting the expression of pro-inflammatory cytokines.

SK and its derivatives reduced the activities of MPO, COX-2 and iNOS in serum

MPO is an enzyme that is largely present in neutrophils, and in monocytes and macrophages in small concentrations (*Davies et al., 2020*) and the high levels of COX-2 (*Chen et al., 2016*) and iNOS (*Pandurangan et al., 2014*) plays a central role in the initiation and propagation of the inflammation. MPO, COX-2 and iNOS concentrations in all the mice groups are shown in Fig. 6A-6C. The MPO activity is an indicator of neutrophil infiltration in inflamed colon tissues. Administration of DSS increased the activity of MPO as compared with normal mice. SK and AK administration at 25 mg/kg markedly reduced the concentrations of MPO, COX-2 and iNOS compared with DSS-induced mice.

SK and its derivatives attenuate the production of COX-2, NF- κ B and NLRP3 in colonic tissues

Immunohistochemical expressions of COX-2, NF- κ B and NLRP3 in all the groups of mice are shown in Fig. 7A-7C. DSS administration caused the increased production of COX-2, NF- κ B and NLRP3 in colonic tissues compared with the control group. Usage of SK and its derivatives reduced the expressions of COX-2, NF- κ B and NLRP3 significantly compared with mice induced with DSS.

SK and its derivatives regulate the COX-2 and iNOS expressions and the activation of NF- κ B in colonic tissues

Transcription factors of the NF- κ B family play a vital role in regulating genes, which can influence the immune and inflammatory response (*Andujar et al., 2012*). Evidence has verified that NF- κ B plays an important role in IBD development. In this disease, NF- κ B and STAT-3 activation is observed in inflamed colonic mucosa, leading to the production of COX-2 and iNOS. The phosphorylation level of STAT-3, NF- κ B p65 and I κ B was detected to evaluate the mechanism of SK. The outcome shown in Fig. 8A-8B suggested that the phosphorylation levels of STAT-3, NF- κ B p65 and I κ B α were markedly upregulated in DSS-treated group compared with the control group. These increases were markedly attenuated by treatment with DSS plus SK and its derivatives at 25 mg/kg, thereby decreasing the generation of COX-2 and iNOS in the colon homogenates (Fig. 8C-8D).

Discussion

L. erythrorhizon is a kind of TCM used for wound healing in ancient times. Its roots are typically rich with a variety of bioactive substances, including natural naphthoquinones, which represent an ongoing source of therapeutic agents (*Qiu et al., 2018*). Among the naphthoquinones isolated from *L. erythrorhizon*, SK has attracted extensive interest due to its multiple biological activities, such as anti-inflammatory (*Yoshida et al., 2017; Yang et al., 2014*), wound healing (*Yao et al.,*

2019), anti-microbial (*Andujar et al., 2013*), anti-ulcer (*Lesuis et al., 2012*), anti-thrombus (*Andrikopoulos et al., 2003*), and anti-cancer activities (*Lin et al., 2018*). Nearly 20 kinds of natural SK derivatives have been identified.

The anti-inflammatory effects of natural naphthoquinones have been extensively investigated. For instance, Fan et al. investigated the effectiveness of hydroxynaphthoquinone mixture (HM) isolated from *A. euchroma* in rats with experimental colitis induced by 2, 4, 6-trinitrobenzene sulfonic acid (TNBS) in 2013 (*Fan et al., 2013*). Their results suggested that HM can remarkably attenuate the clinical and histopathological severity of the TNBS-induced colitis in a dose-dependent manner. In addition, HM can also do the following: alleviate loss of body weight, hematochezia, and inflammation; reduce the macroscopic damage score; and improve the histological signs. Moreover, HM can evidently reduce the inflammatory infiltration, ulcer area, and the severity of goblet cell depletion. Notably, HM, which is isolated from *A. euchroma*, contains seven AK derivatives, including AK, deoxyalkannin, acetylalkannin, β , β -dimethylacrylalkannin, α -methylalkannin, isovalerylalkannin, and β -acetoxylalkannin. Isabel et al. reported that the oral administration of SK can attenuate the DSS-induced acute UC by blocking the activation of two major targets, NF- κ B and STAT-3 (*Andujar et al., 2012*). However, the effectiveness of the commonly used SK and its natural derivatives on acute colitis has not been systematically investigated yet.

In the present study, five representative natural naphthoquinones, namely, naphthoquinone, SK, acetyl-SK, β , β -dimethylacryl-SK and AK, were selected to treat DSS-induced colitis in mice. The administration of these five natural naphthoquinones could ameliorate the clinical severity of the wasting disease, prevent the shortening the length of colorectum, decrease the loss of body weight, improve the appearance of feces, and prevent bloody excrements. Such

beneficial effects were further proved by histological evaluation, as evidenced by the markedly reduced severity and extent of inflamed tissue damage and infiltration of inflammatory cells. Typically, the relative expression of pro-inflammatory cytokines, including IL-6, IL-1 β and TNF- α in serum in colon tissues, suggested that these five natural naphthoquinones had beneficial effects. MPO is a marker of neutrophil infiltration, which has been observed to be activated in several experimental colitis models, including the DSS-induced colitis. Therefore, MPO is an indicator to quantify intestinal inflammation and evaluate the severity of inflammation. In addition, the pro-inflammatory synthases, including iNOS and COX-2, are inflammatory mediators with vital functions during the pathogenesis of UC (*Itzkowitz et al., 2006*). The five natural naphthoquinones could suppress MPO, iNOS and COX-2 activities. Such results were consistent with those found in histological examination, in which the inflammatory extent in colonic tissues was less than that in SK- and AK-treated animals and naphthoquinone-, acetyl-SK- and β , β -dimethylacryl-SK-treated mice. Moreover, results of western blot had further verified that SK can relieve the colitis in mice by inhibiting the activation of NF- κ B and NLRP3 inflammasome and disruption of epithelial TJ proteins. Collectively, these aforementioned results offer insights into the potential use of SK and its derivatives in treating IBD.

Conclusions

Our results suggested that intra-gastric administration of single compound like SK or its derivatives contributed to attenuating the loss of body weight, shortening the colon length, increasing the spleen weight and relieving colonic epithelial tight disruption in mice induced by DSS. Through a number of related pathological experiments, we have explained the underlying mechanism that SK and its derivatives could inhibit the activation of NLRP3 inflammasome and

NF- κ B signaling pathway. These findings provide new evidence for the potential use of SK and its derivatives to treat the inflammatory bowel disease (IBD).

Additional Information and Declarations

Data Availability

The following information was supplied regarding data availability:

The raw data are available in the Supplementary File 3. The raw data is showing H&E data, IHC raw data and Western blot bands.

References

- Ungaro R, Mehandru S, Allen PB, Peyrin-Biroulet L, Colombel JF. 2017. Ulcerative colitis. *Lancet* **359**: 1756-1770 [https://doi.org/10.1016/S0140-6736\(16\)32126-2](https://doi.org/10.1016/S0140-6736(16)32126-2).
- Stone MA, Mayberry JF, Baker R. 2003. Prevalence and management of inflammatory bowel disease: a cross-sectional study from central England. *European journal of gastroenterology & hepatology* **15**:1275-1280 <https://doi.org/10.1097/00042737-200312000-00004>.
- Terzic J, Grivennikov S, Karin E, Karin M. 2010. Inflammation and colon cancer. *Gastroenterology* **138**: 2101-2114 <https://doi.org/10.1053/j.gastro.2010.01.058>.
- Huang LJ, Zhu Q, Lei M, Cao Q. 2009. Current use of immunosuppressive agents in inflammatory bowel disease patients in East China. *World journal of gastroenterology* **15**:3055-3059 <https://doi.org/10.3748/wjg.15.3055>.
- Munk EM, Pedersen L, Floyd A, Norgard B, Rasmussen HH, Sorensen HT. 2004. Inflammatory bowel diseases, 5-aminosalicylic acid and sulfasalazine treatment and risk of acute pancreatitis: a population-based case-control study. *The American journal of gastroenterology* **99**:884-888 <https://doi.org/10.1111/j.1572-0241.2004.04123.x>.

341 **Katz JA. 2004.** Treatment of inflammatory bowel disease with corticosteroids. *Gastroenterology*
342 *clinics of North America* **33**:171-189 [https://doi.org/ 10.1016/j.gtc.2004.02.002](https://doi.org/10.1016/j.gtc.2004.02.002).

343 **Schroder O, Stein J. 2003.** Low dose methotrexate in inflammatory bowel disease: current
344 status and future directions. *The American journal of gastroenterology* **98**:530-537
345 [https://doi.org/ 10.1016/S0002-9270\(02\)06027-6](https://doi.org/10.1016/S0002-9270(02)06027-6).

346 **Khan S, Rupniewska E, Neighbors M, Singer D, Chiarappa J, Obando C. 2019.** Real-world
347 evidence on adherence, persistence, switching and dose escalation with biologics in adult
348 inflammatory bowel disease in the United States: A systematic review. *Journal of clinical*
349 *pharmacy and therapeutics* **44**:495-507 [https://doi.org/ 10.1111/jcpt.12830](https://doi.org/10.1111/jcpt.12830).

350 **Hanauer SB, Feagan BG, Lichtenstein GR, Mayer LF, Schreiber S, Colombel JF,**
351 **Rachmilewitz D, Wolf DC, Olson A, Bao W, Rutgeerts P, Group AIS. 2002.** Maintenance
352 infliximab for Crohn's disease: the ACCENT I randomised trial. *Lancet* **359**:1541-1549
353 [https://doi.org/ 10.1016/S0140-6736\(02\)08512-4](https://doi.org/10.1016/S0140-6736(02)08512-4).

354 **Ma C, Huang V, Fedorak DK, Kroeker KI, Dieleman LA, Halloran BP, Fedorak RN. 2014.**
355 Crohn's disease outpatients treated with adalimumab have an earlier secondary loss of response
356 and requirement for dose escalation compared to infliximab: a real life cohort study. *Journal of*
357 *Crohn's & colitis* **8**:1454-1463 [https://doi.org/ 10.1016/j.crohns.2014.05.007](https://doi.org/10.1016/j.crohns.2014.05.007).

358 **Lichtenstein GR, Feagan BG, Cohen RD, Salzberg BA, Diamond RH, Price S, Langholff W,**
359 **Londhe A, Sandborn WJ. 2012.** Serious infection and mortality in patients with Crohn's
360 disease: more than 5 years of follow-up in the TREAT registry. *The American journal of*
361 *gastroenterology* **107**:1409-1422 [https://doi.org/ 10.1038/ajg.2012.218](https://doi.org/10.1038/ajg.2012.218).

362 **Chen Y, Chen ZY, Chen L, Zhang JY, Fu LY, Tao L, Zhang Y, Hu XX, Shen XC. 2019.**
363 Shikonin inhibits triple-negative breast cancer-cell metastasis by reversing the epithelial-to-

mesenchymal transition via glycogen synthase kinase 3 beta-regulated suppression of beta-catenin signaling. *Biochemical pharmacology* **166**:33-45 <https://doi.org/10.1016/j.bcp.2019.05.001>.

Yoshida LS, Kakegawa T, Yuda Y, Takano-Ohmuro H. 2017. Shikonin changes the lipopolysaccharide-induced expression of inflammation-related genes in macrophages. *Journal of natural medicines* **71**:723-734 <https://doi.org/10.1016/j.bcp.2019.05.001>.

Fan HY, Zhang ZL, Liu K, Yang MY, Lv WH, Che X, Xu H, Song WW. 2013. Effectiveness of a hydroxynaphthoquinone fraction from *Arnebia euchroma* in rats with experimental colitis. *World journal of gastroenterology* **19**:9318-9327 <https://doi.org/10.3748/wjg.v19.i48.9318>.

Papageorgiou VP, Assimopoulou AN, Samanidou VF, Papadoyannis IN. 2006. Analytical methods for the determination of alkannins and shikonins. *Current Organic Chemistry* **10**:583-622 <https://doi.org/10.2174/138527206776055277>.

Farooq SM, Hou Y, Li H, O'Meara M, Wang Y, Li C, Wang JM. 2018. Disruption of GPR35 Exacerbates Dextran Sulfate Sodium-Induced Colitis in Mice. *Digestive diseases and sciences* **63**:2910-2922 <https://doi.org/10.1007/s10620-018-5216-z>.

Yang L, Zhou RY, Tong Y, Chen PF, Shen Y, Miao S, Liu XQ. 2020. Neuroprotection by dihydrotestosterone in LPS-induced neuroinflammation. *Neurobiology of Disease* **140**: 104814-104827. <https://doi.org/10.1016/j.nbd.2020.104814>.

Xu M, Duan XY, Chen QY, Fan H, Hong ZC, Deng SJ, Nan Z, Wu H, Dong YL, Liu YJ, Zhou CZ. 2019. Effect of compound sophorae decoction on dextran sodium sulfate (DSS)-induced colitis in mice by regulating Th17/Treg cell balance. *Biomedicine & Pharmacotherapy* **109**:2396-2408 <https://doi.org/10.1016/j.biopha.2018.11.087>.

Andujar I, Rios JL, Giner RM, Miguel Cerda J, Recio Mdel C. 2012. Beneficial effect of

shikonin on experimental colitis induced by dextran sulfate sodium in BALB/c mice. *Evidence-based complementary and alternative medicine : eCAM* **2012**:271606-271620 <https://doi.org/10.1155/2012/271606>.

Qiu HY, Wang PF, Lin HY, Tang CY, Zhu HL, Yang YH. 2018. Naphthoquinones: A continuing source for discovery of therapeutic antineoplastic agents. *Chemical biology & drug design* **91**:681-690 <https://doi.org/10.1111/cbdd.13141>.

Yang Y, Wang J, Yang Q, Wu S, Yang Z, Zhu H, Zheng M, Liu W, Wu W, He J, Chen Z. 2014. Shikonin inhibits the lipopolysaccharide-induced release of HMGB1 in RAW264.7 cells via IFN and NF-kappaB signaling pathways. *International immunopharmacology* **19**:81-87 <https://doi.org/10.1016/j.intimp.2014.01.003>.

Yao CH, Chen KY, Chen YS, Li SJ, Huang CH. 2019. Lithospermi radix extract-containing bilayer nanofiber scaffold for promoting wound healing in a rat model. *Materials science & engineering C, Materials for biological applications* **96**:850-858 <https://doi.org/10.1016/j.msec.2018.11.053>.

Andujar I, Rios JL, Giner RM, Recio MC. 2013. Pharmacological Properties of Shikonin - A Review of Literature since 2002. *Planta Medica* **79**:1685-1697 <https://doi.org/10.1055/s-0033-1350934>.

Lesuis N, Befrits R, Nyberg F, van Vollenhoven RF. 2012. Gender and the treatment of immune-mediated chronic inflammatory diseases: rheumatoid arthritis, inflammatory bowel disease and psoriasis: an observational study. *BMC Medicine* **10**:82-90. <https://doi.org/10.1186/1741-7015-10-82>.

Davies MJ, Hawkins CL. 2020. The Role of Myeloperoxidase in Biomolecule Modification, Chronic Inflammation, and Disease. *Antioxid Redox Sign.* **32**: 957-981. [10.1089/ars.2020.8030](https://doi.org/10.1089/ars.2020.8030).

410 **Chen L, Teng H, Fang T, Xiao J. 2016.** Agrimonolide from *Agrimonia pilosa*
 411 suppresses inflammatory responses through down-regulation of COX-2/iNOS and
 412 inactivation of NF- κ B in lipopolysaccharide-stimulated macrophages. *Phytomedicine*
 413 **23**:846–855. <https://doi.org/10.1016/j.phymed.2016.03.016>.

414 **Pandurangan AK, Kumar SA, Dharmalingam P, Ganapasam S. 2014.** Luteolin, a
 415 bioflavonoid inhibits azoxymethane-induced colon carcinogenesis: involvement of iNOS and
 416 COX-2. *Pharmacogn. Mag.* **10**: 306–310. <https://doi.org/10.4103/0973-1296.133285>.

417 **Andrikopoulos NK, Kaliora AC, Assimopoulou AN, Papapeorgiou VP. 2003.** Biological
 418 activity of some naturally occurring resins, gums and pigments against in vitro LDL oxidation.
 419 *Phytotherapy Research* **17**:501-507 <https://doi.org/10.1002/ptr.1185>.

420 **Lin HY, Han HW, Sun WX, Yang YS, Tang CY, Lu GH, Qi JL, Wang XM, Yang YH. 2018.**
 421 Design and characterization of alpha-lipoic acyl shikonin ester twin drugs as tubulin and PDK1
 422 dual inhibitors. *European journal of medicinal chemistry* **144**:137-150 <https://doi.org/10.1016/j.ejmech.2017.12.019>.

424 **Itzkowitz SH. 2006.** Molecular biology of dysplasia and cancer in inflammatory bowel disease.
 425 *Gastroenterology clinics of North America* **35**:553-571 <https://doi.org/10.1016/j.gtc.2006.07.002>.

426

Figure 1

SK and its derivatives protect the colon from DSS-induced damage

(A) Macroscopic observation of colon length. (B) Bar graph showing colon length. $**p < 0.01$ versus control group. (C) Pattern of daily weight changes. $*p < 0.05$, $**p < 0.01$ versus DSS-treated group. (D) Disease activity index of mice in each group. $*p < 0.05$, $**p < 0.01$ versus DSS-treated group. (E) Macroscopic observation of spleen size. (F) Bar graph showing spleen weight. Data are shown as mean \pm S.E.M, $*p < 0.05$ versus control group (n=6 per group).




Figure 2

Colon histology and inflammation score of mice in each group (H&E staining 200× and 400×)

(A) The colon histology sections were stained with hematoxylin and eosin. The colon histology sections of DSS-treated group mice revealed sever pathology with observation of submucosal edema with extensive inflammatory cell infiltration. The frame indicates the region magnified in the bottom panel. **(B)** Inflammation scores of each group. Data are shown as mean \pm S.E.M, * p <0.05, ** p <0.01 versus control group; # p <0.05, ## p <0.01 versus DSS-treated group (n=6 per group).




Figure 3

SK and its derivatives relieved DSS-induced colonic epithelial tight junction disruption in mice

(A) Protein expression of ZO-1, VCAM-1, Occludin and Claudin-1 in the colon tissues were analyzed by Western blot. **(B)** Relative protein expression ratios of were determined by densitometry and normalized to GAPDH. Each point represents the mean \pm S.E.M. from three replicates (* p <0.05, ** p <0.01 versus control group; # p <0.05, ## p <0.01 versus DSS-treated group).




Figure 4

SK and its derivatives reduced the level of NLRP3 inflammasome activation in colon tissues

(A) Protein levels of NLRP3, ASC, caspase-1, and IL-1 β were determined by Western blotting. β -actin was used as a control. (B) Relative protein expression ratio was determined by densitometry and normalized to β -actin. Each point represents the mean \pm S.E.M. from three replicates (* p <0.05, ** p <0.01 versus control group; # p <0.05, ## p <0.01 versus DSS-treated group).




Figure 5

SK and its derivatives suppressed pro-inflammatory cytokines and promoted anti-inflammatory cytokine in serum

(A-D) The protein level of cytokines IL-6, IL-1 β , TNF- α and IL-10 in serum were determined by ELISA (DSS: 3.5%; SK, Naphthoquinone, β , β -dimethylacryl-SK, Acetyl-SK, AK: 25 mg/kg; Mesalazine: 200 mg/kg). (E-H) Effects of different concentrations of SK on the levels of cytokines IL-6, IL-1 β , TNF- α and IL-10. Data are shown as mean \pm S.E.M, * p <0.05, ** p <0.01 versus control group; # p <0.05, ## p <0.01 versus DSS-treated group (n=6 per group).




Figure 6

SK and its derivatives reduced the level of COX-2, MPO and iNOS in the serum of mice

The concentrations of COX-2 (**A**), MPO (**B**) and iNOS (**C**) were measured using an enzyme-linked immunosorbent assay kit. Each point represents the mean \pm S.E.M. from three replicates (* p <0.05, ** p <0.01 versus control group; # p <0.05, ## p <0.01 versus DSS-treated group).




Figure 7

SK and its derivatives attenuate the expression of COX-2, NF- κ B and NLRP3

(A) Immuno-histochemical analysis of COX-2 in each group. (B) Immuno-histochemical analysis of NF- κ B in each group. (C) Immuno-histochemical analysis of NLRP3 in each group. The slides were incubated with primary antibody. After the secondary antibody incubation, the slides were developed with DAB and counter-stained with hematoxylin. The images were taken at 400 \times magnification. Data are presented as means \pm S.E.M. from three replicates (* p <0.05, ** p <0.01 versus control group; # p <0.05, ## p <0.01 versus DSS-treated group).




Figure 8

SK and its derivatives modulated COX-2, iNOS expression, and NF- κ B signal pathway in UC mice

(A) Western blot analysis of COX-2 and iNOS levels in colonic tissues in each group. Data shown are representative of three independent experiments. (B) Relative protein expression ratios of COX-2 and iNOS were determined by densitometry. (C) Western blot analysis of proteins expression level of NF- κ B signal pathway. (D) Relative protein expression ratio was determined by densitometry. Each point represents the mean \pm S.E.M. from three replicates (* p <0.05, ** p <0.01 versus control group; # p <0.05, ## p <0.01 versus DSS-treated group).

