

***Monomorium* (Hymenoptera: Formicidae) of the Arabian Peninsula with description of two new species, *M. heggyi* sp. n. and *M. khalidi* sp. n. (#54174)**

1

First submission

Guidance from your Editor

Please submit by **15 Nov 2020** for the benefit of the authors (and your \$200 publishing discount) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Author notes

Have you read the author notes on the [guidance page](#)?



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

! Custom checks

29 Figure file(s)

New species checks



Have you checked our [new species policies](#)?



Do you agree that it is a new species?



Is it correctly described e.g. meets ICZN standard?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

***Monomorium* (Hymenoptera: Formicidae) of the Arabian Peninsula with description of two new species, *M. heggyi* sp. n. and *M. khalidi* sp. n.**

Mostafa Sharaf^{Corresp., 1}, Amr A Mohamed², Brendon E Boudinot³, James K Wetterer⁴, Hathal M Al Dhafer⁵, Francisco Hita Garcia⁶, Abdulrahman S Aldawood⁵

¹ College of Food and Agriculture Sciences, Department of Plant Protection, King Saud University, Riyadh, Riyadh, Saudi Arabia

² Department of Entomology, Cairo University, Giza, Egypt

³ Department of Entomology/Nematology, University of California, Davis, Davis, California, United States

⁴ Wilkes Honors College, Florida Atlantic University, Florida, Florida, USA

⁵ Department of Plant Protection, College of Food and Agriculture Sciences, King Saud University, Riyadh, Riyadh, Saudi Arabia

⁶ Biodiversity and Biocomplexity Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa, Okinawa, Japan

Corresponding Author: Mostafa Sharaf
Email address: antsharaf@gmail.com

We present a revised and updated synoptic list of 44 Arabian *Monomorium* species, including two new species of the *M. salomonis* species-group: *M. heggyi* **sp. n.**, and *M. khalidi* **sp. n.** We propose the following new synonyms: *M. abeillei* André (= *M. wahibiense* Collingwood & Agosti **syn. n.**); *M. areniphilum* Santschi (= *M. fezzanense* Collingwood & Agosti **syn. n.**, = *M. hemame* Collingwood & Agosti **syn. n.** = *M. marmule* Collingwood & Agosti **syn. n.**); *M. bicolor* Emery (= *M. phoenicum* Santschi **syn. n.**); *M. harithe* Collingwood & Agosti (= *M. najrane* Collingwood & Agosti **syn. n.**); *M. niloticum* Emery (= *M. matame* Collingwood & Agosti **syn. n.**); and *M. nitidiventre* Emery (= *M. yemene* Collingwood & Agosti **syn. n.**). An illustrated key and distribution maps are presented for the treated species. Ecological and biological notes are given when available. The majority of Arabian *Monomorium* species (25) are endemic to the peninsula. All except one of the remaining species are more broadly ranging Afrotropical and Palearctic species, supporting the view of Arabia as a biogeographical crossroads between these two regions. *Monomorium floricola* (Jerdon), the sole species of Indomalayan origin, is recorded for the first time from the Arabian Peninsula.

***Monomorium* (Hymenoptera: Formicidae) of the Arabian Peninsula with description of two new species, *M. heggvi* sp. n. and *M. khalidi* sp. n.**

Running head: Revision of Arabian *Monomorium*.

Mostafa R. Sharaf^{a,Ω}, Amr A. Mohamed^b, Brendon E. Boudinot^c, James K. Wetterer^d, Francisco Hita Garcia^e, Hathal M. Al Dhafer^a & Abdulrahman S. Aldawood^a

^a Department of Plant Protection, College of Food and Agriculture Sciences, King Saud University, Riyadh, Kingdom of Saudi Arabia

^b Department of Entomology, Faculty of Science, Cairo University, Giza, Egypt

^c Department of Entomology/Nematology, University of California, Davis, USA.

^d Wilkes Honors College, Florida Atlantic University, 5353 Parkside Drive, Jupiter, FL 33458, USA.

^e Biodiversity and Biocomplexity Unit, Okinawa Institute of Science and Technology Graduate University, Onna-son, Okinawa, Japan.

^Ω corresponding author, e-mail: antsharaf@gmail.com; mosharaf@ksu.edu.sa

ABSTRACT

We present a revised and updated synoptic list of 44 Arabian *Monomorium* species, including two new species of the *M. salomonis* species-group: *M. heggvi* sp. n., and *M. khalidi* sp. n. We propose the following new synonyms: *M. abeillei* André (= *M. wahibiense* Collingwood & Agosti syn. n.); *M. areniphilum* Santschi (= *M. fezzanense* Collingwood & Agosti syn. n., = *M. hemame* Collingwood & Agosti syn. n. = *M. marmule* Collingwood & Agosti syn. n.); *M. bicolor* Emery (= *M. phoenicum* Santschi syn. n.); *M. harithe* Collingwood & Agosti (= *M. najrane* Collingwood & Agosti syn. n.); *M. niloticum* Emery (= *M. matame* Collingwood & Agosti syn. n.); and *M. nitidiventre* Emery (= *M. yemene* Collingwood & Agosti syn. n.). An illustrated key and distribution maps are presented for the treated species. Ecological and biological notes are given when available. The majority of Arabian *Monomorium* species (25) are endemic to the peninsula. All except one of the remaining species are more broadly ranging Afrotropical and Palearctic species, supporting the view of Arabia as a biogeographical crossroads between these two regions. *Monomorium floricola* (Jerdon), the sole species of Indomalayan origin, is recorded for the first time from the Arabian Peninsula.

Keywords. biogeography, faunal list, Middle East, taxonomic revision.

INTRODUCTION

Monomorium is one of the largest genera of ants, with 326 valid species and subspecies, as well as three fossil species (Bolton, 2020). *Monomorium* species predominantly inhabit tropical, subtropical, and warm temperate regions of the world (Bolton, 1987; Bolton, 1994; Brown, 2000). The majority of *Monomorium* species are native to the Afrotropical bioregion (Bolton, 1987; Hita Garcia et al., 2013; Bolton, 2020). In contrast, the Palaearctic and New World faunas are relatively species-poor (Kempf, 1972; Sharaf et al., 2018a; Fernández & Serna, 2019). In total, 56 species have been reported from the Arabian Peninsula (Collingwood, 1985; Collingwood & Agosti, 1996; Aldawood et al., 2010; Aldawood & Sharaf, 2011; Sharaf et al., 2013a, b, c, Sharaf et al., 2014a, b, Sharaf et al., 2015, 2017a, b, 2018a, b). Most are generalized feeders or granivores, but some are leptoecious or parasitic (Ettershank, 1966; Bolton, 1987; Brown, 2000). Notably, *Monomorium* is one of the few genera that includes several very successful cosmopolitan tramp species, such as *M. pharaonis* (L.) and *M. floricola* (Jerdon), *M. latinode* Mayr, and *M. subopacum* (Smith) (Bolton, 1987; Wetterer, 2010a, b).

The genus *Monomorium* has a long and complex taxonomic history (see Heterick 2006 for a full history of the genus). Emery (1922) offered subgeneric names for several *Monomorium* species, based on characters of the antennal club and number of antennomeres, some names were later raised to the generic level. Ettershank (1966) made a bold attempt to organize the global fauna of *Monomorium* and related genera. Bolton (1987) extensively reviewed, refined and keyed the Afrotropical species recognizing 145 species in 8 species-groups including descriptions of 46 new species and 43 new synonyms. The Malagasy *Monomorium* fauna was revised by Heterick (2006) following Bolton's Afrotropical species-groups as a basis for organizing numerous new species. Heterick (2001, 2003, 2009) comprehensively treated the Australian *Monomorium* fauna, whereas DuBois (1986) and Mackay & Mackay (2002) studied the Nearctic fauna. The comprehensive phylogenetic work of the subfamily Myrmicinae (Ward et al., 2015) has adopted substantial taxonomic changes including the resurrection of the genera *Sylophopsis* and *Trichomyrmex* from synonymy under *Monomorium*. Sparks et al. (2014) fully revised the *Monomorium rothsteini* species-complex. Numerous smaller and regional contributions are scattered in the literature including Argentina (Kusnezov, 1949), China (Wu &

Wang, 1995), Colombia (Fernández & Serna, 2019), Fiji (Sarnat & Economo, 2012), Italy (Baroni Urbani, 1964, 1968), Japan (Morisita et al., 1992), Morocco (Barech et al., 2017), New Zealand (Brown, 1958), Polynesia (Wilson & Taylor, 1967), Taiwan (Terayama, 2009), the Iberian Peninsula (Collingwood, 1978), Turkmenistan (Dlussky et al., 1990), and Western Europe (Bernard, 1968).

However, our understanding of the phylogenetic relationships among the currently accepted *Monomorium* species-groups remains far from satisfactory and the overall taxonomic foundation is in a rather moderate stage (Sparks et al., 2019). Moreover, our knowledge about the diversity of the *genus* is still fragmentary for several regions, such as the Mediterranean, the North African-Indian Desert (including what referred as the Saharo-Arabian in some literatures [Holt et al., 2013]) (Vigna Taglianti et al., 1999; Sharaf et al., 2017a), as well as the Indomalayan and the Neotropical realms, which contain large numbers of described infraspecific taxa in need of further reevaluation (Borowiec, 2014). We note that, even the Afrotropical *Monomorium* fauna, which is the most diverse and best studied so far, is in dire need of an updated taxonomic revision with likely more than 100 undescribed species and many valid species needing reevaluation (Bolton, 1987; Hita Garcia, pers. obs.).

The oldest records of *Monomorium* species from the Arabian Peninsula were of *M. carbonarium* (Smith) and *M. niloticum* Emery (Forel). Collingwood (1985) reported 17 ant species from the Kingdom of Saudi Arabia (KSA). Collingwood & Agosti (1996) listed 49 ant species from the entire Arabian Peninsula, including 32 new species. Additional papers have included records from the KSA (Aldawood & Sharaf, 2011; Sharaf & Aldawood, 2013b; Sharaf et al., 2015, 2018 a, b), Oman (Sharaf et al., 2018b; Monks et al., 2019), the Socotra Archipelago (Collingwood, 2004; Sharaf et al., 2017a), the United Arab Emirates (UAE) (Collingwood et al., 2011), and Yemen (Aldawood et al., 2010).

Collingwood (1985) and Collingwood & Agosti (1996) recognized a total of 53 *Monomorium* species on the Arabian Peninsula, of which 17 were described from countries in the region. However, taxonomic status of several species has remained uncertain due to brief descriptions with insufficient differential diagnoses, apparent ambiguities in the taxonomic keys, lack of species-group assignment (Sharaf et al., 2020a, b). The present study aims to clarify the current status of the Arabian *Monomorium* by providing the following:

I. Diagnoses of Arabian *Monomorium* and *Monomorium* species-groups

- II. A synoptic checklist of Arabian *Monomorium*
- III. An illustrated identification key to species based on the worker caste.
- IV. New taxonomic treatments proposing new synonymies and describing two new species.
- V. Biogeographical analyses, including distribution maps

MATERIAL AND METHODS

The species names follow the online catalogue of ants of the world (Bolton 2020). We made digital color images of each species using a Leica DFC450 digital camera with a Leica Z16 APO microscope and LAS (v3.8) software. The images are available online on AntWeb (<http://www.AntWeb.org>) and are accessible through unique specimen identifiers (e.g. CASENT0922288). Distribution maps were made using DIVA-GIS (version 7.5.0.0).

Measurements and indices.

All measurements are in millimeters and follow the standard measurements of Bolton (1987) and Sharaf et al. (2018a).

CI= Cephalic Index ($HW/HL \times 100$).

EI= Eye Index ($EL/HW \times 100$).

EL= Eye Length; maximum diameter of eye in lateral view.

EM= Distance between anterior margin of eye and mandibular insertion in lateral view.

HL= Head Length; maximum length of head, excluding mandibles in full-face view.

HW= Head Width; maximum width of head behind eyes in full-face view.

ML= Mesosoma Length (=Weber Length); length of mesosoma in lateral view); from a point at which pronotum meets cervical shield to posterior base of propodeal lobes or teeth.

PPH= Postpetiole Height; maximum height measured in lateral view.

PPL= Postpetiole Length; maximum length measured in dorsal view.

PPW= Postpetiole Width; maximum width measured in dorsal view.

PTH= Petiole Height; maximum height measured in lateral view.

PTL= Petiole Length; maximum length measured in dorsal view, from anterior margin to posterior margin.

PTW= Petiole Width; maximum width measured in dorsal view.

PW= Pronotal Width; maximum width in dorsal view.

125 SI= Scape Index ($SL/HW \times 100$).
 126 SL= Scape Length, excluding basal neck.
 127 TL= Total Length, sum of lengths of head, mesosoma, petiole, postpetiole and gaster in
 128 profile.

130 Abbreviations of museums.

131 Abbreviations of natural history collections follow Evenhuis (2020). The material on which this
 132 study is based is located and/or was examined at the following institutions:

133 BMNH = The Natural History Museum, London, United Kingdom.

134 CASC = California Academy of Sciences, San Francisco, USA.

135 KSMA = King Saud University Museum of Arthropods, Riyadh, Kingdom of Saudi Arabia.

136 LACM = Los Angeles County Museum of Natural History, Los Angeles, CA, USA.

137 MNHN = Muséum National d'Histoire Naturelle, Paris, France.

138 NHMB = Naturhistorisches Museum, Basel, Switzerland.

139 OUMC = Oxford University Museum, Oxford, U.K.

140 WMLC = World Museum Liverpool, Liverpool, United Kingdom.

141 Nomenclatural acts

142 The electronic version of this article in Portable Document Format (PDF) will represent a
 143 published work according to the International Commission on Zoological Nomenclature (ICZN),
 144 and hence the new names contained in the electronic version are effectively published under that
 145 Code from the electronic edition alone. This published work and the nomenclatural acts it
 146 contains have been registered in ZooBank, the online registration system for the ICZN. The
 147 ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed
 148 through any standard web browser by appending the LSID to the prefix <http://zoobank.org/>. The
 149 LSID for this publication is: urn:lsid:zoobank.org:pub:A7FFDF5C-6CD5-41CA-B106-
 150 BFF6BDFCB258; for *Monomorium heggyi* **sp. n.** is urn:lsid:zoobank.org:act:B57EC2EA-1781-
 151 4C19-ADFE-757C834E2774; and for *Monomorium khalidi* **sp. n.** is
 152 urn:lsid:zoobank.org:act:3B5BB529-D842-4146-B8F7-ACCAC9CD5BA7. The online version
 153 of this work is archived and available from the following digital repositories: PeerJ, PubMed
 154 Central and CLOCKSS.

RESULTS

I. Diagnosis of Arabian *Monomorium*

Members of the genus *Monomorium* can be recognized by combination of the following characters in the worker caste (Bolton, 1987, 1994; DuBois, 1986; Fisher & Bolton, 2016; Sparks *et al.*, 2019): monomorphic to polymorphic; antennae 10–12 segmented (most frequently 12), conspicuous 3-segmented club; mandibles with 3–4 teeth; palp formula 2,2, or 1,2; median clypeal seta conspicuous; median portion of clypeus raised, the raised portion usually longitudinally bicarinate; lateral portions of the clypeus not elevated as shield-like ridges anterior to the antennal toruli; frontal carinae distinct, but absent posterior to the medial arches of the antennal toruli; metanotal groove present, commonly impressed; propodeal dorsum usually unarmed and rounding into the declivity; propodeal spiracle usually circular, located at about the midlength of the sclerite; abdominal segment IV (metasomal III, gastral I) with posttergite overlapping poststernite.

Diagnosis of Arabian *Monomorium* species-groups

The *M. monorium* species-group can be readily recognized by the following combination of characters in the worker caste (Sharaf *et al.* 2018a): monomorphic, with size variation; head longer than broad; mandibles smooth and masticatory margin armed with four teeth; antennae with 10–12 segments, terminating in a three-segmented club; median clypeal portion raised anteriorly and longitudinally bicarinate; anterior clypeal margin without a pair of teeth; eyes present with variable size, located in front of the midlength of the sides in full-face view, and with four or more ommatidia in the longest row; head smooth and shining; metanotal groove well-defined, with distinct cross-ribs; propodeal spiracle circular to subcircular; propodeal dorsum meeting declivity in a rounded angle; promesonotum and propodeal dorsum smooth; body pilosity variable but usually distinct; petiole, postpetiole and gastral tergites usually smooth.

The *M. salomonis* species-group can be diagnosed by the following character states in the worker caste (Bolton, 1987): monomorphic, with minor size variation; palp formula 2,2 or 1,2 in some minute species; mandibles usually sculptured; masticatory margins of mandibles armed with four teeth which decrease in size from apex to base; median clypeal portion raised,

projecting anteriorly; cephalic dorsum usually sculptured, ranging from dense blanketing reticulate-punctuation to feeble superficial reticular patterning; eyes prominent of medium to large size, usually with six or more ommatidia in the longest row; eyes circular to oval in shape; head longer than broad; scapes usually relatively long ($SI > 80$); metanotal groove moderately impressed to absent; metanotal cross-ribs inconspicuous to absent; propodeal spiracle circular to subcircular; propodeum rounded to angular between dorsum and declivity; propodeal dorsum usually sculptured but never transversely striate; petiolar spiracle situated at the node or immediately in front of the anterior face of the node; body pilosity variable in distribution and density, but usually reduced on the head and mesosoma; mesosoma, petiole and postpetiole usually sculptured; first gastral tergite usually shagreenate or finely sculptured.

II. Synoptic checklist of Arabian *Monomorium*

Monomorium monomorium species-group

1. *Monomorium aeyade* Collingwood & Agosti, 1996
2. *Monomorium brunneolucidulum* Collingwood & Agosti, 1996 (**nomen dubium**).
3. *Monomorium carbonarium* (Smith, 1858)
= *Monomorium minutum* Mayr, 1862
4. *Monomorium clavicorne* André, 1881
= *Monomorium clavicorne punicum* Santschi, 1915a
5. *Monomorium exiguum* Forel, 1894
= *Monomorium exiguum bulawayensis* Forel, 1913
= *Monomorium faurei* Santschi, 1915b
= *Monomorium exiguum flavescens* Forel, 1916
= *Monomorium baushare* Collingwood & Agosti, 1996
= *Monomorium qarahe* Collingwood & Agosti, 1996
6. *Monomorium floricola* (Jerdon, 1851)
= *Monomorium poecilum* Roger, 1863
= *Monomorium cinnabari* Roger, 1863
= *Monomorium specularis* Mayr, 1866
= *Monomorium impressum* Smith, 1876
= *Monomorium floreanum* Stitz, 1932

- 217 = *Monomorium angusticlava* Donisthorpe, 1947
- 218 7. *Monomorium holothir* Bolton, 1987
- 219 8. *Monomorium mohammedi* Sharaf & Hita Garcia, 2018
- 220 9. *Monomorium rimae* Collingwood & Agosti, 1996
- 221 10. *Monomorium sarawatense* Sharaf & Aldawood, 2013b
- 222 ***Monomorium salomonis* species-group**
- 223 11. *Monomorium abeillei* André, 1881
- 224 = *Monomorium wahibiense* Collingwood & Agosti, 1996 **syn. n.**
- 225 12. *Monomorium acutinode* Collingwood & Agosti, 1996
- 226 13. *Monomorium areniphilum* Santschi, 1911
- 227 = *Monomorium fezzanense* Collingwood & Agosti, 1996 **syn. n.**
- 228 = *Monomorium hemame* Collingwood & Agosti, 1996 **syn. n.**
- 229 = *Monomorium marmule* Collingwood & Agosti, 1996 **syn. n.**
- 230 = *Monomorium salomonis lepineyi* Santschi, 1934
- 231 = *Monomorium salomonis pullula* Santschi, 1919
- 232 14. *Monomorium asiriense* Collingwood & Agosti, 1996
- 233 15. *Monomorium barbatulum* Mayr, 1877 (New record to KSA)
- 234 16. *Monomorium bicolor* Emery, 1877
- 235 = *Monomorium bicolor aequatoriale* Santschi, 1926
- 236 = *Monomorium bicolor coerulescens* Santschi, 1912
- 237 = *Monomorium bicolor rufibasis* Santschi, 1914
- 238 = *Monomorium bicolor tropicale* Santschi, 1926
- 239 = *Monomorium bicolor uelense* Santschi, 1926
- 240 = *Monomorium phoenicum* Santschi, 1927 **syn. n.**
- 241 17. *Monomorium buettikeri* Collingwood & Agosti, 1996
- 242 18. *Monomorium buxtoni* Crawley, 1920a
- 243 19. *Monomorium carbo* Forel, 1910
- 244 20. *Monomorium dammame* Collingwood & Agosti, 1996
- 245 21. *Monomorium dirie* Collingwood & Agosti, 1996
- 246 22. *Monomorium elghazalyi* Sharaf & Aldawood, 2017
- 247 23. *Monomorium fayfaense* Collingwood & Agosti, 1996

24. *Monomorium gallagheri* Collingwood & Agosti, 1996
25. *Monomorium hanaqe* Collingwood & Agosti, 1996
26. *Monomorium harithe* Collingwood & Agosti, 1996
- = *Monomorium najrane* Collingwood & Agosti, 1996 **syn. n.**
27. *Monomorium hegyi* Sharaf, **sp. n.**
28. *Monomorium jizane* Collingwood & Agosti, 1996
29. *Monomorium khalidi* Sharaf, **sp. n.**
30. *Monomorium knappi* Collingwood & Agosti, 1996
31. *Monomorium luteum* Emery, 1881
32. *Monomorium mahyoubi* Collingwood & Agosti, 1996
33. *Monomorium moathi* Sharaf & Collingwood, 2010
34. *Monomorium niloticum* Emery, 1881
- = *Monomorium matame* Collingwood & Agosti, 1996 **syn. n.**
35. *Monomorium nimihil* Collingwood, 2004
36. *Monomorium nitidiventre* Emery, 1893
- = *Monomorium yemene* Collingwood & Agosti, 1996 **syn. n.**
37. *Monomorium pharaonis* (Linnaeus, 1758)
- = *Monomorium antiguensis* (Fabricius, 1793)
- = *Monomorium domestica* (Shuckard, 1838)
- = *Monomorium contigua* (Smith, 1858)
- = *Monomorium fragilis* (Smith, 1858)
- = *Monomorium minuta* (Jerdon, 1851)
- = *Monomorium vastator* (Smith, 1857)
38. *Monomorium riyadhe* Collingwood & Agosti, 1996
39. *Monomorium salomonis* (Linnaeus, 1758)
- = *Monomorium debilis* (Walker, 1871)
- = *Monomorium salomonis obscuratum* Stitz, 1917
- = *Monomorium thorense* Mayr, 1862
40. *Monomorium subdenticorne* Collingwood & Agosti, 1996
41. *Monomorium subopacum* (Smith, 1858), full synonymy in Heterick (2006)
42. *Monomorium suleyile* Collingwood & Agosti, 1996

43. *Monomorium tumaire* Collingwood & Agosti, 1996

44. *Monomorium venustum* (Smith, 1858)

III. Key to Arabian *Monomorium* based on the worker caste

Note: *M. brunneolucidulum* excluded due to lack of diagnostic characters.

1. Head smooth and glossy (Fig. 1A) ... 2 (*M. monomorium* species-group)

– Head **usually** sculptured, ranging from dense reticulate-punctate, longitudinal striations, to faint reticular patterning (Fig. 1B) ... 10 (*M. salomonis* species-group)

2. Anterior median portion of clypeus strongly concave with two laterally projecting teeth (Fig. 1C) ... 3

– Anterior median portion of clypeus straight or feebly concave without raised ridges (Fig. 1D) ... 4

3. Uniform dark brown to black; metanotal groove broadly and deeply impressed (Fig. 1E) ... *M. carbonarium*

– Bicolored with yellow mesosoma contrasting with the black gaster; metanotal groove shallowly impressed ... *M. rimae*

4. Antenna 12-segmented ... 5

– Antenna 11-segmented ... 7

5. Body pilosity clubbed; mesosoma, petiole and postpetiole distinctly sculptured (Fig. 1F) ... *M. sarawatense*

– Body pilosity simple; mesosoma, petiole and postpetiole smooth and shining (Fig. 2A) ... 6

6. Bicolored, with head and gaster dark brown to black contrasting the yellow or yellow-brown mesosoma and petiole; eyes small (EL 0.21–0.24 x HW) with six ommatidia in longest row, oval in profile (Fig. 2A) ... *M. floricola*

– Uniform yellow; eyes relatively large (EL 0.30–0.32 x HW) with 8-9 ommatidia in longest row, in profile with convex dorsal margin and straight ventral margin (Fig. 2B) ... *M. holothir*

310

311 7. Mesosoma without standing hairs (Fig. 2C) ... 8

312 – Mesosoma with standing hairs (Fig. 2D) ... 9

313

314 8. Eyes larger, with a ring of seven to eight ommatidia encircling a single row of 2 ommatidia,

315 and in profile closer to mandibular insertions (EM 0.05); meso- and metapleuron smooth;

316 petiole and postpetiole smooth and each with one pair of standing hairs (Fig. 2C) ... *M. aeyade*

317 – Eyes smaller, with only 5–6 ommatidia total, and in profile further away from mandibular

318 insertions (EM 0.09–0.11); meso- and metapleuron finely shagreened; petiole and postpetiole

319 superficially shagreened and without standing hairs (Fig. 2E) ... *M. mohammedi*

320

321 9. Mesosoma with two pairs of standing hairs, one on pronotal corners and one on propodeum

322 (Fig. 2F) ... *M. clavicorne*

323 – Mesosoma with several pairs of standing hairs, about 10 pairs (Fig. 2D) ... *M. exiguum*

324

325 10. Underside of head with crowded J-shaped hairs forming a distinct psammophore (Fig. 3A) ...

326 *M. barbatulum*

327 – Underside of head with sparse hairs, psammophore absent ... 11

328

329 11. Mesosoma without standing hairs ... 12

330 – Mesosoma with standing hairs ... 25

331

332 12. Uniform yellow, brown, or yellow-brown, gaster not darker than mesosoma ... 13

333 – Bicolored, with gaster distinctly darker than mesosoma ... 18

334

335 13. Metanotal groove shallowly impressed or indistinct (Fig. 3B) ... 14

336 – Metanotal groove deeply impressed (Fig. 3C) ... 15

337

338 14. Small yellow species (TL 1.7, HL 0.45–0.48, HW 0.34–0.36, SI 92–97); cephalic surface

339 with vestigial or superficial reticular patterning, almost entirely effaced; petiole and postpetiole

- without standing hairs; (Fig. 3B); first gastral tergite completely glabrous, *i.e.*, without standing hairs (Fig. 3B) ... ***M. hegyi* sp. n.**
- Large brown species (TL 2.20–2.75, HL 0.57–0.65, HW 0.51, SI 93–103); cephalic surface with fine and dense reticulate-rugulose sculpture; petiole and postpetiole each with two pairs of standing hairs (Fig. 3D); first gastral tergite always with several pairs of standing hairs (Fig. 3D) ... ***M. harithe***
- 15.** Eyes located nearly at the mid-length of head as seen in full-face view (Fig. 3E); eyes smaller, with 5 ommatidia in longest row); posterior margin of head distinctly concave in full-face view (Fig. 3E) ... ***M. elghazalyi***
- Eyes located nearly behind or at head mid-length as seen in full-face view (Fig. 3F); eyes distinctly larger, with 10–14 ommatidia in longest row); posterior margin of head concave or linear in full-face view ... **16**
- 16.** Underside of head without long, standing hairs (Fig. 4A) ... ***M. dirie***
- Underside of head with numerous pairs of long, standing hairs (Fig. 4B) ... **17**
- 17.** Yellow; scapes just reaching posterior margin of head when laid back; body pilosity abundant over entire body ... ***M. nimihil***
- Brown; scapes surpassing posterior margin of head by about half the length of the pedicel when laid back; body pilosity distinctly reduced over entire surface, mesosoma with a single pair of hairs on propodeum, while the petiole, postpetiole, and gaster are bare ... ***M. salomonis***
- 18.** First gastral tergite without standing hairs ... ***M. moathi***
- First gastral tergite with hairs either scattered on tergite surface or apically on the posterior margin ... **19**
- 19.** Scapes when laid back from their insertions reach or surpass posterior margin of head in full-face view ... **20**
- Scapes when laid back from their insertions fail to reach posterior margin of head in full-face view ... **21**

- 20.** Propodeal dorsum in profile meeting declivity forming two blunt, slightly projecting angled bosses (Fig. 4C) ... *M. subdenticorne*
- Propodeal dorsum in profile meeting declivity in a continuous curve (Fig. 4D) ... *M. bicolor*
- 21.** Petiole in the form of a high triangle in profile with anterior face appears as a continuous line sloping anteriorly (Fig. 4E) ... *M. acutinode*
- Petiole broadly rounded with anterior face sloping downward and then anteriorly to the peduncle (Fig. 4F) ... **22**
- 22.** Head faintly superficial sculptured, slightly shining ... *M. venustum*
- Head densely and finely reticulate to reticulate-shagreenate ... **23**
- 23.** Small (TL 2.3–2.4, HL 0.60–0.63, HW 0.43–0.45, PW 0.30–0.31, ML 0.66–0.70); mesosoma and gaster approximately of the same color, the two not strongly contrasting ... *M. carbo*
- Larger (TL 3.1–3.4, HL 0.70–1.04, HW 0.54–0.88, PW 0.36–0.53, ML 0.88–1.24); mesosoma and gaster conspicuously differently colored, gaster usually darker ... **24**
- 24.** Eyes with 12–14 ommatidia in longest row); metanotal groove deeply impressed (Fig. 5A); postpetiole with two to three pairs of backward directed hairs ... *M. areniphilum*
- Eyes with 9–11 ommatidia in longest row); metanotal groove feebly impressed (Fig. 5B); postpetiole with a single pair of backward directed hairs ... *M. subopacum*
- 25.** Large (TL ≥ 3.8 , HW > 0.75) ... **26**
- Smaller (TL 2.2–3.2, $\leq$ HW 0.67) ... **28**
- 26.** Entirely yellowish; mesosoma rather flat with a shallow oblique metanotal groove (Fig. 5C) ... *M. luteum*
- Gaster dark contrasting with red mesosoma; metanotal groove steeply angled (Fig. 5D) ... **27**

27. Head smooth with superficial sculpture; the first of the three segments forming the club being shorter than the second (Fig. 5E); head in full-face view with feebly but distinctly convex sides; head in full-face view with eyes fail or just break head sides (Fig. 5E) ... *M. niloticum*
- Head completely finely striate (Fig. 5F); the first of the three segments forming the club nearly subequal to the second (Fig. 5F); head in full-face view with straight sides; head in full-face view with eyes break sides (Fig. 5F) ... *M. riyadhe*
28. Mesosoma red, contrasting with dark gaster ... 29
- Mesosoma pale brown to black, concolorous with gaster ... 34
29. Mesosoma hairs restricted to one pair on pronotum ... 30
- Mesosoma with several pairs of hairs scattered over whole dorsum ... 31
30. Head and mesosoma dull red, gaster brown; scapes reaching posterior head margin when laid back (Fig. 6A); cephalic surface with vestigial sculptures; posterior margin of head with a single pair of hairs (Fig. 6A); propodeum in profile with dorsum making an obtuse angle with declivity (Fig. 6B) ... *M. hanage*
- Head and mesosoma bright orange red, gaster black; scapes not reaching posterior head margin when laid (Fig. 6C); cephalic surface densely punctate; posterior margin of head without hairs except for appressed pubescence (Fig. 6C); propodeum in profile with dorsum making a continuous curve with declivity (Fig. 6D) ... *M. jizane*
31. Propodeal dorsum with a single pair of standing hairs ... 32
- Propodeal dorsum with at least three pairs of standing hairs ... 33
32. Scapes when laid back from their insertions reach posterior margin of head in full-face view (Fig. 6E); cephalic surface densely punctate (Fig. 6E); gaster smooth and shining (Fig. 6F) ... *M. fayfaense*
- Scapes when laid back from their insertions fail to reach posterior margin of head in full-face view (Fig. 7A); cephalic surface with vestigial sculptures (Fig. 7A); gaster finely densely shagreened and dull (Fig. 7B) ... *M. knappi*

432

433 **33.** Whole body with abundant fine hairs (Fig. 7C); with head in full-face view outer margins of
434 eyes break head sides (Fig. 7D) ... *M. nitidiventre*

435 – Body pilosity limited and stiff (Fig. 7E); promesonotum with five to six pair of standing hairs,
436 promesonotum and propodeum each with three pairs; petiole and postpetiole each with two-
437 three pairs; with head in full-face view outer margins of eyes fail to break head sides (Fig. 7F)
438 ... *M. khalidi* sp. n.

439

440 **34.** Whole body including gaster densely sculptured and dull ... **35**

441 – Gaster at least more or less shining with superficial sculpture ... **37**

442

443 **35.** Head, mesosoma, and waist segments very light brown or yellow; hairs on mesosoma
444 scattered (Fig. 8A) ... *M. pharaonis*

445 – Head, mesosoma, and waist segments conspicuously of darker brown or black; mesosoma hairs
446 mainly or entirely to pronotum (Fig. 8B) ... **36**

447

448 **36.** Body color brown; larger species (TL 3.1–3.3, HW 0.70); scapes long, surpassing posterior
449 head margin by about the length of the pedicel when laid back (Fig. 8C); head in full-face view
450 with convex sides (Fig. 8C) ... *M. suleyile*

451 – Body color uniformly black; smaller species (TL 2.7–2.9, HW 0.56); scapes shorter, just
452 reaching posterior head margin when laid back of head (Fig. 8D); head in full-face view with
453 parallel sides (Fig. 8D) ... *M. mahyoubi*

454

455 **37.** Whole body glossy, nodes and gaster brilliant ... *M. dammame*

456 – At least mesosoma with close punctate sculpture ... **38**

457

458 **38.** Underside of head with numerous hairs (12–16), the longest exceeding the maximum eye
459 length (Fig. 8E) ... *M. tumaire*

460 – Underside of head with fewer hairs, none as long as maximum eye length (Fig. 8F) ... **39**

461

462 **39.** Head finely and densely punctate, general appearance dull ... **40**

– Head feebly superficially sculptured, relatively but distinctly shining ... **41**

40. Smaller species (TL 2.4–2.7); clypeal border feebly concave to straight with a very small median notch or none (Fig. 9A); mesonotum straight in profile (Fig. 9B) ... *M. gallagheri*

– Larger species (TL > 3.0); clypeus with a distinct anteromedian notch (Fig. 9C); mesonotum distinctly convex in profile (Fig. 9D) ... *M. buxtoni*

41. Scapes distinctly short, surpassing compound eye posterior margins by only about the length of the pedicel ... *M. buettikeri*

– Scapes longer, reaching posterior head margin ... **42**

42. Color dark brown to black; smaller species (TL 2.40–2.75, CI 78–91); metanotal groove shallowly impressed (Fig. 9E) ... *M. abeillei*

– Color light to median brown; larger species (TL 3.2, CI 69–71); metanotal groove distinctly impressed (Fig. 9F) ... *M. asiriense*

IV. New taxonomic treatments

New site records include, when available, geo-coordinates (°N, °E), elevation (m), collection date, collector, and number worker (w) and queen (q) specimens.

Monomorium abeillei André, 1881

(Fig. 10 A–C)

Monomorium abeillei André, 1881: 531 (footnote) (w.) Israel. Palearctic. (MNHN, CASENT0915411). [Image of lectotype worker examined]. [Also described as new by André, 1881: 67.].

Combination in *Monomorium* (*Xeromyrmex*): Emery, 1922: 177; subspecies of *Monomorium salomonis*: Forel, 1910a: 23; Hamann & Klemm, 1967: 413; revived status as species: Collingwood, 1985: 269; Collingwood & Agosti, 1996: 340.

Monomorium wahibiense Collingwood & Agosti, 1996: 357 (w.) Oman. Palearctic.

[NHMB], CASENT0913864. Syntype worker, Oman, Wahiba dunes (21.vii.1985, M. D.

Gallagher) [examined] **syn. n.**

Material examined. KSA: Asir Province: Raydah (18.198, 42.410, 2443m, 22.ii.2014, M.R. Sharaf, 3w); Raydah (18.204, 42.412, 2820m, 21.ii.2014, M.R. Sharaf, 6w); **Riyadh Province:** Huraymila, Buaythiran (25.149, 45.950, 07.ii.2011, M.R. Sharaf, 2w); Rawdhat Khureim (25.383, 47.277, 618m, 02.vi.2013, S. Salman, 6w); Rawdhat Khureim (25.425, 47.235, 579m, 09.i.2015, S. Salman, 12w); Dirab, KSU research station (24.419, 46.654, 568m, 05.xii.2013, S. Salman, 1w); Wadi Hanifa (24.670, 46.654, 657m, 14.ii.2014, S. Salman, 3w); Mezahmyia (24.472, 46.239, 633m, 25.i.2014, S. Salman, 1w); Mezahmyia (24.466, 46.251, 648m, 29.xi.2014, S. Salman, 1w); Al Hayer (24.546, 46.742, 647m, S. Salman, 2w); Al Hayer (24.557, 46.744, 589m, 11.iv.2014, S. Salman, 6w); Runnah (25.571, 46.973, 615m, 12.iv.2014, S. Salman, 1w); Dawademi (24.478, 44.364, 1027m, 18.iv.2014, S. Salman, 1w); Dawademi (24.583, 44.323, 966m, 16.i.2015, S. Salman, 3w); Dawademi (24.538, 44.355, 999m, 16.i.2015, S. Salman, 1w); Afif (23.766, 42.840, 1015m, 19.iv.2014, S. Salman, 6w); Afif (24.302, 43.688, 951m, 19.iv.2014, S. Salman, 1w); Afif (23.900, 42.081, 1052m, 17.i.2015, S. Salman, 15w); Afif (23.957, 42.976, 1059m, 17.i.2015, S. Salman, 1w); Irgah (24.6710, 46.593, 625m, 19.i.2015, S. Salman, 2w); Thadiq (25.294, 45.871, 735m, 26.iv.2014, S. Salman, 1w); Quwayia (24.047, 45.244, 854m, S. Salman, 7w); Shaqra (25.326, 45.233, 710m, 30.v.2014, S. Salman, 9w); Shagra (25.230, 45.319, 703m, 24.i.2015, S. Salman, 1w); Shaqra (25.270, 45.291, 712m, 23.i.2015, 2w); Durma (24.607, 46.130, 646m, 30.i.2015, S. Salman, 1w); Majma'a (25.880, 45.365, 730m, 07.ii.2015, S. Salman, 5w); Kharrarah (24.392, 46.244, 726m, 08.iv.2015, S. Salman, 15w); Al Ghat (26.066, 44.919, 653m, 31.x.2015, S. Salman, 1w); Hawtet Sudeir (25.592, 45.612, 732m, 31.i.2015, S. Salman, 1w); KSU campus (24.737, 46.618, 662m, 29.ii.2012, K. Mahmoud, 1w); Dirab (24.419, 46.654, 804m, 18.ix.2014, S. Salman, 9w); Hareeq (23.614, 46.054, 689m, 22.ii.2015, S. Salman, 1w); Quwayia (24.058, 45.245, 846m, 29.xi.2014, S. Salman, 1w); Salbouxh (25.078, 46.347, 716m, 26.xii.2014, S. Salman, 8w); **Jazan Province:** Sajid Island, Al-Sajid (16.860, 41.932, 05.iii.2017, U. Abuelgheit, 1w).

Remarks. *Monomorium wahibiense* is represented by a single worker deposited in WMLC and accompanied by a red card and handwritten label by C. Collingwood indicating that this specimen represents the syntype. The label's data are consistent with the data for the type in the original description in terms of collecting locality (Oman, Wahiba sand) and collector (M. D. Gallagher), but not the collection date, which we consider a typographical error.

Comparing the mentioned type material with the image of the type material of *M. abeillei* André, we found the two species share the same morphological characters, which can be summarized as follow: scapes relatively short, when laid back from their insertions just reaching posterior head margin; eyes of moderate size with about 10 ommatidia in longest row); cephalic surface between frontal lobes faintly striated; promesonotum and mesonotum forming continuous flat line in profile; mesosoma with single pair of standing hairs on pronotal humeral angles; metanotal groove feebly impressed; petiole with single pair of backward directed hairs; postpetiole with two pairs. Consequently, we propose treating *M. wahibiense* as a junior synonym of *M. abeillei* André.

Geographic Distribution. *Monomorium abeillei* is originally described from Israel and recorded from several countries in the Middle East, including the Arabian Peninsula (KSA, Kuwait, Oman and Yemen) (Collingwood 1985, Collingwood & Agosti 1996), Iran (Paknia et al. 2008), Israel (Vonshak & Ionescu-Hirsch 2009) and North Africa (Borowiec 2014).

Monomorium areniphilum Santschi, 1911

(Fig. 11 A–C)

Monomorium Salomonis var. *areniphila* Santschi, 1911: 84 (w.) Tunisia. Palearctic. [NHMB], CASENT0249829, [Syntype worker, examined].

Emery, 1915: 378 (q.); combination in *Monomorium* (*Xeromyrmex*): Emery, 1922: 177; subspecies of *Monomorium salomonis*: Santschi, 1936a: 50; raised to species: Collingwood, 1985: 269; senior synonym of *Monomorium lepineyi*, *Monomorium pullula*: Bolton, 1987: 336.

Senior synonym of *Monomorium fezzanense* Collingwood & Agosti, 1996: 346 (w.) Saudi Arabia. Afrotropic. [NHMB], Syntype worker, Saudi Arabia, 31 km NW Tabuk (24.iv.1979, CASENT0913557) [examined] **syn. n.**

Senior synonym of *Monomorium hemame* Collingwood & Agosti, 1996: 348 (w.) Kuwait. Palearctic. [WMLC], holotype worker, Kuwait, Umm Al-Hemam (9.III.1988, W. Biittiker, CASENT0922316) [examined]; paratype worker, Saudi Arabia, Uyaynah (01.IV 1976, W. Biittiker, CASENT0913800) [examined] **syn. n.**

Senior synonym of *Monomorium marmule* Collingwood & Agosti, 1996: 349, fig. 21 (w.)

OMAN. Palearctic. Paratype worker, Oman, Minririb (14.i.1986, M.D. Gallagher,

CASENT0913824, NHMB) [examined] **syn. n.**

Remarks. A thorough examination of the type material of *M. fezzanense*, *M. hemame*, *M. marmule*, and *M. areniphilum* yielded no evidence for heterospecificity; they are indistinguishable. All four taxa share the following characters: median portion of anterior clypeal margin shallowly concave; eyes large with 12-15 ommatidia in longest row; promesonotum and anterior portion of mesonotum in profile feebly convex; posterior portion of mesonotum sloping steeply to broadly and deeply impressed metanotal groove; mesosoma without hairs; petiole with single pair of backward directed hairs; postpetiole with two pairs of hairs.

Herein *M. fezzanense*, *M. hemame*, and *M. marmule* are treated as junior synonyms of *M. areniphilum*. It is worth mentioning that in the original description of *M. marmule*, Collingwood & Agosti (1996) gave a brief differential diagnosis with *M. areniphilum* based on variable characters such as the presence of mesosomal pubescence, the petiole and postpetiole color and pilosity.

Geographic Distribution. A species originally described from Tunisia and recorded from most countries of the Arabian Peninsula including KSA, Kuwait, Oman, and Yemen (Collingwood, 1985; Collingwood & Agosti, 1996), and the UAE (Collingwood et al., 2011). It is also reported from North Africa and the Afrotropical Region (Bolton, 1987).

Monomorium barbatulum Mayr, 1877

(Fig. 12 A–C)



Monomorium barbatulum Mayr, 1877: 17 (w.) Kazakhstan. Palearctic.

Material examined. KSA: Riyadh Province: Zulfi (26.367, 44.986, 670m, 18.i.2014, Al Dhafer et al., 1w: CASENT0922263, KSMA).

Geographic Distribution. This species was originally described from Kazakhstan and recorded from Oman (Collingwood, 1985; Collingwood & Agosti, 1996), the UAE (Collingwood et al., 2011), and Israel (Vonshak & Ionescu-Hirsch, 2009). The present material represents a new record to the KSA.

Monomorium bicolor Emery, 1877

(Fig. 13 A–C)

Monomorium bicolor Emery, 1877: 368 (w.) Eritrea. Afrotropic.

Senior synonym of *Monomorium phoenicum* Santschi, 1927e: 242 (w.q.) Lebanon.

Palaearctic. Syntype worker, Lebanon, Beyroth (05.viii.1933, Santschi, CASENT0249831,

NHMB) [examined] **syn. n.**

Material examined. KSA: Asir Province: Almajardah, Wadi Eltalalei (19.003, 41.732, 223m, 10.xi.2012, M.R. Sharaf, 1w); Wadi Shahadan (17.472, 42.856, 452m, 13.xi.2012, M.R. Sharaf, 13w, (1w, CASENT0906396); Allaith, Adam, Wadi Elarj (20.453, 40.816, 450m, 09.xi.2012, M.R. Sharaf, 45w, 1w: CASENT0906395, 1q: CASENT0906394); Wadi Aljora, near Abadan (17.293, 43.070, 465m, 12.xi.2012, M.R. Sharaf, 2w); Almajardah, Wadi Bagara (18.793, 42.019, 436m, 10.xi.2012, M.R. Sharaf, 2w); **Jazan Province:** Abu Arish (17.013, 42.802, 90m, 10.iv.2012, M.R. Sharaf, 11w); Sabia (17.107, 42.650, 43m, 09.iv.2012, M.R. Sharaf, 23w); Jazan (16.97627, 42.61743, 38m, 12.iv.2012, M.R. Sharaf, 12w); **Al Bahah Province:** Dhi Ayn Archeological village (19.930, 41.443, 741m, 18.v.2011, M.R. Sharaf, 3w); Wadi Gonouna (19.429, 41.605, 353m, 12.v.2011, M.R. Sharaf, 10w: KSMA).

Remarks. The type material of *M. bicolor* and *M. phoenicum* are clearly conspecific. They share the same diagnostic characters as follow: scapes relatively long, when laid back from their insertions surpassing posterior head margin by about length of pedicel; head in full-face view with eyes just breaking sides; cephalic surface dull, finely and densely punctate; median anterior clypeal margin distinctly concave; area between frontal carinae finely longitudinally striated; mesosoma without standing hairs; metanotal groove acutely impressed; propodeal dorsum in profile meeting declivity in continuous curve; petiole and postpetiole each with single pair of back directed hairs; first gastral tergite with hairs scattered over tergite surface; biocolored species, with head, mesosoma, petiole and postpetiole yellow-red or yellow-brown, gaster dark brown to black. Herein, we propose *M. phoenicum* as a junior synonym of *M. bicolor*.

Geographic Distribution. *Monomorium bicolor* was originally described from Eritrea and is a widespread species commonly encountered in open, sandy areas through the Afrotropical Region (Bolton, 1987). In the Arabian Peninsula, it is known from the KSA and the UAE (Collingwood, 1985; Collingwood & Agosti, 1996; Collingwood et al., 2011).

Monomorium brunneolucidulum Collingwood & Agosti, 1996

Monomorium brunneolucidulum Collingwood & Agosti, 1996: 343. Oman. Palearctic.

Remarks. In their brief original description of the enigmatic species *M. brunneolucidulum* from Oman, Collingwood & Agosti (1996) neither gave successful diagnostic characters nor illustrations for species recognition. In addition, the type-material is apparently lost. Due to a lack of type material and species diagnostic characters, it is impossible to confirm the identity of the species and it remains as a nomen dubium.

***Monomorium floricola* (Jerdon, 1851)**

(Fig. 14 A–C)

Atta floricola Jerdon, 1851: 107 (w.) India. Indomalaya.

Forel, 1893: 388 (q.m.); Wheeler, 1905: 88 (q.m.); Donisthorpe, 1914: 136

(gynandromorph); Crawley, 1920: 217 (gynandromorph); Wheeler & Wheeler, 1955c:

121 (l.). Combination in *Monomorium*: Mayr, 1879: 671.

Senior synonym of *Monomorium poecilum*: Emery, 1894d: 151.

Senior synonym of *Monomorium specularis*: Mayr, 1879: 671.

Senior synonym of *Monomorium cinnabari*: Wheeler, 1913: 486.

Senior synonym of *Monomorium floreanum*: Brown, in Linsley & Usinger, 1966: 175.

Senior synonym of *Monomorium angusticlava*: Bolton, 1987: 390.

Senior synonym of *Monomorium impressum*: Bolton, 1987: 390.

Senior synonym of *Monomorium floricola furina*: Heterick, 2006: 122.

Senior synonym of *Monomorium floricola philippinensis*: Heterick, 2006: 122.

Material examined. Oman: Dhofar Province: Ayn Dirbat (17.106, 54.453, 207m, 17.xi.2017, M.R. Sharaf, 4w: KSMA, 1w: CASENT0922876, CASC).

Geographic Distribution. This species was originally described from India. It is a successful tramp species of putative Southeast Asian origin that is widely distributed throughout tropical and subtropical regions worldwide (Deyrup et al., 2000; Wetterer, 2010a). The present material represents a new record for Oman and the Arabian Peninsula.

***Monomorium harithe* Collingwood & Agosti, 1996**

(Fig. 15 A–C)

Monomorium harithe Collingwood & Agosti, 1996: 347 (w.) Saudi Arabia. Afrotropic. Holotype worker: Saudi Arabia, desert near Najran (17.533, 44.000, 10.iv.1983, C.A. Collingwood, CASENT0922335, WMLC) [examined] **Syn. n.**

Monomorium najrane Collingwood & Agosti, 1996: 352 (w.) Saudi Arabia. Afrotropic.

Previous records. KSA: Riyadh (24.714, 46.675, 21.i.1980, A.H. Talhouk, 2w); **Yemen:** Taiz (13.578, 44.018, 20.iii.1993, C.A. Collingwood, 2w).

Remarks. *Monomorium harithe* was described from the KSA and Yemen, while *M. najrane* was described from Najran (KSA) near the Saudi-Yemeni borders (Collingwood & Agosti 1996).

The comparison of the type material of the two species reveals a straightforward synonymy. The two species share the following characters: anterior median clypeal margin distinctly concave; scapes distinctly short, when laid back from their insertions failing to reach posterior head margin; mesosoma without standing hairs; metanotal groove feebly impressed but distinct; propodeal dorsum with distinct furrow); mesosoma, petiole and postpetiole finely and densely punctate; petiole and postpetiole each with single pair of back directed hairs; gaster smooth and shining. In addition, the two species share common body measurements (e.g. HW 0.51; SL 0.53).

Geographic Distribution. This Arabian endemic species is only known from the KSA and Yemen (Collingwood & Agosti, 1996).

Monomorium hegyi Sharaf, sp. n.

(Fig. 16 A–C)

Holotype pinned worker. KSA: Al Bahah Province: Shada Al A'la Mountain (19.877, 41.311, 897m, Al Dhafer et al., MRS0261, CASENT0746641, KSMA).

Paratype pinned workers. KSA: Al Baha Province: Shada Al A'la Mountain (19.877, 41.311, 897m, 04.vi.2014, M. R Sharaf, 4w, 1w: CASENT0746641); Shada Al A'la Mountain (19.863, 41.301, 1225m, 08.xii.2014, Al Dhafer et al., 6w, 1w: CASENT0922301); Shada Al A'la Mountain (19.843, 41.312, 1666m, 23.viii.2014, Al Dhafer et al., 1w: CASENT0906391); Shada Al A'la Mountain (19.877, 41.311, 897m, 04.vi.2014, M.R. Sharaf, MRS0261, 2w); Shada Al A'la Mountain (19.877, 41.311, 897m, 23.viii.2014, M.R. Sharaf, MRS0261, 1w: CASENT0746641); Shada Al A'la Mountain (19.863, 41.301, 1225m, 08.xii.2014, Al Dhafer et al., 1w); Shada Al A'la Mountain (19.851, 41.301, 1325m, 15.ii.2014, Al Dhafer et al., 1w); Shada Al A'la Mountain (19.839, 41.310, 15.xi.2015, Al Dhafer et al., 3w: KSMA).

Other Material. Jazan Province: Zabia (17.107, 42.650, 43m, 09.iv.2012, M. R Sharaf, MRS0070, 2w); Abu Arish (17.013, 42.802, 90m, 10.iv.2012, M. R Sharaf, MRS0073, 10w); Jazan (16.97627, 42.61743, 38m, 12.iv.2012, M. R Sharaf, MRS0077, 25w: KSMA, 1w: CASC, 1w: BMNH, 1w: WMLC, 1w: OUMC, 1w: LACM).

Measurements. Holotype: TL 2.01; HL 0.49; HW 0.39; SL 0.42; EL 0.12; EM 0.07; ML 0.59; PW 0.25; PTL 0.11; PTW 0.11; PPL 0.09; PPW 0.09; CI 80; EI 31; SI 108.

Paratype workers: TL 1.50–2.55; HL 0.42–0.49; HW 0.32–0.39; SL 0.28–0.42; EL 0.07–0.12; EM 0.07–0.09; ML 0.45–0.59; PW 0.21–0.31; PTL 0.09–0.14; PTW 0.07–0.11; PPL 0.07–0.09; PPW 0.07–0.09; CI 69–86; EI 21–31; SI 80–108 (n=13).

Diagnosis. *Monomorium hegyi* is diagnosed by the following character combination: scapes when laid back from their insertions just reaching posterior margin of head; head in full-face view with eyes located nearly at midlength; promesonotal outline feebly convex or flat, sloping posteriorly to narrow and shallowly impressed metanotal groove.

Worker. Head. Head in full-face view distinctly longer than broad, with concave posterior margin and shallowly convex sides and feebly concave posterior margin; median portion of clypeus with anterior free margin slightly indented; eyes of moderate size, in profile with convex dorsal sides and straight ventral side, maximum diameter 0.21 x–0.30 x HW, with 7 ommatidia in longest row); head in full-face view eyes located nearly at midlength of head and just breaking sides; scapes when laid back from their insertions just reaching posterior head margin.

Mesosoma. Promesonotal outline feebly convex or flat, sloping posteriorly to narrow and shallowly impressed metanotal groove; propodeal dorsum in profile convex making a continuous curve with propodeal declivity and with defined lateral margins. **Petiole.** Petiole with high

rounded node in profile; subpetiolar process broad and blunt. **Postpetiole.** Postpetiolar node lower than petiolar node in profile and nearly as broad as petiole in dorsal view. **Sculpture.**

Mandibles feebly longitudinally sculptured; cephalic surface with faint vestiges of superficial reticular patterning, almost entirely effaced, area between frontal carinae finely longitudinally striate; clypeus smooth; entire mesosoma, petiole and postpetiole sharply and densely reticulate-punctate; gastral tergites smooth and shining. **Pilosity.** Dorsum of head without standing hairs behind the level of the frontal lobes; several pairs of long hairs on the anterior clypeal margin and on mandibles; antennae with dense appressed pubescence; mesosoma, petiole and postpetiole without standing hairs of any description; first gastral tergite without standing hairs

except for sparse appressed pubescence; pilosity of remaining gastral tergites restricted to the posterior margins. **Color.** Uniformly yellow.

Remarks. *Monomorium heggyi* belongs to the *M. salomonis* species-group (Bolton 1987). It is most similar to *M. rabirium* Bolton, 1987 from Botswana from which it is readily distinguished by the longer scapes (SI 80–108) that reach the posterior head margin in full-face view, the posteriorly shifted eyes located nearly at the midlength of head in full-face view. *Monomorium rabirium* has shorter scapes (SI 92–97) that fail reaching the posterior head margin in full-face view, and eyes conspicuously located in front of midlength of head in full-face view. Among the Arabian species of the *M. salomonis* species-group, *M. heggyi* is superficially similar to *M. elghazalyi* from the Socotra Archipelago from which it can be easily separated by the larger eyes (EI 21–31), the shallowly impressed metanotal groove, and the densely sculptured mesosoma, petiole and postpetiole. *Monomorium elghazalyi* has smaller eyes (EI 19–20), broadly and deeply impressed metanotal groove, and a smooth body surface.

Etymology. The patronymic name honors Essam Heggy, the Egyptian space scientist at NASA.

Habitat. The type locality, Shada Al A’la, is a Nature Reserve (**Fig. 17**) located in the Al Bahah Province in the southwestern KSA at an elevational range of 470–2,222 m. The locality is characterized by relatively high rainfall, diverse habitats, and high biodiversity, as well as by the presence of large areas of terraced fields used for cultivating banana, coffee, figs, and lemon. The region has a diverse range of wild vegetation cover including plants of the Leguminosae (Fabaceae), composites (Asteraceae), and graminoides (Poaceae). Acacia (Fabaceae) and Juniper (Cupressaceae) are the most dominant plants (SWA, 2018, El-Hawagry et al., 2016). Shada Al-A’la harbors a high number of endemic animals including birds, mammals (SWA, 2018) and insects (El-Hawagry et al., 2016).

Geographic Distribution. KSA.

Monomorium khalidi Sharaf, sp. n.

(Fig. 18 A–C)

Holotype pinned worker. KSA: Al Bahah Province: Shada Al A’la (19.839, 41.310, 1563m, 18.x.2014, Al Dhafer et al., CASENT0922288, KSMA).

Three paratype, pinned workers, **KSA: Jazan Province:** Wadi Shahdan (17.452, 42.715, 200m, 13.xi.2012, M.R. Sharaf, MRS0131, CASENT0919810, KSMA); Fayfa, Wadi Al Jora (17.279, 43.062, 419m, 06.iv.2013, M.R. Sharaf, KSMA).

Measurements. Holotype: TL 3.16; HL 0.75; HW 0.67; SL 0.62; EL 0.14; EM 0.25; ML 0.95; PW 0.44; PTL 0.27; PTW 0.21; PPL 0.17; PPW 0.21; CI 89; EI 21; SI 93.

Paratype workers: TL 2.31–2.98; HL 0.59–0.72; HW 0.49–0.59; SL 0.45–0.59; EL 0.14–0.15; EM 0.11–0.18; ML 0.71–0.85; PW 0.32–0.41; PTL 0.28–0.32; PTW 0.11–0.15; PPL 0.11–0.17; PPW 0.08–0.15; CI 81–83; EI 24–29; SI 92–102 (n=3).

Diagnosis. *Monomorium khalidi* can be distinguished by the combination of the following characters: short scape failing to reach posterior head margin in full-face view; abundant mesosomal pilosity; straight outline of promesonotum; densely reticulate-punctate surfaces of head, mesosoma, petiole, and postpetiole; promesonotum dorsally with at least five to six pair of hairs, promesonotum and propodeum each with three pairs.

Worker. Head. Head nearly as long as broad, or little longer than broad with concave posterior margin and feebly convex sides; median portion of clypeus with anterior free margin distinctly concave; eyes of moderate size, in profile view with convex dorsal sides and straight ventral side, maximum diameter 0.20 x HW, with 9-10 ommatidia in longest row); head in full-face view with eyes failing to break head sides; scapes when laid back from their insertions failing to reach posterior margin. **Mesosoma.** Promesonotal outline flat, slopping posteriorly to narrow and feebly impressed metanotal groove; propodeal dorsum flat and short, longitudinally concave, with sharply defined lateral margins. **Petiole.** Petiole with high rounded node in profile.

Postpetiole. Postpetiole as broad as petiole in dorsal view. **Sculpture.** Cephalic surface between and immediately behind frontal lobes finely longitudinally striate; cephalic surface and sides, entire mesosoma, petiole and postpetiole sharply and densely reticulate-punctate; first gastral tergite shagreened and relatively shining. **Pilosity.** Cephalic surface with several pairs of standing hairs behind level of frontal lobes; posterior margin of head with three pairs of standing hairs; underside of head with about five pairs of short hairs; promesonotum dorsally with at least five to six pair of hairs, promesonotum and propodeum each with three pairs; petiole and postpetiole each with two-three pairs of backward directed hairs; first gastral tergite with numerous standing hairs which are evenly distributed over the sclerite in front of the apical

transverse row. **Color.** Bicolored species, head, mesosoma, petiole, postpetiole and appendage light red brown, gaster black.

Remarks. This new species is a member of the *M. salomonis* species-group (Bolton, 1987). *Monomorium khalidi* is closest to *M. junodi* Forel, 1910 from South Africa in terms of the relatively small eyes, the short scapes that fail to reach posterior margin of head, the acute metanotal groove, and densely punctate body surface. The two species have short scapes that fail to reach posterior margin of head in full-face view, metanotal groove feebly impressed; body surface densely punctate; and propodeal longitudinally concave, with sharply defined lateral margins. However, *M. khalidi* can be easily distinguished from *M. junodi* by the following characters: body bicolored with head, mesosoma, petiole, postpetiole and appendages light red-brown contrasting with the black gaster; head in full-face view with eyes failing to break sides; promesonotum dorsally with at least five to six pair of hairs, promesonotum and propodeum each with three pairs; promesonotal outline flat. *Monomorium junodi* is uniformly brown to dark brown, head in full-face view with eyes breaking sides; promesonotum dorsally with two pair of hairs, propodeum without hairs; promesonotal outline feebly but distinctly convex. Among the Arabian *Monomorium* species, *M. khalidi* is superficially similar to *M. nitidiventre* in terms of body size, surface sculpture, eye shape but the former can be readily recognized by the reduced stiff pilosity.

Etymology. The patronymic name honors Khalid Amr (born at 04/11/2012) the son of the second author.

Habitat. The type locality of *M. khalidi* is Shada Al A'la (**Fig. 19**), the same locality where *M. heggyi* was collected.

Geographic Distribution. KSA.

Monomorium niloticum Emery, 1881

(**Fig. 20 A–C**)

Monomorium niloticum Emery, 1881: 533 (w.) Egypt. Palearctic. [MSNG], [Syntype worker, CASENT0905755, image examined].

Combination in *Monomorium* (*Xeromyrmex*): Emery, 1922: 179; subspecies of *Monomorium venustum*: Forel, 1910: 6; Wheeler & Mann, 1916: 170; Stitz, 1917: 346; Finzi, 1936:

175; revived status as species: Santschi, 1936: 37; see also: Collingwood & Agosti, 1996: 352; current subspecies: nominal plus *M. n. gracilicorne*, *M. n. niloticoides*. Senior synonym of *Monomorium matame* Collingwood & Agosti, 1996: 350, fig. 22 (w.) OMAN. Palearctic, Holotype worker, Oman, Wadi Matam (01.II.1986, M.D. Gallagher, CASENT0922325, WMLC) [examined] **Syn. n.**

Material examined. KSA: Asir Province: Jebel Al Habala (18.038, 42.873, W. Buttiker, 1w); Alkhola (13.600, 44.283, 4w, WMLC). **Al Baha Province:** Al Atawla, Al Baha-Taif RD, Wadi Bawa (20.750, 41.247, 1310m, 08.xi.2012, M.R. Sharaf, MRS0099, 7w, 1w: CASENT0906397); Wadi Elzaraeb (20.073, 41.387, 2086m, 09.v.2011, M.R. Sharaf, 3w); Al Mandaq, Wadi Turabah (20.242, 41.263, 1715m, 19.ix.2011, M.R. Sharaf, 6w); Hawtat Bani Tamimc (23.525, 46.845, 19.iv.2008, M.R. Sharaf, 6w); Riyadh city (06.viii.2008, 1w); Wadi Hanifa (24.671, 46.595, 641m, 11.iv.2013, M.R. Sharaf, 11w); Al Mandaq, Wadi Turabah (20.211, 41.288, 1793m, 10.v.2011, M.R. Sharaf, 4w). **Riyadh Province:** Al Hayer (24.280, 46.766, 647m, 10.iii.2011, A.S. Aldawood, 24w); Al Hayer (24.557, 46.744, 589m, 11.iv.2014, S. Salman, 5w); Dawademi (24.557, 44.377, 983m, 18.iv.2014, S. Salman, 6w); Dawademi (24.478, 44.364, 1027m, 18.iv.2014, S. Salman, 4w); Dawademi (24.583, 44.323, 966m, 16.i.2015, S. Salman, 9w); Dawademi (24.538, 44.355, 999m, 16.i.2015, S. Salman, 1w); Afif (23.900, 42.881, 1052m, 17.i.2015, 1w); Afif (23.957, 42.976, 1059m, 17.i.2015, S. Salman, 6w); Dhurma (24.607, 46.130, 646m, 30.i.2015, S. Salman, 8w); Hawtat Bani Tamim (23.454, 46.819, 582m, 19.ii.2015, M.R. Sharaf, 3w); Hawtat Bani Tamim (23.500, 46.850, 612m, 19.ii.2015, M.R. Sharaf, 2w); Hareeq (23.614, 46.054, 689m, 22.ii.2015, S. Salman, 29w); Quwayia (24.070, 45.280, 823m, 03.v.2014, S. Salman, 3w); Majma'a (26.005, 45.019, 730m, 13.ix.2014, S. Salman, 6w, KSMA); Riyadh (23.953, 43.636, x.1979, W. Buttiker, 2w); Wadi Eflah, x.1983, W. Buttiker, 2w); wadi Mawran (22.050833, 46.671944, 10.ii.1985, W. Buttiker, 6w); Riyadh (24.7136, 46.6753, 07.vii.1975, W. Buttiker, 1w); Shoiba (Shuaibah) (20.6295, 39.5624, 06.xii.1983, W. Buttiker, 2w); Wadi Nimar (24.5705, 46.68, v.1983, W. Buttiker, 2w); Harithi (21.28, 40.28, 11.v.1984, W. Buttiker, 1w); Wadi Ellah (09.ix.1986, W. Buttiker, 3w, WMLC); Malham (25.154, 46.282, 711m, 15.ix.2014, S. Salman, 4w); Malham (25.161, 46.229, 742m, 15.ix.2014, S. Salman, 1w); Quwayia (24.058, 45.245, 846m, 29.x.2014, S. Salman, 4w); Quwayia (24.05043, 45.25795, 839m, 29.x.2014, S. Salman, 12w); Quwayia (24.053, 45.262, 836m, 29.xi.2014, S. Salman, 2w); Na'jan (24.026, 47.138, 467m, 13.xii.2014, S. Salman, 9w);

Wadi Al Dawaser (22.778, 44.786, 686m, 20.ii.2015, S. Salman, 3w); Wadi Al Dawaser (20.778, 44.786, 686m, 20.ii.2015, S. Salman, 3w); Qassim, Buraydah (26.216, 44.0414, 633m, 17.ix.2011, Steyaningrum, 9w); Qassim, Buraydah (26.330, 43.979, 623m, 19.x.2013, M.R. Sharaf, 9w); Huraymila (25.1487, 45.950, 815m, 07.ii.2011, M.R. Sharaf, 9w); Dirab Research Station (24.419, 46.654, 570m, 28.ix.2011, B.L. Fisher, 1w, CASENT0260164; Dirab Research Station (24.737, 46.618, 662m, 29.ii.2012, K. Mahmoud, 27w); Naam Dam (23.628, 46.631, 646m, 22.ii.2015, S. Salman, 30w); Dirab (24.409, 46.662, 588m, 30.xii.2009, M.R. Sharaf, 6w); Salbouxh (25.079, 46.347, 689m, 05.xi.2009, M.R. Sharaf, 15w); Salbouxh (25.074, 46.377, 728m, 26.xi.2014, S. Salman, 12w); Ghiyanah (25.074, 46.226, 728m, 26.xii.2014, S. Salman, 3w); Al Hayer (24.280, 46.766, 10.iii.2011, A.S. Aldawood, 10w); Qassim, Buraydah (26.338, 44.024, 643m, 19.x.2013, S. Salman, 2w); Wadi Al Dawaser (20.487, 44.764, 690m, 22.i.2014, S. Salman, 1w, All previous material in KSMA); Wadi Khumra (17.viii.1979, W. Buttiker, CASENT0249836, 1w, NHMB). **Mekkah Province:** Ras Hatibah (21.978, 38.937, 11.i.1983, 2w, WMLC). **OMAN:** Nakhl (23.44696, 57.88062, 364m, 02.iv.2016, M.R. Sharaf, 8w, CASENT0922306); Dhofar, Dhalkout (16.727, 53.249, 623m, 18.xi.2017, M.R. Sharaf, 6w, CASENT0922859, KSMA); no locality (2005, 1w); no locality (xi.1984, 1w); Jebel Akhdar (23.073, 54.662, 1w); Wattayah (23.591, 58.363, 1983, 3w, WMLC). **UNITED ARAB EMIRATES:** Wadi Madaq (25.300, 56.117, 22.xi-02.xii.2010, M. Hauser et al., UAE12977, 1w, CASENT0264568, KSMA); Hatta (24.806, 56.125, iii.1998, A.V. Harten, 2w); Hatta (24.806, 56.125, iii.1995, A.V. Harten, 2w, WMLC). **YEMEN:** Ta'iz (13.578, 44.018, A.V. Harten, 1w); Al Kawd (13.089, 45.365, 1991, A.V. Harten, 2w, WMLC).

Remarks. *Monomorium matame* was described from Oman and KSA based on the worker caste. In the original description Collingwood & Agosti (1996) pointed out the similarity between *M. matame* and *M. niloticum* and used variable characters that were not useful in species recognition. Our examination of the type material of both shows that *M. matame* is not separable from *M. niloticum*. The two species share the following characters: scapes when laid back from their insertions failing to reach posterior head margin; eyes relatively large about 0.30–0.33 x HW); metanotal groove deeply impressed; mesosoma with several pairs of scattered standing hairs; promesonotum with three pairs, mesonotum with two to three, propodeum with a single pair of hairs; bicolored species with head, mesosoma, petiole, postpetiole red-brown contrasting with dark brown to black gaster. Based on the examination of the type images of both species,

we propose to synonymize *M. matame* as a junior synonym of *M. niloticum*, on the basis of morphological similarity.

Geographic Distribution. *Monomorium niloticum* is originally described from Egypt and widely spread in the Arabian Peninsula (Collingwood, 1985; Collingwood & Agosti, 1996; Collingwood et al., 2011, Sharaf et al., 2018b). It is also collected from Israel (Vonshak & Ionescu-Hirsch, 2009), and North Africa (Sharaf, 2006; Borowiec, 2014). *Monomorium niloticum* is one of the most broadly spread myrmicine species throughout several countries of the Arabian Peninsula including the KSA, Oman, UAE, and Yemen (Collingwood, 1985; Collingwood & Agosti, 1996).

Ecological and Biological notes. The broad geographic distribution of the species can be interpreted in the light of the species' diverse habitat preferences, including the deserts, mountainous, and cultivated sites. Several worker series were found nesting in either dry or humid soil beneath rocks in an undisturbed site in the KSA where a broad diverse of plant species exists including *Acacia* (Fabaceae), *Citrus limon* (L.) Osbeck (Rutaceae), *Prunus dulcis* (Mill.) D. A. Webb (Rosaceae), *Juniperus* L. (Cupressaceae), *Mangifera indica* L. (Anacardiaceae), *Ficus* sp. (Moraceae), *Hibiscus* L. (Malvaceae) and *Azadirachta indica* A. Juss. (Meliaceae).

Monomorium nitidiventre Emery, 1893

(Fig. 21 A–C)

Monomorium bicolor subsp. *nitidiventris* Emery, 1893: 256 (w.) Egypt. Palearctic. [MSNG], [Syntype worker, CASENT0904602, image examined].

Mayr, 1901: 7 (q.); Karavaiev, 1911: 5 (m.); combination in *Monomorium* (*Xeromyrmex*):

Wheeler, 1922: 869; subspecies of *Monomorium bicolor*: Santschi, 1938: 39; of

Monomorium subopacum: Santschi, 1927: 245; Menozzi, 1932: 94; Bernard, 1953: 159;

Hamann & Klemm, 1967: 413; raised to species: Collingwood, 1985: 272; see also:

Wheeler & Mann, 1916: 171; Collingwood & Agosti, 1996: 352.

Monomorium yemene Collingwood & Agosti, 1996: 357, fig. 32 (w.) Yemen. Afrotropic.

Holotype worker, Yemen, Taiz (20.x.1991, A. van Harten, NHMB) [Presumed lost];

Paratype worker, Yemen, Zingibar- Shuqrah (13.356, 45.700, 21.iii.1993, C.A.

Collingwood, CASENT0913865, NHMB) [examined] **syn. n.**

Material examined. YEMEN, W. Adem Port, Wadi Tiban, N. W. of Jebel Jihaf (13.198, 44.787, ~3800 ft, 22.x.1937, C.A. Collingwood, from flower of *Adenium* sp., B. M. Exp. To S. W. Arabia, H. Scott & E. B. Britton, B. M. 1938-246, BMNH (E) 1017382, 1w, CASENT0914158, BMNH).

Remarks. The synonymy of *M. yemene* with *M. nitidiventre* is straightforward since both are morphologically similar and indistinguishable. Both share the following key characters: posterior margin emarginated in full-face view); median portion of anterior clypeal margin distinctly concave; metanotal groove deeply impressed; head, mesosoma, petiole, and postpetiole densely reticulate-punctate and covered with abundant pale standing hairs.

Note: However, the locality label of the paratype specimen (CASENT0913865) (Madinat Al shiraq) is not matching the locality data mentioned in the original description but the collecting data and collector are congruous with the description, therefore, this specimen is treated as one of the type material of *M. yemene*.

Geographic Distribution. *Monomorium nitidiventre* is originally described from Egypt and recorded from the KSA, Kuwait and Yemen (Collingwood, 1985; Collingwood & Agosti, 1996).

Monomorium subdenticorne Collingwood & Agosti, 1996

(Fig. 22 A–C)

Monomorium subdenticorne Collingwood & Agosti, 1996: 354, fig. 27 (w.) Yemen. Afrotropic.

Material examined. KSA: Asir Province: Ahad Refedah (18.134, 43.001, 2179m, 23.ii.2015, M. Alharbi, 2w); (KSMA); **Yemen:** Ghaiman, about 9 miles S. E. of San'a (13.933, 44.833, ~8400 ft, 18.ii.1938, B. M. Exp. To S. W. Arabia, H. Scott & E. B. Britton, B. M. 1938-246, BMNH1017456, 1w, CASENT0914318, BMNH).

Geographic Distribution. A species originally known from Yemen and herein collected for the first time from the KSA.

Monomorium venustum (Smith, 1858)

(Fig. 23 A–C)

Myrmica venusta Smith, 1858a: 126 (w.) Syria. Palearctic.

Material examined. KSA: Asir Province: Wadi Asidah (20.417, 41.200, 10.ix.1983, 1480m, W. Buttiker, 1w, WMLC). **Riyadh Province:** Zulfi (26.272, 44.771, 635m, 18.i.2014, S.

Salman, 15w); Mezahmyia (24.466, 46.251, 648m, 29.xi.2014, S. Salman, 10w); Dawademi (24.552, 43.932, 873m, 16.i.2015, S. Salman, 1w); Bijadriyah (24.310, 43.731, 439m, S. Salman, 2w); Afif (23.900, 42.081, 1052m, 17.i.2015, S. Salman, 18w); Shaqra (25.270, 45.291, 712m, 23.i.2015, 63w); Sajir (25.165, 44.601, 750m, 23.i.2015, S. Salman, 6w); Shaqra (25.274, 45.300, 707m, 24.i.2015, S. Salman, 2w, KSMA). **Kuwait:** Kuwait (no date, 2w, WMLC). **Oman:** Wahiba sands (21.438, 58.554, 23.iii.1986, W. Buttiker, 1w, WMLC). **United Arab Emirates:** Jebel Hafit (24.050, 55.767, 27.ii-03.iii.2011, M. Hauser et al., UAE13010, 1w, CASENT0264463); Um Al-Quwain (12. iv-07.vi.2009, M. Hauser et al., UAE12920, 1w, CASENT0264584); Ar-Rafah (25.717, 55.867, 15-30. x.2010, M. Hauser et al., UAE12866, CASENT0264475, KSMA); Wadi Asidah (20.417, 41.200, 10.ix.1983, W. Buttiker, 1w); Riyadh (24.714, 46.675, 18.ii.1975, W. Buttiker, 1w); Sharjah (25.346, 55.421, vii.2003, A.V. Harten, 2w, WMLC). **Yemen:** Sana'a (15.369, 44.191, iii.1990, 1w, WMLC).

Geographic Distribution. *Monomorium venustum* is originally described from Syria and recorded from the KSA, Kuwait, Oman (Collingwood, 1985; Collingwood & Agosti, 1996; Sharaf et al., 2018b), Israel (Vonshak & Ionescu-Hirsch, 2009), and North Africa (Borowiec, 2014).

V. Biogeographical analyses

Species distribution.

(Figs. 24–28)

***Monomorium abeillei* André.** This is a successfully distributed species in the central deserts of the KSA and no material is recorded from any other country in the Arabian Peninsula. The species appears to have a habitat preference confined to the desert areas. Only two records from the southwestern mountains of the KSA (Fig. 24).

***Monomorium niloticum* Emery, and *M. venustum* (Smith) (Figs. 25, 26).** These two species are widely spread through most countries of the Arabian Peninsula except Bahrain, Kuwait and Qatar. The former species has a broader distribution especially in the KSA and seems to be well adapted to inhabit both the desert and mountainous ecosystems of the country. We refer the lack of any record from the mentioned three countries to a deficiency in the sampling efforts, therefore it is likely that one of these two species at least, especially *M. niloticum* will be recorded from these countries.

Monomorium barbatulum Mayr, *M. floricola* (Jerdon). These two species are one of the rarest species of the genus with a single record for each from the Riyadh Province (KSA) and the Dhofar Governorate (Oman) respectively (Fig. 27).

Monomorium bicolor Emery. A mountainous species widely spread in the southwestern mountains of the KSA (Fig. 27).

Monomorium harithe Collingwood & Agosti. This species is only known from KSA and Yemen (Fig. 28).

Monomorium khalidi sp. n. This new species has a distribution restricted to the southwestern mountains of the KSA (Fig. 28).

Monomorium nitidiventre Emery. A species only recorded from Yemen. It is likely the species will be collected from the southwestern mountains of the KSA because of the occurrence of the preferred habitats (Fig. 28).

Monomorium heggyi sp. n. This new species is only recorded from the southwestern mountains of the KSA (Fig. 28).

Monomorium subdenticorne Collingwood & Agosti (Fig. 29). A rare species endemic to the Arabian Peninsula and only recorded from Yemen and the KSA.

Table 1. Biogeography of the Arabian *Monomorium*.

Species	Type locality	Bioregion	Reference
<i>Monomorium abeillei</i>	ISRAEL	Palearctic	1; 2; 3
<i>Monomorium acutinode</i>	OMAN	Palearctic	3
<i>Monomorium aeyade</i>	OMAN	Endemic	3; Sharaf et al. 2018a
<i>Monomorium areniphilum</i>	TUNISIA	Palearctic	1; 3
<i>Monomorium asiriense</i>	KSA	Endemic	3
<i>Monomorium barbatulum</i>	KAZAKHSTAN	Palearctic	1; 3
<i>Monomorium bicolor</i>	ERITREA	Afrotropic	1; 2; 3
<i>Monomorium brunneolucidulum</i>	OMAN	Endemic	3
<i>Monomorium buettikeri</i>	KUWAIT	Endemic	3
<i>Monomorium buxtoni</i>	IRAQ	Palearctic	1; 3

<i>Monomorium carbo</i>	ETHIOPIA	Afrotropic	1; 3
<i>Monomorium carbonarium</i>	MADEIRA	Palearctic	3
<i>Monomorium clavicorne</i>	ISRAEL	Palearctic	3; Sharaf et al. 2018a
<i>Monomorium dammame</i>	KSA	Endemic	3
<i>Monomorium dirie</i>	OMAN	Endemic	3
<i>Monomorium elghazalyi</i>	YEMEN	Endemic	Sharaf et al., 2017a
<i>Monomorium exiguum</i>	ETHIOPIA	Afrotropic	3; Sharaf et al. 2018a
<i>Monomorium fayfaense</i>	KSA	Endemic	3
<i>Monomorium floricola</i>	INDIA	Tramp	Heterick, 2006
<i>Monomorium gallagheri</i>	OMAN	Endemic	3
<i>Monomorium hanaqe</i>	KSA	Endemic	3
<i>Monomorium harithe</i>	KSA	Endemic	3
<i>Monomorium hegyi</i>	KSA	Endemic	
<i>Monomorium holothir</i>	KENYA	Afrotropic	2
<i>Monomorium jizane</i>	KSA	Endemic	3
<i>Monomorium khalidi</i>	KSA	Endemic	
<i>Monomorium knappi</i>	YEMEN	Endemic	3
<i>Monomorium luteum</i>	YEMEN	Afrotropic	1; 3
<i>Monomorium mahyoubi</i>	YEMEN	Endemic	3
<i>Monomorium moathi</i>	YEMEN	Endemic	Sharaf & Collingwood, 2010
<i>Monomorium mohammedi</i>	KSA	Endemic	3; Sharaf et al. 2018a
<i>Monomorium niloticum</i>	EGYPT	Palearctic	1; 3
<i>Monomorium nimihil</i>	YEMEN	Endemic	Collingwood, 2004
<i>Monomorium nitidiventre</i>	EGYPT	Palearctic	1; 3
<i>Monomorium pharaonis</i>	EGYPT	Tramp	1; 3
<i>Monomorium rimae</i>	YEMEN	Endemic	3
<i>Monomorium riyadhe</i>	KSA	Endemic	3
<i>Monomorium salomonis</i>	EGYPT	Palearctic	1; 3
<i>Monomorium sarawatense</i>	KSA	Endemic	3; Sharaf et al. 2018a
<i>Monomorium subdenticorne</i>	YEMEN	Endemic	3

<i>Monomorium subopacum</i>	PORTUGAL	Palearctic	1; 2; 3
<i>Monomorium suleyile</i>	KSA	Endemic	3
<i>Monomorium tumaire</i>	KSA	Endemic	3
<i>Monomorium venustum</i>	SYRIA	Palearctic	1; 3

References: 1 = Collingwood 1985, 2 = Bolton, 1987, 3 = Collingwood & Agosti, 1996

DISCUSSION

Monomorium is one of the most diverse ant genera in the world, but it is rarely the most speciose genus on a regional scale. However, the Arabian *Monomorium* fauna, based on review of previous literature data and on our current work, includes 44 species, making it the most diverse known ant genus of the Arabian Peninsula (Collingwood, 1985; Collingwood & Agosti, 1996; Collingwood et al., 2011; Aldawood & Sharaf, 2011; Sharaf & Aldawood, 2013b; Sharaf et al., 2015, 2017a, 2018 a, b). Currently, the genus represents ~14% of the total number of species reported from the region (312 spp.) (e.g. Collingwood, 1985; Collingwood & Agosti, 1996; Collingwood et al., 2011; Aldawood & Sharaf, 2011; Sharaf et al., 2011, 2013c, 2015, 2017a, b, Sharaf et al., 2018a, b; Sharaf & Aldawood, 2019; Sharaf et al. 2019; 2020). This value is lower than the 20% mentioned by Collingwood & Agosti (1996). This reduction is the result of numerous taxa in previous studies now treated as junior synonyms of other species (Sharaf et al., 2015, 2017a, 2018a; this study).

We found that ~58% of the *Monomorium* species (25/44) appear to be endemic to the Arabian Peninsula. High degrees of endemism have been reported for many of Arabian arthropod groups, including ants in general (Collingwood, 1985; Collingwood & Agosti, 1996; Sharaf et al. 2014a & b; 2017a, 2018b), staphylinid beetles (Assing et al., 2013; Hlaváč et al. 2013), carabid beetles (Abdel-Dayem et al. 2018), termites (Cowie 1989), lepidopterans (Larsen 1984), and pseudoscorpions (Mahnert et al. 2014).

Biogeographically, the biota of the Arabian Peninsula does not constitute a cohesive unit (Larsen, 1984; Sharaf et al., 2020a). Instead, the Arabian Peninsula is often considered to be at the nexus of two terrestrial biogeographic realms, the Palearctic and the Afrotropic. In fact, Olson et al. (2001) places the northern and central Arabian Peninsula in the Palearctic bioregion (along with Europe, northern Africa, Asia north of the Himalayas, and neighboring islands) and the southern and eastern coasts of the Arabian Peninsula in the Afrotropic bioregion (along with

sub-Saharan Africa, southern Iran, southwestern Pakistan, and neighboring islands). Our biogeographic analysis of Arabian *Monomorium* supports this basic division, with 12 Palearctic species more common in the north and central deserts and the five Afrotropic species more common in the southern region and along the coasts.

The majority of the endemic Arabian ant species has faunal similarities with taxa from the Afrotropic bioregion that has been earlier documented by several studies (e.g. Larsen, 1984; Collingwood, 1985; Collingwood & Agosti, 1996; Sharaf & Aldawood, 2011, 2012, 2019; Sharaf et al., 2012a, b, c & 2020 a, b; El-Hawagry et al., 2013, 2017; Hájek & Reiter, 2014). Therefore, it is not surprising that a large proportion of the endemic Arabian *Monomorium* species (13/25) have been found in the mountainous ranges of southwestern KSA that extend to Yemen.

Both new *Monomorium* species reported here, *M. heggyi* and *M. khalidi*, were collected in the Shada Al-A'Ala Nature Reserve (SANR), a protected area consisting of an isolated granite mountain massif in southwestern Saudi Arabia. Its location, elevational range (470–2,222 m) and high rainfall resulting in diverse microclimates and a high biodiversity (SWA 2020). As a unique biodiversity hotspot, the SANR contains ~495 plant species (~22% of the total reported Saudi Arabian flora), including 43% of the threatened plant species and 19 endemic plants (Thomas et al., 2017). The SANR also protects a diverse fauna, including rare and endemic vertebrates, including the griffon vulture (*Gyps fulvus* (Hablitz)), the Arabian leopard (*Panthera pardus nimr* (Hemprich and Ehrenberg)), and the Arabian wolf (*Canis lupus arabs* Pocock) (SWA, 2020). The SANR invertebrate fauna has attracted relatively little attention, but recent insect biodiversity inventories and monitoring research projects conducted by King Saud University Museum of Arthropods resulted in two important faunistic studies that recorded 119 Diptera species (El-Hawagry et al., 2016) and 62 carabid beetle species (Abdel-Dayem et al., 2019). Further studies are planned to be carried out at SANR to explore additional levels of biodiversity.

In addition to the native *Monomorium* species, there are two *Monomorium* known from the Arabian Peninsula that are cosmopolitan tramp species, spread around the world through human commerce: *M. pharaonis* and *M. floricola*. The pharaoh ant, *M. pharoanis*, is a common domestic pest. Although it was first described from Egypt, its original native range is uncertain (Wetterer 2010b). We report the first known Arabian record of *M. floricola*, an Indomalayan species, from a single site in Oman. Although widespread around the world, *M. floricola* is rarely

considered a serious pest. However, because this species is very small, slow moving, cryptically colored, and primarily arboreal, its abundance and ecological importance may be underappreciated (Wetterer, 2010a).

Considering the high degree of endemism encountered, it is likely that the known Arabian *Monomorium* fauna will increase in the future with further exploration of poorly surveyed areas of the Arabian Peninsula, especially the southwestern mountains of the KSA, Yemen, and the mountainous regions of Oman and the UAE. We hope that the present study will serve as a cornerstone of future taxonomic treatments of *Monomorium* in the Arabian Peninsula.

ACKNOWLEDGMENTS

The authors thank Barry Bolton for useful discussions about *Monomorium* over years. We are indebted to the following colleagues: Boris Kondratieff for useful comments, Brian Fisher for kind permission to use *Monomorium* images on AntWeb, Michele Esposito for imaging the new species, Mahmoud Abdel-Dayem for creating the distribution maps, and Ahmed Shams Al ‘Ola for technical assistance with figures. This work was supported by the Researchers Supporting Project number (RSP-2019/107), King Saud University, Riyadh, Saudi Arabia.

REFERENCES

- Abdel-Dayem MS, Elgharbawy AA, Rasool I., Nagel P, Aldhafer HM. 2019. The Carabidae (Coleoptera) of Shada Al-A'Ala Nature Reserve, Southwestern Saudi Arabia, with description of a new species of Paussinae. *Zookeys* **812**: 93–131. doi: 10.3897/zookeys.812.30937
- Aldawood AS, Sharaf MR. 2011. *Monomorium dryhimi* sp. n., a new ant species (Hymenoptera, Formicidae) of the *M. monomorium* group from Saudi Arabia, with a key to the Arabian *Monomorium monomorium*-group. *ZooKeys* **106**: 47–54. doi:10.3897/zookeys.106.1390
- Aldawood AS, Sharaf MR, Collingwood CA. 2010. *Monomorium moathi* sp. n., a new ant species from Yemen (Hymenoptera: Formicidae) related to the *salomonis*-group. *Egyptian Academic Journal of Biological Sciences* **3** (2): 37–42.

- André E. 1881. Catalogue raisonné des Formicides provenant du voyage en Orient de M. Abeille de Perrin et description des espèces nouvelles. *Annales de la Société Entomologique de France* (6)1: 53–78.
- Assing V, Schülke M, Sharaf, MR, Aldawood AS. 2013. On the Staphylinidae of Saudi Arabia, with descriptions of two new species (Insecta: Coleoptera). *Linzer Biologische Beitrage* 45(1):141–154.
- Barech G, Khaldi M, Espadaler X, Cagniant H. 2017. Le genre *Monomorium* (Hymenoptera, Formicidae) au Maghreb (Afrique du Nord): Clé d'identification, avec la redescription de la fourmi *Monomorium major* Bernard, 1953 et nouvelles citations pour l'Algérie. *Boletín de la Sociedad Entomológica Aragonesa* 61: 151–157.
- Baroni Urbani C. 1964. Studi sulla mirmecofauna d'Italia. II. Formiche di Sicilia. *Atti della Accademia gioenia di scienze naturali in Catania* (6) 16: 25–66.
- Baroni Urbani, C. 1968. Studi sulla mirmecofauna d'Italia. V. *Aspetti Ecologici della Riviera del M. Cònero*, *Bolletino di zoologia* 35: 39-76.
- Bernard F. 1953. Les fourmis du Tassili des Ajjer. Pp. 121-250 in: Bernard, F. (ed.) 1953. Mission scientifique au Tassili des Ajjer (1949). Volume I. Recherches zoologiques et médicales. Paris: P. Lechevalier, 302 pp.
- Bernard F. 1968. Faune de l'Europe et du Bassin Méditerranéen. 3. Les fourmis (Hymenoptera Formicidae) d'Europe occidentale et septentrionale. Paris: Masson, 411 pp.
- Bolton B. 1987. A review of the *Solenopsis* genus-group and revision of Afrotropical *Monomorium* Mayr (Hymenoptera: Formicidae). *Bulletin of the British Museum (Natural History). Entomology* 54:263–452.
- Bolton B. 1994. *Identification Guide to the Ant Genera of the World*. Cambridge, Mass, 222 pp.
- Bolton B. 2020. *An online catalog of the ants of the world*. <http://antcat.org>. [accessed 13 July 2020].
- Bolton B, Fisher BL. 2014. The Madagascan endemic myrmicine ants related to *Eutetramorium* (Hymenoptera: Formicidae): taxonomy of the genera *Eutetramorium* Emery, *Malagidris* nom. n., *Myrmisaraka* gen. n., *Royidris* gen. n., and *Vitsika* gen. n. *Zootaxa* 3791:1–99.
- Borowiec L. 2014. Catalogue of ants of Europe, the Mediterranean Basin and adjacent regions (Hymenoptera: Formicidae). *Genus (Wroclaw)* 25 (1–2): 1–340
- Brown WL Jr. 1958. A review of the ants of New Zealand. *Acta Hymenopterologica* 1:1–50.

- 1089 Brown Jr WL. 2000. Diversity of ants. In: Agosti D, Majer J, Alonso E, Schultz TR, eds. *Ants.*
1090 *Standard methods for measuring and monitoring biodiversity*. Biological diversity
1091 handbook series, Washington, D.C.: Smithsonian Institution Press, 45–79, 280 pp.
- 1092 Collingwood CA. 1978. A provisional list of Iberian Formicidae with a key to the worker caste
1093 (Hym. Aculeata). *EOS. Revista Española de Entomología* **52**:65–95.
- 1094 Collingwood CA. 1985. Hymenoptera: Fam. Formicidae of Saudi Arabia. *Fauna of Saudi Arabia*
1095 **7**:230–301.
- 1096 Collingwood CA., Agosti D. 1996. Formicidae (Insecta: Hymenoptera) of Saudi Arabia (part 2).
1097 *Fauna Saudi Arabia* **15**:300–385.
- 1098 Collingwood C, Prince A. 1998. A guide to ants of continental Portugal (Hymenoptera:
1099 Formicidae). *Boletim da Sociedade Portuguesa de Entomologia* Suplemento **5**:1–49.
- 1100 Collingwood C.A, Pohl H, Guesten R, Wranik W, Van Harten A. 2004. The ants (Insecta:
1101 Hymenoptera: Formicidae) of the Socotra Archipelago. *Fauna of Arabia* **20**:473–495.
- 1102 Collingwood CA, Agosti D, Sharaf MR, Van Harten A. 2011. Order Hymenoptera, family
1103 Formicidae. *Arthropod Fauna of the UAE* **4**:405–474.
- 1104 Crawley WC. 1920. A gynandromorph of *Monomorium floricola*, Jerd. *Entomologist's Record*
1105 *and Journal of Variation* **32**:217–218.
- 1106 Deyrup M, Davis L, Cover S. 2000. Exotic ants in Florida. *Transactions of the American*
1107 *Entomological Society* **126**:293–325.
- 1108 Dlussky GM, Soyunov OS, Zabelin SI. 1990. *Ants of Turkmenistan*. [In Russian.]. Ashkabad:
1109 Ylym Press, 273 pp.
- 1110 Donisthorpe H. 1914. Myrmecophilous notes for 1913. *Entomologist's Record and Journal of*
1111 *Variation* **26**:37–45.
- 1112 Donisthorpe H. 1947. Some new ants from New Guinea. *Annals and Magazine of Natural*
1113 *History* (**11**)14:183–197.
- 1114 DuBois MB. 1986. A revision of the native New World species of the ant genus *Monomorium*
1115 (*minimum* group) (Hymenoptera: Formicidae). *University of Kansas Science Bulletin*
1116 **53**:65–119.
- 1117 El-Hawagry MS, Abdel-Dayem MS, Elgharbawy AA, Al Dhafer HM. 2016. A preliminary
1118 account of the fly fauna in Jabal Shada Al-A'la Nature Reserve, Saudi Arabia, with new

- 1119 records and biogeographical remarks (Diptera, Insecta). *ZooKeys* **636**: 107–139. <https://doi.org/10.3897/zookeys.636.9905>
- 1120
- 1121 Emery C. 1877. Catalogo delle formiche esistenti nelle collezioni del Museo Civico di Genova.
- 1122 Parte prima. Formiche provenienti dal Viaggio dei signori Antinori, Beccari e Issel nel
- 1123 Mar Rosso e nel paese dei Bogos. [part]. *Annali del Museo Civico di Storia Naturale*
- 1124 **9**:363–368.
- 1125 Emery C. 1881. Viaggio ad Assab nel Mar Rosso dei Signori G. Doria ed O. Beccari con il R.
- 1126 Avviso "Esploratore" dal 16 novembre 1879 al 26 febbraio 1880. I. Formiche. *Annali*
- 1127 *del Museo Civico di Storia Naturale* **16**:525–535.
- 1128 Emery C. 1893. Voyage de M. E. Simon à l'île de Ceylan (janvier-février 1892). Formicides.
- 1129 *Annales de la Société Entomologique de France* **62**:239–258.
- 1130 Emery C. 1894. Studi sulle formiche della fauna neotropica. VI–XVI. *Bullettino della Società*
- 1131 *Entomologica Italiana* **26**:137–241.
- 1132 Emery C. 1915. Su due formiche della Tripolitania. Bollettino del Laboratorio di Zoologia
- 1133 Generale e Agraria della Reale Scuola Superiore d'Agricoltura. *Portici* **9**:378.
- 1134 Emery C. 1922. Hymenoptera. Fam. Formicidae. Subfam. Myrmicinae. [part]. *Genera*
- 1135 *Insectorum* **174B**:95–206.
- 1136 Ettershank G. 1966. A generic revision of the world Myrmicinae related to *Solenopsis* and
- 1137 *Pheidologeton*. *Australian Journal of Zoology* **14**: 73–171.
- 1138 Evenhuis N. 2020. *The insect and spider collections of the world website*.
- 1139 <http://hbs.bishopmuseum.org/codens/> [accessed 10 Augst 2020].
- 1140 Fabricius JC. 1793. Entomologia systematica emendata et aucta. Secundum classes, ordines,
- 1141 genera, species, adjectis synonymis, locis observationibus, descriptionibus. Tome 2.
- 1142 Hafniae [= Copenhagen]: C. G. Proft, 519 pp.
- 1143 Fernández F, Serna FJ. 2019. Subfamilia Myrmicinae. Fernández, F.; Guerrero, R. J.; Delsinne,
- 1144 T. (eds.) 2019d. *Hormigas de Colombia. Bogotá*: Universidad Nacional de Colombia,
- 1145 1198 pp.
- 1146 Finzi B. 1936. Risultati scientifici della spedizione di S. A.S. il Principe Alessandro della Torre e
- 1147 Tasso nell'Egitto e penisola del Sinai. XI. Formiche. *Bulletin. Société Entomologique*
- 1148 *d'Egypte* **20**:155–210.

- 1149 Fisher BL, Bolton B. 2016. *Ants of Africa and Madagascar: A Guide to the Genera*. Berkeley:
1150 University of California Press, i–ix, 1–503 pp.
- 1151 Forel A. 1893. Formicides de l'Antille St. Vincent, récoltées par Mons. H. H.
1152 Smith. Transactions of the Entomological Society of London 1893:333-418.
- 1153 Forel A. 1907. Formicides du Musée nationale Hongroise. *Annales historico-naturales Musei*
1154 *nationalis hungarici* **5**: 1–42.
- 1155 Forel A. 1910. Glanures myrmécologiques. *Annales de la Société Entomologique de Belgique*
1156 **54**:6–32.
- 1157 Forel A. 1913. Ameisen aus Rhodesia, Kapland usw. (Hym.) gesammelt von Herrn G. Arnold,
1158 Dr. H. Brauns und Anderen. *Deutsche Entomologische Zeitschrift* 1913(Suppl.):203–
1159 225.
- 1160 Forel A. 1916. Fourmis du Congo et d'autres provenances récoltées par MM. Hermann Kohl,
1161 Luja, Mayné, etc. *Revue Suisse de Zoologie* **24**:397–460.
- 1162 Hamann HHF, Klemm W. 1967. Ergebnisse der zoologischen Nubien-Expedition 1962. Teil
1163 XXXIV. Hymenoptera - Formicidae. *Annalen des Naturhistorischen Museums in Wien*
1164 **70**:411–421.
- 1165 Heterick BE. 2001. Revision of the Australian ants of the genus *Monomorium* (Hymenoptera:
1166 Formicidae). *Invertebrate Taxonomy* **15**:353–459.
- 1167 Heterick BE. 2003. Two new Australian *Monomorium* Mayr (Hymenoptera: Formicidae),
1168 including a highly distinctive species. *Australian Journal of Entomology* **42**:249–253.
- 1169 Heterick BE. 2006. A revision of the Malagasy ants belonging to genus *Monomorium* Mayr,
1170 1855. *Proceedings of the California Academy of Sciences* **57**:69–202.
- 1171 Heterick BE. 2009. A guide to the ants of south-western Australia. *Records of the Western*
1172 *Australian Museum Supplement* **76**:1–206.
- 1173 Hita Garcia F, Wiesel E, Fischer G. 2013. The ants of Kenya (Hymenoptera: Formicidae)—
1174 faunal overview, first species checklist, bibliography, accounts for all genera, and
1175 discussion on taxonomy and zoogeography. *Journal of East African Natural History*
1176 **101**:127–222
- 1177 Hlaváč P., Sharaf, MR, Aldawood AS. 2013. New species and record of of Pselaphinae
1178 (Coleoptera: Staphylinidae) from Saudi Arabia. *Zootaxa* **3666** (3): 331–336,
1179 <http://dx.doi.org/10.11646/zootaxa.3666.3.3>

- 1180 Holt BG, Lessard JP, Borregaard MK, Fritz SA, Araújo MB, Dimitrov D, et al. 2013. An update
1181 of Wallace's zoogeographic regions of the world. *Science* **339**: 74–78. pmid:23258408
- 1182 Jerdon TC. 1851. A catalogue of the species of ants found in Southern India. *Madras Journal of*
1183 *Literature and Science* **17**:103–127.
- 1184 Karavaiev V. 1911. Ameisen aus Aegypten und dem Sudan. *Russkoe Entomologicheskoe*
1185 *Obozrenie* **11**:1–12.
- 1186 Kempf WW. 1972. Catálogo abreviado das formigas da região Neotropical. *Studia Entomologica*
1187 **15**:3–344.
- 1188 Kusnezov, N. 1949. El género Monomorium (Hymenoptera, Formicidae) en la Argentina. *Acta*
1189 *Zoologica Lilloana* 7:423-448.
- 1190 Linnaeus C. 1758. *Systema naturae per regna tria naturae, secundum classes, ordines, genera,*
1191 *species, cum characteribus, differentiis, synonymis, locis.* Tomus I. Editio decima,
1192 reformata. Holmiae [= Stockholm]: L. Salvii, 824 pp.
- 1193 Linsley EG, Usinger RL. 1966. Insects of the Galápagos Islands. *Proceedings of the California*
1194 *Academy of Sciences* (4)33:113–196.
- 1195 Mackay W, Mackay E. 2002. *The ants of New Mexico (Hymenoptera: Formicidae).* Lewiston,
1196 New York: Edwin Mellen Press, 400 pp.
- 1197 Mahnert V, Sharaf MR, Aldawood AS. 2014. Further records of Pseudoscorpions (Arachnida,
1198 Pseudoscorpions) from Saudi Arabia. *Zootaxa* **3764** (3): 387–393,
1199 <https://doi.org/10.11646/zootaxa.3764.3.8>
- 1200 Mayr G. 1862. Myrmecologische Studien. *Verhandlungen der Kaiserlich-Königlichen*
1201 *Zoologisch-Botanischen Gesellschaft in Wien* **12**:649–776.
- 1202 Mayr G. 1866. Myrmecologische Beiträge. *Sitzungsberichte der Kaiserlichen Akademie der*
1203 *Wissenschaften in Wien. Mathematisch-Naturwissenschaftliche Classe. Abteilung I*
1204 **53**:484–517.
- 1205 Mayr G. 1877. Formicidae. [In Russian.]. In: Fedchenko, A. P. 1877. Travels in Turkestan. Vol.
1206 2, Div. 5, No. 7. [In Russian.]. *Izvestiya Imperatorskago Obshchestva Lyubitelei*
1207 *Estestvoznaniya Antropologii i Etnografii pri Imperatorskom Moskovskom*
1208 *Universitete* **26**:i-iii, 1-20 (+1).
- 1209 Mayr G. 1879. Beiträge zur Ameisen-Fauna Asiens. *Verhandlungen der Kaiserlich-Königlichen*
1210 *Zoologisch-Botanischen Gesellschaft in Wien* **28**:645–686.

- 1211 Mayr G. 1901. Südafrikanische Formiciden, gesammelt von Dr. Hans Brauns. *Annalen des*
1212 *Kaiserlich-Königlichen Naturhistorischen Museums in Wien* **16**:1–30.
- 1213 Menozzi C. 1932. Missione scientifica del Prof. E. Zavattari nel Fezzan (1931). Hymenoptera-
1214 Formicidae. *Bollettino della Società Entomologica Italiana* **64**:93–95.
- 1215 Monks J, Ross S, Geiser M, De Prins J, Sharaf MR, Wyatt N, Al Rijeibi S & Polaszek A. 2019.
1216 A preliminary survey of the insect fauna of the Hajar Mountain Range, Oman. *Journal of*
1217 *Natural History* **53**: 939–963, <https://doi.org/10.1080/00222933.2019.1611969>
- 1218 Morisita M, Kubota M, Onoyama K, Ogata K, Terayama M, Yamauchi K, Sonobe R, Yamane
1219 S, Kondoh M, Imai HT. 1992. *A guide for the identification of Japanese ants. III.*
1220 *Myrmicinae and supplement to Leptanillinae. (Hymenoptera: Formicidae).* [In
1221 Japanese.]. Tokyo: Myrmecological Society of Japan, 94 pp.
- 1222 Paknia O, Radchenko A, Alipanah H, Pfeiffer M.. 2008. A preliminary checklist of the ants
1223 (Hymenoptera: Formicidae) of Iran. *Myrmecological News* **11**: 151-159.
- 1224 Roger J. 1863a. Die neu aufgeführten Gattungen und Arten meines Formiciden-Verzeichnisses
1225 nebst Ergänzung einiger früher gegebenen Beschreibungen. *Berliner Entomologische*
1226 *Zeitschrift* **7**:131–214.
- 1227 Roger J. 1863b. Verzeichniss der Formiciden-Gattungen und Arten. *Berliner Entomologische*
1228 *Zeitschrift* **7**(Beilage):1–65.
- 1229 Santschi F. 1911. Formicides nouveaux de l'Afrique Mineure (4e note suite). *Bulletin de la*
1230 *Société d'Histoire Naturelle de l'Afrique du Nord* **2**:78–85.
- 1231 Santschi F. 1912. Quelques nouvelles variétés de fourmis africaines. *Bulletin de la Société*
1232 *d'Histoire Naturelle de l'Afrique du Nord* **3**:147–149.
- 1233 Santschi F. 1914. Formicides de l'Afrique occidentale et australe du voyage de Mr. le Professeur
1234 F. Silvestri. *Bollettino del Laboratorio di Zoologia Generale e Agraria della Reale*
1235 *Scuola Superiore d'Agricoltura Portici* **8**:309–385.
- 1236 Santschi F. 1915a. Nouvelles fourmis d'Algérie, Tunisie et Syrie. *Bulletin de la Société*
1237 *d'Histoire Naturelle de l'Afrique du Nord* **6**:54–63.
- 1238 Santschi F. 1915b. Nouvelles fourmis d'Afrique. *Annales de la Société Entomologique de France*
1239 **84**:244–282.
- 1240 Santschi F. 1919. Fourmis nouvelles éthiopiennes. *Revue Zoologique Africaine* (Brussels) **6**:229–
1241 240.

- 1242 Santschi F. 1926. Description de nouveaux Formicides éthiopiens (III^{me} partie). *Revue*
1243 *Zoologique Africaine* (Brussels) **13**:207–267.
- 1244 Santschi F. 1927. Révision myrmécologique. *Bulletin et Annales de la Société Entomologique de*
1245 *Belgique* **67**:240–248.
- 1246 Santschi F. 1934. Mission J. de Lépiney au Soudan Français 1933–1934. (Huitième note.)
1247 Fourmis. *Bulletin de la Société des Sciences Naturelles du Maroc* **14**:33–34.
- 1248 Santschi F. 1936. Étude sur les fourmis du genre *Monomorium* Mayr. *Bulletin de la Société des*
1249 *Sciences Naturelles du Maroc* **16**:32–64.
- 1250 Santschi F. 1938. Quelques nouvelles fourmis d’Egypte. *Bulletin Société Entomologique*
1251 *d’Egypte* **21**:28–44.
- 1252 Sarnat EM, Economo EP. 2012. *The ants of Fiji*. University of California Publications in
1253 Entomology, Volume 132. Berkeley: University of California Press, xiii + 384 pp.
- 1254 Sharaf MR. 2006. *Taxonomic and ecological studies on family Formicidae (Order:*
1255 *Hymenoptera) in Egypt including some protectorates with a study of some insect fauna*
1256 *associated with ant species* [unpublished thesis]. Cairo: Ain Shams University, Faculty of
1257 Science, Entomology Department; 340 pp.
- 1258 Sharaf MR, Aldawood AS. 2013a. First occurrence of the *Monomorium hildebrandti*-group
1259 (Hymenoptera: Formicidae), in the Arabian Peninsula, with description of a new
1260 species *M. kondratieffi* n. sp. *Proceedings of the Entomological Society of Washington*
1261 **115** (1):75–84.
- 1262 Sharaf MR, Aldawood AS. 2013b. *Monomorium sarawatensis* Sharaf & Aldawood, sp. n. Pp.
1263 70-73 in: El-Hawagry, M. S.; Khalil, M. W.; Sharaf, M.R.; Fadl, H. H.; Aldawood,
1264 A.S. 2013. A preliminary study on the insect fauna of Al-Baha Province, Saudi Arabia,
1265 with descriptions of two new species. *ZooKeys* **274**:1–88.
- 1266 Sharaf MR, Aldawood AS. 2013c. The ant genus *Carebara* Westwood in the Arabian Peninsula
1267 (Hymenoptera, Formicidae). *ZooKeys* **357**:67–83. doi:10.3897/zookeys.357.5946
- 1268 Sharaf MR, Al Dhafer HM, Aldawood AS. 2014a. First record of the myrmicine ant genus
1269 *Meranoplus* Smith, 1853 (Hymenoptera: Formicidae) from the Arabian Peninsula with
1270 description of a new species and notes on the zoogeography of southwestern Kingdom
1271 Saudi Arabia. *PLoS ONE* **9**(11):e111298. doi:10.1371/journal.pone.0111298

- 1272 Sharaf MR, Fisher BL, Aldawood AS. 2014b. Notes on ants of the genus *Strumigenys* F. Smith,
1273 1860 (Hymenoptera: Formicidae) in the Arabian Peninsula, with a key to species.
1274 *Sociobiology* **61**(3):293–299. doi:10.13102/sociobiology.v61i3.293-299.
- 1275 Sharaf MR, Collingwood CA, Al Dhafer HM, Al mutairi MS, Aldawood AS.2015. New
1276 synonyms of two Arabian ants of the genus *Monomorium* Mayr, 1855 (Hymenoptera,
1277 Formicidae). *ZooKeys* **505**: 51–58. doi: 10.3897/ zookeys.505.9441
- 1278 Sharaf MR, Fisher BL, Collingwood CA, Aldawood, A.S. 2017a. Ant fauna (Hymenoptera:
1279 Formicidae) of the Socotra Archipelago (Yemen): zoogeography, distribution and
1280 description of a new species. *Journal of Natural History* **51** (5-6):317–378.
1281 10.1080/00222933.2016.1271157
- 1282 Sharaf MR, Akbar, SA, Al Dhafer HM, El-Gharbawy A, Aldawood AS. 2017b. Taxonomy of
1283 the Myrmicine ant genus *Temnothorax* Mayr, 1861 (Formicidae: Myrmicinae) in the
1284 Arabian Peninsula. *European Journal of Taxonomy***280**:1–17. doi:10.5852/ejt.2017.280
- 1285 Sharaf MR, Al Dhafer HM, Aldawood AS, Hita Garcia F. 2018a. Ants of the *Monomorium*
1286 *monomorium* species-group (Hymenoptera: Formicidae) in the Arabian Peninsula with
1287 description of a new species from southwestern Saudi Arabia. *PeerJ* **6**:e4277.
1288 10.7717/peerj.4277
- 1289 Sharaf MR, Fisher BL, Al Dhafer HM, Polaszek A, Aldawood AS. 2018b. Additions to the ant
1290 fauna (Hymenoptera: Formicidae) of Oman: an updated list, new records and a
1291 description of two new species. *Asian Myrmecology* **10**:e010004. 1–38.
1292 doi:10.20362/am.010004
- 1293 Sharaf MR Aldawood AS.2019. Review of the ant genus *Meranoplus* Smith, 1853
1294 (Hymenoptera: Formicidae) in the Arabian Peninsula with description of a new species
1295 *M. mosalahi* sp. n. from Oman. *PeerJ* **7**:e6287. doi:10.7717/peerj.6287
- 1296 Sharaf MR, Aldawood AS, Hita Garcia F. 2019. Review of the Arabian *Crematogaster* Lund
1297 (Hymenoptera, Formicidae), synoptic list, distribution, and description of two new
1298 species from Oman and Saudi Arabia. *ZooKeys* **898**:27–81,
1299 doi:10.3897/zookeys.898.37531
- 1300 Sharaf MR, Aldawood AS, Mohamed, AA, Hita Garcia F. 2020a. The genus *Lepisiota* Santschi,
1301 1926 of the Arabian Peninsula with the description of a new species, *Lepisiota elbazi* sp.

- 1302 nov. from Oman, an updated species identification key, and assessment of zoogeographic
- 1303 affinities. *Journal of Hymenoptera Research* **76**: 127–152, doi:10.3897/jhr.76.50193
- 1304 Sharaf MR, Mohamed AA, Al Dhafer HM, Aldawood AS. 2020b. *Nesomyrmex micheleae*, a
- 1305 new ant species (Hymenoptera: Formicidae) from the Dhofar Governorate, Oman, with a
- 1306 synoptic list, distribution, and a key to the Arabian *Nesomyrmex*. *Journal of Natural*
- 1307 *History* **54**: 351–365. <https://doi.org/10.1080/00222933.2020.1762013>
- 1308 Shuckard WE. 1838. Description of a new species of *Myrmica* which has been found in houses
- 1309 both in the Metropolis and Provinces. *Magazine of Natural History* (2)2:626–627.
- 1310 Smith F. 1857. Catalogue of the hymenopterous insects collected at Sarawak, Borneo; Mount
- 1311 Ophir, Malacca; and at Singapore, by A. R. Wallace. [part]. *Journal and Proceedings*
- 1312 *of the Linnean Society of London. Zoology* **2**:42–88.
- 1313 **Smith F. 1858.** *Catalogue of hymenopterous insects in the collection of the British Museum*. Part
- 1314 VI. Formicidae. London: British Museum, 216 pp.
- 1315 Smith F. 1876. Preliminary notice of new species of Hymenoptera, Diptera, and Forficulidae
- 1316 collected in the island of Rodriguez by the naturalists accompanying the Transit-of-
- 1317 Venus expedition. *Annals and Magazine of Natural History* (4)17:447–451.
- 1318 Sparks KS, Andersen AN, Austin AD. 2014. Systematics of the *Monomorium rothsteini* Forel
- 1319 species complex (Hymenoptera: Formicidae), a problematic ant group in
- 1320 Australia. *Zootaxa* **3893** (4):489–529.
- 1321 Sparks KS., Andersen AN, Austin AD. 2019. A multi-gene phylogeny of Australian
- 1322 *Monomorium* Mayr (Hymenoptera: Formicidae) results in reinterpretation of the genus
- 1323 and resurrection of *Chelaner* Emery. *Invertebrate Systematics* **33**: 225–236.
- 1324 Stitz H. 1917. Ameisen aus dem westlichen Mittelmeergebiet und von den Kanarischen Inseln.
- 1325 *Mitteilungen aus dem Zoologischen Museum in Berlin* **8**:333–353.
- 1326 Stitz H. 1932. Formicidae [of the Wollebaek Galapagos Expedition]. *Nyt Magazin for*
- 1327 *Naturvidenskaberne* **71**:367-372.
- 1328 SWA 2018. *Protected Areas. Saudi Wildlife Commission*. <https://www.swa.gov.sa/en> [accessed
- 1329 13 July 2020].
- 1330 SWA 2020. *Protected Areas. Saudi Wildlife Commission*. <https://www.swa.gov.sa/en> [26-8-
- 1331 2020].

- 1332 Terayama M. 2009. A synopsis of the family Formicidae of Taiwan (Insecta:
1333 Hymenoptera). *Research Bulletin of Kanto Gakuen University*. Liberal Arts **17**:81–266.
- 1334 Thomas J, El-Sheikh MA, Alatar AA. 2017. Endemics and endangered species in the
1335 biodiversity hotspot of the Shada Mountains, Saudi Arabia. *Journal of Arid Land* **9**(1):
1336 109–21. doi: 10.1007/s40333-016-0025-8
- 1337 Vigna Taglianti A, Audisio PA, Biondim, Bologna MA, Carpaneto GM, De Biase A, Fattorini S,
1338 Piattella E, Sindaco R, Venchi A, Zapparolim, 1999. A proposal for a chorotype
1339 classification of the Near East fauna, in the framework of the Western Palaearctic region.
1340 *Biogeographia* **20**: 31–59.
- 1341 Vonshak M, Ionescu-Hirsch A. 2009. A checklist of the ants of Israel (Hymenoptera: Formici-
1342 dae). *Israel Journal of Entomology* **39**: 33–55.
- 1343 Walker F. 1871. *A list of Hymenoptera collected by J. K. Lord, Esq. in Egypt, in the*
1344 *neighbourhood of the Red Sea, and in Arabia, with descriptions of the new species.*
1345 London: E. W. Janson, vi + 59 pp.
- 1346 Ward PS, Brady SG, Fisher BL, Schultz TR. 2015. The evolution of myrmicine ants: phylogeny
1347 and biogeography of a hyperdiverse ant clade (Hymenoptera: Formicidae). *Systematic*
1348 *Entomology* **40**:61–81.
- 1349 Wetterer JK. 2010a. Worldwide spread of the flower ant, *Monomorium floricola* (Hymenoptera:
1350 Formicidae). *Myrmecological News* **13**:19–27.
- 1351 Wetterer JK. 2010b. Worldwide spread of the pharaoh ant, *Monomorium pharaonis*
1352 (Hymenoptera: Formicidae). *Myrmecological News* **13**:115–129
- 1353 Wheeler WM. 1905. The ants of the Bahamas, with a list of the known West Indian
1354 species. *Bulletin of the American Museum of Natural History* **21**:79–135.
- 1355 Wheeler WM. 1913. The ants of Cuba. *Bulletin of the Museum of Comparative Zoology* **54**:477–
1356 505.
- 1357 Wheeler WM. 1922. Ants of the American Museum Congo expedition. New York: *Bulletin of*
1358 *the American Museum of Natural History* **11**:39.
- 1359 Wheeler GC, Wheeler J. 1955. The ant larvae of the myrmicine tribe Solenopsidini. *American*
1360 *Midland Naturalist* **54**:119–141.
- 1361 Wheeler WM, Mann WM. 1916. The ants of the Phillips Expedition to Palestine during 1914.
1362 *Bulletin of the Museum of Comparative Zoology* **60**:167–174.

- 1363 Wilson EO, Taylor RW. 1967b. The ants of Polynesia (Hymenoptera: Formicidae). *Pacific*
1364 *Insects Monograph* **14**: 1–109.
- 1365 Wu J, Wang C. 1995. *The ants of China*. [In Chinese.]. Beijing: China Forestry Publishing
1366 House, x + 214 pp.
- 1367

Figure Legends.

Figure 1. A: head of *M. exiguum* in full-face view, CASENT0217367 (Erin Prado); B: head of *M. khalidi*, **sp. n.** in full-face view, CASENT0922288 (Michele Esposito); C: head of *M. carbonarium* in full-face view, CASENT0902279 (Ryan Perry); D: head of *M. floricola* in full-face view, CASENT0922876 (Michele Esposito); E: body of *M. carbonarium* in profile, CASENT0902279 (Ryan Perry); F: body of *M. sarawatense* in profile, CASENT0280971 (Estella Ortega), www.AntWeb.org.

Figure 2. A: body of *M. floricola* in profile, CASENT0922876 (Michele Esposito); B: body of *M. holothir* in profile, CASENT0902243 (Will Ericson); C: body of *M. aeyade* in profile, CASENT0922329 (Michele Esposito); D: body of *M. exiguum* in profile, CASENT0217367 (Erin Prado); E: body of *M. mohammedi* in profile, CASENT0922351 (Michele Esposito); www.AntWeb.org, F: body of *M. clavicorne* in profile, (Francisco Hita Garcia).

Figure 3. A: head of *M. barbatulum* in profile, CASENT0922263 (Michele Esposito); B: mesosoma of *M. rabirium* in profile, CASENT0746641 (Zach Lieberman); C: mesosoma of *M. elghazalyi* in profile, CASENT0746626 (Michele Esposito); D: petiole and postpetiole of *M. harithe* in profile, CASENT0913802 (Will Ericson); E: head of *M. elghazalyi* in full-face view, CASENT0746626 (Michele Esposito); F head of *M. dirie* in full-face view, CASENT0913571 (Alexandra Westrich), www.AntWeb.org.

Figure 4. A: head of *M. dirie* in profile, CASENT0913571 (Alexandra Westrich); B: head of *M. salomonis* in profile, CASENT0913835 (Will Ericson); C: mesosoma of *M. subdenticorne* in profile, CASENT0914318 (Zach Lieberman); D: mesosoma of *M. bicolor* in profile, CASENT0073615 (Michele Esposito); E: petiole of *M. acutinode* in profile, CASENT0913547 (Will Ericson); F petiole of *M. carbo* in profile, CASENT0249908 (Shannon Hartman), www.AntWeb.org.

Figure 5. A: mesosoma of *M. areniphilum* in profile, CASENT0048600 (Michele Esposito); B: mesosoma of *M. subopacum* in profile, CASENT0064820 (April Nobile); C: mesosoma of *M. luteum* in profile, CASENT0904599 (Will Ericson); D: mesosoma of *M. niloticum* in profile,

CASENT0260164 (Estella Ortega); E: head of *M. niloticum* in full-face view, CASENT0919811 (Michele Esposito); F head of *M. riyadhe* in full-face view, CASENT0922342 (Michele Esposito), www.AntWeb.org.

Figure 6. A: head of *M. hanaqe* in full-face view, CASENT0249834 (Ryan Perry); B: mesosoma of *M. hanaqe* in profile, CASENT0249834 (Ryan Perry); C: head of *M. jizane* in full-face view, CASENT0913806 (Will Ericson); D: propodeum of *M. jizane* in profile, CASENT0913806 (Will Ericson); E: head of *M. fayfaense* in full-face view, CASENT0249833 (Ryan Perry); F gaster of *M. fayfaense* in profile, CASENT0249833 (Ryan Perry), www.AntWeb.org.

Figure 7. A: head of *M. knappi* in full-face view, CASENT0913812 (Will Ericson); B: body of *M. knappi* in profile, CASENT0913812 (Will Ericson); C: body of *M. nitidiventre* in profile, CASENT0904602 (Will Ericson); D: head of *M. nitidiventre* in full-face view, CASENT0904602 (Will Ericson); E: body of *M. khalidi* **sp. n.** in profile, CASENT0922288 (Michele Esposito); F head of *M. khalidi* **sp. n.** in profile, CASENT0922288 (Michele Esposito), www.AntWeb.org.

Figure 8. A: mesosoma of *M. pharaonis* in profile, CASENT0246072 (Andrea Walker); B: mesosoma of *M. buxtoni* in profile, CASENT0902220 (Will Ericson); C: head of *M. suleyile* in full-face view, CASENT0913854 (Zach Lieberman); D: head of *M. mahyoubi* in full-face view, CASENT0913823 (Alexandra Westrich); E: head of *M. tumaire* in profile, CASENT0249858 (Ryan Perry); F head of *M. buettikeri* in profile, CASENT0913565 (Zach Lieberman), www.AntWeb.org.

Figure 9. A: head of *M. gallagheri* in full-face view, CASENT0913582 (Zach Lieberman); B: head of *M. buxtoni* in full-face view, CASENT0902220 (Zach Lieberman); C: body of *M. abeillei* in profile, CASENT0915411 (Will Ericson); D: body of *M. asiriense* in profile, CASENT0913560 (Zach Lieberman), www.AntWeb.org.

Figure 10. *M. abeillei*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0915411 (Will Ericson), www.AntWeb.org.

1430

1431 Figure 11. *M. areniphilum*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1432 CASENT0048600 (Michele Esposito), www.AntWeb.org.

1433

1434 Figure 12. *M. barbatulum*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1435 CASENT0922263 (Michele Esposito), www.AntWeb.org.

1436

1437 Figure 13. *M. bicolor*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1438 CASENT0904601 (Will Ericson), www.AntWeb.org.

1439

1440 Figure 14. *M. floricola*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1441 CASENT0922876 (Michele Esposito), www.AntWeb.org.

1442

1443 Figure 15. *M. harithe*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1444 CASENT0913802 (Will Ericson), www.AntWeb.org.

1445

1446 Figure 16. *M. heggyi* sp. n., A: body in profile; B: body in dorsal view); C: head in full-face
1447 view, CASENT0746641 (Zach Lieberman), www.AntWeb.org.

1448

1449 Figure 17. Shada Al A'la, the type locality of *M. heggyi* **sp. n.** (A. Shams Al Ola).

1450

1451 Figure 18. *M. khalidi* **sp. n.**, A: body in profile; B: body in dorsal view); C: head in full-face
1452 view, CASENT0922288 (Michele Esposito), www.AntWeb.org.

1453

1454 Figure 19. Shada Al A'la, the type locality of *M. khalidi* **sp. n.** (A. Shams Al Ola).

1455

1456 Figure 20. *M. niloticum*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1457 CASENT0905755 (Will Ericson), www.AntWeb.org.

1458

1459 Figure 21. *M. nitidiventre*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1460 CASENT0904602 (Will Ericson), www.AntWeb.org.

1461

1462 Figure 22. *M. subdenticorne*, A: body in profile; B: body in dorsal view); C: head in full-face
1463 view, CASENT0914318 (Zach Lieberman), www.AntWeb.org.

1464 Figure 23. A: *M. venustum*, A: body in profile; B: body in dorsal view); C: head in full-face
1465 view, CASENT0902221 (Will Ericson), www.AntWeb.org.

1466

1467 Figure 24. Distribution map of *M. abeillei*.

1468

1469 Figure 25. Distribution map of *M. niloticum*.

1470

1471 Figure 26. Distribution map of *M. venustum*.

1472

1473 Figure 27. Distribution map of *M. barbatulum*, *M. bicolor*, *M. floricola*.

1474

1475 Figure 28. Distribution map of *M. harithe*, *M. khalidi* **sp. n.**, *M. nitidiventre*, *M. hegyi* **sp. n.**

1476

1477 Figure 29. Distribution map of *M. subdenticorne*.

1478

1479

Figure 1

Figure 1

Figure 1. A: head of *M. exiguum* in full-face view, CASENT0217367 (Erin Prado); B: head of *M. khalidi*, **sp. n.** in full-face view, CASENT0922288 (Michele Esposito); C: head of *M. carbonarium* in full-face view, CASENT0902279 (Ryan Perry); D: head of *M. floricola* in full-face view, CASENT0922876 (Michele Esposito); E: body of *M. carbonarium* in profile, CASENT0902279 (Ryan Perry); F: body of *M. sarawatense* in profile, CASENT0280971 (Estella Ortega), www.AntWeb.org.




Figure 2

A: body of *M. floricola* in profile, CASENT0922876 (Michele Esposito); B: body of *M. holothir* in profile, CASENT0902243 (Will Ericson); C: body of *M. aeyade* in profile, CASENT0922329 (Michele Esposito); D: body of *M. exiguum* i

Key illustrations




Figure 3

A: head of *M. barbatulum* in profile, CASENT0922263 (Michele Esposito); B: mesosoma of *M. rabirium* in profile, CASENT0746641 (Zach Lieberman); C: mesosoma of *M. elghazalyi* in profile, CASENT0746626 (Michele Esposito); D: petiole and postp

Key illustrations




Figure 4

A: head of *M. dirie* in profile, CASENT0913571 (Alexandra Westrich); B: head of *M. salomonis* in profile, CASENT0913835 (Will Ericson); C: mesosoma of *M. subdenticorne* in profile, CASENT0914318 (Zach Lieberman); D: mesosoma of [i]M. bicolo

Key illustrations




Figure 5

A: mesosoma of *M. areniphilum* in profile, CASENT0048600 (Michele Esposito); B: mesosoma of *M. subopacum* in profile, CASENT0064820 (April Nobile); C: mesosoma of *M. luteum* in profile, CASENT0904599 (Will Ericson); D: mesosoma of [i]M. nil

Key illustrations







Figure 6

A: head of *M. hanage* in full-face view, CASENT0249834 (Ryan Perry); B: mesosoma of *M. hanage* in profile, CASENT0249834 (Ryan Perry); C: head of *M. jizane* in full-face view, CASENT0913806 (Will Ericson); D: propodeum of *M. jizane* in

Key illustrations




Figure 7

A: head of *M. knappi* in full-face view, CASENT0913812 (Will Ericson); B: body of *M. knappi* in profile, CASENT0913812 (Will Ericson); C: body of *M. nitidiventre* in profile, CASENT0904602 (Will Ericson); D: head of *M. nitidiventre* in

Key illustrations




Figure 8

A: mesosoma of *M. pharaonis* in profile, CASENT0246072 (Andrea Walker); B: mesosoma of *M. buxtoni* in profile, CASENT0902220 (Will Ericson); C: head of *M. suleyile* in full-face view, CASENT0913854 (Zach Lieberman); D: head of [i]M. mahyoub

Key illustrations




Figure 9

A: head of *M. gallagheri* in full-face view, CASENT0913582 (Zach Lieberman); B: head of *M. buxtoni* in full-face view, CASENT0902220 (Zach Lieberman); C: body of *M. abeillei* in profile, CASENT0915411 (Will Ericson); D: body of [i]M. asirie

Key illustrations




Figure 10

M. abeillei, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0915411 (Will Ericson), www.AntWeb.org . Figure 11. *M. areniphilum*, A: body in profile; B: body in dorsal view); C: head in full-face view,

Monomorium abeillei




Figure 11

Figure 11. *M. areniphilum*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0048600 (Michele Esposito), www.AntWeb.org .

Figure 11. *M. areniphilum*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0048600 (Michele Esposito), www.AntWeb.org .




Figure 12

Figure 12. *M. barbatulum*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0922263 (Michele Esposito), www.AntWeb.org .

Figure 12. *M. barbatulum*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0922263 (Michele Esposito), www.AntWeb.org .




Figure 13

Figure 13. *M. bicolor*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0904601 (Will Ericson), www.AntWeb.org .

Figure 13. *M. bicolor*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0904601 (Will Ericson), www.AntWeb.org .




Figure 14

Figure 14. *M. floricola*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0922876 (Michele Esposito), www.AntWeb.org .

Figure 14. *M. floricola*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0922876 (Michele Esposito), www.AntWeb.org .




Figure 15

Figure 15. *M. harithe*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0913802 (Will Ericson), www.AntWeb.org .

Figure 15. *M. harithe*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0913802 (Will Ericson), www.AntWeb.org .




Figure 16

Figure 16. *M. heggyi* sp. n., A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0746641 (Zach Lieberman), www.AntWeb.org .

Figure 16. *M. heggyi* sp. n., A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0746641 (Zach Lieberman), www.AntWeb.org .




Figure 17

Figure 17. Shada Al A'la, the type locality of *M. heggyi* sp. n. (A. Shams Al Ola).

Figure 17. Shada Al A'la, the type locality of *M. heggyi* **sp. n.** (A. Shams Al Ola).



Figure 18

Figure 18. *M. khalidi* sp. n., A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0922288 (Michele Esposito), www.AntWeb.org .

Figure 18. *M. khalidi* **sp. n.**, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0922288 (Michele Esposito), www.AntWeb.org .




Figure 19

Figure 19. Shada Al A'la, the type locality of *M. khalidi* sp. n. (A. Shams Al Ola).

Figure 19. Shada Al A'la, the type locality of *M. khalidi* **sp. n.** (A. Shams Al Ola).



Figure 20

Figure 20. *M. niloticum*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0905755 (Will Ericson), www.AntWeb.org .

Figure 20. *M. niloticum*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0905755 (Will Ericson), www.AntWeb.org .




Figure 21

Figure 21. *M. nitidiventre*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0904602 (Will Ericson), www.AntWeb.org .

Figure 21. *M. nitidiventre*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0904602 (Will Ericson), www.AntWeb.org .



Figure 22

Figure 22. *M. subdenticorne*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0914318 (Zach Lieberman), www.AntWeb.org .

Figure 22. *M. subdenticorne*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0914318 (Zach Lieberman), www.AntWeb.org .




Figure 23

Figure 23. A: *M. venustum*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0902221 (Will Ericson), www.AntWeb.org .

Figure 23. A: *M. venustum*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0902221 (Will Ericson), www.AntWeb.org .




Figure 24

Figure 24. Distribution map of *M. abeillei*.

Figure 24. Distribution map of *M. abeillei*.




Figure 25

Figure 25. Distribution map of *M. niloticum*.

Figure 25. Distribution map of *M. niloticum*.




Figure 26

Figure 26. Distribution map of *M. venustum*.

Figure 26. Distribution map of *M. venustum*.




Figure 27

Figure 27. Distribution map of *M. barbatulum*, *M. bicolor*, *M. floricola*.

Figure 27. Distribution map of *M. barbatulum*, *M. bicolor*, *M. floricola*.



Figure 28

Figure 28. Distribution map of *M. harithe*, *M. khalidi* sp. n., *M. nitidiventre*, *M. heggyi* sp. n.

Figure 28. Distribution map of *M. harithe*, *M. khalidi* **sp. n.**, *M. nitidiventre*, *M. heggyi* **sp. n.**




Figure 29

Figure 29. Distribution map of *M. subdenticorne*.

Figure 29. Distribution map of *M. subdenticorne*.

