

***Monomorium* (Hymenoptera: Formicidae) of the Arabian Peninsula with description of two new species, *M. heggyi* sp. n. and *M. khalidi* sp. n. (#54174)**

1

First revision

Guidance from your Editor

Please submit by **3 Dec 2020** for the benefit of the authors .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Author notes

Have you read the author notes on the [guidance page](#)?



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

1 Tracked changes manuscript(s)

1 Rebuttal letter(s)

29 Figure file(s)

1 Table file(s)

! Custom checks

New species checks



Have you checked our [new species policies](#)?



Do you agree that it is a new species?



Is it correctly described e.g. meets ICZN standard?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

***Monomorium* (Hymenoptera: Formicidae) of the Arabian Peninsula with description of two new species, *M. heggyi* sp. n. and *M. khalidi* sp. n.**

Mostafa R Sharaf^{Corresp., 1}, Amr A Mohamed², Brendon E Boudinot³, James K Wetterer⁴, Francisco Hita Garcia⁵, Hathal M Al Dhafer⁶, Abdulrahman S Aldawood⁶

¹ Department of Plant Protection, College of Food and Agriculture Sciences, King Saud University, Riyadh, Saudi Arabia

² Department of Entomology, Faculty of Science, Cairo University, Giza, Egypt

³ Department of Entomology/Nematology, University of California, Davis, Davis, California, United States

⁴ Wilkes Honors College, Florida Atlantic University, Florida, Florida, United States

⁵ Biodiversity and Biocomplexity Unit, Okinawa Institute of Science and Technology Graduate University, Onna-son, Okinawa, Japan

⁶ Department of Plant Protection, College of Food and Agriculture Sciences, King Saud University, Riyadh, Riyadh, Saudi Arabia

Corresponding Author: Mostafa R Sharaf

Email address: antsharaf@gmail.com

We present a revised and updated synoptic list of 44 Arabian *Monomorium* species, including two new species of the *M. salomonis* species-group: *M. heggyi* **sp. n.**, and *M. khalidi* **sp. n.** We propose the following new synonyms: *M. abeillei* André (= *M. wahibiense* Collingwood & Agosti **syn. n.**); *M. areniphilum* Santschi (= *M. fezzanense* Collingwood & Agosti **syn. n.**, = *M. hemame* Collingwood & Agosti **syn. n.** = *M. marmule* Collingwood & Agosti **syn. n.**); *M. bicolor* Emery (= *M. phoenicum* Santschi **syn. n.**); *M. harithe* Collingwood & Agosti (= *M. najrane* Collingwood & Agosti **syn. n.**); *M. niloticum* Emery (= *M. matame* Collingwood & Agosti **syn. n.**); and *M. nitidiventre* Emery (= *M. yemene* Collingwood & Agosti **syn. n.**). An illustrated key and distribution maps are presented for the treated species. Ecological and biological notes are given when available. The majority of Arabian *Monomorium* species (24) are endemic to the peninsula. All except one of the remaining species are more broadly ranging Afrotropical and Palearctic species, supporting the view of Arabia as a biogeographical crossroads between these two regions. *Monomorium floricola* (Jerdon), the sole species of Indomalayan origin, is recorded for the first time from the Arabian Peninsula.

***Monomorium* (Hymenoptera: Formicidae) of the Arabian Peninsula with description of two new species, *M. heggvi* sp. n. and *M. khalidi* sp. n.**

Running head: Revision of Arabian *Monomorium*.

Mostafa R. Sharaf^{a,Ω}, Amr A. Mohamed^b, Brendon E. Boudinot^c, James K. Wetterer^d, Francisco Hita Garcia^e, Hathal M. Al Dhafer^a & Abdulrahman S. Aldawood^a

^a Department of Plant Protection, College of Food and Agriculture Sciences, King Saud University, Riyadh, Kingdom of Saudi Arabia

^b Department of Entomology, Faculty of Science, Cairo University, Giza, Egypt

^c Department of Entomology/Nematology, University of California, Davis, USA.

^d Wilkes Honors College, Florida Atlantic University, 5353 Parkside Drive, Jupiter, FL 33458, USA.

^e Biodiversity and Biocomplexity Unit, Okinawa Institute of Science and Technology Graduate University, Onna-son, Okinawa, Japan.

^Ω corresponding author, e-mail: antsharaf@gmail.com; mosharaf@ksu.edu.sa

ABSTRACT

We present a revised and updated synoptic list of 44 Arabian *Monomorium* species, including two new species of the *M. salomonis* species-group: *M. heggvi* sp. n., and *M. khalidi* sp. n. We propose the following new synonyms: *M. abeillei* André (= *M. wahibiense* Collingwood & Agosti syn. n.); *M. areniphilum* Santschi (= *M. fezzanense* Collingwood & Agosti syn. n., = *M. hemame* Collingwood & Agosti syn. n. = *M. marmule* Collingwood & Agosti syn. n.); *M. bicolor* Emery (= *M. phoenicum* Santschi syn. n.); *M. harithe* Collingwood & Agosti (= *M. najrane* Collingwood & Agosti syn. n.); *M. niloticum* Emery (= *M. matame* Collingwood & Agosti syn. n.); and *M. nitidiventre* Emery (= *M. yemene* Collingwood & Agosti syn. n.). An illustrated key and distribution maps are presented for the treated species. Ecological and biological notes are given when available. The majority of Arabian *Monomorium* species (24) are endemic to the peninsula. All except one of the remaining species are more broadly ranging Afrotropical and Palearctic species, supporting the view of Arabia as a biogeographical crossroads between these two regions. *Monomorium floricola* (Jerdon), the sole species of Indomalayan origin, is recorded for the first time from the Arabian Peninsula.

Keywords. biogeography, faunal list, Middle East, taxonomic revision.

INTRODUCTION

Monomorium Mayr is one of the largest genera of ants, with 326 valid species and subspecies, as well as three fossil species (Bolton, 2020). *Monomorium* species predominantly inhabit tropical, subtropical, and warm temperate regions of the world (Bolton, 1987; Bolton, 1994; Brown, 2000). The majority of *Monomorium* species are native to the Afrotropical bioregion (Bolton, 1987; Hita Garcia et al., 2013; Bolton, 2020). In contrast, the Palaearctic and New World faunas are relatively species-poor (Kempf, 1972; Sharaf et al., 2018a; Fernández & Serna, 2019). In total, 56 species have been reported from the Arabian Peninsula (Collingwood, 1985; Collingwood & Agosti, 1996; Aldawood et al., 2010; Aldawood & Sharaf, 2011; Sharaf et al., 2013a, b, c, Sharaf et al., 2014a, b, Sharaf et al., 2015, 2017a, b, 2018a, b). Most are generalized feeders or granivores, but some are leptoecic or parasitic (Ettershank, 1966; Bolton, 1987; Brown, 2000). Notably, *Monomorium* is one of the few genera that includes several very successful cosmopolitan tramp species, such as *M. pharaonis* (L.) and *M. floricola* (Jerdon), *M. latinode* Mayr, and *M. subopacum* (Smith) (Bolton, 1987; Wetterer, 2010a, b).

The genus *Monomorium* has a long and complex taxonomic history (see Heterick 2006 for a full history of the genus). Emery (1922) offered subgeneric names for several *Monomorium* species, based on characters of the antennal club and number of antennomeres, some names were later raised to the generic level. Ettershank (1966) made a bold attempt to organize the global fauna of *Monomorium* and related genera. Bolton (1987) extensively reviewed, refined and keyed the Afrotropical species recognizing 145 species in 8 species-groups including descriptions of 46 new species and 43 new synonyms. The Malagasy *Monomorium* fauna was revised by Heterick (2006) following Bolton's Afrotropical species-groups as a basis for organizing numerous new species. Heterick (2001, 2003, 2009) comprehensively treated the Australian *Monomorium* fauna, whereas DuBois (1986) and Mackay & Mackay (2002) studied the Nearctic fauna. The comprehensive phylogenetic work of the subfamily Myrmicinae (Ward et al., 2015) has adopted substantial taxonomic changes including the resurrection of the genera *Sylophopsis* and *Trichomyrmex* from synonymy under *Monomorium*. Sparks et al. (2014) fully revised the *Monomorium rothsteini* species-complex. Numerous smaller and regional contributions are scattered in the literature including Argentina (Kusnezov, 1949), China (Wu &

Wang, 1995), Colombia (Fernández & Serna, 2019), Fiji (Sarnat & Economo, 2012), Italy (Baroni Urbani, 1964, 1968), Japan (Morisita et al., 1992), Morocco (Barech et al., 2017), New Zealand (Brown, 1958), Polynesia (Wilson & Taylor, 1967), Taiwan (Terayama, 2009), the Iberian Peninsula (Collingwood, 1978), Turkmenistan (Dlussky et al., 1990), and Western Europe (Bernard, 1968).

However, our understanding of the phylogenetic relationships among the currently accepted *Monomorium* species-groups remains far from satisfactory and the overall taxonomic foundation is in a state of major revision at present (Sparks et al., 2019). Moreover, our knowledge about the diversity of the genus is still fragmentary for several regions, such as the Mediterranean, the North African-Indian Desert (including what referred as the Saharo-Arabian in some literatures [Holt et al., 2013]) (Vigna Taglianti et al., 1999; Sharaf et al., 2017a), as well as the Indomalayan and the Neotropical realms, which contain large numbers of described infraspecific taxa in need of further reevaluation (Borowiec, 2014). We note that even the Afrotropical *Monomorium* fauna, which is the most diverse and best studied so far, is in dire need of an updated taxonomic revision with likely more than 100 undescribed species and many valid species needing reevaluation (Bolton, 1987; Hita Garcia, pers. obs.).

The oldest records of *Monomorium* species from the Arabian Peninsula were of *M. carbonarium* (Smith) and *M. niloticum* Emery (Forel 1907). Collingwood (1985) reported 17 valid *Monomorium* species from the Kingdom of Saudi Arabia (KSA) and Collingwood & Agosti (1996) listed 49 valid *Monomorium* species from the entire Arabian Peninsula, including 32 new species. Additional papers have included records from the KSA (Aldawood & Sharaf, 2011; Sharaf & Aldawood, 2013b; Sharaf et al., 2015, 2018 a, b), Oman (Sharaf et al., 2018b; Monks et al., 2019), the Socotra Archipelago (Collingwood et al., 2004; Sharaf et al., 2017a), the United Arab Emirates (UAE) (Collingwood et al., 2011), and Yemen (Aldawood et al., 2010).

Collingwood (1985) and Collingwood & Agosti (1996) recognized a total of 53 *Monomorium* species on the Arabian Peninsula, of which 17 were described from countries in the region. However, the taxonomic status of several species has remained uncertain due to brief descriptions with insufficient differential diagnoses, apparent ambiguities in the taxonomic keys, and lack of species-group assignment (Sharaf et al., 2020a, b). The present study aims to clarify the current status of the Arabian *Monomorium* by providing the following:

I. Diagnoses of Arabian *Monomorium* and *Monomorium* species-groups

- II. A synoptic checklist of Arabian *Monomorium*
- III. An illustrated identification key to species based on the worker caste.
- IV. New taxonomic treatments proposing new synonymies and describing two new species.
- V. Biogeographical analyses, including distribution maps

MATERIAL AND METHODS

The species names follow the online catalogue of ants of the world (Bolton 2020). We made digital color images of each species using a Leica DFC450 digital camera with a Leica Z16 APO microscope and LAS (v3.8) software. The images are available online on AntWeb (<http://www.AntWeb.org>) and are accessible through unique specimen identifiers (e.g. CASENT0922288). Distribution maps were made using DIVA-GIS (version 7.5.0.0).

Measurements and indices.

All measurements are in millimeters and follow the standard measurements of Bolton (1987) and Sharaf et al. (2018a).

CI= Cephalic Index ($HW/HL \times 100$).

EI= Eye Index ($EL/HW \times 100$).

EL= Eye Length; maximum diameter of eye in lateral view.

EM= Eye-Mandible Distance; distance between anterior margin of eye and mandibular insertion in lateral view.

HL= Head Length; maximum length of head, excluding mandibles in full-face view.

HW= Head Width; maximum width of head directly behind eyes in full-face view.

ML= Mesosoma Length (=Weber Length); length of mesosoma in lateral view); from a point at which pronotum meets cervical shield to posterior base of propodeal lobes or teeth.

PpH= Postpetiole Height; maximum height measured in lateral view.

PpL= Postpetiole Length; maximum length of postpetiolar node measured in dorsal view, from anterior margin to posterior margin.

PpW= Postpetiole Width; maximum width measured in dorsal view.

PtH= Petiole Height; maximum height measured in lateral view.

PtL= Petiole Length; maximum length of petiolar node measured in dorsal view, from anterior margin to posterior margin.

125 PtW= Petiole Width; maximum width measured in dorsal view.
 126 PW= Pronotal Width; maximum width in dorsal view.
 127 SI= Scape Index ($SL/HW \times 100$).
 128 SL= Scape Length, excluding basal neck.
 129 TL= Total Length, sum of lengths of head, mesosoma, petiole, postpetiole and gaster in
 130 profile.

131

132 **Abbreviations of museums.**

133 Abbreviations of natural history collections follow Brandão (2000) except for WML that follows
 134 Evenhuis (2020). The material on which this study is based is located and/or was examined at the
 135 following institutions:

136 BMNH = The Natural History Museum, London, United Kingdom.

137 CASC = California Academy of Sciences, San Francisco, USA.

138 KSMA = King Saud University Museum of Arthropods, Riyadh, Kingdom of Saudi Arabia.

139 LACM = Los Angeles County Museum of Natural History, Los Angeles, CA, USA.

140 MNHN = Muséum National d'Histoire Naturelle, Paris, France.

141 NHMB = Naturhistorisches Museum, Basel, Switzerland.

142 OXUM = Hope Entomological Collections, University Museum, Parks Road, OXI 3PW Oxford,
 143 U.K.

144 WML = World Museum Liverpool, Liverpool, U.K.

145 **Nomenclatural acts**

146 The electronic version of this article in Portable Document Format (PDF) will represent a
 147 published work according to the International Commission on Zoological Nomenclature (ICZN),
 148 and hence the new names contained in the electronic version are effectively published under that
 149 Code from the electronic edition alone. This published work and the nomenclatural acts it
 150 contains have been registered in ZooBank, the online registration system for the ICZN. The
 151 ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed
 152 through any standard web browser by appending the LSID to the prefix <http://zoobank.org/>. The
 153 LSID for this publication is: urn:lsid:zoobank.org:pub:A7FFDF5C-6CD5-41CA-B106-
 154 BFF6BDFCB258; for *Monomorium heggyi* **sp. n.** is urn:lsid:zoobank.org:act:B57EC2EA-1781-
 155 4C19-ADFE-757C834E2774; and for *Monomorium khalidi* **sp. n.** is

urn:lsid:zoobank.org:act:3B5BB529-D842-4146-B8F7-ACCAC9CD5BA7. The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central and CLOCKSS.

RESULTS

I. Diagnosis of Arabian *Monomorium*

Members of the genus *Monomorium* can be recognized by combination of the following characters in the worker caste (Bolton, 1987, 1994; DuBois, 1986; Fisher & Bolton, 2016; Sparks *et al.*, 2019): monomorphic to polymorphic; antennae 10–12 segmented (most frequently 12), conspicuous 3-segmented club; mandibles with 3–4 teeth; palp formula 2,2, or 1,2; median clypeal seta conspicuous; median portion of clypeus raised, the raised portion usually longitudinally bicarinate; lateral portions of the clypeus not elevated as shield-like ridges anterior to the antennal toruli; frontal carinae distinct, but absent posterior to the medial arches of the antennal toruli; metanotal groove present, commonly impressed; propodeal dorsum usually unarmed and rounding into the declivity; propodeal spiracle usually circular, located at about the midlength of the sclerite; abdominal segment IV (metasomal III, gastral I) with posttergite overlapping postternite.

Diagnosis of Arabian *Monomorium* species-groups

The *M. monorium* species-group can be readily recognized by the following combination of characters in the worker caste (Sharaf *et al.* 2018a): monomorphic, with size variation; head longer than broad; mandibles smooth and masticatory margin armed with four teeth; antennae with 10–12 segments, terminating in a three-segmented club; median clypeal portion raised anteriorly and longitudinally bicarinate; eyes present with variable size, located in front of the midlength of the sides in full-face view, and with four or more ommatidia in the longest row; head smooth and shining; metanotal groove well-defined, with distinct cross-ribs; propodeal spiracle circular to subcircular; propodeal dorsum meeting declivity in a rounded angle; promesonotum and propodeal dorsum smooth; body pilosity variable but usually distinct; petiole, postpetiole and gastral tergites usually smooth.

The *M. salomonis* species-group can be diagnosed by the following character states in the worker caste (Bolton, 1987): monomorphic, with minor size variation; palp formula 2,2 or 1,2 in

some minute species; mandibles usually sculptured; masticatory margins of mandibles armed with four teeth which decrease in size from apex to base; median clypeal portion raised, projecting anteriorly; cephalic dorsum usually sculptured, ranging from dense blanketing reticulate-punctuation to feeble superficial reticular patterning; eyes prominent of medium to large size, usually with six or more ommatidia in the longest row; eyes circular to oval in shape; head longer than broad; scapes usually relatively long ($SI > 80$); metanotal groove moderately impressed to absent; metanotal cross-ribs inconspicuous to absent; propodeal spiracle circular to subcircular; propodeum rounded to angular between dorsum and declivity; propodeal dorsum usually sculptured but never transversely striate; petiolar spiracle situated at the node or immediately in front of the anterior face of the node; body pilosity variable in distribution and density, but usually reduced on the head and mesosoma; mesosoma, petiole and postpetiole usually sculptured; first gastral tergite usually shagreenate or finely sculptured.

II. Synoptic checklist of Arabian *Monomorium*

Monomorium monomorium species-group

1. *Monomorium aeyade* Collingwood & Agosti, 1996
2. *Monomorium brunneolucidulum* Collingwood & Agosti, 1996 (**nomen dubium**).
3. *Monomorium carbonarium* (Smith, 1858)
= *Monomorium minutum* Mayr, 1862
4. *Monomorium clavicorne* André, 1881
= *Monomorium clavicorne punicum* Santschi, 1915a
5. *Monomorium exiguum* Forel, 1894
= *Monomorium exiguum bulawayensis* Forel, 1913
= *Monomorium faurei* Santschi, 1915b
= *Monomorium exiguum flavescens* Forel, 1916
= *Monomorium baushare* Collingwood & Agosti, 1996
= *Monomorium qarahe* Collingwood & Agosti, 1996
6. *Monomorium floricola* (Jerdon, 1851)
= *Monomorium poecilum* Roger, 1863
= *Monomorium cinnabari* Roger, 1863
= *Monomorium specularis* Mayr, 1866

- 218 = *Monomorium impressum* Smith, 1876
- 219 = *Monomorium floreanum* Stitz, 1932
- 220 = *Monomorium angusticlava* Donisthorpe, 1947
- 221 7. *Monomorium holothir* Bolton, 1987
- 222 8. *Monomorium mohammedi* Sharaf & Hita Garcia, 2018
- 223 9. *Monomorium rimae* Collingwood & Agosti, 1996
- 224 10. *Monomorium sarawatense* Sharaf & Aldawood, 2013b
- 225 ***Monomorium salomonis* species-group**
- 226 11. *Monomorium abeillei* André, 1881
- 227 = *Monomorium wahibiense* Collingwood & Agosti, 1996 **syn. n.**
- 228 12. *Monomorium acutinode* Collingwood & Agosti, 1996
- 229 13. *Monomorium areniphilum* Santschi, 1911
- 230 = *Monomorium fezzanense* Collingwood & Agosti, 1996 **syn. n.**
- 231 = *Monomorium hemame* Collingwood & Agosti, 1996 **syn. n.**
- 232 = *Monomorium marmule* Collingwood & Agosti, 1996 **syn. n.**
- 233 = *Monomorium salomonis lepineyi* Santschi, 1934
- 234 = *Monomorium salomonis pullula* Santschi, 1919
- 235 14. *Monomorium asiriense* Collingwood & Agosti, 1996
- 236 15. *Monomorium barbatulum* Mayr, 1877 (New record to KSA)
- 237 16. *Monomorium bicolor* Emery, 1877
- 238 = *Monomorium bicolor aequatoriale* Santschi, 1926
- 239 = *Monomorium bicolor coerulescens* Santschi, 1912
- 240 = *Monomorium bicolor rufibasis* Santschi, 1914
- 241 = *Monomorium bicolor tropicale* Santschi, 1926
- 242 = *Monomorium bicolor uelense* Santschi, 1926
- 243 = *Monomorium phoenicum* Santschi, 1927 **syn. n.**
- 244 17. *Monomorium buettikeri* Collingwood & Agosti, 1996
- 245 18. *Monomorium buxtoni* Crawley, 1920a
- 246 19. *Monomorium carbo* Forel, 1910
- 247 20. *Monomorium dammame* Collingwood & Agosti, 1996
- 248 21. *Monomorium dirie* Collingwood & Agosti, 1996

22. *Monomorium elghazalyi* Sharaf & Aldawood, 2017
23. *Monomorium fayfaense* Collingwood & Agosti, 1996
24. *Monomorium gallagheri* Collingwood & Agosti, 1996
25. *Monomorium hanaqe* Collingwood & Agosti, 1996
26. *Monomorium harithe* Collingwood & Agosti, 1996
- = *Monomorium najrane* Collingwood & Agosti, 1996 **syn. n.**
27. *Monomorium hegyi* Sharaf, **sp. n.**
28. *Monomorium jizane* Collingwood & Agosti, 1996
29. *Monomorium khalidi* Sharaf, **sp. n.**
30. *Monomorium knappi* Collingwood & Agosti, 1996
31. *Monomorium luteum* Emery, 1881
32. *Monomorium mahyoubi* Collingwood & Agosti, 1996
33. *Monomorium moathi* Sharaf & Collingwood, 2010
34. *Monomorium niloticum* Emery, 1881
- = *Monomorium matame* Collingwood & Agosti, 1996 **syn. n.**
35. *Monomorium nimihil* Collingwood, 2004
36. *Monomorium nitidiventre* Emery, 1893
- = *Monomorium yemene* Collingwood & Agosti, 1996 **syn. n.**
37. *Monomorium pharaonis* (Linnaeus, 1758)
- = *Monomorium antiguensis* (Fabricius, 1793)
- = *Monomorium domestica* (Shuckard, 1838)
- = *Monomorium contigua* (Smith, 1858)
- = *Monomorium fragilis* (Smith, 1858)
- = *Monomorium minuta* (Jerdon, 1851)
- = *Monomorium vastator* (Smith, 1857)
38. *Monomorium riyadhe* Collingwood & Agosti, 1996
39. *Monomorium salomonis* (Linnaeus, 1758)
- = *Monomorium debilis* (Walker, 1871)
- = *Monomorium salomonis obscuratum* Stitz, 1917
- = *Monomorium thorense* Mayr, 1862
40. *Monomorium subdenticorne* Collingwood & Agosti, 1996

41. *Monomorium subopacum* (Smith, 1858), full synonymy in Heterick (2006)

42. *Monomorium suleyile* Collingwood & Agosti, 1996

43. *Monomorium tumaire* Collingwood & Agosti, 1996

44. *Monomorium venustum* (Smith, 1858)

III. Key to Arabian *Monomorium* based on the worker caste

Note: *M. brunneolucidulum* excluded due to lack of diagnostic characters.

1. Head smooth and glossy (Fig. 1A) ... **2** (*M. monomorium* species-group)

– Head sculptured, sculpture ranging from dense reticulate-punctate, longitudinal striations, to faint reticular patterning (Fig. 1B) ... **10** (*M. salomonis* species-group)

2. Anterior median portion of clypeus strongly concave with two laterally projecting teeth (Fig. 1C) ... **3**

– Anterior median portion of clypeus straight or feebly concave without raised ridges (Fig. 1D) ... **4**

3. Uniform dark brown to black; metanotal groove broadly and deeply impressed (Fig. 1E) ... *M. carbonarium*

– Bicolored with yellow mesosoma contrasting with the black gaster; metanotal groove shallowly impressed ... *M. rimae*

4. Antenna 12-segmented ... **5**

– Antenna 11-segmented ... **7**

5. Body pilosity clubbed; mesosoma, petiole and postpetiole distinctly sculptured (Fig. 1F) ... *M. sarawatense*

– Body pilosity simple; mesosoma, petiole and postpetiole smooth and shining (Fig. 2A) ... **6**

6. Bicolored, with head and gaster dark brown to black contrasting the yellow or yellow-brown mesosoma and petiole; eyes small (EL 0.21–0.24 x HW) with six ommatidia in longest row, oval in profile (Fig. 2A) ... *M. floricola*

- Uniform yellow; eyes relatively large (EL 0.30–0.32 x HW) with 8-9 ommatidia in longest row, in profile with convex dorsal margin and straight ventral margin (Fig. 2B) ... *M. holothir*
- 7. Mesosoma without standing hairs (Fig. 2C) ... 8**
- Mesosoma with standing hairs (Fig. 2D) ... **9**
- 8. Eyes larger, with a ring of seven to eight ommatidia encircling a single row of 2 ommatidia, and in profile closer to mandibular insertions (EM 0.05); meso- and metapleuron smooth; petiole and postpetiole smooth and each with one pair of standing hairs (Fig. 2C) ... *M. aeyade***
- Eyes smaller, with only 5–6 ommatidia total, and in profile further away from mandibular insertions (EM 0.09–0.11); meso- and metapleuron finely shagreened; petiole and postpetiole superficially shagreened and without standing hairs (Fig. 2E) ... *M. mohammedi*
- 9. Mesosoma with two pairs of standing hairs, one on pronotal corners and one on propodeum (Fig. 2F) ... *M. clavicorne***
- Mesosoma with several pairs of standing hairs, about 10 pairs (Fig. 2D) ... *M. exiguum*
- 10. Underside of head with crowded J-shaped hairs forming a distinct psammophore (Fig. 3A) ... *M. barbatulum***
- Underside of head without long J-shaped hairs, psammophore absent ... **11**
- 11. Mesosoma without standing hairs ... 12**
- Mesosoma with standing hairs ... **25**
- 12. Uniform yellow, brown, or yellow-brown, gaster not darker than mesosoma ... 13**
- Bicolored, with gaster distinctly darker than mesosoma ... **18**
- 13. Metanotal groove shallowly impressed or indistinct (Fig. 3B) ... 14**
- Metanotal groove deeply impressed (Fig. 3C) ... **15**

14. Small yellow species (TL 1.7, HL 0.45–0.48, HW 0.34–0.36, SI 92–97); cephalic surface with vestigial or superficial reticular patterning, almost entirely effaced; petiole and postpetiole without standing hairs; (Fig. 3B); first gastral tergite completely glabrous, *i.e.*, without standing hairs (Fig. 3B) ... *M. heggyi* sp. n.
- Large brown species (TL 2.20–2.75, HL 0.57–0.65, HW 0.51, SI 93–103); cephalic surface with fine and dense reticulate-rugulose sculpture; petiole and postpetiole each with two pairs of standing hairs (Fig. 3D); first gastral tergite always with several pairs of standing hairs (Fig. 3D) ... *M. harithe*
15. Eyes located nearly at the mid-length of head as seen in full-face view (Fig. 3E); eyes smaller, with 5 ommatidia in longest row; posterior margin of head distinctly concave in full-face view (Fig. 3E) ... *M. elghazalyi*
- Eyes located nearly behind or at head mid-length as seen in full-face view (Fig. 3F); eyes distinctly larger, with 10–14 ommatidia in longest row; posterior margin of head concave or linear in full-face view ... 16
16. Underside of head without long, standing hairs (Fig. 4A) ... *M. dirie*
- Underside of head with numerous pairs of long, standing hairs (Fig. 4B) ... 17
17. Yellow; scapes just reaching posterior margin of head when laid back; body pilosity abundant over entire body ... *M. nimihil*
- Brown; scapes surpassing posterior margin of head by about half the length of the pedicel when laid back; body pilosity distinctly reduced over entire surface, mesosoma with a single pair of hairs on propodeum, while the petiole, postpetiole, and gaster are bare ... *M. salomonis*
18. First gastral tergite without standing hairs ... *M. moathi*
- First gastral tergite with hairs either scattered on tergite surface or apically on the posterior margin ... 19
19. Scapes when laid back from their insertions reach or surpass posterior margin of head in full-face view ... 20

– Scapes when laid back from their insertions fail to reach posterior margin of head in full-face view ... **21**

20. Propodeal dorsum in profile meeting declivity forming two blunt, slightly projecting angled bosses (Fig. 4C) ... *M. subdenticorne*

– Propodeal dorsum in profile meeting declivity in a continuous curve (Fig. 4D) ... *M. bicolor*

21. Petiole in the form of a high triangle in profile with anterior face appears as a continuous line sloping anteriorly (Fig. 4E) ... *M. acutinode*

– Petiole broadly rounded with anterior face sloping downward and then anteriorly to the peduncle (Fig. 4F) ... **22**

22. Head faintly superficial sculptured, slightly shining ... *M. venustum*

– Head densely and finely reticulate to reticulate-shagreenate ... **23**

23. Small (TL 2.3–2.4, HL 0.60–0.63, HW 0.43–0.45, PW 0.30–0.31, ML 0.66–0.70); mesosoma and gaster approximately of the same color, the two not strongly contrasting ... *M. carbo*

– Larger (TL 3.1–3.4, HL 0.70–1.04, HW 0.54–0.88, PW 0.36–0.53, ML 0.88–1.24); mesosoma and gaster conspicuously differently colored, gaster usually darker ... **24**

24. Eyes with 12–14 ommatidia in longest row; metanotal groove deeply impressed (Fig. 5A); postpetiole with two to three pairs of backward directed hairs ... *M. areniphilum*

– Eyes with 9–11 ommatidia in longest row; metanotal groove feebly impressed (Fig. 5B); postpetiole with a single pair of backward directed hairs ... *M. subopacum*

25. Large (TL ≥ 3.8 , HW > 0.75) ... **26**

– Smaller (TL 2.2–3.2, $\leq$ HW 0.67) ... **28**

26. Entirely yellowish; mesosoma rather flat with a shallow oblique metanotal groove (Fig. 5C) ... *M. luteum*

– Gaster dark contrasting with red mesosoma; metanotal groove steeply angled (Fig. 5D) ... **27**

403

404 **27.** Head smooth with superficial sculpture; the first of the three segments forming the club being
405 shorter than the second (Fig. 5E); head in full-face view with feebly but distinctly convex

406 sides; head in full-face view with eyes fail or just break head sides (Fig. 5E) ... *M. niloticum*

407 – Head completely finely striate (Fig. 5F); the first of the three segments forming the club nearly
408 subequal to the second (Fig. 5F); head in full-face view with straight sides; head in full-face

409 view with eyes break sides (Fig. 5F) ... *M. riyadhe*

410

411 **28.** Mesosoma red, contrasting with dark gaster ... **29**

412 – Mesosoma pale brown to black, concolorous with gaster ... **34**

413

414 **29.** Mesosoma hairs restricted to one pair on pronotum ... **30**

415 – Mesosoma with several pairs of hairs scattered over whole dorsum ... **31**

416

417 **30.** Head and mesosoma dull red, gaster brown; scapes reaching posterior head margin when laid
418 back (Fig. 6A); cephalic surface with vestigial sculptures; posterior margin of head with a

419 single pair of hairs (Fig. 6A); propodeum in profile with dorsum making an obtuse angle with
420 declivity (Fig. 6B) ... *M. hanaqe*

421 – Head and mesosoma bright orange red, gaster black; scapes not reaching posterior head margin
422 when laid (Fig. 6C); cephalic surface densely punctate; posterior margin of head without hairs

423 except for appressed pubescence (Fig. 6C); propodeum in profile with dorsum making a
424 continuous curve with declivity (Fig. 6D) ... *M. jizane*

425

426 **31.** Propodeal dorsum with a single pair of standing hairs ... **32**

427 – Propodeal dorsum with at least three pairs of standing hairs ... **33**

428

429 **32.** Scapes when laid back from their insertions reach posterior margin of head in full-face view
430 (Fig. 6E); cephalic surface densely punctate (Fig. 6E); gaster smooth and shining (Fig. 6F) ...

431 *M. fayfaense*

- Scapes when laid back from their insertions fail to reach posterior margin of head in full-face view (Fig. 7A); cephalic surface with vestigial sculpture (Fig. 7A); gaster finely densely shagreened and dull (Fig. 7B) ... *M. knappi*
- 33.** Whole body with abundant fine hairs (Fig. 7C); with head in full-face view outer margins of eyes break head sides (Fig. 7D) ... *M. nitidiventre*
- Body pilosity limited and stiff (Fig. 7E); promesonotum with five to six pair of standing hairs, promesonotum and propodeum each with three pairs; petiole and postpetiole each with two-three pairs; with head in full-face view outer margins of eyes fail to break head sides (Fig. 7F) ... *M. khalidi* sp. n.
- 34.** Whole body including gaster densely sculptured and dull ... **35**
- Gaster at least more or less shining with superficial sculpture ... **37**
- 35.** Head, mesosoma, and waist segments very light brown or yellow; hairs on mesosoma scattered (Fig. 8A) ... *M. pharaonis*
- Head, mesosoma, and waist segments conspicuously of darker brown or black; mesosoma hairs mainly or entirely on pronotum (Fig. 8B) ... **36**
- 36.** Body color brown; larger species (TL 3.1–3.3, HW 0.70); scapes long, surpassing posterior head margin by about the length of the pedicel when laid back (Fig. 8C); head in full-face view with convex sides (Fig. 8C) ... *M. suleyile*
- Body color uniformly black; smaller species (TL 2.7–2.9, HW 0.56); scapes shorter, just reaching posterior head margin when laid back of head (Fig. 8D); head in full-face view with parallel sides (Fig. 8D) ... *M. mahyoubi*
- 37.** Whole body glossy, nodes and gaster brilliant ... *M. dammame*
- At least mesosoma with close punctate sculpture ... **38**
- 38.** Underside of head with numerous hairs (12–16), the longest exceeding the maximum eye length (Fig. 8E) ... *M. tumaire*

– Underside of head with fewer hairs, none as long as maximum eye length (Fig. 8F) ... **39**

39. Head finely and densely punctate, general appearance dull ... **40**

– Head feebly superficially sculptured, relatively but distinctly shining ... **41**

40. Smaller species (TL 2.4–2.7); clypeal border feebly concave to straight with a very small median notch or none (Fig. 9A); mesonotum straight in profile (Fig. 9B) ... ***M. gallagheri***

– Larger species (TL > 3.0); clypeus with a distinct anteromedian notch (Fig. 9C); mesonotum distinctly convex in profile (Fig. 9D) ... ***M. buxtoni***

41. Scapes distinctly short, surpassing compound eye posterior margins by only about the length of the pedicel ... ***M. buettikeri***

– Scapes longer, reaching posterior head margin ... **42**

42. Color dark brown to black; smaller species (TL 2.40–2.75, C1 78–91); metanotal groove shallowly impressed ... ***M. abeillei***

– Color light to median brown; larger species (TL 3.2, CI 69–71); metanotal groove distinctly impressed ... ***M. asiriense***

IV. New taxonomic treatments

New site records include, when available, geo-coordinates (°N, °E), elevation (m), collection date, collector, and number worker (w) and queen (q) specimens.

***Monomorium abeillei* André, 1881**

(Fig. 10 A–C)

Monomorium abeillei André, 1881: 531 (footnote) (w.) Israel. Palearctic. (MNHN, CASENT0915411). [Image of lectotype worker examined]. [Also described as new by André, 1881: 67.].

Combination in *Monomorium* (*Xeromyrmex*): Emery, 1922: 177; subspecies of *Monomorium salomonis*: Forel, 1910a: 23; Hamann & Klemm, 1967: 413; revived status as species: Collingwood, 1985: 269; Collingwood & Agosti, 1996: 340.

Monomorium wahibiense Collingwood & Agosti, 1996: 357 (w.) Oman. Palearctic.

[NHMB], CASENT0913864. Syntype worker, Oman, Wahiba dunes (21.vii.1985, M. D.

Gallagher) [examined] **syn. n.**

Material examined. KSA: Asir Province: Raydah (18.198, 42.410, 2443m, 22.ii.2014, M.R. Sharaf, 3w); Raydah (18.204, 42.412, 2820m, 21.ii.2014, M.R. Sharaf, 6w); **Riyadh Province:** Huraymila, Buaythiran (25.149, 45.950, 07.ii.2011, M.R. Sharaf, 2w); Rawdhat Khureim (25.383, 47.277, 618m, 02.vi.2013, S. Salman, 6w); Rawdhat Khureim (25.425, 47.235, 579m, 09.i.2015, S. Salman, 12w); Dirab, KSU research station (24.419, 46.654, 568m, 05.xii.2013, S. Salman, 1w); Wadi Hanifa (24.670, 46.654, 657m, 14.ii.2014, S. Salman, 3w); Mezahmyia (24.472, 46.239, 633m, 25.i.2014, S. Salman, 1w); Mezahmyia (24.466, 46.251, 648m, 29.xi.2014, S. Salman, 1w); Al Hayer (24.546, 46.742, 647m, S. Salman, 2w); Al Hayer (24.557, 46.744, 589m, 11.iv.2014, S. Salman, 6w); Runnah (25.571, 46.973, 615m, 12.iv.2014, S. Salman, 1w); Dawademi (24.478, 44.364, 1027m, 18.iv.2014, S. Salman, 1w); Dawademi (24.583, 44.323, 966m, 16.i.2015, S. Salman, 3w); Dawademi (24.538, 44.355, 999m, 16.i.2015, S. Salman, 1w); Afif (23.766, 42.840, 1015m, 19.iv.2014, S. Salman, 6w); Afif (24.302, 43.688, 951m, 19.iv.2014, S. Salman, 1w); Afif (23.900, 42.081, 1052m, 17.i.2015, S. Salman, 15w); Afif (23.957, 42.976, 1059m, 17.i.2015, S. Salman, 1w); Irgah (24.6710, 46.593, 625m, 19.i.2015, S. Salman, 2w); Thadiq (25.294, 45.871, 735m, 26.iv.2014, S. Salman, 1w); Quwayia (24.047, 45.244, 854m, S. Salman, 7w); Shaqra (25.326, 45.233, 710m, 30.v.2014, S. Salman, 9w); Shagra (25.230, 45.319, 703m, 24.i.2015, S. Salman, 1w); Shaqra (25.270, 45.291, 712m, 23.i.2015, 2w); Durma (24.607, 46.130, 646m, 30.i.2015, S. Salman, 1w); Majma'a (25.880, 45.365, 730m, 07.ii.2015, S. Salman, 5w); Kharrarah (24.392, 46.244, 726m, 08.iv.2015, S. Salman, 15w); Al Ghat (26.066, 44.919, 653m, 31.x.2015, S. Salman, 1w); Hawtet Sudeir (25.592, 45.612, 732m, 31.i.2015, S. Salman, 1w); KSU campus (24.737, 46.618, 662m, 29.ii.2012, K. Mahmoud, 1w); Dirab (24.419, 46.654, 804m, 18.ix.2014, S. Salman, 9w); Hareeq (23.614, 46.054, 689m, 22.ii.2015, S. Salman, 1w); Quwayia (24.058, 45.245, 846m, 29.xi.2014, S. Salman, 1w); Salbouxh (25.078, 46.347, 716m, 26.xii.2014, S. Salman, 8w); **Jazan Province:** Sajid Island, Al-Sajid (16.860, 41.932, 05.iii.2017, U. Abuelgheit, 1w).

Remarks. *Monomorium wahibiense* is represented by a single worker deposited in WML and accompanied by a red card and handwritten label by C. Collingwood indicating that this specimen represents the syntype. The label's data are consistent with the data for the type in the

original description in terms of collecting locality (Oman, Wahiba sand) and collector (M. D. Gallagher), but not the collection date, which we consider a typographical error. Comparing the mentioned type material with the image of the type material of *M. abeillei* André, we found the two species share the same morphological characters, which can be summarized as follow: scapes relatively short, when laid back from their insertions just reaching posterior head margin; eyes of moderate size with about 10-11 ommatidia in longest row; cephalic surface between frontal lobes faintly striated whereas in some individuals the striations are absent; promesonotum and mesonotum forming continuous flat line in profile; mesosoma with single pair of standing hairs on pronotal humeral angles; metanotal groove feebly impressed; petiole with single pair of backward directed hairs; postpetiole with two pairs. However, the eyes are slightly smaller in *M. abeillei* and the central cephalic sculpture is feebly microreticulate-striolate than in the relatively large-eyed *M. wahibiense* and the superficially sculptured cephalic surface but we consider these two traits as variable characters. Herein, we propose treating *M. wahibiense* as a junior synonym of *M. abeillei* André.

Geographic Distribution. *Monomorium abeillei* is originally described from Israel and recorded from several countries in the Middle East, including the Arabian Peninsula (KSA, Kuwait, Oman and Yemen) (Collingwood 1985, Collingwood & Agosti 1996), Iran (Paknia et al. 2008), Israel (Vonshak & Ionescu-Hirsch 2009) and North Africa (Borowiec 2014).

Monomorium areniphilum Santschi, 1911

(Fig. 11 A–C)

Monomorium Salomonis var. *areniphila* Santschi, 1911: 84 (w.) Tunisia. Palearctic. [NHMB], CASENT0249829, [Syntype worker, examined].

Emery, 1915: 378 (q.); combination in *Monomorium* (*Xeromyrmex*): Emery, 1922: 177; subspecies of *Monomorium salomonis*: Santschi, 1936: 50; raised to species: Collingwood, 1985: 269; senior synonym of *Monomorium lepineyi*, *Monomorium pullula*: Bolton, 1987: 336.

Senior synonym of *Monomorium fezzanense* Collingwood & Agosti, 1996: 346 (w.) Saudi Arabia. Afrotropic. [NHMB], Syntype worker, Saudi Arabia, 31 km NW Tabuk (24.iv.1979, CASENT0913557) [examined] **syn. n.**

Senior synonym of *Monomorium hemame* Collingwood & Agosti, 1996: 348 (w.) Kuwait. Palearctic. [WML], holotype worker, Kuwait, Umm Al-Hemam (9.III.1988, W. Biittiker, CASENT0922316) [examined]; paratype worker, Saudi Arabia, Uyaynah (01.IV 1976, W. Biittiker, CASENT0913800) [examined] **syn. n.**

Senior synonym of *Monomorium marmule* Collingwood & Agosti, 1996: 349, fig. 21 (w.) OMAN. Palearctic. Paratype worker, Oman, Minririb (14.i.1986, M.D. Gallagher, CASENT0913824, NHMB) [examined] **syn. n.**

Remarks. A thorough examination of the type material of *M. fezzanense*, *M. hemame*, *M. marmule*, and *M. areniphilum* yielded no evidence for heterospecificity; they are indistinguishable. All four taxa share the following characters: median portion of anterior clypeal margin shallowly concave; eyes large with 12-15 ommatidia in longest row; promesonotum and anterior portion of mesonotum in profile feebly convex; posterior portion of mesonotum sloping steeply to broadly and deeply impressed metanotal groove; mesosoma without hairs; petiole with single pair of backward directed hairs; postpetiole with two pairs of hairs.

Herein *M. fezzanense*, *M. hemame*, and *M. marmule* are treated as junior synonyms of *M. areniphilum*. It is worth mentioning that in the original description of *M. marmule*, Collingwood & Agosti (1996) gave a brief differential diagnosis with *M. areniphilum* based on variable characters such as the presence of mesosomal pubescence, the petiole and postpetiole color and pilosity.

Geographic Distribution. A species originally described from Tunisia and recorded from most countries of the Arabian Peninsula including KSA, Kuwait, Oman, and Yemen (Collingwood, 1985; Collingwood & Agosti, 1996), and the UAE (Collingwood et al., 2011). It is also reported from North Africa and the Afrotropical Region (Bolton, 1987).

Monomorium barbatulum Mayr, 1877

(Fig. 12 A–C)

Monomorium barbatulum Mayr, 1877: 17 (w.) Kazakhstan. Palearctic.

Material examined. KSA: Riyadh Province: Zulfi (26.367, 44.986, 670m, 18.i.2014, Al Dhafer et al., 1w: CASENT0922263, KSMA).

Geographic Distribution. This species was originally described from Kazakhstan and recorded from Oman (Collingwood, 1985; Collingwood & Agosti, 1996), the UAE (Collingwood et al.,

2011), Turkey (Kiran & Karaman 2020), Israel (Vonshak & Ionescu-Hirsch, 2009). The present material represents a new record to the KSA.

***Monomorium bicolor* Emery, 1877**

(Fig. 13 A–C)

Monomorium bicolor Emery, 1877: 368 (w.) Eritrea. Afrotropic.

Senior synonym of *Monomorium phoenicum* Santschi, 1927e: 242 (w.q.) Lebanon.

Palaearctic. Syntype worker, Lebanon, Beyroth (05.viii.1933, Santschi, CASENT0249831, NHMB) [examined] **syn. n.**

Material examined. KSA: Asir Province: Almajardah, Wadi Eltalalei (19.003, 41.732, 223m, 10.xi.2012, M.R. Sharaf, 1w); Wadi Shahadan (17.472, 42.856, 452m, 13.xi.2012, M.R. Sharaf, 13w, (1w, CASENT0906396); Allaith, Adam, Wadi Elarj (20.453, 40.816, 450m, 09.xi.2012, M.R. Sharaf, 45w, 1w: CASENT0906395, 1q: CASENT0906394); Wadi Aljora, near Abadan (17.293, 43.070, 465m, 12.xi.2012, M.R. Sharaf, 2w); Almajardah, Wadi Bagara (18.793, 42.019, 436m, 10.xi.2012, M.R. Sharaf, 2w); **Jazan Province:** Abu Arish (17.013, 42.802, 90m, 10.iv.2012, M.R. Sharaf, 11w); Sabia (17.107, 42.650, 43m, 09.iv.2012, M.R. Sharaf, 23w); Jazan (16.97627, 42.61743, 38m, 12.iv.2012, M.R. Sharaf, 12w); **Al Bahah Province:** Dhi Ayn Archeological village (19.930, 41.443, 741m, 18.v.2011, M.R. Sharaf, 3w); Wadi Gonouna (19.429, 41.605, 353m, 12.v.2011, M.R. Sharaf, 10w: KSMA).

Remarks. The type material of *M. bicolor* and *M. phoenicum* are clearly conspecific. They share the same diagnostic characters as follow: scapes relatively long, when laid back from their insertions surpassing posterior head margin by about length of pedicel; head in full-face view with eyes just breaking sides; cephalic surface dull, finely and densely punctate; median anterior clypeal margin distinctly concave; area between frontal carinae finely longitudinally striated; mesosoma without standing hairs; metanotal groove acutely impressed; propodeal dorsum in profile meeting declivity in continuous curve; petiole and postpetiole each with single pair of back directed hairs; first gastral tergite with hairs scattered over tergite surface; biocolored species, with head, mesosoma, petiole and postpetiole yellow-red or yellow-brown, gaster dark brown to black. Herein, we propose *M. phoenicum* as a junior synonym of *M. bicolor*.

Geographic Distribution. *Monomorium bicolor* was originally described from Eritrea and is a widespread species commonly encountered in open, sandy areas through the Afrotropical Region

(Bolton, 1987). In the Arabian Peninsula, it is known from the KSA and the UAE (Collingwood, 1985; Collingwood & Agosti, 1996; Collingwood et al., 2011).

***Monomorium brunneolucidulum* Collingwood & Agosti, 1996**

Monomorium brunneolucidulum Collingwood & Agosti, 1996: 343. Oman. Palearctic.

Remarks. In their brief original description of the enigmatic species *M. brunneolucidulum* from Oman, Collingwood & Agosti (1996) neither gave successful diagnostic characters nor illustrations for species recognition. In addition, the type-material is apparently lost. Due to a lack of type material and species diagnostic characters, it is impossible to confirm the identity of the species. Until the type material of this species is available we prefer to treat it as a nomen dubium.

***Monomorium floricola* (Jerdon, 1851)**

(Fig. 14 A–C)

Atta floricola Jerdon, 1851: 107 (w.) India. Indomalaya.

Forel, 1893: 388 (q.m.); Wheeler, 1905: 88 (q.m.); Donisthorpe, 1914: 136

(gynandromorph); Crawley, 1920: 217 (gynandromorph); Wheeler & Wheeler, 1955c:

121 (l.). Combination in *Monomorium*: Mayr, 1879: 671.

Senior synonym of *Monomorium poecilum*: Emery, 1894d: 151.

Senior synonym of *Monomorium specularis*: Mayr, 1879: 671.

Senior synonym of *Monomorium cinnabari*: Wheeler, 1913: 486.

Senior synonym of *Monomorium floreanum*: Brown, in Linsley & Usinger, 1966: 175.

Senior synonym of *Monomorium angusticlava*: Bolton, 1987: 390.

Senior synonym of *Monomorium impressum*: Bolton, 1987: 390.

Senior synonym of *Monomorium floricola furina*: Heterick, 2006: 122.

Senior synonym of *Monomorium floricola philippinensis*: Heterick, 2006: 122.

Material examined. Oman: Dhofar Province: Ayn Dirbat (17.106, 54.453, 207m, 17.xi.2017, M.R. Sharaf, 4w: KSMA, 1w: CASENT0922876, CASC).

Geographic Distribution. This species was originally described from India. It is a successful tramp species of putative Southeast Asian origin that is widely distributed throughout tropical

and subtropical regions worldwide (Deyrup et al., 2000; Wetterer, 2010a). The present material represents a new record for Oman and the Arabian Peninsula.

***Monomorium harithe* Collingwood & Agosti, 1996**

(Fig. 15 A–C)

Monomorium harithe Collingwood & Agosti, 1996: 347 (w.) Saudi Arabia. Afrotropic. Holotype worker: Saudi Arabia, desert near Najran, 17.533, 44.000, 10.iv.1983, C.A. Collingwood, CASENT0922335, WML, [examined] **syn. n.**

Monomorium najrane Collingwood & Agosti, 1996: 352 (w.) Saudi Arabia. Afrotropic, Holotype worker: Saudi Arabia, Najran, semi desert, iv.1984, C.A. Collingwood, CASENT0922335, WML, [image examined] **syn. n.**

Previous records. **KSA:** Riyadh (24.714, 46.675, 21.i.1980, A.H. Talhouk, 2w); **Yemen:** Taiz (13.578, 44.018, 20.iii.1993, C.A. Collingwood, 2w).

Remarks. *Monomorium harithe* was described from the KSA and Yemen, while *M. najrane* was described from Najran (KSA) near the Saudi-Yemeni borders (Collingwood & Agosti 1996). The comparison of the type material of the two species reveals a straightforward synonymy. The two species share the following characters: anterior median clypeal margin distinctly concave; scapes distinctly short, when laid back from their insertions failing to reach posterior head margin; mesosoma without standing hairs; metanotal groove feebly impressed but distinct; propodeal dorsum with distinct furrow); mesosoma, petiole and postpetiole finely and densely punctate; petiole and postpetiole each with single pair of back directed hairs; gaster smooth and shining. In addition, the two species share common body measurements (e.g. HW 0.51; SL 0.53).

Geographic Distribution. This Arabian endemic species is only known from the KSA and Yemen (Collingwood & Agosti, 1996).

***Monomorium hegyi* Sharaf, sp. n.**

(Fig. 16 A–C)

Holotype pinned worker. KSA: Al Bahah Province: Shada Al A'la Mountain (19.877, 41.311, 897m, Al Dhafer et al., MRS0261, CASENT0746641, KSMA).

Paratype pinned workers. KSA: Al Baha Province: Shada Al A'la Mountain (19.877, 41.311, 897m, 04.vi.2014, M. R Sharaf, 4w, 1w: CASENT0746641); Shada Al A'la Mountain (19.863,

41.301, 1225m, 08.xii.2014, Al Dhafer et al., 6w, 1w: CASENT0922301); Shada Al A'la Mountain (19.843, 41.312, 1666m, 23.viii.2014, Al Dhafer et al., 1w: CASENT0906391); Shada Al A'la Mountain (19.877, 41.311, 897m, 04.vi.2014, M.R. Sharaf, MRS0261, 2w); Shada Al A'la Mountain (19.877, 41.311, 897m, 23.viii.2014, M.R. Sharaf, MRS0261, 1w: CASENT0746641); Shada Al A'la Mountain (19.863, 41.301, 1225m, 08.xii.2014, Al Dhafer et al., 1w); Shada Al A'la Mountain (19.851, 41.301, 1325m, 15.ii.2014, Al Dhafer et al., 1w); Shada Al A'la Mountain (19.839, 41.310, 15.xi.2015, Al Dhafer et al., 3w: KSMA).

Other Material. Jazan Province: Zabia (17.107, 42.650, 43m, 09.iv.2012, M. R Sharaf, MRS0070, 2w); Abu Arish (17.013, 42.802, 90m, 10.iv.2012, M. R Sharaf, MRS0073, 10w); Jazan (16.97627, 42.61743, 38m, 12.iv.2012, M. R Sharaf, MRS0077, 25w: KSMA, 1w: CASC, 1w: BMNH, 1w: WML, 1w: OXUM, 1w: LACM).

Measurements. Holotype: TL 2.01; HL 0.49; HW 0.39; SL 0.42; EL 0.12; EM 0.07; ML 0.59; PW 0.25; PTL 0.11; PTW 0.11; PPL 0.09; PPW 0.09; CI 80; EI 31; SI 108.

Paratype workers: TL 1.50–2.55; HL 0.42–0.49; HW 0.32–0.39; SL 0.28–0.42; EL 0.07–0.12; EM 0.07–0.09; ML 0.45–0.59; PW 0.21–0.31; PTL 0.09–0.14; PTW 0.07–0.11; PPL 0.07–0.09; PPW 0.07–0.09; CI 69–86; EI 21–31; SI 80–108 (n=13).

Diagnosis. *Monomorium hegyi* is diagnosed by the following character combination: scapes when laid back from their insertions just reaching posterior margin of head; head in full-face view with eyes located nearly at midlength; promesonotal outline feebly convex or flat, sloping posteriorly to narrow and shallowly impressed metanotal groove.

Worker. Head. Head in full-face view distinctly longer than broad, with concave posterior margin and shallowly convex sides and feebly concave posterior margin; median portion of clypeus with anterior free margin slightly indented; eyes of moderate size, in profile with convex dorsal sides and straight ventral side, maximum diameter 0.21 x– 0.30 x HW, with 7 ommatidia in longest row; head in full-face view eyes located nearly at midlength of head and just breaking sides; scapes when laid back from their insertions just reaching posterior head margin.

Mesosoma. Promesonotal outline feebly convex or flat, sloping posteriorly to narrow and shallowly impressed metanotal groove; propodeal dorsum in profile convex making a continuous curve with propodeal declivity and with defined lateral margins. **Petiole.** Petiole with high rounded node in profile; subpetiolar process broad and blunt. **Postpetiole.** Postpetiolar node lower than petiolar node in profile and nearly as broad as petiole in dorsal view. **Sculpture.**

Mandibles feebly longitudinally sculptured; cephalic surface with faint vestiges of superficial reticular patterning, almost entirely effaced, area between frontal carinae finely longitudinally striate; clypeus smooth; entire mesosoma, petiole and postpetiole sharply and densely reticulate-punctate; gastral tergites smooth and shining. **Pilosity.** Dorsum of head without standing hairs behind the level of the frontal lobes; several pairs of long hairs on the anterior clypeal margin and on mandibles; antennae with dense appressed pubescence; mesosoma, petiole and postpetiole without standing hairs of any description; first gastral tergite without standing hairs except for sparse appressed pubescence; pilosity of remaining gastral tergites restricted to the posterior margins. **Color.** Uniformly yellow.

Remarks. *Monomorium heggyi* belongs to the *M. salomonis* species-group (Bolton 1987). It is most similar to *M. rabirium* Bolton, 1987 from Botswana from which it is readily distinguished by the longer scapes (SI 80–108) that reach the posterior head margin in full-face view, the posteriorly shifted eyes located nearly at the midlength of head in full-face view. *Monomorium rabirium* has shorter scapes (SI 92–97) that fail reaching the posterior head margin in full-face view, and eyes conspicuously located in front of midlength of head in full-face view. Among the Arabian species of the *M. salomonis* species-group, *M. heggyi* is superficially similar to *M. elghazalyi* from the Socotra Archipelago from which it can be easily separated by the larger eyes (EI 21–31), the shallowly impressed metanotal groove, and the densely sculptured mesosoma, petiole and postpetiole. *Monomorium elghazalyi* has smaller eyes (EI 19–20), broadly and deeply impressed metanotal groove, and a smooth body surface.

Etymology. The patronymic name honors Essam Heggy, the Egyptian space scientist at NASA.

Habitat. The type locality, Shada Al A'la, is a Nature Reserve (**Fig. 17**) located in the Al Bahah Province in the southwestern KSA at an elevational range of 470–2,222 m. The locality is characterized by relatively high rainfall, diverse habitats, and high biodiversity, as well as by the presence of large areas of terraced fields used for cultivating banana, coffee, figs, and lemon. The region has a diverse range of wild vegetation cover including plants of the Leguminosae (Fabaceae), composites (Asteraceae), and graminoides (Poaceae). Acacia (Fabaceae) and Juniper (Cupressaceae) are the most dominant plants (SWA, 2018, El-Hawagry et al., 2016). Shada Al-A'Ala harbors a high number of endemic animals including birds, mammals (SWA, 2018) and insects (El-Hawagry et al., 2016).

Geographic Distribution. KSA.

***Monomorium khalidi* Sharaf, sp. n.**

(Fig. 18 A–C)

Holotype pinned worker. KSA: Al Bahah Province: Shada Al A'la (19.839, 41.310, 1563m, 18.x.2014, Al Dhafer et al., CASENT0922288, KSMA).

Three paratype, pinned workers, **KSA: Jazan Province:** Wadi Shahdan (17.452, 42.715, 200m, 13.xi.2012, M.R. Sharaf, MRS0131, CASENT0919810, KSMA); Fayfa, Wadi Al Jora (17.279, 43.062, 419m, 06.iv.2013, M.R. Sharaf, KSMA).

Measurements. Holotype: TL 3.16; HL 0.75; HW 0.67; SL 0.62; EL 0.14; EM 0.25; ML 0.95; PW 0.44; PTL 0.27; PTW 0.21; PPL 0.17; PPW 0.21; CI 89; EI 21; SI 93.

Paratype workers: TL 2.31–2.98; HL 0.59–0.72; HW 0.49–0.59; SL 0.45–0.59; EL 0.14–0.15; EM 0.11–0.18; ML 0.71–0.85; PW 0.32–0.41; PTL 0.28–0.32; PTW 0.11–0.15; PPL 0.11–0.17; PPW 0.08–0.15; CI 81–83; EI 24–29; SI 92–102 (n=3).

Diagnosis. *Monomorium khalidi* can be distinguished by the combination of the following characters: short scape failing to reach posterior head margin in full-face view; abundant mesosomal pilosity; straight outline of promesonotum; densely reticulate-punctate surfaces of head, mesosoma, petiole, and postpetiole; promesonotum dorsally with at least five to six pair of hairs, promesonotum and propodeum each with three pairs.

Worker. Head. Head nearly as long as broad, or little longer than broad with concave posterior margin and feebly convex sides; median portion of clypeus with anterior free margin distinctly concave; eyes of moderate size, in profile view with convex dorsal sides and straight ventral side, maximum diameter 0.20 x HW, with 9-10 ommatidia in longest row); head in full-face view with eyes failing to break head sides; scapes when laid back from their insertions failing to reach posterior margin. **Mesosoma.** Promesonotal outline flat, slopping posteriorly to narrow and feebly impressed metanotal groove; propodeal dorsum flat and short, longitudinally concave, with sharply defined lateral margins. **Petiole.** Petiole with high rounded node in profile.

Postpetiole. Postpetiole as broad as petiole in dorsal view. **Sculpture.** Cephalic surface between and immediately behind frontal lobes finely longitudinally striate; cephalic surface and sides, entire mesosoma, petiole and postpetiole sharply and densely reticulate-punctate; first gastral tergite shagreened and relatively shining. **Pilosity.** Cephalic surface with several pairs of standing hairs behind level of frontal lobes; posterior margin of head with three pairs of standing

hairs; underside of head with about five pairs of short hairs; promesonotum dorsally with at least five to six pair of hairs, promesonotum and propodeum each with three pairs; petiole and postpetiole each with two-three pairs of backward directed hairs; first gastral tergite with numerous standing hairs which are evenly distributed over the sclerite in front of the apical transverse row. **Color.** Bicolored species, head, mesosoma, petiole, postpetiole and appendage light red brown, gaster black.

Remarks. This new species is a member of the *M. salomonis* species-group (Bolton, 1987). *Monomorium khalidi* is closest to *M. junodi* Forel, 1910 from South Africa in terms of the relatively small eyes, the short scapes that fail to reach posterior margin of head, the acute metanotal groove, and densely punctate body surface. The two species have short scapes that fail to reach posterior margin of head in full-face view, metanotal groove feebly impressed; body surface densely punctate; and propodeal longitudinally concave, with sharply defined lateral margins. However, *M. khalidi* can be easily distinguished from *M. junodi* by the following characters: body bicolored with head, mesosoma, petiole, postpetiole and appendages light red-brown contrasting with the black gaster; head in full-face view with eyes failing to break sides; promesonotum dorsally with at least five to six pair of hairs, promesonotum and propodeum each with three pairs; promesonotal outline flat. *Monomorium junodi* is uniformly brown to dark brown, head in full-face view with eyes breaking sides; promesonotum dorsally with two pair of hairs, propodeum without hairs; promesonotal outline feebly but distinctly convex. Among the Arabian *Monomorium* species, *M. khalidi* is superficially similar to *M. nitidiventre* in terms of body size, surface sculpture, eye shape but the former can be readily recognized by the reduced stiff pilosity.

Etymology. The patronymic name honors Khalid Amr (born at 04/11/2012) the son of the second author.

Habitat. The type locality of *M. khalidi* is Shada Al A'la (**Fig. 19**), the same locality where *M. heggyi* was collected.

Geographic Distribution. KSA.

Monomorium niloticum Emery, 1881

(Fig. 20 A–C)

Monomorium niloticum Emery, 1881: 533 (w.) Egypt. Palearctic. [MSNG], [Syntype worker, CASENT0905755, image examined].

Combination in *Monomorium* (*Xeromyrmex*): Emery, 1922: 179; subspecies of *Monomorium venustum*: Forel, 1910: 6; Wheeler & Mann, 1916: 170; Stitz, 1917: 346; Finzi, 1936: 175; revived status as species: Santschi, 1936: 37; see also: Collingwood & Agosti, 1996: 352; current subspecies: nominal plus *M. n. gracilicorne*, *M. n. niloticoides*.

Senior synonym of *Monomorium matame* Collingwood & Agosti, 1996: 350, fig. 22 (w.) OMAN. Palearctic, Holotype worker, Oman, Wadi Matam (01.II.1986, M.D. Gallagher, CASENT0922325, WML) [examined] **syn. n.**

Material examined. KSA: Asir Province: Jebel Al Habala (18.038, 42.873, W. Buttiker, 1w); Alkhola (13.600, 44.283, 4w, WML). **Al Baha Province:** Al Atawla, Al Baha-Taif RD, Wadi Bawa (20.750, 41.247, 1310m, 08.xi.2012, M.R. Sharaf, MRS0099, 7w, 1w: CASENT0906397); Wadi Elzaraeb (20.073, 41.387, 2086m, 09.v.2011, M.R. Sharaf, 3w); Al Mandaq, Wadi Turabah (20.242, 41.263, 1715m, 19.ix.2011, M.R. Sharaf, 6w); Hawtat Bani Tamim (23.525, 46.845, 19.iv.2008, M.R. Sharaf, 6w); Riyadh city (06.viii.2008, 1w); Wadi Hanifa (24.671, 46.595, 641m, 11.iv.2013, M.R. Sharaf, 11w); Al Mandaq, Wadi Turabah (20.211, 41.288, 1793m, 10.v.2011, M.R. Sharaf, 4w). **Riyadh Province:** Al Hayer (24.280, 46.766, 647m, 10.iii.2011, A.S. Aldawood, 24w); Al Hayer (24.557, 46.744, 589m, 11.iv.2014, S. Salman, 5w); Dawademi (24.557, 44.377, 983m, 18.iv.2014, S. Salman, 6w); Dawademi (24.478, 44.364, 1027m, 18.iv.2014, S. Salman, 4w); Dawademi (24.583, 44.323, 966m, 16.i.2015, S. Salman, 9w); Dawademi (24.538, 44.355, 999m, 16.i.2015, S. Salman, 1w); Afif (23.900, 42.881, 1052m, 17.i.2015, 1w); Afif (23.957, 42.976, 1059m, 17.i.2015, S. Salman, 6w); Dhurma (24.607, 46.130, 646m, 30.i.2015, S. Salman, 8w); Hawtat Bani Tamim (23.454, 46.819, 582m, 19.ii.2015, M.R. Sharaf, 3w); Hawtat Bani Tamim (23.500, 46.850, 612m, 19.ii.2015, M.R. Sharaf, 2w); Hareeq (23.614, 46.054, 689m, 22.ii.2015, S. Salman, 29w); Quwayia (24.070, 45.280, 823m, 03.v.2014, S. Salman, 3w); Majma'a (26.005, 45.019, 730m, 13.ix.2014, S. Salman, 6w, KSMA); Riyadh (23.953, 43.636, x.1979, W. Buttiker, 2w); Wadi Eflah, x.1983, W. Buttiker, 2w); wadi Mawran (22.050833, 46.671944, 10.ii.1985, W. Buttiker, 6w); Riyadh (24.7136, 46.6753, 07.vii.1975, W. Buttiker, 1w); Shoiba (Shuaibah) (20.6295, 39.5624, 06.xii.1983, W. Buttiker, 2w); Wadi Nimar (24.5705, 46.68, v.1983, W. Buttiker, 2w); Harithi (21.28, 40.28, 11.v.1984, W. Buttiker, 1w); Wadi Ellah (09.ix.1986, W. Buttiker, 3w, WML);

Malham (25.154, 46.282, 711m, 15.ix.2014, S. Salman, 4w); Malham (25.161, 46.229, 742m, 15.ix.2014, S. Salman, 1w); Quwayia (24.058, 45.245, 846m, 29.x.2014, S. Salman, 4w); Quwayia (24.05043, 45.25795, 839m, 29.x.2014, S. Salman, 12w); Quwayia (24.053, 45.262, 836m, 29.xi.2014, S. Salman, 2w); Na'jan (24.026, 47.138, 467m, 13.xii.2014, S. Salman, 9w); Wadi Al Dawaser (22.778, 44.786, 686m, 20.ii.2015, S. Salman, 3w); Wadi Al Dawaser (20.778, 44.786, 686m, 20.ii.2015, S. Salman, 3w); Qassim, Buraydah (26.216, 44.0414, 633m, 17.ix.2011, Steyaningrum, 9w); Qassim, Buraydah (26.330, 43.979, 623m, 19.x.2013, M.R. Sharaf, 9w); Huraymila (25.1487, 45.950, 815m, 07.ii.2011, M.R. Sharaf, 9w); Dirab Research Station (24.419, 46.654, 570m, 28.ix.2011, B.L. Fisher, 1w, CASENT0260164; Dirab Research Station (24.737, 46.618, 662m, 29.ii.2012, K. Mahmoud, 27w); Naam Dam (23.628, 46.631, 646m, 22.ii.2015, S. Salman, 30w); Dirab (24.409, 46.662, 588m, 30.xii.2009, M.R. Sharaf, 6w); Salbouxh (25.079, 46.347, 689m, 05.xi.2009, M.R. Sharaf, 15w); Salbouxh (25.074, 46.377, 728m, 26.xi.2014, S. Salman, 12w); Ghiyanah (25.074, 46.226, 728m, 26.xii.2014, S. Salman, 3w); Al Hayer (24.280, 46.766, 10.iii.2011, A.S. Aldawood, 10w); Qassim, Buraydah (26.338, 44.024, 643m, 19.x.2013, S. Salman, 2w); Wadi Al Dawaser (20.487, 44.764, 690m, 22.i.2014, S. Salman, 1w, All previous material in KSMA); Wadi Khumra (17.viii.1979, W. Buttiker, CASENT0249836, 1w, NHMB). **Mekkah Province:** Ras Hatibah (21.978, 38.937, 11.i.1983, 2w, WML). **OMAN:** Nakhl (23.44696, 57.88062, 364m, 02.iv.2016, M.R. Sharaf, 8w, CASENT0922306); Dhofar, Dhalkout (16.727, 53.249, 623m, 18.xi.2017, M.R. Sharaf, 6w, CASENT0922859, KSMA); no locality (2005, 1w); no locality (xi.1984, 1w); Jebel Akhdar (23.073, 54.662, 1w); Wattayah (23.591, 58.363, 1983, 3w, WML). **UNITED ARAB EMIRATES:** Wadi Madaq (25.300, 56.117, 22.xi-02.xii.2010, M. Hauser et al., UAE12977, 1w, CASENT0264568, KSMA); Hatta (24.806, 56.125, iii.1998, A.V. Harten, 2w); Hatta (24.806, 56.125, iii.1995, A.V. Harten, 2w, WML). **YEMEN:** Ta'iz (13.578, 44.018, A.V. Harten, 1w); Al Kawd (13.089, 45.365, 1991, A.V. Harten, 2w, WML).

Remarks. *Monomorium matame* was described from Oman and KSA based on the worker caste. In the original description Collingwood & Agosti (1996) pointed out the similarity between *M. matame* and *M. niloticum* and used variable characters that were not useful in species recognition. Our examination of the type material of both shows that *M. matame* is not separable from *M. niloticum*. The two species share the following characters: scapes when laid back from their insertions failing to reach posterior head margin; eyes relatively large about 0.30–0.33 x

HW; metanotal groove deeply impressed; mesosoma with several pairs of scattered standing hairs; promesonotum with three pairs, mesonotum with two to three pairs, propodeum with a single pair of hairs; bicolored species with head, mesosoma, petiole, postpetiole red-brown contrasting with dark brown to black gaster. Based on the examination of the type images of both species, we propose to synonymize *M. matame* as a junior synonym of *M. niloticum*, on the basis of morphological similarity.

Geographic Distribution. *Monomorium niloticum* is originally described from Egypt and widely spread in the Arabian Peninsula (Collingwood, 1985; Collingwood & Agosti, 1996; Collingwood et al., 2011, Sharaf et al., 2018b). It is also collected from Israel (Vonshak & Ionescu-Hirsch, 2009), and North Africa (Sharaf, 2006; Borowiec, 2014). *Monomorium niloticum* is one of the most broadly spread myrmicine species throughout several countries of the Arabian Peninsula including the KSA, Oman, UAE, and Yemen (Collingwood, 1985; Collingwood & Agosti, 1996).

Ecological and Biological notes. The broad geographic distribution of the species can be interpreted in the light of the species' diverse habitat preferences, including the deserts, mountainous, and cultivated sites. Several worker series were found nesting in either dry or humid soil beneath rocks in an undisturbed site in the KSA where a broad diverse of plant species exists including *Acacia* (Fabaceae), *Citrus limon* (L.) Osbeck (Rutaceae), *Prunus dulcis* (Mill.) D. A. Webb (Rosaceae), *Juniperus* L. (Cupressaceae), *Mangifera indica* L. (Anacardiaceae), *Ficus* sp. (Moraceae), *Hibiscus* L. (Malvaceae) and *Azadirachta indica* A. Juss. (Meliaceae).

Monomorium nitidiventre Emery, 1893

(Fig. 21 A–C)

Monomorium bicolor subsp. *nitidiventris* Emery, 1893: 256 (w.) Egypt. Palearctic. [MSNG], [Syntype worker, CASENT0904602, image examined].

Mayr, 1901: 7 (q.); Karavaiev, 1911: 5 (m.); combination in *Monomorium* (*Xeromyrmex*):

Wheeler, 1922: 869; subspecies of *Monomorium bicolor*: Santschi, 1938: 39; of *Monomorium subopacum*: Santschi, 1927: 245; Menozzi, 1932: 94; Bernard, 1953: 159; Hamann & Klemm, 1967: 413; raised to species: Collingwood, 1985: 272; see also: Wheeler & Mann, 1916: 171; Collingwood & Agosti, 1996: 352.

Monomorium yemene Collingwood & Agosti, 1996: 357, fig. 32 (w.) Yemen. Afrotropic.

Holotype worker, Yemen, Taiz (20.x.1991, A. van Harten, NHMB) [Presumed lost];

Paratype worker, Yemen, Zingibar- Shuqrah (13.356, 45.700, 21.iii.1993, C.A.

Collingwood, CASENT0913865, NHMB) [examined] **syn. n.**

Material examined. **YEMEN**, W. Adem Port, Wadi Tiban, N. W. of Jebel Jihaf (13.198, 44.787, ~1158 m, 22.x.1937, C.A. Collingwood, from flower of *Adenium* sp., B. M. Exp. To S. W. Arabia, H. Scott & E. B. Britton, B. M. 1938-246, BMNH (E) 1017382, 1w, CASENT0914158, BMNH).

Remarks. The synonymy of *M. yemene* with *M. nitidiventre* is straightforward since both are morphologically similar and indistinguishable. Both share the following key characters: posterior margin emarginated in full-face view; median portion of anterior clypeal margin distinctly concave; metanotal groove deeply impressed; head, mesosoma, petiole, and postpetiole densely reticulate-punctate and covered with abundant pale standing hairs.

Note: However, the locality label of the paratype specimen (CASENT0913865) (Madinat Al shiraq) is not matching the locality data mentioned in the original description but the collecting data and collector are congruous with the description, therefore, this specimen is treated as one of the type material of *M. yemene*.

Geographic Distribution. *Monomorium nitidiventre* is originally described from Egypt and recorded from the KSA, Kuwait and Yemen (Collingwood, 1985; Collingwood & Agosti, 1996).

Monomorium subdenticorne Collingwood & Agosti, 1996

(Fig. 22 A–C)

Monomorium subdenticorne Collingwood & Agosti, 1996: 354, fig. 27 (w.) Yemen. Afrotropic.

Material examined. KSA: Asir Province: Ahad Refedah (18.134, 43.001, 2179m, 23.ii.2015, M. Alharbi, 2w); (KSMA); **Yemen:** Ghaiman, about 9 miles S. E. of San'a (13.933, 44.833, ~8400 ft, 18.ii.1938, B. M. Exp. To S. W. Arabia, H. Scott & E. B. Britton, B. M. 1938-246, BMNH1017456, 1w, CASENT0914318, BMNH).

Geographic Distribution. A species originally known from Yemen and herein collected for the first time from the KSA.

Monomorium venustum (Smith, 1858)

(Fig. 23 A–C)

Myrmica venusta Smith, 1858: 126 (w.) Syria. Palearctic.

Material examined. KSA: Asir Province: Wadi Asidah (20.417, 41.200, 10.ix.1983, 1480m, W. Buttiker, 1w, WML). **Riyadh Province:** Zulfi (26.272, 44.771, 635m, 18.i.2014, S. Salman, 15w); Mezahmyia (24.466, 46.251, 648m, 29.xi.2014, S. Salman, 10w); Dawademi (24.552, 43.932, 873m, 16.i.2015, S. Salman, 1w); Bijadriyah (24.310, 43.731, 439m, S. Salman, 2w); Afif (23.900, 42.081, 1052m, 17.i.2015, S. Salman, 18w); Shaqra (25.270, 45.291, 712m, 23.i.2015, 63w); Sajir (25.165, 44.601, 750m, 23.i.2015, S. Salman, 6w); Shaqra (25.274, 45.300, 707m, 24.i.2015, S. Salman, 2w, KSMA). **Kuwait:** Kuwait (no date, 2w, WML). **Oman:** Wahiba sands (21.438, 58.554, 23.iii.1986, W. Buttiker, 1w, WML). **United Arab Emirates:** Jebel Hafit (24.050, 55.767, 27.ii-03.iii.2011, M. Hauser et al., UAE13010, 1w, CASENT0264463); Um Al-Quwain (12. iv-07.vi.2009, M. Hauser et al., UAE12920, 1w, CASENT0264584); Ar-Rafah (25.717, 55.867, 15-30. x.2010, M. Hauser et al., UAE12866, CASENT0264475, KSMA); Wadi Asidah (20.417, 41.200, 10.ix.1983, W. Buttiker, 1w); Riyadh (24.714, 46.675, 18.ii.1975, W. Buttiker, 1w); Sharjah (25.346, 55.421, vii.2003, A.V. Harten, 2w, WML). **Yemen:** Sana'a (15.369, 44.191, iii.1990, 1w, WML).

Geographic Distribution. *Monomorium venustum* is originally described from Syria and recorded from the KSA, Kuwait, Oman (Collingwood, 1985; Collingwood & Agosti, 1996; Sharaf et al., 2018b), Israel (Vonshak & Ionescu-Hirsch, 2009), and North Africa (Borowiec, 2014).

V. Biogeographical analysis

Twenty-four species are Arabian endemics. Afrotropical (5) and Palearctic (12) species were also identified (Table 1, Figs. 24–29).

DISCUSSION

Monomorium is one of the most diverse ant genera in the world, but it is rarely the most speciose genus on a regional scale. However, the Arabian *Monomorium* fauna, based on review of previous literature data and on our current work, includes 44 species, making it the most diverse known ant genus of the Arabian Peninsula (Collingwood, 1985; Collingwood & Agosti, 1996; Collingwood et al., 2011; Aldawood & Sharaf, 2011; Sharaf & Aldawood, 2013b; Sharaf

et al., 2015, 2017a, 2018 a, b). Currently, the genus represents ~14% of the total number of species reported from the region (312 spp.) (e.g. Collingwood, 1985; Collingwood & Agosti, 1996; Collingwood et al., 2011; Aldawood & Sharaf, 2011; Sharaf et al., 2011, 2013c, 2015, 2017a, b, Sharaf et al., 2018a, b; Sharaf & Aldawood, 2019; Sharaf et al. 2019; 2020). This value is lower than the 20% mentioned by Collingwood & Agosti (1996). This reduction is the result of numerous taxa in previous studies now treated as junior synonyms of other species (Sharaf et al., 2015, 2017a, 2018a; this study).

We found that ~55% of the *Monomorium* species (24/44) appear to be endemic to the Arabian Peninsula. High degrees of endemism have been reported for many of Arabian arthropod groups, including ants in general (Collingwood, 1985; Collingwood & Agosti, 1996; Sharaf et al. 2014a & b; 2017a, 2018b), staphylinid beetles (Assing et al., 2013; Hlaváč et al. 2013), carabid beetles (Abdel-Dayem et al. 2018), termites (Cowie 1989), lepidopterans (Larsen 1984), and pseudoscorpions (Mahnert et al. 2014).

Biogeographically, the biota of the Arabian Peninsula does not constitute a cohesive unit (Larsen, 1984; Sharaf et al., 2020a). Instead, the Arabian Peninsula is often considered to be at the nexus of two terrestrial biogeographic realms, the Palearctic and the Afrotropic. In fact, Olson et al. (2001) places the northern and central Arabian Peninsula in the Palearctic bioregion (along with Europe, northern Africa, Asia north of the Himalayas, and neighboring islands) and the southern and eastern coasts of the Arabian Peninsula in the Afrotropic bioregion (along with sub-Saharan Africa, southern Iran, southwestern Pakistan, and neighboring islands). Our Geographic distribution data (see also Table 1) of Arabian *Monomorium* supports this basic division, with 12 Palearctic species more common in the north and central deserts and the five Afrotropic species more common in the southern region and along the coasts.

The majority of the endemic Arabian ant species has faunal similarities with taxa from the Afrotropic bioregion that has been earlier documented by several studies (e.g. Larsen, 1984; Collingwood, 1985; Collingwood & Agosti, 1996; Sharaf & Aldawood, 2011, 2012, 2019; Sharaf et al., 2012a, b, c & 2020 a, b; El-Hawagry et al., 2013, 2017; Hájek & Reiter, 2014). Therefore, it is not surprising that a large proportion of the endemic Arabian *Monomorium* species (13/24) have been found in the mountainous ranges of southwestern KSA that extend to Yemen.

Both new *Monomorium* species reported here, *M. heggyi* and *M. khalidi*, were collected in the Shada Al-A'Ala Nature Reserve (SANR), a protected area consisting of an isolated granite mountain massif in southwestern Saudi Arabia. Its location, elevational range (470–2,222 m) and high rainfall resulting in diverse microclimates and a high biodiversity (SWA 2020). As a unique biodiversity hotspot, the SANR contains ~495 plant species (~22% of the total reported Saudi Arabian flora), including 43% of the threatened plant species and 19 endemic plants (Thomas et al., 2017). The SANR also protects a diverse fauna, including rare and endemic vertebrates, including the griffon vulture (*Gyps fulvus* (Hablitz)), the Arabian leopard (*Panthera pardus nimr* (Hemprich and Ehrenberg)), and the Arabian wolf (*Canis lupus arabs* Pocock) (SWA, 2020). The SANR invertebrate fauna has attracted relatively little attention, but recent insect biodiversity inventories and monitoring research projects conducted by King Saud University Museum of Arthropods resulted in two important faunistic studies that recorded 119 Diptera species (El-Hawagry et al., 2016) and 62 carabid beetle species (Abdel-Dayem et al., 2019). Further studies are planned to be carried out at SANR to explore additional levels of biodiversity.

In addition to the native *Monomorium* species, there are two *Monomorium* known from the Arabian Peninsula that are cosmopolitan tramp species, spread around the world through human commerce: *M. pharaonis* and *M. floricola*. The pharaoh ant, *M. pharaonis*, is a common domestic pest. Although it was first described from Egypt, its original native range is uncertain (Wetterer 2010b). We report the first known Arabian record of *M. floricola*, an Indomalayan species, from a single site in Oman. Although widespread around the world, *M. floricola* is rarely considered a serious pest. However, because this species is very small, slow moving, cryptically colored, and primarily arboreal, its abundance and ecological importance may be underappreciated (Wetterer, 2010a).

Considering the high degree of endemism encountered, it is likely that the known Arabian *Monomorium* fauna will increase in the future with further exploration of poorly surveyed areas of the Arabian Peninsula, especially the southwestern mountains of the KSA, Yemen, and the mountainous regions of Oman and the UAE. We hope that the present study will serve as a cornerstone of future taxonomic treatments of *Monomorium* in the Arabian Peninsula.

ACKNOWLEDGMENTS

The authors thank Barry Bolton for useful discussions about *Monomorium* over years. We are indebted to the following colleagues: Boris Kondratieff for useful comments, Brian Fisher for kind permission to use *Monomorium* images on AntWeb, Michele Esposito for imaging the new species, Mahmoud Abdel-Dayem for creating the distribution maps, and Ahmed Shams Al ‘Ola for technical assistance with figures. Special thanks to Brian Heterick and Kadri Kiran for valuable suggestions. The authors extend their appreciation to the Deputyship for Research & Innovation, “Ministry of Education “ in Saudi Arabia for funding this research work through the project no.(IFKSURG-1436-029).

REFERENCES

- Abdel-Dayem MS, Elgharbawy AA, Rasool I., Nagel P, Aldhafer HM. 2019. The Carabidae (Coleoptera) of Shada Al-A'Ala Nature Reserve, Southwestern Saudi Arabia, with description of a new species of Paussinae. *Zookeys* **812**: 93–131. doi: 10.3897/zookeys.812.30937
- Aldawood AS, Sharaf MR. 2011. *Monomorium dryhimi* sp. n., a new ant species (Hymenoptera, Formicidae) of the *M. monomorium* group from Saudi Arabia, with a key to the Arabian *Monomorium monomorium*-group. *ZooKeys* **106**: 47–54. doi:10.3897/zookeys.106.1390
- Aldawood AS, Sharaf MR, Collingwood CA. 2010. *Monomorium moathi* sp. n., a new ant species from Yemen (Hymenoptera: Formicidae) related to the *salomonis*-group. *Egyptian Academic Journal of Biological Sciences* **3** (2): 37–42.
- André E. 1881. Catalogue raisonné des Formicides provenant du voyage en Orient de M. Abeille de Perrin et description des espèces nouvelles. *Annales de la Société Entomologique de France* (6)1: 53–78.
- Assing V, Schülke M, Sharaf, MR, Aldawood AS. 2013. On the Staphylinidae of Saudi Arabia, with descriptions of two new species (Insecta: Coleoptera). *Linzer Biologische Beitrage* **45**(1):141–154.
- Barech G, Khaldi M, Espadaler X, Cagniant H. 2017. Le genre *Monomorium* (Hymenoptera, Formicidae) au Maghreb (Afrique du Nord): Clé d’identification, avec la redescription de la fourmi *Monomorium major* Bernard, 1953 et nouvelles citations pour l’Algérie. *Boletín de la Sociedad Entomológica Aragonesa* **61**: 151–157.

- 1047 Baroni Urbani C. 1964. Studi sulla mirmecofauna d'Italia. II. Formiche di Sicilia. *Atti della*
1048 *Accademia gioenia di scienze naturali in Catania* (6) 16: 25–66.
- 1049 Baroni Urbani, C. 1968. Studi sulla mirmecofauna d'Italia. V. *Aspetti Ecologici della Riviera del*
1050 *M. Cònero, Bolletino di zoologia* 35: 39-76.
- 1051 Bernard F. 1953. Les fourmis du Tassili des Ajjer. Pp. 121-250 in: Bernard, F. (ed.) 1953.
1052 Mission scientifique au Tassili des Ajjer (1949). Volume I. Recherches zoologiques et
1053 médicales. Paris: P. Lechevalier, 302 pp.
- 1054 Bernard F. 1968. Faune de l'Europe et du Bassin Méditerranéen. 3. Les fourmis (Hymenoptera
1055 Formicidae) d'Europe occidentale et septentrionale. Paris: Masson, 411 pp.
- 1056 Bolton B. 1987. A review of the *Solenopsis* genus-group and revision of Afrotropical
1057 *Monomorium* Mayr (Hymenoptera: Formicidae). *Bulletin of the British Museum (Natural*
1058 *History). Entomology* 54:263–452.
- 1059 Bolton B. 1994. *Identification Guide to the Ant Genera of the World*: Cambridge, Mass, 222 pp.
- 1060 Bolton B. 2020. *An online catalog of the ants of the world*. <http://antcat.org>. [accessed 13 July
1061 2020].
- 1062 Fisher BL., Bolton B. 2016. *Ants of Africa and Madagascar, A Guide to the Genera*. Berkeley:
1063 University of California Press, 503 pp.
- 1064 Borowiec L. 2014. Catalogue of ants of Europe, the Mediterranean Basin and adjacent regions
1065 (Hymenoptera: Formicidae). *Genus (Wroclaw)* 25 (1–2): 1–340
- 1066 Brandão, C. R. F. 2000. Major regional and type collections of ants (Formicidae) of the world
1067 and sources for the identification of ant species. Pp. 172-185 in: Agosti, D.; Majer, J. D.;
1068 Alonso, L. E.; Schultz, T. R. (eds.) 2000. *Ants. Standard methods for measuring and*
1069 *monitoring biodiversity*. Washington: Smithsonian Institution Press, xix + 280 pp.
- 1070 Brown WL Jr. 1958. A review of the ants of New Zealand. *Acta Hymenopterologica* 1:1–50.
- 1071 Brown Jr WL. 2000. Diversity of ants. In: Agosti D, Majer J, Alonso E, Schultz TR, eds. *Ants.*
1072 *Standard methods for measuring and monitoring biodiversity*. Biological diversity
1073 handbook series, Washington, D.C.: Smithsonian Institution Press, 45–79, 280 pp.
- 1074 Collingwood CA. 1978. A provisional list of Iberian Formicidae with a key to the worker caste
1075 (Hym. Aculeata). *EOS. Revista Española de Entomología* 52:65–95.
- 1076 Collingwood CA. 1985. Hymenoptera: Fam. Formicidae of Saudi Arabia. *Fauna of Saudi Arabia*
1077 7:230–301.

- 1078 Collingwood CA., Agosti D. 1996. Formicidae (Insecta: Hymenoptera) of Saudi Arabia (part 2).
1079 *Fauna Saudi Arabia* **15**:300–385.
- 1080 **z**
- 1081 Collingwood C.A, Pohl H, Guesten R, Wranik W, Van Harten A. 2004. The ants (Insecta:
1082 Hymenoptera: Formicidae) of the Socotra Archipelago. *Fauna of Arabia* **20**:473–495.
- 1083 Collingwood CA, Agosti D, Sharaf MR, Van Harten A. 2011. Order Hymenoptera, family
1084 Formicidae. *Arthropod Fauna of the UAE* **4**:405–474.
- 1085 Crawley WC. 1920. A gynandromorph of *Monomorium floricola*, Jerd. *Entomologist's Record*
1086 *and Journal of Variation* **32**:217–218.
- 1087 Deyrup M, Davis L, Cover S. 2000. Exotic ants in Florida. Transactions of the American
1088 Entomological Society **126**:293–325.
- 1089 Dlussky GM, Soyunov OS, Zabelin SI. 1990. *Ants of Turkmenistan*. [In Russian.]. Ashkabad:
1090 Ylym Press, 273 pp.
- 1091 Donisthorpe H. 1914. Myrmecophilous notes for 1913. *Entomologist's Record and Journal of*
1092 *Variation* **26**:37–45.
- 1093 Donisthorpe H. 1947. Some new ants from New Guinea. *Annals and Magazine of Natural*
1094 *History* (**11**)14:183–197.
- 1095 DuBois MB. 1986. A revision of the native New World species of the ant genus *Monomorium*
1096 (*minimum* group) (Hymenoptera: Formicidae). *University of Kansas Science Bulletin*
1097 **53**:65–119.
- 1098 El-Hawagry MS, Abdel-Dayem MS, Elgharbawy AA, Al Dhafer HM. 2016. A preliminary
1099 account of the fly fauna in Jabal Shada Al-A'la Nature Reserve, Saudi Arabia, with new
1100 records and biogeographical remarks (Diptera, Insecta). *ZooKeys* **636**: 107–139. <https://doi.org/10.3897/zookeys.636.9905>
1101
- 1102 Emery C. 1877. Catalogo delle formiche esistenti nelle collezioni del Museo Civico di Genova.
1103 Parte prima. Formiche provenienti dal Viaggio dei signori Antinori, Beccari e Issel nel
1104 Mar Rosso e nel paese dei Bogos. [part]. *Annali del Museo Civico di Storia Naturale*
1105 **9**:363–368.
- 1106 Emery C. 1881. Viaggio ad Assab nel Mar Rosso dei Signori G. Doria ed O. Beccari con il R.
1107 Avviso "Esploratore" dal 16 novembre 1879 al 26 febbraio 1880. I. Formiche. *Annali*
1108 *del Museo Civico di Storia Naturale* **16**:525–535.

- 1109 Emery C. 1893. Voyage de M. E. Simon à l'île de Ceylan (janvier-février 1892). Formicides.
1110 *Annales de la Société Entomologique de France* **62**:239–258.
- 1111 Emery C. 1894. Studi sulle formiche della fauna neotropica. VI-XVI. *Bullettino della Società*
1112 *Entomologica Italiana* **26**:137–241.
- 1113 Emery C. 1915. Su due formiche della Tripolitania. Bollettino del Laboratorio di Zoologia
1114 Generale e Agraria della Reale Scuola Superiore d'Agricoltura. *Portici* **9**:378.
- 1115 Emery C. 1922. Hymenoptera. Fam. Formicidae. Subfam. Myrmicinae. [part]. *Genera*
1116 *Insectorum* **174B**:95–206.
- 1117 Ettershank G. 1966. A generic revision of the world Myrmicinae related to *Solenopsis* and
1118 *Pheidologeton*. *Australian Journal of Zoology* **14**: 73–171.
- 1119 Evenhuis N. 2020. *The insect and spider collections of the world website*.
1120 <http://hbs.bishopmuseum.org/codens/> [accessed 10 Augst 2020].
- 1121 Fabricius JC. 1793. Entomologia systematica emendata et aucta. Secundum classes, ordines,
1122 genera, species, adjectis synonymis, locis observationibus, descriptionibus. Tome 2.
1123 Hafniae [= Copenhagen]: C. G. Proft, 519 pp.
- 1124 Fernández F, Serna FJ. 2019. Subfamilia Myrmicinae. Fernández, F.; Guerrero, R. J.; Delsinne,
1125 T. (eds.) 2019d. *Hormigas de Colombia. Bogotá*: Universidad Nacional de Colombia,
1126 1198 pp.
- 1127 Finzi B. 1936. Risultati scientifici della spedizione di S. A.S. il Principe Alessandro della Torre e
1128 Tasso nell'Egitto e penisola del Sinai. XI. Formiche. *Bulletin. Société Entomologique*
1129 *d'Egypte* **20**:155–210.
- 1130 Fisher BL, Bolton B. 2016. *Ants of Africa and Madagascar: A Guide to the Genera*. Berkeley:
1131 University of California Press, i–ix, 1–503 pp.
- 1132 Forel A. 1893. Formicides de l'Antille St. Vincent, récoltées par Mons. H. H.
1133 Smith. Transactions of the Entomological Society of London 1893:333–418.
- 1134 Forel A. 1907. Formicides du Musée nationale Hongroise. *Annales historico-naturales Musei*
1135 *nationalis hungarici* **5**: 1–42.
- 1136 Forel A. 1910. Glanures myrmécologiques. *Annales de la Société Entomologique de Belgique*
1137 **54**:6–32.

- 1138 Forel A. 1913. Ameisen aus Rhodesia, Kapland usw. (Hym.) gesammelt von Herrn G. Arnold,
1139 Dr. H. Brauns und Anderen. *Deutsche Entomologische Zeitschrift* 1913(Suppl.):203–
1140 225.
- 1141 Forel A. 1916. Fourmis du Congo et d'autres provenances récoltées par MM. Hermann Kohl,
1142 Luja, Mayné, etc. *Revue Suisse de Zoologie* **24**:397–460.
- 1143 Hamann HHF, Klemm W. 1967. Ergebnisse der zoologischen Nubien-Expedition 1962. Teil
1144 XXXIV. Hymenoptera - Formicidae. *Annalen des Naturhistorischen Museums in Wien*
1145 **70**:411–421.
- 1146 Heterick BE. 2001. Revision of the Australian ants of the genus *Monomorium* (Hymenoptera:
1147 Formicidae). *Invertebrate Taxonomy* **15**:353–459.
- 1148 Heterick BE. 2003. Two new Australian *Monomorium* Mayr (Hymenoptera: Formicidae),
1149 including a highly distinctive species. *Australian Journal of Entomology* **42**:249–253.
- 1150 Heterick BE. 2006. A revision of the Malagasy ants belonging to genus *Monomorium* Mayr,
1151 1855. *Proceedings of the California Academy of Sciences* **57**:69–202.
- 1152 Heterick BE. 2009. A guide to the ants of south-western Australia. *Records of the Western*
1153 *Australian Museum Supplement* **76**:1–206.
- 1154 Hita Garcia F, Wiesel E, Fischer G. 2013. The ants of Kenya (Hymenoptera: Formicidae)—
1155 faunal overview, first species checklist, bibliography, accounts for all genera, and
1156 discussion on taxonomy and zoogeography. *Journal of East African Natural History*
1157 **101**:127–222
- 1158 Hlaváč P., Sharaf, MR, Aldawood AS. 2013. New species and record of of Pselaphinae
1159 (Coleoptera: Staphylinidae) from Saudi Arabia. *Zootaxa* **3666** (3): 331–336,
1160 <http://dx.doi.org/10.11646/zootaxa.3666.3.3>
- 1161 Holt BG, Lessard JP, Borregaard MK, Fritz SA, Araújo MB, Dimitrov D, Fabre PH, Graham
1162 CH, Graves GR, Jønsson KA, Nogués-Bravo D, Wang Z, Whittaker RJ, Fjeldså J,
1163 Rahbek C. 2013. An update of Wallace's zoogeographic regions of the world. *Science*.
1164 **339**(6115):74–8. doi: 10.1126/science.1228282. pmid:23258408
- 1165 Jerdon TC. 1851. A catalogue of the species of ants found in Southern India. *Madras Journal of*
1166 *Literature and Science* **17**:103–127.
- 1167 Karavaiev V. 1911. Ameisen aus Aegypten und dem Sudan. *Russkoe Entomologicheskoe*
1168 *Obozrenie* **11**:1–12.

- 1169 Kempf WW. 1972. Catálogo abreviado das formigas da região Neotropical. *Studia Entomologica*
1170 **15**:3–344.
- 1171 Kiran K, Karaman C. 2020. Additions to the Ant Fauna of Turkey (Hymenoptera, Formicidae).
1172 *Zoosystema* **42**:285–329.
- 1173 Kusnezov, N. 1949. El género Monomorium (Hymenoptera, Formicidae) en la Argentina. *Acta*
1174 *Zoologica Lilloana* 7:423-448.
- 1175 Linnaeus C. 1758. *Systema naturae per regna tria naturae, secundum classes, ordines, genera,*
1176 *species, cum characteribus, differentiis, synonymis, locis.* Tomus I. Editio decima,
1177 reformata. Holmiae [= Stockholm]: L. Salvii, 824 pp.
- 1178 Linsley EG, Usinger RL. 1966. Insects of the Galápagos Islands. *Proceedings of the California*
1179 *Academy of Sciences* (4)33:113–196.
- 1180 Mackay W, Mackay E. 2002. *The ants of New Mexico (Hymenoptera: Formicidae).* Lewiston,
1181 New York: Edwin Mellen Press, 400 pp.
- 1182 Mahnert V, Sharaf MR, Aldawood AS. 2014. Further records of Pseudoscorpions (Arachnida,
1183 Pseudoscorpions) from Saudi Arabia. *Zootaxa* **3764** (3): 387–393,
1184 <https://doi.org/10.11646/zootaxa.3764.3.8>
- 1185 Mayr G. 1862. Myrmecologische Studien. *Verhandlungen der Kaiserlich-Königlichen*
1186 *Zoologisch-Botanischen Gesellschaft in Wien* **12**:649–776.
- 1187 Mayr G. 1866. Myrmecologische Beiträge. *Sitzungsberichte der Kaiserlichen Akademie der*
1188 *Wissenschaften in Wien. Mathematisch-Naturwissenschaftliche Classe. Abteilung I*
1189 **53**:484–517.
- 1190 Mayr G. 1877. Formicidae. [In Russian.]. In: Fedchenko, A. P. 1877. Travels in Turkestan. Vol.
1191 2, Div. 5, No. 7. [In Russian.]. *Izvestiya Imperatorskago Obshestva Lyubitelei*
1192 *Estestvoznaniya Antropologii i Etnografii pri Imperatorskom Moskovskom*
1193 *Universitete* **26**:i-iii, 1-20 (+1).
- 1194 Mayr G. 1879. Beiträge zur Ameisen-Fauna Asiens. *Verhandlungen der Kaiserlich-Königlichen*
1195 *Zoologisch-Botanischen Gesellschaft in Wien* **28**:645–686.
- 1196 Mayr G. 1901. Südafrikanische Formiciden, gesammelt von Dr. Hans Brauns. *Annalen des*
1197 *Kaiserlich-Königlichen Naturhistorischen Museums in Wien* **16**:1–30.
- 1198 Menozzi C. 1932. Missione scientifica del Prof. E. Zavattari nel Fezzan (1931). Hymenoptera-
1199 Formicidae. *Bollettino della Società Entomologica Italiana* **64**:93–95.

- 1200 Monks J, Ross S, Geiser M, De Prins J, Sharaf MR, Wyatt N, Al Rijeibi S & Polaszek A. 2019.
1201 A preliminary survey of the insect fauna of the Hajar Mountain Range, Oman. *Journal of*
1202 *Natural History* **53**: 939–963, <https://doi.org/10.1080/00222933.2019.1611969>
- 1203 Morisita M, Kubota M, Onoyama K, Ogata K, Terayama M, Yamauchi K, Sonobe R, Yamane
1204 S, Kondoh M, Imai HT. 1992. *A guide for the identification of Japanese ants. III.*
1205 *Myrmicinae and supplement to Leptanillinae. (Hymenoptera: Formicidae).* [In
1206 Japanese.]. Tokyo: Myrmecological Society of Japan, 94 pp.
- 1207 Paknia O, Radchenko A, Alipanah H, Pfeiffer M.. 2008. A preliminary checklist of the ants
1208 (Hymenoptera: Formicidae) of Iran. *Myrmecological News* **11**: 151-159.
- 1209 Roger J. 1863. Die neu aufgeführten Gattungen und Arten meines Formiciden-Verzeichnisses
1210 nebst Ergänzung einiger früher gegebenen Beschreibungen. *Berliner Entomologische*
1211 *Zeitschrift* **7**:131–214.
- 1212 Santschi F. 1911. Formicides nouveaux de l'Afrique Mineure (4e note suite). *Bulletin de la*
1213 *Société d'Histoire Naturelle de l'Afrique du Nord* **2**:78–85.
- 1214 Santschi F. 1912. Quelques nouvelles variétés de fourmis africaines. *Bulletin de la Société*
1215 *d'Histoire Naturelle de l'Afrique du Nord* **3**:147–149.
- 1216 Santschi F. 1914. Formicides de l'Afrique occidentale et australe du voyage de Mr. le Professeur
1217 F. Silvestri. *Bollettino del Laboratorio di Zoologia Generale e Agraria della Reale*
1218 *Scuola Superiore d'Agricoltura Portici* **8**:309–385.
- 1219 Santschi F. 1915a. Nouvelles fourmis d'Algérie, Tunisie et Syrie. *Bulletin de la Société*
1220 *d'Histoire Naturelle de l'Afrique du Nord* **6**:54–63.
- 1221 Santschi F. 1915b. Nouvelles fourmis d'Afrique. *Annales de la Société Entomologique de France*
1222 **84**:244–282.
- 1223 Santschi F. 1919. Fourmis nouvelles éthiopiennes. *Revue Zoologique Africaine* (Brussels) **6**:229–
1224 240.
- 1225 Santschi F. 1926. Description de nouveaux Formicides éthiopiens (III^{me} partie). *Revue*
1226 *Zoologique Africaine* (Brussels) **13**:207–267.
- 1227 Santschi F. 1927. Révision myrmécologique. *Bulletin et Annales de la Société Entomologique de*
1228 *Belgique* **67**:240–248.
- 1229 Santschi F. 1934. Mission J. de Lépiney au Soudan Français 1933–1934. (Huitième note.)
1230 Fourmis. *Bulletin de la Société des Sciences Naturelles du Maroc* **14**:33–34.

- 1231 Santschi F. 1936. Étude sur les fourmis du genre *Monomorium* Mayr. *Bulletin de la Société des*
1232 *Sciences Naturelles du Maroc* **16**:32–64.
- 1233 Santschi F. 1938. Quelques nouvelles fourmis d'Egypte. *Bulletin Société Entomologique*
1234 *d'Egypte* **21**:28–44.
- 1235 Sarnat EM, Economo EP. 2012. *The ants of Fiji*. University of California Publications in
1236 Entomology, Volume 132. Berkeley: University of California Press, xiii + 384 pp.
- 1237 Sharaf MR. 2006. *Taxonomic and ecological studies on family Formicidae (Order:*
1238 *Hymenoptera) in Egypt including some protectorates with a study of some insect fauna*
1239 *associated with ant species* [unpublished thesis]. Cairo: Ain Shams University, Faculty of
1240 Science, Entomology Department; 340 pp.
- 1241 Sharaf MR, Aldawood AS. 2013a. First occurrence of the *Monomorium hildebrandti*-group
1242 (Hymenoptera: Formicidae), in the Arabian Peninsula, with description of a new
1243 species *M. kondratieffi* n. sp. *Proceedings of the Entomological Society of Washington*
1244 **115** (1):75–84.
- 1245 Sharaf MR, Aldawood AS. 2013b. *Monomorium sarawatensis* Sharaf & Aldawood, sp. n. Pp.
1246 70–73 in: El-Hawagry, M. S.; Khalil, M. W.; Sharaf, M.R.; Fadl, H. H.; Aldawood,
1247 A.S. 2013. A preliminary study on the insect fauna of Al-Baha Province, Saudi Arabia,
1248 with descriptions of two new species. *ZooKeys* **274**:1–88.
- 1249 Sharaf MR, Aldawood AS. 2013c. The ant genus *Carebara* Westwood in the Arabian Peninsula
1250 (Hymenoptera, Formicidae). *ZooKeys* **357**:67–83. doi:10.3897/zookeys.357.5946
- 1251 Sharaf MR, Al Dhafer HM, Aldawood AS. 2014a. First record of the myrmicine ant genus
1252 *Meranoplus* Smith, 1853 (Hymenoptera: Formicidae) from the Arabian Peninsula with
1253 description of a new species and notes on the zoogeography of southwestern Kingdom
1254 Saudi Arabia. *PLoS ONE* **9**(11):e111298. doi:10.1371/journal.pone.0111298
- 1255 Sharaf MR, Fisher BL, Aldawood AS. 2014b. Notes on ants of the genus *Strumigenys* F. Smith,
1256 1860 (Hymenoptera: Formicidae) in the Arabian Peninsula, with a key to species.
1257 *Sociobiology* **61**(3):293–299. doi:10.13102/sociobiology.v61i3.293-299.
- 1258 Sharaf MR, Collingwood CA, Al Dhafer HM, Al mutairi MS, Aldawood AS.2015. New
1259 synonyms of two Arabian ants of the genus *Monomorium* Mayr, 1855 (Hymenoptera,
1260 Formicidae). *ZooKeys* **505**: 51–58. doi: 10.3897/ zookeys.505.9441

- 1261 Sharaf MR, Fisher BL, Collingwood CA, Aldawood, A.S. 2017a. Ant fauna (Hymenoptera:
1262 Formicidae) of the Socotra Archipelago (Yemen): zoogeography, distribution and
1263 description of a new species. *Journal of Natural History* **51** (5-6):317–378.
1264 10.1080/00222933.2016.1271157
- 1265 Sharaf MR, Akbar, SA, Al Dhafer HM, El-Gharbawy A, Aldawood AS. 2017b. Taxonomy of
1266 the Myrmicine ant genus *Temnothorax* Mayr, 1861 (Formicidae: Myrmicinae) in the
1267 Arabian Peninsula. *European Journal of Taxonomy* **280**:1–17. doi:10.5852/ejt.2017.280
- 1268 Sharaf MR, Al Dhafer HM, Aldawood AS, Hita Garcia F. 2018a. Ants of the *Monomorium*
1269 *monomorium* species-group (Hymenoptera: Formicidae) in the Arabian Peninsula with
1270 description of a new species from southwestern Saudi Arabia. *PeerJ* **6**:e4277.
1271 10.7717/peerj.4277
- 1272 Sharaf MR, Fisher BL, Al Dhafer HM, Polaszek A, Aldawood AS. 2018b. Additions to the ant
1273 fauna (Hymenoptera: Formicidae) of Oman: an updated list, new records and a
1274 description of two new species. *Asian Myrmecology* **10**:e010004. 1–38.
1275 doi:10.20362/am.010004
- 1276 Sharaf MR Aldawood AS.2019. Review of the ant genus *Meranoplus* Smith, 1853
1277 (Hymenoptera: Formicidae) in the Arabian Peninsula with description of a new species
1278 *M. mosalahi* sp. n. from Oman. *PeerJ* **7**:e6287. doi:10.7717/peerj.6287
- 1279 Sharaf MR, Aldawood AS, Hita Garcia F. 2019. Review of the Arabian *Crematogaster* Lund
1280 (Hymenoptera, Formicidae), synoptic list, distribution, and description of two new
1281 species from Oman and Saudi Arabia. *ZooKeys* **898**:27–81,
1282 doi:10.3897/zookeys.898.37531
- 1283 Sharaf MR, Aldawood AS, Mohamed, AA, Hita Garcia F. 2020a. The genus *Lepisiota* Santschi,
1284 1926 of the Arabian Peninsula with the description of a new species, *Lepisiota elbazi* sp.
1285 nov. from Oman, an updated species identification key, and assessment of zoogeographic
1286 affinities. *Journal of Hymenoptera Research* **76**: 127–152, doi:10.3897/jhr.76.50193
- 1287 Sharaf MR, Mohamed AA, Al Dhafer HM, Aldawood AS. 2020b. *Nesomyrmex micheleae*, a
1288 new ant species (Hymenoptera: Formicidae) from the Dhofar Governorate, Oman, with a
1289 synoptic list, distribution, and a key to the Arabian *Nesomyrmex*. *Journal of Natural*
1290 *History* **54**: 351–365. <https://doi.org/10.1080/00222933.2020.1762013>

- 1291 Shuckard WE. 1838. Description of a new species of *Myrmica* which has been found in houses
1292 both in the Metropolis and Provinces. *Magazine of Natural History* (2)2:626–627.
- 1293 Smith F. 1857. Catalogue of the hymenopterous insects collected at Sarawak, Borneo; Mount
1294 Ophir, Malacca; and at Singapore, by A. R. Wallace. [part]. *Journal and Proceedings*
1295 *of the Linnean Society of London. Zoology* 2:42–88.
- 1296 Smith F. 1858. *Catalogue of hymenopterous insects in the collection of the British Museum*. Part
1297 VI. Formicidae. London: British Museum, 216 pp.
- 1298 Smith F. 1876. Preliminary notice of new species of Hymenoptera, Diptera, and Forficulidae
1299 collected in the island of Rodriguez by the naturalists accompanying the Transit-of-
1300 Venus expedition. *Annals and Magazine of Natural History* (4)17:447–451.
- 1301 Sparks KS, Andersen AN, Austin AD. 2014. Systematics of the *Monomorium rothsteini* Forel
1302 species complex (Hymenoptera: Formicidae), a problematic ant group in
1303 Australia. *Zootaxa* 3893 (4):489–529.
- 1304 Sparks KS., Andersen AN, Austin AD. 2019. A multi-gene phylogeny of Australian
1305 *Monomorium* Mayr (Hymenoptera: Formicidae) results in reinterpretation of the genus
1306 and resurrection of *Chelaner* Emery. *Invertebrate Systematics* 33: 225–236.
- 1307 Stitz H. 1917. Ameisen aus dem westlichen Mittelmeergebiet und von den Kanarischen Inseln.
1308 *Mitteilungen aus dem Zoologischen Museum in Berlin* 8:333–353.
- 1309 Stitz H. 1932. Formicidae [of the Wollebaek Galapagos Expedition]. *Nyt Magazin for*
1310 *Naturvidenskaberne* 71:367-372.
- 1311 SWA 2018. *Protected Areas. Saudi Wildlife Commission*. <https://www.swa.gov.sa/en> [accessed
1312 13 July 2020].
- 1313 SWA 2020. *Protected Areas. Saudi Wildlife Commission*. <https://www.swa.gov.sa/en> [26-8-
1314 2020].
- 1315 Terayama M. 2009. A synopsis of the family Formicidae of Taiwan (Insecta:
1316 Hymenoptera). *Research Bulletin of Kanto Gakuen University. Liberal Arts* 17:81–266.
- 1317 Thomas J, El-Sheikh MA, Alatar AA. 2017. Endemics and endangered species in the
1318 biodiversity hotspot of the Shada Mountains, Saudi Arabia. *Journal of Arid Land* 9(1):
1319 109–21. doi: 10.1007/s40333-016-0025-8
- 1320 Vigna Taglianti A, Audisio PA, Biondim, Bologna MA, Carpaneto GM, De Biase A, Fattorini S,
1321 Piattella E, Sindaco R, Venchi A, Zapparolim, 1999. A proposal for a chorotype

- classification of the Near East fauna, in the framework of the Western Palaearctic region.
- Biogeographia* **20**: 31–59.
- Vonshak M, Ionescu-Hirsch A. 2009. A checklist of the ants of Israel (Hymenoptera: Formici-
- dae). *Israel Journal of Entomology* **39**: 33–55.
- Walker F. 1871. *A list of Hymenoptera collected by J. K. Lord, Esq. in Egypt, in the*
- neighbourhood of the Red Sea, and in Arabia, with descriptions of the new species.*
- London: E. W. Janson, vi + 59 pp.
- Ward PS, Brady SG, Fisher BL, Schultz TR. 2015. The evolution of myrmicine ants: phylogeny
- and biogeography of a hyperdiverse ant clade (Hymenoptera: Formicidae). *Systematic*
- Entomology* **40**:61–81.
- Wetterer JK. 2010a. Worldwide spread of the flower ant, *Monomorium floricola* (Hymenoptera:
- Formicidae). *Myrmecological News* **13**:19–27.
- Wetterer JK. 2010b. Worldwide spread of the pharaoh ant, *Monomorium pharaonis*
- (Hymenoptera: Formicidae). *Myrmecological News* **13**:115–129
- Wheeler WM. 1905. The ants of the Bahamas, with a list of the known West Indian
- species. *Bulletin of the American Museum of Natural History* **21**:79–135.
- Wheeler WM. 1913. The ants of Cuba. *Bulletin of the Museum of Comparative Zoology* **54**:477–
- 505.
- Wheeler WM. 1922. Ants of the American Museum Congo expedition. New York: *Bulletin of*
- the American Museum of Natural History* **1139**.
- Wheeler GC, Wheeler J. 1955. The ant larvae of the myrmicine tribe Solenopsidini. *American*
- Midland Naturalist* **54**:119–141.
- Wheeler WM, Mann WM. 1916. The ants of the Phillips Expedition to Palestine during 1914.
- Bulletin of the Museum of Comparative Zoology* **60**:167–174.
- Wilson EO, Taylor RW. 1967b. The ants of Polynesia (Hymenoptera: Formicidae). *Pacific*
- Insects Monograph* **14**: 1–109.
- Wu J, Wang C. 1995. *The ants of China*. [In Chinese.]. Beijing: China Forestry Publishing
- House, x + 214 pp.

Figure Legends.

Figure 1. A: head of *M. exiguum* in full-face view, CASENT0217367 (Erin Prado); B: head of *M. khalidi*, **sp. n.** in full-face view, CASENT0922288 (Michele Esposito); C: head of *M. carbonarium* in full-face view, CASENT0902279 (Ryan Perry); D: head of *M. floricola* in full-face view, CASENT0922876 (Michele Esposito); E: body of *M. carbonarium* in profile, CASENT0902279 (Ryan Perry); F: body of *M. sarawatense* in profile, CASENT0280971 (Estella Ortega), www.AntWeb.org.

Figure 2. A: body of *M. floricola* in profile, CASENT0922876 (Michele Esposito); B: body of *M. holothir* in profile, CASENT0902243 (Will Ericson); C: body of *M. aeyade* in profile, CASENT0922329 (Michele Esposito); D: body of *M. exiguum* in profile, CASENT0217367 (Erin Prado); E: body of *M. mohammedi* in profile, CASENT0922351 (Michele Esposito); www.AntWeb.org, F: body of *M. clavicorne* in profile, (Francisco Hita Garcia).

Figure 3. A: head of *M. barbatulum* in profile, CASENT0922263 (Michele Esposito); B: mesosoma of *M. rabirium* in profile, CASENT0746641 (Zach Lieberman); C: mesosoma of *M. elghazalyi* in profile, CASENT0746626 (Michele Esposito); D: petiole and postpetiole of *M. harithe* in profile, CASENT0913802 (Will Ericson); E: head of *M. elghazalyi* in full-face view, CASENT0746626 (Michele Esposito); F head of *M. dirie* in full-face view, CASENT0913571 (Alexandra Westrich), www.AntWeb.org.

Figure 4. A: head of *M. dirie* in profile, CASENT0913571 (Alexandra Westrich); B: head of *M. salomonis* in profile, CASENT0913835 (Will Ericson); C: mesosoma of *M. subdenticorne* in profile, CASENT0914318 (Zach Lieberman); D: mesosoma of *M. bicolor* in profile, CASENT0073615 (Michele Esposito); E: petiole of *M. acutinode* in profile, CASENT0913547 (Will Ericson); F petiole of *M. carbo* in profile, CASENT0249908 (Shannon Hartman), www.AntWeb.org.

Figure 5. A: mesosoma of *M. areniphilum* in profile, CASENT0048600 (Michele Esposito); B: mesosoma of *M. subopacum* in profile, CASENT0064820 (April Nobile); C: mesosoma of *M. luteum* in profile, CASENT0904599 (Will Ericson); D: mesosoma of *M. niloticum* in profile,

CASENT0260164 (Estella Ortega); E: head of *M. niloticum* in full-face view, CASENT0919811 (Michele Esposito); F head of *M. riyadhe* in full-face view, CASENT0922342 (Michele Esposito), www.AntWeb.org.

Figure 6. A: head of *M. hanaqe* in full-face view, CASENT0249834 (Ryan Perry); B: mesosoma of *M. hanaqe* in profile, CASENT0249834 (Ryan Perry); C: head of *M. jizane* in full-face view, CASENT0913806 (Will Ericson); D: propodeum of *M. jizane* in profile, CASENT0913806 (Will Ericson); E: head of *M. fayfaense* in full-face view, CASENT0249833 (Ryan Perry); F gaster of *M. fayfaense* in profile, CASENT0249833 (Ryan Perry), www.AntWeb.org.

Figure 7. A: head of *M. knappi* in full-face view, CASENT0913812 (Will Ericson); B: body of *M. knappi* in profile, CASENT0913812 (Will Ericson); C: body of *M. nitidiventre* in profile, CASENT0904602 (Will Ericson); D: head of *M. nitidiventre* in full-face view, CASENT0904602 (Will Ericson); E: body of *M. khalidi* **sp. n.** in profile, CASENT0922288 (Michele Esposito); F head of *M. khalidi* **sp. n.** in profile, CASENT0922288 (Michele Esposito), www.AntWeb.org.

Figure 8. A: mesosoma of *M. pharaonis* in profile, CASENT0246072 (Andrea Walker); B: mesosoma of *M. buxtoni* in profile, CASENT0902220 (Will Ericson); C: head of *M. suleyile* in full-face view, CASENT0913854 (Zach Lieberman); D: head of *M. mahyoubi* in full-face view, CASENT0913823 (Alexandra Westrich); E: head of *M. tumaire* in profile, CASENT0249858 (Ryan Perry); F head of *M. buettikeri* in profile, CASENT0913565 (Zach Lieberman), www.AntWeb.org.

Figure 9. A: head of *M. gallagheri* in full-face view, CASENT0913582 (Zach Lieberman); B: head of *M. buxtoni* in full-face view, CASENT0902220 (Zach Lieberman); C: body of *M. abeillei* in profile, CASENT0915411 (Will Ericson); D: body of *M. asiriense* in profile, CASENT0913560 (Zach Lieberman), www.AntWeb.org.

Figure 10. *M. abeillei*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0915411 (Will Ericson), www.AntWeb.org.

1413

1414 Figure 11. *M. areniphilum*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1415 CASENT0048600 (Michele Esposito), www.AntWeb.org.

1416

1417 Figure 12. *M. barbatulum*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1418 CASENT0922263 (Michele Esposito), www.AntWeb.org.

1419

1420 Figure 13. *M. bicolor*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1421 CASENT0904601 (Will Ericson), www.AntWeb.org.

1422

1423 Figure 14. *M. floricola*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1424 CASENT0922876 (Michele Esposito), www.AntWeb.org.

1425

1426 Figure 15. *M. harithe*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1427 CASENT0913802 (Will Ericson), www.AntWeb.org.

1428

1429 Figure 16. *M. heggyi* sp. n., A: body in profile; B: body in dorsal view); C: head in full-face
1430 view, CASENT0746641 (Zach Lieberman), www.AntWeb.org.

1431

1432 Figure 17. Shada Al A'la, the type locality of *M. heggyi* **sp. n.** (A. Shams Al Ola).

1433

1434 Figure 18. *M. khalidi* **sp. n.**, A: body in profile; B: body in dorsal view); C: head in full-face
1435 view, CASENT0922288 (Michele Esposito), www.AntWeb.org.

1436

1437 Figure 19. Shada Al A'la, the type locality of *M. khalidi* **sp. n.** (A. Shams Al Ola).

1438

1439 Figure 20. *M. niloticum*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1440 CASENT0905755 (Will Ericson), www.AntWeb.org.

1441

1442 Figure 21. *M. nitidiventre*, A: body in profile; B: body in dorsal view); C: head in full-face view,
1443 CASENT0904602 (Will Ericson), www.AntWeb.org.

1444

1445 Figure 22. *M. subdenticorne*, A: body in profile; B: body in dorsal view); C: head in full-face
1446 view, CASENT0914318 (Zach Lieberman), www.AntWeb.org.

1447

1448 Figure 23. A: *M. venustum*, A: body in profile; B: body in dorsal view); C: head in full-face
1449 view, CASENT0902221 (Will Ericson), www.AntWeb.org.

1450

1451 Figure 24. Distribution map of *M. abeillei*.

1452

1453 Figure 25. Distribution map of *M. barbatulum*, *M. bicolor*, *M. floricola*.

1454

1455 Figure 26. Distribution map of *M. harithe*, *M. khalidi* **sp. n.**, *M. nitidiventre*, *M. hegyi* **sp. n.**

1456

1457 Figure 27. Distribution map of *M. niloticum*.

1458

1459 Figure 28. Distribution map of *M. subdenticorne*.

1460

1461 Figure 29. Distribution map of *M. venustum*.

Figure 1

Figure 1

Figure 1. A: head of *M. exiguum* in full-face view, CASENT0217367 (Erin Prado); B: head of *M. khalidi*, **sp. n.** in full-face view, CASENT0922288 (Michele Esposito); C: head of *M. carbonarium* in full-face view, CASENT0902279 (Ryan Perry); D: head of *M. floricola* in full-face view, CASENT0922876 (Michele Esposito); E: body of *M. carbonarium* in profile, CASENT0902279 (Ryan Perry); F: body of *M. sarawatense* in profile, CASENT0280971 (Estella Ortega), www.AntWeb.org.




Figure 2

A: body of *M. floricola* in profile, CASENT0922876 (Michele Esposito); B: body of *M. holothir* in profile, CASENT0902243 (Will Ericson); C: body of *M. aeyade* in profile, CASENT0922329 (Michele Esposito); D: body of *M. exiguum* i

Key illustrations




Figure 3

Fig3.jpg

Key illustrations




Figure 4

A: head of *M. dirie* in profile, CASENT0913571 (Alexandra Westrich); B: head of *M. salomonis* in profile, CASENT0913835 (Will Ericson); C: mesosoma of *M. subdenticorne* in profile, CASENT0914318 (Zach Lieberman); D: mesosoma of [i]M. bicolo

Key illustrations




Figure 5

A: mesosoma of *M. areniphilum* in profile, CASENT0048600 (Michele Esposito); B: mesosoma of *M. subopacum* in profile, CASENT0064820 (April Nobile); C: mesosoma of *M. luteum* in profile, CASENT0904599 (Will Ericson); D: mesosoma of [i]M. nil

Key illustrations




Figure 6

A: head of *M. hanage* in full-face view, CASENT0249834 (Ryan Perry); B: mesosoma of *M. hanage* in profile, CASENT0249834 (Ryan Perry); C: head of *M. jizane* in full-face view, CASENT0913806 (Will Ericson); D: propodeum of *M. jizane* in

Key illustrations




Figure 7

A: head of *M. knappi* in full-face view, CASENT0913812 (Will Ericson); B: body of *M. knappi* in profile, CASENT0913812 (Will Ericson); C: body of *M. nitidiventre* in profile, CASENT0904602 (Will Ericson); D: head of *M. nitidiventre* in

Key illustrations




Figure 8

A: mesosoma of *M. pharaonis* in profile, CASENT0246072 (Andrea Walker); B: mesosoma of *M. buxtoni* in profile, CASENT0902220 (Will Ericson); C: head of *M. suleyile* in full-face view, CASENT0913854 (Zach Lieberman); D: head of [i]M. mahyoub

Key illustrations




Figure 9

A: head of *M. gallagheri* in full-face view, CASENT0913582 (Zach Lieberman); B: head of *M. buxtoni* in full-face view, CASENT0902220 (Zach Lieberman); C: body of *M. abeillei* in profile, CASENT0915411 (Will Ericson); D: body of [i]M. asirie

Key illustrations




Figure 10

Fig10.jpg

M. abeillei, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0915411 (Will Ericson), www.AntWeb.org . Figure 11. *M. areniphilum*, A: body in profile; B: body in dorsal view); C: head in full-face view,




Figure 11

Fig11.jpg

Figure 11. *M. areniphilum*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0048600 (Michele Esposito), www.AntWeb.org .




Figure 12

Fig12.jpg

Figure 12. *M. barbatulum*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0922263 (Michele Esposito), www.AntWeb.org .




Figure 13

Fig13.jpg

Figure 13. *M. bicolor*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0904601 (Will Ericson), www.AntWeb.org .




Figure 14

Fig14.jpg

Figure 14. *M. floricola*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0922876 (Michele Esposito), www.AntWeb.org .




Figure 15

Fig15.jpg

Figure 15. *M. harithe*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0913802 (Will Ericson), www.AntWeb.org .




Figure 16

Fig16.jpg

Figure 16. *M. heggyi* sp. n., A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0746641 (Zach Lieberman), www.AntWeb.org .




Figure 17

Fig17.JPG

Figure 17. Shada Al A'la, the type locality of *M. hegyi* **sp. n.** (A. Shams Al Ola).



Figure 18

Fig18.jpg

Figure 18. *M. khalidi* **sp. n.**, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0922288 (Michele Esposito), www.AntWeb.org .




Figure 19

Fig19.jpg

Figure 19. Shada Al A'la, the type locality of *M. khalidi* **sp. n.** (A. Shams Al Ola).



Figure 20

Fig20.jpg

Figure 20. *M. niloticum*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0905755 (Will Ericson), www.AntWeb.org .




Figure 21

Fig21.jpg

Figure 21. *M. nitidiventre*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0904602 (Will Ericson), www.AntWeb.org .



Figure 22

Fig22.jpg

Figure 22. *M. subdenticorne*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0914318 (Zach Lieberman), www.AntWeb.org .




Figure 23

Fig23.jpg

Figure 23. A: *M. venustum*, A: body in profile; B: body in dorsal view); C: head in full-face view, CASENT0902221 (Will Ericson), www.AntWeb.org .




Figure 24

Fig24.png

Distribution map of *M. abeillei*.




Figure 25

Fig25.png

Distribution map of *M. barbatulum*, *M. bicolor*, *M. floricola*.



Figure 26

Fig26.png

Distribution map of *M. harithe*, *M. khalidi* **sp. n.**, *M. nitidiventre*, *M. heggyi* **sp. n.**




Figure 27

Fig27.png

Distribution map of *M. niloticum*.




Figure 28

Fig28.png

Distribution map of *M. subdenticorne*.




Figure 29

Fig29.png

Distribution map of *M. venustum*.




Table 1(on next page)

Table1.docx

Biogeography of the Arabian *Monomorium*. References: 1 = Collingwood 1985, 2 = Bolton, 1987, 3 = Collingwood & Agosti, 1996

Table 1. Biogeography of the Arabian *Monomorium*. References: 1 = Collingwood 1985, 2 = Bolton, 1987, 3 = Collingwood & Agosti, 1996

3

Species	Type locality	Bioregion	Reference	Distribution map*
<i>M. abeillei</i>	ISRAEL	Palearctic	1; 2; 3	Fig. 24
<i>M. acutinode</i>	OMAN	Palearctic	3	-
<i>M. aeyade</i>	OMAN	Endemic	3; Sharaf et al. 2018a	-
<i>M. areniphilum</i>	TUNISIA	Palearctic	1; 3	-
<i>M. asiriense</i>	KSA	Endemic	3	-
<i>M. barbatulum</i>	KAZAKHSTAN	Palearctic	1; 3	Fig. 25
<i>M. bicolor</i>	ERITREA	Afrotropic	1; 2; 3	Fig. 25
<i>M. brunneolucidulum</i>	OMAN	unknown	3	-
<i>M. buettikeri</i>	KUWAIT	Endemic	3	-
<i>M. buxtoni</i>	IRAQ	Palearctic	1; 3	-
<i>M. carbo</i>	ETHIOPIA	Afrotropic	1; 3	-
<i>M. carbonarium</i>	MADEIRA	Palearctic	3	-
<i>M. clavicorne</i>	ISRAEL	Palearctic	3; Sharaf et al. 2018a	-
<i>M. dammame</i>	KSA	Endemic	3	-
<i>M. dirie</i>	OMAN	Endemic	3	-
<i>M. elghazalyi</i>	YEMEN	Endemic	Sharaf et al., 2017a	-
<i>M. exiguum</i>	ETHIOPIA	Afrotropic	3; Sharaf et al. 2018a	-
<i>M. fayfaense</i>	KSA	Endemic	3	-
<i>M. floricola</i>	INDIA	Tramp	Heterick, 2006	Fig. 25
<i>M. gallagheri</i>	OMAN	Endemic	3	-
<i>M. hanaqe</i>	KSA	Endemic	3	-
<i>M. harithe</i>	KSA	Endemic	3	Fig. 26
<i>M. hegyi</i>	KSA	Endemic		Fig. 26
<i>M. holothir</i>	KENYA	Afrotropic	2	-

<i>M. jizane</i>	KSA	Endemic	3	-
<i>M. khalidi</i>	KSA	Endemic		Fig. 26
<i>M. knappi</i>	YEMEN	Endemic	3	-
<i>M. luteum</i>	YEMEN	Afrotropic	1; 3	-
<i>M. mahyoubi</i>	YEMEN	Endemic	3	-
<i>M. moathi</i>	YEMEN	Endemic	Sharaf & Collingwood, 2010	-
<i>M. mohammedi</i>	KSA	Endemic	3; Sharaf et al. 2018a	-
<i>M. niloticum</i>	EGYPT	Palearctic	1; 3	Fig. 27
<i>M. nimihil</i>	YEMEN	Endemic	Collingwood et al. 2004	-
<i>M. nitidiventre</i>	EGYPT	Palearctic	1; 3	Fig. 26
<i>M. pharaonis</i>	EGYPT	Tramp	1; 3	-
<i>M. rimae</i>	YEMEN	Endemic	3	-
<i>M. riyadhe</i>	KSA	Endemic	3	-
<i>M. salomonis</i>	EGYPT	Palearctic	1; 3	-
<i>M. sarawatense</i>	KSA	Endemic	3; Sharaf et al. 2018a	-
<i>M. subdenticorne</i>	YEMEN	Endemic	3	Fig. 28
<i>M. subopacum</i>	PORTUGAL	Palearctic	1; 2; 3	-
<i>M. suleyile</i>	KSA	Endemic	3	-
<i>M. tumaire</i>	KSA	Endemic	3	-
<i>M. venustum</i>	SYRIA	Palearctic	1; 3	Fig. 29

* Maps were created for species with coordinate records.

5

6