

Habitat suitability of cetaceans in the Gulf of Mexico using an ecological niche modeling approach

M. Rafael Ramírez-León¹, María C. García-Aguilar^{Corresp., 2}, Alfonsina E. Romo-Curiel², Zurisaday Ramírez-Mendoza², Arturo Fajardo-Yamamoto², Oscar Sosa-Nishizaki²

¹ Posgrado en Ecología Marina, Centro de Investigación Científica y de Educación Superior de Ensenada, Baja California, Ensenada, Baja California, Mexico

² Departamento de Oceanología Biológica, Centro de Investigación Científica y de Educación Superior de Ensenada, Baja California, Ensenada, Baja California, Mexico

Corresponding Author: María C. García-Aguilar
Email address: gaguilar@cicese.mx

Background. The Gulf of Mexico (GOM) is a semi-enclosed sea where the waters of the United States, Mexico and Cuba converge. At least 21 species of cetaceans inhabit it. The only mysticete (baleen whale) is found in the northeast (U.S. waters). The distribution of the 20 species of odontocetes (toothed cetaceans) is well understood in U.S. waters, but practically unknown in Mexican and Cuban waters. In this study we used sighting data from several odontocete species to construct habitat suitability maps in order to identify geographical regions suitable for high diversity throughout the GOM. **Methods.** Historical datasets of georeferenced sightings from across the GOM were used to implement the maximum entropy algorithm (MaxEnt) to model the habitat suitability of each species. Five environmental predictors were used, selected for their influence over the occurrence of cetaceans: two oceanographic predictors (sea surface temperature and chlorophyll-a concentration), and three bathymetric predictors (depth, slope, and distance to 200-m isobath). A spatial approach based on the habitat suitability maps was used to identify the suitable regions. **Results.** Only 12 species were modeled, which were the ones with the minimum sample size required. The models performed well, showing good discriminatory power and slight overfitting. Overall, depth, minimum sea surface temperature, and bottom slope were the most contributing predictor in the models. High suitability areas of 10 species were located on the continental slope, and four suitable regions were identified: (1) the Mississippi Canyon and the Louisiana-Texas slope in the northern GOM, (2) the west Florida slope in the east-northeastern GOM, (3) the Rio Grande slope in the west-northwestern GOM, and (4) Tamaulipas-Veracruz in the west-southwestern GOM.

Habitat suitability of cetaceans in the Gulf of Mexico using an ecological niche modeling approach.

M. Rafael Ramírez-León¹, María C. García-Aguilar², Alfonsina E. Romo-Curiel², Zurisaday Ramírez-Mendoza², Arturo Fajardo-Yamamoto², Oscar Sosa Nishizaki²

¹Posgrado en Ecología Marina, Centro de Investigación Científica y de Educación Superior de Ensenada, Baja California, Ensenada, Baja California, Mexico.

²Departamento de Oceanología Biológica, Centro de Investigación Científica y de Educación Superior de Ensenada, Baja California, Ensenada, Baja California, Mexico.

Corresponding Author:

María C. García-Aguilar

Carretera Ensenada-Tijuana N° 3918, Ensenada, Baja California, 22860, Mexico.

Email address: gaguilar@cicese.mx

Abstract

Background. The Gulf of Mexico (GOM) is a semi-enclosed sea where the waters of the United States, Mexico and Cuba converge. At least 21 species of cetaceans inhabit it. The only mysticete (baleen whale) is found in the northeast (U.S. waters). The distribution of the 20 species of odontocetes (toothed cetaceans) is well understood in U.S. waters, but practically unknown in Mexican and Cuban waters. In this study we used sighting data from several odontocete species to construct habitat suitability maps in order to identify geographical regions suitable for high diversity throughout the GOM.

Methods. Historical datasets of georeferenced sightings from across the GOM were used to implement the maximum entropy algorithm (MaxEnt) to model the habitat suitability of each species. Five environmental predictors were used, selected for their influence over the occurrence of cetaceans: two oceanographic predictors (sea surface temperature and chlorophyll-*a* concentration), and three bathymetric predictors (depth, slope, and distance to 200-m isobath). A spatial approach based on the habitat suitability maps was used to identify the suitable regions.

Results. Only 12 species were modeled, which were the ones with the minimum sample size required. The models performed well, showing good discriminatory power and slight overfitting. Overall, depth, minimum sea surface temperature, and bottom slope were the most contributing predictor in the models. High suitability areas of 10 species were located on the continental slope, and four suitable regions were identified: (1) the Mississippi Canyon and the Louisiana-Texas slope in the northern GOM, (2) the west Florida slope in the east-northeastern GOM, (3) the Rio Grande slope in the west-northwestern GOM, and (4) Tamaulipas-Veracruz in the west-southwestern GOM.

Conclusions. We were able to detect four geographic regions in the GOM where a high diversity of odontocetes is expected, all located on the continental slope. Although the methodology to identify them (spatial overlap) is a very conservative approach, it is useful for conservation and management purposes. The paucity of data did not allow all species to be modeled, which highlights the importance of establishing transboundary monitoring programs.

Keywords: Odontocetes, species distribution, MaxEnt, segregated distribution.

Introduction

Understanding species' geographic distribution patterns and related environmental factors is a central topic of population ecology (Guisan & Zimmermann, 2000). Environmental factors include both abiotic conditions that influence the physiological response (e.g., temperature) determined in turn by the species' adaptive responses and the interspecific interactions (e.g., prey availability) (Soberón & Peterson, 2005; Peterson et al., 2011). Cetaceans are a group of fully aquatic mammals whose anatomical, morphological and physiological adaptations have allowed them to colonize a wide variety of aquatic habitats (Katona & Whitehead, 1988). Nevertheless, their distribution is usually explained in terms of the abundance of prey, primarily controlled by dynamic oceanographic conditions (e.g., sea surface temperature and mesoscale processes), as well as by physiographic features (e.g., bottoms depth and slope) (Kenney et al., 1997; Forcada, 2018).

The Gulf of Mexico (GOM) is a semi-enclosed sea connected to the Atlantic Ocean and Caribbean Sea. It has a 1.6 million km² surface and includes the Exclusive Economic Zones (EEZ) of Mexico (which represent ~55% of the total surface of the GOM), the United States

(~40%), and Cuba (~5%) (De Lanza Espino & Gómez-Rojas, 2004). There are at least 21 species of cetaceans that inhabit the GOM, including one mysticete, the Bryde's whale (*Balaenoptera edeni*), which is distributed exclusively in the northeastern GOM (Soldevilla et al., 2017), and 20 odontocetes (Würsig, 2017).

The distribution of cetaceans in the northern GOM (i.e., the U.S. EEZ) has been extensively studied. Based on sighting records, Maze-Foley & Mullin (2006) divided the cetaceans into two communities: (1) the continental shelf community, which includes the bottlenose dolphin (*Tursiops truncatus*), the Atlantic spotted dolphin (*Stenella frontalis*), and the Bryde's whale, and (2) the continental slope community, which comprises the remaining species, although the rough-toothed dolphin (*Steno bredanensis*) can be found in both. More recently, Roberts et al. (2016) used density surface models to describe the spatial distribution of cetaceans in the northern GOM. Overall, their results are consistent with those of Maze-Foley & Mullin (2006), although they highlight the importance of the continental slope and submarine canyons, such as the Mississippi Canyon, as areas of high density of cetaceans. In contrast, the distribution of cetaceans in the southern GOM (i.e., the EEZs of Mexico and Cuba) is poorly understood. In fact, only one study has covered this region, but it was conducted by extrapolating data from the northern GOM (Mannocci et al., 2017).

Besides the ecological relevance of the GOM, it is an important economic area where fishing, tourism, and the hydrocarbon industry generate billions of dollars annually (Karnauskas et al., 2013), and it is a key transportation region (Shepard et al., 2013). Given its economic importance, the GOM ecosystem is under increasing anthropogenic pressure, threatening cetacean populations (Roberts et al., 2016). To determine the extent and impact of these hazards and to optimize threat mitigation and conservation measures, it is necessary to have accurate

predictions of their distribution on a broader scale; that is, at the ecosystem level. However, the latter is complicated given the limited data on cetaceans in the southern GOM (Ramírez-León et al., 2020), but one way to achieve this is to use ecological niche models (ENM).

ENM are statistical tools that define the distribution of suitable habitats of a species based on its ecological requirements (Peterson et al., 2011). The rationale is that the records, which are discontinuous in nature, are related to environmental and/or spatial characteristics (environmental predictors) to predict the suitable areas of the species in unsampled locations; therefore, the maps produced are spatially continuous, showing the regions where greater aggregation is expected (Franklin, 2010). Thus, areas of high habitat suitability are defined as those sites where ideal (or favorable) conditions exist for a species' long-term subsistence (Peterson & Soberón, 2012).

Our objective was to estimate the habitat suitability of the odontocetes of the GOM to identify those geographical regions that could support a high diversity of these cetaceans. The analysis included historical datasets of georeferenced sightings (presence-only data) recorded in both the south and north of the GOM. We used the maximum entropy (MaxEnt) modeling approach (Phillips, Anderson & Schapire, 2006), and five environmental predictors were selected based on their documented importance in determining the occurrence of cetaceans, either directly or indirectly: sea surface temperature, chlorophyll-*a* concentration, bottom depth and slope, and distance to the 200-m isobath (e.g., Praca et al., 2009; Fernandez et al., 2018; Pace et al., 2018).

Materials and Methods

Study and modeling area

The modeling area was not restricted to the GOM, due to the high movement capacity of cetaceans, and because there are no physical barriers in the marine environment for them. The area was extended to include the warm-temperate and tropical oceanic provinces of the northwest Atlantic Ocean (Fig. 1a; Spalding et al., 2007).

The physiography of the GOM is complex and consists of 13 physiographic sub-provinces (Fig 1b). The continental shelf (≤ 200 m deep) can be very narrow, as the Tamaulipas-Veracruz shelf, or extensive, like the Yucatan and Florida shelves. The continental slope extends from the 200 m continental shelf break to 2,800 m depth, and there are vast canyons, such as the Mississippi Canyon. The oceanic zone extends beyond the slope up to the abyssal plain, where depths $> 3,500$ m are reached (Bouma & Roberts, 1990; Monreal-Gómez, Salas-de León & Velasco-Méndez, 2004). The GOM oceanic waters have oligotrophic conditions that contrast with the eutrophic coastal regions, which receive a high nutrient input by river discharges, mainly in the northern GOM (Biggs, 1992; Lohrenz et al., 1999; Muller-Karger et al., 2015).

Presence-only data

Historical georeferenced sightings (presence-only data) of odontocetes were compiled. We discarded the Blainville's beaked whale (*Mesoplodon densirostris*), Gervais's beaked whale (*M. europaeus*), killer whale (*Orcinus orca*), and Fraser's dolphin (*Lagenodelphis hosei*) from our study because sightings of these species are infrequent (Würsig, 2017). The presence-only data of the remaining 16 species (Table 1; Supplemental Data S1) were collected from the literature (e.g., peer-reviewed articles, thesis, and technical reports), and digital databases of the Sistema Nacional de Información Sobre la Biodiversidad (SNIB; <http://www.snib.mx/>; CONABIO, 2016) and Ocean Biogeographic Information System Spatial Ecological Analysis of Megavertebrate

Populations (OBIS-SEAMAP; <http://seamap.env.duke.edu/>; Halpin et al., 2006). To reduce the sampling bias (the north of the GOM is oversampled relative to the south) and the spatial autocorrelation that negatively affects model performance (Boria et al., 2014; Varela et al., 2014), we filtered our databases (one per species) using the spThin package (Aiello-Lammens et al., 2015) in R software (R Core Team, 2019). The *thin* function uses a random approach to return a dataset with the maximum number of records for a given distance restriction, which in this study it was defined by the average daily displacement of each species (Table 1; Supplemental Information S2).

Environmental predictors

Five environmental predictors were selected based on previous knowledge about the environmental factors that influence the cetaceans' occurrence (e.g., Praca et al., 2009; Fernandez et al., 2018; Pace et al., 2018). The selected predictors included both oceanographic and bathymetric variables. Used oceanographic predictors were the sea surface temperature (SST, °C) and chlorophyll-*a* concentration (Chl-*a*, mg/m³), included in three metrics: mean, minimum, and maximum. Data of both variables were downloaded from the Ocean Color portal (<https://oceancolor.gsfc.nasa.gov/>; NASA, 2018) of the MODIS-Aqua sensor for the period July 2002 –December 2018. The data are at an L3 processing level with a spatial resolution of 0.041° (~4 km). Weekly values (8-d composite) were downloaded and averaged across the 16 years with available data. Bathymetric predictors were depth (*D*, m), bottom slope (*S*, degrees), and distance to the 200-m isobath (*D*₂₀₀, m). The first was acquired from the General Bathymetric Chart of the Ocean (GEBCO; <https://www.gebco.net/>; IOC & IHO, 2018) with a spatial resolution of 0.008° (~1 km); the other two were calculated from the depth using the raster

package (Hijmans et al., 2013) in R software. These bathymetric predictors were re-projected at a spatial resolution of 0.041°.

The co-linearity among environmental predictors was evaluated using the Pearson correlation coefficient (ρ) (Dormann et al., 2012; Cruz-Cárdenas et al., 2014). If $\rho \geq 0.70$ (Dormann et al., 2013), a principal component analysis was performed (Supplemental Information S3) to determine which of the correlated predictors should be discarded.

Habitat suitability modeling

We used the MaxEnt algorithm (Phillips, Anderson & Schapire, 2006) to predict the habitat suitability of odontocetes in the GOM. MaxEnt assumes that the species are distributed uniformly (i.e., the maximum entropy distribution) on the modeling area, and the environmental values constrain this distribution at the presence of records locations (Phillips, Anderson & Schapire, 2006; Phillips et al., 2017). The habitat suitability modeling for each species was conducted using the ENMeval package (Muscarella et al., 2014, 2016) in R. We built models with a random sample of 10,000 background points (i.e., points not registered as occurrence records in the modeling area that are contrasted with the occurrence positions) and select the *Linear*, *Quadratic*, and *Hinge* features of the MaxEnt algorithm. The cross-validation of the models was done using the block method that splits the presence data into four bins, three as training data and one as test data, based on the latitude and longitude lines that divided the occurrence localities (Muscarella et al., 2014).

The performance of each model was evaluated using the area under the receiver-operator curve (AUC), which measures the discriminatory ability of each model, and the omission rate (OR), which indicates the proportion of test localities that fall into cells not predicted as suitable

(Philips, Anderson & Schapire, 2006). An AUC of 1 indicates perfect discrimination between sites where the species is present or absent, and an AUC < 0.5 indicates that the model performance is less than a random assumption (Elith et al., 2006). We used the 10-percentile training omission rate (OR_{10}) because it is less sensitive to the outlier presence locations (Radosavljevic & Anderson, 2014). Omission rates greater than the expected value of 0.1 (or 10%) suggest model overfitting (Peterson et al., 2011; Radosavljevic & Anderson, 2014). Finally, we used the contribution percentages returned by each MaxEnt model to evaluate the contribution of each environmental predictor (Phillips, Anderson & Schapire, 2006).

We selected the logistic output and obtained the habitat suitability for each $0.041^\circ \times 0.041^\circ$ cell of the modeling area, which was expressed in an interval between 0 (unsuitable conditions) and 1 (highly suitable conditions). In this study, high suitability areas were defined as those sites (cells) with suitability values ≥ 0.6 (Kaschner et al., 2011). A spatial approach based on the habitat suitability maps was used to identify suitable regions for cetaceans; that is, regions capable of supporting a high diversity of cetaceans (i.e., suitable regions) were defined as regions where the high suitability areas of at least seven species overlap.

Results

Habitat suitability was modeled for only twelve species (Table 1), which were those that after filtering had the minimum sample required (≥ 30 presence records; Wisz et al., 2008). The pygmy sperm whale (*Kogia breviceps*), pygmy killer whale (*Feresa attenuata*), false killer whale (*Pseudorca crassidens*), and melon-head whale (*Peponocephala electra*) were excluded due to small sample size (Table 1).

The models showed a good degree of discriminatory ability based on the AUC scores, which ranged from 0.74 (the pantropical spotted dolphin, *Stenella attenuata*, model) to 0.91 (the bottlenose dolphin model) (Table 2). On the other hand, the OR₁₀ value was close to the expected value in some models, such as the rough-toothed dolphin, but in others it was higher, as in the spinner dolphin (*Stenella longirostris*) model (Table 2), suggesting some degree of overfitting.

The environmental predictors used in each model differ (Table 2; Supplemental Information S4). The bottlenose dolphin models had the fewest predictors, while the Risso's dolphin (*Grampus griseus*), Atlantic spotted dolphin, pantropical spotted dolphin and spinner dolphin models had the most. Slope was included in 11 models, while both depth and minimum-SST in 10 (Table 2). However, in terms of contribution, depth was the most important environmental predictor, with a contribution of > 25% in seven models, followed by the minimum-SST, which had an important contribution in five models.

High suitability areas for 10 species were located on the continental slope (Figs. 2-4). The pantropical spotted dolphin seems to be the species with the widest distribution, potentially occupying the entire continental slope. High suitability areas for the sperm whale (*Physeter macrocephalus*), short-finned pilot whale (*Globicephala macrorhynchus*), and spinner dolphin were found on the inner continental slope, while for the rough-toothed dolphin they were located both on the outer continental shelf and on the slope. Dwarf sperm whale (*Kogia sima*), Risso's dolphin, and Cuvier's beaked whale (*Ziphius cavirostris*) models show almost continuous high suitability along the northern continental slope, from Florida to Louisiana-Texas, even extending to the Rio Grande and Tamaulipas-Veracruz slopes. High suitability areas for the Clymene dolphin (*Stenella clymene*) were found on the northern slope, with small patches on the Tamaulipas-Veracruz slope. Striped dolphin (*Stenella coeruleoalba*) model identified as high

suitability areas some patches on the Tamaulipas-Veracruz slope and on the bay of Campeche.

The Atlantic spotted dolphin model indicates that high suitability areas are located from the continental shelf to the inner slope of the entire GOM, whereas the model of the bottlenose dolphin points to the continental shelf, from Florida to the Tamaulipas-Veracruz.

The main region of high diversity of odontocetes was located between the Mississippi Canyon and the Louisiana-Texas slope (Fig. 5). Other suitable regions were identified on the west Florida slope and on the western continental slope, between the Rio Grande and Tamaulipas-Veracruz slopes

Discussion

ENM are powerful tools for generating spatially explicit maps of species' habitat suitability. We used the MaxEnt approach to model the habitat suitability of the GOM odontocetes, using data from both the north and south, and managed to identify regions where high diversity can be expected. We decided to use this approach because it allows the development of reliable models of the potential distribution based on presence-only data, although it is important to emphasize that these models do not represent the probability of the presence of a species. However, due to the paucity of data, we were able to model only 12 of the 20 species present in the GOM. Furthermore, because we use historical sighting records, the resulting maps are integrated images that show no temporal variations, and no biotic interaction were considered. Biotic interactions might improve habitat suitability models, however, require abundance demographic data at population over time (Anderson, 2017).

All models had a good discrimination power, with AUC values > 0.70 , indicating that the results are reliable, and can be used in planning management and conservation measures (Elith et

al., 2006; Raes & Aguirre-Gutiérrez, 2018). On the other hand, the OR_{10} values were higher than the expected value; however, they are within the range reported in other studies (e.g., Kramer-Schadt et al., 2013; Arthur, Morrison & Morey, 2019). High OR_{10} values suggest overfitting, which could be due to sampling bias and/or noise in the presence data (Anderson & Gonzalez, 2011; Merow et al., 2014). In our study area, the south of the GOM is under-sampled compared to the north. We attempted to reduce this bias using the spatial filtering to minimize the omission error, but it may not have been completely successful, especially in the case of the spinner dolphin model.

Among the used environmental predictors, depth was the most important, followed by minimum SST and bottom slope. These results are not surprising since it is well known that these variables influence the occurrence of cetaceans directly; for example, some species display relatively persistent bathymetric associations (Yen, Sydeman & Hyrenbach, 2004; Harvey et al., 2017), but mainly indirectly by playing a determining role in the availability, distribution, and abundance of their prey (Davis et al., 2002; MacLeod, 2009; Forcada, 2018). In fact, previous studies have shown that the distribution of several species of cetaceans of the GOM is strongly related to depth (e.g., Baumgartner, 1997; Davis et al., 1998; Baumgartner et al., 2001).

Our results are consistent with the segregated distribution of cetaceans proposed by Maze-Foley & Mullin (2006) for the northern GOM. The two dolphin species of the shelf community, the Atlantic spotted dolphins and the bottlenose dolphins, use different habitats. High suitability areas of the Atlantic spotted dolphins were located on the outer continental shelf and the inner slope, while the bottlenose dolphin has coastal habitats, occupying shallower waters; actually, it is the only species that inhabits lagoons, estuaries, and bays (e.g., Mullin et al., 1990; Griffin & Griffin, 2003; Martínez-Serrano et al., 2011). The continental slope community is composed of

the remaining species (Maze-Foley & Mullin, 2006; present study), although densities of these can vary seasonally, at least in the northern GOM (Roberts et al., 2016; Mannocci et al., 2017). The aggregation of multiple species reveals important biological regions capable of supporting a high cetacean diversity (Harvey et al., 2017). Ten of the 12 species modeled showed habitat suitability areas along the continental slope, consistent with the relatively high diversity of cetaceans observed on the northern GOM continental slope (Davis et al., 2002; Maze-Foley & Mullin, 2006; Roberts et al., 2016). The exploitation of different types of habitat and prey allows this co-existence (Bearzi, 2005; Schick et al., 2011). The continental slope of the GOM covers a large area and presents underwater canyons (Bouma & Roberts, 1990), where the main prey (e.g., cephalopods) of the deep-diving species can accumulate (Biggs, Leben & Ortega-Ortiz, 2000; O'Hern & Biggs, 2009; Moors-Murphy, 2014). On the other hand, the species that primarily feed on epipelagic prey preferentially use the upper layers of the water column, where mesoscale structures occur (Davis et al., 1998, 2002).

We identify four suitable regions. The most notable was located in the north, encompassing the Mississippi Canyon and the Louisiana-Texas slopes, consistent with that previously reported for the north of the GOM (e.g., Mullin & Fulling, 2006; Roberts et al., 2016). The other regions were located on the west Florida slope (east-northeast of the GOM), the Rio Grande slope (west-northwestern of the GOM), and the Tamaulipas-Veracruz slope (west-southwestern of the GOM). All these regions are characterized by their high primary productivity. In the north, productivity is directly influenced by the input of nutrients from the Mississippi and Atchafalaya rivers (Lohrenz et al., 1999), while in the west by the Grande and Pánuco rivers (Salmerón-García et al., 2011). The plumes of nutrient-rich waters are transported through the continental shelf (Del Castillo et al., 2001; Morey et al., 2003; Zavala-Hidalgo, Morey & O'Brien, 2003),

reaching the slope by the interactions of anticyclonic-cyclonic eddies (Toner, 2003; Martínez-López & Zavala-Hidalgo, 2009). However, the largest region suitable for cetaceans was located in the north for two possible reasons. First, the large nutrient input from the Mississippi River into the shelf ecosystem favors huge phytoplankton blooms on spatial scales of tens to hundreds of kilometers (Lohrenz et al., 1997). Second, the continental slope in this region is extremely wide, which could favor the convergence of a greater number of cetacean species.

Conclusions

We identified areas of high suitability for 12 species of odontocetes in the GOM through the implementation of an ENM. Unfortunately, the paucity of data did not allow modeling all the species, which highlights the importance to establish transboundary research and monitoring programs between the U.S., Cuba, and Mexico to improve knowledge on the cetaceans of the GOM. Even so, we were able to detect four geographic regions where a high diversity of odontocetes is expected, all located on the continental slope. These suitable regions were identified using a spatial overlay, which although it is a very conservative approach (Harvey et al., 2017), it can be useful to detect areas where to focus conservation efforts (Tolimieri et al., 2015).

Acknowledgments

This is a contribution of the Gulf of Mexico Research Consortium (CIGoM). We acknowledge PEMEX's specific request to the Hydrocarbon Fund to address the environmental effects of oil spills in the Gulf of Mexico. We thank Paloma Ladrón de Guevara, Georgina Castro, and Fernanda Urrutia for helping to obtain and compile the occurrence data; Aimie Moulin and Rigel Zaragoza

for helping to process the environmental layers, and Fabricio Villalobos and Carlos Yañez-Arenas for technical support; Sharon Herzka for key logistic support. We appreciate the suggestions of Ladd Irvine, Jason Roberts and one anonymous reviewer, which greatly improved the manuscript.

References

- Aiello-Lammens ME, Boria RA, Radosavljevic A, Vilela B, Anderson RP. 2015. spThin: an R package for spatial thinning of species occurrence records for use in ecological niche models. *Ecography* 38:541–545. DOI: 10.1111/ecog.01132.
- Anderson RP. 2017. When and how should biotic interactions be considered in models of species niches and distributions? *Journal of Biogeography* 44:8-17. DOI: 10.1111/jbi.12825
- Anderson RP, Gonzalez I. 2011. Species-specific tuning increases robustness to sampling bias in models of species distributions: An implementation with Maxent. *Ecological Modelling* 222:2796–2811. DOI: 10.1016/j.ecolmodel.2011.04.011.
- Arthur FH, Morrison III WR, Morey AC. 2019. Modeling the potential range expansion of larger grain borer, *Prostephanus truncatus* (Coleoptera: Bostrichidae). *Scientific Reports* 9: 6862. DOI: 10.1038/s41598-019-42974-5
- Baird RW, Schorr GS, Webster DL, Mahaffy SD, McSweeney DJ, Hanson MB, Andrews RD. 2009. *Movements of satellite-tagged Cuvier's and Blainville's beaked whales in Hawaii' i: evidence for an offshore population of Blainville's beaked whales*. U.S. Department of Commerce.
- Baird RW, Schorr GS, Webster D I., McSweeney DJ, Hanson MB, Andrews RD. 2010. Movements and habitat use of satellite-tagged false killer whales around the main Hawaiian Islands. *Endangered Species Research* 10:107–121. DOI: 10.3354/esr00258.

- Baird RW, Schorr GS, Webster DL, McSweeney DJ, Hanson MB, Andrews RD. 2011. Movements of two satellite-tagged pygmy killer whales (*Feresa attenuata*) off the island of Hawai'i. *Marine Mammal Science* 27:E332–E337. DOI: 10.1111/j.1748-7692.2010.00458.x.
- Baird RW, Webster DL, Aschettino JM, Verbeck D, Mahaffy SD. 2012. *Odontocete movements off the island of Kau'I: Results of satellite tagging and photo-identification efforts in January 2012*. Cascadia Research Collective.
- Baumgartner MF. 1997. The distribution of Risso's dolphin (*Grampus griseus*) with respect to the physiography of the northern Gulf of Mexico. *Marine Mammal Science* 13:614–638.
- Baumgartner MF, Mullin KD, May LN, Leming TD. 2001. Cetacean habitats in the northern Gulf of Mexico. *Fishery Bulletin* 99:219–239.
- Bearzi M. 2005. Dolphin sympatric ecology. *Marine Biology Research* 1:165–175. DOI: 10.1080/17451000510019132.
- Biggs DC. 1992. Nutrients, plankton, and productivity in a warm-core ring in the western Gulf of Mexico. *Journal of Geophysical Research: Oceans* 97:2143–2154.
- Biggs DC, Leben RR, Ortega-Ortiz JG. 2000. Ship and satellite studies of mesoscale circulation and sperm whale habitats in the northeast Gulf of Mexico during GulfCet II. *Gulf of Mexico Science* 18:15–22.
- Boria RA, Olson LE, Goodman SM, Anderson RP. 2014. Spatial filtering to reduce sampling bias can improve the performance of ecological niche models. *Ecological Modelling* 275:73–77. DOI: 10.1016/j.ecolmodel.2013.12.012.
- Bouma AH, Roberts HH. 1990. Northern Gulf of Mexico continental slope. *Geo-Marine Letters* 10:177–181. DOI: 10.1007/BF02431064.

CONABIO (Comisión Nacional para el Conocimiento y Uso de la Biodiversidad). 2016. Sistema Nacional de Información sobre Biodiversidad de México. <http://www.snib.mx/> (accessed on 1 April 2016)

Cruz-Cárdenas G, López-Mata L, Villaseñor JL, Ortiz E. 2014. Potential species distribution modeling and the use of principal component analysis as predictor variables. *Revista Mexicana de Biodiversidad* 85:189–199. DOI: 10.7550/rmb.36723.

Davis RW, Fargion GS, May N, Leming TD, Baumgartner M, Evans WE, Hansen LJ, Mullin K. 1998. Physical habitat of cetaceans along the continental slope in the north-central and western Gulf of Mexico. *Marine Mammal Science* 14:490–507.

Davis RW, Ortega-Ortiz JG, Ribic CA, Evans WE, Biggs DC, Ressler PH, Cady RB, Leben RR, Mullin KD, Würsig B. 2002. Cetacean habitat in the northern oceanic Gulf of Mexico. *Deep Sea Research Part I: Oceanographic Research Papers* 49:121–142.

Davis RW, Worthy GAJ, Würsig B, Lynn SK, Townsend FI. 1996. Diving behavior and at-sea movements of an Atlantic spotted dolphin in the Gulf of Mexico. *Marine Mammal Science* 12:569–581. DOI: 10.1111/j.1748-7692.1996.tb00069.x.

De Lanza Espino G, Gómez-Rojas JC. 2004. Physical and chemical characteristics of the Gulf of Mexico. In: Pisanty I, Ezcurra E, Whithers K, Nipper M eds. *Environmental analysis of the Gulf of Mexico*. México, D.F. Harte Research Institute for Gulf of Mexico Studies Special Publication Series N°1, 41–61.

Del Castillo CE, Coble PG, Conmy RN, Müller-Karger FE, Vanderbloemen L, Vargo GA. 2001. Multispectral in situ measurements of organic matter and chlorophyll fluorescence in seawater: documenting the intrusion of the Mississippi River plume in the West Florida Shelf. *Limnology and Oceanography* 46:1836–1843. DOI: 10.4319/lo.2001.46.7.1836.

- Dormann CF, Elith J, Bacher S, Buchmann C, Carl G, Carré G, Marquéz JRG, Gruber B, Lafourcade B, Leitão PJ, Münkemüller T, McClean C, Osborne PE, Reineking B, Schröder B, Skidmore AK, Zurell D, Lautenbach S. 2013. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* 36:27–46. DOI: 10.1111/j.1600-0587.2012.07348.x.
- Dormann CF, Schymanski SJ, Cabral J, Chuine I, Graham C, Hartig F, Kearney M, Morin X, Römermann C, Schröder B. 2012. Correlation and process in species distribution models: bridging a dichotomy. *Journal of Biogeography* 39:2119–2131.
- Elith J, Graham CH, Anderson RP, Dudík M, Ferrier S, Guisan A, Hijmans RJ, Huettmann F, Leathwick JR, Lehmann A, Li J, Lohmann LG, Loiselle BA, Manion G, Moritz C, Nakamura M, Nakazawa Y, McC. Overton J, Peterson AT, Phillips SJ, Richardson K, Scachetti-Pereira R, Schapire RE, Soberón J, Williams S, Wisz MS, Zimmermann NE. 2006. Novel methods improve prediction of species' distributions from occurrence data. *Ecography* 29:129–151.
- Fernandez M, Yesson C, Gannier A, Miller PI, Azevedo JMN. 2018. A matter of timing: how temporal scale selection influences cetacean ecological niche modelling. *Marine Ecology Progress Series* 595:217–231. DOI: 10.3354/meps12551.
- Franklin J. 2010. *Mapping species distribution: spatial inference and prediction*. Cambridge University Press.
- Forcada J. 2018. Distribution. In: Würsig B, Thewissen JGM, Kovacs KM eds. *Encyclopedia of Marine Mammals*. Elsevier, 259–262. DOI: 10.1016/B978-0-12-804327-1.00106-0.
- Gannier A. 1999. Diel variations of the striped dolphin distribution on the French Riviera (Northwestern Mediterranean Sea). *Aquatic Mammals* 25:123–134.

- Griffin RB, Griffin NJ. 2003. Distribution, habitat partitioning, and abundance of Atlantic spotted dolphins, bottlenose dolphins, and loggerhead sea turtles on the eastern Gulf of Mexico continental shelf. *Gulf of Mexico Science* 21:23–34.
- Guisan A, Zimmermann NE. 2000. Predictive habitat distribution models in ecology. *Ecological Modelling* 135:147–186.
- Halpin PN, Read AJ, Best BD, Hyrenbach KD, Fujioka E, Coyne MS, Crowder LB, Freeman SA, Spoerri C. 2006. OBIS-SEAMAP: developing a biogeographic research data commons for the ecological studies of marine mammals, seabirds, and sea turtles. *Marine Ecology Progress Series* 316:239–246. DOI: 10.3354/meps316239.
- Harvey GKA, Nelson TA, Fox CH, Paquet PC. 2017. Quantifying marine mammal hotspots in British Columbia, Canada. *Ecosphere* 8: e01884. DOI: 10.1002/ecs2.1884
- Hijmans RJ, van Etten J, Mattiuzzi M, Sumner M, Greenberg J, Lamigueiro O, Bevan A, Racine E, Shortridge A. 2013. Raster package in R.
- IOC (Intergovernmental Oceanographic Commission), IHO (International Hydrographic Organization). 2018. GEBCO Digital Atlas published by the British Oceanographic Data Centre on behalf of IOC and IHO. <http://www.gebco.net/> (accessed 1 March 2018)
- Irvine AB, Scott MD, Wells RS, Kaufmann JH. 1981. Movements and activities of the Atlantic bottlenose dolphin, *Tursiops truncatus*, near Sarasota, Florida. *Fishery Bulletin* 79:671–688.
- Karnauskas M, Schirripa MJ, Kelble CR, Cook GS, Craig JK. 2013. *Ecosystem Status Report for the Gulf of Mexico*. U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southeast Fisheries Science Center.

- 426 Kaschner K, Tittensor DP, Ready J, Gerrodette T, Worm B. 2011. Current and Future Patterns of
427 Global Marine Mammal Biodiversity. PLOS ONE 6:e19653. DOI:
428 10.1371/journal.pone.0019653.
- 429 Katona S, Whitehead H. 1988. Are cetacean ecologically important? *Oceanography and Marine*
430 *Biology: An Annual Review* 26:553–568.
- 431 Kenney RD, Scott GP, Thompson TJ, Winn HE. 1997. Estimates of prey consumption and
432 trophic impacts of cetaceans in the USA northeast continental shelf ecosystem. *Journal of*
433 *Northwest Atlantic Fishery Science* 22:155–171.
- 434 Kramer-Schadt S, Niedballa J, Pilgrim JD, Schröder B, Lindenborn J, Reinfelder V, Stillfried M,
435 Heckmann I, Scharf AK, Augeri DM, Cheyne SM, Hearn AJ, Ross J, Macdonald DW,
436 Mathai J, Eaton J, Marshall AJ, Semiadi G, Rustam R, Bernard H, Alfred R, Samejima
437 H, Duckworth JW, Breitenmoser-Wuersten C, Belant JL, Hofer H, Wilting A. 2013. The
438 importance of correcting for sampling bias in MaxEnt species distribution models.
439 *Diversity and Distributions* 19:1366-1379. DOI: 10.1111/ddi.12096
- 440 Lohrenz SE, Fahnenstiel GL, Redalje DG, Lang GA, Chen X, Dagg MJ. 1997. Variations in
441 primary production of northern Gulf of Mexico continental shelf waters linked to nutrient
442 inputs from the Mississippi River. *Marine Ecology Progress Series* 155: 45 – 54.
443 DOI:10.3354/meps155045
- 444 Lohrenz SE, Wiesenburg DA, Arnone RA, Chen X. 1999. What controls primary production in
445 the Gulf of Mexico? In: Kumpf H, Steidinger K, Sherman K eds. *The Gulf of Mexico*
446 *Large Marine Ecosystem: Assessment, Sustainability and Management*. Malden, M.A.:
447 Blackwell Science, 151–170.

- MacLeod CD. 2009. Global climate change, range changes and potential implications for the conservation of marine cetaceans: a review and synthesis. *Endangered Species Research* 7:125–136.
- Mannocci L, Roberts JJ, Miller DL, Halpin PN. 2017 Extrapolating cetacean densities to quantitatively assess human impacts on populations in the high seas. *Conservation Biology* 31: 601-614. DOI: 10.1111/cobi.12856.
- Martínez-López B, Zavala-Hidalgo J. 2009. Seasonal and interannual variability of cross-shelf transports of chlorophyll in the Gulf of Mexico. *Journal of Marine Systems* 77:1–20. DOI: 10.1016/j.jmarsys.2008.10.002.
- Martínez-Serrano I, Serrano A, Heckel G, Schramm Y. 2011. Distribution and home range of bottlenose dolphins (*Tursiops truncatus*) off Veracruz, Mexico. *Ciencias Marinas* 37:379–392.
- Maze-Foley K, Mullin KD. 2006. Cetaceans of the oceanic northern Gulf of Mexico: distributions, group sizes and interspecific associations. *Journal of Cetacean Research and Management* 8:203–213.
- McAlpine DF. 2018. Pygmy and dwarf sperm whales: *Kogia breviceps* and *K. sima*. In: *Encyclopedia of marine mammals*. Elsevier, 786–788.
- Merow C, Smith MJ, Edwards Jr, TC, Guisan A, McMahon SM, Normand S, Thuiller W, Wüest RO, Zimmermann NE, Elith J. 2014. What do we gain from simplicity versus complexity in species distribution models? *Ecography* 37:1267-1281. DOI: 10.1111/ecog.00845
- Monreal-Gómez MA, Salas-de León DA, Velasco-Méndola H. 2004. The hydrodynamics of the Gulf of Mexico. In: Pisanty I, Ezcurra E, Whithers K, Nipper M eds. *Environmental*

analysis of the Gulf of Mexico. México, D.F.: Harte Research Institute for Gulf of Mexico Studies Special Publication Series N°1, 2–17.

Moors-Murphy HB. 2014. Submarine canyons as important habitat for cetaceans, with special reference to the Gully: A review. *Deep Sea Research Part II: Topical Studies in Oceanography* 104:6–19. DOI: 10.1016/j.dsr2.2013.12.016.

Morey SL, Schroeder WW, O'Brien JJ, Zavala-Hidalgo J. 2003. The annual cycle of riverine influence in the eastern Gulf of Mexico basin. *Geophysical Research Letters* 30. DOI: 10.1029/2003GL017348.

Muller-Karger FE, Smith JP, Werner S, Chen R, Roffer M, Liu Y, Muhling B, Lindo-Atichati D, Lamkin J, Cerdeira-Estrada S, Enfield DB. 2015. Natural variability of surface oceanographic conditions in the offshore Gulf of Mexico. *Progress in Oceanography* 134:54–76. DOI: 10.1016/j.pocean.2014.12.007.

Mullin KD, Fulling GL. 2004. Abundance of cetaceans in the oceanic northern Gulf of Mexico, 1996–2001. *Marine Mammal Science* 20:787–807.

Mullin KD, Lohofener RR, Hoggard W, Roden CL, Rogers CM. 1990. Abundance of bottlenose dolphins, *Tursiops truncatus*, in the coastal Gulf of Mexico. *Northeast Gulf Science* 11:113–122.

Muscarella R, Galante PJ, Soley-Guardia M, Boria RA, Kass JM, Uriarte M, Anderson RP. 2014. ENM eval: An R package for conducting spatially independent evaluations and estimating optimal model complexity for Maxent ecological niche models. *Methods in Ecology and Evolution* 5:1198–1205.

Muscarella R, Galante P, Soley-Guardia M, Boria R, Kass J, Uriarte M, Anderson R. 2016. Package "ENMeval: Automated runs and evaluations of ecological niche models".

NASA Goddard Space Flight Center, Ocean Ecology Laboratory, Ocean Biology Processing Group. 2018. Moderate-resolution Imaging Spectroradiometer (MODIS) Aqua. NASA OB.DAAC, Greenbelt, MD. <https://oceancolor.gsfc.nasa.gov/l3/> (accessed 15 August 2018).

O'Hern JE, Biggs DC. 2009. Sperm whale (*Physeter macrocephalus*) habitat in the Gulf of Mexico: satellite observed ocean color and altimetry applied to small-scale variability in distribution. *Aquatic Mammals* 35:358–366. DOI: 10.1578/AM.35.3.2009.358.

Olson PA. 2018. Pilot Whales: *Globicephala melas* and *G. macrorhynchus*. In: Würsig B, Thewissen JGM, Kovacs KM eds. *Encyclopedia of Marine Mammals (Third Edition)*. Academic Press, 701–705. DOI: 10.1016/B978-0-12-804327-1.00194-1.

Pace DS, Arcangeli A, Mussi B, Vivaldi C, Ledon C, Lagorio S, Giacomini G, Pavan G, Ardizzone G. 2018. Habitat suitability modeling in different sperm whale social groups: sperm whale habitat suitability modeling. *The Journal of Wildlife Management* 82:1062–1073. DOI: 10.1002/jwmg.21453.

Perrin WF. 2018. Spinner dolphin: *Stenella longirostris*. In: Würsig B, Thewissen JGM, Kovacs KM eds. *Encyclopedia of Marine Mammals (Third Edition)*. Academic Press, 925–928. DOI: 10.1016/B978-0-12-804327-1.00243-0.

Peterson AT, Soberón J. 2012. Species distribution modeling and ecological niche modeling: getting the concepts right. *Natureza & Conservação* 10:102–107. DOI: 10.4322/natcon.2012.019.

Peterson AT, Soberón J, Pearson RG, Anderson RP, Martínez-Meyer E, Nakamura M, Araújo MB. 2011. *Ecological niches and geographic distributions*. Princeton University Press.

Phillips SJ, Anderson RP, Dudík M, Schapire RE, Blair ME. 2017. Opening the black box: an open-source release of Maxent. *Ecography* 40:887–893. DOI: 10.1111/ecog.03049.

Phillips SJ, Anderson RP, Schapire RE. 2006. Maximum entropy modeling of species geographic distributions. *Ecological Modelling* 190:231–259. DOI: 10.1016/j.ecolmodel.2005.03.026.

Praca E, Gannier A, Das K, Laran S. 2009. Modelling the habitat suitability of cetaceans: Example of the sperm whale in the northwestern Mediterranean Sea. *Deep Sea Research Part I: Oceanographic Research Papers* 56:648–657. DOI: 10.1016/j.dsr.2008.11.001.

R Core Team. 2019. R: A language and environment for statistical computing.

Radosavljevic A, Anderson RP. 2014. Making better Maxent models of species distributions: complexity, overfitting and evaluation. *Journal of Biogeography* 41:629–643. DOI: 10.1111/jbi.12227.

Raes N, Aguirre-Gutiérrez J. 2018. Modeling framework to estimate and project species distributions space and time. In: Hoorn C, Perrigo A, Antonelli A eds. *Mountains, Climate and Biodiversity*. Wiley Blackwell, 309-320.

Ramírez-León MR, García-Aguilar MC, Aguayo-Lobo A, Fuentes-Allen I, Sosa-Nishizaki O. 2020. What do we know about cetaceans in the Mexican waters of the Gulf of Mexico? A review. *Aquatic Mammals* 46:623–632. DOI 10.1578/AM.46.6.2020.623

Roberts JJ, Best BD, Mannocci L, Fujioka E, Halpin PN, Palka DL, Garrison LP, Mullin KD, Cole TVN, Khan CB, McLellan WA, Pabst DA, Lockhart GG. 2016. Habitat-based cetacean density models for the U.S. Atlantic and Gulf of Mexico. *Scientific Reports* 6. DOI: 10.1038/srep22615.

- 537 Salmerón-García O, Zavala-Hidalgo J, Mateos-Jasso A, Romero-Centeno R. 2011.
538 Regionalization of the Gulf of Mexico from space-time chlorophyll-a concentration
539 variability. *Ocean Dynamics* 61:439–448. DOI: 10.1007/s10236-010-0368-1.
- 540 Shepard AN, Valentine JF, D’Elia CF, Yoskowitz DW, Dismukes DE. 2013. Economic impact
541 of Gulf of Mexico ecosystem goods and services and integration into restoration
542 decision-making. *Gulf of Mexico Science* 31:2. DOI: 10.18785/goms.3101.02
- 543 Schick RS, Halpin PN, Read AJ, Urban DL, Best BD, Good CP, Roberts JJ, LaBrecque EA,
544 Dunn C, Garrison LP, Hyrenbach KD, McLellan WA, Pabst DA, Palka DL, Stevick P.
545 2011. Community structure in pelagic marine mammals at large spatial scales. *Marine*
546 *Ecology Progress Series* 434:165–181. DOI: 10.3354/meps09183.
- 547 Scott MD, Chivers SJ. 2009. Movements and diving behavior of pelagic spotted dolphins.
548 *Marine Mammal Science* 25:137–160. DOI: 10.1111/j.1748-7692.2008.00241.x.
- 549 Soberón J, Peterson AT. 2005. Interpretation of models of fundamental ecological niches and
550 species' distributional areas. *Biodiversity Informatics* 2:1–10.
- 551 Soldevilla MS, Hildebrand JA, Frasier KE, Aichinger-Dias L, Martinez A, Mullin KD, Rosel PE,
552 Garrison LP. 2017. Spatial distribution and dive behavior of Gulf of Mexico Bryde's
553 whales: potential risk of vessel strikes and fisheries interactions. *Endangered Species*
554 *Research* 32:533–550. DOI: 10.3354/esr00834.
- 555 Spalding MD, Fox HE, Allen GR, Davidson N, Ferdaña ZA, Finlayson M, Halpern BS, Jorge
556 MA, Lombana A, Lourie SA, Martin KD, McManus E, Molnar J, Recchia CA, Robertson
557 J. 2007. Marine ecoregions of the world: A bioregionalization of coastal and shelf areas.
558 *BioScience* 57:573–583. DOI: 10.1641/B570707.

- 559 Tolimieri N, Shelton AO, Feist BE, Simon V. 2015. Can we increase our confidence about the
- 560 locations of biodiversity 'hotspots' by using multiple diversity indices? *Ecosphere* 6:290.
- 561 DOI: 10.1890/ES14-00363.1.
- 562 Toner M. 2003. Chlorophyll dispersal by eddy-eddy interactions in the Gulf of Mexico. *Journal*
- 563 *of Geophysical Research* 108:3105. DOI: 10.1029/2002JC001499.
- 564 Varela S, Anderson RP, García-Valdés R, Fernández-González F. 2014. Environmental filters
- 565 reduce the effects of sampling bias and improve predictions of ecological niche models.
- 566 *Ecography* 37:1084–1091. DOI: 10.1111/j.1600-0587.2013.00441.x.
- 567 Wells RS, Early GA, Gannon JG, Lingenfelter RG, Sweeney P. 2008. *Tagging and tracking of*
- 568 *Rough-toothed Dolphins (Steno bredanensis) from the March 2005 mass stranding in the*
- 569 *Florida Keys*. U.S. Department of Commerce, National Oceanic and Atmospheric
- 570 Administration, National Marine Fisheries Service, Southeast Fisheries Science Center.
- 571 Wells RS, Manire CA, Byrd L, Smith DR, Gannon JG, Fauquier D, Mullin KD. 2009.
- 572 Movements and dive patterns of a rehabilitated Risso's dolphin, *Grampus griseus*, in the
- 573 Gulf of Mexico and Atlantic Ocean. *Marine Mammal Science* 25:420–429. DOI:
- 574 10.1111/j.1748-7692.2008.00251.x.
- 575 Whitehead H. 2018. Sperm whale: *Physeter macrocephalus*. In: Würsig B, Thewissen JGM,
- 576 Kovacs KM eds. *Encyclopedia of Marine Mammals (Third Edition)*. Academic Press,
- 577 919–925. DOI: 10.1016/B978-0-12-804327-1.00242-9.
- 578 Wisz MS, Hijmans RJ, Li J, Peterson AT, Graham CH, Guisan A, NCEAS Predicting Species
- 579 Distributions Working Group†. 2008. Effects of sample size on the performance of
- 580 species distribution models. *Diversity and Distributions* 14:763–773. DOI:
- 581 10.1111/j.1472-4642.2008.00482.x.

- Würsig B. 2017. Marine Mammals of the Gulf of Mexico. In: Ward CH ed. *Habitats and Biota of the Gulf of Mexico: Before the Deepwater Horizon oil spill, Volume 2: Fish resources, fisheries, sea turtles, avian resources, marine mammals, diseases and mortalities*. New York, NY: Springer Open, 1489–1588.
- Yen PPW, Sydeman WJ, Hyrenbach KD. 2004. Marine birds and cetacean associations with bathymetric habitats and shallow-water topographies: implications for trophic transfer and conservation. *Journal of Marine Systems* 50: 79-99. DOI: 10.1016/j.jmarsys.2003.09.015.
- Zavala-Hidalgo J, Morey SL, O'Brien JJ. 2003. Seasonal circulation on the western shelf of the Gulf of Mexico using a high-resolution numerical model. *Journal of Geophysical Research* 108. DOI: 10.1029/2003JC001879.

Table 1(on next page)

Georeferenced records for 16 odontocetes from the Gulf of Mexico

Total number of georeferenced sightings of 16 odontocetes in the Gulf of Mexico, period of the presence data, filtering distance (average daily displacement in km), and number of sightings used in modeling (sample size). Modeled species are marked with *.

Table 1. Total number of georeferenced sightings of 16 odontocetes in the Gulf of Mexico, period of the presence data, filtering distance (average daily displacement in km), and number of sightings used in modeling (sample size). Modeled species are marked with *.

Scientific name	Common name	Period	Total sightings	Filtering distance	Used sightings (n)
<i>Physeter macrocephalus</i> *	Sperm whale	1978 - 2017	810	90 ¹	70
<i>Kogia breviceps</i>	Pygmy sperm whale	1992 - 2011	51	75 ²	17
<i>Kogia sima</i> *	Dwarf sperm whale	1990 - 2011	319	75 ²	37
<i>Ziphius cavirostris</i> *	Cuvier's beaked whale	1990 - 2017	88	25 ³	39
<i>Feresa attenuata</i>	Pygmy killer whale	1990 - 2008	24	70 ⁴	16
<i>Peponocephala electra</i>	Melon-headed whale	1992 - 2011	70	70 ⁵	25
<i>Pseudorca crassidens</i>	False killer whale	1986 - 2017	96	70 ⁶	25
<i>Globicephala macrorhynchus</i> *	Short-finned pilot whale	1984 - 2017	200	80 ⁷	61
<i>Steno bredanensis</i> *	Rough-toothed dolphin	1983 - 2017	90	90 ⁸	37
<i>Grampus griseus</i> *	Risso's dolphin	1990 - 2017	330	80 ⁹	54
<i>Stenella frontalis</i> *	Atlantic spotted dolphin	1979 - 2015	1,557	70 ¹⁰	128

5 **Table 1.** Continuation.

Scientific name	Common name	Period	Total sightings	Filtering distance	Used sightings (n)
<i>Stenella attenuata</i> *	Pantropical spotted dolphin	1983 - 2012	800	90 ¹¹	93
<i>Stenella coeruleoalba</i> *	Striped dolphin	1992 - 2005	76	90 ¹²	35
<i>Stenella longirostris</i> *	Spinner dolphin	1983 - 2012	126	80 ¹³	41
<i>Stenella clymene</i> *	Clymene dolphin	1990 - 1998	108	70 ¹²	37
<i>Tursiops truncatus</i> *	Bottlenose dolphin	1971 - 2017	3,778	35 ¹⁴	305

6 Sources: ¹Whitehead (2018), ²McAlpine, (2018), ³Baird et al. (2009), ⁴Baird et al. (2011), ⁵Baird et al. (2012), ⁶Baird et al. (2010),
7 ⁷Olson (2018), ⁸Wells et al. (2008), ⁹Wells et al. (2009), ¹⁰Davis et al. (1996), ¹¹Scott & Chivers (2009), ¹²Gannier (1999), ¹³Perrin
8 (2018), ¹⁴Irvine et al. (1981)

Table 2 (on next page)

Statistics of each model and contribution percentages of each environmental predictor

Values of the area under the receiver operator curve (AUC) and of the 10-percentile training omission rate (OR_{10}), and percent of contribution of the environmental predictors in each model. Environmental predictors: SST_m = mean sea surface temperature, SST_{min} = minimum sea surface temperature, SST_{max} = maximum sea surface temperature, $Chl-a_m$ = mean chlorophyll-*a* concentration, $Chl-a_{min}$ = minimum chlorophyll-*a* concentration, $Chl-a_{max}$ = maximum chlorophyll-*a* concentration, D = depth, S = slope, D_{200} = distance to the 200-m isobath.

Table 2. Values of the area under the receiver operator curve (AUC) and of the 10-percentile training omission rate (OR_{10}), and percent of contribution of the environmental predictors in each model. Environmental predictors: SST_m = mean sea surface temperature, SST_{min} = minimum sea surface temperature, SST_{max} = maximum sea surface temperature, $Chl-a_m$ = mean chlorophyll-*a* concentration, $Chl-a_{min}$ = minimum chlorophyll-*a* concentration, $Chl-a_{max}$ = maximum chlorophyll-*a* concentration, D = depth, S = slope, D_{200} = distance to the 200-m isobath.

Species	AUC	OR	Environmental predictors								
			SST_m	SST_{min}	SST_{max}	$Chl-a_m$	$Chl-a_{min}$	$Chl-a_{max}$	D	S	D_{200}
Sperm whale	0.83	0.21		17.34				7.23	38.25	9.47	27.71
Dwarf sperm whale	0.86	0.28		45.57	14.59			7.96	28.81	3.06	
Cuvier's beaked whale	0.85	0.24		35.77	1.34				41.51	4.92	16.46
Short-finned pilot whale	0.83	0.18		15.25		5.47			18.00	18.76	42.53
Rough-toothed dolphin	0.83	0.13		57.93	1.10	25.45				7.45	8.07
Risso's dolphin	0.87	0.27		39.06	1.23	12.86			29.60	17.25	
Atlantic spotted dolphin	0.83	0.20		6.95		4.33			74.38	1.54	12.80
Pantropical spotted dolphin	0.74	0.25		24.87	7.60	24.83			27.68	8.59	6.44

7 **Table 2.** Continuation

Specie	AUC	OR	Environmental predictors								
			SST _m	SST _{min}	SST _{max}	Chl- <i>a</i> _m	Chl- <i>a</i> _{min}	Chl- <i>a</i> _{max}	<i>D</i>	<i>S</i>	<i>D</i> ₂₀₀
Striped dolphin	0.81	0.25	21.28		13.91	37.04			16.85	10.91	
Spinner dolphin	0.80	0.38		8.88	4.75	28.27			10.04	15.44	32.62
Clymene dolphin	0.88	0.16	44.88		18.27			18.37		14.44	4.03
Bottlenose dolphin	0.91	0.16		20.18	1.07	2.91			75.84		

8

Figure 1

Study and modeling area

(A) Geographic extension of the modeling area in the northwestern Atlantic Ocean. (B) Gulf of Mexico and physiographic sub-provinces: 1. West Florida shelf, 2. West Florida slope, 3. Mississippi-Alabama shelf, 4. Mississippi Canyon, 5. Louisiana-Texas shelf, 6. Louisiana-Texas slope, 7. Rio Grande slope, 8. Tamaulipas-Veracruz shelf, 9. Tamaulipas-Veracruz slope, 10. Bay of Campeche, 11. Bank of Campeche, 12. Yucatan shelf, 13. Campeche terrace, and 14. Sigsbee plain.





Figure 2

Habitat suitability map

Habitat suitability of (A) sperm whale, (B) dwarf sperm whale, (C) Cuvier's beaked whale, and (D) Short-finned pilot whale. In the scale bar, reds indicates high habitat suitability values (≥ 0.6) encompassed by the solid line, and light yellow indicates low habitat suitability values.




Figure 3

Habitat suitability map

Habitat suitability of (A) rough-toothed dolphin, (B) Risso's dolphin, (C) Atlantic spotted dolphin, and (D) pantropical spotted dolphin. In the scale bar, reds indicates high habitat suitability values (≥ 0.6) encompassed by the solid line, and light yellow indicates low habitat suitability values.




Figure 4

Habitat suitability map

Habitat suitability of (A) striped dolphin, (B) spinner dolphin, (C) clymene dolphin, and D) bottlenose dolphin. In the scale bar, reds indicates high habitat suitability values (≥ 0.6) encompassed by the solid line, and light yellow indicates low habitat suitability values.




Figure 5

Suitable regions map

Map of the overlap of the suitable habitat of cetaceans in the Gulf of Mexico. The solid line delimits the highly suitable regions with a high diversity of cetaceans ≥ 7 species.

