

# Genes for degradation and utilization of uronic acid-containing polysaccharides of a marine bacterium *Catenovulum* sp. CCB-QB4

Go Furusawa<sup>Corresp., 1</sup>, Nor Azura Azami<sup>1</sup>, Aik-Hong Teh<sup>1</sup>

<sup>1</sup> Centre for Chemical Biology, Universiti Sains Malaysia, Bayan Lepas, Penang, Malaysia

Corresponding Author: Go Furusawa  
Email address: furusawa@usm.my

## Background

Oligosaccharides from polysaccharides containing uronic acids are known to have many useful bioactivities. Thus, polysaccharide lyases (PLs) and glycoside hydrolases (GHs) involved in producing the oligosaccharides have attracted interest in both medical and industrial settings. The numerous polysaccharide lyases and glycoside hydrolases involved in producing the oligosaccharides were isolated from soil and marine microorganisms. Our previous report demonstrated that an agar-degrading bacterium, *Catenovulum* sp. CCB-QB4, isolated from a coastal area of Penang, Malaysia, possessed 183 glycoside hydrolases and 43 polysaccharide lyases in the genome. We expected that the strain might be degraded and use uronic acid-containing polysaccharides as a carbon source, indicating that the strain has a potential for a source of novel genes for degrading the polysaccharides.

## Methods

To confirm the expectation, the QB4 cells were cultured in artificial seawater media with uronic acid-containing polysaccharides, namely alginate, pectin (and saturated galacturonate), ulvan, and gellan gum, and the growth was observed. The genes involved in degradation and utilization of uronic acid-containing polysaccharides were explored in the QB4 genome using CAZy analysis and BlastP analysis.

## Results

The QB4 cells were capable of using these polysaccharides as a carbon source, and especially, the cells exhibited a robust growth in the presence of alginate. 28 PLs and 22 GHs related to degradation of these polysaccharides were found in the QB4 genome based on the CAZy database. 11 polysaccharide lyases and 16 glycoside hydrolases contained lipobox motif, indicating that these enzymes play an important role in degrading the polysaccharides. 14 of 28 polysaccharide lyases were classified into ulvan lyase, and the QB4 genome possessed the most abundant ulvan lyase genes in the CAZy database. Besides, genes involved in uronic acid metabolisms were also present in the genome. These results were consistent with cell growth. In the pectin metabolic pathway, the strain had genes for three different pathways. However, the growth experiment using saturated galacturonate exhibited that the strain can only use the pathway related to unsaturated galacturonate.

**Genes for degradation and utilization of uronic acid-containing polysaccharides of a  
marine bacterium *Catenovulum* sp. CCB-QB4**

**Go Furusawa\* Nor Azula Binti Azami, and Aik-Hong Teh**

Author's institutional affiliation:

Centre for Chemical Biology, Universiti Sains Malaysia, 10 Persiaran Bukit Jambul, 11900  
Bayan Lepas, Penang, Malaysia

Contact information;

Go Furusawa

Centre for Chemical Biology, Universiti Sains Malaysia, 10 Persiaran Bukit Jambul, 11900  
Bayan Lepas, Penang, Malaysia

E-mail: [furusawa@usm.edu.my](mailto:furusawa@usm.edu.my)

## Abstract

## Background

Oligosaccharides from polysaccharides containing uronic acids are known to have many useful bioactivities. Thus, polysaccharide lyases (PLs) and glycoside hydrolases (GHs) involved in producing the oligosaccharides have attracted interest in both medical and industrial settings. The numerous polysaccharide lyases and glycoside hydrolases involved in producing the oligosaccharides were isolated from soil and marine microorganisms. Our previous report demonstrated that an agar-degrading bacterium, *Catenovulum* sp. CCB-QB4, isolated from a coastal area of Penang, Malaysia, possessed 183 glycoside hydrolases and 43 polysaccharide lyases in the genome. We expected that the strain might be degraded and use uronic acid-containing polysaccharides as a carbon source, indicating that the strain has a potential for a source of novel genes for degrading the polysaccharides.

## Methods

To confirm the expectation, the QB4 cells were cultured in artificial seawater media with uronic acid-containing polysaccharides, namely alginate, pectin (and saturated galacturonate), ulvan, and gellan gum, and the growth was observed. The genes involved in degradation and utilization of uronic acid-containing polysaccharides were explored in the QB4 genome using CAZy analysis and BlastP analysis.

## Results

The QB4 cells were capable of using these polysaccharides as a carbon source, and especially, the cells exhibited a robust growth in the presence of alginate. 28 PLs and 22 GHs related to degradation of these polysaccharides were found in the QB4 genome based on the CAZy database. 11 polysaccharide lyases and 16 glycoside hydrolases contained lipobox motif, indicating that these enzymes play an important role in degrading the polysaccharides. 14 of 28 polysaccharide lyases were classified into ulvan lyase, and the QB4 genome possessed the most abundant ulvan lyase genes in the CAZy database. Besides, genes involved in uronic acid metabolisms were also present in the genome. These results were consistent with cell growth. In the pectin metabolic pathway, the strain had genes for three different pathways. However, the growth experiment using saturated galacturonate exhibited that the strain can only use the pathway related to unsaturated galacturonate.

# Introduction

Uronic acids are a class of sugar acids oxidized the hydroxyl group on C6 of aldoses. Uronic acids including D-glucuronic acid, D-galacturonic acid, D-mannuronic acid, L-guluronic acid, and L-iduronic acid are components of polysaccharides produced by animals (heparin), terrestrial plants (pectin), seaweed (alginate and ulvan), and bacteria (gellan gum) (de Lederkremer and Marino, 2003). In general, these polysaccharides such as pectin, alginate, ulvan, and gellan gum, have broad potential in many applications due to their excellent properties of biocompatibility, non-toxic, immunogenicity, availability, and relatively low cost (Morelli. A. and Chiellini, 2010; Venkatesan et al., 2015; Rahman et al., 2017).

Pectin is a polymer with a linear structure characterized by a backbone consisting of a few hundred to thousand D-galacturonic acid units linked together by  $\alpha$ -(1→4)-glycosidic linkages. It found in the cell walls of the plant and intracellular layer of plant cells, mainly fruits, such as apples, oranges, and lemons (Mudgil, 2017). Pectin contains a significant amount of neutral sugar, typically L-rhamnose, L-arabinose, D-galactose, D-xylose, and D-glucose which linked to the hydroxyl groups on the number 2 and 3 carbons of the main chain. Pectin attracted attention due to its gelling capabilities (Penhasi and Meidan, 2015). Pectin is widely used as a thickener and stabilizing agent in the food industry (Munarin et al., 2012).

Alginate, also known as alginic acid is an unbranched polymer composed of  $\beta$ -D-mannuronic acid (M) and  $\alpha$ -L-guluronic acid (G), which are covalently (1-4)-linked. The residues are randomly arranged into MM-, GG- and MG-blocks (Meng and Liu, 2013). Alginate is distributed widely in the cell wall of marine brown agar and has long been used in the industry such as medical field, fabric, food, and beverage industries as thickening, gel-forming, and colloidal stabilizing agents (Liakos et al., 2013; Martău et al., 2019).

Ulvan is water-soluble polysaccharides found in the cell wall of green algae (*Ulva* and *Enteromorpha*) composed mainly of 3-sulfated rhamnose (Rha3S), glucuronic acid (GlcA), iduronic acid (IdoA), and xylose (Xyl) (Kim et al., 2011). Ulvan has attracted pharmaceutical and medical applications for its anti-viral, anti-coagulant, and anti-proliferative activities towards cancer cells and its immune-stimulating properties (Alves, Sousa & Reis, 2013). In addition, ulvan also showed to be an activator of plant defense and an inducer of plant resistance (Alves, Sousa & Reis, 2013). Therefore, it is a good potential for agricultural applications.

Gellan gum is an exopolysaccharide produced from non-pathogenic, Gram-negative bacterium, *Sphingomonas elodea* (earlier *Pseudomonas elodea*) using aerobic fermentation (Vendrusculo et al., 1994). Gellan gum consisting of repeating tetrasaccharide units of glucose, glucuronic acid, and rhamnose residues in a 2:1:1 ratio: [ $\rightarrow$ 3)- $\beta$ -D-glucose-(1 $\rightarrow$ 4)- $\beta$ -D-glucuronic acid-(1 $\rightarrow$ 4)- $\beta$ -D-glucose-(1 $\rightarrow$ 4)- $\alpha$ -L-rhamnose-(1 $\rightarrow$ )] (Jansson et al., 1983). Gellan gum has been used in medicine, pharmaceutical formulations, cosmetics, or tissue engineering. As a biocompatibility polysaccharide, gellan gum is used in contact with or inside the body. Besides, gellan gum also useful in the food and biotechnology industry as immobilization of enzymes and yeast cells (Iurciuc (Tincu) et al., 2015).

These polysaccharides could also be alternative sustainable sources for fermentative biofuel production (John et al., 2011). The PLs and GHs play an important role in the saccharification of the polysaccharides in the process of biofuel production (Edwards et al., 2011; Takeda et al., 2011; Li et al., 2015). For example, endo- and exolytic alginate lyases from *Saccharophagus degradans* A1 were co-displayed on the yeast cell surface, and the co-displaying yeasts were able to effectively produce monosaccharides (Takagi et al., 2016b). On the other hand, oligosaccharides from uronic acids generated by PLs are known to have multiple biological activities. For instance, alginate oligosaccharides stimulate the growth of human endothelial (Kawada et al., 1997) and keratinocytes cells (Kawada et al., 1999). In addition, the oligosaccharides also promote the growth and root elongation of rice and barley (Tomoda et al., 1994; Hien et al., 2000). Mandalari and colleagues reported that pectin oligosaccharides (POS) have probiotic effects through the improvement of bifidobacteria and lactobacillus (Mandalari et al., 2007). Besides, POS inhibited inflammation, fibrosis formation, as well as cancer progression, transformation, and metastasis (Bonnin Garnier & Ralet, 2014). Thus, PLs and GHs have attracted considerable interest in both academic and commercial spheres.

Many bacterial species, such as the genera *Agrobacterium*, *Bacillus*, *Cellulophaga*, *Clostridium*, *Erwinia*, *Flammeovirga*, *Flavobacterium*, *Microbulbifer*, *Psuedoalteromonas*, *Pseudomonas*, *Saccharophagus*, *Sphingomonas*, *Vibrio*, *Xanthomonas*, and *Zobellia*, are known as bacteria that are capable of degrading uronic acid-containing polysaccharides, mainly alginate and pectin (Liu et al., 2019; Dubey et al., 2016; Takagi et al., 2016). The genus *Catenovulum* consisted of three species, *C. agarivorans*, *C. maritimum*, and *C. sediminis*, was also known to degrade agar (all three strains) and alginate (*C. maritimum*, and *C. sediminis*). Besides that,

*Catenovulum* sp. LP was able to produce an ulvan lyase (Li et al., 2015; Qiao et al., 2020). However, the degradation and utilization pathway of uronic acid-containing polysaccharides containing alginate, gellan gum, pectin, and ulvan by the genus *Catenovulum* are poorly understood. Our group isolated *Catenovulum* sp. CCB-QB4 (referred to hereafter as QB4) from Queens Bay of Penang, Malaysia, and the complete genome sequence was reported (Lau et al., 2019). From the study, it was reported that the QB4 genome contained 183 GHs and 43 PLs. Based on the information, this bacterium is predicted to have the ability to utilize many polysaccharides. To confirm the expectation, in this study, the cell growth of QB4 in the presence of polysaccharides containing uronic acids, namely, alginate, pectin, ulvan, and gellan gum, was confirmed using a shake flask fermentation method. In addition, genes involved in degrading and utilizing these polysaccharides were explored in the QB4 genome. This study is the first report to describe the degradation and probable metabolic pathways for utilization of four different uronic acid-containing polysaccharides in the genus *Catenovulum*.

## Materials and Methods

### Strain and chemicals

The QB4 cells (Lau et al., 2019) was cultured using high nutrient artificial seawater medium (H-ASWM) condition [0.5% tryptone, 2.4% (w/v) artificial sea salt mix (Marine Enterprises International), 10mM HEPES, pH 7.6], as reported by Furusawa and co-workers, 2015 (Furusawa et al., 2015). 0.1 % of polygalacturonic acid [pectic acid] (Nacalai Tesque), sodium alginate (Sigma-Aldrich), Gelzan<sup>TM</sup> CM (gellan gum) (Sigma-Aldrich), and ulvan were used as carbon source throughout the research.

### Purification of ulvan

The ulvan was purified according to the method described by Tabarsa et al. (2012) with slight modification. 5 g of the dry powder of *Ulva pertusa* was dissolved and stirred in 100 mL of water and stirred at 65 °C for 3 h. The mixture was cooled and centrifuged at 10,000 g for 20 min at 15 °C using Sorvall<sup>TM</sup> RC 6 Plus Centrifuge (ThermoFisher Scientific). 4 volume of cold isopropanol was added into the supernatant to precipitate crude polysaccharide. The solution was left at 4 °C

for overnight. The crude ulvan was harvested by filtration, washed with 70 % isopropanol several times, and dried overnight at 60 °C.

# **Determination of bacterial growth**

The QB4 cells were inoculated in 10 mL of H-ASWM broth and cultured overnight at 30 °C. 0.1 mL of the cell suspension was inoculated into 100 mL of H-ASWM medium with 0.2% of each uronic acid as the carbon sources. Media with and without 0.2% glucose (Fisher Scientific) were used as positive and negative controls, respectively.

Each flask was inoculated with 0.1 mL of pre-cultured bacterial cell suspension and incubated at 30 °C with an agitation speed of 200 rpm on the orbital shaker. To monitor the bacterial growth, the colony-forming unit (CFU) counts were conducted by culturing the cells on H-ASWM agar plates because the optical density measurement was not suitable for the sample with gelzan in which the broth was solidified by calcium ion contained in H-ASWM medium. 100 µl of the cell suspension from each sample was collected every 3 hours. The suspension was diluted into 900 µl of H-ASWM medium. After that, sequential 10-fold serial dilutions were made, and 100µl of aliquots of each dilution were plated on H-ASWM agar plates. Colonies were scored after incubation for 48 h at 30 °C, and the growth curve was constructed. The experiment was performed in triplicates. The generation time (G) was estimated by the method described by Aparna et al. based on the growth curve (Aparna et al., 2020). The number of generation (n) was determined by the method of  $n = (\log b - \log B)/\log 2$ , where B and b are the number of bacteria at the beginning of a time interval and at the end of the time interval, respectively. The generation time (G) was determined by the method of  $G = t/n$ , where t is the time interval (min).

To confirm the utilization of saturated galacturonate, the QB4 cells were culture in an H-ASWM medium with 0.2% saturated galacturonate. Media with and without 0.2% glucose was used as positive and negative controls, respectively. To monitor the bacterial growth, the optical density (OD<sub>600nm</sub>) was measured at 3 h intervals for 30 h using UV spectrophotometer UV-1800 (Shimadzu). The experiment was performed in triplicates.

# **Genomic analysis of *Catenovulum* sp. CCB-QB4**

The complete genome of QB4 deposited at GenBank under the accession number CP026604-CP026605 was determined by Lau et al. (Lau et al., 2019). Genes involved in the carbohydrate-

active enzyme (CAZymes) in QB4 were predicted using dbCAN pipelines (Yin et al., 2012). Several enzymes related to uronic acid metabolism in QB4 were predicted by Blastp at the National Center for Biotechnology Information (NCBI) server (Bethesda, MD, U.S.A.) and were found using the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway database (Kanehisa and Goto, 2000). Finally, the amino acid sequence similarity of all enzymes was confirmed by using Blastp analysis with Protein Data Base (PDB). Signal peptide prediction was conducted by Lipop 1.0 (Juncker et al., 2003) and SignalP 5.0 (Armenteros, et al., 2017). For sequence alignment of DEH reductase, the amino acid sequence of A1-R and A1-R' from *Sphingomonas* sp. A1 and DEH reductase from *Saccharophagus degradans* 2-40 were obtained from GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>). The sequence alignment with 4 sequences was conducted by ClustalW (Thompson et al., 1994) at PRABI Lyon-Gerland ([https://npsa-prabi.ibcp.fr/cgi-bin/npsa\\_automat.pl?page=/NPSA/npsa\\_clustalw.html](https://npsa-prabi.ibcp.fr/cgi-bin/npsa_automat.pl?page=/NPSA/npsa_clustalw.html)).

## Results

### Growth confirmation of QB4 using uronic acids.

To confirm the growth of QB4 in the presence of uronic acids, the cells of QB4 were cultured in H-ASWM with alginate, pectin, ulvan, and gelzan. As shown in figure 1, the cells of QB4 showed good growth in the presence of the uronic acid-containing polysaccharides and glucose except for negative control. The cells of QB4 reached the early stationary phase at 9 h in H-ASWM with alginate and gelzan. The generation time of alginate and gelzan was 76.76 and 74.07 min. However, the cell number of the sample with gelzan was 5.7 times lower than that of the sample with alginate. On the other hand, the cells cultured by pectin and glucose reached the early stationary phase at 18 h. Besides that, in the case of using ulvan, the cells reached stationary phase at 24 h, and the sample exhibited the highest cell number at 30 h in the experiment. The generation time of the samples with glucose, pectin, and ulvan was 91.37, 95.74, and 85.10 min, which were longer than that of the samples with alginate and gelzan. Thus, QB4 cells exhibited a robust growth in the presence of uronic acids. Subsequently, we focused on genes for degradation and utilization of uronic acids in the genome of QB4.

### Genes for alginate degradation and utilization



Alginate is a major polysaccharide found in the cell wall of brown algae and consisted of guluronate and mannuronate arranged as 1,4-linked polysaccharides. First, alginate is degraded into oligomeric or monomeric units by alginate lyases. Eight alginate lyase genes were found in the QB4 genome (Table 1). Ad1-PL6, Ad2-PL6 and Ad3-PL6, Ad4-PL7, Ad5-PL7, Ad6-PL7, Ad7-PL7, and Ad8-PL17 were classified into family PL6, PL7, and PL17 based on the CAZy database, respectively. As a result of Blastp search with Protein Data Bank (PDB), Ad proteins were similar to Alygc from *Paraglaciecola chathamensis* (Ad1-PL6 and Ad3-PL6), AlyQ from *Persicobacter* sp. CCB-QB2 (Ad2-PL6, Ad4-PL7, and Ad5-PL7), alginate lyase of *Klebsiella pneumoniae* (Ad6-PL7 and Ad7-PL7) and *Saccharophagus degradans* 2-40 (Ad8-PL17) (Table S1). LipoP 1.0 (Juncker et al., 2003) was used to predict the signal peptides and their type of each enzyme. Table 1 demonstrated that Ad1-PL6, Ad2-PL6, Ad4-PL7 and Ad5-PL7 possessed type I signal peptide, which was cleaved by signal peptidase I, and Ad6-PL7, Ad7-PL7 and Ad8-PL17 have type II lipoprotein signal peptide which was cleaved by signal peptidase II. The lipoprotein signal peptide referred to as “lipobox” plays an important role in anchoring the protein on the outer surface of the cell membrane after secretion and modification of N-terminal cysteine residue (Pugsley et al., 1986; Hutcheson et al., 2011). This suggested that Ad6-PL7, Ad7-PL7, and Ad8-PL17 may localize on the cell surface. In contrast, Ad1-PL6, Ad2-PL6, Ad4-PL7, and Ad5-PL7 may be released into the culture medium. On the other hand, Ad3-PL6, which does not possess any signal peptides, may locate at the cytoplasm.

Takase et al. (Takase et al., 2010) described that an exotype oligoalginate lyase, A1-IV, from *Sphingomonas* sp. A1 degraded oligoalginates into monosaccharides, which are then nonenzymatically converted to 4-deoxy-L-erythro-5-hexoseulose uronic acid (DEH). Takagi and colleagues also reported that an alginate lyase, Alg7K, from *Saccharophagus degradans* showed exolytic activity and produced monosaccharides from oligoalginates (Takagi et al., 2016a). The amino acid sequence of the alginate lyase, Ad6-PL7, showed high similarity (77.3 %) to that of Alg7K, suggesting that Ad6-PL7 might mediate hydrolysis of oligoalginate to produce monomers. The DEH is converted to 2-keto-3-deoxy-D-gluconate (KDG) by NADH or NADPH-dependent DEH reductase (Takase et al., 2010; Takase et al., 2014; Kim et al., 2016). A result of the BLASTp search demonstrated that the SDR family oxidoreductase, Au1, found in the QB4 genome (Table 1) showed high similarity to NADH (A1-R', 60.9 %) and NADPH (A1-R, 50.8 %)-dependent DFH reductase. In addition, the gene alignments of DEH reductases showed that the

TGXXXGX motif and catalytic triad (Ser, Tyr, and Lys) are highly conserved in SDR family enzymes (Takase et al., 2014) were conserved in the SDR family oxidoreductase (Fig. S1). These results indicated that Au1 might function as DEH reductase. KDG kinase catalyzes the conversion of KDG to 2-keto-3-deoxy-phosphogluconate (KDGP), and then KDGP is converted into D-glyceraldehyde-3-phosphate and pyruvate by 2-dehydro-3-deoxy-phosphogluconate aldolase via Entner -Doudoroff pathway. Two of the genes encoding the KDG kinase, Au2 and Au3, and the aldolase, Au4, and Au5, were found in the genome of QB4 (Table 1). Finally, pyruvate produced by the pathway goes into further metabolic pathways for generating energy. The five Au proteins also demonstrated a high degree of amino acid sequence similarity (>50 %) to DEH reductase A1-R' from *Sphingomonas* sp. A1 (Au1), KDG kinase *Shigella flexneri* (Au2 and 3), 2-keto-3-deoxy-6-phosphogluconate aldolase from *Thermotoga maritima* (Au4), and KdpG aldolase from *Escherichia coli* (Au5) (Table S1). As mentioned above, QB4 possessed many alginate lyases and enzymes responsible for alginate utilization. This is consistent with robust growth of QB4 in the presence of alginate, as shown in Fig. 1.

## Genes for pectin degradation and utilization

As a first step, pectin is depolymerized by pectin lyases or polygalacturonases. Pectin and pectate lyases are classified into five families of PLs. Although pectin lyases attack highly methyl-esterified pectin, pectate lyases specifically attack non-methylated polygalacturonate or methylated pectin with a very low degree (Hugouvieux-Cotte-Pattat et al., 2014). These enzymes degrade pectin to unsaturated pectic-oligosaccharides and disaccharides.

In the CAZy database, seven pectate lyases belonging to family PL1 (Pd1\_PL1, Pd2\_PL1, Pd3\_PL1, Pd4\_PL1, and Pd5\_PL1), PL3 (Pd6\_PL3), and PL10 (Pd7\_PL10) family pectate lyases were found in the genome of QB4. The result of Blastp search using PDB database on Pd proteins also demonstrated that these Pd proteins were similar (>45 %) to pectate lyase (Pd1\_PL1, Pd2\_PL1, Pd3\_PL1, Pd5\_PL1, Pd6\_PL3, and Pd7\_PL10) and pectinesterase (Pd4\_PL1) from other bacterial species (Table S2). CBM13 found in Pd1\_PL1 and Pd6\_PL3 and CBM35 found in Pd2\_PL1 and Pd3\_PL1 were capable of binding to multi-ligands, such as arabinan, arabinoxylan, and pectin (Fujimoto, 2013; Dhillon et al., 2018). However, CBM77 contained in Pd2\_PL1 recognized homogalacturonan (Fujimoto 2013). Pectin methylesterase (PME) domain from family 8 Carbohydrate Esterase (CE8) was found in Pd4\_PL1 and

Pd5\_PL1. PMEs produce de-esterified homogalacturonan by catalyzing the de-esterification of the methoxyl group of the pectin; as a result, the products are effectively degraded by pectate lyases (Kashyap et al., 2001). Besides, although Pd1\_PL1, Pd3\_PL1, Pd4\_PL1, and Pd7\_PL10 possessed type I signal peptide, Pd5\_PL1 and Pd6\_PL3 had lipobox, indicating that both two lyases localize on the cell surface. Hence, these results suggested that Pd5\_PL1 plays an important role in the pectin degradation of QB4 due to its de-esterification and depolymerization activities and its localization.

The unsaturated disaccharides generated by the pectate lyases are converted to 5-keto-4-deoxyuronate (DKI) by two distinct processes. One is that the disaccharides are degraded by oligogalacturonate lyase belonging to family PL22, and then the product,  $\Delta$ -4,5-unsaturated galacturonate, is linearized into DKI by KdgF (Hobbs et al., 2016). The other is that unsaturated galacturonyl hydrolases belonging to GH105 degrades the disaccharides and directly releases DKI (Hobbs et al., 2019). In the QB4 genome, no PL22 enzymes were found in the genome. However, three GH105 proteins, Pd8\_GH105, Pd9\_GH105, and Pd10\_GH105, were present in the genome (Table 2). These enzymes showed a high degree of similarity (49 ~ 67 %) on unsaturated rhamnogalacturonyl hydrolases (YteR) based on Blastp search with PDB database. These results indicated that oligogalacturonate might directly convert to DKI by GH105 proteins in QB4. DKI is converted to 2-keto-3-deoxygluconate (KDG) by two enzymes, a DKI isomerase (KduI) and a 2-dehydro-3-deoxy-D-gluconate 5-dehydrogenase (KduD) (Hobbs et al., 2019). The pathway analysis based on KEGG showed that four of KduI (Pu1, Pu2, Pu3, and Pu4) and KduD (Pu5, Pu6, Pu7, and Pu8) were present in the genome of QB4. Blastp search with PDB database also exhibited that these enzymes showed a high degree of similarity (63 ~ 71 %) on KduD and KduI from *Enterococcus faecalis*, *Escherichia coli*, and a pectolytic bacterium, *Pectobacterium carotovorum* (Table S2). These results indicated that QB4 use pectin as a carbon source for cell growth and is consistent with the result shown in Fig. 1.

On the other hand, the pathway analysis also showed the other pathway using saturated galacturonate (monosaccharide) produced by polygalacturonases. As mentioned above, although QB4 possesses seven pectate lyases, only two polygalacturonases (GH28), Pd11\_GH28 and Pd12\_GH28, which were highly similar (>66 %) to polygalacturonase from pectolytic bacteria, such as *Erwinia carotovora* and *Thermotoga maritima* (Table S2), were found in the genome. Based on SignalP 5.0 analysis, it was found that Pd11\_GH28 and Pd12\_GH28 contained Tat and

type I signal peptide, respectively. Saturated galacturonate is converted to D.tagaturonate by glucuronate isomerase. Three glucuronate isomerases, Pu9, Pu10, and Pu11, which were highly similar (> %) to glucuronate isomerase of *Salmonella enterica* subsp. *enterica* serovar Typhimurium and uronate isomerase of *Caulobacter vibrioides* CB15 (Table S2), were found in the QB4 genome. The conversion of D.tagaturonate to KDG occurs through two distinctly different pathways. One is that the process consists of two steps involving tagaturonate reductase and altronate hydrolase and one intermediate, D-altronate (Richard and Hilditch, 2009). The other is that tagaturonate is converted via three steps involving tagaturonate epimerase, fructuronate reductase, and mannonate dehydratase and two intermediates, D-fructuronate and D-mannonate (Valk et al., 2020). As shown in Table 2, QB4 possessed all genes involved in the two different pathways, namely tagaturonate reductase (Pu12), altronate hydrolase (Pu13), tagaturonate epimerase (Pu14), fructuronate reductase (Pu15), and mannoate dehydratase (Pu16). Homologous enzymes of these proteins were also found in the PDB database (Table S2). This result suggested that QB4 is also capable of using saturated galacturonate as a carbon source. To confirm which pathway is functioning when QB4 cells grew in H-ASWM broth with polygalacturonic acid, the QB4 cells were cultured in H-ASWM broth with polygalacturonic acid or saturated galacturonate. As shown in Figure 2, although the cells with glucose demonstrated robust growth, the cells with were unable to grow in the broth with saturated galacturonate as well as the negative control in the incubation period. This result indicated that the unsaturated galacturonate utilization pathway is the main pathway to utilize polygalacturonic acid of QB4. Interestingly, genes for all pectate lyases, polygalacturonases, and one set of KduI (Pu4), KduD (Pu8), glucuronate isomerase (Pu11), and tegaturonate reductase were located in a plasmid ([https://www.ncbi.nlm.nih.gov/nuccore/NZ\\_CP026605.1](https://www.ncbi.nlm.nih.gov/nuccore/NZ_CP026605.1)) found in QB4 (Table 2).

### **Genes for ulvan degradation and utilization**

Ulvan lyases classified into 5 families, PL24, PL25, PL28, PL 37, and PL40 in the CAZy database (Li et al., 2020) were isolated from several marine bacteria, such as genera, *Alteromonas*, *Pseudoalteromonas*, *Formosa*, and *Nonlabens* (Kopel et al., 2016; Qin et al., 2018; Ulaganathan et al., 2018a; Ulaganathan et al., 2018b; Reisky et al., 2019). Ulvan lyases cleave between L-rhamnose 3-sulfate (Rha3S) and D-glucuronic acid (GlcA) or L-iduronic acid (IdoA). Fourteen ulvan lyases in QB4 were classified into three PLs families, PL24 (Ud1\_PL24,

Ud2\_PL24, Ud3\_PL24, Ud4\_PL24, Ud5\_PL24, Ud6\_PL24, Ud7\_PL24, Ud8\_PL24, and Ud9\_PL24), PL25 (Ud10\_PL25, Ud11\_PL25, Ud12\_PL25, and Ud13\_PL25) and PL40 (Ud14\_PL40). Ud proteins belonging to the PL24 family were highly similar (>62 %) to short ulvan lyase of *Alteromonas* sp. LOR, and Ud proteins belonging to the PL25 family were highly similar (>57 %) to ulvan lyase-PL25 of *Pseudoalteromonas* sp. PLSV (Table S3). The number of the ulvan lyases was obviously higher than that of *F. agariphila* (three PL40 proteins) (Reisky et al., 2019) and *Alteromonas* sp. LOR (one PL24 and one nonclassified protein) (Foran et al., 2017). Ud6\_PL24 and Ud9\_PL24 and the enzymes belonging to PL25 possessed lipobox. Although Ud2\_PL24, Ud3\_PL24, Ud4\_PL24, Ud5\_PL24, and Ud8\_PL24 contained type I signal peptide, the remaining enzymes did not have any signal peptides. Although only one pectin lyase had a lipobox, many ulvan lyases possessed lipobox as well as alginate lyases, indicating that alginate and ulvan degradation activity of QB4 may be more effective than pectin degradation of the strain.

The unsaturated uronyl residue at the non-reducing end of the oligosaccharides produced by the ulvan lyases may be released by unsaturated glucuronyl hydrolases (GH105), because five putative unsaturated glucuronyl hydrolases, Ud15\_GH105, Ud16\_GH105, Ud17\_GH105, Ud18\_GH105, and Ud19\_GH105 were found in the QB4 genome and showed similarity on unsaturated 3s-rhamnoglycuronyl hydrolase of *Nonlabens ulvanivorans*, which is known as a ulvan-degrading bacterium, based on BLASTp search (Table 3; Table S3) (Collén et al., 2014). After forming DKI by the enzymes, the following process may be the same as pectin utilization. In the ulvan-degrading process of *F. agariphila*, Rha3S-Xyl-Rha3S was also the main product by ulvan lyases (Reisky et al., 2019). First, Rha3S at the non-reducing end is desulfated by sulfatases. The amino acid sequences of two sulfatases, Ulu1 and Ulu2, in the QB4 genome showed high similarity (77.3 and 86.2 %) to the sulfatases, WP.032096151.1 and WP.632096147.1, located into ulvan utilization loci of *Alteromonas* sp. LOR (Foran et al., 2017), indicating that the two sulfatases may involve desulfation of the Rha3S. The Rha3 will remove the Rha3-Xyl-Rha3S by  $\alpha$ -L-rhamnosidase. Five  $\alpha$ -L-rhamnosidases classified into GH78 family, Ud20\_GH78, Ud21\_GH78, Ud22\_GH78, Ud23\_GH78, and Ud24\_GH78, were found in the QB4 genome and showed similarity on rhamnosidase from *Bacillus* sp. GL1, *Dictyoglomus thermophilum*, *Streptomyces avermitilis*, and *Bacteroides thetaiotaomicron* (Table S3). Ud21\_GH78, Ud22\_GH78, Ud23\_GH78, and Ud24\_GH78 contained lipobox, suggesting that

these enzymes are located on the cell surface. In addition, Ud20\_GH78 and Ud22\_GH78 possessed the CBM67 domain, which binds L-rhamnose in a calcium-dependent manner (Fujimoto et al., 2013), suggesting that the two  $\alpha$ -L-rhamnosidases are the main components to generate Rha3.

Rha3 metabolic pathway was described by Reisky et al., 2019.  $\alpha$ -L-rhamnose is converted to  $\beta$ -L-rhamnose by L-rhamnose mutarotase. Next, isomerization of the  $\beta$ -L-rhamnose to L-rhamnulose is catalyzed by rhamnose isomerase. The product is converted to L-rhamnulose-1-phosphate by pentulose/hexulose kinase (rhamnulokinase), and subsequently, the L-rhamnulose-1-phosphate is cleaved into L-lactaldehyde and dihydroxyacetone phosphate by rhamnulose aldolase. Finally, the L-lactaldehyde is converted to pyruvate by aldehyde dehydrogenase and lactate dehydrogenase. Table 3 displayed that the corresponding genes (Ulu3, Ulu4, Ulu5, Ulu6, Ulu7, and Ulu8) were found in the QB4 genome, and these proteins showed similarity on corresponding proteins in the PDB database (Table S3). This result indicated that Rha3 is metabolized by QB4 cells. The genes involving the metabolic pathway downstream of the mutarotation in *F. agariphila* form a gene cluster (Reisky et al., 2019). However, the genes in QB4 were randomly distributed in the genome.

Xyl will be released by  $\beta$ -xylosidase classified into GH3 and GH43 families. Based on the CAZy database, seven GH43 (Ud25\_GH43, Ud26\_GH43, Ud27\_GH43, Ud28\_GH43, Ud29\_GH43, Ud30\_GH43, and Ud31\_GH43) proteins were found in the QB1 genome while *F. agariphila* possessed two each of GH3 and GH43 proteins (Reisky et al., 2019). Ud25\_GH43, Ud29\_GH43, and Ud31\_GH43 were similar to glycoside hydrolases from *Zobellia galactanivorans* (50.00 %) and *Halothermothrix orenii* H 168 (45.75 and 39.10 %) (Table S3). The D-xylose is converted to D-xylulose-5-phosphate via xylose isomerase and xylulose kinase, and then the product is passed to the pentose phosphate pathway. Two genes encoding xylose isomerase (Ulu9) and xylulose kinase (Ulu10) were found in the QB4 genome based on RAST server annotation and formed a gene cluster with transcription repressor, *xyIR* (Ulu11) (Table 3). These results suggested that QB4 cells may use not only glucuronic acid but also L-rhamnose and D-xylose as carbon sources. In addition, the presence of numerous genes encoding ulvan lyase may promote strong ulvan degradation and its utilization. Actually, the QB4 cells exhibited a robust cell growth in the presence of ulvan as well as that in the presence of alginate, as shown in Figure 1.

## Genes for gellan gum degradation and utilization

Gellan gum is depolymerized gellan lyases classified into PL33 family proteins. Gellan lyases are found in several bacterial species, such as *Bacillus* sp. GL1, *Geobacillus* stearothermophilus 98, and *Opitutaceae bacterium* TAV5 (Hashimoto et al., 1997; Derekova et al., 2006; Helbert et al., 2019). Gellan gum is degraded to tetrasaccharides composed of [ $\rightarrow$ 3)- $\beta$ -D-glucose-(1 $\rightarrow$ 4)- $\beta$ -D-glucuronic acid-(1 $\rightarrow$ 4)- $\beta$ -D-glucose-(1 $\rightarrow$ 4)- $\alpha$ -L-rhamnose-(1 $\rightarrow$ ] by gellan lyases. The tetrasaccharides are completely degraded to monosaccharides by  $\beta$ -D-glucosidase (GH1, GH2, GH3, GH5, GH9, GH16, GH30, GH39, and GH116), unsaturated glucuronyl hydrolase (GH88 and GH105), and  $\alpha$ -L-rhamnosidase (GH78) (Hashimoto et al., 2003).

Figure 1 demonstrated that QB4 might also use gellan gum as a carbon source for its growth. However, gellan lyases were not found in the genome. As shown in Table 3, QB4 possessed five unsaturated glucuronyl hydrolases and five  $\alpha$ -L-rhamnosidases (Table 3). The hydrolysates, DEH, and  $\alpha$ -L-rhamnose, may be metabolized by the pathway described above. In addition, one  $\beta$ -D-glucosidase classified into the GH1 family (Gd1-GH1) was also found in the genome, and Gd1-GH1 showed high similarity (63.23 %) on  $\beta$ -glucosidase A of *Hungateiclostridium thermocellum* in the PDB database (Table 4; Table S3).  $\beta$ -D-glucose produced by the  $\beta$ -D-glucosidases is converted to  $\beta$ -D-glucose-6-phosphate by glucokinase, and then the product is converted to  $\beta$ -D-fructose-6-phosphate, which is an intermediate of glycolysis, by glucose-6-phosphate isomerase. One of these genes, Gu1 and Gu2, were found in the QB4 genome (Table 4) and were similar on glucokinase and glucose-6-phosphate Isomerase of *E. coli* in PDB database, respectively (Table S4), indicating that  $\beta$ -D-glucose is converted to  $\beta$ -D-fructose-6-phosphate and is metabolized by glycolysis.

## Discussion

Polysaccharides from plants, seaweed, and bacteria, especially their oligosaccharides, have attracted considerable attention due to various biological activities. Thus, polysaccharide lyases and glycosyl hydrolases were isolated from microorganisms, including fungi and bacteria. The genus *Catenovulum* have been known as bacteria that can degrade agar and polysaccharide containing uronic acids, such as alginate, pectin, and ulvan (Cui et al., 2014; Li et al., 2015; Lee et al., 2019; Liu et al., 2019). Our previous report demonstrated that many genes encoding polysaccharide degrading enzymes were present in the QB4 genome (Lau et al., 2019). In this

study, expectedly, QB4 cells were able to degrade and used the polysaccharides as a carbon source for their growth (Fig. 1). Especially, the QB4 cells exhibited a robust growth in the presence of alginate and ulvan, suggesting that polysaccharides from marine algae are effectively used by QB4 cells rather than that of plants and bacteria. The growth profile of QB4 was different from *S. degradans* of which the cells reached the early stationary phase at 9 h with alginate, glucose, and pectin (Takagi et al., 2016a).

Three pathways of pectin metabolisms were known in bacteria. One is the polygalacturonate pathway metabolized unsaturated disaccharides. The pathway was found in bacteria, such as *Escherichia coli* and phytopathogenic enterobacterium, *Erwinia chrysanthemi* (Chatterjee et al., 1985; Richard and Hilditch 2009). The other two pathways metabolize saturated galacturonate through uronate isomerase (isomerase pathway) (Richard and Hilditch, 2009) or tagaturonate epimerase (epimerase pathway) (Rodionova et al., 2012). Table 2 displayed that the QB4 possessed genes involved in the three pathways. However, QB4 cells did not grow in the presence of saturated galacturonate (Fig. 2). Although a pectin-degrading marine bacterium, *Pseudoalteromonas* sp. PS47, also has genes for the epimerase pathway, the strain was unable to grow on saturated galacturonate (Hobbs et al., 2019). Hobbs and colleagues proposed that GH28s of the strain are periplasmic proteins and, thus, that saturated galacturonates would be produced in the periplasm (Hobbs et al., 2019). It was suggested that the strain might be unable to transport extracellular saturated galacturonate. On the other hand, although Pd11\_GH28 containing Tat signal peptide would be exported into periplasm through Tat (for twin-arginine translocation) system (Stanley et al., 2000), Pd12\_GH28 would be secreted to the outside of the cells via Sec system, suggesting that saturated galacturonate might be generated in the culture broth in contrast to the strain PS47. However, the QB4 did not grow using saturated galacturonate. Thus, it seems like the BQ4 cells are also unable to transport extracellular saturated galacturonate as well as the strain PS47. In other cases, some investigators reported that the isomerase pathway is not crucial for utilizing pectin in *Dickeya dadantii* (formerly *E. chrysanthemi*) (Hugouvieux-Cotte-Pattat et al., 1996; Pédrón et al., 2018). It was known that *D. dadantii* belonging to the class Gammaproteobacteria, which is a plant pathogen and pectinolytic bacterium, also possessed the polygalacturonate pathway and the isomerase pathway (Hugouvieux-Cotte-Pattat et al., 1996). Even though the enzyme production of the isomerase pathway was not impaired by a mutation in genes of the polygalacturonate pathway (*kduD* and



*kduI*), these mutants did not exhibit their growth on polygalacturonate (Hugouvieux-Cotte-Pattat et al., 1996). In addition, the transcriptomic analysis of *D. dadantii* during the early colonization on the plant leaf demonstrated that genes involved in the polygalacturonate pathway were upregulated in the condition (Pédrón et al., 2018). These results suggested that the isomerase pathway was not the main pathway for utilizing pectin in the bacterium. The third pathway, epimerase pathway, involving the conversion of D-tagatose to D-fructuronate by tagatose epimerase (UxaE), was found in the hyperthermophilic bacterium, *Thermotoga maritima*, belonging to the phylum Thermotoga. However, the pathway was not found in other bacteria, such as *Escherichia* and *Bacillus* (Kuivanen et al., 2019). *uxaE* and other epimerase pathway genes, such as hexuronate catabolism regulator (*uxaR*), fructuronate reductase (*uxaD*), predicted D-mannonate utilization enzyme (*gntE*) constituted a regulon regulated by the GntR-like transcription factor UxaR in the genus *Thermotoga* (Rodionova et al., 2012). A regulon is a gene cluster or operon that is regulated by the same regulatory protein. However, Table 2 demonstrated that genes encoding Pu14, Pu15, and Pu16 were scattered in the chromosome and plasmid, suggesting that Pu14, Pu15, and Pu16 might not constitute a regulon. In addition, figure 2 demonstrated that the saturated galacturonate, which is a substrate of the epimerase pathway, was not used by QB4 cells. These results suggested that the epimerase pathway might not function in QB4.

Figure 1 demonstrated that the QB4 cells exhibited a robust growth using ulvan as a carbon source. Qiao et al. (2020) fermented *Catenovulum* sp. LP using the shaking-flask method containing 1.2 % purified ulvan, of which the concentration was 6 times-higher than that of the QB4. The bacterial culture reached the stationary phase at 36 h incubation period (Qiao et al., 2020) that was quite slower than that of the QB4. Table S5 showed that the QB4 genome contained 14 ulvan lyase genes, which is the largest number compared to *Siansivirga zeaxanthinifaciens* CC-SAMT-1 (8 genes) *Tamlana* sp. UJ94 (7 genes), *Polaribacter* sp. BM10 (7 genes), and *Wenyingzhunangia fucanilytica* CZ1127 (7 genes) in the CAZy database (Table S5). The numerous genes in the QB4 genome may be crucial to efficiently degrade ulvan and to stimulate robust growth.

Figure 1 demonstrated that the QB4 cells were capable of using gelzan as a carbon source, however, no gallan lyases found in the QB4 genome. It was known that *Bacillus* sp. GL1 and *G. stearothermophilus* 98, which possessed all genes for gellan gum degradation and utilization,

were able to use gellan gum as a sole carbon source (Hashimoto et al., 1998; Derekova et al., 2006). On the other hand, although *Paludisphaera borealis* PX4 belonging to the order *Planctomycetales* does not have any gellan lyases, the bacterium was capable of degrading gellan gum (Ivanova et al., 2017). The authors suggested that an unsaturated glucuronyl hydrolase and two  $\alpha$ -L-rhamnosidases of the bacterium are involving degrading gellan gum. Tables 3 and 4 showed that QB4 cells possess five unsaturated glucuronyl hydrolases, five  $\alpha$ -L-rhamnosidases, and one  $\beta$ -D-glucosidase. We expected that these hydrolases play a critical role in degrading gellan gum in QB4 cells. On the other hand, the lack of gellan lyases may cause lower degradation efficiency. Thus, the cell growth with gellan gum was not robust compared to those with other uronic acids.

## Conclusion

The agar-degrading bacterium, *Catenovulum* sp. CCB-QB4, used uronic acids, including alginate, pectin ulvan, and gellan gum as carbon sources for its growth. Especially, QB4 cells exhibited a robust cell growth in the presence of alginate and ulvan from seaweed. In gene analysis based on the CAZy database, a large number of polysaccharide lyases and hydrolases involved in degrading these uronic acids were found in the QB4 genome. Many alginate lyases, ulvan lyases, unsaturated glucuronyl hydrolases,  $\alpha$ -L-rhamnosidase, and  $\beta$ -xylosidase contained lipobox, indicating that QB4 cells can effectively degrade alginate and ulvan and uptake the oligosaccharides. Of course, genes for metabolizing the uronic acids were also present in the QB4. These results were suggested that QB4 will become a source of novel uronic acid degradation enzymes.

## Conflict of interest

The authors do not have any conflict of interest in relation to the work described.

# Reference

- Alves, A., Sousa, R.A., and Reis, R.L. 2013. A practical perspective on ulvan extracted from green algae. *J. Appl. Phycol.* **25**, 407-424.
- Aparna, S., Parvathi, A., and Kaniyassery, A. 2020. Isolation and characterization of a moderately halophilic *Marinobacter* phage-host system from the Arabian Sea. *Environ. Monit. Assess.* 192, 199
- Armenteros, J.J.A., Tsirigos, K.D., Sønderby, C.K., Petersen, T.N., Winther, O., Brunak, S., von Heijne, G., and Nielsen, H. 2019. SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nat. Biotechnol.* **37**, 420-423.
- Bonnin, E., Garnier, C., and Ralet, M.-C. 2014. Pectin-modifying enzymes and pectin-derived materials: applications and impacts. *Appl. Microbiol. Biotechnol.* **98**, 519-532.
- Chatterjee, A.K., Thurn, K.K., and Tyrell, D.J. 1985. Isolation and characterization of Tn5 insertion mutants of *Erwinia chrysanthemi* that are deficient in polygalacturonate catabolic enzymes oligogalacturonate lyase and 3-deoxy-D-glycero-2,5-hexodiulosonate dehydrogenase. *J. Bacteriol.* **162**, 708-714.
- Collén, P.N., Jeudy, A., Sassi, J.-F., Groisillier, A., Czjzek, M., Coutinho, P.M., and Helbert, W. 2014. A novel unsaturated  $\beta$ -glucuronyl hydrolase involved in ulvan degradation unveils the versatility of stereochemistry requirements in family GH105. *J. Biol. Chem.* **289**, 6199-6211.
- Cui, F., Dong, S., Shi, X., Zhao, X., and Zhang, X.-H. 2014. Overexpression and characterization of a novel thermostable  $\beta$ -agarase YM01-3, from marine bacterium *Catenovulum agarivorans* YM01T. *Mar. Drugs* **12**, 2731-2747.
- de Lederkremer, R.M., and Marino, C. 2003. Acids and Other Products of Oxidation of Sugars. *Adv. Carbohydr. Chem. Biochem.* **58**, 199-306.

557 Derekova, A., Sjöholm, C., Mandeva, R., Michailova, L., and Kambourova, M. 2006. Biosynthesis  
558 of a thermostable gellan lyase by newly isolated and characterized strain of *Geobacillus*  
559 *stearothermophilus* 98. *Extremophiles* **10**, 321-326.

560 Dhillon, A., Sharma, K., Rajulapati, V., and Goyal, A. 2018. The multi-ligand binding first family  
561 35 Carbohydrate Binding Module (CBM35) of *Clostridium thermocellum* targets  
562 rhamnogalacturonan I. *Arch. Biochem. Biophys.* **654**, 194-208.

563 Dubey, A.K., Yadav, S., Kumar, M., Anand, G., and Yadav, D. 2016. Molecular biology of  
564 microbial pectate lyase: a review. *Br. Biotechnol. J.* **13**, 1-26.

565 Edwards, M.C., Henriksen, E.D., Yomano, L.P., Gardner, B.C., Sharma, L.N., Ingram, L.O., and  
566 Peterson, J.D. 2011. Addition of genes for cellobiase and pectinolytic activity in *Escherichia coli*  
567 for fuel ethanol production from pectin-rich lignocellulosic biomass. *Appl. Environ. Microbiol.*  
568 **77**, 5184-5191.

569 Foran, E., Buravenkov, V., Kopel, M., Mizrahi, N., Shoshani, S., Helbert, W., and Banin, E. 2017.  
570 Functional characterization of a novel “ulvan utilization loci” found in *Alteromonas* sp. LOR  
571 genome. *Algal Res.* **25**, 39-46.

572 Fujimoto, Z. 2013, Structure and function of carbohydrate-binding module families 13 and 42 of  
573 glycoside hydrolases, comprising a  $\beta$ -trefoil fold. *Biosci. Biotechnol. Bioch.* **77**, 1363-1371.

574 Furusawa, G., Lau, N.-S., Shu-Chien, A.C., Jaya-Ram, A., and Amirul, A.-A.A. 2015.  
575 Identification of polyunsaturated fatty acid and diterpenoid biosynthesis pathways from draft  
576 genome of *Aureispira* sp. CCB-QB1. *Mar. Genomics* **19**, 39-44.

577 Hashimoto, W., Miyake, O., Nankai, H., and Murata, K. 2003. Molecular identification of an  $\alpha$ -L-  
578 rhamnosidase from *Bacillus* sp. strain GL1 as an enzyme involved in complete metabolism of  
579 gellan. *Arch. Biochem Biophys.* **415**, 235-244.

580 Hashimoto, W., Miki, H., Nankai, H., Sato, N., Kawai, S., and Murata, K. 1998. Molecular Cloning  
581 of Two Genes for  $\beta$ -d-Glucosidase in *Bacillus* sp. GL1 and Identification of One as a Gellan-  
582 Degrading Enzyme. *Arch. Biochem Biophys.* **360**, 1-9.

583 Hashimoto, W., Maesaka, K., Sato, N., Kimura, S., Yamamoto, K., Kumagai, H., and Murata, K.  
584 1997. Microbial System for Polysaccharide Depolymerization: Enzymatic Route for Gellan  
585 Depolymerization by *Bacillus* sp. GL1. *Arch. Biochem Biophys.* **339**: 17-23.

586 Helbert, W., Poulet, L., Drouillard, S., Mathieu, S., Loiodice, M., Couturier, M., Lombard, V.,  
587 Terrapon, N., Turchetto, J., Vincentelli, R., et al. 2019. Discovery of novel carbohydrate-active

enzymes through the rational exploration of the protein sequences space. *Proc. Natl. Acad. Sci. USA*. **116**, 6063-6068.

Hien, N.Q., Nagasawa, N., Tham, L.X., Yoshii, F., Dang, V.H., Mitomo, H., Makuuchi, K., and Kume, T. 2000. Growth-promotion of plants with depolymerized alginates by irradiation. *Radiat. Phys. Chem.* **59**, 97-101.

Hobbs, J.K., Hettle, A.G., Vickers, C., and Boraston, A.B. 2019. Biochemical reconstruction of a metabolic pathway from a marine bacterium reveals its mechanism of pectin depolymerization. *Appl. Environ. Microbiol.* **85**, e02114-02118.

Hobbs, J.K., Lee, S.M., Robb, M., Hof, F., Barr, C., Abe, K.T., Hehemann, J.-H., McLean, R., Abbott, W.A., and Boraston, A.B. 2016. KdgF, the missing link in the microbial metabolism of uronate sugars from pectin and alginate. *Proc. Natl. Acad. Sci. USA*. **113**, 6188-6193.

Hugouvieux-Cotte-Pattat, N., Condemine, G., and Shevchik, V.E. 2014. Bacterial pectate lyases, structural and functional diversity. *Environ. Microbiol. Rep.* **6**, 427-440.

Hutcheson, S.W., Zhang, H., and Suvorov, M. 2011. Carbohydrase systems of *Saccharophagus degradans* degrading marine complex polysaccharides. *Mar. Drugs* **9**, 645-665.

Iurciuc (Tincu), C.E., Lungu, C., Martin, P., and Popa, M. 2015. Gellan. pharmaceutical, medical and cosmetic applications. *Cellulose Chem. Technol.* **51**, 187-202.

Jansson, P.-E., Lindberg, B., and Sandford, P.A. 1983. Structural studies of gellan gum, an extracellular polysaccharide elaborated by *Pseudomonas elodea*. *Carbohydr. Res.* **124**, 135-139.

John, R.P., Anisha, G., Nampoothiri, K.M., and Pandey, A. 2011. Micro and macroalgal biomass: a renewable source for bioethanol. *Bioresour. Technol.* **102**, 186-193.

Juncker, A.S., Willenbrock, H., von Heijne, G., Brunak, S., Nielsen, H., and Krogh, A. 2003. Prediction of lipoprotein signal peptides in Gram-negative bacteria. *Protein Sci.* **12**, 1652-1662.

Kanehisa, M., and Goto, S. 2000. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* **28**, 27-30.

Kashyap, D.R., Vohra, P.K., Chopra, S., and Tewari, R. 2001. Applications of pectinases in the commercial sector: a review. *Bioresour. Technol.* **77**, 215-227.

Kawada, A., Hiura, N., Tajima, S., and Takahara, H. 1999. Alginate oligosaccharides stimulate VEGF-mediated growth and migration of human endothelial cells. *Arch. Dermatol. Res.* **291**, 542-547.

618 Kawada, A., Hiura, N., Shiraiwa, M., Tajima, S., Hiruma, M., Hara, K., Ishibashi A., and Takahara,  
619 H. 1997. Stimulation of human keratinocyte growth by alginate oligosaccharides, a possible  
620 co-factor for epidermal growth factor in cell culture. *FEBS lett.* **408**, 43-46.

621 Kim, D.H., Wang, D., Yun, E.J., Kim, S., Kim, S.R., and Kim, K.H. 2016. Validation of the  
622 metabolic pathway of the alginate-derived monomer in *Saccharophagus degradans* 2-40T by gas  
623 chromatography–mass spectrometry. *Process Biochem.* **51**, 1374-1379.

624 Kim, S.-K., Thomas, N.V., and Li, X. 2011. Anticancer compounds from marine macroalgae and  
625 their application as medicinal foods. *Adv. Food Nutr. Res.* **64**, 213-24.

626 Kopel, M., Helbert, W., Belnik, Y., Buravenkov, V., Herman, A., and Banin, E. 2016. New family  
627 of ulvan lyases identified in three isolates from the Alteromonadales order. *J. Biol. Chem.* **291**,  
628 5871-5878.

629 Lau, N.-S., Tan, W.-R., Furusawa, G., and Amirul, A.A. 2019. Complete genome sequence of the  
630 novel agarolytic *Catenovulum*-like strain CCB-QB4. *Mar. Genomics* **43**, 50-53.

631 Lee, C.H., Lee, C.-R., and Hong, S.K. 2019. Biochemical characterization of a novel cold-adapted  
632 agarotetraose-producing  $\alpha$ -agarase, AgaWS5, from *Catenovulum sediminis* WS1-A. *Appl.*  
633 *Microbiol. Biotechnol.* **103**, 8403–8411.

634 Li, Y., Huang, Z., Qiao, L., Gao, Y., Guan, H., Hwang, H., Aker, W.G., and Wang, P. 2015.  
635 Purification and characterization of a novel enzyme produced by *Catenovulum* sp. LP and its  
636 application in the pre-treatment to *Ulva prolifera* for bio-ethanol production. *Process Biochem.*  
637 **50**, 799-806.

638 Li, Q., Hu, F., Zhu, B., Ni, F., and Yao, Z. 2020. Insights into ulvan lyase: review of source,  
639 biochemical characteristics, structure and catalytic mechanism. *Crti. Rev. Biotechnol.* **40**, 432-441.

640 Liakos, I., Rizzello, L., Bayer, I.S., Pompa, P.P., Cingolani, R., and Athanassiou, A. 2013.  
641 Controlled antiseptic release by alginate polymer films and beads. *Carbohydr Polym.* **92**, 176-183.

642 Liu, J., Liu, Z., Jiang, C., and Mao, X. 2019. Biochemical Characterization and Substrate  
643 Degradation Mode of a Novel  $\alpha$ -Agarase from *Catenovulum agarivorans*. *J. Agric. Food Chem.*  
644 **67**, 10373-10379.

645 Liu J., Yang, S., Li, X., Yan, Q., Reaney, M.J.T., and Jiang, Z. 2019. Alginate oligosaccharides:  
646 production, biological activities, and potential applications. *Compr. Rev, Food Sci. F.* **18**, 1859-  
647 1881.

648 Mandalari, G., Palop, C.N., Tuohy, K., Gibson, G.R., Bennett, R.N., Waldron, K.W., Bisignano,  
649 G., Narbad, A., and Faulds, C.B. 2007. In vitro evaluation of the prebiotic activity of a pectic  
650 oligosaccharide-rich extract enzymatically derived from bergamot peel. *Appl. Microbiol.*  
651 *Biotechnol.* **73**, 1173-1179.

652 Martău, G.A., Mihai, M., and Vodnar, D.C. 2019. The use of chitosan, alginate, and pectin in the  
653 biomedical and food sector-biocompatibility, bioadhesiveness, and biodegradability. *Polymers* **11**,  
654 1-28.

655 Meng, S., and Liu, Y. 2013. Alginate block fractions and their effects on membrane fouling. *Water*  
656 *Res.* **47**, 6618-6627.

657 Morelli, A. and Chiellini, F. 2010. Ulvan as a new type of biomaterial from renewable resources:  
658 functionalization and hydrogel preparation. *Macromol. Chem. Phys* **211**, 821-832.

659 Mudgil, D. 2017. The Interaction Between Insoluble and Soluble Fiber. In Samaan, R.A. (ed)  
660 Dietary Fiber for the Prevention of Cardiovascular Disease, Academic Press, USA, pp. 35-59.

661 Munarin, F., Tanzi, M.C., and Petrini, P. 2012. Advances in biomedical applications of pectin gels.  
662 *Int. J. Biol. Macromol.* **51**, 681-689.

663 Pédrón, J.P., Chapelle, E., Alunni, B., and van Gijsegem, F. 2018. Transcriptome analysis of the  
664 *Dikeya dadantii* PecS regulon during the early stages of interaction with *Arabidopsis thaliana*.  
665 *Mol. Plant Pathol.* **19**, 647-663

666 Penhasi, A., and Meidan, V.M. 2015. Preparation and characterization of in situ ionic cross-linked  
667 pectin films: unique biodegradable polymers. *Carbohydr. Polym.* **15**, 254-260.

668 Pugsley, A.P., Chapon, C., and Schwartz, M. 1986. Extracellular pullulanase of *Klebsiella*  
669 *pneumoniae* is a lipoprotein. *J. Bacteriol.* **166**, 1083-1088.

670 Qiao, L., Yang, X., Xie, R., Du, C., Chi, Y., Zhang, J., and Wang, P. 2020. Efficient production of  
671 ulvan lyase from *Ulva prolifera* by *Catenovulum* sp. LP based on stage-controlled fermentation  
672 strategy. *Algal Res.* **46**, 101812.

673 Qin, H.-M., Xu, P., Guo, Q., Cheng, X., Gao, D., Sun, D., Zhu, Z., and Li, F. 2018. Biochemical  
674 characterization of a novel ulvan lyase from *Pseudoalteromonas* sp. strain PLSV. *RSC Adv.* **8**,  
675 2610-2615.

676 Qiao, L., Yang, X., Xie, R., Du, C., Chi, Y., Zhang, J., and Wang, P. 2020. Efficient production of  
677 ulvan lyase from *Ulva Prolifera* by *Catenovulum* sp. LP based on stage-controlled fermentation  
678 strategy. *Algal Res.* **46**, 101812.

679 Rahman, N., Dafader, N.C., and Banu, P. 2017. Preparation and Property Analysis of  
680 Biodegradable Packaging Film from Alginate, Starch and Citric acid. *J. Polym. Sci. Technol.* **2**,  
681 20-35.

682 Reisky, L., Prechoux, A., Zühlke, M.-K., Bäumgen, M., Robb, C.S., Gerlach, N., Roret, T.,  
683 Stanetty, C., Larocque, R., Michel, G., et al., 2019. A marine bacterial enzymatic cascade degrades  
684 the algal polysaccharide ulvan. *Nat. Chem. Biol.* **15**, 803-812.

685 Richard, P., and Hilditch, S. 2009. D-galacturonic acid catabolism in microorganisms and its  
686 biotechnological relevance. *Appl. Microbiol. Biotechnol.* **82**, 597-604.

687 Rodionova, I., Scott, D.A., Grishin, N.V., Osterman, A.L., and Rodionov, D.A. 2012.  
688 Tagaturonate-fructuronate epimerase UxaE, a novel enzyme in the hexuronate catabolic network  
689 in *Thermotoga maritima*. *Environ. Microbiol.* **14**, 2920-2934.

690 Stanley, N.R., Palmer, T., and Berks, B.C. 2000. The twin arginine consensus motif of Tat signal  
691 peptides is involved in Sec-independent protein targeting in *Escherichia coli*. *J. Biol. Chem.* **275**,  
692 11591-11596.

693 Tabarsa, M., Han, J.H., Kim C.Y., and You, S.G. 2012. Molecular characteristics and  
694 immunomodulatory activities of water-soluble sulfated polysaccharides from *Ulva pertusa*. *J.*  
695 *Med. Food* **15**, 135-144.

696 Takagi, T., Morisaka, H., Aburaya, S., Tatsukami, Y., Kuroda, K., and Ueda, M. 2016a. Putative  
697 alginate assimilation process of the marine bacterium *Saccharophagus degradans* 2-40 based on  
698 quantitative proteomic analysis. *Mar. Biotechnol.* **18**, 15-23.

699 Takagi, T., Yokoi, T., Shibata, T., Morisaka, H., Kuroda, K., and Ueda, M. 2016b. Engineered  
700 yeast whole-cell biocatalyst for direct degradation of alginate from macroalgae and production of  
701 non-commercialized useful monosaccharide from alginate. *Appl. Microbiol. Biotechnol.* **100**,  
702 1723-1732.

703 Takase, R., Ochiai, A., Mikami, B., Hashimoto, W., and Murata, K. 2010. Molecular identification  
704 of unsaturated uronate reductase prerequisite for alginate metabolism in *Sphingomonas* sp. A1.  
705 *Biochim Biophys Acta.* **1804**, 1925-1936.

706 Takase, R., Mikami, B., Kawai, S., Murata, K., and Hashimoto, W. 2014. Structure-based  
707 conversion of the coenzyme requirement of a short-chain dehydrogenase/reductase involved in  
708 bacterial alginate metabolism. *J. Biol. Chem.* **289**, 33198-33214.



- 709 Takeda, H., Yoneyama, F., Kawai, S., Hashimoto, W., and Murata, K. 2011. Bioethanol  
710 production from marine biomass alginate by metabolically engineered bacteria. *Energy Environ.*  
711 *Sci.* **4**, 2575-2581.
- 712 Thompson, J.D., Higgins, D.G., and Gibson, T.J. 1994. CLUSTAL W: improving the sensitivity  
713 of progressive multiple sequence alignment through sequence weighting, position-specific gap  
714 penalties and weight matrix choice. *Nucleic Acids Res.* **22**, 4673-4680.
- 715 Tomoda, Y., Umemura, K., and Adachi, T. 1994. Promotion of barley root elongation under  
716 hypoxic conditions by alginate lyase-lysate (ALL). *Biosci. Biotechnol. Biochem.* **58**, 202-203.
- 717 Ulaganathan, T., Banin, E., Helbert, W., and Cygler, M. 2018a. Structural and functional  
718 characterization of PL28 family ulvan lyase NLR48 from *Nonlabens ulvanivorans*. *J. Biol. Chem.*  
719 **293**, 11564-11573.
- 720 Ulaganathan, T., Helbert, W., Kopel, M., Banin, E., and Cygler, M. 2018b. Structure–function  
721 analyses of a PL24 family ulvan lyase reveal key features and suggest its catalytic mechanism. *J.*  
722 *Biol. Chem.* **293**, 4026-4036.
- 723 Valk, L.C., Luttik, M.A., de Ram, C., Pabst, M., van den Broek, M., van Loosdrecht, M.C.M., and  
724 Pronk, J.T. 2020. A Novel D-Galacturonate Fermentation Pathway in *Lactobacillus suebicus*  
725 Links Initial Reactions of the Galacturonate-Isomerase Route With the Phosphoketolase Pathway.  
726 *Front. Microbiol.* **10**, 3027.
- 727 Vendrusculo, C.T., Pereira, J.L., and Scamparini, A.R.P. 1994. Gellan gum: production and  
728 properties. In Nishinari, K. and Doi, E. (eds) *Food hydrocolloids: structure, properties and*  
729 *function*. Springer, Boston, MA, USA, pp. 91-95.
- 730 Venkatesan, J., Lowe, B., Anil, S., Manivasagan, P., Kheraif, A.A.A., Kang, K.-H., and Kim, S.-  
731 K. 2015. Seaweed polysaccharides and their potential biomedical applications. *Starch-Stärke* **67**,  
732 381-390.
- 733 Xie, W., Lin, B., Zhou, Z., Lu, G., Lun, J., Xia, C., Li, S., and Hu, Z. 2013. Characterization of a  
734 novel  $\beta$ -agarase from an agar-degrading bacterium *Catenovulum* sp. X3. *Appl. Microbiol.*  
735 *Biotechnol.* **97**, 4907-4915.
- 736 Yin, Y., Mao, X., Yang, J., Chen, X., Mao, F., and Xu, Y. 2012. dbCAN: a web resource for  
737 automated carbohydrate-active enzyme annotation. *Nucleic Acids Res.* **40**, W445-W451.

# Figure legends


Figure 1. Growth confirmation of QB4 cultured with four different uronic acid-containing polysaccharides, alginate (Alg), pectin (Pct), Ulvan (Ulv), and Gelzan (Gel). The growth was measured by cfu/mL. Media with and without 0.2% glucose was used as positive (Glu) and negative controls (NC), respectively. All data shown are mean values from three replicate experiments. Error bars denote the standard deviation of triplicate samples.

Figure 2. Growth confirmation of QB4 cultured with saturated galacturonate (Sat. Gal). The growth was measured as optical density ( $OD_{600nm}$ ). Media with and without 0.2% glucose was used as positive (Glu) and negative controls (NC), respectively. All data shown are mean values from three replicate experiments. Error bars denote the standard deviation of triplicate samples.

# Figure 1

Growth confirmation of QB4 cultured with four different uronic acid-containing polysaccharides, alginate (Alg), pectin (Pct), Ulvan (Ulv), and Gelzan (Gel).


The growth was measured by cfu/mL. Media with and without 0.2% glucose was used as positive (Glu) and negative controls (NC), respectively. All data shown are mean values from three replicate experiments. Error bars denote the standard deviation of triplicate samples.



# Figure 2

Growth confirmation of QB4 cultured with saturated galacturonate (Sat. Gal).

The growth was measured as optical density ( $OD_{600nm}$ ). Media with and without 0.2% glucose was used as positive (Glu) and negative controls (NC), respectively. All data shown are mean values from three replicate experiments. Error bars denote the standard deviation of triplicate samples.



**Table 1** (on next page)

Genes involving in alginate metabolism.

Table 1. Genes involving in alginate metabolism.

Alginate-degrading enzymes				
Abbreviation	Function	CAZy	Sp.	GenBank
Ad1_PL6	Alginate lyase	PL6, CBM32, CBM32	Type I	WP_108601791.1
Ad3_PL6	Alginate lyase	PL6, CBM16	-	WP_108605000.1
Ad4_PL7	Alginate lyase	PL7, CBM32	Type I	WP_159084278.1
Ad5_PL7	Alginate lyase	PL7, CBM32, CBM32	Type I	WP_108602212.1
Ad6_PL7	Alginate lyase	PL7	Lipobox	WP_108602720.1
Ad7_PL7	Alginate lyase	PL7	Lipobox	WP_108603319.1
Ad8_PL17	Alginate lyase	PL17	Lipobox	WP_108601502.1
Alginate-utilization proteins				
Abbreviation	Function	GenBank		
Au1	DEH reductase	WP_108601028.1		
Au2	KDG kinase	WP_108601027.1		
Au3	KDG kinase	WP_108603513.1		
Au4	2-dehydro-3-deoxyphosphogluconate aldolase	WP_108602314.1		
Au5	2-dehydro-3-deoxyphosphogluconate aldolase	WP_108603883.1		

Sp. indicates signal peptide.

## **Table 2**(on next page)

Genes involving in pectin metabolism.

Table 2. Genes involving in pectin metabolism.

Pectin-degrading enzymes				
Abbreviation	Function	CAZy	Sp.	GenBank
Pd1_PL1	Pectate Lyase (Plasmid)	PL1, CBM13	Type I	WP_108605188.1
Pd2_PL1	Pectate Lyase (Plasmid)	PL1, CBM35, CBM77	-	WP_108605291.1
Pd3_PL1	Pectate Lyase (Plasmid)	PL1, CBM35	Type I	AWB69198
Pd4_PL1	Pectate Lyase (Plasmid)	PL1, CE8	Type I	WP_159084287.1
Pd5_PL1	Pectate Lyase (Plasmid)	PL1, CE8	Lipobox	WP_108605272.1
Pd6_PL3	Pectate Lyase (Plasmid)	PL3, CBM13	Lipobox	WP_108605187.1
Pd7_PL10	Pectate Lyase (Plasmid)	PL10	Type I	WP_108605295.1
Pd8_GH105	Unsaturated galacturonyl hydrolases (Plasmid)	GH105	Lipobox	WP_108605228.1
Pd9_GH105	Unsaturated galacturonyl hydrolases (Plasmid)	GH105	Type I	WP_108605230.1
Pd10_GH105	Unsaturated galacturonyl hydrolases (Plasmid)	GH105	Lipobox	WP_108605292.1
Pd11_GH28	Polygalacturonases (Plasmid)	GH28	Tat	WP_108605238.1
Pd12_GH28	Polygalacturonases (Plasmid)	GH28	Type I	WP_108605277.1
Pectin-utilization proteins				
Abbreviation	Function	GenBank		
Pu1	5-dehydro-4-deoxy-D-glucuronate isomerase (KduI)	WP_108601704.1		
Pu2	5-dehydro-4-deoxy-D-glucuronate isomerase (KduI)	WP_108601929.1		
Pu3	5-dehydro-4-deoxy-D-glucuronate isomerase (KduI)	WP_108602166.1		
Pu4	5-dehydro-4-deoxy-D-glucuronate isomerase (KduI) (Plasmid)	WP_108605242.1		
Pu5	2-dehydro-3-deoxy-D-gluconate 5-dehydrogenase (KduD)	WP_108601158.1		
Pu6	2-dehydro-3-deoxy-D-gluconate 5-dehydrogenase (KduD)	WP_108601705.1		
Pu7	2-dehydro-3-deoxy-D-gluconate 5-dehydrogenase (KduD)	WP_108602165.1		
Pu8	2-dehydro-3-deoxy-D-gluconate 5-dehydrogenase (KduD) (Plasmid)	WP_108605243.1		
Pu9	Glucuronate isomerase	WP_108601474.1		
Pu10	Glucuronate isomerase	WP_108604964.1		
Pu11	Glucuronate isomerase (Plasmid)	WP_108605248.1		
Pu12	Tagaturonate reductase	WP_108601943.1		
Pu13	Altronate hydrolase	WP_108601944.1		
Pu14	Tagaturonate epimerase (Plasmid)	WP_108605252.1		
Pu15	Fructuronate reductase	WP_108602164.1		
Pu16	Mannoate dehydratase	WP_108601469.1		



34 Plasmid indicated that the gene is present in plasmid, not the genome.

**Table 3**(on next page)

Genes involving in ulvan metabolism.

Table 3. Genes involving in ulvan metabolism.

Ulvan-degrading enzymes

Abbreviation	Function	CAZy	Sp.	GenBank
Ud1_PL24	Ulvan lyase	PL24		WP_108601530.1
Ud2_PL24	Ulvan lyase	PL24		Type I WP_108601531.1
Ud3_PL24	Ulvan lyase	PL24		Type I WP_108604943.1
Ud4_PL24	Ulvan lyase	PL24		Type I WP_108601549.1
Ud5_PL24	Ulvan lyase	PL24		Type I WP_108601554.1
Ud6_PL24	Ulvan lyase	PL24		Lipobox WP_108602230.1
Ud7_PL24	Ulvan lyase	PL24, CBM32		WP_108604992.1
Ud8_PL24	Ulvan lyase	PL24		Type I WP_108602276.1
Ud9_PL24	Ulvan lyase	PL24		Lipobox WP_108602290.1
Ud10_PL25	Ulvan lyase	PL25		Lipobox WP_108601547.1
Ud11_PL25	Ulvan lyase	PL25		Lipobox WP_108601636.1
Ud12_PL25	Ulvan lyase	PL25		Lipobox WP_108601685.1
Ud13_PL25	Ulvan lyase	PL25		Lipobox WP_108602265.1
Ud14_PL40	Ulvan lyase	PL40		WP_108602237.1
Ud15_GH105	Unsaturated glucuronyl hydrolase	GH105		Lipobox WP_108601620.1
Ud16_GH105	Unsaturated glucuronyl hydrolase	GH105		Lipobox WP_108601696.1
Ud17_GH105	Unsaturated glucuronyl hydrolase	GH105		Lipobox WP_108601928.1
Ud18_GH105	Unsaturated glucuronyl hydrolase	GH105		Lipobox WP_108602225.1
Ud19_GH105	Unsaturated glucuronyl hydrolase	GH105		Lipobox WP_108602277.1
Ud20_GH78	$\alpha$ -L-rhamnosidase	GH78, CBM67		Type I WP_108601540.1
Ud21_GH78	$\alpha$ -L-rhamnosidase	GH78		Lipobox WP_108604942.1
Ud22_GH78	$\alpha$ -L-rhamnosidase	GH78, CBM67		Lipobox WP_108601635.1
Ud23_GH78	$\alpha$ -L-rhamnosidase	GH78		Lipobox WP_108602268.1
Ud24_GH78	$\alpha$ -L-rhamnosidase	GH78		Lipobox WP_108602275.1
Ud25_GH43	Putative $\beta$ -xylosidase	GH43		Type I WP_108601365.1
Ud26_GH43	Putative $\beta$ -xylosidase	GH43		Lipobox WP_108601561.1
Ud27_GH43	$\beta$ -xylosidase	GH43		Type I WP_108601631.1
Ud28_GH43	$\beta$ -xylosidase	GH43		Lipobox WP_108602133.1
Ud29_GH43	$\beta$ -xylosidase	GH43		Lipobox WP_159084088.1
Ud30_GH43	$\beta$ -xylosidase	GH43		Lipobox WP_108602377.1

34	Ud31_GH43	$\beta$ -xylosidase	GH43	Lipobox	WP_108602397.1
35	Ulvan-utilization proteins				
36	Abbreviation	Function	GenBank		
37	Ulu1	Sulfatase	WP_108601682.1		
38	Ulu2	Sulfatase	WP_108601697.1		
39	Ulu3	L-rhamnose mutarotase	WP_108601622.1		
40	Ulu4	Rhamnose isomerase	WP_108601699.1		
41	Ulu5	Pentulose/hexulose kinase (rhamnulokinase)	WP_108602148.1		
42	Ulu6	Rhamnulose aldolase	WP_108604932.1		
43	Ulu7	Aldehyde dehydrogenase	WP_108601159.1		
44	Ulu8	Lactate dehydrogenase	WP_108602249.1		
45	Ulu9	Xylose isomerase	WP_108604489.1		
46	Ulu10	Xylulose kinase	WP_108604490.1		
47	Ulu11	Transcriptional regulatory protein XylR	WP_10860449.1		

**Table 4**(on next page)

Genes involving in ulvan metabolism.

Table 3. Genes involving in ulvan metabolism.

Gellan-degrading enzymes				
Abbreviation	Function	CAZy	Sp.	GenBank
Gd1_GH1	$\beta$ -D-glucosidase	GH1		WP_159084202.1
Gellan-utilization proteins				
Abbreviation	Function			GenBank
Gu1	Glucokinase			WP_108603884.1
Gu2	Glucose-6-phosphate isomerase			WP_108604264.1