

Soil fertility evaluation and spatial distribution characteristics of grassland in Qilian mountain nature reserve - Tibetan Plateau (#50167)

1

First submission

Guidance from your Editor

Please submit by **3 Sep 2020** for the benefit of the authors (and your \$200 publishing discount) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Author notes

Have you read the author notes on the [guidance page](#)?



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

2 Figure file(s)

1 Table file(s)

1 Raw data file(s)



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Soil fertility evaluation and spatial distribution characteristics of grassland in Qilian mountain nature reserve

Qiang Li ^{Corresp., 1}, Junying Yang ¹, Wenhao Guan ¹, Zhigang Liu ¹, guoxing he ¹, Degang Zhang ¹, Xiaoni Liu ¹

¹ Gansu Agricultural University, Lanzhou, China

Corresponding Author: Qiang Li

Email address: 1245524440@qq.com

The study assessed the overall soil characteristics of Qilian mountain grassland and rated soil nutrient status with the China second national soil census grading standard. The Nemerow index method was used to evaluate the soil fertility of different grassland types, and GIS was used to analyse the soil nutrients spatial distribution, provide a data basis for the ecological protection and restoration of the grassland. The study indicated that the content of soil organic matter, total N, total K, total P and available K were graded at level 2 (high) or above with content of 4.99-131.52, 0.63-4.97, 15.36-21.72, 0.81-1.69 and 0.21-1.06 g kg⁻¹ respectively for most regions. Soil available P was graded as level 4 with the contents of 6.79-24.27 mg kg⁻¹. While the soil bulk density, total porosity and pH were 0.77-1.32 g cm⁻³, 35.36-58.83% and 7.63-8.54, respectively. The soil comprehensive fertility index was in a ranking order of Temperate steppe > Alpine meadow > Alpine steppe > Upland meadow > Alpine desert > Lowland meadow > Temperate Desert Steppe > Temperate Desert, and the areas with high, medium and low soil fertility accounted for 63.19%, 34.24% and 2.57% of the total grassland area of Qilian mountain soil fertility for different grassland types had different main limiting factors. The grassland soil fertility of Qilian mountain was generally at the mid-upper level. Considering the main limiting factors of different type grasslands and spatial distribution, the typical grasslands and meadows may need to apply acidic phosphate fertilizers, and desert grasslands to apply compound fertilizers of nitrogen and phosphorus to improve soil comprehensive fertility to improve grassland productivity

Soil fertility evaluation and spatial distribution characteristics of grassland in Qilian mountain nature reserve-Tibetan Plateau

Qiang Li; Junyin Yang; Wenhao Guan; Zhigang Liu; Guoxing He, Degang Zhang; Xiaoni Liu*
(College of Grassland Science, Gansu Agricultural University/Key Laboratory of Grassland Ecosystem of the Ministry of Education, Lanzhou 730070, China)

Abstract: The study assessed the overall soil characteristics of Qilian mountain grassland and rated soil nutrient status with the China second national soil census grading standard. The Nemerow index method was used to evaluate the soil fertility of different grassland types, and GIS was used to analyse the soil nutrients spatial distribution, provide a data basis for the ecological protection and restoration of the grassland. The study indicated that the content of soil organic matter, total N, total K, total P and available K were graded at level 2 (high) or above with content of 4.99-131.52, 0.63-4.97, 15.36-21.72, 0.81-1.69 and 0.21-1.06 g kg⁻¹ respectively for most regions. Soil available P was graded as level 3 with the contents of 6.79-24.27 mg kg⁻¹. While the soil bulk density, total porosity and pH were 0.77-1.32 g cm⁻³, 55.36-58.83% and 7.63-8.54, respectively. The soil comprehensive fertility index was in a ranking order of Temperate steppe > Alpine meadow > Alpine steppe > Upland meadow > Alpine desert > Lowland meadow > Temperate Desert Steppe > Temperate Desert, and the areas with high, medium and low soil fertility accounted for 63.19%, 34.24% and 2.57%. of the total grassland area of Qilian mountain soil fertility for different grassland types had different main limiting factor. The grassland soil fertility of Qilian mountain was generally at the mid-upper level. Considering the main limiting factors of different type grasslands and spatial distribution, the typical grasslands and meadows may need to apply acidic phosphate fertilizers, and desert grasslands to apply compound fertilizers of nitrogen and phosphorus to improve soil comprehensive fertility to improve grassland productivity.

Keyword: Grasslands; soil fertility; Qilian natural reserve; spatial distribution characteristics;

Soil fertility has a directly impact on the health of grasslands and is also influenced by grassland vegetation (Hao et al., 2020). Without human disturbance, the growth and distribution of grassland vegetation is strongly affected by soil fertility apart from climate (Wang et al., 2016; Harpole et al., 2007). Soil fertility not only affects the growth of grassland vegetation, but also affects the grassland ecosystem health (Ma et al., 2019). Therefore, good understanding and objective evaluation of soil fertility characteristic is of great significance to restoration and reconstruction of regional vegetation, and improvement of fragile grassland ecosystem (Su et al., 2019; Jin et al., 2018).

Grassland soil fertility plays a key role in supporting grassland ecosystem services (Clanet, 1980; Hu et al., 2018; Qu et al., 2015). Soil organic matter, nitrogen, phosphorus, potassium, soil density and pH are important components of soil fertility, while their content and spatial distribution directly affect grassland vegetation productivity (Wuest, 2015; Li et al., 2014). Many methods have been used for soil fertility evaluation, including Nemerow index method (Hua et al., 2018; Shahab et al., 2013), AHP (Keshavarzi et al., 2020; Sousa et al., 2012), correlation coefficient method, principal component analysis, etc. Among those evaluation methods, Nemerow index method has been well recognized due to it can avoid the influence of subjective factors, which could highlight the influence of the worst factor of soil attribute factors on soil fertility (Hao et al., 2012). So Nemerow comprehensive index method reflects the law of the smallest factor of plant growth in ecology, which can improve the credibility of the evaluation results (An et al., 2015; Zhou et al., 2017).

* Corresponding author. TEL:13809318054
E-mail address: liuxn@gsau.edu.cn (Liu Xiaoni).

Comprehensive evaluation of soil fertility combining with geographic information system (GIS) has been widely used to assess spatial distribution characteristics of soil nutrients, which is helpful to explore the relationship between soil nutrients and environmental factors (Wang et al., 2007; Peng et al., 2013; Nie et al., 2016; Wang et al., 2009).

The Qilian mountain natural reserve is one of the most sensitive regions under global warming and an important ecological security barrier in northwestern of China (Wang et al., 2001). Grassland ecosystem is the largest ecological system in Qilian mountain natural reserve, which plays an important role in maintaining biodiversity, water conservation and ecological balance of the natural reserve (Li et al., 2019). In last few decades climate change, human activities and mismanagement have severely damaged the Qilian mountain grassland ecosystem. Understanding the current status of grassland soil in Qilian mountain is of great significance to the health and sustainable development of grassland ecosystems. There are many types of grasslands in Qilian mountain. Due to differences in terrain, rainfall and temperature, the distribution of the same type of grassland is very patchy and has discontinuities and irregularities. Previous studies found that different grassland types have large differences in soil nutrients due to the differences in vegetation types and utilization methods (Yayiah et al., 2019; Chen et al., 2020). Many studies have been carried out on the soil of degraded grassland in Qilian mountain (Cheng et al., 2019; Wang et al., 2018; Chen et al., 2016). However, there are few studies on grassland soil fertility and its spatial distribution characteristics in Qilian mountain. Therefore, the aims of this study were 1) to investigate the soil of different type grasslands in Qilian mountain natural reserve, 2) to analyze the distribution characteristics of soil fertility index and 3) to determine the limiting factors for grassland soil fertility in Qilian mountain nature reserve, which can help to provide scientific insight for improving grassland ecological services.

1 Materials and methods

1.1 Study area

The study sites were located in the Qilian mountain nature reserve in Gansu province, China (94°10'-103°04'E, 35°50'-39°19'N). From southeast to northwest at horizontal direction, there are four vegetation zones in the order of forest, shrub, grassland and desert. At vertical direction, from low to high altitude, there are three vegetation belts: grassland belt, forest belt and alpine meadow grassland belt. The main types of soil are mountain gray cinnamon soil, subalpine meadow soil, alpine meadow soil and alpine cold desert soil. The annual precipitation varies from 100 to 500mm, mostly occurring during June to September. The average annual temperature is from -0.6 to 2.0°C; the average annual relative humidity is from 20% to 70%; the average annual evaporation is about 1200-1400 mm; and the frost-free period is about 90-120 days (<http://www.qilianshan.com.cn>).

1.2 Sites selection and Sample collection

This study sites were mainly located on the Qilian mountain natural reserve in Gansu province, China. The grassland types were Temperate steppe (TS), Alpine meadow (AM), Alpine steppe (AS), Upland meadow (UM), Alpine desert (AD), Temperate desert steppe (TDS), Lowland meadow (LM), Temperate desert (TD) (Table 1).

The sampling time of this study was from July to August 2019, when the plants were in full bloom. The central points of the typical distribution area of the above 8 types of grasslands (AM, TS, LM, AS, UM, TDS, AD and TD)

were selected as the sampling sites. A 200-meter sample line was randomly set for each sample site and the sample spots were set for every 20-meter interval. Four soil samples were taken at each sample spot using soil drill at a depth of 0-10 cm, 10-20 cm and 20-30 cm respectively. The four soil samples from each layer were mixed as one sample for that layer. The samples were put into a sample bag and taken back to a laboratory for air-drying for further test. Three soil samples were taken from each layer at each sample site. Meanwhile, a 100 cm³ soil clod was taken using ring knife method at each soil layer to measure soil bulk density and total porosity.

Table 1 Information of the sample sites

Type Grassland	Altitude m	longitude and latitude	Main plant species	Coverage %
Lowland meadow (LM)	1364	39°40'35.02"N 99°8'45.09"E	<i>Phragmites australis</i> (Cav.) Trin. ex Steud, <i>Achnatherum splendens</i> , <i>Sophora alopecuroides</i> L.	48.33
Upland meadow (UM)	3114	37°11'36.47"N 102°43'42.73"E	<i>Potentilla anserina</i> L., <i>Poa annua</i> L., <i>Elymus nutans</i> Griseb., <i>Melissilus ruthenicus</i> (L.) Peschkova, <i>Artemisia annua</i> L..	81.67
Alpine meadow (AM)	2977	37°10'48.66"N 102°47'13.83"E	<i>Polygonum viviparum</i> L., <i>Kobresia myosuroides</i> (Villars) Fiori, <i>Melissilus ruthenicus</i> (L.) Peschkova, <i>Artemisia annua</i> Linn., <i>Saussurea japonica</i> DC.	85.00
Temperate steppe (ST)	2817	37°22'13.68"N 102°40'44.93"E	<i>Poa annua</i> L., <i>Kobresia myosuroides</i> (Villars) Fiori, <i>Stipa capillata</i> Linn., <i>Potentilla anserina</i> L. , <i>Artemisia annua</i> Linn.	85.00
Alpine steppe (AT)	3735	39°16'32.99"N 97°42'52.57"E	<i>Stipa purpurea</i> , <i>Kobresia myosuroides</i> (Villars) Fiori, <i>Poa annua</i> L., <i>Potentilla anserina</i> L., <i>Androsace umbellata</i>	85.00
Temperate desert Steppe (TDS)	2139	38°57'57.23"N 99°47'41.95"E	<i>Sympegma regelii</i> Bunge, <i>Salsola collina</i> Pall., <i>Allium polyrhizum</i> Turcz, <i>Stipa capillata</i> Linn., <i>Ajania nematoloba</i>	43.75
Temperate Desert (TD)	1358	39°29'29.11"N 99°18'45.00"E	<i>Nitraria tangutorum</i> Bobr, <i>Nitraria sphaerocarpa</i> Maxim, <i>Suaeda glauca</i> (Bunge) Bunge, <i>Sympegma regelii</i> Bunge	31.67
Alpine desert (AD)	4290	39°15'34.39"N 97°45'6.70"E	<i>Rhodiola rosea</i> L., <i>Saussurea japonica</i> DC., <i>Kobresia myosuroides</i> (Villars) Fiori	28.33

1.3 Sample analyses

Soil bulk density was determined by the Core method (Dong et al., 2012). Total porosity was determined by Water immersion weighing method (Soil Physics institute, 1978). Soil samples were air-dried at room temperature, and visible roots and other debris in the soil were removed. Each soil sample was sieved through a 2-mm sieve. Soil organic matter was determined by the Walkley–Black method (Nelson and Sommers, 1996). Total N was determined by the Kjeldahl acid digestion method (Foss Kjeltex 8400, FOSS, DK) (Nelson and Sommers, 1996). Total P was measured by Mo-Sb colorimetry (UV-2102C, UNICO, Shanghai, China) (Nelson and Sommers, 1996). Total K was measured by Sodium hydroxide melting- Flame photometer method (Nelson and Sommers, 1996). Available N was determined by the Mo-Sb-Ascorbic acid colorimetric method after acid digestion (Nelson and Sommers, 1996). Available K was determined by Ammonium acetate-Flame photometer method (Nelson and Sommers, 1996).

1.4 Evaluation of soil fertility

1.4.1 Evaluation of individual indicators of soil fertility

This study used the China second soil census standard (Table 1) to rank the grassland soil organic matter, total N, total P, total K, available P, available N, pH, bulk density and total porosity indicators of Qilian mountain grassland, and to compare the differences between different grassland types (Zhou et al., 2017).

Table 1 Classification criteria used for soil nutrients

Grades	SOC g kg ⁻¹	Total N g kg ⁻¹	Total P g kg ⁻¹	Total K g kg ⁻¹	Available P mg kg ⁻¹	Available K mg kg ⁻¹	Comment
1	>40	>2.0	>1.0	>25	>40	>200	Very high
2	30-40	1.5-2.0	0.8-1.0	20-25	20-40	150-200	High
3	20-30	1.0-1.5	0.6-0.8	15-20	10-20	100-150	Upper
4	10-20	0.75-1.0	0.4-0.6	10-15	5-10	50-100	Mid-low
5	6-10	0.5-0.75	0.2-0.4	5-10	3-5	30-50	Low
6	<6	<0.5	<0.2	<5	<3	<30	Very low

1.4.2 Comprehensive soil fertility evaluation

The improved Nemerow index method was used to comprehensively evaluate the soil fertility quality of Qilian mountain (Zhou et al., 2017; Zhou et al., 2018). According to the grading standards of soil properties in the China second soil census standard (Table 2), the selected index parameters were standardized to eliminate the dimensional difference between the parameters. The standardized treatment methods are as follows:

When the attribute value belongs to the level of low, $c_i \leq x_a$, $F_i = c_i / x_a$ ($F_i \leq 1$) (1)

When the attribute value belongs to the level of upper, $x_a < c_i \leq x_c$, $F_i = 1 + (c_i - x_a) / (x_c - x_a)$ ($1 < F_i \leq 2$) (2)

When the attribute value belongs to the level of high, $x_c < c_i \leq x_p$, $F_i = 2 + (c_i - x_c) / (x_p - x_c)$ ($2 < F_i \leq 3$) (3)

When the attribute value belongs to the level of very high, $c_i > x_p$, $F_i = 3$ (4)

In the above formulas, F_i is the attribute division coefficient, c_i is the measured value of the attribute, and x_a , x_c , and x_p are the classification indexes.

The improved Nemerow index method was then used to comprehensively evaluate the grassland soil fertility in Qilian mountain. The revised Nemerow formula is as follows:

$$F = \sqrt{\frac{F_i^2 + F_{\min}^2}{2}} \cdot \left(\frac{n-1}{n}\right)$$

Where F is the soil comprehensive fertility index, F_i is the average value of each sub-fertility index, $F_{\min}$ is the minimum value of each sub-fertility index, and n is the number of participating indicators.

Table 3 Grading criterion for various soil properties in the Nemerow grading method

Soil properties	Soil bulk density	total porosity	pH	SOM	Total N	Total P	Total K	Available P	Available K
Classification	x_p	0.95	0.50	7	30	2.00	1.0	25	200
index of	x_c	1.10	0.40	8	20	1.50	0.6	15	100
Nemerow	x_a	1.25	0.30	9	10	0.75	0.4	5	50

1.5 Soil comprehensive fertility index spatial distribution

Analysis method based on multiple regression and residues (AMMRR) had been widely used in many studies for grassland spatial interpolation (Liu et al., 2012; Guo et al., 2011). This method is more accurate than many other interpolating methods and can also effectively avoids systematic errors. In this paper, based on the comprehensive fertility index determined by the improved Nemerow index method, and the ArcGIS10.2.2

(Provide a reference here) was used to conduct the spatial analyses including extracting the center points of different grassland types (Figure 1), assigning values for grassland types, performing AMMRR interpolation, and drawing the Qilian mountain grassland soil fertility index spatial distribution. The comprehensive fertility index was divided into low (<1.50), medium ($1.50-2.00$), and high (>2.00).




Fig.1 based on ArcGIS10.2.2 extract the center points of different grassland types patches spatial distribution

1.6 Statistical analyses

Statistical analyses were conducted using SPSS (version 19.0 SPSS Inc., Chicago, IL, USA). All results were presented as mean and standard deviations. One-way ANOVA and least significant difference (LSD) tests were declared at $P < 0.05$.

2 Results

2.1 Characteristics of grassland soil fertility indexes

The soil bulk density of Qilian mountain grassland was $0.77-1.32 \text{ g cm}^{-3}$, with an average value of 1.01 g cm^{-3} and a CV of 17.88% (Table 4). The total porosity was 35.36-58.83%, with an average value of 48.25% and a CV of 16.38%. The pH was 7.63-8.54, with an average value of 8.07 and a CV of 4.71%. The total N was $0.63-4.97 \text{ g kg}^{-1}$, with an average value of 2.38 g kg^{-1} and a CV of 75.49%. The total P was $0.81-1.69 \text{ g kg}^{-1}$, with an average value of 1.18 g kg^{-1} and a CV of 26.67%. The total K was $15.36-21.72 \text{ g kg}^{-1}$, with an average value of 19.92 g kg^{-1} and a CV of 9.88%. The available P was $6.79-24.27 \text{ mg kg}^{-1}$, with an average value of 12.81 mg kg^{-1} and a CV of 43.09%. The

available K was 0.21-1.06 g kg⁻¹ with an average value of 0.40 g kg⁻¹ and a CV of 68.00%. The soil organic matter was 4.99-131.52 g kg⁻¹, with an average value of 51.23 g kg⁻¹ and a CV of 95.32%.

Table 4. Descriptive statistics of grassland soils in Qilian mountain nature reserve

Item	MIN	MAX	Mean	SD	CV%
Soil bulk density g cm ⁻³	0.77	1.32	1.01	0.18	17.88
Total porosity %	35.36	58.83	48.25	7.90	16.38
pH	7.63	8.54	8.07	0.38	4.71
Total N g kg ⁻¹	0.63	4.97	2.38	1.8	75.49
Total P g kg ⁻¹	0.81	1.69	1.18	0.31	26.67
Total K g kg ⁻¹	15.36	21.72	19.92	1.965	9.88
Available P mg kg ⁻¹	6.79	24.27	12.81	5.52	43.09
Available K g kg ⁻¹	0.21	1.06	0.40	0.27	68.00
Soil organic matter g kg ⁻¹	4.99	131.52	51.23	48.83	95.32

2.2 Soil physical and chemical features in different type grasslands

The soil fertility index for different type grasslands were shown in Table 5. Except for total K, there were significant difference ($P < 0.05$) between all other soil fertility indexes of different grassland types. Soil bulk density was in a ranking order of desert type > meadow type > steppe type, or AM > TS > LM > AS > UM > TDS > AD > TD. The total porosity was in a ranking order of TS > AS > AM > UM > TDS > LM > AD > TD. The pH was in a ranking order of TD > LM > TDS > UM > AD > AS > AM > TS. Total N was in a ranking order of AM > TS > AS > UM > AD > TDS > LM > TD. The soil organic matter was in ranking order of TS > AM > AS > UM > AD > TDS > LM > TD. The total P was in a ranking order of AM > UM > LM > TS > AS > TD > AD > TDS. The available P was in a ranking order of LM > TS > AM > AS > UM > TD > TDS > AD. The available K was in a ranking of LM > UM > AS > TD > TS > AM > AD > TDS.

168 Table 5 Soil physical and chemical properties in different Grassland types in Qilian mountain Nature Reserve

Grassland Type	Soil bulk density g cm ⁻³	Total porosity %	pH	Total N g kg ⁻¹	Total P g kg ⁻¹	Total K g kg ⁻¹	Available P mg kg ⁻¹	Available K g kg ⁻¹	SOM g kg ⁻¹
LM	1.05± 0.09bc	43.65±4.83cd	8.51±0.04a	0.64±0.10d	1.36±0.09bc	19.47±2.66a	24.27±3.55a	1.06±0.91a	12.67±1.63cd
UM	0.95±0.07cd	48.73±2.06bc	7.97±0.24b	2.01±0.51c	1.53±0.23ab	15.36±1.34a	12.34±2.97b	0.45±0.06b	36.87±16.45c
AM	0.83± 0.09de	51.59±5.45b	7.76±0.26d	4.81±0.13a	1.69±0.16a	21.72±0.99a	13.73±7.54ab	0.30±0.15bc	116.46±28.35a
TS	0.77± 0.03df	54.98±1.92ab	7.63±0.10e	4.97±0.78a	1.21±0.08cd	21.18±1.02a	14.94±5.69ab	0.30±0.06c	131.52±14.33a
AS	0.91± 0.05d	58.83±2.50a	7.83±0.03bc	3.36±0.35b	0.99±0.26de	20.60±0.87a	13.65±6.95ab	0.35±0.08bc	65.56±20.49b
TDS	1.14± 0.09b	52.53±1.06b	8.50±0.03a	0.77±0.12d	0.81±0.16e	20.37±2.43a	7.96±0.65b	0.21±0.03c	13.18±1.94cd
TD	1.32± 0.06 a	35.36±4.69e	8.54±0.05a	0.63±0.08d	0.98±0.06de	20.76±0.69a	8.81±2.22b	0.32±0.05bc	4.99±0.99d
AD	1.11± 0.06b	40.32±2.18de	7.84±0.04bc	1.88±0.07c	0.90±0.06e	19.99±1.47a	6.79±0.97b	0.24±0.04c	28.65±1.90cd

169 Note: Data are presented as the mean±SD; Different small letters in the same row mean significant difference at 0.05 level. TS, Temperate steppe; AM, Alpine meadow; AS,
 170 Alpine steppe; UM, Upland meadow; AD, Alpine desert; TDS, Temperate Desert Steppe; LM, Lowland meadow; TD, Temperate Desert.

171

172

173

2.3 Soil physical and chemical spatial distribution

The soil physical and chemical spatial distributions were showed in Fig.2. Soil bulk density in most areas was 0.75-0.94 g cm⁻³. Total porosity in most areas was 50-60%. pH in most areas was 8-9. The SOM in most areas was 30-134 g kg⁻¹. The total N in most areas was 2.0-5.0 g kg⁻¹. The total P in most areas was 1.0-1.75 g kg⁻¹. The total K in most areas was 15-25 g kg⁻¹. The Available P in most areas was 10-20 mg kg⁻¹. The Available K in most areas was 0.3-1.5 g kg⁻¹.




Fig. 2 Spatial distributions of soil physical and chemical characteristics.

2.4 Soil comprehensive fertility index

The soil comprehensive fertility index of different type grasslands ranged from 1.14 to 2.32 (Table 6). The soil comprehensive fertility index was significantly higher in AM, UM, AS and TS than in AD, significantly higher in AD than in LM and TDS, and significantly higher in TDS than in TD, but no significant differences were found among others. The soil comprehensive fertility index was in a ranking order of TS > AM > AS > UM > AD > LM > TDS > TD.

The soil fertility of Qilian mountain grassland was at a moderate or high level (Fig.3). In terms of spatial distribution, the soil comprehensive fertility index was higher in eastern and western of Qilian mountain, and the soil fertility in the central region was moderate. There are only a few areas where the soil fertility of the grassland was low, distributed in the marginal regions of the western and central regions. The areas with high, medium and low soil fertility accounted for 63.19%, 34.24% and 2.57% of the total grassland area of Qilian mountain.

Table 6 Comprehensive evaluation of different Grassland Types in Qilian mountain Nature Reserve soil fertility using Nemerow index

Grassland Type	Soil bulk density	Total porosity	pH	F_i						$\overline{F_i}$	F
				Total N	Total P	Total K	Available P	Available K	SOM		
LM	2.67	2.37	1.49	0.85	3.00	2.45	3.00	3.00	1.27	2.23	1.50c
UM	3.00	2.87	2.03	3.00	3.00	2.34	2.23	3.00	3.00	2.69	2.12a
AM	3.00	3.00	2.24	3.00	3.00	2.67	2.37	3.00	3.00	2.81	2.26a
TS	3.00	3.00	2.37	3.00	3.00	2.62	2.49	3.00	3.00	2.83	2.32a
AS	3.00	3.00	2.17	3.00	2.98	2.56	2.37	3.00	3.00	2.79	2.22a
TDS	1.73	3.00	1.50	1.03	2.53	2.53	1.59	3.00	1.32	2.02	1.43c
TD	1.06	2.54	1.46	0.84	2.95	2.58	1.76	3.00	0.50	1.74	1.14d
AD	1.93	2.03	2.16	2.76	2.75	2.50	1.36	3.00	2.87	2.37	1.72b

Note: TS, Temperate steppe; AM, Alpine meadow; AS, Alpine steppe; UM, Upland meadow; AD, Alpine desert; TDS, Temperate Desert Steppe; LM, Lowland meadow; TD, Temperate Desert.

2.5 limiting factors for grassland soil comprehensive fertility

The different type grasslands soil fertility had different major limiting factors (Table 6). For example, the pH, total N and SOM were the main factors limiting soil fertility in LM, and pH and available P were the main factors limiting soil fertility in UM, AM, TS and AS. The Soil bulk density, pH, total N, SOM and available P were the main factors limiting soil fertility in TD and TDS. Soil bulk density and available P were the main factors limiting soil fertility in AD. The limiting factors for grassland soil comprehensive fertility of Qilian mountain grassland was showed in Fig.3.




Fig.3 Spatial distribution of grassland soil comprehensive fertility index and limiting factors for grassland soil comprehensive fertility.

3 Discuss

Soil organic matter is closely related to soil fertility and soil health. Nitrogen, phosphorus and potassium provide essential nutrients for plant growth and development, and are the main components of soil nutrients (Zhang et al.,2013; Zhou et al., 2016). The content of soil organic matter, total N, total P and available K were graded as

level 2 (high) or above for Qilian mountain grasslands according to the classification of China second soil census standards, with content of soil organic matter, total N, total P and available K of 4.99-131.52, 0.63-4.97, 0.81-1.69 and 0.21-1.06 g kg⁻¹ for most regions. Available P was graded as level 4 with the contents of 6.79-24.27 mg kg⁻¹. Ours study found that the difference between the maximum value and the minimum value of each fertility index was quite large, which means there was a background difference in the distribution of soil nutrients This finding was similar to the results obtained by Cui et al. (2010) and Zhou et al. (2017) for urban green space. Soil density suitable for plant growth is generally within 1.14 to 1.26 g cm⁻³ (Reisinger et al., 1988). The average soil bulk density of grassland in Qilian mountain was 1.01 g cm⁻³, with most of the area between 0.75-1.14 g cm⁻³. The grassland soil comprehensive fertility index of Qilian mountain decreases from east to west. The spatial distribution and succession of grassland types were the main factors affecting grassland soil fertility. From west to east, there are desert, typical grassland and meadow grassland types in Qilian mountain mainly. CV is an indicator of degree of dispersion of the sample. CV <10% means weak variation, 10-100% means medium variation and > 100% means strong variation. The results of ours study indicated that, except for soil pH and total K, which were weak variations, the other nutrient indicators were all moderately variable.

Grassland type is influenced by climate, vegetation and soil (Hu et al., 1978). Soil as the substrate of grassland, its physical and chemical properties of different types of grasslands provide important insight to understand grassland degradation and evolution (Gou et al., 2019; Li et al., 2019). Zhang et al (2019) found that the contents of total N, organic carbon and soluble organic carbon of different alpine types of grassland were in an order of alpine meadow> alpine meadow grassland> alpine grassland> alpine desert, and the differences between alpine types of grasslands were significant. This study observed that the ranking of the different types of grassland was desert type > meadow type > steppe type for soil bulk density, which suggested the changes of total porosity were opposite to soil bulk density. Furthermore, the total N, soil organic matter and soil comprehensive fertility index in different grassland types had significant differences. Since soil nutrients were mainly derived from the decomposition of animals, plants, microbial residues, litters, root exudates and soil parent materials, spatial heterogeneity of soil fertility distribution in different types of grassland observed in this study indicated these grasslands were influenced through the different climate and vegetation (Wei et al., 2018). Soil organic matter and nitrogen mainly came from the decomposition of organic matter, but moisture and temperature were the dominant factors controlling the decomposition rate of organic matter. This was why Ren and Hu (2008) used rainfall and temperature accumulated as a first-class classification index to classify grassland types in CSCS.

The rational construction of comprehensive evaluation factors was the key content of comprehensive evaluation of soil fertility, which directly determined the rationality and objectivity of evaluation results (Chen et al., 2019; Science - Soil Science, 2019). In many studies, the evaluation indicators of soil fertility mainly focused on nutrients such as soil organic matter, nitrogen, phosphorus and potassium (Chen et al., 2019; Science-Soil Science, 2019; Yu et al., 2018). In our study, the evaluation indexes of soil fertility incorporated the soil bulk density and total porosity, which could reflect the status of soil fertility from different angles soil compactness, permeability, infiltration performance and water holding capacity were also reflected by soil bulk density and total porosity (Garrigues et al., 2012). The modified Nemero formula highlights the effect of the minimum factor on soil fertility, reflecting the law

of the smallest factor of plant growth in ecology (An et al., 2015), and the soil minimum factor can be judged according to the minimum value of the F_i in the Nemero formula. In our study, the different type grasslands soil fertility had different main limiting factors. Such as pH, total N and SOM were the main factors limiting soil fertility in LM, and pH and available P was the main factors limiting soil fertility in UM, AM, TS and AS. The Soil bulk density, pH, total N, SOM and available P were the main factors limiting soil fertility in TD and TDS. Soil bulk density and available P were the main factors limiting soil fertility in AD. The Nemero index method can objectively reflect the comprehensive fertility characteristics of grassland soil, but many studies have not analyzed the spatial distribution characteristics of soil fertility in depth (Bao et al., 2012; Fan, et al., 2012). Our research combined GIS and soil science to draw a spatial distribution map of grassland soil fertility in Qilian mountain, which more intuitively reflected the distribution of grassland soil fertility. In our study, the areas with high, medium and low soil fertility accounted for the total grassland area of Qilian mountain was 63.19%, 34.24% and 2.57%.

Grassland was an important foundation for the construction of the Qilian mountain ecosystem. Based on the research results, the actual distribution of grassland types, Reasonable management could promote benign and sustainable development of grassland ecosystems

4 Conclusions

The results of soil nutrients and their spatial distribution of grasslands in Qilian mountain showed that, except for the low-available P content, other nutrient indicators had reached level 2 and above in China's second soil census standard, while soil bulk density was relatively low and pH was relatively high. The soil comprehensive fertility index was in a ranking order of TS > AM > AS > UM > AD > LM > TDS > TD, and the areas with high, medium and low soil fertility accounted for 63.19%, 34.24% and 2.57% of the total grassland area of Qilian mountain. The different type grasslands soil fertility had different main limiting factors. The grassland soil fertility of Qilian mountain was generally at the mid-upper level. The appropriate management methods should be adopted to improve soil fertility combining with the main limiting factors, to improve grassland ecological service function.

Acknowledgements

This study was supported by National Natural Science Foundation of China (31160475; 61401439); A new round of grassland Reward and subsidy Benefit evaluation and grassland ecological evaluation in Gansu Province(XZ20191225)..

Reference

- Abdalla, K., Mutema, M., Chivenge, P., Everson, C., & Chaplot, V. 2018. Grassland degradation significantly enhances soil CO₂ emission. *Catena*, 167, 284–292.
- An K, Xie X P, Zhang H Z, et al. 2015. Spatial pattern and impact factors of soil fertility in West Lake Scenic Area. *Chinese Journal of Ecology*, 34(4): 1091-1096. (in Chinese) DOI: 10.13292/j.1000-4890.20150304.020
- Bao Y X, Xu M G, Li F T, et al. 2012. Evaluation Method on Soil Fertility Under Long-Term Fertilization[J]. *scientia agricultura sinica*.
- Bao Y X, Xu M G, Lu F T, et al. 2012. Evaluation method on soil fertility under long-term fertilization. *Scientia Agricultura Sinica*, 45(20): 4197-4204. (in Chinese) DOI: CNKI: SUN: ZNYK.0.2012- 20-012
- Chen S, Lin B W, Li Y Q, et al. 2019. Spatial and temporal changes of soil properties and soil fertility evaluation in a large grain-production area of subtropical plain, China. *Geoderma*, 357: 113937. <https://doi.org/10.1016/j.geoderma.2019.113937>

- Chen Shuai, Lin Bowen, Li Yanqiang, et al. 2020. Spatial and temporal changes of soil properties and soil fertility evaluation in a large grain-production area of subtropical plain, China. *Geoderma*, <https://doi.org/10.1016/j.geoderma.2019.113937>
- Chen, L F, He, Z B, Zhu X, Du J, Yang J J, & Li J. 2016. Impacts of afforestation on plant diversity, soil properties, and soil organic carbon storage in a semi-arid grassland of northwestern China. *CATENA*, 147, 300–307. [Doi: 10.1016/j.catena.2016.07.009](https://doi.org/10.1016/j.catena.2016.07.009)
- Cheng T Y, Jia Y F, Wang J A, et al. 2019. Soil Conservation Function of Qilian Mountains National Nature Reserve Based on InVEST Model. *Arid Zone Research*. (in Chinese) <http://kns.cnki.net/kcms/detail/65.1095.x.20191121.0940.004.html>
- Clanet, V. 1980. In increasing or maintaining the fertility of grassland and herds: phosphoric acid plays a key role [J]. *Fourrages Actualites Supplement A Lelevage Bovin Ovin Caprin*, 9-13.
- Cui X X, Gao Y, Lu Y Z. 2010. Spatial variability of soil fertility in Daxing District of Beijing. *Transactions of the CSAE*, 26 (9): 327-333. (in Chinese) [DOI: CNKI:SUN:NYGU.0.2010-09-056](https://doi.org/10.1016/j.geoderma.2019.113937).
- Dong S K, Wen L, Li Y Y, Wang X X, Zhu L, Li X Y. 2012. Soil-quality effects of grassland degradation and restoration on the Qinghai-Tibetan Plateau. *Soil Science Society of America Journal*, 76, 2256–2264.
- Fan H R, Li Y P, Wu S X. 2012. Study on the Comprehensive Evaluation of the Soil Fertility Quality of the Green Land in Qinhuangdao[J]. *Advanced Materials Research*, 573-574:191-194. [DOI: 10.4028/www.scientific.net/AMR.573-574.191](https://doi.org/10.4028/www.scientific.net/AMR.573-574.191)
- Fayiah Moses, Dong Shikui, Li Yu, et al. 2019. The relationships between plant diversity, plant cover, plant biomass and soil fertility vary with grassland type on Qinghai-Tibetan Plateau. *Agriculture, Ecosystems & Environment*, 286, 106659. [doi:10.1016/j.agee.2019.106659](https://doi.org/10.1016/j.agee.2019.106659)
- Garrigues E, Corson M S, Angers D A, et al. 2012. Soil quality in life cycle assessment: Towards development of an indicator[J]. *Ecol Indic*, 18: 434-442. [Doi: 10.1016/j.ecolind.2011.12.014](https://doi.org/10.1016/j.ecolind.2011.12.014).
- Gou X, Hu J, Chen Y, Wei X, Du Z, Zhou Q. 2019. The effect of artificial vegetation recovery on the soil nutrients and enzyme activities in sub humid desert land on the southeast Qinghai-Tibetan Plateau, China. *Ecological Engineering*, 139, 105528. [Doi: 10.1016/j.eco.leng.2019.06.023](https://doi.org/10.1016/j.eco.leng.2019.06.023)
- Guo J, Ren Z G, Liu X N. 2011. An improved method for spatial interpolation of meteorological data based on GIS modules-A case study in Gansu Province, 31(4):41-45. (in Chinese)
- Harpole W S, Potts D L, Suding K N. 2007. Ecosystem responses to water and nitrogen amendment in a California grassland. *Global change biology*, 13(11): 2341-2348.
- Hao A H, Xue X A, Peng F, et al. 2020. different vegetation and soil degradation characteristics of a typical grassland in the Qinghai-Tibetan Plateau. *Acta Ecologica Sinica*, 40(3). (in Chinese) <http://kns.cnki.net/kcms/detail/11.2031.Q.20191120.0936.052.html>
- Hu S J, Hu R, Pu Y L, et al. 2018. Influence of ecological restoration on soil biological fertility in desertified grassland[J]. *Pratacultural Science*.
- Hua S, Li X, Jinting Z, et al. 2018. Evaluation of Heavy Metal Pollution in the Soils around the Brownfield Based on the Modified Nemero Index Method[J]. *Environmental Protection Science*.
- Hu Z Z, Zhang P J, Nan Z B, et al. 1978. Grassland types in Gansu Province. *Journal of Gansu Agricultural University*, [Doi: 10.13432/j.cnki.jgsau.1978.01.001](https://doi.org/10.13432/j.cnki.jgsau.1978.01.001). <http://www.qilianshan.com.cn/html/1/271/160/168/index.html>
- Jin H F, Shi D M, Chen Z F, et al. 2018. Evaluation indicators of cultivated layer soil quality for red soil slope farmland based on cluster and PCA analysis. *Transactions of the Chinese Society of Agricultural Engineering*, 34(7): 155- 164. (in Chinese) [Doi: 10.11975/j.issn.1002-6819.2018.07.020](https://doi.org/10.11975/j.issn.1002-6819.2018.07.020)
- Keshavarzi A, Tuffour H O, Bagherzadeh A, et al. 2020. Using fuzzy-AHP and parametric technique to assess soil fertility status in Northeast of Iran[J]. *Journal of Mountain ence*, 17(4):931-948. [DOI: 10.1007/s11629-019-5666-6](https://doi.org/10.1007/s11629-019-5666-6)
- Li J, Min Q W, Li W H, et al. 2014. Spatial variability analysis of soil nutrients based on GIS and geostatistics: a case study of Yisa township, Yunnan, China. *Journal of Resources and Ecology*, 5(4) :348-355.
- Li W, Jia X, Li M, Wu H. 2019. Insight into the vertical characteristics of dissolved organic matter in 5-m soil profiles under different land-use types on the Loess Plateau. *Science of The Total Environment*, 692: 613-621. [Doi:10.1016/j.scitotenv.2019.07.339](https://doi.org/10.1016/j.scitotenv.2019.07.339)
- Li Y G, Li Z X, Feng Q, Wei W, et al. 2019. Research on the development of the ecological protection of the Qilian Mountains based on ecological redline. *Acta Ecologica Sinica*, 39(7): 2343 – 2352. (in Chinese) [Doi: 10.5846/stxb201802250383](https://doi.org/10.5846/stxb201802250383).

- Liu X N, Guo J, Ren Z G, et al. 2012. Chinese rangeland CSCS classification based on optimal simulation for spatial distribution of meteorological factors[J]. Transactions of the Chinese Society of Agricultural Engineering (Transactions of the CSAE), 28(9): 222—229. (in Chinese) doi:10.3969/j.issn.1002-6819.2012.09.037.
- Ma Jian, Liu Xiande, Li Guang, et al. 2019.Evaluation on soil fertility quality of Picea crassifolia forest in middle Qilian Mountains. Arid Land Geography. (in Chinese) <http://kns.cnki.net/kcms/detail/65.1103.X.20191008.1740.015.html>
- Mu S, Zhou S, Chen Y, et al. 2013. Assessing the impact of restoration-induced land conversion and management alternatives on net primary productivity in Inner Mongolian grassland, China[J]. Global & Planetary Change, 108:29-41. DOI: 10.1016/j.gloplacha.2013.06.007.
- Nelson, D. W., & Sommers, L. E. 1996. Total carbon, organic carbon, and organic matter. In D. L. Sparks, A. L. Page, P. A. Helmke, R. H. Loeppert, P. N. Soltanpour, M. A. Tabatabai, C. T. Johnston, & M. E. Sumner (Eds.), Methods of soil analysis. Part 3: Chemical methods (pp. 961–1010). Madison: Soil Science Society of America, Inc., American Society of Agronomy, Inc.
- Nie Y, Yu J, Wu Y , et al. 2016. A Comprehensive Evaluation of Soil Fertility of Cultivated Land: A GIS-Based Soil Basic Niche-Fitness Model[J]. Communications in Soil Science & Plant Analysis, 00103624.2016.1146748. DOI: 10.1080/00103624.2016.1146748.
- Peng G, Bing W, Guangpo G, et al. 2013. Spatial distribution of soil organic carbon and total nitrogen based on GIS and geostatistics in a small watershed in a Hilly area of Northern China. PLoS ONE, 8(12): e83592. doi:10.1371/journal.pone.0083592.
- Qu T B, Du W C, Yuan X, et al. 2016. Impacts of Grazing Intensity and Plant Community Composition on Soil Bacterial Community Diversity in a Steppe Grassland[J]. PLoS ONE, 11(7): e0159680. DOI: 10.1371/journal.pone.0159680
- Ren J Z, Hu Z Z, Zhao J, Zhang D G, Hou F J, Lin H L, Mu X D. 2008. A Grassland classification system and its application in China[J]. The Rangeland Journal, 30: 199-209.
- Science-Soil Science. 2019. Studies from University of Ege Reveal New Findings on Soil Science (Evaluation of soil fertility in citrus planted areas by geostatistics analysis method) [J]. Science Letter. <https://kns.cnki.net/KCMS/detail/detail.aspx.dbcode>.
- Si J H, Feng Q, Yu T F, et al. 2009. Spatial heterogeneity of soil nutrients in Ejina oasis. Chinese Journal of Ecology, 28 (12): 2600-2606. (in Chinese) DOI: <http://ir.casnw.net/handle/362004/10023>
- Shahab H, Emami H, Haghnia G H , et al. 2013, Pore Size Distribution as a Soil Physical Quality Index for Agricultural and Pasture Soils in Northeastern Iran[J]. Pedosphere, (03):42-50. DOI: CNKI:SUN:TRQY.0.2013-03-007
- Soil Physics institute, Nanjing institute of Soil Science, Chinese Academy of Sciences. 1978. Soil physical properties determination method[M]. Beijing: Science Press.
- Sousa G G D, Azevedo B M D, André Henrique Pinheiro Albuquerque, et al. 2012. Características agronômicas do amendoimzeiro sob irrigação com águas salinas em solo com biofertilizantes[J]. Agro@mbiente On-line. DOI: 10.18227/1982-8470ragro.v6i2.708
- Su T T, Ma H B, Zhou Y, et al. 2019. Response of typical steppe grassland soil physical and chemical properties to various ecological restoration measures in the Ningxia Loess Hill Region. Acta Prataculturae Sinica, 28(4): 34-46. (in Chinese) Doi: 10.11686/cyxb2018247
- Wei W, Zhou J J, Shi R Z, et al. 2018. Relationship between Root Distribution Characteristics and Soil Factors in Different Grassland Communities in Tibet. Chinese Journal of Grassland, 40(6): 33-38. (in Chinese) Doi: 10.16742/j.zgcdxb.2018-06-05.
- Wang, Q., Yang, Q., Guo, H., Xiao, X., Jin, H., Li, L., Wu, Q. 2018. Hydrothermal variations in soils resulting from the freezing and thawing processes in the active layer of an alpine grassland in the Qilian Mountains, northeastern Tibetan Plateau. Theoretical and Applied Climatology. Doi:10.1007/s00704-018-2529-y
- Wang G H, Ren J Z, Zhang Z H. 2001. A Study on the population diversity of plant community, in Hexi mountain-oasis-desert area: General features. ACTA PRATACULTURAE SINICA, 10(1): 1-12. Doi: 1004-5759(2001)01-0001-12
- Wang Z Q, Zhang Y Z, Yang Y, et al. 2016. quantitative assess the driving forces on the grassland degradation in the Qinghai-Tibetan Plateau, in China. Ecological Informatics, 33: 32-44.
- Wang Z, Fu Q, Jiang Q. 2009. Comprehensive evaluation of soil fertility based on GIS and attribute recognition model[J]. Transactions of the Chinese Society of Agricultural Engineering, 25(6):76-80.
- Wang S H, Li M Cn, Su M. 2007. Research on comprehensive evaluation to land consolidation supported by GIS: a case study of Shaoshan city in Hunan province[C]// Geoinformatics: Geospatial Information Technology & Applications. International Society for Optics and Photonics, DOI: 10.1117/12.765491.

- Wuest S B. 2015. Seasonal Variation in Soil Bulk Density, Organic Nitrogen, Available Phosphorus, and pH[J]. Soil Science Society of America Journal, 79(4). DOI: [10.2136/sssaj2015.02.0066](https://doi.org/10.2136/sssaj2015.02.0066)
- Yu P J, Han D L, Huang Y X, et al. 2018. Soil quality assessment under different land uses in an alpine grassland. Catena, 171: 280-287. <https://doi.org/10.1016/j.catena.2018.07.021>.
- Zhang L N, Li J, Fan P, et al. 2013. Distribution of soil N P K nutrient content in deep soil profile of typical apple orchards on the Loess Plateau. Acta Ecologica Sinica, 33(6):1907-1915. (in Chinese) DOI: [CNKI:SUN:STXB.0.2013-06-026](https://doi.org/10.1016/j.actaes.2013.06.026)
- Zhang M M, Chen W, Lin I, et al. 2019. A study of soil nutrient characteristics and soil soluble organic carbon levels in different types of alpine grassland in Qinghai Province. Acta Prataculturae Sinica, 28(3): 20-08. (in Chinese) Doi: [10.11686/cyxb2018540](https://doi.org/10.11686/cyxb2018540).
- Zhou W, Wang W J, Zhang B, et al. 2017. Soil fertility evaluation for urban forests and green spaces in Changchun City. Acta Ecologica Sinica, 37(4) :1211-1220. DOI: 10.5846/stxb 20160 4180723
- Zhou Y, Biswas A, Ma Z Q, et al. 2016. revealing the scale- specific controls of soil organic matter at large scale in Northeast and North China Plain. Geoderma, (271): 71-79.

Table 1(on next page)

nformation of the sample sites, Classification criterion of soil nutrients, Grading criterion for various soil properties in the Nemerow grading method, Criteria for determining the organic matter, total nitrogen and bulk density of grassland soi

nformation of the sample sites, Classification criterion of soil nutrients,Grading criterion for various soil properties in the Nemerow grading method, Criteria for determining the organic matter, total nitrogen and bulk density of grassland soils with different degradation degrees,Descriptive statistics in various studied parameters of grassland soil in Qilian Mountain Nature Reserve

Table1. information of the sample sites

Type grassland	Altitude m	longitude and latitude	Main plant species	Coverage %
Lowland meadow (LM)	1364	39°40'35.02"N 99°8'45.09"E	<i>Phragmites australis</i> (Cav.) Trin. ex Steud, <i>Achnatherum splendens</i> , <i>Sophora alopecuroides</i> L.	48.33
Upland meadow (UM)	3114	37°11'36.47"N 102°43'42.73"E	<i>Potentilla anserina</i> L., <i>Poa annua</i> L., <i>Elymus nutans</i> Griseb., <i>Melissilus ruthenicus</i> (L.) Peschkova, <i>Artemisia annua</i> L..	81.67
Alpine meadow (AM)	2977	37°10'48.66"N 102°47'13.83"E	<i>Polygonum viviparum</i> L., <i>Kobresia myosuroides</i> (Villars) Fiori, <i>Melissilus ruthenicus</i> (L.) Peschkova, <i>Artemisia annua</i> Linn., <i>Saussurea japonica</i> DC.	85.00
Temperate steppe (ST)	2817	37°22'13.68"N 102°40'44.93"E	<i>Poa annua</i> L., <i>Kobresia myosuroides</i> (Villars) Fiori, <i>Stipa capillata</i> Linn., <i>Potentilla anserina</i> L. , <i>Artemisia annua</i> Linn.	85.00
Alpine steppe (AT)	3735	39°16'32.99"N 97°42'52.57"E	<i>Stipa purpurea</i> , <i>Kobresia myosuroides</i> (Villars) Fiori, <i>Poa annua</i> L., <i>Potentilla anserina</i> L., <i>Androsace umbellata</i>	85.00
Temperate Desert Steppe (TDS)	2139	38°57'57.23"N 99°47'41.95"E	<i>Sympegma regelii</i> Bunge, <i>Salsola collina</i> Pall., <i>Allium polyrhizum</i> Turcz., <i>Stipa capillata</i> Linn., <i>Ajania nematoloba</i>	43.75
Temperate Desert (TD)	1358	39°29'29.11"N 99°18'45.00"E	<i>Nitraria tangutorum</i> Bobr, <i>Nitraria sphaerocarpa</i> Maxim, <i>Suaeda glauca</i> (Bunge) Bunge, <i>Sympegma regelii</i> Bunge	31.67
Alpine desert (AD)	4290	39°15'34.39"N 97°45'6.70"E	<i>Rhodiola rosea</i> L., <i>Saussurea japonica</i> DC., <i>Kobresia myosuroides</i> (Villars) Fiori	28.33

Table2 Classification criterion of soil nutrients

Grades	SOC g kg ⁻¹	Total N g kg ⁻¹	Total P g kg ⁻¹	Total K g kg ⁻¹	Available P mg kg ⁻¹	Available K mg kg ⁻¹	Comment
1	>40	>2.0	>1.0	>25	>40	>200	Very high
2	30-40	1.5-2.0	0.8-1.0	20-25	20-40	150-200	High
3	20-30	1.0-1.5	0.6-0.8	15-20	10-20	100-150	Upper
4	10-20	0.75-1.0	0.4-0.6	10-15	5-10	50-100	Mid-low
5	6-10	0.5-0.75	0.2-0.4	5-10	3-5	30-50	Low
6	<6	<0.5	<0.2	<5	<3	<30	Very low

Table 3 Grading criterion for various soil properties in the Nemerow grading method

Soil properties		Soil bulk density	total porosity	pH	SOM	Total N	Total P	Total K	Available P	Available K
Classification	x _p	1.14	0.50	7	30	2.00	1.0	25	20	200
index of	x _c	1.04	0.40	8	20	1.50	0.6	20	10	100
Nemorow	x _a	0.94	0.30	9	10	0.75	0.4	5	5	50

Table4 Descriptive statistics in various studied parameters of grassland soil in Qilian Mountain Nature Reserve

Item	MIN	MAX	Mean	SD	CV%
Soil bulk density g cm ⁻³	0.77	1.32	1.01	0.18	17.88
Total porosity %	35.36	58.83	48.25	7.90	16.38
pH	7.63	8.54	8.07	0.38	4.71
Total N g kg ⁻¹	0.63	4.97	2.38	1.8	75.49

Total P g kg ⁻¹	0.81	1.69	1.18	0.31	26.67
Total K g kg ⁻¹	10.24	14.48	13.28	1.31	9.88
Available P mg kg ⁻¹	6.79	24.27	12.81	5.52	43.09
Available K g kg ⁻¹	0.21	1.06	0.40	0.27	68.00
Soil organic matter g kg ⁻¹	4.99	131.52	51.23	48.83	95.32

15

16 Table5 Soil physical and chemical properties in different Grassland Types in Qilian Mountain Nature Reserve

Grassland Type	Soil bulk density g cm ⁻³	Total porosity %	pH	Total N g kg ⁻¹	Total P g kg ⁻¹	Total K g kg ⁻¹	Available P mg kg ⁻¹	Available K g kg ⁻¹	SOM g kg ⁻¹
LM	1.05±0.09bc	43.65±4.83cd	8.51±0.04a	0.64±0.10d	1.36±0.09bc	12.98±1.77a	24.27±3.55a	1.06±0.91a	12.67±1.63cd
UM	0.95±0.07cd	48.73±2.06bc	7.97±0.24b	2.01±0.51c	1.53±0.23ab	10.24±0.89a	12.34±2.97b	0.45±0.06b	36.87±16.45c
AM	0.83±0.09de	51.59±5.45b	7.76±0.26d	4.81±0.13a	1.69±0.16a	14.48±0.66a	13.73±7.54ab	0.30±0.15bc	116.46±28.35a
TS	0.77±0.03df	54.98±1.92ab	7.63±0.10e	4.97±0.78a	1.21±0.08cd	14.12±0.68a	14.94±5.69ab	0.30±0.06c	131.52±14.33a
AS	0.91±0.05d	58.83±2.50a	7.83±0.03bc	3.36±0.35b	0.99±0.26de	13.73±0.58a	13.65±6.95ab	0.35±0.08bc	65.56±20.49b
TDS	1.14±0.09b	52.53±1.06b	8.50±0.03a	0.77±0.12d	0.81±0.16e	13.51±1.62a	7.96±0.65b	0.21±0.03c	13.18±1.94cd
TD	1.32±0.06 a	35.36±4.69e	8.54±0.05a	0.63±0.08d	0.98±0.06de	13.84±0.46a	8.81±2.22b	0.32±0.05bc	4.99±0.99d
AD	1.11±0.06b	40.32±2.18de	7.84±0.04bc	1.88±0.07c	0.90±0.06e	13.33±0.98a	6.79±0.97b	0.24±0.04c	28.65±1.90cd

17 Note: Data are presented as the mean±SD; Different small letters in the same row mean significant difference at 0.05 level. TS, Temperate steppe; AM, Alpine meadow; AS,
 18 Alpine steppe; UM, Upland meadow; AD, Alpine desert; TDS, Temperate Desert Steppe; LM, Lowland meadow; TD, Temperate Desert.

19
 20
 21

22

23

24 Table 6 Comprehensive evaluation of different Grassland Types in Qilian Mountain Nature Reserve soil
25 fertility via Nemerow index

26

27 Note: TS, Temperate steppe; AM, Alpine meadow; AS, Alpine steppe; UM, Upland meadow; AD, Alpine desert; TDS, Temperate Desert Steppe;
28 LM, Lowland meadow; TD, Temperate Desert.

28

29

Figure 1

Fig.1 based on ArcGIS10.2.2 extract the center points of different grassland type patches spatial distribution

Fig.1 based on ArcGIS10.2.2 extract the center points of different grassland type patches spatial distribution




Figure 2

Fig.2 Soil physical and chemical spatial distribution

Fig.2 Soil physical and chemical spatial distribution




Fig.2 Soil physical and chemical spatial distribution