

# Evaluation potential of PGPR to protect tomato against *Fusarium* wilt and promote plant growth

**Rizwana begum Syed Nabi** <sup>Equal first author, 1, 2</sup>, **Raheem Shahazad** <sup>Equal first author, 3</sup>, **Rupesh Tayade** <sup>4</sup>, **Muhammad Shahid** <sup>1</sup>, **Adil Hussain** <sup>5</sup>, **Muhammad Waqas Ali** <sup>6</sup>, **Byung-Wook Yun** <sup>Corresp. 1</sup>

<sup>1</sup> Laboratory of Plant Functional Genomics, School of Applied Biosciences, Kyungpook National University, Daegu, South Korea

<sup>2</sup> Department of Southern Area Crop Science, National Institute of Crop Science, Rural; Development Administration, Miryang, South Korea

<sup>3</sup> Department of Horticulture, The University of Haripur, Haripur, Haripur, Pakistan

<sup>4</sup> Laboratory of Plant Breeding, School of Applied Biosciences, Kyungpook National University, Daegu, South Korea

<sup>5</sup> Department of Agriculture, Abdul Wali Khan University, Mardan, Pakistan

<sup>6</sup> School of Biosciences, University of Birmingham, Edgbaston, Birmingham, United Kingdom

Corresponding Author: Byung-Wook Yun  
Email address: bwyun@knu.ac.kr

Soilborne fungal diseases are most common among vegetable crops and have major implications for crop yield and productivity. Eco-friendly sustainable agriculture practices that can overcome biotic and abiotic stresses are of prime importance. In this study, we evaluated the ability of plant growth-promoting rhizobacterium (PGPR) *Bacillus aryabhatai* strain SRB02 to control the effects of tomato wilt disease caused by *Fusarium oxysporum* f. sp. *lycopersici* (strain KACC40032) and promote plant growth. *In vitro* bioassays showed significant inhibition of fungal growth by SRB02. Inoculation of susceptible and tolerant tomato cultivars in the presence of SRB02 showed significant protection of the cultivar that was susceptible to infection and promotion of plant growth and biomass production in both of the cultivars. Further analysis of SRB02-treated plants revealed a significantly higher production of amino acids following infection by *F. oxysporum*. Analysis of plant defense hormones after inoculation by the pathogen revealed significantly higher accumulation of salicylic acid (SA), with a concomitant reduction in jasmonic acid (JA). These results indicate that *B. aryabhatai* strain SRB02 reduces the effects of *Fusarium* wilt disease in tomato by modulating endogenous phytohormones and amino acid levels.

# Evaluation potential of PGPR to protect tomato against *Fusarium* wilt and promote plant growth

Rizwana Begum Syed Nabi <sup>1,2#</sup>, Raheem Shahzad <sup>3</sup>, Rupesh Tayade <sup>4#</sup>, Muhammad Shahid <sup>1</sup>, Adil Hussain <sup>5</sup>, Muhammad Waqas Ali <sup>6</sup>, Byung-Wook Yun <sup>\*1</sup>

<sup>1</sup> Laboratory of Plant Functional Genomics, School of Applied Biosciences, Kyungpook National University, Daegu 41566, Republic of Korea; [ruhii.syed@gmail.com](mailto:ruhii.syed@gmail.com) (R.B.S.N); [shahidariswat@gmail.com](mailto:shahidariswat@gmail.com) (M.S); [bwyun@knu.ac.kr](mailto:bwyun@knu.ac.kr) (B.W)

<sup>2</sup> Department of Southern Area Crop Science, National Institute of Crop Science, Rural; Development Administration, Miryang 50424, Republic of Korea; [rizwananabi@korea.kr](mailto:rizwananabi@korea.kr) (R.B.S.N)

<sup>3</sup> Department of Horticulture, The University of Haripur, Haripur, Pakistan [raheemshahzad@ymail.com](mailto:raheemshahzad@ymail.com) (R.S);

<sup>4</sup> Laboratory of Plant Breeding, School of Applied Biosciences, Kyungpook National University, Daegu 41566, Republic of Korea; [rupesh.tayade@knu.ac.kr](mailto:rupesh.tayade@knu.ac.kr) (R.T)

<sup>5</sup> Department of Agriculture, Abdul Wali Khan University, Mardan Pakistan, [adilhussain@awkum.edu.pk](mailto:adilhussain@awkum.edu.pk) (A.H);

<sup>6</sup> School of Biosciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK; [waqasali3515@yahoo.com](mailto:waqasali3515@yahoo.com) (W.A)

\* Correspondence: [bwyun@knu.ac.kr](mailto:bwyun@knu.ac.kr); (B.W)

**Abstract:** Soilborne fungal diseases are most common among vegetable crops and have major implications for crop yield and productivity. Eco-friendly sustainable agriculture practices that can overcome biotic and abiotic stresses are of prime importance. In this study, we evaluated the ability of plant growth-promoting rhizobacterium (PGPR) *Bacillus aryabhattai* strain SRB02 to control the effects of tomato wilt disease caused by *Fusarium oxysporum* f. sp. *lycopersici* (strain KACC40032) and promote plant growth. *In vitro* bioassays showed significant inhibition of fungal growth by SRB02. Inoculation of susceptible and tolerant tomato cultivars in the presence of SRB02 showed significant protection of the cultivar that was susceptible to infection and promotion of plant growth and biomass production in both of the cultivars. Further analysis of SRB02-treated plants revealed a significantly higher production of amino acids following infection by *F. oxysporum*. Analysis of plant defense hormones after inoculation by the pathogen revealed significantly higher accumulation of salicylic acid (SA), with a concomitant reduction in jasmonic acid (JA). These results indicate that *B. aryabhattai* strain SRB02 reduces the effects of *Fusarium* wilt disease in tomato by modulating endogenous phytohormones and amino acid levels.

**Keywords:** PGPR; *Bacillus aryabhattai*; SRB02; *Fusarium oxysporum*; Tomato wilt; Tomato; Plant growth

37

# 38 1. Introduction

39 Tomato (*Solanum lycopersicum* L.) is the second most economically important edible vegetable after potato from  
40 the *Solanaceae* family and is widely cultivated and consumed around the world (Hanson & Yang, 2016). Tomato is  
41 used as a model plant for investigating the genetics and molecular aspects of disease resistance mechanisms. The  
42 tomato crop is under threat worldwide owing to biotic and abiotic stresses that have caused significant reductions in  
43 yield and productivity. One reason is that tomato is a host for nearly 200 species of plant pathogens, including fungi,  
44 bacteria, nematodes, viruses, and others that infect plants at all developmental stages (Stout, Kurabchew & Leite,  
45 2017), reducing both yield and quality.

46 Vascular wilt is one of the most important fungal diseases of tomato and occurs wherever these crops are grown.  
47 This disease is caused by the soilborne fungus *Fusarium oxysporum* f. sp. *lycopersici* (FOL). Three different  
48 pathotypes have been identified so far, which can be further classified into three races, 1, 2, and 3, based on various  
49 pathogenicity features during infection in tomato. Being soilborne, it is omnipresent and is very hard to get rid of once  
50 introduced into the cropping system. If infection occurs at the nursery or seedling stage, plants simply die back,  
51 whereas severe losses can occur if the disease appears in the field after transplantation. The fungus can spread in  
52 different ways, such as through the transport of infested soil, irrigation water, infected plants and transplants, and  
53 seeds (Jones et al., 2014). Infection occurs via the roots, causing serious vascular damage and wilting of the plant that  
54 subsequently leads to cell death. In severe infections, more than 80% of crop loss has been reported (Worku & Sahe,  
55 2018). Some studies have reported the applicability of protective fungicides as a possible remedy against the different  
56 strains of the pathogen. However, the use of chemicals in agriculture has not only raised serious concerns regarding  
57 human health and environmental hazards but is also considered responsible for the development of strains that are  
58 resistant to these widely used agrochemicals (Zouari et al., 2016). Hence, eco-friendly alternates to chemical measures  
59 are needed.

60 Biological control of plant pathogens has been of great interest to researchers. Apart from pathogenic microbes,  
61 plants also have symbiotic or mutualistic interactions with a wide range of soilborne microbes, which protect plants  
62 from pathogens either directly or by inducing resistant mechanisms (Pieterse et al., 2014). These microbes associate  
63 with the plant roots and help enhance growth-related attributes by improving the uptake of essential ions and minerals,  
64 atmospheric nitrogen fixation, and protection from pathogens (Lugtenberg & Kamilova, 2009). These growth-  
65 promoting bacteria are mainly isolated from the rhizosphere of the plants. These microbes are commonly known as  
66 plant growth-promoting rhizobacteria (PGPR) (Kloepper, Lifshitz & Zablotovicz, 1989; Backer et al., 2018) and  
67 include organisms such as *Pseudomonas* spp. Other microbes are known as plant growth-promoting endophytic  
68 bacteria, plant growth-promoting fungi, or biocontrol fungi (BCF), including *Trichoderma* spp. and *Sebacinales* spp.  
69 These can play a role in plant growth and can stimulate plant immune systems (Shoreish, Harman & Mastouri, 2010;  
70 Singh et al., 2019). Endophytes are widely dispersed and can be found in diverse environments including the tropics,  
71 temperate zone, aquatics, xerophytics and deserts, tundra, geothermal soils, rainforests, mangroves, and coastal  
72 forests. They inhabit plant tissues such as endosperms, roots, leaves, stems, flowers, and fruits (Singh et al., 2017).

Generally, plant growth promotion may occur owing to the regulation of the plant hormonal system, modifications in root architecture, production of siderophores, solubilization of soil minerals, activation of secondary mechanisms of plant defense, and production of biochemicals (Pupin & Nahas, 2014; Backer et al., 2018).

PGPRs and endophytes have a non-pathogenic symbiotic life cycle associated with their host plant tissues; these endophytes can be easily isolated from plant tissues (Arnold & Lutzoni, 2007; Costa et al., 2012). Seeds are the source of vertical dispersal of numerous seed-borne endophytes, or PGPRs (Ernst et al., 2003). Along with the alleviation of biotic stress in plants, these PGPRs have been reported to help mitigate a wide range of abiotic stresses as well (Shahzad et al., 2017a). Independent studies have reported the ameliorating effects of PGPRs on plant growth and fungal diseases in tomato and sunflower (Shittu et al., 2009; Waqas et al., 2015). In addition, studies have revealed the remediation abilities of PGPRs in soil contaminated with heavy metals (Jing, He & Yang, 2007; Bilal et al., 2018). All of these impacts of PGPRs make them widely attractive as biofertilizers and soil microbe mediators (Backer et al., 2018; Rosier, Medeiros & Bais, 2018). The positive effects of PGPRs on plant growth attributes are well known, but the exact molecular mechanism(s) behind them have not yet been clearly demonstrated.

PGPRs affect plant growth by either direct or indirect means. The direct promotion of plant growth occurs by a synthesis of complex compounds by the microbes—for instance, phytohormones such as indole-3-acetic acid (IAA), gibberellic acid (GA3), zeatin, and abscisic corrosive (ABA)—or by incremental nutrient accessibility by nitrogen fixation from the surrounding climate, thereby providing supplements for mineral solubilization (Glick, 1995; Bhardwaj et al., 2014). The indirect method of plant growth promotion takes place when PGPRs get involved in reducing the negative effects of one or more phytopathogenic microbes or fungi. This occurs by the production of substantial antagonistic substances or by inducing resistance in plants against the pathogens; for instance, the production of siderophores, hydrogen cyanide (HCN), hydrolytic proteins, etc. (Glick, 1995; Mahmood, Gupta & Kaiser, 2009).

The role of antifungal PGPRs as biological control agents to control plant diseases has been widely examined. PGPRs are considered either extracellular, including the genera *Agrobacterium*, *Arthrobacter*, *Azotobacter*, *Azospirillum*, *Bacillus*, *Burkholderia*, *Chromobacterium*, *Erwinia*, *Flavobacterium*, *Micrococcous*, *Pseudomonas*, and *Serratia*, or intracellular, including the genera *Allorhizobium*, *Bradyrhizobium*, *Mesorhizobium*, and *Rhizobium* (Martínez-Viveros et al., 2010; Gouda et al., 2018). The fact that rhizospheric bacteria *Bacillus aryabhattai* strain B8W22 was previously identified and isolated from cryotubes used for collecting air samples from the earth stratosphere (Shivaji et al., 2009) indicates that these bacteria have cosmic ancestry. Moreover, different strains of the bacterium were isolated from the rhizosphere in South Korea, India, and Tibet (Pailan et al., 2015; Lee et al., 2015; Yun et al., 2016). The plant growth-promoting ability of *B. aryabhattai* was initially reported by (Lee et al., 2012), who demonstrated growth promotion in *Xanthium italicum* plants. Similarly, Ramesh et al. (2014) (Ramesh et al., 2014) reported on *B. aryabhattai* contributions to plant growth by enhancing the mobilization and bio-fortification of zinc in soybean and wheat. More recently, *B. aryabhattai* strain SRB02 has been found to play a role in oxidative and nitrosative stress tolerance and promotion of growth in soybean plants by modulating the production of phytohormones (Park et al., 2017a). In addition, *B. aryabhattai* strains also show the ability for biosynthesis of thermostable alkaline phosphatase, anti-leukemic tumor-inhibiting L-asparaginase enzyme (Gill et al., 2013; Singh et

al., 2014), and degradation of pesticides (Pailan et al., 2015). In additions, various species of *Bacillus* have been identified as plant growth promoting bacteria as well as biocontrol agents against various pathogenic fungi (Compant et al., 2005; Shahzad et al., 2017a). Plant growth promoting rhizosphere bacteria employ a variety of strategies to facilitate plant growth and survival under pathogenic attack by both direct and indirect mechanisms. The most common direct mechanisms are phytohormone production, the acquisition of nutrients, and the control of pathogens through various means, for example, through the synthesis of hydrolytic enzymes, antifungal compounds, lipopeptides, or antibiotics. The indirect mechanisms include protection by triggering of specific defense-related pathways, particularly the induction of systemic resistance (ISR) against pathogens and pests (Khan et al., 2012; Martínez-Hidalgo et al., 2015) and the release of bacterial volatile compounds (Bernier et al., 2011). However, many environmental factors influence the biological control potential of PGPR by either predisposing pathogens to microbial antagonism, regulating the growth or production of metabolites by specific antagonists, or modulating disease development and consequently the level of disease suppression achieved.

From our literature survey, it is evident that except for some reports in crops (*Xanthium italicum*, soybean, rice, tomato, and wheat) there is a lack of information about the growth-promoting activity of *B. aryabhattai* and its role in tolerance to biotic and abiotic stress in other plant species (Viljoen et al., 2019; Yoo et al., 2019). In this study, we evaluated the plant growth-promoting abilities of *B. aryabhattai* SRB02 in tomato cultivars inoculated with phytopathogenic fungus *FOL*.

## 2. Materials and Methods

### 2.1. Growth of PGPR and *FOL*

*B. aryabhattai* SRB02 was isolated previously from the rhizosphere of a soybean field in the Chungcheong buk-do region of South Korea (Park et al., 2017b). Bacteria were cultured on LB agar or in broth (AppliChem, Darmstadt, Germany) media at 28 °C for 24 h. *F. oxysporum* f. sp. *lycopersici* strain KACC 40032 was obtained from the Korean Agricultural Culture Collection (KACC, <http://genebank.rda.go.kr>) and grown on potato dextrose agar plates at 28 °C for 7 d. The antifungal activity of *B. aryabhattai* SRB02 against *FOL* was evaluated following the protocol of (Shahzad et al., 2017a).

Briefly, a 0.5 cm<sup>2</sup> disc of active fungal mycelia of *FOL* was placed at the center of a 90 mm disposable plastic Petri dish (SPL, Korea) containing LB agar (Becton, Dickinson and Company, France). The overnight bacterial culture of *B. aryabhattai* SRB02 was aseptically streaked around the fungal disc at equal distances in a square pattern. For the untreated control, a fungal disc was placed on LB agar, as mentioned earlier, but instead of *B. aryabhattai* SRB02, only sterile water was streaked. For comparison, the effects of fungal growth inhibition of organic acids against the pathogen were also evaluated. All of the plates were incubated at 28 °C for 7 d. After the incubation period, the inhibition zone was measured and the percent inhibition was calculated according to the following formula.

$$\text{Inhibition \%} = \frac{(\text{diameter of fungus on control plate} - \text{diameter of fungus on SRB02 co-cultured plate})}{\text{diameter of fungus on control plate}} \times 100$$

### 2.2. Screening of tomato varieties for resistance to *FOL*

In the current study, tomato seeds of four Korean cultivars (IT 252842-13 (Cultivar-1), IT 252869-14 (Cultivar-2), IT 260627-16 (Cultivar-3), IT 259462-15 (Cultivar-4) ) were selected for their response to the pathogen. Seeds were sterilized with 2.5% sodium hypochlorite for 10 min and kept on wet paper towels inside Petri plates in an incubator at 25 °C for 5 d. Horticultural soil, distilled water, and pots were autoclaved at 121 °C for 20 min. Uniformly germinated seeds were transferred to separate trays filled with sterilized horticultural soil (Soil and Fertilizer Technology, Korea). After one week, uniformly grown seedlings were transplanted to big pots with the dimensions (LxWxH)-3.5 x 3 x 3 inches and volume 85-90gm. Plants were allowed to acclimatize for a few days, and the experimental treatments were set up in triplicates, with each replicate containing at least six plants. The fungal spore suspension of *FOL* strain KACC 40032 was prepared according to the protocol described by (Lichtenzweig et al., 2006). Control plants were treated with distilled water, and plants were allowed to grow for 5 d. Plants to be treated with the pathogen were inoculated by applying a spore suspension ( $10^6$  conidia/mL) to the exposed roots of tomato plants. The roots were then covered with soil. Plants were allowed to grow at relatively high humidity of  $80 \pm 2\%$ . After 14 d of growth under the conditions mentioned above, the inoculated plants were assessed on the basis of symptomatology (severity of plant wilting) and growth.

### 2.3. *In planta* biocontrol assessment

After the screening test, two cultivars (resistant and susceptible, one each) were selected based on disease symptoms and growth under biotic stress. Seeds of the selected cultivars were surface-sterilized, germinated, and grown before being transplanted to pots as mentioned previously. The plants were allowed to acclimatize for a few days, and the experimental treatments were set up in triplicate, with each replicate containing at least six plants. SRB02 was applied to plants by soil drenching with 10 mL SRB02 broth culture ( $4 \times 10^8$  cfu/mL) in the root zone. The fungal spore suspension *FOL* strain KACC 40032 was prepared as mentioned previously. Control plants were treated with distilled water, and plants were allowed to grow for 5 d. Plants to be treated with the pathogen were inoculated by applying spore suspension ( $10^6$  conidia/mL) to the exposed roots of tomato plants. The roots were then covered with soil. The plants were allowed to grow at relatively high humidity of  $80 \pm 2\%$  because to further exploit the pathogenic impact of fungus. Data were recorded on growth parameters such as plant height (PH), root length (RL), fresh weight (FW), dry weight (DW), and chlorophyll content (Chl. Cont.) to determine the response of plants to infection in the presence or absence of SRB02. For fresh plant biomasses, the plants were uprooted, carefully washed and frozen in liquid nitrogen, and then transferred to storage at  $-80$  °C until further analysis.

### 2.4. Extraction and quantification of amino acid content

The plant amino acids were extracted according to the protocol described by Khan et al. (2017)(Khan et al., 2017), with some modifications. Briefly, the freeze-dried whole plant samples were ground to homogenate, and 100 mg powdered samples were hydrolyzed under a vacuum in 6N HCl at 110 °C followed by 80 °C for 24 h. The dried residue was suspended in 0.02N HCl and filtered through a 0.45 µm filter. The amino acids were then quantified using an automatic amino acid analyzer (Hitachi, Japan; L-8900). The experiments were conducted in triplicate, and each replicate was comprised of six plants. The amino acid concentration was determined using relevant standards. These

standard known as amino acid standard mixture solution (type H) used for automatic amino acid investigation was procured through Wako Pure Chemical Industries Ltd (Japan), and used for endogenous amino acids assessment.

## 2.5. Jasmonic acid quantification

For the quantification of endogenous jasmonic acid (JA) content, the optimized protocol described by McCloud and Baldwin (1997) was used. Briefly, homogenized powder (0.3 g) from the immediately freeze-dried whole plant samples was suspended in extraction buffer (70:30 v/v acetone and 50 mM citric acid), and 25 ng JA internal standard ([9, 10-2H<sup>2</sup>]-9, 10-dihydro-JA) was also added to the suspension. The extract suspension was kept overnight at room temperature for evaporation of highly volatile organic solvents and to retain the less-volatile fatty acids. The subsequent aqueous phase was filtered and then extracted with 30 mL diethyl ether three times. The collective extracts were subsequently loaded onto a solid-phase extraction cartridge (500 mg of sorbent, aminopropyl). In addition, 7.0 mL of trichloromethane and 2-propanol (2:1 v/v) were used to wash the loaded cartridges. Then, the exogenous JA and relevant standard were eluted with 1 mL of diethyl ether and acetic acid (98:2 v/v). Following evaporation, the samples were esterified and analyzed by GCMS (6890N network GC system) and a 5973 network mass selective detector (Agilent Technologies, Palo Alto, CA, USA) in the relevant ion mode. The relevant ion mode was selected for JA determination. The ion fragment was examined at  $m/z = 83$  AMU, corresponding to the base peaks of JA and [9, 10-2H<sup>2</sup>]-9, 10-dihydro-JA. The endogenous JA values were determined from the peak areas with respect to relevant standards.

## 2.6. Salicylic acid (SA) quantification

The SA of SRB02-treated tomato plants was extracted and quantified according to the protocol described by (Enyedi et al., 1992; Seskar, Shulaev & Raskin, 1998). Immediately freeze-dried whole plant tissues were homogenized, and 0.2 g of homogenate powder was used for the extraction using 90% and 100% methanol. The pellets were dried and re-suspended in 2.5 mL 5% trichloroacetic acid (TCA) and further partitioned with ethyl acetate, cyclopentane, and isopropanol (ratio of 100:99:1, v/v). The upper organic layer containing free SA was used for air-drying with nitrogen gas. The dry SA was again suspended in 1 mL 70% methanol and subjected to high-performance liquid chromatography (HPLC) using a Shimadzu device outfitted with a fluorescence indicator (Shimadzu RF-10AxL) with excitation at 305 nm and emission at 365 nm filled with a C18 reverse-phase HPLC column (HP Hypersil ODS, particle size 5 µm, pore size 120 Å, Waters). The flow rate was maintained at 1.0 mL/min.

## 2.7. Statistical analysis

All experiments were replicated three times, and each replicate was comprised of six plants. Data were statistically evaluated with Duncan multiple range tests and *t*-tests where appropriate, using SAS version 9.2 software (Cary, NC, USA).

# 3. Results

## 3.1. In vitro antifungal assay

The *in vitro* antifungal activity of PGPR *B. aryabhattai* SRB02 was assessed against pathogenic *Fusarium oxysporum* in dual culture. The results revealed that the PGPR *B. aryabhattai* SRB02 significantly inhibited the growth of pathogenic *F. oxysporum*, as shown in Figure S1.

### 3.2. Response of tomato cultivars under pathogenic infection by *F. oxysporum*

To determine the response of four tomato cultivars, the plants were challenged with a spore suspension of the pathogen. The pathogen was applied to the exposed roots of tomato plants and incubated under higher relative humidity to create a conducive environment for successful infection. After 14 d of inoculation, the cultivars revealed a differential level of tolerance to the pathogen (Figure 1). The plant tolerance level was determined based on the symptomatology (severity of plant wilting). In the susceptible plants, clear symptoms of wilting were evident. Susceptible plants were also observed with retarded growth as compared to the tolerant plants. Based on the plant growth attributes and resistance level, as shown in Figure 1, the most tolerant and susceptible tomato cultivars were selected for further experiments.

### 3.3. Plant growth-promoting and ameliorative effects of *B. aryabhattai* SRB02 against FOL

Based on screening, the plant-growth-promoting and biocontrol efficiency of PGPR *B. aryabhattai* SRB02 against a virulent strain of *F. oxysporum* was investigated in both the selected tolerant and susceptible varieties (Figure 2). *B. aryabhattai* SRB02 significantly promoted plant growth, and interestingly reduced the disease in both tolerant and susceptible tomato cultivars (Figure 2).

The growth-related traits of the disease-tolerant plants were significantly improved when applied with SRB02 alone. The plant height (PH) was improved by 37.4%, while RL was improved by 26.8% as compared to the water-treated control plants. Other traits including seedling FW, seedling DW, and chlorophyll content were also improved by 15.3%, 23.3%, and 5.8%, respectively. A similar trend was also observed when the tolerant plants were treated with the pathogen and SRB02 together, compared to the pathogen-treated plants. The PH, RL, seedling FW, DW, and chlorophyll content were improved by 124%, 6.4%, 15.8%, 42.3%, and 39.7%, respectively, compared to the plants inoculated with the pathogen alone. SRB02 also improved the growth attributes of the disease-susceptible plants with or without co-treatment by the pathogen. The PH of the disease-susceptible plants was improved by 14.1% with the application of SRB02 in comparison with the water-treated control plants; however, the increase in PH was significantly greater (105.7%) in plants treated with SRB02 and *F. oxysporum* combined as compared to the pathogen-treated plants. Likewise, other traits were also improved in plants treated with PGPR alone as compared to the water-treated plants and also in the PGPR and pathogen co-treated susceptible plants in comparison with the plants treated with the pathogen alone. The RL, seedling FW, seedling DW, and chlorophyll content were improved by 9% and 44.5%, 10.4% and 32.6%, 3%, and 24.6%, and 4% and 61.3% in plants treated with SRB02 alone and in plants co-treated with PGPR and the pathogen, respectively (Figure 2, Table 1).

### 3.4. *B. aryabhattai* regulates defense against *F. oxysporum* by modulating defense-related hormones in tomato

Measurement of basal and induced levels of the plant defense-related hormones SA and JA following inoculation with *FOL* in the absence or presence of SRB02 revealed strict regulation of plant defense responses in SRB02-treated plants due to the regulation of the synthesis of both of these hormones (Figures 3 and 4). Interestingly, these results were observed in both the resistant and susceptible cultivars, indicating the high utility of SRB02 for field use even in susceptible crops. More specifically, SRB02-treated infected plants (tolerant and susceptible) produced significantly lower JA (11.10 % and 10.30 %, respectively) compared to control plants (Figure 3). Even the SRB02-treated plants in the absence of *FOL* accumulated lower JA (6.92% and 17.91%).

Furthermore, SRB02 treatment with *F. oxysporum*-inoculated plants of the tolerant cultivar accumulated 48.48% more SA compared to plants not treated with the PGPR. More interestingly, the response of the *F. oxysporum*-inoculated plants of the susceptible cultivar was more robust in the presence of SRB02, as these plants produced 74.60% more SA as compared to plants not treated with PGPR (Figure 4). However, no significant differences in SA accumulation were observed in SRB02-treated plants of either tolerant and susceptible cultivars in the absence of *F. oxysporum*.

### 3.5. *B. aryabhattai* SRB02 regulates amino acids in plants with or without biotic stress

The current study showed that *B. aryabhattai* SRB02 regulates amino acids in both disease-tolerant and susceptible tomato plants in the presence or absence of *F. oxysporum* (Table 2). Under pathogenic infection by *F. oxysporum*, *B. aryabhattai* SRB02 inoculation significantly enhanced aspartic acid (115.57% and 147.48%), threonine (123.18% and 118.56%), serine (123.13% and 158.91%), glutamic acid (4.86% and 157.89%), glycine (131.82% and 143.58%), alanine (99.61% and 109.67%), valine (98.13% and 74.62%), methionine (239.06% and 172.93%), isoleucine (42.60% and 97.82%), leucine (103.21% and 58.03%), tyrosine (138.45% and 65.45%), phenylalanine (39.86% and 34.16%), lysine (113.15% and 98.03%), histidine (98.42% and 111.74%), arginine (108.69% and 157.18%), and proline (90.09% and 115.25%) in disease-susceptible and tolerant tomato plants, respectively (Table 2). Only cysteine was decreased by 9.65% and 21.82% in *B. aryabhattai* SRB02 applied to susceptible and tolerant plants, respectively (Table 2).

Likewise, in the absence of the pathogen, *B. aryabhattai* SRB02 significantly enhanced aspartic acid (3.35% and 24.98%), threonine (32.99% and 118.56%), glutamic acid (4.86% and 157.89%), glycine (30.78% and 143.58%), alanine (29.70% and 4.65%), cysteine (44.22% and 60.20%), valine (21.91% and 31.81%), methionine (132.35% and 31.17%), isoleucine (97.76% and 29.48%), leucine (36.52% and 32.40%), phenylalanine (114.92% and 77.20%), histidine (22.81% and 41.48%), and arginine (35.98% and 23.95%) in the susceptible and tolerant plants, respectively (Table 2). However, *B. aryabhattai* SRB02 showed an increase in serine-intolerant (9.73%) plants and a decrease of 13.22% in susceptible plants. Tyrosine was increased by 45.03% only in the tolerant plants when applied with PGPR, while it was decreased by 51.52% in susceptible plants. Similarly, lysine was increased (35.12%) in the PGPR-applied tolerant plants, while no significant difference was observed in the susceptible plants. In contrast, proline was increased by 35.77% only in susceptible plants, while no significant difference was recorded in intolerant plants when challenged with PGPR (Table 2).

## 5. Discussion

The use of microbial-based techniques in the management of plant diseases has gained significant attention in recent years. In particular, PGPRs and their interactions with the plants under biotic or abiotic stress are gaining importance, with the ultimate aim of improvement in protection of crops and increases in agricultural production. These biocontrol approaches are eco-friendly and are becoming very popular, reliable, and long-lasting. Plant growth improvement by PGPR is one of the outstanding characteristics of these naturally occurring microbes. The improvements in plant growth and its ameliorating abilities with regard to plant diseases are determined by the interactions between the host plant and PGPR (Vejan et al., 2016). PGPR improves plant growth and health by direct or indirect mechanisms that can overcome diseases. The plant growth-promoting activity of PGPR bacteria has been reviewed in detail by d Santoyo et al. (2016) (Xia et al., 2015; Santoyo et al., 2016). *Bacillus* and *Pseudomonas* species are widely known as invaluable resources for plant growth promotion and the suppression of disease symptoms (Sundaramoorthy & Balabaskar, 2013; Chaves-López et al., 2015). Over the last few decades, several studies have reported on the beneficial aspects of *Bacillus* spp. as biocontrol and biofertilizer agents; e.g., *Bacillus licheniformis*, *Bacillus subtilis*, *Bacillus cereus*, *Bacillus pumilus*, and *Bacillus amyloliquefaciens* (Pane & Zaccardelli, 2015; Han et al., 2016). The plant growth promotion and other beneficial aspects of *Bacillus* strains can be attributed to their ability to enhance the production of phytohormones such as auxin (IAA), and gibberellic acid (Gamalero & Glick, 2011).

A wide range of plant species are infected by pathogens, including the diverse genera of *Alternaria*, *Botrytis*, *Fusarium*, and *Rhizoctonia*. These pathogens result in severe losses to crop yield and productivity, thereby posing a threat to food security. *F. oxysporum* is a devastating fungal pathogen that attacks the vascular system and causes severe damages to tomato crops across the globe. Conversely, microbes, or PGPRs, found in the rhizosphere of plants are directly associated with roots and are a vital source for plant growth promotion and suppression of soilborne plant pathogens such as *F. oxysporum*. To isolate and evaluate the beneficial role of PGPR, an appropriate *in vitro* experimental setup is required. Shahzad et al. (2017) (Shahzad et al., 2017a) reported plant growth promotion by endophytic bacteria RWL-1 against the pathogenic infection by *FOL* in tomato. In addition, it was recently reported that *B. aryabhattai* SRB02 plays a role in oxidative and nitrosative stress tolerance and promotes the growth of soybean and rice plants by modulating the production of phytohormones (Park et al., 2017a). However, it was not clear whether *B. aryabhattai* SRB02 could be used to rescue the plants from biotic stress. Hence, in the present study, we subjected disease-tolerant and susceptible tomato plants to the PGPR *B. aryabhattai* SRB02 in the presence and absence of a virulent strain of *FOL*, hypothesizing that SRB02 would rescue the plants from the disease and improve their growth under stress conditions. Prior to inoculation by the pathogen, tomato plants were treated with a cell suspension of *B. aryabhattai* SRB02. The SRB02 application improved the disease tolerance level of the infected plants. In a previous study by Shahzad et al. (2017) (Shahzad et al., 2017a), PGPRs were shown to enhance plant growth, reduce infection by the pathogen, and result in improved disease tolerance.

The present study showed that under pathogenic infection, the PGPR association rescued the plants from disease and enhanced plant growth and biomass. This result might occur by restricting the pathogenic fungus, enhancing nutrient uptake, and producing phosphate solubilization substances, or by induction of phytohormonal biosynthesis.

The present findings further strengthen the role of *Bacillus* species as a PGPR and biocontrol agent, as reported by numerous researchers, against diverse diseases in various plant species, such as root wilting, damping off, fusarium wilt, ring rot, and charcoal rot in tomato, soybean, banana, apple, and common bean, respectively (Yu et al., 2002; Vitullo et al., 2012; Wang & Fobert, 2013; Chen et al., 2016; Torres et al., 2016). The current findings also indicate that PGPR strains producing bioactive components may suppress the negative effects of pathogenesis and biotic stress in infected plants. In addition, our study also confirmed and exhibited similar results to previous reports that organic acids, as one of the many components produced by *Bacillus* species, can help rescue the plant from the disease. Moreover, PGPR produces siderophores and organic acids, which mitigate the negative effects of pathogen-infected sunflower plants (Waqas et al., 2015). From these studies, it is evident that biotic stress-related ameliorative effects are commonly regulated by endogenous phytohormones such as SA and JA. Under normal and stress conditions, phytohormone signaling and crosstalk play a vital role in plant growth and development. Accordingly, in the present study, we found that inoculation with PGPR *B. aryabhatai* SRB02 extensively modulated the endogenous levels of JA and SA. Our findings are in conformity with previously elucidated phytohormonal regulation; i.e., increased SA (Figure 4) and reduced JA (Figure 3) with PGPR, as revealed by independent studies (Khan et al., 2015; Waqas et al., 2015; Shahzad et al., 2016, 2017b; Ali et al., 2017) comparing plants in the presence or absence of biotic stress.

**Author Contributions:** RBSN performed the experiments and analyzed the data. RBSN, RT, and RS drafted the manuscript. MS, WA, and RBSN prepared illustrations, figures, tables, and references. B-WY, AH edited the manuscript. B-WY contributed critical comments to the draft and approved the manuscript. All authors contributed to the article and approved the submitted version.

**Funding:** This research was supported by Kyungpook National University Research Fund, 2019.

**Conflicts of Interest:** The authors declare no conflict of interest

# References

- Ali MW, Kim ID, Bilal S, Shahzad R, Saeed MT, Adhikari B, Nabi RBS, Kyo JR, Shin DH. 2017. Effects of bacterial fermentation on the biochemical constituents and antioxidant potential of fermented and unfermented soybeans using probiotic bacillus subtilis (KCTC 13241). *Molecules* 22. DOI: 10.3390/molecules22122200.
- Arnold AE, & Lutzoni F. 2007. Diversity and host range of foliar fungal endophytes: are tropical leaves biodiversity hotspots? *Ecology* 88:541–549.
- Backer R, Rokem JS, Ilangumaran G, Lamont J, Praslickova D, Ricci E, Subramanian S, Smith DL. 2018. Plant growth-promoting rhizobacteria: Context, mechanisms of action, and roadmap to commercialization of biostimulants for sustainable agriculture. *Frontiers in Plant Science* 871:1–17. DOI: 10.3389/fpls.2018.01473.
- Bernier SP, Létoffé S, Delepierre M, Ghigo JM. 2011. Biogenic ammonia modifies antibiotic resistance at a distance in physically separated bacteria. *Molecular Microbiology* 81:705–716. DOI: 10.1111/j.1365-2958.2011.07724.x.
- Bhardwaj D, Ansari MW, Sahoo RK, Tuteja N. 2014. Biofertilizers function as key player in sustainable agriculture by improving soil fertility, plant tolerance and crop productivity. *Microbial Cell Factories* 13:1–10. DOI: 10.1186/1475-2859-13-66.
- Bilal S, Shahzad R, Khan AL, Kang SM, Imran QM, Al-Harrasi A, Yun BW, Lee IJ. 2018. Endophytic microbial consortia of phytohormones-producing fungus paecilomyces formosus lhl10 and bacteria sphingomonas sp. lkl1 to glycine max l. regulates physio-hormonal changes to attenuate aluminum and zinc stresses. *Frontiers in Plant Science* 9:1–18. DOI: 10.3389/fpls.2018.01273.
- Chaves-López C, Serio A, Gianotti A, Sacchetti G, Ndagijimana M, Ciccarone C, Stellarini A, Corsetti A, Paparella A. 2015. Diversity of food-borne Bacillus volatile compounds and influence on fungal growth. *Journal of Applied Microbiology* 119:487–499. DOI: 10.1111/jam.12847.

- 364 Chen G, Ma C, Mukherjee A, Musante C, Zhang J, White JC, Dhankher OP, Xing B. 2016. *Tannic acid alleviates*  
365 *bulk and nanoparticle Nd<sub>2</sub>O<sub>3</sub> toxicity in pumpkin: A physiological and molecular response*. DOI:  
366 10.1080/17435390.2016.1202349.
- 367 Compant S, Duffy B, Nowak J, Clément C, Barka EA. 2005. Use of plant growth-promoting bacteria for biocontrol  
368 of plant diseases: Principles, mechanisms of action, and future prospects. *Applied and Environmental*  
369 *Microbiology* 71:4951–4959. DOI: 10.1128/AEM.71.9.4951-4959.2005.
- 370 Costa LEDO, Queiroz MVD, Borges AC, Moraes CAD, & Araújo EFD. 2012. Isolation and characterization of  
371 endophytic bacteria isolated from the leaves of the common bean (*Phaseolus vulgaris*). *Microbiology* 0:1562–  
372 1575.
- 373 Enyedi AJ, Yalpani N, Silverman P, Raskin I. 1992. Localization, conjugation, and function of salicylic acid in tobacco  
374 during the hypersensitive reaction to tobacco mosaic virus. *Proceedings of the National Academy of Sciences of*  
375 *the United States of America* 89:2480–2484. DOI: 10.1073/pnas.89.6.2480.
- 376 Ernst M, Mendgen KW, Wiersel SGR, Phytopathologie L, Biologie F, Konstanz U, Konstanz D-. 2003. Endophytic  
377 Fungal Mutualists : Seed-Borne Stagonospora Spp . Enhance Reed Biomass Production in Axenic Microcosms.  
378 16:580–587.
- 379 Gamalero E, Glick BR. 2011. *Bacteria in Agrobiolgy: Plant Nutrient Management*. DOI: 10.1007/978-3-642-21061-  
380 7.
- 381 Gill SS, Hasanuzzaman M, Nahar K, Macovei A, Tuteja N. 2013. Importance of nitric oxide in cadmium stress  
382 tolerance in crop plants. *Plant Physiology and Biochemistry* 63:254–261. DOI: 10.1016/j.plaphy.2012.12.001.
- 383 Glick BR. 1995. The enhancement of plant growth by free-living bacteria. *Canadian Journal of Microbiology* 41:109–  
384 117. DOI: 10.1139/m95-015.
- 385 Gouda S, Kerry RG, Das G, Paramithiotis S, Shin HS, Patra JK. 2018. Revitalization of plant growth promoting

- 386 rhizobacteria for sustainable development in agriculture. *Microbiological Research* 206:131–140. DOI:  
387 10.1016/j.micres.2017.08.016.
- 388 Han R, Wang H, Hu ZZ, Kumar A, Li W, Long LN, Schemm JKE, Peng P, Wang W, Si D, Jia X, Zhao M, Vecchi  
389 GA, LaRow TE, Lim YK, Schubert SD, Camargo SJ, Henderson N, Jonas JA, Walsh KJE. 2016. An assessment  
390 of multimodel simulations for the variability of Western North Pacific tropical cyclones and its association with  
391 ENSO. *Journal of Climate* 29:6401–6423. DOI: 10.1175/JCLI-D-15-0720.1.
- 392 Hanson PM, Yang RY. 2016. Epidural Analgesia Inhibits the Renin and Aldosterone Response to Surgery. *World*  
393 *Vegetable Center. Ekin J.* 2:1–10. DOI: 10.1111/j.1399-6576.1979.tb01450.x.
- 394 Jing Y de, He Z li, Yang X e. 2007. Role of soil rhizobacteria in phytoremediation of heavy metal contaminated soils.  
395 *Journal of Zhejiang University. Science. B.* 8:192–207. DOI: 10.1631/jzus.2007.B0192.
- 396 Jones JB, Zitter TA, Momol MT, Miller SA. 2014. Compendium of tomato diseases second edition. :176.
- 397 Khan AL, Hussain J, Al-Harrasi A, Al-Rawahi A, Lee IJ. 2015. Endophytic fungi: Resource for gibberellins and crop  
398 abiotic stress resistance. *Critical Reviews in Biotechnology* 35:62–74. DOI: 10.3109/07388551.2013.800018.
- 399 Khan N, Mishra A, Nautiyal CS. 2012. *Paenibacillus lentimorbus* B-30488 r controls early blight disease in tomato  
400 by inducing host resistance associated gene expression and inhibiting *Alternaria solani*. *Biological Control*  
401 62:65–74. DOI: 10.1016/j.biocontrol.2012.03.010.
- 402 Khan AL, Waqas M, Asaf S, Kamran M, Shahzad R, Bilal S, Khan MA, Kang SM, Kim YH, Yun BW, Al-Rawahi  
403 A, Al-Harrasi A, Lee IJ. 2017. Plant growth-promoting endophyte *Sphingomonas* sp. LK11 alleviates salinity  
404 stress in *Solanum pimpinellifolium*. *Environmental and Experimental Botany* 133:58–69. DOI:  
405 10.1016/j.envexpbot.2016.09.009.
- 406 Kloepper JW, Lifshitz R, Zablotovicz RM. 1989. Free-living bacteria inocula for enhancing crop productivity . *Trends*  
407 *Biotechnol* 7 : 39-44. 7799:39–44. DOI: 10.1016/0167-7799(89)90057-7.

- 408 Lee TJ, Chau HM, Thi T, Ha T, Yi GH, Park DJ, Kim H, Kim NS, Park CK, Cho WD. 2012. Yield and Fruit  
409 Characteristics of Korean Tomato (*Lycopersicon esculentum* Mill.) Cultivars Established at Different Cropping  
410 Seasons in Hanoi, Vietnam. *Korean Journal of International Agriculture* 24:309–315.
- 411 Lee J-Y, Jun N-R, Yoon D, Shin C, Baik I. 2015. Association between dietary patterns in the remote past and telomere  
412 length. *European Journal of Clinical Nutrition* 69:1048–1052. DOI: 10.1038/ejcn.2015.58.
- 413 Lichtenzweig J, Anderson J, Thomas G, Oliver R. 2006. Inoculation and growth with soil borne pathogenic fungi. :1–  
414 10.
- 415 Lugtenberg B, Kamilova F. 2009. Plant-Growth-Promoting Rhizobacteria. DOI:  
416 10.1146/annurev.micro.62.081307.162918.
- 417 Mahmood T, Gupta KJ, Kaiser WM. 2009. Cadmium Stress Stimulates Nitric Oxide Production By Wheat Roots.  
418 *Pak. J. Bot* 41:1285–1290.
- 419 Martínez-Hidalgo P, García JM, Pozo MJ. 2015. Induced systemic resistance against *Botrytis cinerea* by  
420 *Micromonospora* strains isolated from root nodules. *Frontiers in Microbiology* 6:1–11. DOI:  
421 10.3389/fmicb.2015.00922.
- 422 Martínez-Viveros O, Jorquera MA, Crowley DE, Gajardo G, Mora ML. 2010. Mechanisms and practical  
423 considerations involved in plant growth promotion by Rhizobacteria. *Journal of Soil Science and Plant Nutrition*  
424 10:293–319. DOI: 10.4067/S0718-95162010000100006.
- 425 McCloud ES, Baldwin IT. 1997. Herbivory and caterpillar regurgitants amplify the wound-induced increases in  
426 jasmonic acid but not nicotine in *Nicotiana sylvestris*. *Planta* 203:430–435. DOI: 10.1007/s004250050210.
- 427 Pailan S, Gupta D, Apte S, Krishnamurthi S, Saha P. 2015. Degradation of organophosphate insecticide by a novel  
428 *Bacillus aryabhatai* strain SanPS1, isolated from soil of agricultural field in Burdwan, West Bengal, India.  
429 *International Biodeterioration and Biodegradation* 103:191–195. DOI: 10.1016/j.ibiod.2015.05.006.

- 430 Pane C, Zaccardelli M. 2015. Evaluation of *Bacillus* strains isolated from solanaceous phylloplane for biocontrol of  
431 *Alternaria* early blight of tomato. *Biological Control* 84:11–18. DOI: 10.1016/j.biocontrol.2015.01.005.
- 432 Park YG, Mun BG, Kang SM, Hussain A, Shahzad R, Seo CW, Kim AY, Lee SU, Oh KY, Lee DY, Lee IJ, Yun BW.  
433 2017a. *Bacillus aryabhattai* SRB02 tolerates oxidative and nitrosative stress and promotes the growth of soybean  
434 by modulating the production of phytohormones. *PLoS ONE* 12:1–28. DOI: 10.1371/journal.pone.0173203.
- 435 Park YG, Mun BG, Kang SM, Hussain A, Shahzad R, Seo CW, Kim AY, Lee SU, Oh KY, Lee DY, Lee IJ, Yun BW.  
436 2017b. *Bacillus aryabhattai* SRB02 tolerates oxidative and nitrosative stress and promotes the growth of soybean  
437 by modulating the production of phytohormones. *PLoS ONE* 12:1–28. DOI: 10.1371/journal.pone.0173203.
- 438 Pieterse CMJ, Zamioudis C, Berendsen RL, Weller DM, Van Wees SCM, Bakker PAHM. 2014. Induced Systemic  
439 Resistance by Beneficial Microbes. *Annual Review of Phytopathology* 52:347–375. DOI: 10.1146/annurev-  
440 phyto-082712-102340.
- 441 Pupin B, Nahas E. 2014. Microbial populations and activities of mangrove, restinga and Atlantic forest soils from  
442 Cardoso Island, Brazil. *Journal of Applied Microbiology* 116:851–864. DOI: 10.1111/jam.12413.
- 443 Ramesh A, Sharma SK, Sharma MP, Yadav N, Joshi OP. 2014. Inoculation of zinc solubilizing *Bacillus aryabhattai*  
444 strains for improved growth, mobilization and biofortification of zinc in soybean and wheat cultivated in  
445 Vertisols of central India. *Applied Soil Ecology* 73:87–96. DOI: 10.1016/j.apsoil.2013.08.009.
- 446 Rosier A, Medeiros FHV, Bais HP. 2018. Defining plant growth promoting rhizobacteria molecular and biochemical  
447 networks in beneficial plant-microbe interactions. *Plant and Soil* 428:35–55. DOI: 10.1007/s11104-018-3679-  
448 5.
- 449 Santoyo G, Moreno-Hagelsieb, Orozco-Mosqueda G del C, Ma Glick BR. 2016. Plant growth-promoting bacterial  
450 endophytes. *Microbiological Research* 183:92–99. DOI: 10.1016/j.micres.2015.11.008.
- 451 Seskar M, Shulaev V, Raskin I. 1998. Endogenous methyl salicylate in pathogen-inoculated tobacco plants. *Plant*

452       *Physiology* 116:387–392. DOI: 10.1104/pp.116.1.387.

453   Shahzad R, Khan AL, Bilal S, Asaf S, Lee IJ. 2017a. Plant growth-promoting endophytic bacteria versus pathogenic  
454       infections: An example of *Bacillus amyloliquefaciens* RWL-1 and *Fusarium oxysporum* f. sp. *lycopersici* in  
455       tomato. *PeerJ* 2017:e3107. DOI: 10.7717/peerj.3107.

456   Shahzad R, Khan AL, Bilal S, Asaf S, Lee IJ. 2017b. Plant growth-promoting endophytic bacteria versus pathogenic  
457       infections: An example of *Bacillus amyloliquefaciens* RWL-1 and *Fusarium oxysporum* f. sp. *lycopersici* in  
458       tomato. *PeerJ* 2017:e3107. DOI: 10.7717/peerj.3107.

459   Shahzad R, Waqas M, Khan AL, Asaf S, Khan MA, Kang SM, Yun BW, Lee IJ. 2016. Seed-borne endophytic *Bacillus*  
460       *amyloliquefaciens* RWL-1 produces gibberellins and regulates endogenous phytohormones of *Oryza sativa*.  
461       *Plant Physiology and Biochemistry* 106:236–243. DOI: 10.1016/j.plaphy.2016.05.006.

462   Shittu HO, Castroverde DCM, Nazar RN, Robb J. 2009. Plant-endophyte interplay protects tomato against a virulent  
463       *Verticillium*. *Planta* 229:415–426. DOI: 10.1007/s00425-008-0840-z.

464   Shivaji S, Chaturvedi P, Begum Z, Pindi PK, Manorama R, Padmanaban DA, Shouche YS, Pawar S, Vaishampayan  
465       P, Dutt CBS, Datta GN, Manchanda RK, Rao UR, Bhargava PM, Narlikar J V. 2009. *Janibacter hoylei* sp. nov.,  
466       *Bacillus isronensis* sp. nov. and *Bacillus aryabhatai* sp. nov., isolated from cryotubes used for collecting air  
467       from the upper atmosphere. *International Journal of Systematic and Evolutionary Microbiology* 59:2977–2986.  
468       DOI: 10.1099/ijs.0.002527-0.

469   Shoresh M, Harman GE, Mastouri F. 2010. Induced Systemic Resistance and Plant Responses to Fungal Biocontrol  
470       Agents. *Annual Review of Phytopathology* 48:21–43. DOI: 10.1146/annurev-phyto-073009-114450.

471   Singh M, Kumar A, Singh R, Deo K. 2017. Endophytic bacteria : a new source of bioactive compounds. *3 Biotech*  
472       7:1–14. DOI: 10.1007/s13205-017-0942-z.

473   Singh M, Singh D, Gupta A, Pandey KD, Singh PK, Kumar A. 2019. Plant Growth Promoting Rhizobacteria. In:

- 474 *PGPR Amelioration in Sustainable Agriculture*. Elsevier, 41–66. DOI: 10.1016/B978-0-12-815879-1.00003-3.
- 475 Singh N, Srivastava G, Talat M, Raghubanshi H, Srivastava ON, Kayastha AM. 2014. Cicer a-galactosidase  
476 immobilization onto functionalized graphene nanosheets using response surface method and its applications.  
477 *Food Chemistry* 142:430–438. DOI: 10.1016/j.foodchem.2013.07.079.
- 478 Stout MJ, Kurabchew H, Leite GLD. 2017. *Host-Plant Resistance in Tomato*. Elsevier Inc. DOI: 10.1016/B978-0-12-  
479 802441-6.00009-7.
- 480 Sundaramoorthy S, Balabaskar and P. 2013. Evaluation of Combined Efficacy of *Pseudomonas fluorescens* and  
481 *Bacillus subtilis* in Managing Tomato Wilt Caused by *Fusarium oxysporum* f. sp. *lycopersici* (Fol). *Plant*  
482 *Pathology Journal* 12:154–161. DOI: 10.3923/ppj.2013.154.161.
- 483 Torres MJ, Brandan CP, Petroselli G, Erra-Balsells R, Audisio MC. 2016. Antagonistic effects of *Bacillus subtilis*  
484 subsp. *subtilis* and *B. amyloliquefaciens* against *Macrophomina phaseolina*: SEM study of fungal changes and  
485 UV-MALDI-TOF MS analysis of their bioactive compounds. *Microbiological Research* 182:31–39. DOI:  
486 10.1016/j.micres.2015.09.005.
- 487 Vejan P, Abdullah R, Khadiran T, Ismail S, Nasrulhaq Boyce A. 2016. Role of plant growth promoting rhizobacteria  
488 in agricultural sustainability-A review. *Molecules* 21:1–17. DOI: 10.3390/molecules21050573.
- 489 Viljoen JJF, Labuschagne N, Fourie H, Sikora RA. 2019. Biological control of the root-knot nematode *Meloidogyne*  
490 *incognita* on tomatoes and carrots by plant growth-promoting rhizobacteria. *Tropical Plant Pathology* 44:284–  
491 291. DOI: 10.1007/s40858-019-00283-2.
- 492 Vitullo D, Di Pietro A, Romano A, Lanzotti V, Lima G. 2012. Role of new bacterial surfactins in the antifungal  
493 interaction between *Bacillus amyloliquefaciens* and *Fusarium oxysporum*. *Plant Pathology* 61:689–699. DOI:  
494 10.1111/j.1365-3059.2011.02561.x.
- 495 Wang L, Fobert PR. 2013. *Arabidopsis* Clade I TGA Factors Regulate Apoplastic Defences against the Bacterial

496 Pathogen *Pseudomonas syringae* through Endoplasmic Reticulum-Based Processes. *PLoS ONE* 8:1–13. DOI:  
 497 10.1371/journal.pone.0077378.

498 Waqas M, Khan AL, Hamayun M, Shahzad R, Kim YH, Choi KS, Lee IJ. 2015. Endophytic infection alleviates biotic  
 499 stress in sunflower through regulation of defence hormones, antioxidants and functional amino acids. *European*  
 500 *Journal of Plant Pathology* 141:803–824. DOI: 10.1007/s10658-014-0581-8.

501 Worku M, Sahe S. 2018. Review on Disease Management Practice of Tomato Wilt Caused *Fusarium oxysporum* in  
 502 Case of Ethiopia. *Journal of Plant Pathology & Microbiology* 9:9–12. DOI: 10.4172/2157-7471.1000460.

503 Xia J, Niu S, Ciais P, Janssens IA, Chen J, Ammann C, Arain A, Blanken PD, Cescatti A, Bonal D, Buchmann N,  
 504 Curtis PS, Chen S, Dong J, Flanagan LB, Frankenberg C, Georgiadis T, Gough CM, Hui D, Kiely G, Li J, Lund  
 505 M, Magliulo V, Marcolla B, Merbold L, Montagnani L, Moors EJ, Olesen JE, Piao S, Raschi A, Roupsard O,  
 506 Suyker AE, Urbaniak M, Vaccari FP, Varlagin A, Vesala T, Wilkinson M, Weng E, Wohlfahrt G, Yan L, Luo  
 507 Y. 2015. Joint control of terrestrial gross primary productivity by plant phenology and physiology. *Proceedings*  
 508 *of the National Academy of Sciences of the United States of America* 112:2788–2793. DOI:  
 509 10.1073/pnas.1413090112.

510 Yoo SJ, Weon HY, Song J, Sang MK. 2019. Induced tolerance to salinity stress by halotolerant bacteria *Bacillus*  
 511 *aryabhattai* H19-1 and *B. Mesonae* H20-5 in tomato plants. *Journal of Microbiology and Biotechnology*  
 512 29:1124–1136. DOI: 10.4014/jmb.1904.04026.

513 Yu GY, Sinclair JB, Hartman GL, Bertagnolli BL. 2002. Production of iturin A by *Bacillus amyloliquefaciens*  
 514 suppressing *Rhizoctonia solani*. *Soil Biology and Biochemistry* 34:955–963. DOI: 10.1016/S0038-  
 515 0717(02)00027-5.

516 Yun B, Skelly MJ, Yin M, Yu M, Mun B, Lee S, Hussain A, Spoel SH, Loake GJ. 2016. Nitric oxide and S -  
 517 nitrosogluthathione function additively during plant immunity. *New Phytologist* 211:516–526. DOI:  
 518 10.1111/nph.13903.


519 Zouari I, Jlaiel L, Tounsi S, Trigui M. 2016. Biocontrol activity of the endophytic *Bacillus amyloliquefaciens* strain  
 520 CEIZ-11 against *Pythium aphanidermatum* and purification of its bioactive compounds. *Biological Control*  
 521 100:54–62. DOI: 10.1016/j.biocontrol.2016.05.012.

522

523

# Figure 1


Figure 1. Response of four tomato cultivars against *F. oxysporum* f. sp. *Lycopersici*. Cultivar 1 showed tolerance, cultivar 2 was moderately tolerant, and cultivars 3 and 4 were susceptible based upon the symptomatology.



# Figure 2

Figure 2. Effect of SRB02 on susceptible and tolerant tomato plants under inoculation with pathogenic *F. oxysporum* f. sp. *lycopersici*. The effect of SRB02 on tomato disease-tolerant plants


(A) and disease-susceptible plants (B) to ameliorate disease symptoms and promote growth parameter SRB02 improved the roots of both tolerant and susceptible plants in the presence and absence of the pathogen (C).



# Figure 3

Figure 3. Regulation of jasmonic acid accumulation in plants.

A. Under pathogenic stress by *Fusarium oxysporum*, *B. aryabhattai* SRB02 significantly reduced the endogenous JA accumulation level in both susceptible and tolerant plants. Conversely, a reduction i



# **Figure 4**(on next page)

Figure 4. Regulation of salicylic acid accumulation in plants. In the presence of the pathogen, *B. aryabhattai* SRB02 significantly enhanced SA accumulation in both tolerant and susceptible plants, while in the absence of the pathogen, the plants displayed

Salicylic acid (ng/g dry weight)



Tolerant

Susceptible


Control

SRB02

F. oxy

SRB02 + F. oxy

125  
100  
75  
50  
25  
0



**Table 1** (on next page)

Table 1. Effects of SRB02 on growth parameters of tomato plants under control and biotic stress conditions

**Table 1. Effects of SRB02 on growth parameters of tomato plants under control and biotic stress conditions**

Variety	Treatment	PH (cm)	RL (cm)	FW (g)	DW (g)	Chl.Cont. (SPAD)
<b>Tolerant</b>	Control	28.73 ± 0.40b	15.16 ± 0.34b	14.38 ± 0.05b	6.09 ± 0.25b	28.38 ± 0.37b
	SRB02	39.48 ± 0.96a	19.23 ± 0.64a	16.58 ± 1.00a	7.51 ± 0.60a	30.03 ± 0.40a
	<i>F. oxysporum</i>	17.50 ± 0.51b	14.25 ± 0.48b	13.59 ± 0.01b	5.18 ± 0.10b	23.38 ± 0.38b
	SRB02 + <i>F. oxy</i>	39.20 ± 0.88a	15.17 ± 0.34a	15.74 ± 0.42a	7.37 ± 0.27a	32.67 ± 0.48a
<b>Susceptible</b>	Control	29.88 ± 0.45b	18.75 ± 0.43b	16.11 ± 0.18b	6.89 ± 0.49b	27.11 ± 0.34b
	SRB02	34.10 ± 0.71a	20.43 ± 0.58a	14.42 ± 0.04a	7.10 ± 0.08a	28.17 ± 1.42a
	<i>F. oxysporum</i>	14.63 ± 0.62b	13.41 ± 0.76b	10.83 ± 0.57b	4.88 ± 0.18b	25.11 ± 1.07b
	SRB02 + <i>F. oxy</i>	30.10 ± 0.53a	19.38 ± 0.37a	14.36 ± 0.27a	6.08 ± 0.08a	40.51 ± 0.98a

Notes: PH, plant height; RL, root length; FW, fresh weight; DW, dry weight; Chl. Cont., chlorophyll content. Data represent means ± SD of six replicates from three independent experiments. Each values in columns followed by different letters are significantly different at  $P \geq 0.05$ .

## Table 2 (on next page)

Table 2. Regulation of Amino acids (mg/g DW) in tomato plants applied with SBR02 in the presence and absence of *F. oxysporum* f. sp. *lycopersici*.

1 **Table 2.** Regulation of Amino acids (mg/g DW) in tomato plants applied with SBR02 in the presence and absence of *F. oxysporum* f. sp. *lycopersici*.

Cultivar	Treatment	Asp	Thr	Met	ILE	Ser	Glu	Leu	Tyr	Gly	Phe	Lys	Cys	Val	His	Arg	Ala	Pro
Tolerant	Control	4.93	4.84	0.69	4.95	4.83	12.5	9.91	3.30	5.73	3.38	6.63	0.36	6.10	2.67	4.49	6.31	9.87
		±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
		0.12b	0.27b	0.17b	0.22b	0.11b	0.32b	0.11b	0.60a	0.22b	0.32b	2.69b	0.02a	0.21b	0.32b	0.45b	0.21b	1.15b
	SRB02	6.16	4.71	0.91	6.41	5.30	16.1	13.13	1.60	5.80	5.99	7.73	0.57	8.05	3.78	5.56	6.60	13.4
		±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
		0.07a	0.27a	0.08a	0.24a	0.14a	0.33a	0.4a	0.15b	0.16a	0.14a	1.52a	0.03a	0.14a	0.31a	0.47a	0.18a	0.74a
	<i>F. oxy</i>	2.96	3.05	0.32	2.51	2.70	7.36	6.21	2.29	3.24	5.92	5.29	0.53	3.54	1.25	2.82	4.06	4.67
		±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
		0.12b	0.27b	0.06b	0.44b	0.14b	0.35b	0.20b	0.2b	0.14b	0.18b	0.62b	0.05a	0.42b	0.13b	0.3b	0.06b	0.56b
	SRB02 + <i>F. oxy</i>	7.33	6.66	0.88	4.98	7.01	18.99	9.82	3.79	7.90	7.94	10.49	0.41	6.18	2.66	7.26	8.52	10.0
		±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
		0.46a	0.22a	0.02a	0.09a	0.22a	0.2a	0.26a	0.25a	0.15a	0.19a	2.26a	0.01b	0.19a	0.33a	0.6a	0.34a	0.7a
Susceptible	Control	6.14	4.41	0.73	3.02	5.90	7.77	9.28	2.50	5.31	3.57	8.44	0.39	6.30	2.96	4.67	6.36	11.5
		±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
		0.07b	0.38b	0.1b	0.14b	0.09b	0.8b	0.24b	0.30a	0.26b	0.24b	0.38b	0.04a	0.36b	0.09b	0.32b	0.43b	1.09b
	SRB02	6.34	5.87	1.71	5.99	5.12	14.1	12.67	3.60	6.95	7.69	11.4	0.55	7.68	3.63	6.35	8.25	12.5
		±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
		0.13a	0.12a	0.38a	0.14a	0.11a	0.69a	0.54a	0.20b	0.1a	0.36a	1.08a	0.01a	0.29a	0.44a	0.17a	0.22a	0.56a
	<i>F. oxy</i>	3.45	2.94	0.43	4.58	2.90	16.87	6.19	1.60	3.28	5.60	5.33	0.69	3.95	1.22	3.15	4.20	5.55
		±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
		0.47b	0.13b	0.05b	0.35b	0.13b	0.15b	0.30b	0.30a	0.2b	0.29b	0.36b	0.07a	0.16b	0.28b	0.23b	0.21b	0.38b
	SRB02 + <i>F. oxy</i>	7.45	6.57	1.47	6.54	6.48	17.69	12.59	3.90	7.60	7.83	11.37	0.62	7.83	2.43	6.57	8.38	10.55
		±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
		0.23a	0.47a	0.20a	0.44a	0.38a	0.22a	0.38a	0.5b	0.37a	0.27a	0.64a	0.04b	0.19a	0.17a	0.39a	0.25a	0.38a

2 **Notes:** Asp, Aspartic acid; Thr, Threonine; Met, Methionine; ILE, Isoleucine; Ser, Serine; Glu, Glutamic acid; Leu, Leucine; Tyr,  
3 Tyrosine; Gly, Glycine; Phe, Phenylalanine; Lys, Lysine; Cys, Cysteine; Val, Valine; His, Histidine; Arg, Arginine; Ala, Alanine; Pro,  
4 Proline. Each value represents means ± SD of three independent experiments. Each values in columns followed by different letters are  
5 significantly different at  $P \geq 0.05$ .