

Injures and molting interference in a trilobite from the Cambrian Furongian of South China (#55705)

1

First submission

Guidance from your Editor

Please submit by **19 Dec 2020** for the benefit of the authors (and your \$200 publishing discount) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

3 Figure file(s)



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Injures and molting interference in a trilobite from the Cambrian Furongian of South China

Ruiwen Zong^{Corresp. 1}

¹ State Key Laboratory of Biogeology and Environmental Geology, China University of Geosciences, Wuhan, China

Corresponding Author: Ruiwen Zong
Email address: zongruiwen@cug.edu.cn

An injured *Shergoldia laevigata* Zhu, Hughes & Peng 2007 (Trilobita, Asaphida) was collected from the Furongian of Guangxi, South China. The wounds occurred in the left pleural region possessing two marked V-shaped gaps in the thorax. It led to substantial transversal shortening of the left pleural segments, with barely perceptible traces of healing. This malformation is interpreted as a sub-lethal attack from unknown predator. The morphology of the wounds indicated that the predatory structure might have been the frontal appendage(s) of larger arthropod or arthropod-like organism. There were overlapped somites located in the front of the wounds, and slightly dislocated thoracic segments on the left part of the thorax, suggesting that the trilobite had experienced difficulties during the molting process. Freshly moulted trilobite had dragged forward the old exuvia causing the irregular arrangement of somites. This unusual trilobite specimen indicates that the injuries interfered with molting and other living activities.

Injures and molting interference in a trilobite from the Cambrian Furongian of South China

Ruiwen Zong

State Key Laboratory of Biogeology and Environmental Geology, China University of Geosciences, Wuhan, Hubei, China

Corresponding Author:

Ruiwen Zong

Lumo Road No. 388, Wuhan, Hubei, 430074, China

Email address: zongruiwen@cug.edu.cn

Abstract

An injured *Shergoldia laevigata* Zhu, Hughes & Peng 2007 (Trilobita, Asaphida) was collected from the Furongian of Guangxi, South China. The wounds occurred in the left pleural region possessing two marked V-shaped gaps in the thorax. It led to substantial transversal shortening of the left pleural segments, with barely perceptible traces of healing. This malformation is

interpreted as a sub-lethal attack from unknown predator. The morphology of the wounds indicated that the predatory structure might have been the frontal appendage(s) of larger arthropod or arthropod-like organism. There were overlapped somites located in the front of the wounds, and slightly dislocated thoracic segments on the left part of the thorax, suggesting that the trilobite had experienced difficulties during the molting process. Freshly moulted trilobite had dragged forward the old exuvia causing the irregular arrangement of somites. This unusual trilobite specimen indicates that the injuries interfered with molting and other living activities.

Introduction

Numerous trilobite exoskeleton deformities have been documented, including abnormal healing, hyperplasia, deformation, and missing or fractured somites. The causes of these deformities are usually thought to be injuries, developmental disorders and diseases (*Owen, 1985; Babcock, 1993; Pates et al., 2017; Bicknell & Pates, 2020*). The evaluation of injury caused by predator attack is most useful for proving the interactions between predators and trilobites, and for reconstructing the food chain and ecological structure in deep time (*Klomp maker et al., 2019*). Furthermore, such predatorial injuries are used to elucidate behavioral information (*Babcock & Robison, 1989; Babcock, 1993; Pates et al., 2017; Bicknell, Paterson & Hopkins, 2019*). The wounds caused by predators have mainly been detected on the edges of trilobites, especially in the thoraces and pygidia, and are generally considered to have been non-lethal (*Babcock, 2003, 2007*). Although numerous studies have evaluated injured trilobites (e.g., *Owen, 1985; Rudkin, 1985; Babcock, 1993, 2003, 2007; Zhu et al., 2007; Bicknell & Paterson, 2018; Bicknell & Pates, 2020; Bicknell & Holland, 2020*), the predators remain unidentified. The shapes of the trilobite wounds suggest that some predators may have been radiodonts (*Babcock & Robison, 1989; Babcock, 1993; Nedin, 1999; Bicknell & Pates, 2020; Bicknell & Holland,*

2020). Other predator candidates include cephalopods, echinoderms, fish, and other larger arthropods (Bruton, 1981; Briggs & Collins, 1988; Babcock, 1993; Fatka, Budil & Grigar, 2015).

It is also unclear whether injured trilobites experienced certain interferences with their activities of living. Although it has been inferred that injuries did interfere with daily activities, direct fossil records are rare. Herein, I discuss an injured *Shergoldia laevigata* Zhu, Hughes & Peng 2007 (Zhu, Hughes & Peng, 2007) from the Cambrian (Furongian) of Jingxi, Guangxi, South China. The exoskeleton wounds suggest that the predatory structure might have been the frontal appendages of larger arthropod or arthropod-like organism. In addition, the findings indicate that these injuries would have caused difficulties during the trilobite molting process, but did not cause molting failure.

Materials & Methods

The described *Shergoldia laevigata* specimen, housed in the State Key Laboratory of Biogeology and Environmental Geology, China University of Geoscience (Wuhan), was discovered from the Cambrian (Furongian)-aged Sandu Formation of Guole Town, Jingxi County, Guangxi Zhuang Autonomous Region, South China (Fig. 1) (Zhu, Hughes & Peng, 2007). The Sandu Formation is represented by thin-middle bedded calcareous mudstones, siltstones, and argillaceous banded limestones, which formed most probably in the uppermost part of the continental slope (Lerosey-Aubril, Zhu & Ortega-Hernández, 2017). The Sandu Formation is richly fossiliferous, containing abundant, well-preserved articulated trilobites (Han et al., 2000; Zhu, 2005; Zhu, Hughes & Peng, 2007, 2010), non-trilobite arthropods (Lerosey-Aubril, Ortega-Hernández & Zhu, 2013; Lerosey-Aubril, Zhu & Ortega-Hernández, 2017),

echinoderms (*Han & Chen, 2008; Chen & Han, 2013; Zamora et al., 2017; Zamora, Zhu & Lefebvre, 2013; Zhu, Zamora & Lefebvre, 2014*), brachiopods, graptolites (*Zhan et al., 2010*), hyolithids, cnidarians, algae, and some exceptionally preserved soft-bodied fossils (*Zhu et al., 2016*).

Trilobites from the Sandu Formation have been classified into at least 25 genera so far (*Zhu, 2005; Zhu, Hughes & Peng, 2007, 2010*); however, only six abnormal specimens have been found in this formation (five *Tamdaspis jingxiensis* and one *Guangxiaspis guangxiensis*) (*Zhu, 2005; Zhu et al., 2007; Zong, 2020*). The injured *Shergoldia laevigata* was collected from the grey-yellow calcareous mudstones. The age of the specimen is limited to the *Probinacunaspis nasalis*–*Peichiashania hunanensis* Zone of the Furongian Jiangshanian (*Peng, 2009; Zhu et al., 2016*).

The fossil in Fig. 2C was whitened with magnesium oxide powder, and all photographs were captured using a Nikon D5100 camera with a Micro-Nikkor 55 mm F3.5 lens. All figures were completed in CorelDRAW X7.

Results

The injured *Shergoldia laevigata* is preserved as a nearly complete dorsal exoskeleton (30.5 mm long) without librigena; such configuration was considered as an exuvia (*Daley & Drage, 2016; Drage, 2019*). The posterior of the cranium overlies the first three thoracic segments, especially on the left side of the body (Fig. 2). The malformation is present on the left part of exoskeleton, while the medial and right sections are normal. There are two wounds: one on the third to fourth thoracic segments, and one on the seventh thoracic segment. The left thoracic segments are shorter than those on the right side and show limited healing. The left pleural

segments are only about half as long as the normal segments because of the first wound; the most seriously damaged part is the contact site of the two thoracic segments, where there is a V-shaped wound. The second wound caused the left pleural part of the seventh thoracic segment to be half as long as normal, with the wound presenting as an asymmetric V-shape. The lack of regeneration (e.g., swelling, elongation, or bending of the thoracic tergites) suggests that these malformations occurred during the most recent post- or inter-molt stage.

Discussion

Possible origin of the wounds and potential predatory structure

Trilobites that are malformed due to predatory attacks have typical V-, U-, or W-shaped wounds (*Owen, 1985; Babcock, 1993; Pratt, 1998; Jago & Haines, 2002; Zamora et al., 2011; Pates et al., 2017; Bicknell & Paterson, 2018; Bicknell & Pates, 2020*), with a few showing in irregularly shaped wounds (*Fatka, Budil & Grigar, 2015*), in addition to certain signs of healing or regeneration (*Rudkins, 1979; Pates et al., 2017*). In the present specimen, the two wounds with traces of healing were classified as the result of predator attack. The two wounds have a similar degree of healing without any regeneration, suggesting that these injuries may have been incurred at the same time.

The most superior Cambrian predators are considered to have been the radiodonts, especially anomalocaridids and amplexobeluids, as their frontal appendages and oral cone were extremely effective predatory structures (*Whittington & Briggs, 1985; Babcock, 1993; Zamora et al., 2011*). These Cambrian predators preyed on mineralized and non-mineralized trilobites (*Nedin, 1999*). Cambrian arthropods or arthropod-like organisms with gnathobases are also considered possible predators, similar to the modern horseshoe crab (*Bicknell et al., 2018a*). Some amplexobeluid genera have been documented with gnathobase-like structures (*Cong et*

al., 2017, 2018), suggesting that amplexobeluid radiodonts may have been predators of Cambrian trilobites (Bicknell & Pates, 2020). In addition, some trilobites and predatory arthropods with reinforced gnathobasic spines on the protopodal sections of their walking legs are also considered as potential predators (Bruton, 1981; Conway Morris & Jenkins, 1985; Zacaï, Vannier & Lerosey-Aubril, 2016; Bicknell et al., 2018a, b; Bicknell & Holland, 2020; Bicknell, Paterson & Hopkins, 2019).

Despite these predatory advantages, Cambrian animals with gnathobases did not cause the wounds in the *Shergoldia laevigata* specimen. Gnathobases have a slight size gradation of spines along the gnathal edge (Stein, 2013), or a saw-toothed pattern with spines of alternating sizes (Bicknell et al., 2018a). However, the *Shergoldia laevigata* specimen had two V-shaped wounds and was missing one or two separate thoracic segments, which rules out the gnathobases of predators as a cause of injury. These wounds were instead cause by predatory structure similar to the frontal appendages of radiodonts. It is likely that the exoskeleton of the trilobite was attacked by a radiodont with variously sized ventral spines of the frontal appendages, which produced the different types of injuries (Fig. 3A-B). Ventral spines of various sizes have been discovered on the frontal appendages of some radiodonts, such as *Anomalocaris* (Wang, Huang & Hu, 2013). The two wounds may have been caused by the larger ventral spines of the frontal appendages, while the smaller ventral spines may have not damaged the *Shergoldia laevigata*. While the arthropods *Aglaspella sanduensis* (Lerosey-Aubril, Ortega-Hernández & Zhu, 2013) and *Glypharthrus trispinicaudatus*, *Mollisonia*-like arthropods, unnamed aglaspidid-like arthropods, and *Perspica*-like bivalve arthropods (Zhu et al., 2016; Lerosey-Aubril, Zhu & Ortega-Hernández, 2017) were discovered in the Sandu Formation at the same site, none of these have sufficiently strong frontal appendages. Therefore, the tracking down the predator who

attacked **on** studied *Shergoldia laevigata* specimen is still unsuccessful; it is not excluded that other, so far unknown arthropods or arthropod-like organisms with strong frontal appendages occurred in Sandu Formation.

Interference with the molting of trilobite

Pervious studies have reported abundant injured trilobites and presented the possible identity of the predators, including information about their behavior (*Babcock, 1993, 2007; Pates et al., 2017; Bicknell, Paterson & Hopkins, 2019*). However, there are few direct fossil records showing that injury has disturbed the molting of trilobites (*Šnajdr, 1985*). The trilobite specimen from Guangxi has an apparent overlap of tergites along with the wounds that are mainly present in the posterior of the cranidium and the front of the thorax, especially in the left part of the exoskeleton. The anterior margin of the third thoracic segment with part of the wound was covered by the second thoracic segment without breaking or being lost (*Fig. 2A-D*), **indicating that the wound formed before the overlap of the tergites**. The first thoracic segment covered most of the left pleural segment of the second thoracic segment, as well as the anterior margin of the right pleural segment. Similarly, most of the first thoracic segment was covered by the posterior area of the fixigena, particularly on its left side. The recuperative terminal of the posterior area of the fixigena covered the second thoracic segment (*Fig. 2E*). In addition, the left pleural segments of the fourth to eighth thoracic segments presented an interlaced arrangement, i.e. the anterior margin of the fourth thoracic segment extended upon the third thoracic segment, and the seventh extended upon the sixth (*Fig. 2*), while there was a typical imbricated arrangement in the right pleural region. **All abnormal arrangements of the segments appeared near the wound**, and the overlapped part of the somites was only located before the most serious wound (on the third to fourth thoracic segments). Based on the above mentioned descriptions, it is speculated that all of

the irregular patterns were caused by molting of *Shergoldia laevigata*. Namely, the new exoskeleton could not be smoothly separated from the old one due to the unbalanced body with wounds (Drage, 2019). The trilobite dragged forward the old shell to get rid of the exuvia, which led to the overlap of somites and the dislocated arrangement of thoracic segments, especially near the injuries (Fig. 3). Some previous studies have reported cases of failed molting of a trilobite (McNamara & Rudkin, 1984) and the arthropod *Marrella* (García-Bellido & Desmond, 2004), in which the new exoskeletons were preserved under the old exuvia. However, none of the fragments of the new exoskeleton were found under or near the exuvia of *Shergoldia laevigata*, which implies that the molting might not have failed. Although the wounds complicated the molting process, which, however, was successful and the molted trilobite left away.

Conclusions

The *Shergoldia laevigata* specimen has substantially shorter pleural segments of the third to fourth and seventh thoracic segments, with signs of lightly healing in the wounds incurred during a sub-lethal predator attack. The degree of healing in both wounds and the distribution of the wounds show that they were caused simultaneously by the same predatory structure. Based on the morphology of the wounds, the structure may have been the frontal appendage of larger arthropod or arthropod-like organism. The conspicuous overlapping of the somites and dislocated arrangement of the thoracic segments, especially in the left pleural region and near the wounds, shows that the injured *Shergoldia laevigata* encountered certain obstacles during the molting process. The trilobite dragged the old exuvia forward, which lead to the irregular arrangement of the somites. Such configuration can demonstrate that even provisionally healed injury can cause certain complication of the molting process in trilobites.

Acknowledgements

The author is very grateful to Guangchun Zeng (from Guangxi), Qi Liu (from Hunan) and Yonggang Tang (from Shandong) for their help in the field work and collection of specimens.

References

- Babcock LE. 1993.** Trilobite malformations and the fossil record of behavioral asymmetry. *Journal of Paleontology*, **67**: 217–229 DOI 10.1017/S0022336000032145.
- Babcock LE. 2003.** Trilobites in Palaeozoic predator-prey systems, and their role in reorganization of early Paleozoic ecosystems. In: Kelly PH, Kowalewski M, Hansen TA, eds. *Predator–Prey Interactions in the Fossil Record*. New York: Springer, 55–92
- Babcock LE. 2007.** Role of malformations in elucidating trilobite paleobiology: a historical synthesis. In: Mikulic DG, Landing E, Kluesendorf J, eds. *Fabulous Fossils—300 Years of Worldwide Research on Trilobites*. New York: The University of State of New York, 3–18
- Babcock LE, Robison RA. 1989.** Preferences of Palaeozoic predators. *Nature* **337**: 695–696 DOI 10.1038/337695c0.
- Bicknell RDC, Holland B. 2020.** Injured trilobites within a collection of dinosaurs: Using the Royal Tyrrell Museum of Palaeontology to document Cambrian predation. *Palaeontologia Electronica* **23**: a33 DOI 10.26879/1087.
- Bicknell RDC, Ledogar JA, Wroe S, Gutzler BC, Watson III WH, Paterson JR. 2018a.** Computational biomechanical analyses demonstrate similar shell-crushing abilities in modern and ancient arthropods. *Proceedings of the Royal Society B: Biological Sciences* **285**: 20181935. DOI 10.1098/rspb.2018.1935.

199 **Bicknell RDC, Paterson JR. 2018.** Reappraising the early evidence of durophagy and drilling
200 predation in the fossil record: implications for escalation and the Cambrian explosion.
201 *Biological Reviews* **93**: 754–784 DOI 10.1111/brv.12365.

202 **Bicknell RDC, Paterson JR, Caron JB, Skovsted CB. 2018b.** The gnathobasic spine
203 microstructure of Recent and Silurian chelicerates and the Cambrian arthropodan *Sidneyia*:
204 functional and evolutionary implications. *Arthropod Structure & Development* **47**: 12–24
205 DOI 10.1016/j.asd.2017.12.001.

206 **Bicknell RDC, Paterson JR, Hopkins MJ. 2019.** A trilobite cluster from the Silurian Rochester
207 Shale of New York: predation patterns and possible defensive behavior. *American Museum*
208 *Novitates* **39**: 1–16 DOI 10.1206/3937.1.

209 **Bicknell RDC, Pates S. 2020.** Exploring abnormal Cambrian-aged trilobites in the Smithsonian
210 collection. *PeerJ* **8**: e8453 DOI 10.7717/peerj.8453.

211 **Briggs DEG, Collins D. 1988.** A Middle Cambrian chelicerate from Mount Stephen, British
212 Columbia. *Palaeontology* **31**: 779–798.

213 **Bruton DL. 1981.** The arthropod *Sidneyia inexpectans*, Middle Cambrian, Burgess Shale,
214 British Columbia. *Philosophical Transactions of the Royal Society of London. Series B,*
215 *Biological Sciences* **295**: 619–656 DOI 10.1098/rstb.1981.0164.

216 **Chen GY, Han NR. 2013.** New materials of Stylophora from the Upper Cambrian of the Jingxi
217 area, Guangxi, South China. *Acta Palaeontologica Sinica* **52**: 288–293.

218 **Cong PY, Daley AC, Edgecombe GD, Hou, XG. 2017.** The functional head of the Cambrian
219 radiodontan (stem-group Euarthropoda) *Amplectobelua symbrachiata*. *BMC Evolutionary*
220 *Biology* **17**: 208. DOI 10.1186/s12862-017-1049-1.

221 **Cong PY, Edgecombe GD, Daley AC, Guo J, Pates S, Hou XG. 2018.** New radiodonts with
222 gnathobase-like structures from the Cambrian Chengjiang biota and implications for the
223 systematics of Radiodonta. *Papers in Palaeontology* **4**: 605–621 DOI 10.1002/spp2.1219.

224 **Conway Morris S, Jenkins RJF. 1985.** Healed injuries in Early Cambrian trilobites from South
225 Australia. *Alcheringa* **9**: 167–177 DOI 10.1080/03115518508618965.

226 **Daley AC, Drage HB. 2016.** The fossil record of ecdysis, and trends in the moulting behaviour
227 of trilobites. *Arthropod Structure & Development* **45**: 71–96 DOI
228 10.1016/j.asd.2015.09.004.

229 **Drage HB. 2019.** Quantifying intra- and interspecific variability in trilobite moulting behaviour
230 across the Palaeozoic. *Palaeontologia Electronica* **22.2.34**: 1–39 DOI 10.26879/940.

231 **Fatka O, Budil P, Grigar L. 2015.** A unique case of healed injury in a Cambrian trilobite.
232 *Annales de Paléontologie* **101**: 295–299 DOI 10.1016/j.annpal.2015.10.001.

233 **García-Bellido DC, Collins DH. 2004.** Moulting arthropod caught in the act. *Nature* **429**: 40
234 DOI 10.1038/429040a.

235 **Han NR, Chen GY. 2008.** New stylophorans (Echinodermata) from the Upper Cambrian of
236 Guangxi, South China. *Science China Series D: Earth Sciences* **51**: 181–186 DOI
237 10.1007/s11430-008-0003-x.

238 **Han NR, Tang L, Wei RS, Wang GB. 2000.** Stratigraphy of Upper Cambrian from Guole,
239 Jingxi, Guangxi. *Journal of Guilin Institute of Technology* **20**: 350–355.

240 **Jago JB, Haines PW. 2002.** Repairs to an injured early Middle Cambrian trilobite, Elkedra area,
241 Northern Territory. *Alcheringa* **26**: 19–21 DOI 10.1080/03115510208619241.

242 **Klomp maker AA, Kelley PH, Chattopadhyay D, Clements JC, Huntley JW, Kowalewski**
243 **M. 2019.** Predation in the marine fossil record: studies, data, recognition, environmental

244 factors, and behavior. *Earth-Science Reviews*, **194**: 472–520 DOI
 245 10.1016/j.earscirev.2019.02.020.

246 **Lerosey-Aubril R, Ortega-Hernández J, Zhu, XJ. 2013.** The first aglaspidid *sensu stricto*
 247 from the Cambrian of China (Sandu Formation, Guangxi). *Geological Magazine* **150**: 565–
 248 571 DOI 10.1017/S0016756812001045.

249 **Lerosey-Aubril R, Zhu XJ, Ortega-Hernández J. 2017.** The Vicissicaudata revisited--insights
 250 from a new aglaspidid arthropod with caudal appendages from the Furongian of China.
 251 *Scientific Reports* **7**: 11117 DOI 10.1038/s41598-017-11610-5.

252 **Mcnamara KJ, Rudkin DM. 1984.** Techniques of trilobite exuviations. *Lethaia* **17**: 153–173.

253 **Nedin C. 1999.** *Anomalocaris* predation on nonmineralized and mineralized trilobites. *Geology*
 254 **27**: 987–990 DOI 10.1130/0091-7613(1999)027%3C0987:aponam%3E2.3.co;2.

255 **Owen AW. 1985.** Trilobite abnormalities. *Transactions of the Royal Society of Edinburgh:*
 256 *Earth Sciences* **76**: 255–272 DOI 10.1017/s0263593300010488.

257 **Pates S, Bicknell RDC, Daley AC, Zamora S. 2017.** Quantitative analysis of repaired and
 258 unrepaired damage to trilobites from the Cambrian (Stage 4, Drumian) Iberian Chains, NE
 259 Spain. *Palaaios* **32**: 750–761 DOI 10.2110/palo.2017.055.

260 **Peng SC. 2009.** Review on the studies of Cambrian trilobite faunas from Jiangnan slope belt,
 261 South China, with notes on Cambrian correlation between South and North China. *Acta*
 262 *Palaeontologica Sinica* **48**: 437–452.

263 **Pratt BR. 1998.** Probable predation on Upper Cambrian trilobites and its relevance for the
 264 extinction of soft-bodied Burgess Shale-type animals. *Lethaia* **31**: 73–88 DOI
 265 10.1111/j.1502-3931.1998.tb00493.x.

266 **Rudkin DM. 1979.** Healed injuries in *Ogygopsis klotzi* (Trilobita) from the Middle Cambrian of
267 British Columbia. *Royal Ontario Museum, Life Sciences Occasional Paper* 32: 1–8.

268 **Rudkin DM. 1985.** Exoskeletal abnormalities in four trilobites. *Canadian Journal of Earth*
269 *Sciences* **22**: 479–483 DOI 10.1139/e85-047.

270 **Šnajdr M. 1985.** Anomalous exoskeletons of Bohemian encrinurine trilobites. *Vestník*
271 *Úsredního Ústavu Geologického* **60**: 303–306.

272 **Stein M. 2013.** Cephalic and appendage morphology of the Cambrian arthropod *Sidneyia*
273 *inexpectans*. *Zoologischer Anzeiger* **253**: 164–178 DOI 10.1016/j.jcz.2013.05.001.

274 **Wang YY, Huang DY, Hu SX. 2013.** New anomalocardid frontal appendages from the
275 Guanshan biota, eastern Yunnan. *China Science Bulletin* **58**: 3937–3942 DOI
276 10.1007/s11434-013-5908-x.

277 **Whittington HB, Briggs DEG. 1985.** The largest Cambrian animal, Anomalocaris, Burgess
278 Shale, British Columbia. *Philosophical Transactions of the Royal Society of London. Series*
279 *B, Biological Sciences* **309**: 569–609 DOI 10.1098/rstb.1985.0096.

280 **Zacai A, Vannier J, Lerosey-Aubril R. 2016.** Reconstructing the diet of a 505-million-year-old
281 arthropod: *Sidneyia inexpectans* from the Burgess Shale fauna. *Arthropod Structure &*
282 *Development* **45**: 200–220 DOI 10.1016/j.asd.2015.09.003.

283 **Zamora S, Mayoral E, Esteve J, Gámez-Vintaned JA, Santos A. 2011.** Exoskeletal
284 abnormalities in paradoxid trilobites from the Cambrian of Spain, and a new type of bite
285 trace. *Bulletin of Geosciences* **86**: 665–673 DOI 10.3140/bull.geosci.1275.

286 **Zamora S, Sumrall C, Zhu, XJ, Lefebvre B. 2017.** A new stemmed echinoderm from the
287 Furongian of China and the origin of Glyptocystitida (Blastozoa, Echinodermata).
288 *Geological Magazine* **154**: 465–475 DOI 10.1017/S001675681600011X.

289 **Zamora S, Zhu XJ, Lefebvre B. 2013.** A new Furongian (Cambrian) echinoderm-Lagerstätte
290 from the Sandu Formation (South China). *Cahiers de Biologie Marine* **54**: 565–569.

291 **Zhan RB, Jin JS, Rong JY, Zhu XJ, Han NR. 2010.** Late Cambrian brachiopods from Jingxi,
292 Guangxi Province, South China. *Alcheringa* **34**: 99–133 DOI
293 10.1080/03115510903522872.

294 **Zhu XJ. 2005.** Trilobite faunas from Cambrian Upper Furongian of Guangxi with special notes
295 on malformation, dimorphism and the function of eye ridges. Ph.D. Thesis, Nanjing
296 Institute of Geology and Palaeontology, Chinese Academy of Sciences, Nanjing.

297 **Zhu XJ, Hughes NC, Peng SC. 2007.** On a new species of *Shergoldia* Zhang, Jell, 1987
298 (Trilobita), the family Tsinaniidae and the order Asaphida. *Memoirs of the Association of*
299 *Australasian Palaeontologists* **34**: 243–253.

300 **Zhu XJ, Hughes NC, Peng SC. 2010.** Ventral structure and ontogeny of the late Furongian
301 (Cambrian) trilobite *Guangxiaspis guangxiensis* Zhou, 1977 and the diphyletic origin of the
302 median suture. *Journal of Paleontology* **84**: 493–504 DOI 10.1666/09-070.1.

303 **Zhu XJ, Peng SC, Du SX, Hu YS. 2007.** Ontogeny and malformation of *Tamdaspis jingxiensis*
304 sp. nov. (trilobite, Cambrian) from Jingxi, Guangxi, China. *Acta Palaeontologica Sinica* **46**:
305 225–231.

306 **Zhu XJ, Peng SC, Zamora S, Lefebvre B, Chen GY. 2016.** Furongian (upper Cambrian)
307 Guole Konservat-Lagerstätte from South China. *Acta Geologica Sinica (English Edition)* **90**:
308 30–37 DOI 10.1111/1755-6724.12640.

309 **Zhu XJ, Zamora S, Lefebvre B. 2014.** Morphology and palaeoecology of a new edrioblastoid
310 (Edrioasteroidea) from the Furongian of China. *Acta Palaeontologica Polonica* **59**: 921–
311 926 DOI 10.4202/app.2012.0116.

312 **Zong RW. 2020.** Abnormalities in early Paleozoic trilobites from central and eastern China.
 313 *Palaeoworld* **in press** DOI 10.1016/j.palwor.2020.07.003.

Figure 1

(A) Map of fossil locality at Guole Town, Jingxi County, Guangxi, South China; (B) Stratigraphic sketch showing relative position and age of the Sandu Formation.




Figure 2

Malformed trilobite *Shergoldia laevigata* from the Cambrian Furongian of Jingxi, Guangxi (Specimen No. CUG-GJ-2015-01).

(A) The uncoated specimen; (B) close-up of abnormality in box in figure (A); (C) the specimen whitened by the magnesium powder; (D) a sketch of the figure (A); (E) the picture after recovery of the cranidium and the first three thoracic segments, showing the superposed relationship between the posterior area of the fixigena and thoracic segments.




Figure 3

Reconstruction of injured of studied *Shergoldia laevigata* specimen and possible predatory structure of unknown predator from the Cambrian Furongian of Jingxi, Guangxi.

(A–B) Predatory frontal appendage with variable-sized ventral spines attacking on *Shergoldia laevigata*, and leading to damage in the exoskeleton. (C) *Shergoldia laevigata* drag forward the old shell during molting, because of the deformation of the exoskeleton. Such condition leads to the overlap of somites and dislocated arrangement of thoracic segments.

