

Injuries and molting interference in a trilobite from the Cambrian (Furongian) of South China (#55705)

1

First revision

Guidance from your Editor

Please submit by **6 Mar 2021** for the benefit of the authors .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

1 Tracked changes manuscript(s)

1 Rebuttal letter(s)

3 Figure file(s)



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Injuries and molting interference in a trilobite from the Cambrian (Furongian) of South China

Ruiwen Zong^{Corresp. 1}

¹ State Key Laboratory of Biogeology and Environmental Geology, China University of Geosciences, Wuhan, China

Corresponding Author: Ruiwen Zong
Email address: zongruiwen@cug.edu.cn

An injured *Shergoldia laevigata* Zhu, Hughes & Peng 2007 (Trilobita, Asaphida) was collected from the Furongian of Guangxi, South China. The injuries occurred in the left thoracic pleurae possessing two marked V-shaped gaps. It led to substantial transverse shortening of the left pleural segments, with barely perceptible traces of healing. This malformation is interpreted as a sub-lethal attack from an unknown predator. The morphology of the injuries and the spatial and temporal distribution of predators indicated that the predatory structure might have been the ganathobase or ganathobase-like structure of a larger arthropod. There were overlapped somites located in the front of the injuries, and slightly dislocated thoracic segments on the left part of the thorax, suggesting that the trilobite had experienced difficulties during the molting process. The freshly molted trilobite had dragged forward the old exuvia causing the irregular arrangement of somites. This unusual trilobite specimen indicates that the injuries interfered with molting.

Injuries and molting interference in a trilobite from the Cambrian (Furongian) of South China

Ruiwen Zong

State Key Laboratory of Biogeology and Environmental Geology, China University of Geosciences, Wuhan, Hubei, China

Corresponding Author:

Ruiwen Zong

Lumo Road No. 388, Wuhan, Hubei, 430074, China

Email address: zongruiwen@cug.edu.cn

Abstract

An injured *Shergoldia laevigata* Zhu, Hughes & Peng 2007 (Trilobita, Asaphida) was collected from the Furongian of Guangxi, South China. The injuries occurred in the left thoracic pleurae possessing two marked V-shaped gaps. It led to substantial transverse shortening of the left pleural segments, with barely perceptible traces of healing. This malformation is interpreted as a sub-lethal attack from an unknown predator. The morphology of the injuries and the spatial and temporal distribution of predators indicated that the predatory structure might have been the ganathobase or ganathobase-like structure of a larger arthropod. There were overlapped somites located in the front of the injuries, and slightly dislocated thoracic segments on the left

part of the thorax, suggesting that the trilobite had experienced difficulties during ~~the~~ molting process. The freshly molted trilobite had dragged forward the old exuvia causing the irregular arrangement of somites. This unusual trilobite specimen indicates that the injuries interfered with molting.

Introduction

Numerous trilobite exoskeleton deformities have been documented, including abnormal healing, hyperplasia, deformation, and missing or fractured somites. The causes of these deformities are usually thought to be injuries, developmental disorders, and diseases (*Owen, 1985; Babcock, 1993; Pates et al., 2017; Bicknell & Pates, 2020*). The evaluation of injuries caused by predator attack is useful for presenting the interactions between predators and trilobites, and for reconstructing the food web and ecological structure in deep time (*Klompaker et al., 2019*). Furthermore, such predatorial injuries are used to uncover behavioral information (*Babcock & Robison, 1989; Babcock, 1993; Pates et al., 2017; Bicknell, Paterson & Hopkins, 2019*). The injuries caused by predators have mainly been detected on the edges of trilobites, especially in the thoraces and pygidia, and are generally considered to have been non-lethal (*Babcock, 2003, 2007*), while cephalic attacks are more often fatal (*Pates & Bicknell, 2019*). Although numerous studies have evaluated injured trilobites (e.g., *Owen, 1985; Rudkin, 1985; Babcock, 1993, 2003, 2007; Zhu et al., 2007; Schoenemann, Clarkson & Høyberget, 2017; Bicknell & Paterson, 2018; Bicknell & Pates, 2020; Bicknell & Holland, 2020; Zong, 2020*), most predators remain unidentified, except some carnivores with trilobite fragments in their guts or coprolites (*Vannier & Chen, 2005; Vannier, 2012; Zacaï, Vannier & Lerosey-Aubril, 2016; Bicknell & Paterson, 2018; Kimmig & Pratt, 2018*). The shapes of the Cambrian trilobite injuries suggest that some predators may have been radiodonts (*Babcock & Robison, 1989; Babcock,*

1993; Nedin, 1999). Other predator candidates include cephalopods, echinoderms, fish, and other larger arthropods (Bruton, 1981; Briggs & Collins, 1988; Babcock, 1993; Fatka, Budil & Grigar, 2015; Jago, García-Bellido & Gehling, 2016; Bicknell & Paterson, 2018; Bicknell et al., 2018a; Zhai et al., 2019).

Moreover, although it has been inferred that injuries did interfere with daily activities of trilobites, direct fossil records are rare (Šnajdr, 1985). Herein, I discuss an injured *Shergoldia laevigata* Zhu, Hughes & Peng 2007 from the Cambrian (Furongian) of Jingxi, Guangxi, South China. The exoskeletal injuries suggest that the predatory structure might have been the gnathobase or gnathobase-like structure of a larger arthropod. In addition, the findings indicate that these injuries would have caused difficulties for trilobite during molting, but did not cause molting failure.

Materials & Methods

The described *Shergoldia laevigata* specimen, housed in the State Key Laboratory of Biogeology and Environmental Geology, China University of Geoscience (Wuhan), was discovered from the Cambrian (Furongian)-aged Sandu Formation of Guole Town, Jingxi County, Guangxi Zhuang Autonomous Region, South China (Fig. 1) (Zhu, Hughes & Peng, 2007). The Sandu Formation is represented by calcareous mudstones, siltstones, and argillaceous banded limestones, which formed most probably in the uppermost part of the continental slope (Lerosey-Aubril, Zhu & Ortega-Hernández, 2017). The Sandu Formation is richly fossiliferous, containing abundant, well-preserved articulated trilobites (Han et al., 2000; Zhu, 2005; Zhu, Hughes & Peng, 2007, 2010), non-trilobite arthropods (Lerosey-Aubril, Ortega-Hernández & Zhu, 2013; Lerosey-Aubril, Zhu & Ortega-Hernández, 2017), echinoderms (Han & Chen, 2008; Chen & Han, 2013; Zamora et al., 2017; Zamora, Zhu & Lefebvre, 2013; Zhu, Zamora

& Lefebvre, 2014), brachiopods, graptolites (Zhan et al., 2010), hyolithids, cnidarians, algae, and some exceptionally preserved soft-bodied fossils (Zhu et al., 2016).

Trilobites from the Sandu Formation have been classified into at least 25 genera (Zhu, 2005; Zhu, Hughes & Peng, 2007, 2010); however, only six abnormal specimens have been documented from this formation (five *Tamdaspis jingxiensis* Zhu et al., 2007 and one *Guangxiaspis guangxiensis* Zhou in Zhou et al, 1977) (Zhu, 2005; Zhu et al., 2007; Zong, 2020). The injured *Shergoldia laevigata* was collected from the grey-yellow calcareous mudstones. The specimen is from the *Probinacunaspis nasalis*–*Peichiashania hunanensis* Zone of the Furongian, Jiangshanian (Peng, 2009; Zhu et al., 2016).

The fossil in Fig. 2C was whitened with magnesium oxide powder, and all photographs were captured using a Nikon D5100 camera with a Micro-Nikkor 55 mm F3.5 lens.

Results

The injured *Shergoldia laevigata* is preserved as a nearly complete dorsal exoskeleton (30.5 mm long) without librigena, suggesting an exuvia (Daley & Drage, 2016; Drage, 2019). The posterior of the cranidium overlies the first two thoracic segments, this is most pronounced on the left side (Fig. 2). In addition, the first thoracic segment covered most of the left pleural segment of the second thoracic segment, as well as the anterior margin of the right pleural segment. Similarly, most of the first thoracic segment was covered by the posterior area of the fixigena, particularly on its left side. Moreover, the left pleural segments of the fourth to eighth thoracic segments presented an interlaced arrangement, i.e. the anterior margin of the fourth thoracic segment extended upon the third thoracic segment, and the seventh extended upon the sixth (Fig. 2), while there was a typical imbricated arrangement in the right pleural region.

The malformation is on the left part of the exoskeleton, while the medial (axial) and right sections are undamaged. The left thoracic segments are shorter than those on the right side and show limited healing. There are two injuries: one on the third to fourth thoracic segments, and one on the seventh thoracic segment. Two pleural segments are truncated by 3.3 mm because of the first injury; the most seriously damaged part is the contact site of the two thoracic segments, where there is a V-shaped injury. The second injury truncates the left pleural section of the seventh thoracic segment by 3.5 mm, ~~with the injury presenting as an asymmetric V-shape.~~

Discussion

Possible origin of the injuries and potential predatory structure

Trilobites that are malformed due to predatory attacks have typically V-, U-, or W-shaped injuries (*Owen, 1985; Babcock, 1993; Pratt, 1998; Jago & Haines, 2002; Zamora et al., 2011; Pates et al., 2017; Bicknell & Paterson, 2018; Bicknell & Pates, 2020*), with a few showing in irregularly shaped injuries (*Fatka, Budil & Grigar, 2015*). Furthermore, there is occasionally signs of healing or regeneration (*Rudkin, 1979; Mcnamara & Tuura, 2011; Pates et al., 2017*). In the present specimen, the injuries have traces of healing and are therefore considered evidence of a predatory attack. The two injuries have a similar degree of healing without any regeneration, suggesting that these injuries may have been incurred in the same inter-molt stage.

In the past, the most suspicious Cambrian predators are considered to have been the radiodonts, especially anomalocaridids and amplexobeluids, as their frontal appendages and oral cone were extremely effective predatory structures (*Whittington & Briggs, 1985; Babcock, 1993; Zamora et al., 2011*). Cambrian arthropods or arthropod-like organisms with gnathobases are also considered possible predators, similar to the modern horseshoe crab (*Bicknell et al., 2018a, 2021*). Some amplexobeluid genera have been documented with gnathobase-like structures

(*Cong et al., 2017, 2018*), suggesting that amplexobeluid radiodonts may have been predators of Cambrian trilobites (*Bicknell & Pates, 2020*). In addition, some trilobites and predatory arthropods with reinforced gnathobasic spines on the protopodal sections of their walking legs are also considered as potential predators (*Bruton, 1981; Conway Morris & Jenkins, 1985; Zacaï, Vannier & Lerosey-Aubril, 2016; Bicknell et al., 2018a, b; Bicknell, Paterson & Hopkins, 2019; Bicknell & Holland, 2020*).

However, ~~so far,~~ the youngest amplexobeluid and anomalocaridid are from the Drumian and Guzhangian, respectively (*Lerosey-Aubril et al., 2014, 2020*). Most Furongian and Ordovician radiodonts belong to the family Hurdidiidae, ~~which do not have endites of alternating size, and all members of this family are considered to be~~ sediment sifters or suspension feeders (*Daley, Budd & Caron, 2013; Daley et al., 2013; Lerosey-Aubril & Pates, 2018; Van Roy, Daley & Briggs, 2015; Pates et al., 2020*). Moreover, no radiodonts were discovered in the Sandu Formation. So, the radiodonts unlikely cause the injuries in the *Shergoldia laevigata* specimen. Gnathobases have a slight size gradation of spines along the gnathal edge (*Stein, 2013*), or a saw-toothed pattern with spines of alternating sizes (*Bicknell et al., 2018a*). These spines may caused smaller injuries, that is missing one or two separate thoracic segments, on the edges of trilobites.

The arthropods *Aglaspella sanduensis* (*Lerosey-Aubril, Ortega-Hernández & Zhu, 2013*) and *Glypharthrus trispinicaudatus*, *Mollisonia*-like arthropods, unnamed aglaspideid-like arthropods, *Perspicaris*-like bivalve arthropods (*Zhu et al., 2016; Lerosey-Aubril, Zhu & Ortega-Hernández, 2017*), and some larger trilobites (*Zhu, 2005*) were discovered in the Sandu Formation at the same site. Therefore, the predator who attacked the studied *Shergoldia laevigata* specimen maybe one of these arthropods.

Interference with the molting of trilobite

Previous studies have reported abundant injured trilobites and presented the possible identity of the predators, including information about their behavior (*Babcock, 1993, 2007; Pates et al., 2017; Bicknell & Paterson, 2018; Bicknell, Paterson & Hopkins, 2019; Pates & Bicknell, 2019*). However, there are few direct fossil records showing that injury has disturbed the molting of trilobites (*Šnajdr, 1985*). The studied specimen has an apparent overlap of somites along with the injuries that are mainly present in the posterior of the cranidium and the front of the thorax, especially in the left part of the exoskeleton. The anterior margin of the injured third thoracic segment was covered by the unbroken second thoracic segment (*Fig. 2A-D*), indicating that the injury formed before the overlap of the somites. Bottom currents can also caused the overlap and even disruption of trilobite somites, the Sandu Formation formed in a relatively calm environment (*Lerosey-Aubril, Zhu & Ortega-Hernández, 2017*), although there are overlapping segments on the exuvia and carcass of *Shergoldia laevigata* in the same horizon, their thoracic segments still maintain imbricated arrangement (*Zhu, Hughes & Peng, 2007*). There is also overlap of thoracic segments on the exuvia of uninjured trilobites (*Daley & Drage, 2016*), but it is difficult to determine whether it was caused by molting or other abiotic factors. In contrast to the above two cases, in addition to the overlap of somites in the studied specimen, the left thoracic segments are presented an interlaced arrangement rather than imbricated (*Fig. 2*), which seems not to be caused by bottom currents and is rather likely caused by the active behavior of trilobite. Moreover, all abnormal arrangements of the segments appeared near the injury, and the overlapped part of the somites was only located before the most serious injury (on the third to fourth thoracic segments). It is speculated that all of the irregular patterns were caused by post-injury molting of *Shergoldia laevigata*. Namely, the new exoskeleton could not be smoothly separated from the old one due to the unbalanced body with injuries (*Drage, 2019; Drage et al.,*

2019). The trilobite dragged forward the old shell to get rid of the exuvia, which led to the overlap of somites and the dislocated arrangement of thoracic segments, especially near the injuries (Fig. 3). Some previous studies have reported cases of failed molting of a trilobite (McNamara & Rudkin, 1984) and other ecdysozoans (García-Bellido & Desmond, 2004; Drage & Daley, 2016; Yang et al., 2019), in which the new exoskeletons were preserved under the old exuvia. However, none of the fragments of the new exoskeleton were found under or near the exuvia of *Shergoldia laevigata*, which implies that the molting might not have failed. Although the injuries complicated the molting process, it was successful and the molted trilobite moved away.

Conclusions

The *Shergoldia laevigata* specimen has substantially shorter pleural segments of the third to fourth and seventh thoracic segments, with signs of lightly healing in the injuries incurred during a sub-lethal predator attack. The degree of healing in both injuries and the distribution of the injuries show that they may have been caused in the same inter-molt stage. Based on the morphology of the injuries and the spatial and temporal distribution of predators, the predatory structure may have been the gnathobase or gnathobase-like structure of a larger arthropod. The conspicuous overlapping of the somites and dislocated arrangement of the thoracic segments, especially in the left pleural region and near the injuries, shows that the injured *Shergoldia laevigata* encountered certain obstacles during the molting process. The trilobite dragged the old exuvia forward, which lead to the irregular arrangement of the somites. Such configuration can demonstrate that even provisionally healed injury can cause certain complication of the molting process in trilobites.

Acknowledgements

I appreciate much the constructive and critical comments from Russell Bicknell, Stephen Pates, John Foster and one anonymous reviewer, which aided in the further improvement of the manuscript. I would like to thank Guangchun Zeng (from Guangxi), Qi Liu (from Hunan) and Yonggang Tang (from Shandong) for their help in the field work and collection of specimens.

References

- Babcock LE. 1993.** Trilobite malformations and the fossil record of behavioral asymmetry. *Journal of Paleontology*, **67**: 217–229 DOI 10.1017/S0022336000032145.
- Babcock LE. 2003.** Trilobites in Palaeozoic predator-prey systems, and their role in reorganization of early Paleozoic ecosystems. In: Kelly PH, Kowalewski M, Hansen TA, eds. *Predator–Prey Interactions in the Fossil Record*. New York: Springer, 55–92.
- Babcock LE. 2007.** Role of malformations in elucidating trilobite paleobiology: a historical synthesis. In: Mikulic DG, Landing E, Kluessendorf J, eds. *Fabulous Fossils—300 Years of Worldwide Research on Trilobites*. New York: The University of State of New York, 3–18.
- Babcock LE, Robison RA. 1989.** Preferences of Palaeozoic predators. *Nature* **337**: 695–696 DOI 10.1038/337695c0.
- Bicknell RDC, Holland B. 2020.** Injured trilobites within a collection of dinosaurs: Using the Royal Tyrrell Museum of Palaeontology to document Cambrian predation. *Palaeontologia Electronica* **23**: a33 DOI 10.26879/1087.
- Bicknell RDC, Holmes JD, Edgecombe GD, Lasso SR, Ortega-Hernández J, Wroe S, Paterson JR. 2021.** Biomechanical analyses of Cambrian euarthropod limbs reveal their effectiveness in mastication and durophagy. *Proceedings of the Royal Society B: Biological Sciences* **288**: 20202075. DOI 10.1098/rspb.2020.2075.

- 205 **Bicknell RDC, Ledogar JA, Wroe S, Gutzler BC, Watson III WH, Paterson JR. 2018a.**
206 Computational biomechanical analyses demonstrate similar shell-crushing abilities in
207 modern and ancient arthropods. *Proceedings of the Royal Society B: Biological Sciences*
208 **285**: 20181935. DOI 10.1098/rspb.2018.1935.
- 209 **Bicknell RDC, Paterson JR. 2018.** Reappraising the early evidence of durophagy and drilling
210 predation in the fossil record: implications for escalation and the Cambrian explosion.
211 *Biological Reviews* **93**: 754–784 DOI 10.1111/brv.12365.
- 212 **Bicknell RDC, Paterson JR, Caron JB, Skovsted CB. 2018b.** The gnathobasic spine
213 microstructure of Recent and Silurian chelicerates and the Cambrian arthropodan *Sidneyia*:
214 functional and evolutionary implications. *Arthropod Structure & Development* **47**: 12–24
215 DOI 10.1016/j.asd.2017.12.001.
- 216 **Bicknell RDC, Paterson JR, Hopkins MJ. 2019.** A trilobite cluster from the Silurian Rochester
217 Shale of New York: predation patterns and possible defensive behavior. *American Museum*
218 *Novitates* **39**: 1–16 DOI 10.1206/3937.1.
- 219 **Bicknell RDC, Pates S. 2020.** Exploring abnormal Cambrian-aged trilobites in the Smithsonian
220 collection. *PeerJ* **8**: e8453 DOI 10.7717/peerj.8453.
- 221 **Briggs DEG, Collins D. 1988.** A Middle Cambrian chelicerate from Mount Stephen, British
222 Columbia. *Palaeontology* **31**: 779–798.
- 223 **Bruton DL. 1981.** The arthropod *Sidneyia inexpectans*, Middle Cambrian, Burgess Shale,
224 British Columbia. *Philosophical Transactions of the Royal Society of London. Series B,*
225 *Biological Sciences* **295**: 619–656 DOI 10.1098/rstb.1981.0164.
- 226 **Chen GY, Han NR. 2013.** New materials of Stylophora from the Upper Cambrian of the Jingxi
227 area, Guangxi, South China. *Acta Palaeontologica Sinica* **52**: 288–293.

228 **Cong PY, Daley AC, Edgecombe GD, Hou, XG. 2017.** The functional head of the Cambrian
229 radiodontan (stem-group Euarthropoda) *Amplectobelua symbrachiata*. *BMC Evolutionary*
230 *Biology* **17**: 208. DOI 10.1186/s12862-017-1049-1.

231 **Cong PY, Edgecombe GD, Daley AC, Guo J, Pates S, Hou XG. 2018.** New radiodonts with
232 gnathobase-like structures from the Cambrian Chengjiang biota and implications for the
233 systematics of Radiodonta. *Papers in Palaeontology* **4**: 605–621 DOI 10.1002/spp2.1219.

234 **Conway Morris S, Jenkins RJF. 1985.** Healed injuries in Early Cambrian trilobites from South
235 Australia. *Alcheringa* **9**: 167–177 DOI 10.1080/03115518508618965.

236 **Daley AC, Budd GE, Caron JB. 2013.** Morphology and systematics of the anomalocaridid
237 arthropod *Hurdia* from the Middle Cambrian of British Columbia and Utah. *Journal of*
238 *Systematic Palaeontology* **11**: 743–787 DOI 10.1080/14772019.2012.732723.

239 **Daley AC, Drage HB. 2016.** The fossil record of ecdysis, and trends in the moulting behaviour
240 of trilobites. *Arthropod Structure & Development* **45**: 71–96 DOI
241 10.1016/j.asd.2015.09.004.

242 **Daley AC, Paterson JR, Edgecombe GD, García-Bellido DC, Jago JB. 2013.** New anatomical
243 information on *Anomalocaris* from the Cambrian Emu Bay Shale of South Australia and a
244 reassessment of its inferred predatory habits. *Palaeontology* **56**: 971–990 DOI
245 10.1111/pala.12029.

246 **Drage HB. 2019.** Quantifying intra- and interspecific variability in trilobite moulting behaviour
247 across the Palaeozoic. *Palaeontologia Electronica* **22.2.34**: 1–39 DOI 10.26879/940.

248 **Drage HB, Daley AC. 2016.** Recognising moulting behaviour in trilobites by examining
249 morphology, development and preservation: Comment on Błażejowski et al. 2015.
250 *BioEssays* **38**: 981–990 DOI 10.1002/bies.201600027.

251 **Drage HB, Vandenbroucke TRA, Van Roy P, Daley AC. 2019.** Sequence of post-moult
252 exoskeleton hardening preserved in a trilobite mass moult assemblage from the Lower
253 Ordovician Fezouata Konservat-Lagerstätte, Morocco. *Acta Palaeontologica Polonica* **64**:
254 261–273 DOI 10.4202/app.00582.2018.

255 **Fatka O, Budil P, Grigar L. 2015.** A unique case of healed injury in a Cambrian trilobite.
256 *Annales de Paléontologie* **101**: 295–299 DOI 10.1016/j.annpal.2015.10.001.

257 **García-Bellido DC, Collins DH. 2004.** Moulting arthropod caught in the act. *Nature* **429**: 40
258 DOI 10.1038/429040a.

259 **Han NR, Chen GY. 2008.** New stylophorans (Echinodermata) from the Upper Cambrian of
260 Guangxi, South China. *Science China Series D: Earth Sciences* **51**: 181–186 DOI
261 10.1007/s11430-008-0003-x.

262 **Han NR, Tang L, Wei RS, Wang GB. 2000.** Stratigraphy of Upper Cambrian from Guole,
263 Jingxi, Guangxi. *Journal of Guilin Institute of Technology* **20**: 350–355.

264 **Jago JB, García-Bellido DC, Gehling JG. 2016.** An early Cambrian chelicerate from the Emu
265 Bay shale, south Australia. *Palaeontology* **1**–14 DOI 10.1111/pala.12243.

266 **Jago JB, Haines PW. 2002.** Repairs to an injured early Middle Cambrian trilobite, Elkedra area,
267 Northern Territory. *Alcheringa* **26**: 19–21 DOI 10.1080/03115510208619241.

268 **Kimmig J, Pratt BR. 2018.** Coprolites in the Ravens Throat River Lagerstätte of northwestern
269 Canada: implications for the Middle Cambrian food web. *Palaos* **33**: 125–140 DOI
270 10.2110/palo.2017.038.

271 **Klomp maker AA, Kelley PH, Chattopadhyay D, Clements JC, Huntley JW, Kowalewski**
272 **M. 2019.** Predation in the marine fossil record: studies, data, recognition, environmental

273 factors, and behavior. *Earth-Science Reviews*, **194**: 472–520 DOI
 274 10.1016/j.earscirev.2019.02.020.

275 **Lerosey-Aubril R, Hegna TA, Babcock LE, Bonino E, Kier C. 2014.** Arthropod appendages
 276 from the Weeks Formation Konservat-Lagerstätte: new occurrences of anomalocaridids in
 277 the Cambrian of Utah, USA. *Bulletin of Geosciences* **89**: 269–282 DOI
 278 10.3140/bull.geosci.1442.

279 **Lerosey-Aubril R, Kimmig J, Pates S, Skabelund J, Weug A, Ortega-Hernández J. 2020.**
 280 New exceptionally preserved panarthropods from the Drumian Wheeler
 281 Konservat-Lagerstätte of the House Range of Utah. *Papers in Palaeontology* **in press** DOI
 282 10.1002/spp2.1307.

283 **Lerosey-Aubril R, Ortega-Hernández J, Zhu, XJ. 2013.** The first aglaspidid *sensu stricto*
 284 from the Cambrian of China (Sandu Formation, Guangxi). *Geological Magazine* **150**: 565–
 285 571 DOI 10.1017/S0016756812001045.

286 **Lerosey-Aubril R, Pates S. 2018.** New suspension-feeding radiodont suggests evolution of
 287 microplanktivory in Cambrian macronekton. *Nature communications* **9**: 3774 DOI
 288 10.1038/s41467-018-06229-7.

289 **Lerosey-Aubril R, Zhu XJ, Ortega-Hernández J. 2017.** The Vicissicaudata revisited--insights
 290 from a new aglaspidid arthropod with caudal appendages from the Furongian of China.
 291 *Scientific Reports* **7**: 11117 DOI 10.1038/s41598-017-11610-5.

292 **Mcnamara KJ, Rudkin DM. 1984.** Techniques of trilobite exuviations. *Lethaia* **17**: 153–173.

293 **McNamara KJ, Tuura ME. 2011.** Evidence for segment polarity during regeneration in the
 294 Devonian asteropygine trilobite *Greenops widderensis*. *Journal of Paleontology* **85**: 106–
 295 110 DOI 10.2307/23019504.

296 **Nedin C. 1999.** *Anomalocaris* predation on nonmineralized and mineralized trilobites. *Geology*
 297 **27:** 987–990 DOI 10.1130/0091-7613(1999)027%3C0987:aponam%3E2.3.co;2.

298 **Owen AW. 1985.** Trilobite abnormalities. *Transactions of the Royal Society of Edinburgh:*
 299 *Earth Sciences* **76:** 255–272 DOI 10.1017/s0263593300010488.

300 **Pates S, Bicknell RDC. 2019.** Elongated thoracic spines as potential predatory deterrents in
 301 olenelline trilobites from the lower Cambrian of Nevada. *Palaeogeography,*
 302 *Palaeoclimatology, Palaeoecology* **519:** 295–306 DOI 10.31233/osf.io/2jr9c.

303 **Pates S, Bicknell RDC, Daley AC, Zamora S. 2017.** Quantitative analysis of repaired and
 304 unrepaired damage to trilobites from the Cambrian (Stage 4, Drumian) Iberian Chains, NE
 305 Spain. *Palaos* **32:** 750–761 DOI 10.2110/palo.2017.055.

306 **Pates S, Botting JP, McCobb LM, Muir LA. 2020.** A miniature Ordovician hurdiid from
 307 Wales demonstrates the adaptability of Radiodonta. *Royal Society Open Science* **7:** 200459
 308 DOI 10.1098/rsos.200459.

309 **Peng SC. 2009.** Review on the studies of Cambrian trilobite faunas from Jiangnan slope belt,
 310 South China, with notes on Cambrian correlation between South and North China. *Acta*
 311 *Palaeontologica Sinica* **48:** 437–452.

312 **Pratt BR. 1998.** Probable predation on Upper Cambrian trilobites and its relevance for the
 313 extinction of soft-bodied Burgess Shale-type animals. *Lethaia* **31:** 73–88 DOI
 314 10.1111/j.1502-3931.1998.tb00493.x.

315 **Rudkin DM. 1979.** Healed injuries in *Ogygopsis klotzi* (Trilobita) from the Middle Cambrian of
 316 British Columbia. *Royal Ontario Museum, Life Sciences Occasional Paper* **32:** 1–8.

317 **Rudkin DM. 1985.** Exoskeletal abnormalities in four trilobites. *Canadian Journal of Earth*
 318 *Sciences* **22:** 479–483 DOI 10.1139/e85-047.

319 **Schoenemann B, Clarkson ENK, Høyberget M. 2017.** Traces of an ancient immune system –
 320 how an injured arthropod survived 465 million years ago. *Scientific Reports* **7**: 40330 DOI
 321 10.1038/srep40330.

322 **Šnajdr M. 1985.** Anomalous exoskeletons of Bohemian encrinurine trilobites. *Vestník*
 323 *Ústředního Ústavu Geologického* **60**: 303–306.

324 **Stein M. 2013.** Cephalic and appendage morphology of the Cambrian arthropod *Sidneyia*
 325 *inexpectans*. *Zoologischer Anzeiger* **253**: 164–178 DOI 10.1016/j.jcz.2013.05.001.

326 **Van Roy P, Daley AC, Briggs DE. 2015.** Anomalocaridid trunk limb homology revealed by a
 327 giant filter-feeder with paired flaps. *Nature* **522**: 77–80 DOI 10.1038/nature14256.

328 **Vannier J. 2012.** Gut contents as direct indicators for trophic relationships in the Cambrian
 329 marine ecosystem. *PLoS One* **7**: e52200 DOI 10.1371/journal.pone.0052200.

330 **Vannier J, Chen JY. 2005.** Early Cambrian food chain: new evidence from fossil aggregates in
 331 the Maotianshan Shale biota, SW China. *Palaios* **20**: 3–26 DOI 10.2110/palo.2003.p03-40.

332 **Whittington HB, Briggs DEG. 1985.** The largest Cambrian animal, *Anomalocaris*, Burgess
 333 Shale, British Columbia. *Philosophical Transactions of the Royal Society of London. Series*
 334 *B, Biological Sciences* **309**: 569–609 DOI 10.1098/rstb.1985.0096.

335 **Yang J, Ortega-Hernández J, Drage HB, Du KS, Zhang XG. 2019.** Ecdysis in a stem-group
 336 euarthropod from the early Cambrian of China. *Scientific Reports* **9**: 5709 DOI
 337 10.1038/s41598-019-41911-w.

338 **Zacai A, Vannier J, Lerosey-Aubril R. 2016.** Reconstructing the diet of a 505-million-year-old
 339 arthropod: *Sidneyia inexpectans* from the Burgess Shale fauna. *Arthropod Structure &*
 340 *Development* **45**: 200–220 DOI 10.1016/j.asd.2015.09.003.

- Zamora S, Mayoral E, Esteve J, Gámez-Vintaned JA, Santos A. 2011.** Exoskeletal abnormalities in paradoxid trilobites from the Cambrian of Spain, and a new type of bite trace. *Bulletin of Geosciences* **86**: 665–673 DOI 10.3140/bull.geosci.1275.
- Zamora S, Sumrall C, Zhu, XJ, Lefebvre B. 2017.** A new stemmed echinoderm from the Furongian of China and the origin of Glyptocystitida (Blastozoa, Echinodermata). *Geological Magazine* **154**: 465–475 DOI 10.1017/S001675681600011X.
- Zamora S, Zhu XJ, Lefebvre B. 2013.** A new Furongian (Cambrian) echinoderm-Lagerstätte from the Sandu Formation (South China). *Cahiers de Biologie Marine* **54**: 565–569.
- Zhai DY, Edgecombe GD, Bond AD, Mai HJ, Hou XG, Liu Y. 2019.** Fine-scale appendage structure of the Cambrian trilobitomorph *Naraoia spinosa* and its ontogenetic and ecological implications. *Proceedings of the Royal Society B: Biological Sciences* **286**: 20192371 DOI 10.1098/rspb.2019.2371.
- Zhan RB, Jin JS, Rong JY, Zhu XJ, Han NR. 2010.** Late Cambrian brachiopods from Jingxi, Guangxi Province, South China. *Alcheringa* **34**: 99–133 DOI 10.1080/03115510903522872.
- Zhou TM, Liu YR, Meng XS, Sun ZH. 1977.** Class Trilobita, In: *Palaeontological atlas of Central and South China. 1, Early Palaeozoic*. Beijing: Geological Publishing House, 104–266.
- Zhu XJ. 2005.** Trilobite faunas from Cambrian Upper Furongian of Guangxi with special notes on malformation, dimorphism and the function of eye ridges. Ph.D. Thesis, Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, Nanjing.

- 362 **Zhu XJ, Hughes NC, Peng SC. 2007.** On a new species of *Shergoldia* Zhang, Jell, 1987
363 (Trilobita), the family Tsinaniidae and the order Asaphida. *Memoirs of the Association of*
364 *Australasian Palaeontologists* **34**: 243–253.
- 365 **Zhu XJ, Hughes NC, Peng SC. 2010.** Ventral structure and ontogeny of the late Furongian
366 (Cambrian) trilobite *Guangxiaspis guangxiensis* Zhou, 1977 and the diphyletic origin of the
367 median suture. *Journal of Paleontology* **84**: 493–504 DOI 10.1666/09-070.1.
- 368 **Zhu XJ, Peng SC, Du SX, Hu YS. 2007.** Ontogeny and malformation of *Tamdaspis jingxiensis*
369 sp. nov. (trilobite, Cambrian) from Jingxi, Guangxi, China. *Acta Palaeontologica Sinica* **46**:
370 225–231.
- 371 **Zhu XJ, Peng SC, Zamora S, Lefebvre B, Chen GY. 2016.** Furongian (upper Cambrian)
372 Guole Konservat-Lagerstätte from South China. *Acta Geologica Sinica (English Edition)* **90**:
373 30–37 DOI 10.1111/1755-6724.12640.
- 374 **Zhu XJ, Zamora S, Lefebvre B. 2014.** Morphology and palaeoecology of a new edrioblastoid
375 (Edrioasteroidea) from the Furongian of China. *Acta Palaeontologica Polonica* **59**: 921–
376 926 DOI 10.4202/app.2012.0116.
- 377 **Zong RW. 2020.** Abnormalities in early Paleozoic trilobites from central and eastern China.
378 *Palaeoworld* **in press** DOI 10.1016/j.palwor.2020.07.003.

Figure 1

(A) Map of fossil locality at Guole Town, Jingxi County, Guangxi, South China; (B) Stratigraphic sketch showing relative position and age of the Sandu Formation.




Figure 2

Malformed trilobite *Shergoldia laevigata* from the Cambrian Furongian of Jingxi, Guangxi (Specimen No. CUG-GJ-2015-01).

(A) Uncoated specimen; (B) close-up of abnormality in box in figure (A); (C) specimen whitened by the magnesium powder; (D) sketch of the figure (A); (E) picture after recovery of the cranidium and the first three thoracic segments, showing the superposed relationship between the posterior area of the fixigena and thoracic segments.




Figure 3

Reconstruction of injured of studied *Shergoldia laevigata* specimen from the Cambrian Furongian of Jingxi, Guangxi.

(A–B) Predator attack on *Shergoldia laevigata*, and leading to damage in the exoskeleton. (C) *Shergoldia laevigata* drag forward the old shell during molting, because of the deformation of the exoskeleton. Such condition leads to the overlap of somites and dislocated arrangement of thoracic segments.

