

Stable isotope compositions of precipitation over Central Asia (#55326)

1

First revision

Guidance from your Editor

Please submit by **25 Mar 2021** for the benefit of the authors .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

1 Tracked changes manuscript(s)

1 Rebuttal letter(s)

8 Figure file(s)

1 Table file(s)




Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor






 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).





Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).





BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Stable isotope compositions of precipitation over Central Asia

Junqiang Yao ^{Corresp., 1}, Xinchun Liu ^{Corresp., 1}, Wenfeng Hu ²

¹ Institute of Desert Meteorology, China Meteorological Administration, Urumqi, Xinjiang, China

² Fuyang Normal University, Fuyang, Xinjiang, China

Corresponding Authors: Junqiang Yao, Xinchun Liu
Email address: yaojq@idm.cn, liuxch@idm.cn

Central Asia is one of the driest regions in the world with a unique water cycle and a complex moisture transport process. However, there is little information on the precipitation $\delta^{18}\text{O}$ content in Central Asia. We compiled a precipitation $\delta^{18}\text{O}$ database from 47 meteorological stations across Central Asia to reveal its spatial-temporal characteristics. We determined the **relationships** between precipitation $\delta^{18}\text{O}$ and environmental variables and investigated the relationship between $\delta^{18}\text{O}$ and large-scale atmospheric circulation. The Central Asia meteoric water line was established as $\delta^2\text{H} = 7.30 \delta^{18}\text{O} + 3.12$ ($R^2=0.95$, $n=727$, $p<0.01$), and precipitation $\delta^{18}\text{O}$ ranged from +2‰ to -25.4‰ with a mean of -8.7‰. The precipitation $\delta^{18}\text{O}$ over Central Asia was related to environmental variables. The $\delta^{18}\text{O}$ had a significant positive correlation with temperature, and the $\delta^{18}\text{O}$ -temperature gradient ranged from 0.28‰/°C to 0.68‰/°C. However, the dependence of $\delta^{18}\text{O}$ on precipitation was unclear; a significant precipitation effect was only observed at the Zhangye and Teheran stations, showing $\delta^{18}\text{O}$ -precipitation gradients of 0.20‰/mm and -0.08‰/mm, respectively. Latitude was always significantly correlated with $\delta^{18}\text{O}$, when considering geographical controls on $\delta^{18}\text{O}$, and both latitude **and longitude** were significantly correlated with $\delta^{18}\text{O}$ in winter, with $\delta^{18}\text{O}/\text{LAT}$ and $\delta^{18}\text{O}/\text{ALT}$ **gradients** of -0.42‰/° and -0.001‰/m, respectively. The relationship between $\delta^{18}\text{O}$ and large-scale atmospheric circulation suggested that the moisture in Central Asia is mainly transported by westerly circulation and is indirectly affected by the Indian monsoon. Meanwhile, the East Asian monsoon may affect the precipitation $\delta^{18}\text{O}$ content in westerly and monsoon transition regions. These results improve our understanding of the precipitation $\delta^{18}\text{O}$ and moisture transport in Central Asia, as well as the paleoclimatology and hydrology

processes in Central Asia.

Stable isotope compositions of precipitation over

Central Asia

Junqiang Yao¹, Xinchun Liu¹, Wenfeng Hu²

1. Institute of Desert Meteorology, China Meteorological Administration, Urumqi 830002, China;

2. Fuyang Normal University, Fuyang 236000, China

Junqiang Yao ^{1*}

Institute of Desert Meteorology, China Meteorological Administration, Urumqi 830002

Email address: yaojq@idm.cn (J.Y); yaojq1987@126.com (J.Y)

Xinchun Liu ^{1*}

Institute of Desert Meteorology, China Meteorological Administration, Urumqi 830002

Email address: liuxinchun2001@163.com (X.L)

Abstract

Central Asia is one of the driest regions in the world with a unique water cycle and a complex moisture transport process. However, there is little information on the precipitation $\delta^{18}\text{O}$ content in Central Asia. We compiled a precipitation $\delta^{18}\text{O}$ database from 47 meteorological stations across Central Asia to reveal its spatial-temporal characteristics. We determined the relationships between precipitation $\delta^{18}\text{O}$ and environmental variables and investigated the relationship between $\delta^{18}\text{O}$ and large-scale atmospheric circulation. The Central Asia meteoric water line was established as $\delta^2\text{H} = 7.30 \delta^{18}\text{O} + 3.12$ ($R^2=0.95$, $n=727$, $p<0.01$), and precipitation $\delta^{18}\text{O}$ ranged from +2‰ to -25.4‰ with a mean of -8.7‰. The precipitation $\delta^{18}\text{O}$ over Central Asia was related to environmental variables. The $\delta^{18}\text{O}$ had a significant positive correlation with temperature, and the $\delta^{18}\text{O}$ -temperature gradient ranged from 0.28‰/°C to 0.68‰/°C. However, the dependence of $\delta^{18}\text{O}$ on precipitation was unclear; a significant precipitation effect was only observed at the Zhangye and Teheran stations, showing $\delta^{18}\text{O}$ -precipitation gradients of 0.20‰/mm and -0.08‰/mm, respectively. Latitude was always significantly correlated with $\delta^{18}\text{O}$, when considering geographical controls on $\delta^{18}\text{O}$, and both latitude and longitude were significantly correlated with $\delta^{18}\text{O}$ in winter, with $\delta^{18}\text{O}/\text{LAT}$ and $\delta^{18}\text{O}/\text{ALT}$ gradients of -0.42‰/° and -0.001‰/m, respectively. The relationship between $\delta^{18}\text{O}$ and large-scale atmospheric circulation suggested that the moisture in Central Asia is mainly transported by westerly circulation and is indirectly affected by the Indian monsoon. Meanwhile, the East Asian monsoon may affect the precipitation $\delta^{18}\text{O}$ content in westerly and monsoon transition regions. These results improve our understanding of the precipitation $\delta^{18}\text{O}$ and moisture transport in Central Asia, as well as the paleoclimatology and hydrology processes in Central Asia.

Introduction

Stable water isotopes, including $\delta^{18}\text{O}$, and $\delta^2\text{H}$, are critical indicators of global and regional water cycles and paleoclimatic investigations improve our understanding of hydrological and atmospheric processes (*Dansgaard, 1964; Craig and Gordon, 1965; Song et al., 2007; Sophocleous, 2002; Zhang et al., 2004; Yao et al., 2013*). Precipitation is a critical variable of the global hydrological cycles. The isotope composition of precipitation from different sources varies and can be used as natural tracers to determine the sources and moisture transport of water vapor as well as water cycle processes in different climatic regions (*Yamanaka et al., 2007; Li et al., 2012; Dansgaard, 1953; Chen, 2014; Zhang and Wang, 2016*).

In 1961, the International Atomic Energy Agency (IAEA), in conjunction with the World Meteorological Organization (WMO), initiated the Global Network of Isotopes in Precipitation (GNIP) on a global scale to monitor the stable isotope composition in monthly precipitation. The number of observation stations has since gradually increased. In terms of the spatial distribution, the GNIP observation stations are mainly concentrated in low-latitude wet regions, with few stations inland and in arid regions. For example, there are only ten GNIP stations that record $\delta^{18}\text{O}$ in Central Asia and surrounding regions, including three stations in the arid region of Northwest China (i.e., the Urumqi, Hetian, and Zhangye stations) (*Gu, 2012*). However, most recording stopped in the mid-1990s (*Liu et al., 2014*) and information on the stable isotope composition of precipitation in Central Asia and the surrounding regions is limited.

Since the 1980s, China has become expert in the observation and investigation of stable isotope compositions in precipitation, mostly focusing on the Tibetan Plateau (*Gao et al., 2011; Tian et al., 1997; Tian et al., 2001; Yao et al., 2006*). Similar to the GNIP database, the Tibetan Network for Isotopes in Precipitation (TNIP) was established to observe and monitor the Tibetan Plateau and the surrounding regions; the China Network of Isotopes in Precipitation and Rivers

(CHIRP) was developed for nationwide observation (*Yao et al., 2009; Liu et al., 2014*). In 2004, the Chinese Network of Isotopes in Precipitation (CHNIP) was started by the Chinese Ecosystem Research Network (CERN) (*Song et al., 2007*). However, there were only two CHNIP stations in Northwest China (the Fukang and Cele stations). An observation network was established in 2012 around the Tianshan Mountains (TSNIP) to investigate the types of precipitation isotopes in the high mountains over Central Asia, (*Wang et al., 2016*).

Few studies have investigated precipitation isotopes around the Tibetan Plateau and Northwest China. Using observations and simulations, *Yao et al. (2013)* systematically examined the spatial-temporal distribution of precipitation $\delta^{18}\text{O}$ over the Tibetan Plateau and determined its relationship with the mechanisms of the westerlies, Indian monsoons, and transitions in between.

Central Asia is one of the largest arid regions in the world and includes five Central Asian countries and the arid region of Northwest China. The topography is complex, with desert and oases coexisting. Precipitation is the main source of water for the mountainous regions but is unevenly distributed (*Chen, 2012; Yao et al., 2014*). The moisture source, transport path, and atmospheric processes are complex and are particularly sensitive to climate change in the unique mountain-basin structure of Central Asia (*Chen, 2014*). Few studies have reported the changes and mechanisms of moisture transport processes in Central Asia (*Huang et al., 2015, 2018; Shi et al., 2008; He et al., 2016; Chen et al., 2010; Yao et al., 2020*). Previous studies have mainly focused on the effects of spatial-temporal characteristics, compositions, and climatic controls on precipitation isotopes on a regional scale, such as in the Tianshan Mountains and at Urumqi station (*Yao et al., 1999; Kong et al., 2013, 2016; Liu et al., 2015; Wang et al., 2016, 2017; Li et al., 2017; Zhang and Wang, 2018*). However, only a few studies have been conducted on the composition of stable isotopes in the precipitation over the Central Asian region.

Central Asia has a unique water cycle and complex moisture transport processes. We compiled a database of information from 47 stations to evaluate precipitation $\delta^{18}\text{O}$ over Central Asia to reveal its spatial-temporal characteristics. We determined the relationships between precipitation $\delta^{18}\text{O}$ and local factors, such as geography (latitude and altitude) and climatic parameters (temperature and precipitation). We also investigated the relationship between $\delta^{18}\text{O}$ and large-scale atmospheric circulation.

Materials & Methods

We studied the precipitation isotopes data ($\delta^{18}\text{O}$ and $\delta^2\text{D}$) from Central Asia to comprehensively evaluate the processes and characteristics of the composition of the stable isotopes in the precipitation of that region.

From the observed monthly precipitation, $\delta^{18}\text{O}$ and $\delta^2\text{D}$ was obtained from 47 observation stations over Central Asia, including 12 stations of the GNIP, 22 stations of the TSNIP (*Wang et al., 2016*), two TNIP stations (*Yao et al., 2013*), two CHNIP stations (*Liu et al., 2014*), and nine stations from reference studies (*Figure 1 and Table 1*).

The Kabul station had the longest data record, with monthly data starting in the 1960s and continuous data for 1962-1964, 1967-1975, and 1982-1989. The Urumqi station kept monthly data starting in 1986 and had continuous data for 1961-1976, 1979-1981, 1986-1987, and 2000-2004. Among the stations from which data was collected, five GNIP stations (Teheran, Kabul, Astrakhan, Urumqi, and Zhangye stations) took samples for more than 10 years and data were available for interannual variability studies (*Figure 1*). Furthermore, eight GNIP stations (Teheran, Kabul, Saratov, Barabinsk, Astrakhan, Urumqi, Hetian, and Zhangye stations) were sampled for more than 30 months and data were available to evaluate the relationship of precipitation $\delta^{18}\text{O}$ with temperature and precipitation over Central Asia (*Figure 1*). Spatial patterns of observed precipitation $\delta^{18}\text{O}$ over Central Asia were available from all stations. All



precipitation $\delta^{18}\text{O}$ data were employed with respect to the Vienna standard mean ocean water (VSMOW) and were shown in precipitation amount-weighted values. In addition, meteorological variables, including precipitation amount and air temperature, were recorded at each observation station.

The precipitation $\delta^{18}\text{O}$ was determined by atmospheric characteristics and their dynamics. We selected three atmospheric characteristic indices to discuss their effects on the precipitation $\delta^{18}\text{O}$ over Central Asia, including the westerly circulation index (WCI), East Asian summer monsoon index (EASMI), and Indian monsoon index (IMI). Monthly EASMI and IMI were obtained from Dr. Jianping Li's webpage (<http://ljp.gcess.cn/dct/page/65610>) (Li et al., 2002; 2003).

Temperature averages are calculated for monthly/annual analyses, and the corresponding precipitation amounts are calculated as monthly/annual data for each observation station. A stepwise linear regression analysis technique was employed to fit the precipitation $\delta^{18}\text{O}$ with geographical parameters. The following geographical parameters were considered: altitude (ALT, m), latitude (LAT, °N), and longitude (LON, °E). The Pearson correlation coefficient was used to investigate the relationship between precipitation $\delta^{18}\text{O}$ and meteorological variables or atmospheric circulation indices.

Results and Discussion

Local Meteoric Water Lines (MWL)

The local meteoric water lines (MWLs) of precipitation, $\delta^2\text{H}$, and $\delta^{18}\text{O}$, provide important information on the water cycle, water vapor sources, and water transport (Jones et al. 2007; Lutz et al. 2011). A Central Asia MWL (CAMWL) was established as $\delta^2\text{H} = 7.30\delta^{18}\text{O} + 3.12$ ($R^2=0.95$, $n=727$, $p<0.01$) (Figure 2) based on 727 precipitation groups. The slope of the CAMWL was slightly lower than that of the global MWL (GMWL), which was eight (Craig,



1961) and the Chinese MWL of 7.48 (Liu et al., 2014). Differences in the GMWL slope often occur because of deviations in humidity at the source of moisture or evaporation. Central Asia is located in the Eurasian hinterland, and has a significant variation in the annual cycle and alternating dry or wet seasons. A low LMWL slope is associated with non-equilibrium conditions that affect falling raindrops during dry conditions (Liu et al., 2014), leading to the potential for significant sub-cloud evaporation.

The precipitation $\delta^{18}\text{O}$ over Central Asia ranged from +2‰ to -25.4‰ with a mean of -8.7‰, and $\delta^2\text{H}$ ranged from -4.2 ‰ to -191.4‰ with a mean of -67.1 ‰.

Seasonal variations in precipitation $\delta^{18}\text{O}$

The arid Central Asian region is affected by monsoons and westerlies, resulting in annual differences in climatic variables and precipitation $\delta^{18}\text{O}$. The variation in the annual air temperature represents a continental climate. The maximum and minimum temperatures occur in July and January, respectively. In this study, the temperature variation was unimodal, and the precipitation amount represented two distribution types (Figure 2). The three precipitation patterns observed in different regions are shown in Figure 3. (1) Maximum precipitation in occurred in the summer, with summer precipitation accounting for 41.4% of the total precipitation. These stations (including Hetian, Zhangye, Barabinsk, and Urumqi stations) are mainly distributed in northern Central Asia and are primarily influenced by the intensity and location of the westerly circulation. (2) Maximum precipitation values in winter and spring, with winter and spring precipitation accounting for 39.6% and 37.8% of the total precipitation, respectively, while the summer precipitation accounted for only 6.4% of the total precipitation. These stations (including the Kabul and Teheran stations) are mainly located in southern Central Asia and are primarily influenced by the Indian monsoon. (3) A well-distributed seasonal precipitation pattern was observed at Astrakhan and Saratov stations. These are present in

northwestern Central Asia and are primarily influenced by airflow from the Arctic. This result also highlighted the complexity of the spatial–temporal variations in precipitation in Central Asia.

The annual temperature and precipitation influenced the annual precipitation $\delta^{18}\text{O}$ cycle. The maximum $\delta^{18}\text{O}$ occurred from June to August (JJA). Maximum precipitation $\delta^{18}\text{O}$ at the Urumqi, Hetian, Barabinsk, and Teheran stations were -5.4‰, 0‰, 2‰, and -0.3‰ in August, respectively. The maximum values at the Zhangye and Kabul stations occurred in July. The maximum values occurred in June and September, respectively, at the Saratov and Astrakhan stations. All minimum values occurred from December to February (DJF). From January to July, the $\delta^{18}\text{O}$ values in precipitation were continuously high but gradually declined from August to December. The maximum and minimum values appeared in the summer and winter, respectively. Therefore, the $\delta^{18}\text{O}$ content in precipitation was higher in summer than in winter, and the seasonal pattern of the $\delta^{18}\text{O}$ in precipitation reflected the climate regime during the annual cycle. In winter, air masses were relatively cold and dry and the amount of precipitation was very low, while in summer, the opposite was true.

Spatial characteristics of precipitation $\delta^{18}\text{O}$

Figure 4 shows the spatial distribution of precipitation $\delta^{18}\text{O}$ throughout the year, and for summer and winter seasons at each station in Central Asia. The maximum value of the annual mean $\delta^{18}\text{O}$ (-1.47‰) was at Cele station, located in southern Xinjiang, and the minimum value (-18.6‰) was observed at Hami station. For the summer months, the maximum $\delta^{18}\text{O}$ value (2.8 ‰) was observed at Cele station, and the minimum value (-13.4‰) was observed at Altay station, located in northeastern Central Asia. The maximum $\delta^{18}\text{O}$ value (-7.7 ‰) during the winter months was observed at the Teheran station, located in southern Central Asia, and the minimum value (-35.3‰) was observed at the Dabancheng station.

There were remarkable spatial differences in precipitation $\delta^{18}\text{O}$ over Central Asia. The spatial distribution of precipitation $\delta^{18}\text{O}$ was mainly affected by the thermodynamics of water vapor condensation during the Rayleigh fractionation process and included meteorological elements, water vapor transport, and geographic factors (Dansgaard, 1964; Yurtsever and Gat, 1981; Rozanski et al., 2003).

Several studies have verified that altitude and latitude are the main geographic factors that affect changes in temperature and water vapor condensation, and can be referred to as geographic factor effects of precipitation $\delta^{18}\text{O}$ (Liu et al., 2009; Yao et al., 2013). The correlation of precipitation $\delta^{18}\text{O}$ with latitude, altitude, and longitude was investigated using partial correlation analysis, and the correlation coefficients (CC) were 0.53 ($p < 0.05$), 0.32 ($p < 0.05$), and 0.17, respectively, indicating that latitude is the most significant factor. In summer, latitude was significantly correlated with precipitation $\delta^{18}\text{O}$ ($\text{CC} = 0.54$, $p < 0.05$), while both latitude and longitude were significantly correlated with precipitation $\delta^{18}\text{O}$ in winter ($\text{CC} = 0.31$, $p < 0.05$; $\text{CC} = 0.67$, $p < 0.01$, respectively). For the annual $\delta^{18}\text{O}$, the $\delta^{18}\text{O}/\text{latitude}$ gradient was $-0.42\text{‰}/^\circ$, implying that the precipitation $\delta^{18}\text{O}$ was reduced by approximately 0.42‰ for every one-degree variation in latitude, which was larger than the gradient in China ($-0.22\text{‰}/^\circ$). Similarly, the $\delta^{18}\text{O}/\text{altitude}$ gradient was $-0.001\text{‰}/\text{m}$, which was lower than the global value of $-0.0022\text{‰}/\text{m}$ and the value of $-0.0016\text{‰}/\text{m}$ in China (Liu et al., 2009; Bowen and Wilkinson, 2002). In Central Asia, altitudes range from -18 to $4,200$ m, with a complex topography and various climatic characteristics.

Several researchers have constructed models to explain the relationship between precipitation $\delta^{18}\text{O}$ and geographic factors (Liu et al., 2009; Yao et al., 2013), including the Bowen and Wilkinson (BW) model (Bowen and Wilkinson, 2002). To confirm the effects of altitude and

latitude on precipitation $\delta^{18}\text{O}$ over Central Asia, we evaluated the relationship between precipitation $\delta^{18}\text{O}$ and geographical factors using stepwise regression analysis.

In Central Asia, geographical controls on precipitation $\delta^{18}\text{O}$ were best expressed by a stepwise linear regression, including altitude (ALT , m), latitude (LAT , °N) and longitude (LON , °E):

$$\delta^{18}\text{O} = (-0.179) \times LAT + (-0.022) \times LON + 0.00085 \times ALT \quad (R^2 = 0.51, p < 0.05) \quad (1)$$

The best model accounting for JJA $\delta^{18}\text{O}$ included both altitude (ALT , m) and latitude (LAT , °N):

$$\delta^{18}\text{O} = (-0.632) \times LAT + (-0.002) \times ALT + 23.851 \quad (R^2 = 0.64, p < 0.05) \quad (2)$$

The DJF $\delta^{18}\text{O}$ model was expressed by linear regression as:

$$\delta^{18}\text{O} = (-0.482) \times LAT + (-0.280) \times LON + (-0.001) \times ALT + 23.891 \quad (R^2 = 0.73, p < 0.01) \quad (3)$$

Our model comprehensively explained the effect of altitude, latitude, and longitude on precipitation $\delta^{18}\text{O}$ over Central Asia, and revealed that the variation in precipitation $\delta^{18}\text{O}$ depended on geographical factors at each station. These results showed that our models were suitable for use in Central Asia.

We used the aforementioned models to obtain the spatial distribution of precipitation $\delta^{18}\text{O}$ over Central Asia (Figure 5). The annual precipitation $\delta^{18}\text{O}$ gradually decreased as the latitude increased (Figure 5a). In the summer, higher $\delta^{18}\text{O}$ values were observed in southwest Central Asia and lower $\delta^{18}\text{O}$ values were observed in northern Central Asia and the Pan-Third Polar region (including the Tibetan Plateau, Iranian Plateau, and Tianshan Mountains) (Figure 5b). This pattern shows the effects of latitude and altitude on precipitation $\delta^{18}\text{O}$. However, precipitation $\delta^{18}\text{O}$ in Central Asia was higher than that in the Tibetan Plateau (Figure 5b). This was mainly due to the transport of moist air originating from the Arabian Sea and the Bay of

Bengal into Central Asia, which resulted in more summer precipitation (Zhao *et al.*, 2014). A strong anti-cyclonic pattern was noted in the Arabian Sea, Indian subcontinent, and the Bay of Bengal, and was associated with a strong southerly flow controlling the Indian subcontinent and extending north up to the valley between the Iranian Plateau and Tibetan Plateau regions. Furthermore, the valley between the Iranian Plateau and the Tibetan Plateau is below 1,500 m, and the anomalous southerlies can transport moisture into Central Asia (Zhao *et al.*, 2014; Zhao and Zhang, 2016). The cyclonic pattern observed in Central Asia was associated with this pattern, and an anomalous southwesterly wind may have transported moisture into Central Asia. Tian *et al.* (2001) indicated that there was a different moisture source between the northern and southern Tibetan Plateau, and that the northern limit of the summer monsoon was north of the Yarlung Zangbo River located in the middle of the Tibetan Plateau.

In the winter, precipitation $\delta^{18}\text{O}$ gradually decreased with increasing longitude (Figure 5c). The $\delta^{18}\text{O}$ values were higher in Central Asia. The moisture transported from the Eurasian continent and the Mediterranean Sea into northern Central Asia generated more winter precipitation and westerly wind was needed to transport moist air into inland regions.

Correlation between precipitation $\delta^{18}\text{O}$ and meteorological variables

We investigated correlations between monthly precipitation $\delta^{18}\text{O}$, monthly air temperature (Figure 6), and precipitation at eight stations in Central Asia (Figure 7). As shown in the figure, monthly precipitation $\delta^{18}\text{O}$ at each station had a significant positive correlation with temperature (CC = 0.51–0.96, $p < 0.01$). Moreover, the latitude effect with $\delta^{18}\text{O}$ gradually increased as the temperature increased. The effect of temperature was the strongest at the Barabinsk station, followed by those at the Urumqi, Hotan, Zhangye, Saratov, and Kabul stations, while it was the weakest at the Astrakhan and Teheran stations. The gradients between $\delta^{18}\text{O}$ and air temperature ranged from 0.28‰/°C at the Saratov and Teheran stations to 0.68‰/°C at the Hetian station.

These results were similar to global mid- and high-latitudes with gradients of 0.55‰/°C (Rozanski et al., 1993) and elsewhere in China with gradients of 0.36‰/°C (Gu, 2011).

Significant positive correlations between precipitation $\delta^{18}\text{O}$ and precipitation were observed at four stations (Zhangye, Barabinsk, Hetian, and Urumqi), and only one station had a significant negative correlation (Teheran station). Correlations for other stations were not significant. These results reveal an indeterminant dependence of precipitation $\delta^{18}\text{O}$ on precipitation, which is in accordance with the insignificant effect of precipitation in inland regions (Gu, 2012). The $\delta^{18}\text{O}$ –precipitation gradients ranged from -0.08‰/mm (Teheran station) to 0.20 ‰/mm (Zhangye station). In addition, the Zhangye and Teheran stations had a significant precipitation effect as a result of their location on the edge of the East Asian monsoon and Indian monsoon regions, respectively.

The Urumqi station has the longest and most systematic GNIP observations in Central Asia, with an observation period spanning from 1986 to 2003. Based on a subset of selected predictors, we established a model using stepwise regression analysis and the corresponding monthly temperature and precipitation data. The $\delta^{18}\text{O}$ model in Urumqi station was expressed by linear regression as:

$$\delta^{18}\text{O} = (-0.009) \times P + 0.417 \times T - 15.53 \quad (R^2 = 0.69, p < 0.01) \quad (4)$$

We reconstructed and estimated the monthly precipitation $\delta^{18}\text{O}$ time series from 1985 to 2013 at Urumqi station (Figure 8a). The reconstructed precipitation $\delta^{18}\text{O}$ was high-positively correlated with the observed values at Urumqi stations ($R = 0.83$, $p < 0.01$), and the results clearly depicted the seasonal cycle of precipitation $\delta^{18}\text{O}$, except for some values during extreme cold or hot months. Furthermore, the reconstructed precipitation $\delta^{18}\text{O}$ values for most spring and autumn seasons are similar to the observations. This indicates that basic fractionation mechanisms can be reflected using the reconstructed model. The reconstructed model indicates that the temperature

effect of precipitation $\delta^{18}\text{O}$ values was reflected in Urumqi Station. Several previous studies confirmed the temperature effect of precipitation isotopes existed in arid Central Asia (Yao et al., 1999, 2013; Wang et al., 2016). Thus, the selected control factors and established regressions can be used to reconstruct the long-term variation in precipitation $\delta^{18}\text{O}$, and also act as a proxy of an historic environment. In High Asia, the climatic controls on precipitation isotopes was used as a base for ice core sampling (Tian and Yao, 2016). In Central Asia, the $\delta^{18}\text{O}$ of ice cores were in good agreement with temperatures at nearby stations, which indicates its value in paleoclimate reconstruction using ice cores (Tian et al., 2006; Zhang and Wang, 2018).

Relationship of precipitation $\delta^{18}\text{O}$ with general atmospheric circulation

To reveal the relationship between precipitation $\delta^{18}\text{O}$ and large-scale atmospheric circulation, we analyzed the correlation between precipitation $\delta^{18}\text{O}$ at each station using long-term observation data and atmospheric circulation (e.g., IMI, SASMI, and WCI).

Our results showed that the precipitation $\delta^{18}\text{O}$ at the Zhangye station was positively correlated with the EASMI ($R = 0.62$, $p < 0.05$), indicating that the East Asian summer monsoon has an important effect on precipitation $\delta^{18}\text{O}$ in Zhangye. It is located on the eastern side of the arid northwest China and at the northeastern edge of the Tibetan Plateau as well as at the transition zone between the East Asian monsoon and westerlies. The East Asian monsoon can affect the precipitation $\delta^{18}\text{O}$ in the westerly and monsoon transition regions.

The precipitation $\delta^{18}\text{O}$ was negatively correlated with the IMI at both the Urumqi and Teheran stations ($R = 0.57$ and 0.41 , respectively, $p < 0.05$), revealing an important effect of the Indian monsoon on precipitation $\delta^{18}\text{O}$. In general, the stronger the monsoon, the lower the $\delta^{18}\text{O}$ value, and vice versa. The Teheran station is close to the Indian Ocean and is one of the main moisture paths for the Indian monsoon moving towards the north. Therefore, precipitation $\delta^{18}\text{O}$

at the Teheran station was mainly influenced by the Indian monsoon. However, Urumqi is located in the Asian hinterland and is indirectly influenced by the Indian monsoon. At Urumqi, precipitation $\delta^{18}\text{O}$ is affected by an anomalous moisture transport path through a multi-step process. A weakened Indian monsoon can cause an anomalous cyclone in the middle and upper troposphere in Central Asia, resulting in cooling, which is directly correlated with the increased precipitation in Xinjiang (Zhao *et al.*, 2014). Furthermore, the Indian monsoon affects moisture transport from the Bay of Bengal and the Arabian Sea to Xinjiang through a two-step process (Zhao *et al.*, 2014). The extreme summer precipitation in northern Xinjiang was controlled by moisture sources originating from the Indian Ocean, which was closely related to a stronger meridional circulation (Huang *et al.* 2017). The anomalous circulation can also transport some moisture from the Indian Ocean along the eastern periphery of the Tibetan Plateau to North Xinjiang (Huang *et al.*, 2017).

The moisture in Central Asia is mainly transported by the westerly circulation from the North Atlantic Ocean and the Eurasian continent. The westerly circulation is one of the most important factors affecting precipitation $\delta^{18}\text{O}$ over Central Asia. However, we found a weak positive correlation between precipitation $\delta^{18}\text{O}$ and the WCI index. This may be related to the moisture transported by the westerly circulation, which originates from the Atlantic and Arctic oceans and dissipates along the way. The water vapor transported by the westerly is exhausted by the time it reaches Central Asia because of substantial evapotranspiration and evaporation below the clouds in arid regions.

The precipitation d-excess parameter can provide supplementary information to precipitation $\delta^{18}\text{O}$ and $\delta^2\text{H}$ values, and it is largely controlled by the moisture source and water transport paths. Thus, it may be the most important indicator to represent the regional moisture source and atmospheric circulation (Merlivat and Jouzel, 1979). In addition, the precipitation d-

excess is closely related to the re-evaporation of raindrops during condensation and precipitation, and is affected by temperature and relative humidity during evaporation (Merlivat and Jouzel, 1979; Jouzel et al., 1982, 1997; Wang et al., 2014). In Central Asia, the precipitation d-excess values ranged from +21.66 ‰ to -8.5‰, with an average of 8.5 ‰. The temporal variations in precipitation d-excess shows more positive values in spring and more negative values in summer and the opposite pattern for $\delta^{18}\text{O}$. This variation is in agreement with variations in the conditions under which the moisture source evaporates. There is a similar seasonal pattern in precipitation d-excess and $\delta^{18}\text{O}$ of Central Asia and that of the westerly climate regime zone (Yao et al., 2009). However, there is a dissimilarity with the annual cycle in Lhasa, which is affected by southwest monsoon air mass (Wang et al., 2014). This demonstrates that the moisture transported to Central Asia was derived predominantly from the westerlies and the polar air masses. Moreover, the high precipitation d-excess values showed that recycled moisture derived from local sources makes a significant contribution to precipitation, especially in Central Asia. Yao et al (2020) also suggested that the warming and increased moisture content of the atmosphere contributed to the local moisture cycle and increased precipitation recycling in eastern Central Asia.

Conclusions

We established the CAMWL and analyzed the spatial–temporal characteristics of precipitation $\delta^{18}\text{O}$ and its relationship with meteorological variables and geographical factors based on the precipitation $\delta^{18}\text{O}$ values observed at 47 stations over Central Asia, with information from GNIP, TNIP, CHNIP, and TSNIP databases and reference studies. In addition, we revealed the relationship between precipitation $\delta^{18}\text{O}$ and large-scale atmospheric circulation. We made the following conclusions:

(1) The CAMWL was established as $\delta^2\text{H} = 7.30\delta^{18}\text{O} + 3.12$ ($R^2=95$, $p<0.01$) for 727 groups of monthly precipitation $\delta^{18}\text{O}$ over Central Asia. The precipitation $\delta^{18}\text{O}$ over Central Asia ranged from +2‰ to -25.4‰ with a mean of -8.7‰.

(2) The precipitation $\delta^{18}\text{O}$ over Central Asia was related to meteorological factors. It had a significant positive correlation with temperature, with $\delta^{18}\text{O}$ -temperature gradients ranging from 0.28‰/°C to 0.68‰/°C. However, the dependence of $\delta^{18}\text{O}$ on precipitation was unclear, with a significant effect on precipitation observed only at the Zhangye and Teheran stations, showing $\delta^{18}\text{O}$ -precipitation gradients of 0.20‰/mm and -0.08‰/mm, respectively.

(3) In summer, the latitude was significantly correlated with precipitation $\delta^{18}\text{O}$ ($CC = 0.54$, $p < 0.05$), while in winter, both latitude and longitude were significantly correlated. The gradient of $\delta^{18}\text{O}/\text{LAT}$ and $\delta^{18}\text{O}/\text{ALT}$ were -0.42‰/° and -0.001‰/m, respectively.

(4) The precipitation $\delta^{18}\text{O}$ at the Zhangye station showed a significantly positive correlation with the EASMI, and a negative correlation with the IMI at the Urumqi and Teheran stations. In addition, there was a weakly positive correlation between $\delta^{18}\text{O}$ and the WCI. Our results suggest that the moisture in Central Asia is mainly transported by westerly circulation and is indirectly affected by the Indian monsoon. Furthermore, the East Asian monsoon can affect the precipitation $\delta^{18}\text{O}$ in westerly and monsoon transition regions.

References



- Bowen G J, Wilkinson B H. 2002.** Spatial distribution of $\delta^{18}\text{O}$ in meteoric precipitation. *Geology* 30(4): 315—318 DOI [10.1130/0091-7613\(2002\)030<0315:SDOOIM>2.0.CO;2](https://doi.org/10.1130/0091-7613(2002)030<0315:SDOOIM>2.0.CO;2)
- Bowen, G. J. and Revenaugh, J. 2003.** Interpolating the isotopic composition of modern meteoric precipitation. *Water Resour. Res* 39(10): 1299 DOI [10.1029/2003WR002086](https://doi.org/10.1029/2003WR002086)
- Coplen T B. 1993.** Uses of environmental isotopes. In: Alley W B, ed. *Regional Ground Water*

Quality. New York: Van Nostrand Reinhold, 227—254

Chen Fahu, Yu Zicheng, Yang Meilin et al. 2008. Holocene moisture evolution in arid central Asia and its out- of- phase relationship with Asian monsoon history. *Quaternary Science Reviews* 27(3/4): 351-364 DOI [10.1016/j.quascirev.2007.10.017](https://doi.org/10.1016/j.quascirev.2007.10.017)

Chen Xi. Reveal and analysis of evaporation in Central area of Asia. Beijing: China Meteorological Press, 2012 (in Chinese).

Chen Yaning. Water resources research in arid regional of northwest in China. Beijing: China Science Press, 2014. (in Chinese)

Cai Ying, Song Minhong, Qian Zhenag'an. 2015. Reanalyses of Precipitation Circulation and Vapor Transportation of Severe Dry and Wet Events in Summer in Arid Region of Northwest China. *Plateau Meteorology*, 34 (3): 597-610 (in Chinese).

Craig, H., and L. I. Gordon. 1965. Deuterium and oxygen 18 variations in the ocean and the marine atmosphere, in *Stable Isotopes in Oceanographic Studies and Paleotemperatures*, edited by E. Tongiorgi, V. Lischi, Pisa, Italy, pp. 9 – 130.

Chen Zhong-xiao, Cheng Jun, Guo Pin-wen, et al. 2010. Distribution characters and its control factors of stable isotope in precipitation over China. *Trans Atmos Sci* 33 (6): 667-679 (in Chinese).

Chen, F.H., Chen, J.H., Holmes, J., Boomer, I., Austin, P., Gates, J.B., Wang, N.L., Brooks, S.J., Zhang, J.W., 2010. Moisture changes over the last millennium in arid Central Asia: a review, synthesis and comparison with monsoon region. *Quat. Sci. Rev* 29: 1055–1068 DOI [10.1016/j.quascirev.2010.01.005](https://doi.org/10.1016/j.quascirev.2010.01.005)

Huang, W., Feng, S., Liu, C., Chen, J., Chen, J.H., Chen, F.H., 2018. Changes of climate regimes during the last millennium and the twenty-first century simulated by the Community Earth System Model. *Quat. Sci. Rev.* 180: 42–56 DOI [10.1016/j.quascirev.2018.08.017](https://doi.org/10.1016/j.quascirev.2018.08.017)

[10.1016/j.quascirev.2017.11.019](https://doi.org/10.1016/j.quascirev.2017.11.019)

Dansgaard W. 1953. The abundance of ^{18}O in atmospheric water and water vapor. *Tellus* 5(4): 461-469

Dansgaard W. 1964. Stable isotopes in precipitation. *Tellus* 16(4): 436—468 [DOI](https://doi.org/10.1111/j.2153-3490.1964.tb00181.x)

[10.1111/j.2153-3490.1964.tb00181.x](https://doi.org/10.1111/j.2153-3490.1964.tb00181.x)

Dai Xingang, Li Weijing, Ma Zhuguo. 2006. The variation characteristics of water vapor source in Xinjiang in recent decades. *Progress in Natural Science* 16(12): 1651-1656 (in Chinese).

Feng F, Li ZQ, Zhang MJ, et al., 2013. Deuterium and oxygen 18 in precipitation and atmospheric moisture in the upper Urumqi River Basin, eastern Tianshan Mountains. *Environmental Earth Sciences*, 68(4): 1199–1209 [DOI 10.1007/s12665-012-1820-y](https://doi.org/10.1007/s12665-012-1820-y).

Harvey F E, Welker J M. 2000. Stable isotopic composition of precipitation in the semi-arid north-central portion of the US Great Plains. *Journal of Hydrology* 238: 90-109 [DOI 10.1016/S0022-1694\(00\)00316-4](https://doi.org/10.1016/S0022-1694(00)00316-4)

He Qing, Yuan Yujiang, Zhao Yong, et al. 2016. Investigation on climate change in Central Asia. Beijing: China Meteorological Press (in Chinese).

Huang W, Feng S, Chen J, et al. 2015. Physical Mechanisms of Summer Precipitation Variations in the Tarim Basin in Northwestern China. *Journal of Climate* 28(9): 3579-3591 [DOI 10.1175/JCLI-D-14-00395.1](https://doi.org/10.1175/JCLI-D-14-00395.1)

Huang Wei, Shiqiao, C., Chengling, X., & Zhiping, Z. 2017. Moisture sources of extreme summer precipitation events in north xinjiang and their relationship with atmospheric circulation. *Advances in Climate Change Research* 8(1): 12-17 [DOI 10.1016/j.accre.2017.02.001](https://doi.org/10.1016/j.accre.2017.02.001)

Gu Weizu. 2012. Isotope hydrology. Beijing: China Science Press (in Chinese).

- 418 **Gao, J., V. Masson-Delmotte, T. Yao, L. Tian, C. Risi, and G. Hoffmann. 2011.** Precipitation
419 water stable isotopes in the south Tibetan Plateau: Observations and modeling, *J. Clim.*, 24,
420 3161 – 3178 DOI [10.1175/2010JCLI3736.1](https://doi.org/10.1175/2010JCLI3736.1).
- 421 **Jones MD, Leng MJ, Arrowsmith C. 2007.** Local δD and $\delta^{18}O$ variability in UK rainfall.
422 *Hydrol Earth Syst Sci* 4:2403–2423 DOI [10.5194/hessd-4-2403-2007](https://doi.org/10.5194/hessd-4-2403-2007)
- 423 **Jouzel, J., Merlivat, L., Lorius, C., 1982.** Deuterium excess in an east Antarctic ice core
424 suggests higher relative humidity at the oceanic surface during the last glacial maximum.
425 *Nature* 299: 688-691 DOI [10.1038/299688a0](https://doi.org/10.1038/299688a0)
- 426 **Jouzel, J., Froehlich, K., Schotterer, U., 1997.** Deuterium and oxygen-18 in present-day
427 precipitation: data and modeling. *Hydrological Science* 42: 747-763 DOI
428 [10.1080/02626669709492070](https://doi.org/10.1080/02626669709492070)
- 429 **Kong, Y., Pang, Z. and Froehlich, K. 2013.** Quantifying recycled moisture fraction in
430 precipitation of an arid region using deuterium excess. *Tellus* 65B, 19251. DOI
431 [10.3402/tellusb.v65i0.19251](https://doi.org/10.3402/tellusb.v65i0.19251)
- 432 **Li Xiaofei, Zhang Mingjun, Li Yaju, et al. 2012.** Characteristics of $\delta^{18}O$ in Precipitation and
433 Moisture Transports over the Arid Region in Northwest China. *Chinese Journal of*
434 *Environmental Science* 33(3): 711-719 (in Chinese).
- 435 **Liu Xiaokang, Rao Zhiguo, Zhang Xiaojian, et al. 2015.** Variations in the oxygen isotopic
436 composition of precipitation in the Tianshan Mountains region and their significance for the
437 Westerly circulation. *Acta Geographica Sinica* 70(1): 97-109 (in Chinese).
- 438 **Liu Jianrong, Song Xianfang, Yuan Guofu, Sun Xiaomin & Yang Lihu. 2014.** Stable
439 isotopic compositions of precipitation in China, *Tellus B: Chemical and Physical*
440 *Meteorology*, 66:1, 22567 DOI: [10.3402/tellusb.v66.22567](https://doi.org/10.3402/tellusb.v66.22567)
- 441 **Li Hui, Zhou Hongfei. 2007.** Variation characteristics of δD and $\delta^{18}O$ stable isotopes in the

- precipitation of Urumqi. *Journal of Arid Land Resources and Environment* 21(9): 46-50 (in Chinese).
- Li, J. P., and Q. C. Zeng. 2003.** A new monsoon index and the geographical distribution of the global monsoons. *Adv. Atmos. Sci.* 20: 299-302
- Li, J. P., and Q. C. Zeng. 2002.** A unified monsoon index. *Geophysical Research Letters* 29(8): 1274 DOI:10.1029/2001GL013874.
- Liu Zhongfang, Tian Lide, Yao Tandong. 2009.** Spatial distribution of $\delta^{18}\text{O}$ in precipitation over China. *Chinese Science Bulletin* 54(6): 804-811 (in Chinese).
- Liu Jianfang, Song Xianfang, Yuanguofu, et al. 2008.** Characteristics of ^{18}O in Precipitation over Northwest China and Its Water Vapor Sources. *Acta Geographica Sinica* 63(1): 12-22 (in Chinese).
- Liu JR, Song XF, Yuan GF, et al., 2014a.** Stable isotopic compositions of precipitation in China. *Tellus B: Chemical and Physical Meteorology*, 66(1): 22567 DOI 10.3402/tellusb.v66. 22567
- Lutz A, Thomas JM, Panorska A. 2011.** Environmental controls on stable isotope precipitation values over Mali and Niger, West Africa. *Environ Earth Sci* 62:1749–1759 DOI 10.1007/s12665-010.0655-7
- Merlivat, L., Jouzel, J., 1979.** Global climatic interpretation of the deuterium-oxygen 18 relationship for precipitation. *Journal of Geophysical Research* 84: 5029-5033 DOI 10.1029/JC084iC08p05029
- Rozanski K, Araguas-Araguas L, Gonfiantini R. 1993.** Isotopic patterns in modern global precipitation. In: Swart P, Lohmann K C, McKenzie J A, et al, eds. *Climate Change in Continental Isotopic Records*. Washington D C: American Geophysical Union, 1—36.
- Shi Yuguang, Sun Zhaobo. 2008.** Climate characteristics of water vapor transportation and its

variation over Xinjiang. Plateau Meteorology 27(2): 310-319 (in Chinese).

Song Xianfang, Liu Jianrong, Sun Xiaomin et al. 2007. Establishment of Chinese Network of Isotopes in Precipitation (CHNIP) based on CERN. Advances in Earth Science 22(7): 738-747 (in Chinese).

Sophocleous M. 2002. Interactions between groundwater and surface water: The state of the science. Hydrogeology Journal 10(1): 52-67.

Tian LD, Yao TD, Li Z, et al., 2006. Recent rapid warming trend revealed from the isotopic record in Muztagata ice core, eastern Pamirs. Journal of Geophysical Research: Atmospheres, 111(D13): D13103. DOI 10.1029/2005JD006249.

Tian LD, Yao TD, 2016. High-resolution climatic and environmental records from the Tibetan Plateau ice cores. Chinese Science Bulletin 61(9): 926–937 DOI 10.1360/N972015-00779 (in Chinese)

Kong, Y., Pang, Z., & Froehlich, K. 2013. Quantifying recycled moisture fraction in precipitation of an arid region using deuterium excess. Tellus B: Chemical and Physical Meteorology 65(1): 19251 DOI 10.3402/tellusb.v65i0.19251

Kong, Y., & Pang, Z. 2016. A positive altitude gradient of isotopes in the precipitation over the Tianshan Mountains: Effects of moisture recycling and sub-cloud evaporation. Journal of Hydrology 542: 222-230 DOI 10.1016/j.jhydrol.2016.09.007

Sun CJ, Li WH, Chen YN, et al., 2015. Isotopic and hydrochemical composition of runoff in the Urumqi River, Tianshan Mountains, China. Environmental Earth Sciences 74(2): 1521–1537 DOI 10.1007/s12665-015-4144-x.

Tian, L., T. Yao, Z. Yang, and J. Pu. 1997. A 4-year' s observation of $\delta^{18}\text{O}$ in precipitation on the Tibetan Plateau, J. Glacio and Geocryol 19: 32–36.

Tian, L., V. Masson-Delmotte, M. Stievenard, T. Yao, and J. Jouzel. 2001. Tibetan Plateau

summer monsoon northward extent revealed by measurements of water stable isotopes, J. Geophys. Res. 106(D22): 28,081–28,088, DOI 10.1029/2001 JD900186.

Wang Keli, Jiang Hao, Zhao Hongyan. 2005. Atmospheric water vapour transport from westerly and monsoon over the Northwest China. *Advances in Water Science* 16(3): 432-438 (in Chinese).

Wang, S., Zhang, M., Hughes, C. E., Zhu, X., Dong, L., Ren, Z., & Chen, F. 2016. Factors controlling stable isotope composition of precipitation in arid conditions: An observation network in the Tianshan Mountains, central Asia. *Tellus B: Chemical and Physical Meteorology* 68(1): 26206 DOI 10.3402/tellusb. v68.26206

Wang, S., Zhang, M., Crawford, J., Hughes, C. E., Du, M., & Liu, X. 2017. The effect of moisture source and synoptic conditions on precipitation isotopes in arid central Asia. *Journal of Geophysical Research: Atmospheres* 122(5): 2667-2682 DOI 10.1002/2015JD024626

Wang SJ, Zhang MJ, Che YJ, et al., 2016. Influence of below-cloud evaporation on deuterium excess in precipitation of arid Central Asia and its meteorological controls. *Journal of Hydrometeorology* 17(7): 1973–1984 DOI 10.1175/JHM-D-15-0203.1

Wang, X., Li, Z., Tayier, R., & Wang, S. 2015. Characteristics of atmospheric precipitation isotopes and isotopic evidence for the moisture origin in Yushugou River basin, Eastern Tianshan Mountains, China. *Quaternary international* 380: 106-115 DOI 10.1016/j.quaint.2014.12.023

Welker J M, White J C, Vachon R, et al. 2002. High resolution spatial patterns of the isotopic characteristics of precipitation across the U.S. during El Nino and La Nina phases. *EOS Trans* 83: 47

Yamanaka T, Tsujimura M, Oyunbaatar D et al. 2007. Isotopic variation of precipitation

- over eastern Mongolia and its implication for the atmospheric water cycle. Journal of Hydrology 333: 21-34 DOI 10.1016/j.jhydrol.2006.07.022
- Yao, T., X. Guo, L. Thompson, K. Duan, N. Wang, J. Pu, B. Xu, X. Yang, and W. Sun. 2006b.** $\delta^{18}\text{O}$ record and temperature change over the past 100 years in ice cores on the Tibetan Plateau, Sci. China Ser. D 49(1): 1–9 DOI:10.1007/s11430-004-5096-2.
- Yao, T., et al. 2013.** A review of climatic controls on $\delta^{18}\text{O}$ in precipitation over the Tibetan Plateau: Observations and simulations, Rev. Geophys. 51: 525 – 548 DOI:10.1002/rog.20023.
- Yao Tandong. 2009.** Spatial distribution of $\delta^{18}\text{O}$ in precipitation over China. Chinese Science Bulletin 54(15): 2123-2123 (in Chinese).
- Yao TD, Masson V, Jouzel J, et al., 1999.** Relationships between $\delta^{18}\text{O}$ in precipitation and surface air temperature in the Urumqi River Basin, East Tianshan Mountains, China. Geophysical Research Letters 26(23): 3473–3476. DOI: 10.1029/1999GL006061.
- Yao Junqiang, Liu Zhihui, Yang Qing, et al. 2014.** Temperature variability and its possible causes in the typical basins of the arid Central Asia in recent 130 years. Acta Geographica Sinica 69(3):291- 302 (in Chinese).
- Yang Meixue, Yao Tandong, Tian Lide., et al. 2004.** Analysis of Precipitation From Different Water Vapor Sources in Tibetan Plateau. Scientia Geographica Sinica 24(4): 426-431 (in Chinese).
- Yurtsever Y, Gat J R. 1981.** Atmospheric waters. In: Gat J R, Gonfiantini R, eds. Stable Isotope Hydrology: Deuterium and Oxygen-18 in the Water Cycle. Vienna: International Atomic Energy Association 103—139
- Yao Junqiang, Yang Qing, Han Xueyun, et al. 2016.** Quantifying recycled moisture fraction in precipitation of Tianshan Mountain, Central Asia. Desert and oases meteorology 10 (6) (in

Chinese).

Yao, J., Chen, Y., Zhao, Y., Guan, X., Mao, W., & Yang, L. 2020. Climatic and associated

atmospheric water cycle changes over the Xinjiang, China. *Journal of Hydrology*, 124823

[DOI org/10.1016/j.jhydrol.2020.124823](https://doi.org/10.1016/j.jhydrol.2020.124823)

Zhao Y., A N Huang, Y Zhou, et al. 2014. Impact of the middle and upper tropospheric cooling

over Central Asia on the summer rainfall in the Tarim Basin. *J. Climate* 27: 4721-4732 [DOI](https://doi.org/10.1175/JCLI-D-13-00456.1)

[10.1175/JCLI-D-13-00456.1](https://doi.org/10.1175/JCLI-D-13-00456.1)

Zhao, Y., & Zhang, H. 2015. Impacts of sst warming in tropical indian ocean on cmip5 model-

projected summer rainfall changes over central asia. *Climate Dynamics* 46(9-10): 1-16 [DOI](https://doi.org/10.1007/s00382-015-2765-0)

[10.1007/s00382-015-2765-0](https://doi.org/10.1007/s00382-015-2765-0)

Zhang Xinping, Liu Jingmiao, Tian Lide et al. 2004. Variations of (18O in precipitation along

vapour transport paths over Asia. *Acta Geographica Sinica* 59(5): 699-708 (in Chinese).

Zhang, M., & Wang, S. 2018. Precipitation isotopes in the Tianshan Mountains as a key to

water cycle in arid central Asia. *Sciences in Cold and Arid Regions* 10(1): 27-37 [DOI](https://doi.org/10.3724/SP.J.1226.2018.00027)

[10.3724/SP.J.1226.2018.00027](https://doi.org/10.3724/SP.J.1226.2018.00027)

Zhang MJ, Wang SJ, 2016. A review of precipitation isotope studies in China: basic pattern

and hydrological process. *Journal of Geographical Sciences* 26(7): 921–938 [DOI](https://doi.org/10.1007/s11442-016-1307-y)

[10.1007/s11442-016-1307-y](https://doi.org/10.1007/s11442-016-1307-y)

Table 1(on next page)

Summary data of the precipitation sampling stations over Central Asia

GNIP = Global Network of Isotopes in Precipitation; TNIP = Tibetan Network for Isotopes in Precipitation; CHNIP = Chinese Network of Isotopes in Precipitation; TSNIP = Tianshan Network of Isotopes in Precipitation; and TGS = Tianshan Glaciological Station, Chinese Academy of Sciences.

1

Table 1. Summary data of the precipitation sampling stations over Central Asia

No	Station	Latitude (°N)	Longitude (°E)	Altitude (m)	Precipitation (mm)	Temperature (°C)	Database
1	Urumqi	43.78	87.62	918	304	7.4	GNIP
2	Hetian	37.13	79.93	1375	209	9.1	GNIP
3	Zhangye	38.93	100.43	1483	154	7.8	GNIP
4	Kabul	34.57	69.21	1860	330	11.6	GNIP
5	Barabinsk	55.33	78.37	120	389	1	GNIP
6	Astrakhan	46.25	48.03	-18	228	10.5	GNIP
7	Saratov	51.56	46.03	166	490	6.6	GNIP
8	Teheran	35.68	51.32	1200	227	16.7	GNIP
9	Tashkent	41.27	69.27	428	478	13.9	GNIP
10	Teheran East	35.74	51.58	1350	297	15.4	GNIP
11	Novosibirsk	55.03	82.9	162	479	2.5	GNIP
12	Telavi	41.93	45.48	590	679	13.1	GNIP
13	Taxkorgen	37.77	75.27	3100	115	1.6	TNIP
14	Delingha	37.37	97.37	2981	186	2.2	TNIP
15	Fukang	44.29	87.93	460	167	7.5	CHNIP
16	Cele	37.02	80.73	1306	51	12.9	CHNIP
17	Yining	43.95	81.33	662.5	298.9	9.5	TSNIP
18	Jinghe	44.62	82.9	320.1	112.1	8.2	TSNIP
19	Kuytun	44.4	84.87	562	183.5	8.5	TSNIP
20	Shihezi	44.32	86.05	442.9	226.9	7.8	TSNIP
21	Caijiahu	44.2	87.53	440.5	153.8	6.5	TSNIP
22	Qitai	44.02	89.57	793.5	200.9	5.4	TSNIP
23	Wuqia	39.72	75.25	2175.7	188.7	7.7	TSNIP
24	Akqi	40.93	78.45	1984.9	237.7	6.8	TSNIP
25	Bayinbuluke	43.03	84.15	2458	280.5	-4.2	TSNIP
26	Baluntai	42.73	86.3	1739	220.4	7	TSNIP
27	Balikun	43.57	93.05	1677.2	230.5	2.7	TSNIP
28	Yiwu	43.27	94.7	1728.6	104.4	4.2	TSNIP
29	Aksu	41.17	80.23	1103.8	80.4	10.8	TSNIP
30	Baicheng	41.78	81.9	1229.2	136.6	8.2	TSNIP
31	Kuqa	41.72	82.97	1081.9	76.7	11.3	TSNIP
32	Luntai	41.78	84.25	976.1	78.6	11.6	TSNIP
33	Korla	41.75	86.13	931.5	59.2	12	TSNIP
34	Kumux	42.23	88.22	922.4	59.9	9.8	TSNIP
35	Dabancheng	43.35	88.32	1103.5	76.7	6.9	TSNIP
36	Turpan	42.93	89.2	34.5	15.4	15.1	TSNIP
37	Shisanjianfang	43.22	91.73	721.4	22.6	12.5	TSNIP

38	Hami	42.82	93.52	737.2	43.7	10.3	TSNIP
39	UG1	43.1	86.84	3693	460	-5.6	TGS
40	Zongkong	43.11	86.89	3404	400	-4.6	TGS
41	Daxigou	43.11	86.86	3539	458	-5	TGS
42	Altay	47.73	88.08	735	193	4.5	Tian et al., 2001
43	Gaoshan	43.1	86.83	3545	390	-4.3	Kong et al., 2013
44	Houxia	43.28	87.8	2100	424	1.5	Kong et al., 2013
45	Yuejinqiao	43.12	87.05	2526	330	-2.5	Yao et al., 1999
46	Yingxiongqiao	44.37	87.2	1920	210	2.3	Sun et al., 2015
47	Yushugou	43.08	93.95	1670	94.3	18.6	Wang et al., 2015

2 **Note:** GNIP = Global Network of Isotopes in Precipitation; TNIP = Tibetan Network for Isotopes in Precipitation; CHNIP =
3 Chinese Network of Isotopes in Precipitation; TSNIP = Tianshan Network of Isotopes in Precipitation; and TGS = Tianshan
4 Glaciological Station, Chinese Academy of Sciences.

5

Figure 1

Locations of the sampling stations over Central Asia




Figure 2

Linear δD - $\delta^{18}O$ relationships (CAMWL) based on all precipitation measurements over Central Asia




Figure 3

Seasonal patterns of observed precipitation $\delta^{18}\text{O}$, precipitation amount (P), and temperature (T) at different stations. For each station, data were averaged over the observation periods.




Figure 4

Spatial patterns of annual (top), JJA (middle), and DJF (bottom) observed precipitation $\delta^{18}\text{O}$ at meteorological stations over Central Asia.




Figure 5

Spatial distribution of annual (a), JJA (b), and DJF (c) mean estimated $\delta^{18}\text{O}$ in precipitation based on the model using spatial variables






Figure 6

Regression of observed precipitation $\delta^{18}\text{O}$ with temperature over Central Asia











Figure 7

Regression of observed precipitation $\delta^{18}\text{O}$ with amount of precipitation over Central Asia.











Figure 8

Reconstruction of monthly $\delta^{18}\text{O}$ time series during 1986–2013 based on the regression model established for Urumqi station.

