

# New insight into the divergent responses of plants to warming in the context of root endophytic bacterial and fungal communities

Xiaoting Wei<sup>1</sup>, Fengyan Jiang<sup>1</sup>, Bing Han<sup>1</sup>, Hui Zhang<sup>1</sup>, Ding Huang<sup>Corresp., 1</sup>, Xinqing Shao<sup>1, 2, 3</sup>

<sup>1</sup> College of Grassland Science and Technology, China Agricultural University, Beijing, China

<sup>2</sup> Key Laboratory of Restoration Ecology of Cold Area in Qinghai province, Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xining, China

<sup>3</sup> Qinghai Provincial Key Laboratory of Adaptive Management on Alpine Grassland, Xining, China

Corresponding Author: Ding Huang

Email address: huangding@cau.edu.cn

Plant adaptation under climate changes is critical to the maintenance of terrestrial ecosystem structure and function. Studying the response of the endophytic community to climate warming is a novel way to reveal the mechanism of host environmental adaptability because of the prominent role endophytes play in host nutrient acquisition and stress tolerance. However, host performance was generally neglected in previous relevant research, which limits our understanding of the relationships between the endophytic community and host responses to climate warming. The present study selected two plants with different responses to climate warming. *Elymus nutans* is more suitable for growing in warm environments at low altitude compared to *Kobresia pygmaea*. *K. pygmaea* and *E. nutans* were sampled along an altitude gradient in the natural grassland of Qinghai-Tibet Plateau, China. Root endophytic bacterial and fungal communities were analyzed using high throughput sequencing. The results revealed that hosts growing in more suitable habitats held higher endophytic fungal diversity. Elevation and host identity significantly affected the composition of the root endophytic bacterial and fungal community. 16S rRNA functional prediction demonstrated that hosts that adapted to lower temperatures recruited endophytic communities with higher abundance of genes related to cold resistance. Hosts that were more suitable for warmer and drier environments recruited endophytes with higher abundance of genes associated with nutrient absorption and oxidation resistance. We associated changes in the endophytic community with hosts adaptability to climate warming and suggested a synchronism of endophytic communities and hosts in environmental adaptation.

**New insight into the divergent responses of plants to warming in the context of root  
endophytic bacterial and fungal communities**

Xiaoting Wei <sup>1</sup>, Fengyan Jiang <sup>1</sup>, Bing Han <sup>1</sup>, Hui Zhang <sup>1</sup>, Ding Huang <sup>1\*</sup> and Xinqing Shao <sup>1,2,3</sup>

<sup>1</sup> College of Grassland Science and Technology, China Agricultural University, Beijing, China

<sup>2</sup> Key Laboratory of Restoration Ecology of Cold Area in Qinghai province, Northwest Institute  
of Plateau Biology, Chinese Academy of Sciences, Xining, Qinghai Province, China

<sup>3</sup> Qinghai Provincial Key Laboratory of Adaptive Management on Alpine Grassland, Xining,  
Qinghai Province, China

**Corresponding author:**

Ding Huang

Email address: [huangding@cau.edu.cn](mailto:huangding@cau.edu.cn)

# Abstract

Plant adaptation under climate changes is critical to the maintenance of terrestrial ecosystem structure and function. Studying the response of the endophytic community to climate warming is a novel way to reveal the mechanism of host environmental adaptability because of the prominent role endophytes play in host nutrient acquisition and stress tolerance. However, host performance was generally neglected in previous relevant research, which limits our understanding of the relationships between the endophytic community and host responses to climate warming. The present study selected two plants with different responses to climate warming. *Elymus nutans* is more suitable for growing in warm environments at low altitude compared to *Kobresia pygmaea*. *K. pygmaea* and *E. nutans* were sampled along an altitude gradient in the natural grassland of Qinghai-Tibet Plateau, China. Root endophytic bacterial and fungal communities were analyzed using high throughput sequencing. The results revealed that hosts growing in more suitable habitats held higher endophytic fungal diversity. Elevation and host identity significantly affected the composition of the root endophytic bacterial and fungal community. 16S rRNA functional prediction demonstrated that hosts that adapted to lower temperatures recruited endophytic communities with higher abundance of genes related to cold resistance. Hosts that were more suitable for warmer and drier environments recruited endophytes with higher abundance of genes associated with nutrient absorption and oxidation resistance. We associated changes in the endophytic community with hosts adaptability to climate warming and suggested a synchronism of endophytic communities and hosts in environmental adaptation.

43

## 44 **Introduction**

45 Plant endophytes are microorganisms that colonize the intercellular space of plants and  
 46 establish a harmonious association with their host plants (*Odelade and Babalola, 2019*). The  
 47 genome of the host and its microbiome are considered a unit according to the hologenome theory  
 48 (*Zilber-Rosenberg and Rosenberg, 2008*). Studies on plant-associated microbiota recently  
 49 exhibited phenomenal expansion in the areas of natural and agricultural ecosystems due to their  
 50 great potential in improving host fitness, including nutrient acquisition (*van der Heijden, 2016*),  
 51 stress tolerance (*Liu et al., 2020*) and disease defense (*Dini-Andreote, 2020; Carrión, 2019*).  
 52 Endophytic community diversity was closely associated with multiple ecosystem functions  
 53 (*Laforest-Lapointe et al., 2017*).

54 Recently, great attention has been paid to the environmental problems caused by climate  
 55 warming in terrestrial ecosystems, such as the loss of biodiversity, changes in vegetation  
 56 composition and community succession. Climate changes also affect plant ranges and allow  
 57 some species to expand or narrow their ranges (*Morin, Viner & Chuine, 2008*). The formation of  
 58 new interspecific relationships between plant species under climate change may drive these  
 59 ecological processes (*Little et al., 2016*), and endophytes may modulate the interspecific  
 60 interactions and vegetation composition (*Porras-Alfaro, 2011; Jackrel, 2020*).

61 Plant endophytic communities are dynamic and variable among seasons (*Li, Yang & Zhao,*  
 62 *2005*) and environments (*Laforest-Lapointe, 2016*) and reflect the adaptability of plants to  
 63 changing environments. The recent “cry for help” hypothesis suggests that plants selectively

recruit beneficial microbes from their surroundings under biotic or abiotic stress (Neal *et al.* 2012; Liu *et al.* 2019). *Pseudomonas* colonization in the roots of the desert plant *Alhagi sparsifolia* increased under drought conditions and promoted drought resistance (Zhang *et al.*, 2020). Different plant species show unique abilities in the process of recruitment. For example, most plants form symbioses with arbuscular mycorrhizal fungi (AMF) to increase phosphorus availability. *Arabidopsis thaliana*, which lose the ability of symbiosing with AMF, recruit Helotiales and *Colletotrichum tofieldiae* under phosphorus-limited conditions (Almario *et al.*, 2017; Hiruma *et al.*, 2016).

However, studies on the impacts of climate change on plant endophyte communities are insufficient and many findings are conflicting. Endophytes and their hosts are “communities of interest”. Previous studies focused only on endophytic communities and host responses to climate change were generally neglected (Qian *et al.*, 2018; Cordier *et al.*, 2012), which limits our understanding of the ecological significance of endophytic community changes. Host selection is critical because plant species (Yao *et al.*, 2019; Laforest-Lapointe, Messier & Kembel, 2016) and compartments (leaf, stem, root) of the same plant greatly affect endophytic community composition (Qian *et al.*, 2019). Fujimura also concluded that the conflicting results between studies on the effects of warming on endophytic communities may due to differences in host responses (Fujimura, Egger & Henry, 2008). Therefore, plant species that are sensitive to climate changes should be selected to examine the responses of endophyte communities and improve our understanding of plant-microbe interactions and elucidate the mechanisms of plant adaptation to climate change from the perspective of endophytes.

The Qinghai-Tibet Plateau is the highest plateau in the world, and it is experiencing rapid climate warming at rate more than twice the global level (Ma *et al.*, 2017), which threatens biodiversity and ecosystem function in this area. *Kobresia pygmaea* and *Elymus nutans* are two common plants in the Qinghai-Tibet Plateau that respond differently to climate warming. *Kobresia pygmaea* is a carpet-like sedge species that is highly adaptable to high-altitude environments, and it primarily grows on the alpine meadow of the Qinghai-Tibet Plateau between altitudes of 3 000 m and 6 000 m (Wu *et al.*, 2017; Miehe *et al.*, 2008). The *K. pygmaea* alpine meadow, which is dominated by *K. pygmaea*, covers 17% of the total area of the Qinghai-Tibet Plateau (Miehe *et al.*, 2008), which makes it an important species in maintaining the stability of the local ecosystem. *Elymus nutans* is a dominant and constructive grass of meadow steppe, and it is characterized by strong drought, cold and pest resistance. It is widely distributed and survives at elevations ranging from 450 m to 4 500 m. *E. nutans* is better suited to low altitudes and warmer habitats than *K. pygmaea*, and an increase in temperature from high to low elevations promotes the growth of *E. nutans* (Qi *et al.*, 2020). *K. pygmaea* primarily grows at high altitudes in humid and cold habitats (Wu *et al.*, 2017). Warming increased the proportion of *Elymus nutans* and decreased the proportion of *K. pygmaea* in alpine grassland ecosystems (Liu *et al.*, 2018; Niu *et al.*, 2019).

The present study examined the adaptability of plants to climate warming by analyzing changes in endophytic community diversity and composition. We used elevation gradients where climatic conditions, vegetation and soil characteristics show regular changes over short geographic distances to investigate the responses of terrestrial ecosystems to climate warming

(Korner, 2007). *K. pygmaea* and *E. nutans* were sampled along an elevation gradient in the Qinghai-Tibet Plateau, and the root endophytic bacterial and fungal communities were analyzed based on the high-throughput sequencing. The innovation of this study lies in the selection of the two hosts with opposing responses to climate warming to elucidate how endophytic communities change during the promotion or inhibition of host growth by climate warming. We addressed the following questions: 1) how endophytic community diversity in the roots of *K. pygmaea* and *E. nutans* changed with climate warming (decreasing elevation); 2) whether climate warming affected endophytic community composition in *K. pygmaea* and *E. nutans*; and 3) how changes in the endophytic community explain host responses to climate warming.

## Materials and Methods

### Study site and sampling

Fieldwork was performed in August 2019 in the middle of the Qilian Mountains (37.68° N, 100.75° E) near Qilian County, which is located in the northeastern part of Qinghai Province, China. With an average elevation over 3 000 m, this area experiences a typical plateau continental climate. The mean annual and growing season temperatures (from June to August) are 1°C and 13°C, respectively. The annual mean precipitation is 480 mm, with 80% occurring during the growing season. The climate and vegetation composition change significantly with elevation in the Qilian Mountains. We sampled at elevations of 3 350 m, 3 460 m, 3 570 m, 3 680 m, and 3 800 m above sea level (a.s.l.) on the south slope of the mountain. The temperature decreases and the precipitation increases with increasing altitude in this area (Jin et al., 2017; Li et al., 2018). According to the vegetation investigation, the grassland was classified as alpine

127 meadows at elevations of 3 350 m, 3 460 m, and 3 570 m and alpine shrub meadows at  
128 elevations of 3 680 m and 3 800 m.

129 We defined three plots at each elevation, with each plot 20 m away from the other plots. The  
130 roots of *K. pygmaea* and *E. nutans* were sampled around each plot. *K. pygmaea* always  
131 aggregates in grasslands and forms slightly yellow green patches because of its dense network of  
132 roots, which are produced clonally. Therefore, patches of *K. pygmaea* and clumps of *E. nutans*  
133 were dug using a sterilized shovel from a depth of 10 cm. We reserved the belowground parts,  
134 including the soil and roots, when sampling, and separated the roots after transported to the  
135 laboratory. We reserved the belowground parts, including the soil and roots, when sampling, and  
136 separated the roots after transport to the laboratory. Five individuals of each plant species per  
137 plot composed one sample, which resulted in 30 root samples ( $2 \text{ species} \times 5 \text{ elevations} \times 3$   
138 replicates). All samples were placed in labeled sterile plastic bags and immediately stored in an  
139 incubator with dry ice. Samples were transported to the laboratory and stored at  $-80^{\circ}\text{C}$  until  
140 further processing. The roots of *K. pygmaea* and *E. nutans* were separated in the laboratory using  
141 sterilized scissors and brushes then transferred to 50-mL sterile centrifuge tubes containing 30  
142 mL of 0.01 M sterilized PBS (136 mM NaCl, 8 mM  $\text{Na}_2\text{HPO}_4$ , 2 mM  $\text{KH}_2\text{PO}_4$ , 2.6 mM KCl, pH  
143 7.2). The tubes were placed on a shaking platform (180 rpm) for 20 min to remove the soil  
144 attached to the root. The roots were rinsed in deionized water and surface sterilized according to  
145 the following steps: immersion in sterile water twice for 30 s, 75% ethanol for 1 min, 3.25%  
146 sodium hypochlorite for 3 min, 75% ethanol for 30 s, sterile water for 1 min, and a final rinse  
147 with sterile water. The surface-sterilized roots were stored in a freezer at  $-80^{\circ}\text{C}$ .



# **Genomic DNA extraction, amplification and purification**

Endophytic genomic DNA was extracted from 0.5 g of frozen surface-sterilized roots using the MOBIO PowerSoil® DNA Isolation Kit (MOBIO Laboratories, Carlsbad, CA, USA). The integrity and purity of the extracted DNA were tested on 1% agarose gels, and a NanoDrop One was used to measure the concentration and purity of the extracted DNA. The 16S rRNA gene V5-V7 region was amplified using the primers 799F (5'-AACMGGATTAGATACCCCKG-3') and 1193R (5'-ACGTCATCCCCACCTTCC-3') with a 12-bp barcode (*Horton et al., 2014*). The primer set 799F-1193R reduces co-amplification levels of mitochondrial gene during endophytic 16S rRNA gene PCR amplification (*Wang et al., 2018*). The internal transcribed spacer 1(ITS1) region of fungal rRNA was amplified using the forward primer 5'-CTTGGTCATTTAGAGGAAGTAA-3' and the reverse primer 5'-GCTGCGTTCTTCATCGATGC-3' (*Horton et al., 2014*). Primers were constructed by Invitrogen (Carlsbad, CA, USA). A 50-μL reaction containing 25 μL of 2x PremixTaq (Takara Biotechnology, Dalian Co. Ltd., China), 1 μL of each primer (10 mM) and 3 μL of DNA template (20 ng/μL) was used for PCR amplification in a BioRad S1000 (Bio-Rad Laboratory, CA). The following thermal cycling process was used: 5 min at 94°C for initialization; 30 cycles of denaturation at 94°C for 30 s; annealing at 52°C for 30 s; extension at 72°C for 30 s; and final elongation at 72°C for 10 min. The PCR products of 3 replicates of each sample were mixed, and 1% agarose gel electrophoresis was used to detect the quality of the PCR products. The products were mixed in equidensity ratios according to GeneTools analysis software (version 4.03.05.0, SynGene). An EZNA Gel Extraction Kit (Omega, USA) was used for purification of the mixed

169 PCR products.

# 170 **Library preparation and sequencing**

171 Libraries were processed with the NEBNext® Ultra™ DNA Library Prep Kit for Illumina®  
 172 (New England Biolabs, USA). The libraries were assessed using a Qubit® 2.0 fluorometer  
 173 (Thermo Scientific) and an Agilent Bioanalyzer 2100 system, and sequenced on the Illumina  
 174 HiSeq 2500 platform to generate 250-bp paired-end reads. High-quality clean reads were  
 175 obtained following the Trimmomatic quality control process (V0.33,  
 176 <http://www.usadellab.org/cms/?page=trimmomatic>). The clean paired-end reads from which the  
 177 barcodes and primers were removed, were merged using FLASH (V1.2.11,  
 178 <https://ccb.jhu.edu/software/FLASH/>). Effective clean tags were retained after filtering the  
 179 spliced sequences using Trimmomatic software.

# 180 **OTU cluster and species annotation**

181 Usearch software (V8.0.1517, <http://www.drive5.com/usearch/>) was applied for sequence  
 182 analysis. Sequences with  $\geq 97\%$  similarity were assigned to the same operational taxonomic unit  
 183 (OTU). Singletons were removed using usearch  
 184 ([http://www.drive5.com/usearch/manual/chimera\\_formation.html](http://www.drive5.com/usearch/manual/chimera_formation.html)) after OTU clustering, and the  
 185 chimera sequences were detected and removed using the UCHIME *de novo* algorithm  
 186 ([http://www.drive5.com/usearch/manual/uchime\\_algo.html](http://www.drive5.com/usearch/manual/uchime_algo.html)). The Greengenes  
 187 (<http://greengenes.secondgenome.com/>) and UNITE V8.0 databases were consulted based on the  
 188 RDP Classifier algorithm and the assign\_taxonomy.py script  
 189 ([http://qiime.org/scripts/assign\\_taxonomy.html](http://qiime.org/scripts/assign_taxonomy.html)) in QIIME for taxonomic annotation (the

confidence threshold was set to 0.8). OTUs that were annotated as chloroplasts or mitochondria (16S rRNA) and could not be annotated at the kingdom level were removed.

## Data analysis

The Shannon index of the endophytic bacterial and fungal communities was calculated using QIIME (V1.9.1), and the Kruskal-Wallis test at the  $P < 0.05$  level was used to examine the effects of elevation and species on the diversity of endophytic bacterial and fungal community. Nonmetric multidimensional scaling (NMDS) analysis based on Bray-Curtis distance was performed to visualize the dissimilarities of endophytic bacterial and fungal community composition in different species and at different elevations. Permutational multivariate analysis of variance (PerMANOVA) using the “Anosim” function in R (3.6.1) with 999 random permutations was performed to compare differences in endophytic community structure between plant species and elevations. The correlations of elevation and the relative abundances of the top 20 genera were analyzed using the Spearman method. Samples from different elevations were combined for further analysis of the differences in endophytic bacterial and fungal communities between the two plant species. The linear discriminant analysis (LDA) effect size (LEfSe) method with a threshold of 4 was used to investigate indicator species that exhibited significant differences in relative abundance between plant species. KEGG functional analysis of endophytic bacteria was performed using Phylogenetic Investigation of Communities by Reconstruction of Unobserved States (PICRUSt) (Langille *et al.*, 2013). We also analyzed genes associated with cold resistance, antioxidant enzymes (catalase, peroxidase and superoxide dismutase), nitrogen metabolism (nitrogen fixation, nodulation protein, nitrogen regulatory

protein, and nitrate reductase) and phosphorus metabolism (phosphatase).  $\text{Ca}^{2+}$  signaling (Monroy, Sarhan & Dhindsa, 1993), cold shock proteins, HSP20 family proteins (Elkelish et al., 2020), glutathione (Liu et al., 2020), glycerol-3-phosphate acyltransferases (Gomes et al., 2000), glycine betaine (Annunziata et al., 2019; Chen and Murata, 2011), glycosyltransferases (Shi et al., 2020), protein kinases (histidine kinase, serine protein kinase, and tyrosine kinase) (Martín and Busconi, 2001), trehalose (Liu et al., 2020), nitric oxide (Farnese et al., 2016), and antioxidant enzymes (Baier et al., 2019) have been reported to improve plant cold resistance.

## Results

### Diversity of endophytic bacterial and fungal communities

The rarefaction curves of the Shannon index reached a saturation plateau in all samples, which suggested that the sampling was sufficient to obtain most of the OTUs (Fig. S1). We found a higher Shannon index of the root endophytic bacterial and fungal communities in *K. pygmaea* than *E. nutans*. Elevation significantly affected the Shannon index of root endophytic bacteria in *K. pygmaea*, but not *E. nutans* (Fig. 1A). The Kruskal-Wallis test revealed that elevation had a significant effect on the Shannon index of root endophytic fungi in *E. nutans* ( $P = 0.021$ ) but not *K. pygmaea* ( $P > 0.05$ ). The Shannon index of endophytic fungi in *E. nutans* roots decreased significantly with increasing altitude, but an inconspicuous increasing trend was observed for *K. pygmaea* roots (Fig. 1B). There was no significant difference in the Shannon index of root endophytic fungi between *K. pygmaea* and *E. nutans* at lower elevations (3 350-3 570 m). The Shannon index of root endophytic fungi was much higher in *K. pygmaea* than *E. nutans* at higher elevations (3 680 and 3 800 m).

# Community compositions of endophytic bacteria and fungi

The NMDS ordination (Fig. 2) and analysis of similarities (ANOSIM) (Table 1) showed that endophytic bacterial and fungal community composition in *K. pygmaea* was significantly different than *E. nutans* ( $R=0.78$ ,  $P=0.001$ ;  $R=0.58$ ,  $P=0.001$ ) (Fig. 2A, 2B). Elevation also obviously influenced root endophytic bacterial and fungal communities in *E. nutans* ( $R=0.42$ ,  $P=0.001$ ;  $R=0.84$ ,  $P=0.001$ ) (Fig. 2C, 2D) and *K. pygmaea* ( $R=0.76$ ,  $P=0.001$ ;  $R=0.98$ ,  $P=0.001$ ) (Fig. 2E, 2F). The root endophytic bacterial community composition in *K. pygmaea* was more sensitive to elevation than *E. nutans*, which was validated by the R value of ANOSIM (Table 1). Greater the R values indicate a more significant influence of altitude on community composition. The endophytic fungal community was more sensitive to elevation than the bacterial community for each species. Plant identity had a greater influence on the endophytic bacterial community than the fungal community.

Proteobacteria was the predominant phylum in *E. nutans* and *K. pygmaea* roots (Fig. S2A). The average relative abundance of Proteobacteria was 58% in *K. pygmaea* roots, which was much lower than *E. nutans* (80%). Actinobacteria was the subdominant phylum in *K. pygmaea* roots and accounted for 30% of the bacterial reads, which was much higher than its abundance in *E. nutans* roots (1.4%). There were significant differences in root endophytic bacterial community composition at the order level between *K. pygmaea* and *E. nutans* (Fig. 3A). For *E. nutans* roots, Pseudomonadales was the most abundant order, with the relative abundance ranging from 36% to 69% along the elevation gradient. The endophytic bacterial community composition was more uniform in *K. pygmaea* roots than *E. nutans* roots. Rhizobiales accounted

for 20% of the total community abundance, and there was no obvious difference between elevations. Pseudomonadales was highly abundant (50%) at the elevation of 3 800 m, and Pseudonocardiales showed low abundance at this elevation.

Most endophytic fungi in all samples belonged to ascomycetes, with average 76% for *E. nutans* roots and 61% for *K. pygmaea* (Fig. S2B). Helotiales were the dominant order in *E. nutans* roots and were more abundant at high altitudes, which was different from that in *K. pygmaea* (Fig. 3B). Pleosporales were more abundant in *K. pygmaea* roots than *E. nutans* roots, especially at lower altitudes (3 350-3 570 m). Agaricales were enriched in *K. pygmaea* roots at higher altitudes (3 680 m and 3 800 m) and more abundant in *E. nutans* roots (Fig. 3B). The relative abundance of Hypocreales in *K. pygmaea* roots were 18.55% and 8.1% at 3 350 m and 3 460 m, respectively, which was higher than *E. nutans* roots at the same elevations (Fig. 3B).

We compared the abundances of the top 20 OTUs which covered more than 50% of the total reads in all samples at different altitudes and found that plants tended to be enriched with specific endophytes at different elevations (Fig. S3A-D). Species with significant differences in relative abundance between *K. pygmaea* and *E. nutans* were analyzed according to LEfSe (Fig. 4). The results indicated that bacterial species belonging to Actinobacteria (Frankiales, Pseudonocardiales and Micromonosporales) and Rhizobiales (Rhizobiaceae) were highly enriched in *K. pygmaea* roots, and Gammaproteobacteria (Burkholderiaceae), Pseudomonadales (Pseudomonadaceae) and Flavobacteriales (Flavobacteriaceae) were highly enriched in *E. nutans* roots (Fig. 4A). Fungal species belonging to Dothideomycetes (Pleosporales), Eurotiomycetes (Chaetothyriales), and Sordariomycetes (Coniochaetales, Hypocreales) were highly enriched in

*K. pygmaea* roots, and Helotiales (Hyaloscyphaceae, Helotiaceae) and Tremellales (Bulleribasidiaceae) were highly enriched in *E. nutans* roots (Fig. 4B).

The correlations of the relative abundances of the top 20 genera and elevation were analyzed (Table 2). For bacterial communities, the relative abundances of *Pantoea*, *Sphingomonas*, and *Paenibacillus* in *E. nutans* roots showed positive correlations with elevation. The relative abundances of *Pseudomonas*, *Pantoea*, *Serratia*, and *Paenibacillus* in *K. pygmaea* roots positively correlated with elevation, and *Rhizobium*, *Acidibacter*, *Actinophytocola*, and *Cryptosporangium* negatively correlated with elevation. For fungal communities, the relative abundances of *Mortierella* and *Cistella* showed negative correlations with elevation in *E. nutans* roots. The relative abundances of *Mycena*, *Mortierella*, *Ilyonectria*, *Sebacina*, and *Leptosphaeria* showed positive correlations with elevation in *K. pygmaea* roots.

The results of 16S rRNA functional prediction indicated that the endophytic bacterial community in *E. nutans* roots had a higher abundance of genes associated with nitrogen fixation, phosphatase activity and antioxidase activity and a lower abundance of genes associated with cold resistance than *K. pygmaea* roots (Fig. 5). Genes associated with nutrient absorption, including nitrogen fixation, nodulation, and phosphatase activity, were enriched in *K. pygmaea* roots at higher elevations. These genes were enriched at lower elevations in *E. nutans* roots, (Fig. S4).

## Discussion

### Diversity of the endophytic community along an elevation gradient

Most studies on the influence of climate change on plant associated microbial communities

overlooked host responses and did not link the responses of hosts and endophytes (*Qian et al., 2018; Cordier et al., 2012*). Therefore, the significance of changes in the endophytic community of the host in response to climate change were not well explained. Two plant species showing contrasting responses to climate warming were selected in the present study to fill in this knowledge gap. The increase in endophytic fungal diversity in *E. nutans* and decrease in *K. pygmaea* with warming (i.e., decreasing elevation) indicated that the endophytic fungal diversity of plants growing in suitable environments was higher than plants growing in unsuitable habitats. Our findings demonstrated that changes in endophytic fungal diversity under climate warming were closely related to host adaption, and endophytic diversity decreased when the climate became unfavorable for plant growth. A previous study indicated that endophytic diversity in sugar maple seedlings was higher within its natural range than at the edge of species' elevational range (*Wallace, Laforest-Lapointe & Kembel, 2018*), which was consistent with the results of our study. We hypothesized that the loss of endophytic diversity in adverse environments was related to host selection. The symbiotic relationships between the host and some endophytes break down, and some endophytes that are beneficial to host survival are selectively enriched (*Werner et al., 2018*).

Plant adaptability to environmental changes is very complex and affected by many factors. We just found a phenomenon that fungal diversity was higher in the habitat that suitable for host growth. However, no direct cause-and-effect relationship between fungal diversity and host adaptation was demonstrated in the present study. Whether higher endophytic diversity contributes to host growth is worthy to study in the future.



# **Species and elevation significantly affected the root endophytic bacterial and fungal communities**

Previous studies indicated that plant endophytes showed significant host preferences (*Toju, Kurokawa & Kenta, 2019; Yao et al., 2019*). Different plant species sharing similar environments recruit different microbial communities in roots (*Aleklett et al., 2015*), which may be related to the metabolic characteristics of the host. Because root exudates contain important factors that shape endophytic microbiome assembly, such as salicylic acid (*Lebeis et al., 2015*) and jasmonic acid (*Carvalhais et al., 2015*). Environmental parameters are also key factors that cause changes in the endophytic community (*Carper et al., 2018; Bei et al., 2019; Chen et al., 2019*). Climate characteristics, soil nutrients and vegetation composition change along altitude gradients, which may cause variations in the endophytic community (*Cai et al., 2020; Zarraonaindia et al., 2015*). Plant species and altitude significantly affected the root endophytic bacterial and fungal community composition in the present study. The bacterial community composition was affected more strongly by plant species, and elevation more strongly affected the fungal community. Previous studies also concluded that fungal communities were more susceptible to geographic distance than bacterial communities (*Coleman-Derr et al., 2016; Meiser, Bálint & Schmitt, 2014*).

Among the endophytic bacteria, Proteobacteria was the predominant phylum in *E. nutans* and *K. pygmaea* roots, which was consistent with many other studies involving different plant species (*Bulgarelli et al., 2012; Beckers et al., 2017; Carrell and Frank, 2015*). However, *K. pygmaea* roots had a higher abundance of Actinobacteria than *E. nutans* roots. At the order level, the relative abundances of Pseudonocardiales, Frankiales and Rhizobiales in *K. pygmaea* roots

were much higher than *E. nutans* roots. However, *E. nutans* roots had a higher abundance of Pseudomonadales. These root endophytes promote host plant growth and help plants resist biotic and abiotic stress (Hasegawa, et al., 2006; Santoyo et al., 2016). *Pseudomonas* participates in nitrogen fixation (Yan et al., 2008) and iodoacetamide (IAA) synthesis (Taghavi et al., 2009), which promote plant growth. In addition to symbiosis with leguminous plants, *Rhizobium* acts as an endophyte of nonleguminous plants to promote nitrogen and phosphorus uptake and improve the photosynthetic rate (Yanni et al., 2001; Chi, 2006). Helotiales, which promote phosphorus absorption in nonmycorrhizal plants (Almario et al., 2017), were enriched in *E. nutans* roots. Some genera, such as *Mortierella*, showed positive correlations with elevation in *K. pygmaea* but negative correlations with elevation in *E. nutans*, which also reflected the plant selectivity.

# **Functional differences of root endophytic bacteria in *K. pygmaea* and *E. nutans***

Chilling stress is an important environmental factor that affects plant growth and geographical distribution (Lv et al., 2018), which is especially true on the Qinghai-Tibet Plateau. A previous study also demonstrated that temperature and altitude factors primarily affected the distribution of alpine meadow grass (Lu et al., 2020). Symbiosis with bacteria that could improve the cold tolerance of the host represents a survival strategy in alpine regions with high elevations (Acuna-Rodriguez et al., 2020; Tiryaki, Aydin & Atici, 2019). Beirinckx et al. also indicated that the root microbiome promoted maize growth under chilling conditions (Beirinckx et al., 2020). The present study provided new insight into the role of endophytic bacteria in host adaptation to climate warming in terms of bacterial functional genes associated with cold resistance and nutrient absorption.

The study area in the present study exhibited a cold and wet climate at high altitudes and a warm and dry climate at low altitudes. Temperature is the main factor affecting plant growth at high elevations. Increased recruitment of bacteria with cold resistance ability in *K. pygmaea* roots made this plant more adaptable to high altitude than *E. nutans* although the abundance of genes associated with nutrient uptake was higher in *E. nutans* than *K. pygmaea*. Because the expression of these genes and the enzymes activity involved in nutrient absorption are restricted by low temperatures. Therefore, symbiosis with high abundance of these endophytes in *E. nutans* cost plant carbohydrates without getting the expected return.

The outcomes of symbiosis with microbiomes are environmentally dependent (Rubin *et al.*, 2020). For example, arbuscular mycorrhizal fungi significantly enhance phosphorus absorption and promote plant growth in phosphorus deficient soils, but the contribution of mycorrhizal pathways to phosphorus uptake in plants is weakened under high P soil condition (Chu *et al.*, 2020). Climate warming relieves cold stress at high elevations and exacerbates heat and drought stress at lower elevations (Reinhardt *et al.*, 2011; Moyes *et al.*, 2013). With climate warming or decreasing altitude, plant growth is freed from the cold limitation. Selectively symbiosis with bacteria that have the ability to facilitate host nutrient absorption, such as nitrogen fixation and phosphorus solubilization, in *E. nutans* roots made this plant more adaptable to warming environments than *K. pygmaea*.

Endophytes do not always establish harmonious symbioses with plant hosts. They also cause host disease and other adverse effects. There is a 'balance of antagonisms' relationship between fungal and plant partners (Schulz and Boyle 2005). We only focused on the "good" side of

endophytes for their importance in maintaining a stable symbiotic relationship. Exploring beneficial roles of endophytes to their hosts in special environments such as the Qinghai-Tibet Plateau contributes a lot to the development and application of microbial resources. However, functional prediction alone is not sufficient to directly demonstrate the role of endophytic bacteria in host responses to climate changes. Further studies should concentrate more on the interactions between endophytic community and host to rich our understanding of responses of terrestrial ecosystem to global warming.

## Conclusions

The present study demonstrated that the diversity of the endophytic community was higher in hosts growing in habitats that were conducive to its growth. *K. pygmaea*, with higher endophytic diversity and greater abundance of genes associated with cold resistance, was suited for growth in cold areas at high altitudes, and its growth was suppressed by warming. Higher endophytic diversity and greater abundance of genes associated with nutrient absorption and oxidation resistance in warmer environments (lower altitude) may contribute to the growth of *E. nutans* under global warming (Fig. 6). We also found that the relative abundances of the same taxa in different hosts showed different correlations with elevation, which demonstrated that the ability to recruit endophytes differed between the hosts when the habitat changed. Our study highlights the importance of plant endophytes in the responses of terrestrial ecosystems to

400 climate change.

# 401 Acknowledgements

402 We would like to thank Lu Yu, Wan-tong Zhang for their contribution to the field sampling;  
403 Chang-lin Xu for his help in plant species identification.

# 404 Conflict of interest:

405 The authors declare there are no competing interests

406

407

408

409

410

# 411 References

412 **Acuna-Rodriguez IS, Newsham KK, Gundel PE, Torres-Diaz C, Molina-Montenegro MA.**

413 2020. Functional roles of microbial symbionts in plant cold tolerance. *Ecology Letters* **23**:  
414 1034-1048.

415 **Aleklett K, Leff JW, Fierer N, Hart M.** 2015. Wild plant species growing closely connected in  
416 a subalpine meadow host distinct root-associated bacterial community. *Peer J.* **3**: e804.

417 **Almario J, Jeena G, Wunder J, Langen G, Zuccaro A, Coupland G, Bucher M.** 2017. Root-  
418 associated fungal microbiota of nonmycorrhizal *Arabis alpina* and its contribution to plant  
419 phosphorus nutrition. *PNAS* **114**: 9403-9412.

420 **Annunziata MG, Ciarmiello LF, Woodrow P, Dell'Aversana E, Carillo P.** 2019. Spatial and

temporal profile of glycine betaine accumulation in plants under abiotic stresses. *Frontiers in Plant Science*. **10**: 230.

**Baier M, Bittner A, Prescher A, van Buer J.** 2019. Preparing plants for improved cold tolerance by priming. *Plant Cell Environment*. **42**: 782-800.

**Beckers B, Beeck MOD, Weyens N, Boerjan W, Vangronsveld J.** 2017. Structural variability and niche differentiation in the rhizosphere and endosphere bacterial microbiome of field-grown poplar trees. *Microbiome* **5**: 25.

**Bei Q, Gerald M, Wu XH, Christoph M, Werner L.** 2019. Metatranscriptomics reveals climate change effects on the rhizosphere microbiomes in European grassland. *Soil Biology and Biochemistry* **138**: 107604.

**Beirinckx S, Viaene T, Haegeman A, Debode J, Amery F, Vandenabeele S.** 2020. Tapping into the maize root microbiome to identify bacteria that promote growth under chilling conditions. *Microbiome* **8**: 54.

**Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad N, Assenza F, Rauf P, Huettel B, Reinhardt R, Schmelzer E, Peplies J, Gloeckner FO, Amann R, Eickhorst T, Schulze-Lefert P.** 2012. Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial microbiota. *Nature* **488**: 91-95.

**Cai ZQ, Wang XB, Sreetama B, Gao Q.** 2020. Distinct factors drive the assembly of quinoa-associated microbiomes along elevation. *Plant Soil* **448**: 55-69.

**Carper DL, Carrell AA, Kueppers LM, Frank AC.** 2018. Bacterial endophyte communities in *Pinus flexilis* are structured by host age, tissue type, and environmental factors. *Plant Soil*

**428:** 335-352.

**Carrell AA, Frank AC.** 2015. Bacterial endophyte communities in the foliage of coast redwood and giant sequoia. *Frontiers in Microbiology* **6**: 1008.

**Carrión VJ, Perez-Jaramillo J, Cordovez V, Tracanna V, de Hollander M, Ruiz-Buck D, Mendes LW, Van IJcken W, Gomez-Exposito R, Elsayed SS, Mohanraju P, Arifah A, van der Oost J, Paulson JN, Mendes R, van Wezel GP, Medema MH, Raaijmakers JM.** 2019. Pathogen-induced activation of disease-suppressive functions in the endophytic root microbiome. *Science* **366**: 606-612.

**Carvalhais LC, Dennis PG, Badri DV, Kidd BN, Vivanco JM, Schenk PM.** 2015. Linking jasmonic acid signaling, root exudates, and rhizosphere microbiomes. *Molecular Plant-Microbe Interactions* **28**: 1049-1058.

**Chen P, Zhao M, Tang F, Hu Y, Peng X, Shen S.** 2019. The effect of environment on the microbiome associated with the roots of a native woody plant under different climate types in China. *Applied Microbiology and Biotechnology* **103**: 3899-3913.

**Chen TH, Murata N.** 2011. Glycinebetaine protects plants against abiotic stress: mechanisms and biotechnological applications. *Plant Cell Environment* **34**: 1-20.

**Chi F.** 2006. Migration of rhizobia in plants and proteome analysis of their interaction. Beijing: Graduate School of Chinese Academy of Sciences. *Doctor Disseration*.

**Chu Q, Zhang L, Zhou J, Yuan L, Chen F, Zhang F, Feng G, Rengel Z.** 2020. Soil plant-available phosphorus levels and maize genotypes determine the phosphorus acquisition efficiency and contribution of mycorrhizal pathway. *Plant Soil* **449**:357-371

- Coleman-Derr D, Desgarennnes D, Fonseca-Garcia C, Gross S, Clingenpeel S, Woyke T, North G, Visel A, Partida-Martinez LP, Tringe SG.** 2016. Plant compartment and biogeography affect microbiome composition in cultivated and native *Agave* species. *New Phytologist* **209**: 798-811.
- Cordier T, Robin C, Capdevielle X, Fabreguettes O, Desprez-Loustau ML, Vacher C.** 2012. The composition of phyllosphere fungal assemblages of European beech (*Fagus sylvatica*) varies significantly along an elevation gradient. *New Phytologist* **196**: 510-519.
- Dini-Andreote F.** 2020. Endophytes: the second layer of plant defense. *Trends Plant Science* **25**: 319-322.
- Elkelish A, Qari SH, Mazrou YSA, Abdelaal KAA, Hafez YM, Abu-Elsaoud AM, Hafez YM, Abu-Elsaoud AM, Batiha GE, El-Esawi MA, El Nahhas N.** 2020. Exogenous ascorbic acid induced chilling tolerance in tomato plants through modulating metabolism, osmolytes, antioxidants, and transcriptional regulation of catalase and heat shock proteins. *Plants* **9**: 431.
- Farnese FS, Menezes-Silva PE, Gusman GS, Oliveira JA.** 2016. When bad guys become good ones: The key role of reactive oxygen species and nitric oxide in the plant responses to abiotic stress. *Frontiers in Plant Science* **7**: 471.
- Fujimura KE, Egger KN, Henry GH.** 2008. The effect of experimental warming on the root-associated fungal community of *Salix arctica*. *ISME J* **2**: 105-114.
- Gomes E, Jakobsen MK, Axelsen KB, Geisler M, Palmgren MG.** 2000. Chilling tolerance in *Arabidopsis* involves ALA1, a member of a new family of putative aminophospholipid



translocases. *Plant Cell* **12**: 2441-2453.

**Hasegawa S, Meguro A, Shimizu M, Nishimura T, Kunoh H.** 2006. Endophytic actinomycetes and their interactions with host plants. *Actinomycetologica* **20**: 72-81.

**Hiruma K, Gerlach N, Sacristán S, Nakano RT, Hacquard S, Kracher B, Neumann U, Ramírez D, Bucher M, O’Connell RJ, Schulze-Lefert P.** 2016. Root endophyte *Colletotrichum tofieldiae* confers plant fitness benefits that are phosphate status dependent. *Cell* **165**: 464-474.

**Horton MW, Bodenhausen N, Beilsmith K, Meng D, Muegge BD, Subramanian S, Vetter MM, Vilhjalmsen BJ, Nordborg M, Gordon JJ, Bergelson, J.** 2014. Genome-wide association study of *Arabidopsis thaliana* leaf microbial community. *Nat Commun.* **5**:5320.

**Jackrel SL, Schmidt KC, Cardinale BJ, Denef VJ.** 2020. Microbiomes reduce their host’s sensitivity to interspecific interactions. *MBio.* **11**: e02657-19.

**Jin CW, Zhao YG, Li XS, Zhi JJ, Zhang GL.** 2017. Spatial distribution and environmental factors affecting mattic epipedon at different developmental levels in alpine meadows in the middle of Qilian Mountains. *Acta Ecologica Sinica* **37**: 6732-6742.

**Korner C.** 2007. The use of “altitude” in ecological research. *Trends Ecology and Evolution* **22**: 569-574.

**Laforest-Lapointe I, Messier C, Kembel SW.** 2016. Host species identity, site and time drive temperate tree phyllosphere bacterial community structure. *Microbiome* **4**: 27.

**Laforest-Lapointe I, Paquette A, Messier C, Kembel SW.** 2017. Leaf bacterial diversity mediates plant diversity and ecosystem function relationships. *Nature* **546**: 145-147.

- Langille MGI, Zaneveld J, Caporaso JG, McDonald D, Knights D, JA Reyes, Clemente JC, Burkepile DE, Vega Thurber RL, Knight R. 2013. Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. *Nature Biotechnology* **31**: 814.
- Lebeis SL, Paredes SH, Lundberg DS, Breakfield N, Gehring J, McDonald M, Malfatti S, del Rio D, Glavina T, Jones CD, Tringe SG. 2015. Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Science* **349**: 860-864.
- Li L, Yang A, Zhao Z. 2005. Seasonality of arbuscular mycorrhizal symbiosis and dark septate endophytes in a grassland site in southwest China. *FEMS Microbiol Ecology* **54**: 367-373.
- Li YQ, Wang XY, Niu YY, Lian J, Luo YQ, Chen YP. 2018. Spatial distribution of soil organic carbon in the ecologically fragile Horqin Grassland of northeastern China. *Geoderma* **325**: 102-109.
- Little CJ, Wheeler JA, Sedlacek J, Cortés AJ, Rixen C. 2016. Small-scale drivers: the importance of nutrient availability and snowmelt timing on performance of the alpine shrub *Salix herbacea*. *Oecologia* **180**: 1015-1024.
- Liu F, Zhang X, Cai B, Pan D, Fu X, Bi H, Ai X. 2020. Physiological response and transcription profiling analysis reveal the role of glutathione in H<sub>2</sub>S-induced chilling stress tolerance of cucumber seedlings. *Plant Science* **291**: 110363.
- Liu H, Brettell LE. 2019. Plant defense by VOC-induced microbial priming. *Trends Plant Science* **24**: 187-189.
- Liu H, Brettell LE, Qiu Z, Singh BK. 2020. Microbiome-mediated stress resistance in plants.

- Trends Plant Science*. <https://doi.org/10.1016/j.tplants.2020.03.014>.
- Liu H, Mi Z, Lin L, Wang Y, Zhang Z, Zhang F.** 2018. Shifting plant species composition in response to climate change stabilizes grassland primary production. *PNAS* **115**: 4051-4056.
- Lv XZ, Li HZ, Chen XX, Xiang X, Guo ZX, Yu JQ, Zhou YH.** 2018. The role of calcium-dependent protein kinase in hydrogen peroxide, nitric oxide and ABA-dependent cold acclimation. *Journal of Experimental Botany* **69**: 4127-4139.
- Ma ZY, Liu HY, Mi ZR, Zhang ZH, Wang YH, Xu W, Jiang L, He JS.** 2017. Climate warming reduces the temporal stability of plant community biomass production. *Nature Communications* **8**:15378.
- Martín ML, Busconi L.** 2001. A rice membrane-bound Calcium-dependent protein kinase is activated in response to low temperature. *Plant Physiology*. **125**: 1442-1449.
- Martinez-Garcia LB, De Deyn GB, Pugnaire FI, Kothamasi D, van der Heijden MGA.** 2017. Symbiotic soil fungi enhance ecosystem resilience to climate change. *Global Change Biology* **23**: 5228-5236.
- Meiser A, Bálint M, Schmitt I.** 2014. Meta-analysis of deep-sequenced fungal communities indicates limited taxon sharing between studies and the presence of biogeographic patterns. *New Phytologist* **201**: 623-635.
- Miehe G, Miehe S, Kaiser K, Liu JQ, Zhao XQ.** 2008. Status and dynamics of *Kobresia pygmaea* ecosystem on the Tibetan plateau. *Ambio*. **37**: 272-279.
- Monroy AF, Sarhan F, Dhindsa RS.** 1993. Cold-induced change in freezing tolerance, protein phosphorylation, and gene expression. *Plant Physiology* **102**: 1127-1135.

- 547 **Morin X, Viner D, Chuine I. 2008.** Tree species range shifts at a continental scale: new  
548 predictive insights from a process-based model. *Journal of Ecology* **96**:784-794
- 549 **Moyes AB, Castanha C, Germino MJ, Kueppers LM. 2013.** Warming and the dependence of  
550 limber pine (*Pinus flexilis*) establishment on summer soil moisture within and above its  
551 current elevation range. *Oecologia* **171**: 271-282.
- 552 **Neal AL, Ahmad S, Gordon-Weeks R, Ton J. 2012.** Benzoxazinoids in root exudates of maize  
553 attract *Pseudomonas putida* to the rhizosphere. *PLoS One* **7**: e35498.
- 554 **Niu Y, Yang S, Zhou J, Chu B, Ma S, Zhu H. 2019.** Vegetation distribution along mountain  
555 environmental gradient predicts shifts in plant community response to climate change in  
556 alpine meadow on the Tibetan Plateau. *Science of Total Environment*. **650**: 505-514.
- 557 **Odelade KA, Babalola OO. 2019.** Bacteria, fungi and archaea domains in rhizospheric soil and  
558 their effects in enhancing agricultural productivity. *International Journal of Environmental*  
559 *Research and Public Health* **16**: 3873.
- 560 **Porras-Alfaro A, Bayman P. 2011.** Hidden fungi, emergent properties: endophytes and  
561 microbiomes. *Annual Review of Phytopathology* **49**: 291-315.
- 562 **Qi J, Liu WH, Jiao T, Hamblin A. 2020.** Variation in morphological and physiological  
563 characteristics of wild *Elymus nutans* ecotypes from different altitudes in the northeastern  
564 Tibetan Plateau. *Journal of Sensors* **11**: 1-11.
- 565 **Qian X, Chen L, Guo XM, He D, Shi MM, Zhang DX. 2018.** Shifts in community  
566 composition and co-occurrence patterns of phyllosphere fungi inhabiting *Mussaenda*  
567 *shikokiana* along an elevation gradient. *Peer J* **6**: e5767.

- Qian X, Li HZ, Wang YL, Wu BW, Wu MS, Chen L, Qian X.** 2019. Leaf and root endospheres harbor lower fungal diversity and less complex fungal co-occurrence patterns than rhizosphere. *Frontiers in Microbiology* **10**: 15.
- Reinhardt K, Castanha C, Germino MJ, Kueppers LM.** 2011. Ecophysiological variation in two provenances of *Pinus flexilis* seedlings across an elevation gradient from forest to alpine. *Tree Physiology* **31**: 615-625.
- Rubin RL, Jones AN, Hayer M, Shuman-Goodier ME, Andrews LV, Hungate BA.** 2020. Opposing effects of bacterial endophytes on biomass allocation of a wild donor and agricultural recipient. *FEMS Microbiology Ecology*. doi:10.1093/femsec/fiaa012.
- Santoyo G, Moreno-Hagelsieb G, Orozco-Mosqueda MC, Glick BR.** 2016. Plant growth-promoting bacterial endophytes. *Microbiological Research* **183**: 92-99.
- Schulz B, Boyle C.** 2005. The endophytic continuum. *Mycological Research* **109**: 661-686.
- Shi Y, Phan H, Liu Y, Cao S, Zhang Z, Chu C, Schläppi MR.** 2020. Glycosyltransferase OsUGT90A1 helps protect the plasma membrane during chilling stress in rice. *Journal of Experimental Botany* **71**: 2723-2739.
- Taghavi S, Garafola C, Monchy S, Newman L, Hoffman A., Weyens N, Barac T, Vangronsveld J, van der Lelie D.** 2009. Genome survey and characterization of endophytic bacteria exhibiting a beneficial effect on growth and development of poplar trees. *Applied Environmental Microbiology* **75**: 748-757.
- Tiryaki D, Aydin I, Atici O.** 2019. Psychrotolerant bacteria isolated from the leaf apoplast of cold-adapted wild plants improve the cold resistance of bean (*Phaseolus vulgaris* L.) under

low temperature. *Cryobiology* **86**: 111-119.

**Toju H, Kurokawa H, Kenta T.** 2019. Factors influencing leaf and root-associated communities of bacteria and fungi across 33 plant orders in a grassland. *Front Microbiol* **10**. DOI: 10.3389/fmicb.2019.00241

**van der Heijden MGA, de Bruin S, Luckerhoff L, van Logtestijn RSP, Schlaeppi K.** 2016. A widespread plant-fungal-bacterial symbiosis promotes plant biodiversity, plant nutrition and seedling recruitment. *ISME J* **10**: 389-399.

**Wang F, Men X, Zhang G, Liang K, Xin Y, Wang J, Li A, Zhang H, Liu H, Wu L.** 2018. Assessment of 16S rRNA gene primers for studying bacterial community structure and function of aging flue-cured tobaccos. *AMB Express* **8**: 182.

**Wang Q, Zhang ZH, Du R, Wang SP, Duan JC, Iler AM.** 2019. Richness of plant communities plays a larger role than climate in determining responses of species richness to climate change. *Journal of Ecology* **107**: 1944-1955.

**Wallace J, Laforest-Lapointe I, Kembel SW.** 2018. Variation in the leaf and root microbiome of sugar maple (*Acer saccharum*) at an elevational range limit. *Peer J* **6**: e5293.

**Wu L, Wang M, OUYANG H, Cheng SK, Song MH.** 2017. Spatial distribution modelling of *Kobresia pygmaea* (Cyperaceae) on the Qinghai--Tibetan Plateau. *Journal of Resource Ecology* **8**:20-29.

**Yan YL, Yang J, Dou Y, Chen M, Ping SZ, Peng JP, Lu W, Zhang WP, Yan ZY, Li HQ, Liu W, He S, Geng LZ, Zhang XB, Yang F, Yu HY, Zhan YH, Li DH, Lin ZL, Wang YP, Elmerich C, Lin M, Jin Q.** 2008. Nitrogen fixation island and rhizosphere competence

traits in the genome of root-associated *Pseudomonas stutzeri* A1501. *PNAS* **105**: 7564-7569.

**Yanni YG, Rizk YR, Abd El-Fattah FK, Squartini A, Corich V, Giacomini A, de Bruijn F, Rademaker J, Maya-Flores J, Ostrom P, Vega-Hernande M, Hollingsworth RI, Martinez-Molina E, Mateos P, Velazquez E, Wopereis J, Triplett E, Umali-Garcia M, Anarna JA, Rolfe BG, Ladha JK, Hill J, Mujoo R, Ng PK, Dazzo FB.** 2001. Beneficial Plant Growth-Promoting Association of *Rhizobium leguminosarum* bv. *trifolii* with Rice Roots. *Australian journal of plant physiology* **28**: 845-870.

**Yao H, Sun X, He C, Maitra P, Li XC, Guo LD.** 2019. Phyllosphere epiphytic and endophytic fungal community and network structures differ in a tropical mangrove ecosystem. *Microbiome* **7**: 57.

**Zarraonaindia I, Owens SM, Weisenhorn P, West K, Hampton-Marcell J, Lax S, Bokulich NA, Mills DA, Martin G, Taghavi S.** 2015. The soil microbiome influences grapevine-associated microbiota. *MBio* **6**: e02527-14.

**Zhang L, Zhang W, Li Q, Cui R, Wang Z, Wang Y, Zhang YZ, Ding W, Shen XH.** 2020. Deciphering the root endosphere microbiome of the desert plant *Alhagi sparsifolia* for drought resistance-promoting bacteria. *Applied and Environmental Microbiology* **86**: e02863-19.

**Zilber-Rosenberg I, Rosenberg E.** 2008. Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiol Reviews* **32**: 723-735.

# **Table 1**(on next page)

Table 1. Significance test of community composition differences among elevations and hosts determined by permutational multivariate analysis of variance using distance matrices (PERMANOVA) according to Bray-Curtis distances.



**Table 1.** Significance test of community composition differences among elevations and hosts determined by permutational multivariate analysis of variance using distance matrices (PERMANOVA) according to Bray-Curtis distances.

			Anosim	
			<i>R</i>	<i>P</i>
Bacteria	Elevation	<i>E. nutans</i>	0.416	0.001
		<i>K. pygmaea</i>	0.756	0.001
	Host species	<i>E. nutans</i> vs <i>K. pygmaea</i>	0.78	0.001
Fungi	Elevation	<i>E. nutans</i>	0.84	0.001
		<i>K. pygmaea</i>	0.984	0.001
	Host species	<i>E. nutans</i> vs <i>K. pygmaea</i>	0.579	0.001

**Table 2**(on next page)

Spearman correlation analysis between elevation and the relative abundance of bacterial and fungal at genus level.


**Table 2.** Spearman correlation analysis between elevation and the relative abundance of bacterial and fungal at genus level.

	Genus	<i>K. pygmaea</i>		<i>E. nutans</i>	
		<i>R</i>	<i>P</i>	<i>R</i>	<i>P</i>
Bacteria	<i>Pseudomonas</i>	<b>0.731</b>	<b>0.002</b>	0.153	0.587
	<i>Pantoea</i>	<b>0.633</b>	<b>0.011</b>	<b>0.556</b>	<b>0.031</b>
	<i>Rhizobium</i>	<b>-0.546</b>	<b>0.035</b>	0.011	0.969
	<i>Acidibacter</i>	<b>-0.676</b>	<b>0.006</b>	-0.207	0.458
	<i>Actinophytocola</i>	<b>-0.524</b>	<b>0.045</b>	0.022	0.938
	<i>Cryptosporangium</i>	<b>-0.676</b>	<b>0.006</b>	-0.016	0.954
	<i>Serratia</i>	<b>0.535</b>	<b>0.040</b>	0.491	0.063
	<i>Sphingomonas</i>	-0.262	0.346	<b>0.644</b>	<b>0.010</b>
	<i>Paenibacillus</i>	<b>0.742</b>	<b>0.002</b>	<b>0.764</b>	<b>0.001</b>
Fungi	<i>Mycena</i>	<b>1.000</b>	<b>0.000</b>	-0.700	0.188
	<i>Mortierella</i>	<b>0.900</b>	<b>0.037</b>	<b>-0.900</b>	<b>0.037</b>
	<i>Cistella</i>	0.700	0.188	<b>-0.900</b>	<b>0.037</b>
	<i>Ilyonectria</i>	<b>0.900</b>	<b>0.037</b>	-0.800	0.104
	<i>Sebacina</i>	<b>0.900</b>	<b>0.037</b>	-0.700	0.188
	<i>Leptosphaeria</i>	<b>0.975</b>	<b>0.005</b>	-0.616	0.269

# Figure 1

Shannon index of root endophytic bacteria (A) and fungi (B) in *K. pygmaea* and *E. nutans* at different elevations.


Averages  $\pm$  SE are displayed (n=3). Different letters indicate significant differences between elevations



# Figure 2

Non-metric multidimensional scaling (NMDS) ordination of the community composition of endophytic bacteria and fungi based on Bray-Curtis distances.

A and B indicated fungal and bacterial community composition in *E. nutans* and *K. pygmaea* root regardless of elevations; C and D indicated fungal and bacterial community composition at different elevations in *E. nutans* root; E and F indicated fungal and bacterial community composition at different elevations in *K. pygmaea* root. Plant species were distinguished by colors, and elevations were distinguished by shapes.




# Figure 3

Relative abundance of bacterial (a) and fungal taxa (b) at the order level.

ER *E. nutans* root; KR *K. pygmaea* root; the numbers represent elevations. For example, ER3350 indicate *E. nutans* root at the elevation of 3 350 m; KR3350 indicate *K. pygmaea* root at the elevation of 3 350 m. The same as below.






# Figure 4


Cladograms of LEfSe showing bacterial and fungal taxa with significant differences in relative abundance between *K. pygmaea* and *E. nutans*.

The filled circles from inside to outside indicate the taxonomic levels with phylum, class, order, family, genus, and species. Circles or nodes shown in color corresponding to different plant species represented a significantly more abundant group. Yellow circles indicate species with no significant differences in relative abundance.



# Figure 5

Differences in relative abundance of functional genes associated with cold resistance, nitrogen absorption, phosphatase and antioxidase between *K. pygmaea* and *E. nutans*.



# Figure 6

Frame diagram of response of endophytic bacterial and fungal community to altitude

The proportion of *Elymus nutans* in vegetation community as well as its endophytic fungal diversity decreased with elevation increasing. While, the proportion of *Kobresia pygmaea* as well as its endophytic fungal diversity showed a slight increase trend with elevation increasing. The Shannon index of root bacterial and fungal community in *Kobresia pygmaea* were higher than that in *Elymus nutans*. Besides, bacteria in *Kobresia pygmaea* root had more genes associated with cold resistance, and in *Elymus nutans* root had more genes associated with nutrient absorption.

