

Similar seed dispersal systems between native and alien plant species with local frugivorous birds in a coastal seawall forest (#55858)

1

First submission

Guidance from your Editor

Please submit by **4 Jan 2021** for the benefit of the authors (and your \$200 publishing discount) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

7 Figure file(s)

2 Table file(s)

5 Raw data file(s)

! Custom checks

Vertebrate animal usage checks



Have you checked the authors [ethical approval statement](#)?



Were the experiments necessary and ethical?



Have you checked our [animal research policies](#)?

Field study



Have you checked the authors [field study permits](#)?



Are the field study permits appropriate?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Similar seed dispersal systems between native and alien plant species with local frugivorous birds in a coastal seawall forest

Bin Liu^{1,2}, Guohai Wang¹, Yuting An², Dandan Xue², Libo Wang², Changhu Lu^{Corresp. 1}

¹ College of Biology and the Environment, Nanjing Forestry University, Nanjing, Jiangsu, China

² Management Bureau of Dafeng Milu National Nature Reserve, Yancheng, Jiangsu, China

Corresponding Author: Changhu Lu

Email address: luchanghu@njfu.com.cn

Frugivorous birds play an important role in seed dispersal. Many alien plant species usually need local birds for their seed dispersal and established population in a new habitat. Alien plant species which produced similar fruits with the native species may be easier to attract local birds, and the new established mutualism systems would be similar with local ones. In autumn of 2018 and 2019, we selected an alien plant species, *Phytolacca Americana*, and a native plant species, *Cayratia japonica* in a coastal seawall forest to study their seed dispersal pattern. The fruit characteristics, local frugivorous birds foraging behaviors on the fruits, seedlings microhabitats and seed germination rates of both plant species were examined to find out if the seed dispersal system of alien species was similar to that of native one. Our results showed that the fruit type, color and ripening period of *Phytolacca americana* were much similar to those of *Cayratia japonica*, which were attracting the visitation of local birds. The positive correlation between the percent fruit ripening rate and the percent fruit missing rate of both plant species showed that local frugivorous birds have the potential to disperse the alien plant sufficiently to enable its spread in the coastal seawall forest (General Line Model, $P < 0.01$). 11 bird species consumed fruits of both alien species and native species in the study period. Bird foraging behaviors as feeding frequency (U -test, $P > 0.05$), feeding duration (t -test, $P > 0.05$) and first stop distance (t -test, $P > 0.05$) also were similar and showed that a stable relationship in the process of seed dispersal had been established between the alien plant and local frugivorous birds in the study site. The larger amount of the fruits of alien plant carried each time by birds even indicated that *Phytolacca americana* was a little more effective in its seed dispersal than that of *Cayratia japonica* (t -test, $P < 0.01$). Seedlings of alien plant preferred forest gap microhabitat (t -test, $P < 0.01$). Bird digestion treatment could promote seed germination success in both plant species. Our study suggests that in a narrow coastal seawall forest the alien plant species establish its population successfully relying on the similar seed dispersal systems with the local one.

Similar seed dispersal systems between native and alien plant species with local frugivorous birds in a coastal seawall forest

Bin Liu^{1,2}, Guohai Wang¹, Yuting An², Dandan Xue², Libo Wang², Changhu Lu¹

¹ College of Biology and the Environment, Nanjing Forestry University, Nanjing, Jiangsu, China

² Management Bureau of Dafeng Milu National Nature Reserve, Yancheng, Jiangsu, China

Corresponding Author:

Changhu Lu¹

Longpan road 159, Nanjing, Jiangsu, 210037, China

Email address: luchanghu@njfu.com.cn

Abstract

Frugivorous birds play an important role in seed dispersal. Many alien plant species usually need local birds for their seed dispersal and established population in a new habitat. Alien plant species which produced similar fruits with the native species may be easier to attract local birds, and the new established mutualism systems would be similar with local ones. In autumn of 2018 and 2019, we selected an alien plant species, *Phytolacca Americana*, and a native plant species, *Cayratia japonica* in a coastal seawall forest to study their seed dispersal pattern. The fruit characteristics, local frugivorous birds foraging behaviors on the fruits, seedlings microhabitats and seed germination rates of both plant species were examined to find out if the seed dispersal system of alien species was similar to that of native one. Our results showed that the fruit type, color and ripening period of *Phytolacca americana* were much similar to those of *Cayratia japonica*, which were attracting the visitation of local birds. The positive correlation between the percent fruit ripening rate and the percent fruit missing rate of both plant species showed that local frugivorous birds have the potential to disperse the alien plant sufficiently to enable its spread in the coastal seawall forest (General Line Model, $P < 0.01$). 11 bird species consumed fruits of both alien species and native species in the study period. Bird foraging behaviors as feeding frequency (U -test, $P > 0.05$), feeding duration (t -test, $P > 0.05$) and first stop distance (t -test, $P > 0.05$) also were similar and showed that a stable relationship in the process of seed dispersal had been established between the alien plant and local frugivorous birds in the study site. The larger amount of the fruits of alien plant carried each time by birds even indicated that *Phytolacca americana* was a little more effective in its seed dispersal than that of *Cayratia japonica* (t -test, $P < 0.01$). Seedlings of alien plant preferred forest gap microhabitat (t -test, $P < 0.01$). Bird digestion treatment could promote seed germination success in both plant species.

Our study suggests that in a narrow coastal seawall forest the alien plant species establish its population successfully relying on the similar seed dispersal systems with the local one.

Introduction

Seed dispersal was critical for maintenance of plant population (Butler et al., 2007; Wilson & Downs, 2012), and alien invasions (Schierenbeck, 2004; Buckley et al., 2006). Some plant species relied on wind, water and other external forces to spread their seeds far away, while more others were dependent on frugivorous birds for seed dispersal (Moles et al., 2005; Prado, 2013). A reciprocal relationship between plant and frugivorous birds had been established in the evolution process (Lu, 2003; Li et al., 2019). Plants provided necessary nutrients for birds, in return, birds dispersed seeds to other habitats and improved plant regeneration (Camargo et al., 2019; Morán-López et al., 2020; Galindo-González, Guevara & Sosa, 2000). Bird-mediated seed dispersal was the most important pattern in the ecosystem, and birds became the most common seed disperser in the world (Saavedra et al., 2014; Wang & Ives, 2017).

Frugivorous birds play a key role in seed dispersal of many plant species (Schleuning et al., 2011; Duan, Goodale & Quan, 2013; Lovas-Kiss, 2020), including invasive alien plants (Thabethe et al., 2015). It was urgent for alien plant species which relied on birds for seed dispersal, to establish a rapid adaptation mechanism to settle down, build population and spread rapidly after being carried to a new habitat. Alien plants with fleshy fruits should establish effective dispersal interactions with local frugivorous birds (Drummond, 2005; Kollmann et al., 2007). Or else, they would be failed to settle in a new environment or greatly reduced potential for spreading in the competition with native plants (Aslan, 2011).

Relationship between alien plants and local birds who feed on their fruits and dispersal their seeds have been carried in many researches (Renne et al., 2002; Gosper, Stansbury & Vivian-Smith, 2005). The adaptability of most alien plant species was based on the variation seed dispersal systems to native plant species. For example, a study in a coastal belt forest in South Africa showed that fleshy fruit introduced plants that were effectively dispersed were small-seed, open habitats species with longer fruiting length (Bitanni et al., 2020). A study based on proportion of introduced plant fruits removed by birds in northern California, USA showed that all three non-native species, whose fruits differed greatly in type and appearance with native plants, had been assimilated into local bird diets, and there was no limitation for them to dispersal by birds (Aslan, 2011). In secondary forest sites on Bonin island, Japan, a study showed that the introduced plant fruits can dispersed differently by both native and non-native bird species, and new mutualistic relationships involving native and introduced birds and plants had been established again in a seed-dispersal system (Kawakami, Frederiksen & Vestergaard, 2009). But if the seed dispersal system established by alien plant species and local frugivorous birds are the similar to native one has not been detailed reported yet.

An alien fleshy fruited plant species *Phytolacca americana* and a native one *Cayratia japonica* were found coexist in a coastal seawall forest in the Yellow Sea wetland, Jiangsu province, China. The forest was a long corridor with artificial trees, which could provide

migration channels, foraging places and breeding sites for kinds of birds (Liu et al., 2020). The seed dispersal of *Phytolacca americana* had been reported both in China and its native land (Mcdonnell et al., 1984; Li et al., 2019), but not the plant species *Cayratia japonica*. Fruits of both the plant species were found to be feed by groups of frugivorous birds in autumn in the coastal seawall forest in previous observations, the forest was thus an excellent field site to study the seed dispersal system of native and introduced plant species with local frugivorous birds.

Both *Cayratia japonica* and *Phytolacca americana* were the dominate fruit plants in autumn in the study site, and we select these two species to examine the mutualistic systems established between the alien plant species and the local frugivorous birds, and whether the alien plant and the native plant have the similar seed dispersal system based on fruit characteristics, local frugivorous birds foraging behaviors, microhabitats selected by seedlings and seed germination rates.

Materials & Methods

Ethics statement

Field studies were conducted under the permission from the Administrative Bureau of Dafeng Milu National Nature Reserve.

Study site

The field work was conducted in Dongchuan coastal seawall forest in Dafeng Milu National Nature Reserve, Jiangsu Province, China (32°59'N~33°03'N, 120°47'E~120°53'E). The forest was planted in the 1990s, mainly used for windbreak and dike consolidation. The region is characterized by a transition climate between subtropical and warm temperate. Mean annual precipitation in the area is approximately 1068 mm (concentrated in summer), and mean annual temperature is 14.1 °C. The vegetation is dominated by *Robinia pseudoacacia*, with *Metasequoia glyptostroboides* and *Populus canadensis* slightly. The width of the forest is about 20~50m. The habitat in one side of the forest is intertidal zone with the main vegetation *Spartina alterniflora*, *Suaeda glauca*, *Phragmites communis* and *Imperata cylindrica*, the other side was farmland and culture pond. Many open forest gaps were exposed after some trees were blown every year by typhoon. Many bird species were living in the forest such as Light-vented Bulbul *Pycnonotus sinensis*, Common Blackbird *Turdus merula* and Black-billed Magpie *Pica pica*. From April to May in spring and September to October in autumn, many passerines passed through the forest to complete their migration life cycle, mainly including warblers and flycatchers. In autumn, there were also a small number of other fruits species in the study area beside *Cayratia japonica* and *Phytolacca americana*, such as *Lycium barbarum*, *Sapium sebiferum*, *Ligustrum lucidum*, and *Paederia scandens*.

Field observation

Fruits and seedlings investigation

From September to October, 2018, the two focal fruit plant species were inspected every 50m along the coastal seawall forest. The fruit characteristics, fruit height and microhabitat selecting were recorded. Fruit size were measured by a caliper based on 20 mature fruits each species. The density of each species was investigated in both study site and another site about 8 km away, representing a younger age of stand, in order to compare the density difference of these two fruit plants between older or younger age of stand. The number of ripe fruits of each species were investigated every 10 days, to calculated the rates of ripe and missing fruits. Seedlings of *Phytolacca americana* and *Cayratia japonica* were found along the coastal seawall forest. The microhabitats (forest gap or understory) which the seedlings located were also recorded.

Bird feeding behaviors

Behaviors of birds feeding fruits of the two plant species in the forest was observed from September to October in 2018 and 2019. Focal individual sampling was used to quantify visitation and fruit removal by birds at focal stands of each study species. Because most bird foraging occurs at dusk and dawn, observations were carried from 6 am. to 9 am., or from 4 pm. to 6 pm in the sunny day. Each observation continued until the focal bird disappeared. We counted the number of fruits swallowed, dropped, pecked and taken in flight by each bird species and the lasting foraging time in every observation. The first stopping distance of the focal bird was measured by a laser range finder. We calculated the whole feeding frequency number of each bird on each plant species in the study period. The average feeding duration per time on each plant species by birds were also recorded (Li et al., 2019).

Seed germination trials

The cumulative percentage germination success of defecated seeds by local frugivorous birds was compared with that of hand-cleaned seeds and intact fruits. All of seeds and fruits were collected in October 2018 in the study site. Defecated seeds from each individual bird, hand-cleaned seeds and the intact fruits were planted in separate bowls with a diameter of 10 cm. All bowls were then placed in a laboratory balcony and water daily in the first month. Seeds were considered germinated when the seedlings emerged through the soil surface. Seed germinated number was recorded at least once every 15 days, from early June to middle August. Once counted, seedlings were removed from the tray.

Data analysis

We compared the fruit height and fruit size of each plant species using *t*-test. *T*-test was also used to compare the density in older and younger age of stand of each plant species and to compare the microhabitats selecting of seedlings. We used a generalized linear model to test the correlation between fruit ripening rate and fruit missing rate each species. Total feeding frequency of each bird species on each plant species, averaged feeding duration per time, feeding fruit amount per time and first stopping distance of focal bird were compared by *U*-test or *t*-test. ANOVA tests were used to compare germination success of defected seeds to hand-cleaned

seeds and intact fruits. All analyses were performed using SPSS 26.0 software, and Origin75 software. The level of statistical significance was set at $P < 0.05$.

Results

Fruit characteristics of plant species

In autumn, *Phytolacca americana* and *Cayratia japonica* were the most common fruiting species in the coastal seawall forest. Both fruit types were berry. When was ripe, the fruits' color changed to black or purple black. There was a significant difference in the fruit size between the species (t -test, $t = 8.252$, $P < 0.01$, $n = 40$) (Table 1).

Phytolacca americana mainly grew in the forest gap habitat, while *Cayratia japonica* was widely distributed in the understory along the seawall forest. The fruit ripening period of the two plant species basically overlapped, both ripening from late August to late October. There was a significant difference in the height of fruits distribution between the two species (U -test, $Z = -5.420$, $P < 0.05$, $n = 40$). The tree density of *Phytolacca americana* was higher in older age of stand, while the tree density of *Cayratia japonica* was higher in younger age of stand.

Fruits feeding and Seed dispersal by frugivorous birds

As time going on, the fruits of the two plant species gradually ripen, and fruit color changed from green to black or purple black. Ripen fruits were constantly eaten by frugivores, and in study period, no other animal groups except the birds were found to eat the fruits of *Phytolacca americana* or *Cayratia japonica*. The relationship between percent fully ripen fruits and the percent of fruit missing caused by birds feeding was a significant positive correlation for both plant species (General Line Model, $P < 0.01$) (Fig. 1 A & B). It showed that the more ripen fruits, the greater proportion of missing fruit eaten by birds.

11 bird species were observed feeding on the fruits of *Phytolacca americana* or *Cayratia japonica* in study period. They are Light-vented Bulbul, Common Blackbird, Black-billed Magpie, White-cheeked Starling *Sturnus cineraceus*, Black-napped Oriole *Oriolus chinensis*, Grey-backed Thrush *Turdus hortulorum*, Vinous-throated Parrotbill *Paradoxornis webbianus*, Great Tit *Parus major*, Blue-and-white Flycatcher *Cyanoptila cyanomelana*, Black-tailed Hawfinch *Eophona migratoria* and Black-faced Bunting *Emberiza spodocephala*. 7 of all bird species feed on both the fruits of the two plants. However, 3 species, the Blue-and-white Flycatcher, Vinous-throated Parrotbill, and Black-faced Bunting feed only on the fruits of *Phytolacca americana* (Fig.2).

Light-vented Bulbul and Black-billed Magpie had the most feeding frequency on *Phytolacca americana* and *Cayratia japonica*, respectively. The longest foraging time on the fruits of *Phytolacca americana* and *Cayratia japonica* were Light-vented Bulbul and White-cheeked Starling, respectively. White-cheeked Starling was the species whose first stopping distance longest after feeding on both plants (Table 2). But there were no significant differences in bird feeding frequency (U -test, $P > 0.05$), feeding duration (t -test, $df = 16$, $P > 0.05$) and first stopping distance (t -test, $t = 1.401$, $P > 0.05$) between *Phytolacca americana* and *Cayratia japonica*.

However, the bird feeding amount each time on *Phytolacca americana* significantly larger than on *Cayratia japonica* (t -test, $df=16$, $P<0.01$) (Table 2).

Seedlings microhabitat

300 *Phytolacca americana* seedlings and 280 *Cayratia japonica* seedlings were found around 15 mother trees respectively in the study period. There were differences in microhabitat characteristics between the two plant species. Seedlings of *Phytolacca americana* preferred forest gap habitat, and much more seedlings were found in the forest gap habitat than in the understory habitat (t -test, $t=5.684$, $P<0.01$) (Fig. 3, A). However, seedlings of *Cayratia japonica* did not show obvious preference for microhabitat selection (t -test, $t=0.092$, $P>0.05$) (Fig. 3, B).

Seed germination rate

Seed germination experiments were carried from 20th, June to 20th, August. Seed germination rate of intact fruits was lower than both hand-cleaned seeds and defected seeds of both plant species. The final seed germination rates of *Phytolacca americana* with three different controls were as follows: 79% for defected seeds, 60% for hand-cleaned seeds and 13% for intact fruits (Fig. 4, A). And the final seed germination rates of *Cayratia japonica* were 87%, 71% and 18%, respectively (Fig. 4, B). There was extremely significant difference of seed germination rates with different controls between *Phytolacca americana* (ANOVA: $F_{2,15}=13.283$, $P<0.01$) and *Cayratia japonica* (ANOVA: $F_{2,15}=4.591$, $P<0.05$). The results showed that local frugivorous birds' digestion could promote seed germination success in both plant species.

Discussion

Similarity of the fruits characters of the two plant species

Bird-mediated seed dispersal is important for the establishment of many fleshy-fruited plant species (Wenny, 2001). Many alien plants with fleshy fruits are able to establish effective dispersal interactions with local birds, based on the attracting fruit characters to native plants (Gosper, Stansbury & Vivian-Smith, 2005). Our study showed that the fruit characteristics e.g. fruit type, fruit shape, ripen fruit color, fruit ripening period of alien *Phytolacca americana* were much similar to native *Cayratia japonica* (Table 1). This similarity indicates that the alien species *Phytolacca americana* have great potentiality to disperse its seeds by local frugivorous birds, which have established a positive reaction with the native plant species *Cayratia japonica*.

Some studies have found that small fruits and seeds appear to be removed and dispersed more effectively than larger ones (Rey et al., 1997), and birds prefer visiting trees with larger crop size (Sallabanks, 1993; Korine, Kalko & Herre, 2000). A study in a coastal belt forest in South Africa, which found that fleshy-fruited invasion plant species effectively dispersed by birds were small-seed, open habitat species with longer fruiting length than native plant species (Bitani et al., 2020). If the former conclusion is true, and based on the fact that the fruit size of *Phytolacca americana* was smaller but containing more seeds than *Cayratia japonica*, the alien plant species

Phytolacca americana may be removed and dispersed more effectively by local frugivorous birds.

The forest gap microhabitat formed by light disturbance were more suitable for the growth of alien plant species (Williamson & Fitter, 1996; Ridenour & Callaway, 2001). Our study showed that the density of *Phytolacca americana* was higher in older age of stand in the forest, where more forest gap formed by typhoon disturbance every summer in the study site. The result suggests that the alien plant species *Phytolacca americana* prefers forest gap microhabitat in the study site.

Similar seed dispersal system

A fruit species can be consumed by a variety of potential bird dispersers, meanwhile, a potential bird disperser can remove a variety of fruit species (Dugger et al., 2018; Schneiberg et al., 2020). 7 of 11 local frugivorous birds consumed fruits of both the plant species in our study, which suggested that a stable relationship have already been established by the alien plant and local birds. Besides the 7 species, other 3 bird species were found only consume the alien species, showing that it had a wider niche to attracting more local bird species, according to the study by Levey, Gosper and Whelan (Whelan & Willson, 1994; Levey & Martinez del Rio, 2001).

Pycnonotus sinensis was a small and common frugivorous bird species in the study site according to the field observation. In autumn, the ripening fruits were the main food resource to them in the coastal seawall forest, where is a lack of insect resources. We usually observed 3-5 individuals of *Pycnonotus sinensis* foraging on the trees of *Phytolacca americana* with a large crop sizes and purple black ripen fruits color. Therefore, the *Pycnonotus sinensis* had a highest intensity interaction with the alien species.

There was no significant difference in bird feed frequency, feed duration, and first stop distance between the alien and native plant species. The results showed that there was no obvious "preference" in the selection and utilization of the seeds of these two plants by local birds, which reflected that the simulation degree of introduced plant species *Phytolacca americana* to native species *Cayratia japonica* was highly similar. But the higher efficiency of seed dispersal of *Phytolacca americana* by birds was showed according to the higher feed amount each time. We found that the frugivores feeding on the fruits were most of resident birds. The contribution of migratory birds to seed dispersal were small according to the lower feed frequency, shorter feed duration of the three migratory species, Black-napped Oriole, Grey-backed Thrush and Blue-and-white Flycatcher. So, the resident birds were the main contributors to seed dispersal of *Phytolacca americana* and *Cayratia japonica*. This might be due to the fact that the birds migrating through the seawall forest in the study site in autumn were mostly insectivorous flycatchers and warblers (Liu et al., 2020).

Effects of birds on seed germination rate

Birds' ingestion may increase seed germination rates of different plant species, especially the seeds of many invasive plant species have been shown to benefit from ingestion by frugivorous birds (Combs et al., 2011; Carrion-Tacuri et al., 2012; Czarnecka, Orlowski & Karg, 2012; Thabethe et al., 2015).

Our result showed that the seed germination rates of both plant species were lower from intact fruits than from ingested and manually de-pulped seeds (Fig.4), which consist to the results of Barnea, Yom-Tov & Friedman (1990), Yagihashi, Hayashida & Miyamoto (1999) and Traveset (1998). This suggested that local frugivorous birds' ingestion similarly and positively affected germination success of alien plant and native plant species in the study site. The local birds were legitimate seed dispersal agents of the fleshy-fruited alien plant. The factors of higher seed germination rate of ingested and manually de-pulped seeds may attribute to the removal of pulp or appropriate seed coat abrasion, but more trials were necessary to prove this conclusion. The final cumulative percentage germination of native plant species was much similar to alien plant species under three different controls (ingested seeds: 87% VS 79%; manually de-pulp seeds: 71% VS 60%; intact fruits: 18% VS 13%). This suggested that the seed dispersal pattern in seed germination of alien plant species *Phytolacca americana* was similar to native plant species *Cayratia japonica*.

We also found that seedlings of the alien plant species tended to occupy the microhabitat of forest gap which was suitable for seedling growth. But there was no preference tendency in microhabitat selecting by native plant species seedlings. This showed that seedlings of alien plant species had good developmental potential to successful population regeneration in the coastal seawall forest.

Conclusions

The alien species *Phytolacca americana* has a much similar fruit characteristics with the native plant species *Cayratia japonica* in autumn in the coastal seawall forest. A seed dispersal system between the alien plant species and local frugivorous birds has been established, and it is similar to the native one. Seeds of *Phytolacca americana* after ingestion by local frugivorous birds could promote the germination rate, and its seedling was more likely to occupy the microhabitat of forest gap than *Cayratia japonica* seedlings. All of these eventually enabled the alien plant species *Phytolacca americana* to stay and rapidly expand in the coastal seawall forest in our study site.

Acknowledgements

We are grateful to Guohai Wang for helping us in processing the data. We also thank the staffs of the Dafeng Milu National Nature Reserve for their contributions in the field.

References

- Aslan CE. 2011. Implications of newly-formed seed-dispersal mutualisms between birds and introduced plants in northern California, USA. *Biological Invasions*, 13:2829–2845.
- Barnea A, Yom-Tov Y, Friedman J. 1990. Differential germination of two closely related species of *Solanum* in response to bird ingestion. *Oikos*, 57: 222-228.
- Bitani N, Ehlers Smith DA, Ehlers Smith YC, et al. 2020. Functional traits vary among fleshy-fruited invasive plant species and their potential avian dispersers. *Acta Oecologica*, DOI: 10.1016/j.actao.2020.103651.
- Butler DW, Green RJ, Lamb D, et al. 2007. Biogeography of seed-dispersal syndromes, life forms and seed sizes among woody rain forest plants in Australia's subtropics. *Journal of Biogeography*, 34(10): 1736-1750.
- Buckley YM, Anderson S, Catterall CP, et al. 2006. Management of plant invasions mediated by frugivore interactions. *Journal of Applied Ecology*, 43: 848-857.
- Camargo P, Rodrigues S, Piratelli AJ, et al. Interhabitat variation in diplochory: Seed dispersal effectiveness by birds. *Perspectives in Plant Ecology, Evolution and Systematics*, 2019, 38: 48-57.
- Carrion-Tacuri J, Berjano R, Guerrero G, et al. 2012. Predation on seeds of invasion *Lantana camara* by Darwin' finches in Galapagos Islands. *Wilson Journal Ornithology*, 124: 338-344.
- Combs JK, Reichard SH, Groom MJ, et al. 2011. Invasive competitor and native seed predators contribute to rarity of the narrow endemic *Astragalus sinuatus* Piper. *Ecological Applications*, 21: 2498-2509.
- Czarnecka J, Orlowski G, Karg J. 2012. Endozoochorous dispersal of alien and native plants by two palearctic avian frugivores with special emphasis on invasive giant goldenrod *Solidago gigantea*. *Central European Journal of Biology*, 7: 895-901.
- Drummond BA. 2005. The selection of native and invasive plants by frugivorous birds in Maine. *Northeast Naturalist*, 12: 33–44.
- Duan Q, Goodale E, Quan RC. 2013. The effect of color on fruit selection in six Tropical Asian birds. *The Condor*, 115(3): 623–629.
- Dugger PJ, Blendinger PG, Böhning-Gaese K, et al. 2018. Seed-dispersal networks are more specialized in the Neotropics than in the Afrotropics. *Global Ecology and Biogeography*, 28(2): 248-261.
- Galindo-González J, Guevara S, Sosa VJ. Bat-and bird-generated seed rains at isolated trees in pastures in a tropical rainforest. *Conservation Biology*, 2000, 14(6): 1693-1703.
- Gosper CR, Stansbury CD, Vivian-Smith G. 2005. Seed dispersal of fleshy-fruited invasive plants by birds: contributing factors and management options. *Diversity and Distributions*, 11:549-558.
- Kawakami K, Mizusawa L, Higuchi H. 2009. Re-established mutualism in a seed-dispersal system consisting of native and introduced birds and plants on the Bonin Islands, Japan. *Ecological Research*, 24: 741-748.

- 353 Kollmann J, Frederiksen L, Vestergaard P, et al. 2007. Limiting factors for seedling emergence
354 and establishment of the invasive non-native *Rosa rugosa* in a coastal dune system. *Biol*
355 *Invasions*, 9: 31-42.
- 356 Korine C, Kalko EKV, Herre EA. 2000. Fruit characteristics and factors affecting fruit removal
357 in a Panamanian community of strangler figs. *Oecologia*, 123:560-568.
- 358 Leishman MR, Wright IJ, Moles AT, et al. 2000. The evolutionary ecology of seed size. In:
359 Fenner M (ed) *Seeds: the ecology of regeneration in plant communities*, 2nd edn. CABI
360 Publishing, Wallingford, pp 31–57.
- 361 Levey DJ, Martínez del Rio C. 2001. It takes guts (and more) to eat fruit: lessons from avian
362 nutritional ecology. *Auk*, 118:819-831.
- 363 Li N, Wang Z, Zhang S, et al. 2019. Importance of bird traits for seed dispersal patterns of co-
364 fruiting trees in a patchy forest. *Integrative Zoology*, 14: 470-478.
- 365 Liu B, Xu P, Xue DD, et al. 2020. Nest site selection and niche differentiation of *Eophona*
366 *migratoria* and *Dicrurus macrocercus* in Yancheng Coastal Seawall Forest. *Chinese Journal*
367 *of Ecology*, 39(1): 186-193.
- 368 Lovas-Kiss A, Vincze O, Kleyheeg E, et al. 2020. Seed mass, hardness, and phylogeny explain
369 the potential for endozoochory by granivorous waterbirds. *Ecology and Evolution*, 10:1413-
370 1424.
- 371 Lu C. 2003. Biology of mistletoe (*Viscum coloratum*) and its seed dispersal frugivorous birds.
372 *Acta Ecologica Sinica*, 23(4): 834-839.
- 373 McDonnell MJ, Cheplick G, Armesto JJ, et al. 1984. Bird-dispersal of *Phytolacca americana* L.
374 and the influence of fruit removal on subsequent fruit development. *American Journal of*
375 *Botany*, 71(7): 895-901.
- 376 Moles AT, Acherly DD, Webb CO, et al. 2005. A brief history of seed size. *Science*, 307: 576-
377 580.
- 378 Morán-López T, González-Castro A, Morales JM, et al. Behavioural complementarity among
379 frugivorous birds and lizards can promote plant diversity in island ecosystems. *Functional*
380 *Ecology*, 2020, 34:182-193.
- 381 Prado FA. 2013. Feeding behavior, bird visitation, and seed dispersal in *guarea macrophylla* and
382 *trichilia quadrijuga*. *Ornitologia Neotropical*, 24: 459-468.
- 383 Renne IJ, Barrow WC Jr, Johnson Randall LA, et al. 2002. Generalized avian dispersal
384 syndrome contributes to Chinese tallow tree (*Sapium sebiferum*, Euphorbiaceae)
385 invasiveness. *Diversity and Distributions*, 8:285-295.
- 386 Rey PJ, Gutiérrez JE, Alcántara J, et al. 1997. Fruit size in wild olives: implications for avian
387 seed dispersal. *Functional Ecology*, 11:611-618.
- 388 Ridenour WM, Callaway RM. 2001. The relative importance of allelopathy in interference: the
389 effects of an invasive weed on a native bunchgrass. *Oecologia*, 126 :444-450.
- 390 Saavedra F, Hensen I, Beck SG, et al. 2014. Functional importance of avian seed dispersers
391 changes in response to human induced forest edges in a tropical seed-dispersal networks.
392 *Oecologia*, 176: 837-848.

- Sallabanks R. 1993. Hierarchical mechanisms of fruit selection by an avian frugivore. *Ecology*, 74: 1326-1336.
- Schierenbeck KA. 2004. Japanese Honeysuckle (*Lonicera japonica*) as an invasive species; History, ecology, and context. *Critical Reviews in Plant Sciences*, 23(5): 391-400.
- Schleuning M, Blüthgen N, Florchinger M, et al. 2011. Specialization and interaction strength in a tropical plant-frugivore network differ among forest strata. *Ecology*, 92(1): 26–36.
- Schneiberg I, Boscolo D, Devoto M, et al. 2020. Urbanization homogenizes the interactions of plant-frugivore bird networks. *Urban Ecosystems*, 23: 457-470.
- Traveset A. 1998. Effect of seed passage through vertebrate frugivores' guts on germination: a review. *Perspectives in Plant Ecology, Evolution and Systematics*, 1:151-190.
- Thabethe V, Wilson A, Hart LA, et al. 2015. Ingestion by an invasive parakeet species reduces germination success of invasive alien plants relative to ingestion by indigenous turaco species in South Africa. *Biological Invasions*, 17: 3029-3039.
- Wang B, Ives AR. 2017. Tree-to-tree variation in seed size and its consequences for seed dispersal versus predation by rodents. *Oecologia*, 183: 751-762.
- Wenny DG. 2001. Advantages of seed dispersal: a re-evaluation of directed dispersal. *Evolutionary Ecology Research*, 3:51-74.
- Whelan CJ, Willson MF. 1994. Fruit choice in migrating North American birds: field and aviary experiments. *Oikos*, 71:137-151.
- Willamson M, Fitter A. 1996. The varying success of invaders. *Ecology*, 77: 1661-1666.
- Wilson AL, Downs CT. 2012. Knysna Turacos (*Tauraco corythaix*) do not improve seed germination of ingested fruit of some indigenous South African tree species. *South African Journal Botany*, 78:55-62.
- Wu JX, Delparte DM, Hart PJ. 2014. Movement patterns of a native and non-native frugivore in Hawaii and implications for seed dispersal. *Biotropica*, 46(2): 175-182.
- Yagihashi T, Hayashida M, Miyamoto T. 1999. Effects of bird ingestion on germination of two *Prunus* species with different fruit-ripening seasons. *Ecological Research*, 14(1):71-76.

Table 1(on next page)

Fruit characteristics of the two plant species and their distribution in Yancheng seawall forest.

1

Fruit characteristics	<i>Cayratia japonica</i>	<i>Phytolacca americana</i>
Type	Berry	Berry
Size (mm)	10.0±0.9	7.9±0.7
Shape and Color	Round, black	Round, Purple black
Ripening period	August to October	August to October
Fruit height (m)	0.2±0.1	1.3±0.3
Tree density (individuals/m ²)	0.5±0.3	0.1±0.1

2

Table 2 (on next page)

List of bird species consuming fruits of native and alien plant species, followed by foraging statistics.

CY: *Cayratia japonica*; PA: *Phytolacca americana* Feeding pattern: S: Swallow; P: Pecking
Seasonal type: R: Resident birds; T: Travel birds; W: Winter birds

Bird species	Feeding frequency (times)		Feeding time (s)		Feeding amount (ind.)		First stopping distance(m)		Feeding pattern	Seasonal type
	CY	PA	CY	PA	CY	PA	CY	PA		
<i>Pycnonotus sinensis</i>	4.0	65.0	10.0	14.2	1.5	4.1	9.3	13.4	S	R
<i>Turdus merula</i>	3.0	18.0	6.7	7.3	1.3	3.7	25.0	28.6	S	R
<i>Cyanoptila cyanomelana</i>	0	11.0	0	5.2	0	1.8	0	8.3	S	T
<i>Sturnus cineraceus</i>	11.0	9.0	26.9	10.1	1.9	4.6	75.0	70.0	S	R
<i>Pica pica</i>	35.0	9.0	16.7	10.0	2.1	4.2	29.9	23.3	S	R
<i>Oriolus chinensis</i>	12.0	0	11.9	0	2.6	0	31.9	0	S	T
<i>Turdus hortulorum</i>	6.0	4.0	14.5	7.8	1.8	2.8	42.5	19.3	S	T
<i>Paradoxornis webbianus</i>	0	17.0	0	5.8	0	3.6	0	6.7	S/P	R
<i>Parus major</i>	1.0	13.0	5.0	6.8	1.0	2.2	12.0	12.2	S/P	R
<i>Eophona migratoria</i>	13.0	4.0	20.8	10.0	1.7	4.5	26.0	15.0	S/P	R
<i>Emberiza spodocephala</i>	0	13.0	0	11.6	0	2.0	0	7.2	S/P	W

- 1 CY: *Cayratia japonica*; PA: *Phytolacca americana*
- 2 Feeding pattern: S: Swallow; P: Pecking
- 3 Seasonal type: R: Resident birds; T: Travel birds; W: Winter birds
- 4

Figure 1

The relationship between percent fully ripen fruits and percent of fruit missing caused by bird foraging.

(A) *Phytolacca americana*.




Figure 2

The relationship between percent fully ripen fruits and percent of fruit missing caused by bird foraging.

(B) *Cayratia japonica*




Figure 3

Correspondence relationship between avian frugivores and fruiting plants based on feeding behaviors.

Widths of connecting lines denote the relative number of observed interactions (wider represents higher intensity interaction) Avian frugivores: a. *Pycnonotus sinensis*; b. *Pica pica*; c. *Turdus merula*; d. *Sturnus cineraceus*; e. *Eophona migratoria*; f. *Paradoxornis webbianus*; g. *Parus major*; h. *Emberiza spodocephala*; i. *Oriolus chinensis*; j. *Cyanoptila cyanomelana*; k. *Turdus hortulorum*




Figure 4

The microhabitat selecting of seedlings of the two plant species.

(A) *Phytolacca americana*




Figure 5

The microhabitat selecting of seedlings of the two plant species.

(B)*Cayratia japonica*




Figure 6

Comparison of the seed germination rates of the two plant species with different controls.

(A) *Phytolacca americana*




Figure 7

Comparison of the seed germination rates of the two plant species with different controls.

(B) *Cayratia japonica*

