

Between darkness and light: spring habitats provide new perspectives for modern researchers on groundwater biology

Raoul Manenti ^{Corresp., 1, 2}, **Beatrice Piazza** ¹

¹ Department of Environmental Science and Policy, Università degli Studi di Milano, Milano, Lombardia, Italy

² Parco Regionale del Monte Barro, Laboratorio di Biologia Sotterranea "Enrico Pezzoli", Galbiate, Italy

Corresponding Author: Raoul Manenti
Email address: raoul.manenti@unimi.it

Springs are interfaces between groundwater and surface habitats and may play an important role in the study of subterranean animals. In this systematic evidence review and meta-analysis, we explore whether observations of stygobionts in springs are relevant and more common than observations of epigean animals in groundwater. We searched the Web of Science database for papers on groundwater fauna and spring fauna. For each paper we found, we recorded whether the paper reported the occurrence of typical stygobionts in springs, of surface animals in groundwater, or of the same taxa in both habitats. If so, we recorded how many such species were reported. We also recorded the scientific discipline of each study and the year of publication. Our search yielded 342 papers. A considerable number of these papers reported stygobionts in springs: 20% of papers dealing with groundwater fauna and 16% of papers dealing with spring fauna reported the occurrence of stygobionts in spring habitats. Both the number of papers that mentioned stygobionts in springs, and the number of stygobiont species that were documented in springs, were higher than equivalent measures for the occurrence of surface fauna underground. We also detected a positive relationship between year of publication and the number of reports of stygofauna in springs. To broaden the insights from biological research on underground environments, we suggest that springs should be considered not only as simple sampling points of stygobionts but also as core stygobiont habitats.

Between darkness and light: spring habitats provide new perspectives for modern researchers on groundwater biology

Raoul Manenti^{1,2}, Beatrice Piazza¹

¹ 1Department of Environmental Science and Policy, Università degli Studi di Milano, via Celoria, 26, 20133 Milano Italy

² 2 Laboratorio di Biologia Sotterranea “Enrico Pezzoli”, Parco Regionale del Monte Barro, Galbiate, Italy.

Corresponding Author:

Raoul Manenti¹

via Celoria, 26, 20133 Milano Italy

Email address: raoul.manenti@unimi.it

Abstract

Springs are interfaces between groundwater and surface habitats and may play an important role in the study of subterranean animals. In this systematic evidence review, we explore whether observations of stygobionts in springs are relevant and more common than observations of epigean animals in groundwater.

We searched the Web of Science database for papers on groundwater fauna and spring fauna. For each paper we found, we recorded whether the paper reported the occurrence of typical stygobionts in springs, of surface animals in groundwater, or of the same taxa in both habitats. If so, we recorded how many such species were reported. We also recorded the scientific discipline of each study and the year of publication.

Our search yielded 342 papers. A considerable number of these papers reported stygobionts in springs: 20% of papers dealing with groundwater fauna and 16% of papers dealing with spring fauna reported the occurrence of stygobionts in spring habitats. Both the number of papers that mentioned stygobionts in springs, and the number of stygobiont species that were documented in springs, were higher than equivalent measures for the occurrence of surface fauna underground. We also detected a positive relationship between year of publication and the number of reports of stygofauna in springs.

To broaden the insights from biological research on underground environments, we suggest that springs should be considered not only as simple sampling points of stygobionts but also as core stygobiont habitats.

Introduction

The zoologist Lazăr Botoșăneanu (1998) defined springs as the “doors on River Styx,” the river of the Greek mythological underworld. Other biologists who study subterranean environments and groundwaters similarly consider springs to be openings that allow them to see the inhabitants of an otherwise inaccessible environment (Culver et al. 2012; Fiasca et al. 2014; Galassi et al. 2014; Pipan & Culver 2012; Pipan et al. 2012). This view of spring habitats as windows into a different environment is particularly true in non-karst areas, where the lack of caves prevents human exploration of the subterranean realm and springs are often the only way to access groundwater organisms (Manenti & Pezzoli 2019). Springs are interfaces between groundwater and surface freshwaters with a great variety of connectivity, permeability and biodiversity (Gibert et al. 1997; Kresic 2010; Springer & Stevens 2009). The interplay of both subterranean and epigeal habitat features that characterises springs (Alfaro & Wallace 1994; Cantonati et al. 2006), makes challenging to define the spatial boundaries of these ecotones (Gibert et al. 1997).

In some cases the transition between surface and subterranean habitats, may also be abrupt; indeed, the magnitude of this transition strongly depends on the morphology of the spring and can be mutable with daylight. Some springs represent an abrupt shift from the subterranean environment to the surface, whereas others, like the natural emitting caves (such as caves from which subterranean streams flow outside) of artificial draining galleries (such as galleries built to collect groundwater), represent extended ecotonal environments (Balland 1992; White 2019). The border between the subterranean and surface environment can be particularly distinct during daytime,

when it is strictly demarcated by the sun. Aside from sunlight, the differences that distinguish subterranean and surface environments in a spring, even across the few meters or centimetres that may characterize a spring with a sudden interface (Fig. 1), include the availability of trophic resources, the density of potential predators, and microclimate conditions (Barzaghi et al. 2017; MacAvoy et al. 2016; Manenti et al. 2013; Von Fumetti & Nagel 2011).

Because springs are border habitats, it can be difficult for biological studies to consider springs in their entirety; this difficulty has limited the potential for insights from springs to drive stronger advances in different fields of research. For example, studies that focus on springs often only consider a surface perspective and neglect the role played by groundwater (Manenti & Pezzoli 2019), whereas in karst areas, scientists studying the subterranean environment see springs as “access points” that can be used to sample the groundwater fauna living in different subterranean, underwater environments, such as the phreatic zone of karst aquifers (Malard et al. 2002). This latter view reflects the scarce consideration that is often given to springs and may limit a more general understanding of the ecological role of border habitats. As some studies have already suggested, transition zones are important for regulating ecosystem processes and the flow of organic matter and organisms between surface (epigean) and underground (hypogean) habitats (Moseley 2009; Plenet & Gibert 1995; Prous et al. 2004). With this opinion paper based on a systematic review of the recent scientific literature, we aim to stimulate a change in the conception of and in the approach to springs by studies dealing with stygobionts and groundwater fauna. Particularly we want to underline that springs have the potential to reveal general patterns related to the zoology of stygobionts.

Stygobionts are obligate groundwater-dwellers; the etymology of the word “stygobiont” reflects the fact that these species, and stygofauna more broadly, are “of the River Styx.” These organisms

have evolved adaptations specific to the underground freshwater habitats in which they spend their entire life cycle (Trajano & De Carvalho 2017). Stygobionts often exhibit morphological features associated with their underground habitat. These characteristics, such as blindness and depigmentation, are commonly referred to as troglomorphisms (Pipan & Culver 2012; Romero 2009), and they limit stygobionts' ability to exploit surface environments. However, at night, the constraints are generally less clear and surface borders (i.e., springs) may become more permeable by stygofauna; for example, some samplings of springs' fauna have reported during night the occurrence of organisms considered to be strict stygobionts (Bressi et al. 1999; Fišer 2019; Manenti & Barzaghi 2020; Manenti & Barzaghi In press), as it has been observed also for the twilight zones of terrestrial caves (Mammola & Isaia 2018). One such case is that of *Stygobromus* amphipods, which are believed to regularly leave hypotelminorheic habitats to feed (Culver et al. 2006; Culver & Pipan 2014). Nevertheless, these findings are often viewed as exceptions or accidental events, and the use of springs is rarely mentioned as a trait of stygobiont biology. Observations of surface animals in caves have been similarly overlooked in the past (Sket 2008) and improperly seen as accidental; such "accidental" observations have recently been described for both groundwater and terrestrial subterranean habitats (Ficetola et al. 2018; Lunghi et al. 2014a; Manenti 2014).

Stygobionts are the main focus of subterranean biology and are usually studied using two distinct approaches. The first approach includes intense taxonomic investigations focused on the discovery and description of new taxa. The second approach views caves as powerful natural laboratories for evolutionary, ecological and behavioural studies on their inhabitants (Culver & Pipan 2014; Culver & Pipan 2019). The idea of caves as natural laboratories, first postulated by the speleologist Édouard-Alfred Martel (1894), has been espoused for more than one hundred years of subterranean

studies (Poulson & White 1969) and recently updated and broadened in the current context of biological research (Mammola 2019; Mammola et al. 2020) . However, of the relatively large number of caves that were effectively used as laboratories during last century (Vandel 1964), few remain active. In addition, the outcomes of studies on stygofauna in these caves is rarely compared to insights obtained from studies of surface freshwater organisms. This is partially due to the characteristic features of stygobionts: they are often difficult to sample in deep subterranean environments and springs can be seen as useful sampling points for them, using a combination of different methods depending on the spring morphology and hydrology (Malard et al. 2002). Particularly in springs are recommended the sampling of drifting stygobionts during spates, samplings of the spring benthic layer, and bed sediments and also the use of artificial substrates and baited traps (Malard et al. 2002).

Moreover, due to their long life cycles and low fertility, stygobionts are difficult to raise in an experimental settings. Would including springs and other surface/underground border habitats in studies on subterranean biology increase the understanding of how the constraints of the hypogean environment affect the phenotypic responses and genetic makeup of stygobionts? The rationale of this paper takes origin from this question and is to suggest that a substantial inclusion of springs (and other border habitats between underground and surface) in studies on subterranean biology, can increase the understanding of principles governing exploitation and adaptation to hypogean environments.

In this paper, we investigate the perspectives of modern researchers on considering springs not only as simple sampling points, but also as core stygobiont habitats that can broaden the insights obtained from biological studies of underground environments. We specifically performed a systematic review of the recent scientific literature to understand i) the relevance of previous

observations of typical stygobionts in springs; ii) if these observations vary according with study discipline and the year of publication; and iii) if these observations are more common than observations of epigean animals (i.e. aquatic surface species) in caves. By demonstrating that typical stygofauna are observed in springs more commonly than usually thought, we propose that, at least in some cases, the exploitation of border habitats be considered a non-negligible aspect of stygofauna ecology.

Survey methodology

To avoid bias, many scientific fields have largely started to favour the use of systematic evidence reviews (Acreman et al. 2020). We therefore performed a systematic review to find focused data that addressed our three aims (Table 1). For this review, we followed the Preferred Reporting Items for Systematic Reviews (PRISMA) guidelines (Page & Moher 2017), and we searched the Web of Science database for peer-reviewed papers on both stygofauna and fauna living in spring habitats (Fig. 2). The Web of Science database contains metadata for peer-reviewed scientific articles published since 1965. We used two search strings designed to find all articles in the database that might contain observations of fauna in both caves and springs. Our search was conducted in May 2020 from Milano, Italy, using the keywords “groundwater fauna” (GF) and “spring fauna freshwater” (SFF) and searching them by topics. For the search we used a ASUS K501 PC and the Google Chrome browser, after having emptied its cache box.

One of us (BP) initially screened the articles that met our search criteria by discarding articles that were not clearly related to our study aims. She rejected articles about botany, palaeontology, geology, and all their associated subdisciplines (paleoecology, stratigraphy, geomorphology, etc.), as well as articles about subterranean environments or groundwater that did not mention animals. The articles found using the key words “spring fauna freshwater” were more difficult to screen; for the most part, the authors of these articles did not specify if their study species were part of

stygo fauna or not. She therefore discarded these articles only if they were not related to our study aims (e.g., papers about estuaries, palaeontology, or related topics) or if the authors provided clear evidence that the study species were not cave-dwellers.

She additionally discarded several articles that dealt strictly with agricultural sciences, biogeochemical cycles, the impacts of various pollutants (crude oils, perchlorate, etc.) on groundwater, or other environments strongly connected to groundwater (i.e., all surface water environments), but did not mention the finding of stygo fauna or epigean fauna. She discarded also articles that addressed single species or taxa that are not stygo fauna or typical spring fauna and have no hypogean representatives (e.g., Rechulicz 2011 treated *Pseudorasbora parva* and Vilenica et al. 2016 treated mayflies). Articles concerning terrestrial environments, estuaries, swamps, mangroves, streams, rivers, lakes, and all saltwater environments, were similarly discarded (see Table 1 for more detailed information on the article selection procedure). After this first screening, she performed a second selection procedure in which she removed any articles that were unavailable or were written in a language other than English.

From the papers she collected the information listed in Table 1, including the typology of the study, distinguishing between ecology, taxonomy behaviour, conservation and fauna assessment and considering that the same paper could belong to multiple categories. Moreover, she assessed the country and the continent of origin of the data provided by the papers.

Statistical analyses

To assess the relationships between features of the selected documents and the occurrence of stygo fauna in springs, we built a series of generalized linear models (GLMs) with binomial error distributions.

First, we assessed if the fact that a paper reported the occurrence of stygobionts in springs, the occurrence of surface fauna in groundwaters, or the contemporary occurrence of a stygobiont in both groundwaters and springs, was related to the paper's field of study. Reported occurrences were used as the dependent variable, and the study disciplines (ecology, taxonomy, faunal assessment, and conservation) were used as fixed factors (Equation 1). We similarly built GLMs with the same dependent variable but with publication year and the search term as independent variables (Equation 2).

Stygofauna found in springs OR Surface fauna underground OR

Occurrence of the same taxa in Same taxa both in springs and groundwaters ~ Ecology + Taxonomy + Fauna.assessment + Conservation + Behaviour, family = binomial.

(eqn 1)

Stygofauna found in springs OR Surface fauna underground OR

Occurrence of the same taxa in Same taxa both in springs and groundwaters ~

Year of publication + as.factor(key research term), family = binomial.

(eqn 2)

Second, using only the papers selected with the GF search term we built GLMs to test two distinct hypotheses: (1) that the number of mentions of stygobiont species in springs is higher than the number of mentions of surface species in groundwater; and (2) that missing information is different among the two situations. Both hypotheses were tested only for papers selected with the keyword “groundwater fauna” to avoid biases associated with the fact that studies found using the keyword “spring fauna freshwater” may not have sampled underground habitats. For the first test (Equation 3), we used the number of species mentioned by each paper as a dependent variable, including

both stygofauna found in springs and surface fauna found underground. The type of observation (stygofauna in springs vs. surface fauna in groundwater) was used as a fixed factor.

$$\text{Number of species mentioned} \sim \text{Typology of observation, family} = \text{nbinom2.}$$

(eqn 3)

For the second test (Equation 4), we defined the dependent variable as whether it was possible to assess the number of species mentioned in a study, including both types of observations (stygofauna in springs and surface fauna in groundwater). The type of observation was used as a fixed factor, as before.

$$\text{NAns occurrence} \sim \text{Typology of observation, family} = \text{nbinom2.}$$

(eqn 4)

Third, we also built a series of GLMs to test if the fact that a paper reported the occurrence of stygobionts in springs, the occurrence of surface fauna in groundwaters, or the contemporary occurrence of a stygobiont in both groundwaters and springs, was related to the continent of sampling of the springs.

For all models we checked models assumptions by verifying the absence of multicollinearity issues though VIF calculation and plotting residuals versus fitted values, versus each covariate; To avoid overdispersion bias in the models used for the second step of analysis, we built both models using a type 2 negative binomial error distribution in the package glmmTMB (Brooks et al. 2017).

We used a likelihood ratio test to assess the significance of all the fixed factors included in each GLM (Bolker et al. 2008). All analyses were performed in the R 3.6.3 environment.

224

225 **Results**

226 We retrieved 824 potentially relevant papers after removing duplicate articles. After removing
 227 articles based on the first selection criteria described above, there were 415 potentially relevant
 228 documents. After the second selection procedure, we obtained 342 papers: 275 derived from the
 229 search term “groundwater fauna” (GF) and 67 from the search term “spring fauna freshwater”
 230 (SFF). Many papers found using the “groundwater fauna” search did not specify the sampling site
 231 for the taxa considered, and many papers found using the “spring fauna freshwater” search did not
 232 clearly identify if they sampled stygofauna or not. Overall, 57 papers (representing 19% of the
 233 papers with information on sampling habitat) reported the occurrence of stygofauna in springs, 37
 234 (11.7%) reported the occurrence of typical surface fauna underground, and 33 (11%) reported the
 235 same taxa in both springs and groundwater (Table 2). With respect to our search terms, 20% of
 236 papers dealing with GF and 16% of papers dealing with SFF described the occurrence of
 237 stygobionts in spring habitats.

238 There were 45,375 mentions of species across all papers we retrieved. Of these, 138 were
 239 stygobionts observed/sampled in springs and 46 were surface species observed in subterranean
 240 habitats.

241 The study disciplines covered by the papers were mainly ecology (196 papers) and faunistic
 242 assessments (177 papers). Four papers were behavioural studies, and 24 papers addressed
 243 conservation issues. There were 194 papers that encompassed multiple ad hoc definitions that we
 244 established in the methods.

245 Our first analysis revealed that faunal assessments are significantly more likely to report the
 246 occurrence of surface fauna in groundwater, whereas taxonomic studies are more likely to report
 247 the occurrence of the same taxon in both environments (Table 3). We did not detect any

relationship between the discipline of a paper (ecology, taxonomy, etc.) and the reported occurrence of stygofauna in springs. However, we did detect a positive relationship between the year of publication and the reports of stygofauna in springs ($\chi^2=4.53$, $P=0.03$). Papers selected using the SFF search term were significantly less likely to report the occurrence of surface taxa in groundwaters ($\chi^2=4.09$, $P=0.04$).

Most data of the papers came from Europe (59.76%) and Oceania (27.9 %); we detected no significant relationships between the continent of sampling and the occurrence of stygobionts in springs, the occurrence of surface fauna in groundwaters, and the contemporary occurrence of a stygobiont in both groundwaters and springs.

GLMs performed on papers selected using the GF search term revealed that the number of mentions of stygobiont species in springs is higher than the number of mentions of surface fauna underground ($\chi^2=4.19$, $P=0.04$). However, there is also less information available on whether stygofauna have been observed in springs compared to whether surface species have been recorded in groundwaters ($\chi^2=14.08$, $P<0.001$).

Discussion and perspectives

Our systematic review revealed that there are more papers about stygofauna available on Web of Science than there are papers addressing fauna and springs. Because the word “spring” is a homograph with multiple meanings, our initial search retrieved many papers that were ultimately discarded because they did not discuss fauna and spring habitats. Preliminary literature searches performed using synonyms of “spring” and/or terms that define specific spring habitats, such as “sources” or “seepage,” resulted in fewer papers. Most of these papers were already included in our analysis; however, the few that were not could be used in future study with a larger set of papers. Someone could disagree as it is likely that our research missed some papers (as an example we are sure that we missed at least 15 papers on aquatic cave-dwelling salamanders and 10 papers

on the taxonomy of strictly subterranean planarians) and that further keywords should have been added, for example: stygob*, ecoton*, hypogean, subterranean etc, but they would have increased the number of papers dealing with stygofauna without significant increase in the number of papers related to spring fauna. The large difference in the number of papers obtained with the two search terms, GF and SFF, underscores the fact that the fauna of spring habitats have received much less attention not only than the inhabitants of lakes and streams/ivers, as already pointed out by previous studies (Cantonati et al. 2011), but also than stygofauna.

We limited our review to articles archived in Web of Science; this approach was more conservative because it included only relatively recent papers published in indexed, high-impact journals that perform selective peer review. An analogous review could be performed using the Google Scholar database or a more exhaustive search of grey literature in online and physical repositories. It is possible that the older, descriptive papers archived in these databases may have reported stygofauna in springs, but it is also possible that some form of bias could arise from using older literature that has not been rigorously peer reviewed. The effects of database selection should therefore be investigated in the future.

Using both GF and SFF as search terms, we found papers that mentioned the occurrence of stygofauna in springs, of typical surface fauna in groundwaters, and of the same taxa in both environments. The number of papers that reported stygofauna in springs, as well as the number of stygobiont species that were documented in springs, represented only a fraction of the total papers and documented species but were nevertheless non-negligible. This pattern was not linked to any specific field of study; though taxonomic studies were non-significantly more likely to report stygofauna in springs.

294 Springs have been recognized as relevant habitats for studying stygobionts since the beginning of
 295 the 19th century. Most subterranean biologists devoted at least some of their studies to spring
 296 habitats (Culver et al. 2012; Culver et al. 2014; Vandel 1920), and Albert Vandel, the founder of
 297 the “Laboratoire Souterrain de Moulis” (France), one of the most popular subterranean laboratories
 298 in the world (Botosaneanu 1980), stated in 1920 that a systematic study of spring habitats could
 299 furnish important insights for solving some of the evolutionary questions posed by cave-dwelling
 300 animals (Vandel 1920). However, this concept appears only in Vandel’s conclusions and is not
 301 further developed; the idea that springs are just sampling points in non-karst areas largely prevails
 302 throughout the rest of the paper (Vandel 1920).

303 Stygobionts are known to colonize the mixed assemblages of organisms residing in springs via
 304 emigration and drift from groundwaters (Malard et al. 2009; Malard et al. 2002; Manenti &
 305 Barzaghi In press). Typical stygobionts may be more or less permanently detected outside the
 306 spring outlet, where they can exploit different microhabitats (Malard et al. 2002; Mathieu et al.
 307 1994). This is especially true when there is a stable supply of immigrants from karst groundwater
 308 (Mathieu et al. 1999). Our results revealed that there is a positive significant correlation with the
 309 reporting of typical stygobionts in springs and the year of publication; this means that, with respect
 310 to older studies, researchers are paying more attention when reporting data on sample collection
 311 habitats, regardless of their study discipline (ecology, taxonomy, conservation, or faunal
 312 assessment). However, in the papers that we collected, the occurrence of stygobionts in springs
 313 was often reported as either an effect of the sampling method or an occasional finding. None of
 314 the papers assessed patterns in the use of springs by stygobionts. This is true also for some papers
 315 that we missed with our search but that are well known in spring literature. As an example (Rouch
 316 1986) defined “the hemorrhage” the flow of stygofauna pushed out from aquifers during high

discharge periods through springs, erroneously considering this only as a passive mechanism. Also the case of the olm (*Proteus anguinus*) is challenging with both mentions of cases of passive drift of individuals from groundwaters (Aljančič 2019) and hypotheses of active exploitations of spring habitats (Bressi et al. 1999). In more recent times, some papers were devoted to spring discharge and the passive presence of stygobionts being flushed from "conductive" or "capacitive" aquifers (Di Lorenzo et al. 2005); other large-scale ecological surveys of springs demonstrated that in mountain areas, where species richness of stygobionts is usually poor due to the effect of Quaternary Glaciations, their occurrence seems low or occasional in springs (Stoch et al. 2011), suggesting that the geographical location of springs matters and could be more in depth considered in future systematic reviews dealing with springs. Springs are also being studied with recent and 'modern' approaches like DNA metabarcoding techniques and eDNA that can allow to detect the presence of stygobionts in springs (West et al. 2020; White et al. 2020; Yonezawa et al. 2020) and be used in the future to assess the patterns that determine this occurrence.

The occurrence of a stygobiont species, or a species that is strictly linked to a hypogean groundwater habitat for its life cycle, in an epigean spring habitat, underlines a contradiction that might reflect the human conceptual limit of understanding borders. The human perception of limits and boundaries may be biased, as humans may recognize or emphasize abrupt distinctions when they do not exist (Pirni 2016; Sturz & Bodily 2016). Our results demonstrate that, at least for some stygobionts, border habitats and adjacent areas are an important part of the range and biology of stygofauna, and a proper consideration of these habitats in subterranean biology studies could provide larger perspectives. For example, stygobiont populations or individuals that exploit springs more or less permanently are likely exposed to different constraints and advantages than populations or individuals that exploit deeper aquifers. Selective pressures may therefore act

differently, at least for the individuals living in springs or at the interface between subterranean and epigeal habitats.

For example, different species and/or populations of the genus *Niphargus*, which shows typical features of stygobionts including depigmentation and the absence of eyes, have the unique ability to detect light (Fišer et al. 2016). This ability has been associated with the need to distinguish the border between surface and subterranean environments and avoid risky surface habitats (Fišer et al. 2016) where UV rays may be dangerous for a depigmented animal. However, surface habitats may also be advantageous by furnishing higher trophic resources and, at night, they are not exposed to UV light. Several studies have reported *Niphargus* amphipods in border habitats (Fiser et al. 2007; Manenti & Pezzoli 2019; Marković et al. 2018). Is light perception the same between individuals from borders and individuals from deeper aquifers? Are there evolutionary adaptations for exploiting not only deep subterranean habitats but also border habitats at the interface with the surface? These questions are applicable to all stygobionts that are recurrently found in springs. Considering border habitats in addition to deeper subterranean environments therefore has the potential to double the insights obtained from studies of stygobionts. These insights could be used not only to disentangle evolution from the adaptations to the selective pressures of groundwater habitats but also to characterize the physiological responses stimulated by the interaction with different environmental conditions.

Our results further demonstrate that stygofauna are reported in springs more frequently than surface fauna are reported in groundwater, in terms of both number of papers and overall numbers of species. In recent years, a growing body of literature has shown that even the occurrence of surface species in caves is often not accidental (Lunghi et al. 2014b; Lunghi et al. 2017), a finding that has important implications for the communities of shallow subterranean habitats (Kozel et al.

2019; Lunghi 2018; Lunghi et al. 2020; Salvidio et al. 2020; Silva et al. 2020). If stygofauna occur in springs and adjacent microhabitats more commonly than surface fauna occur underground, it is likely that, at least for some stygobionts, the use of the surface environment is not accidental. Further systematic reviews and analyses of the literature on spring fauna could be performed to investigate the countries where the largest number of studies on springs were carried out, the most studied taxa and the most studied functional traits.

A parallel can be made between springs and marine caves which can further support the idea that springs are just an ecotone that should also be studied from an ecological viewpoint (Romero, *in litteris*). For example, there are sea fish species that enter and exit marine caves playing a significant role in those environments' ecology. That is the case with the cardinal fish *Apogon imberbis*. This is a small-sized fish distributed along the eastern Atlantic coast from Morocco to the Gulf of Guinea, including the Azores. It can be found as solitary or forming schools and is common in small crevices to marine caves, where they can be found in large densities. They show no troglomorphisms, yet they play a significant role in transferring organic material to these marine caves as mysid crustaceans do (Romero, *in litteris*). Like bats and Dolichopoda cave crickets (Mammola & Isaia 2018), they tend to stay in the shelters during the day and leave the caves at night, presumably for feeding (Bussotti et al. 2003).

The occurrence of stygobionts in springs could affect both the dynamics of boundary habitats and, at the level of the whole stygobiont population, the intrinsic traits of the species. There are several different perspectives for how a stronger conceptual inclusion of springs in subterranean research may provide additional insights on subterranean biology. First, springs may favour intraspecific variation that could be assessed by comparative experimental studies, which would benefit studies of intraspecific dynamics between boundaries and deep areas. Second, springs can inform studies

of the processes that promote adaptation to and colonization of border habitats, as research on springs could be used to distinguish possible phenotypic plasticity from local adaptations. Third, given the view of springs as useful laboratories, devoting space and infrastructure at the entrance to subterranean environments could provide important experimental opportunities.

Conclusions

Even if the transitional and ecotonal role of springs is known and studied since several decades, and the term Groundwater Dependent Ecosystems applied to springs (Fattorini et al. 2020; Rohde et al. 2017) allows to study the connected network of surface and subterranean ecosystems following the 'holistic' approach suggested by (Linke et al. 2019), these concepts are rarely translated in ecological and evolutionary studies dealing with groundwater animals. The results of our systematic review broadly suggest that springs and other boundaries with surface environments should be considered and investigated as part of subterranean habitats and of the biology of at least some stygobionts. Studies of groundwater environments and stygobiont biology that do not consider springs may furnish only a limited perspective on subterranean environments, because they could exclude *a priori* a potential source of selective pressures for groundwater-dwelling animals. The study of groundwater-adapted organisms in subterranean aquifers has the potential to reveal new insights in several scientific fields (Pipan & Culver 2013; Reboleira et al. 2011), but the study of the boundaries of groundwater environments, such as springs, is not only equally important, but even necessary to understand the zoology ecology and evolution of groundwater fauna.

Acknowledgements

We are grateful to Prof. David Culver and Prof. Aldemaro Romero Jr. for comments and suggestions on an early draft of this manuscript. The comments of Stefano Mammola and of an anonymous reviewer improved the quality of the manuscript.

References

- Acreman M, Hughes KA, Arthington AH, Tickner D, and Duenas MA. 2020. Protected areas and freshwater biodiversity: a novel systematic review distils eight lessons for effective conservation. *Conservation Letters* 13. 10.1111/conl.12684
- Alfaro C, and Wallace M. 1994. Origin and classification of springs and historical review with current applications. *Environmental Geology* 24:112-124.
- Aljančič G. 2019. History of research on *Proteus anguinus* Laurenti 1768 in Slovenia. . *Folia Biologica et Geologica* 60:39-69.
- Balland D. 1992. *Les eaux cachées. Études géographiques sur les galeries drainantes souterraines*. Paris: Departement de Géographie, Univers Sorbonne.
- Barzaghi B, Ficetola GF, Pennati R, and Manenti R. 2017. Biphase predators provide biomass subsidies in small freshwater habitats: A case study of spring and cave pools. *Freshwater Biology* 62:1637–1644. DOI: 10.1111/fwb.12975
- Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, and White JS. 2008. Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology and Evolution* 24:127 - 135.
- Botosaneanu L. 1980. Karstic ecosystem - 10 years of hydrobiological studies at CNR underground laboratory in Moulis (France) *Hydrobiologia* 69:21-24. 10.1007/bf00016530

431 Botosaneanu L. 1998. Sources: aux portes du Styx (Springs: gates to and from the Styx). In:
 432 Botosaneanu L, ed. *Studies in Crenobiology- The biology of springs and springbooks*
 433 Leiden: Bachuys Publishers, 229-250.

434 Bressi N, Aljancic M, and Lapini L. 1999. Notes on presence and feeding of *Proteus anguinus*
 435 Laurenti, 1768 outside caves. *Rivista di Idrobiologia* 38:431 - 435.

436 Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ,
 437 Machler M, and Bolker BM. 2017. glmmTMB Balances Speed and Flexibility Among
 438 Packages for Zero-inflated Generalized Linear Mixed Modeling. *R Journal* 9:378-400.

439 Bussotti S, Guidetti P, and Belmonte G. 2003. Distribution patterns of the cardinal fish, *Apogon*
 440 *imberbis*, in shallow marine caves in southern Apulia (SE Italy). *Italian Journal of*
 441 *Zoology* 70:153-157.

442 Cantonati M, Gerecke R, and Bertuzzi E. 2006. Springs of the Alps – sensitive ecosystems to
 443 environmental change: from biodiversity assessments to long-term studies.
 444 *Hydrobiologia* 562:59–96.

445 Cantonati M, Gerecke R, Juttner I, and Cox EJ. 2011. Springs: neglected key habitats for
 446 biodiversity conservation Introduction to the Special Issue. *Journal of Limnology* 70:1-1.

447 Culver D, Pipan T, and Gottstein S. 2006. Hypotelminorheic-a unique freshwater habitat.
 448 *Subterranean Biology* 4:1.

449 Culver DC, Holsinger JR, and Feller DJ. 2012. The Fauna of Seepage Springs and Other Shallow
 450 Subterranean Habitats in the Mid-Atlantic Piedmont and Coastal Plain. *Northeastern*
 451 *Naturalist* 19:1-42. 10.1656/045.019.m901

452 Culver DC, and Pipan T. 2014. *Shallow Subterranean Habitats: Ecology, Evolution, and*
 453 *Conservation*. New York, NY, USA: Oxford University Press.

- 454 Culver DC, and Pipan T. 2019. *The biology of caves and other subterranean habitats*. New
455 York: Oxford University Press.
- 456 Culver DC, Pipan T, Culver DC, and Pipan T. 2014. *Seepage springs and the hypotelminorheic*
457 *habitat*.
- 458 Di Lorenzo T, Cipriani D, Bono P, Rossini L, De Laurentiis P, Fiasca B, Pantani C, and Galassi
459 DMP. 2005. Dynamics of groundwater copepod assemblages from the Mazzoccolo
460 karstic spring (central Italy). *Meiofauna Marina* 14:97-103.
- 461 Fattorini S, Fiasca B, Di Lorenzo T, Di Cicco M, and Galassi DMP. 2020. A new protocol for
462 assessing the conservation priority of groundwater-dependent ecosystems. *Aquatic*
463 *Conservation-Marine and Freshwater Ecosystems* 30:1483-1504. 10.1002/aqc.3411
- 464 Fiasca B, Stoch F, Olivier M, Maazouzi C, Petitta M, Di Cioccio A, and Galassi DMP. 2014.
465 The dark side of springs: what drives small-scale spatial patterns of subsurface
466 meiofaunal assemblages? *Journal of Limnology* 73:71 - 80.
- 467 Ficetola GF, Lunghi E, Canedoli C, Padoa-Schioppa E, Pennati R, and Manenti R. 2018.
468 Differences between microhabitat and broad-scale patterns of niche evolution in
469 terrestrial salamanders. *Scientific Reports* 8:10575.
- 470 Fišer C. 2019. *Niphargus*—A model system for evolution and ecology. In: White WB, Culver
471 DC, and Pipan T, eds. *Encyclopedia of Caves*. Cambridge, MA, USA: Academic Press,
472 746-755.
- 473 Fiser C, Zaksek V, Zagmajster M, and Sket B. 2007. Taxonomy and biogeography of *Niphargus*
474 *steueri* (Crustacea : Amphipoda). *Limnology* 8:297-309. 10.1007/s10201-007-0221-5

475 Fišer Z, Novak L, Lustrik R, and Fiser C. 2016. Light triggers habitat choice of eyeless
 476 subterranean but not of eyed surface amphipods. *Science of Nature* 103:7.
 477 10.1007/s00114-015-1329-9

478 Galassi DMP, Lombardo P, Fiasca B, Di Cioccio A, Di Lorenzo T, Petitta M, and Di Carlo P.
 479 2014. Earthquakes trigger the loss of groundwater biodiversity. *Scientific Reports* 4.
 480 Gibert J, Mathieu J, and Fournier F. 1997. *Groundwater/surface ecotones: biological and*
 481 *hydrological interactions and masnagement options*. Cambridge, UK: University Press.

482 Kozel P, Pipan T, Mammola S, Culver DC, and Novak T. 2019. Distributional dynamics of a
 483 specialized subterranean community oppose the classical understanding of the preferred
 484 subterranean habitats. *Invertebrate Biology* 138. 10.1111/ivb.12254

485 Kresic N. 2010. Types and classifications of springs. In: Kresic N, and Stevanovic Z, eds.
 486 *Groundwater hydrology of springs Engineering, theory, management, and sustainability*.
 487 Amsterdam, The Netherlands.: Elsevier Press, 31-86.

488 Linke S, Turak E, Asmyhr MG, and Hose G. 2019. 3D conservation planning: Including aquifer
 489 protection in freshwater plans refines priorities without much additional effort. *Aquatic*
 490 *conservation*.

491 Lunghi E. 2018. Ecology and life history of *Meta bourneti* (Araneae: Tetragnathidae) from
 492 Monte Albo (Sardinia, Italy). *Peerj* 6. 10.7717/peerj.6049

493 Lunghi E, Ficetola GF, Zhao YH, and Manenti R. 2020. Are the Neglected Tipuloidea Crane
 494 Flies (Diptera) an Important Component for Subterranean Environments? *Diversity-Basel*
 495 12. 10.3390/d12090333

496 Lunghi E, Manenti R, and Ficetola GF. 2014a. Do cave features affect underground habitat
497 exploitation by non-troglobite species? *Acta Oecologica* 55:29 - 35.
498 10.1016/j.actao.2013.11.003.

499 Lunghi E, Manenti R, and Ficetola GF. 2014b. Do cave features affect underground habitat
500 exploitation by non-troglobite species? *Acta Oecologica-International Journal of*
501 *Ecology* 55:29-35. 10.1016/j.actao.2013.11.003

502 Lunghi E, Manenti R, and Ficetola GF. 2017. Cave features, seasonality and subterranean
503 distribution of non-obligate cave dwellers. *Peerj* 5:e3169. DOI: 10.7717/peerj.3169

504 MacAvoy SE, Braciszewski A, Teng E, and Fong DW. 2016. Trophic plasticity among spring
505 vs. cave populations of Gammarus minus: examining functional niches using stable
506 isotopes and C/N ratios. *Ecological Research* 31:589-595. 10.1007/s11284-016-1359-6

507 Malard F, Boutin C, Camacho AI, Ferreira D, Michel G, Sket B, and Stoch F. 2009. Diversity
508 patterns of stygobiotic crustaceans across multiple spatial scales in Europe. *Freshwater*
509 *Biology* 54:756-776.

510 Malard F, Dole-Olivier MJ, Mathieu J, and Stoch F. 2002. *Sampling Manual for the Assessment*
511 *of Regional Groundwater Biodiversity*. Lyon: PASCALIS Project.

512 Mammola S. 2019. Finding answers in the dark: caves as models in ecology fifty years after
513 Poulson and White. *Ecography* 42:1331-1351. 10.1111/ecog.03905

514 Mammola S, Amorim IR, Bichuette ME, Borges PAV, Cheeptham N, Cooper SJB, Culver DC,
515 Deharveng L, Eme D, Ferreira RL, Fiser C, Fiser Z, Fong DW, Griebler C, Jeffery WR,
516 Jugovic J, Kowalko JE, Lilley TM, Malard F, Manenti R, Martinez A, Meierhofer MB,
517 Niemiller ML, Northup DE, Pellegrini TG, Pipan T, Protas M, Reboleira A, Venarsky

MP, Wynne JJ, Zigmajster M, and Cardoso P. 2020. Fundamental research questions in subterranean biology. *Biological Reviews* 95:1855-1872. 10.1111/brv.12642

Mammola S, and Isaia M. 2018. Day-night and seasonal variations of a subterranean invertebrate community in the twilight zone. *Subterranean Biology* 27:31-51. 10.3897/subtbiol.27.28909

Manenti R. 2014. Role of cave features for aquatic troglobiont fauna occurrence: effects on "accidentals" and troglomorphic organisms distribution. *Acta Zoologica Academiae Scientiarum Hungaricae* 60:257-270.

Manenti R, and Barzaghi B. 2020. Is landscape of fear of macroinvertebrate communities a major determinant of mesopredator and prey activity? *Knowledge and Management of Aquatic Ecosystems*. 10.1051/kmae/2019050

Manenti R, and Barzaghi B. In press. Diel activity of *Niphargus* amphipods in spring habitats. *Crustaceana*.

Manenti R, and Pezzoli E. 2019. Think of what lies below, not only of what is visible above, or: a comprehensive zoological study of invertebrate communities of spring habitats. *The European Zoological Journal*, 86:272-279. 10.1080/24750263.2019.1634769

Manenti R, Siesa ME, and Ficetola GF. 2013. Odonata occurrence in caves: active or accidentals? A new case study. *Journal of Cave and Karst Studies* 75:205-209. 10.4311/2012LSC0281

Marković V, Novaković B, Ilić M, and Nikolić V. 2018. Epigean *Niphargids* in Serbia: New Records of *Niphargus valachicus* Dobreaanu & Manolache, 1933 (Amphipoda: Niphargidae), with Notes on its Ecological Preferences. *Acta Zoologica Bulgarica* 70:45-50.

- Mathieu J, Essafi K, and Chergui H. 1999. Spatial and temporal variations of stygobite Amphipod populations in interstitial aquatic habitats of karst floodplain interfaces in France and Morocco. *Annales De Limnologie-International Journal of Limnology* 35:133-139. 10.1051/limn/1999018
- Mathieu J, Essafichergui K, and Jeannerod F. 1994. A gradient of interstitial Niphargus rhenorhodanensis populations in 2 karst floodplain transition zones of the French Jura. *Hydrobiologia* 286:129-137. 10.1007/bf00006244
- Moseley M. 2009. Are all caves ecotones. *Cave and Karst Science* 36:53-58.
- Niemiller ML, Porter ML, Keany J, Gilbert H, Fong DW, Culver DC, Hobson CS, Kendall KD, Davis MA, and Taylor SJ. 2018. Evaluation of eDNA for groundwater invertebrate detection and monitoring: a case study with endangered Stygobromus (Amphipoda: Crangonyctidae). *Conservation Genetics Resources* 10:247-257. 10.1007/s12686-017-0785-2
- Page MJ, and Moher D. 2017. Evaluations of the uptake and impact of the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) Statement and extensions: a scoping review. *Systematic Reviews* 6. 10.1186/s13643-017-0663-8
- Pipan T, and Culver DC. 2012. Convergence and divergence in the subterranean realm: a reassessment. *Biological Journal of the Linnean Society* 107:1-14. DOI 10.1111/j.1095-8312.2012.01964.x
- Pipan T, and Culver DC. 2013. Forty years of epikarst: what biology have we learned? *International Journal of Speleology* 42:215-223. 10.5038/1827-806x.42.3.5
- Pipan T, Fiser C, Novak T, and Culver DC. 2012. Fifty Years of the Hypotelminorheic: What Have We Learned? *Acta Carsologica* 41:275-285.

564 Pirni A. 2016. Space and Anthropology of Limit: A Philosophical Perspective. *Frontiers in*
565 *Astronomy and Space Sciences* 3. 10.3389/fspas.2016.00022

566 Plenet S, and Gibert J. 1995. A comparison of surface water/ground water interface zones in
567 fluvial and karstic systems. *Comptes Rendus de l'Académie des sciences* 318:499-509.

568 Poulson TL, and White WB. 1969. The cave environment. *Science* 165:971–981.

569 Prous X, Ferreira RL, and Martins RP. 2004. Ecotone delimitation: Epigean–hypogean transition
570 in cave ecosystems. *Austral Ecology*, 29:374-382.

571 Reboleira A, Borges PAV, Goncalves F, Serrano ARM, and Oromi P. 2011. The subterranean
572 fauna of a biodiversity hotspot region - Portugal: an overview and its conservation.
573 *International Journal of Speleology* 40:23-37. 10.5038/1827-806x.40.1.4

574 Rechulicz J. 2011. Monitoring of the Topmouth Gudgeon, *Pseudorasbora Parva* (Actinopterygii:
575 Cypriniformes: Cyprinidae) in a Small Upland Ciemiega River, Poland. *Acta*
576 *Ichthyologica Et Piscatoria* 41:193-199.

577 Rohde MM, Froend R, and Howard J. 2017. A Global Synthesis of Managing Groundwater
578 Dependent Ecosystems Under Sustainable Groundwater Policy. *Groundwater* 55:293-
579 301. 10.1111/gwat.12511

580 Romero A. 2009. *Cave biology*. New York: Cambridge University Press.

581 Rouch R. 1986. Sur l'écologie des eaux souterraines dans le karst. *Stygologia* 2:352-398.

582 Salvidio S, Costa A, Oneto F, and Pastorino MV. 2020. Variability of A Subterranean Prey-
583 Predator Community in Space and Time. *Diversity-Basel* 12. 10.3390/d12010017

584 Silva MS, Iniesta LFM, and Ferreira RL. 2020. Invertebrates diversity in mountain Neotropical
585 quartzite caves: which factors can influence the composition, richness, and distribution of
586 the cave communities? *Subterranean Biology* 33:23-43. 10.3897/subtbiol.33.46444

587 Sket B. 2008. Can we agree on an ecological classification of subterranean animals? *Journal of*
588 *Natural History* 42:1549-1563. DOI: 10.1080/00222930801995762

589 Springer AE, and Stevens LE. 2009. Spheres of discharge of springs. *Hydrogeology Journal*
590 17:83-93. 10.1007/s10040-008-0341-y

591 Stoch F, Gerecke R, Pieri V, Rossetti G, and Sambugar B. 2011. Exploring species distribution
592 of spring meiofauna (Annelida, Acari, Crustacea) in the south-eastern Alps. *Journal of*
593 *Limnology* 70:65.

594 Sturz BR, and Bodily KD. 2016. Detecting the perception of illusory spatial boundaries:
595 Evidence from distance judgments. *Cognition* 146:371-376.
596 10.1016/j.cognition.2015.10.015

597 Thomsen PF, Kielgast J, Iversen LL, Wiuf C, Rasmussen M, Gilbert MTP, Orlando L, and
598 Willerslev E. 2012. Monitoring endangered freshwater biodiversity using environmental
599 DNA. *Molecular Ecology* 21:2565-2573.

600 Trajano E, and De Carvalho MR. 2017. Towards a biologically meaningful classification of sub-
601 terranean organisms: a critical analysis of the Schiner-Racovitza system from a historical
602 perspective, difficulties of its application and implications for conservation. *Subterranean*
603 *Biology* 22:1–26.

604 Vandel A. 1920. Sur la faune des sources. *Bulletin De La Societe Zoologique De France* 45:177
605 - 183.

606 Vandel A. 1964. *Biospeleologie: la biologie des animaux cavernicoles*. Paris: Gauthiers-Villars.

607 Vilenica M, Previsic A, Kucinic M, Gattolliat JL, Sartori M, and Mihaljevic Z. 2016.
608 Distribution and Autecology of Mayflies (Insecta, Ephemeroptera) in a Mediterranean
609 River in the Western Balkans. *Entomological News* 126:19-35.

- 610 Von Fumetti S, and Nagel P. 2011. A first approach to a faunistic crenon typology based on
- 611 functional feeding groups. *Journal of Limnology* 70:147-154.
- 612 West KM, Richards ZT, Harvey ES, Susac R, Greal A, and Bunce M. 2020. Under the karst:
- 613 detecting hidden subterranean assemblages using eDNA metabarcoding in the caves of
- 614 Christmas Island, Australia. *Scientific Reports* 10. 10.1038/s41598-020-78525-6
- 615 White NE, Guzik MT, Austin AD, Moore GI, Humphreys WF, Alexander J, and Bunce M. 2020.
- 616 Detection of the rare Australian endemic blind cave eel (*Ophisternon candidum*) with
- 617 environmental DNA: implications for threatened species management in subterranean
- 618 environments. *Hydrobiologia* 847:3201-3211. 10.1007/s10750-020-04304-z
- 619 White WB. 2019. Springs. In: White WB, Culver DC, and Pipan T, eds. *Encyclopedia of Caves*.
- 620 Cambridge, MA, USA: Academic Press, 1031 -1040.
- 621 Yonezawa S, Nakano T, Nakahama N, Tomikawa K, and Isagi Y. 2020. Environmental DNA
- 622 reveals cryptic diversity within the subterranean amphipod genus *Pseudocrangonyx*
- 623 Akatsuka & Komai, 1922 (Amphipoda: Crangonyctoidea: Pseudocrangonyctidae) from
- 624 central Japan. *Journal of Crustacean Biology* 40:479-483. 10.1093/jcbiol/ruaa028

Table 1(on next page)

Table 1

Table 1. Search terms and inclusion/exclusion criteria used to describe published evidence of stygobionts in springs and to answer specific questions.

1 **Table 1**

Categories	Restrictions applied
Number of species mentioned	If clearly stated for all taxa considered in the study
Stygofauna found in springs	If clearly stated that the species found in springs are stygobites
Number of stygofaunal species in springs	If the number of stygobite species found in springs is clearly stated for all taxa considered in the study
Surface fauna found underground	If clearly stated that the species found underground are of epigeal origin
Number of surface species found underground	If the number of epigeal species found underground is clearly stated for all taxa considered in the study
Species found both in caves and springs	If clearly stated that the species found both in caves and springs are epigeal or stygobites
Number of species in both (caves and springs)	If the number of stygobites or epigeal species found in both is clearly stated for all taxa
Ecology	Yes/no, depending on whether the paper provides original ecological information (habitat of occurrence, environmental drivers etc..)
Taxonomy	Yes/no, depending on whether the paper provides original taxonomic data
Behavior	Yes/no, depending on whether the paper tests/reports original behavioral information/observations
Conservation	Yes/no, depending on whether the paper explores original conservation/restoration problems or actions
Faunal assessment	Yes/no, depending on whether the paper is mainly devoted to assess faunal composition of spring/groundwater habitat

2 Table 1. Search terms and inclusion/exclusion criteria used to describe published evidence of
 3 stygobionts in springs and to answer specific questions.

4

5

Table 2 (on next page)

Table 2

Table 2. Number of papers reporting observations of stygofauna in springs, of surface fauna in groundwaters, and of the same taxa in both environments. Papers are divided based on the key words used for the systematic review: GF, groundwater fauna; SFF, spring freshwater fauna.

Table 2

		Total	GF	SFF
Stygofauna in springs	YES	57	49	8
	NO	235	195	40
	Information missing	50	31	19
Surface fauna underground	YES	37	34	3
	NO	278	220	58
	Information missing	27	21	6
Same taxa both in springs and groundwaters	YES	33	30	3
	NO	266	214	52
	Information missing	43	31	12

Table 2. Number of papers reporting observations of stygofauna in springs, of surface fauna in groundwaters, and of the same taxa in both environments. Papers are divided based on the key words used for the systematic review: GF, groundwater fauna; SFF, spring freshwater fauna.

Table 3(on next page)

Table 3

Table 3. The reported occurrence of stygofauna in springs, of surface fauna in groundwaters, and of the same taxa in both habitats shown as a function of study discipline. Relationships were assessed using generalized linear models (GLMs) followed by a likelihood ratio test. Significant relationships are reported in bold.

1 **Table 3**

	Research discipline	Estimate	SE	χ^2	P
Stygofauna in springs	Ecology	0.27	0.34	0.64	0.42
	Taxonomy	0.75	0.39	3.67	0.06
	Faunal assessment	0.24	0.35	0.48	0.49
	Conservation	0.67	0.53	1.50	0.22
	Behavior	-13.84	834.76	0.98	0.32
Surface fauna underground	Ecology	0.02	0.41	0.00	0.96
	Taxonomy	0.27	0.50	0.29	0.59
	Faunal assessment	1.09	0.45	6.62	0.01
	Conservation	0.51	0.59	0.68	0.41
	Behavior	1.30	1.20	0.94	0.33
Same taxa both in springs and groundwaters	Ecology	0.09	0.43	0.04	0.84
	Taxonomy	0.91	0.45	3.96	0.04
	Faunal assessment	0.06	0.44	0.02	0.89
	Conservation	0.77	0.60	1.45	0.23
	Behavior	-13.15	839.90	0.51	0.48

2 Table 3. The reported occurrence of stygofauna in springs, of surface fauna in groundwaters, and
3 of the same taxa in both habitats shown as a function of study discipline. Relationships were
4 assessed using generalized linear models (GLMs) followed by a likelihood ratio test. Significant
5 relationships are reported in bold.

6

Figure 1

Figure 1

Figure 1. Diagram of a spring showing differences between surface and groundwater habitats during day (A) and night (B). During night the border between surface waters and groundwaters softens and stygobionts can move outside interacting with surface invertebrates and vertebrates including predators (here represented by a diurnal predator fish and a nocturnal predator salamander larva). White silhouettes represent stygobionts, black silhouettes represent potential predators (fish and salamanders), and brown silhouettes surface aquatic invertebrates. Drawing is modified from Andrea Melotto and Benedetta Barzaghi (unpublished).




Figure 2

Figure 2

Figure 2 PRISMA 2020 flow diagram for the systematic reviews which included search of Web of Science database only.

