



Gene selection for studying frugivore-plant interactions: a review and an example using Queensland fruit fly in tomato

Shirin Roohigohar, Anthony R. Clarke and Peter J. Prentis

School of Biology and Environmental Science, Queensland University of Technology (QUT), Brisbane, Queensland, Australia

ABSTRACT

Fruit production is negatively affected by a wide range of frugivorous insects, among them tephritid fruit flies are one of the most important. As a replacement for pesticide-based controls, enhancing natural fruit resistance through biotechnology approaches is a poorly researched but promising alternative. The use of quantitative reverse transcription PCR (RT-qPCR) is an approach to studying gene expression which has been widely used in studying plant resistance to pathogens and non-frugivorous insect herbivores, and offers a starting point for fruit fly studies. In this paper, we develop a gene selection pipe-line for known induced-defense genes in tomato fruit, *Solanum lycopersicum*, and putative detoxification genes in Queensland fruit fly, *Bactrocera tryoni*, as a basis for future RT-qPCR research. The pipeline started with a literature review on plant/herbivore and plant/pathogen molecular interactions. With respect to the fly, this was then followed by the identification of gene families known to be associated with insect resistance to toxins, and then individual genes through reference to annotated *B. tryoni* transcriptomes and gene identity matching with related species. In contrast for tomato, a much better studied species, individual defense genes could be identified directly through literature research. For *B. tryoni*, gene selection was then further refined through gene expression studies. Ultimately 28 putative detoxification genes from cytochrome P450 (P450), carboxylesterase (CarE), glutathione S-transferases (GST), and ATP binding cassette transporters (ABC) gene families were identified for *B. tryoni*, and 15 induced defense genes from receptor-like kinase (RLK), D-mannose/L-galactose, mitogen-activated protein kinase (MAPK), lipoxygenase (LOX), gamma-aminobutyric acid (GABA) pathways and polyphenol oxidase (PPO), proteinase inhibitors (PI) and resistance (R) gene families were identified from tomato fruit. The developed gene selection process for *B. tryoni* can be applied to other herbivorous and frugivorous insect pests so long as the minimum necessary genomic information, an annotated transcriptome, is available.

Submitted 12 November 2020
Accepted 21 June 2021
Published 5 August 2021

Corresponding author
Shirin Roohigohar,
s.roohigohar@qut.edu.au,
shirinroohigohar@gmail.com

Academic editor
Daniel Silva

Additional Information and
Declarations can be found on
page 17

DOI 10.7717/peerj.11762

© Copyright
2021 Roohigohar et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Agricultural Science, Ecology, Entomology, Genetics, Molecular Biology

Keywords Quantitative reverse transcription PCR, Gene expression, Fruit induced-defense, Frugivorous insect, Detoxification genes, Tephritidae

INTRODUCTION

Tephritid fruit flies are globally significant pests of horticulture (Aluja & Mangan, 2008; Qin et al., 2015). Frugivorous tephritids lay their eggs into fruit, where the resultant larvae hatch and feed, causing yield loss (Hafsi et al., 2016). With a global trend in trying to reduce the use of pesticides for insect pest control (Ricroch, 2019), alternative management strategies for fruit flies are needed (Sarwar, 2015). Plant breeding for fruit fly resistance is one such option, and while this might be achieved through traditional selection methods (Choudhary et al., 2018; Medjkouh et al., 2018), it is more likely to be achieved in the modern era through manipulation of the plant's defense genes using biotechnological tools (Kumar et al., 2020).

At the phenotype level, numerous studies have documented how variation between fruit, at the species, variety and ripening-stage levels, can impact on fruit fly offspring survival (Aluja, Diaz-Fleischer & Arredondo, 2004; Nunes et al., 2015; Roohigohar, Prentis & Clarke, 2020). Some of these studies have also correlated fruit fly offspring performance with fruit traits such as peel toughness (Diaz-Fleischer & Aluja, 2003; Rattanapun, Amornsak & Clarke, 2009), amount of peel oils and secondary chemicals (Papachristos, Papadopoulos & Nanos, 2008; Papachristos et al., 2009), or forming calluses in 'Hass' and 'Sharwil' avocados around *Anastrepha* sp. egg clusters using a combination of chemically and mechanically induced resistance mechanisms in fruit (Aluja et al., 2014). However, the amount of genotypic data available to help understand fruit fly/fruit interactions to progress biotechnology-based plant defense breeding is sparse and limited to a molecular study of induced defenses of green olive drupes against the olive fruit fly, *Bactrocera oleae* (Corrado et al., 2012; Grasso et al., 2017). This lack of genetic information is in contrast to insect folivory research (Dugé de Bernonville et al., 2017; Gloss, Abbot & Whiteman, 2019; Subramanyam et al., 2019), and also plant pathogen research where there is a growing body of molecular data on fruit defense (Alkan et al., 2015; Rao & Nandineni, 2017; Baba et al., 2019).

Different analytical techniques can be applied to the study of molecular interactions between fruit and frugivorous insects. Corrado et al. (2012) and Grasso et al. (2017) applied comparative transcriptomic and proteomics in their studies, which provides a comprehensive overview of the molecular and protein responses associated with the interaction. However, a limitation of this approach, particularly in non-model organisms (such as crop pest species), is the inability to ascribe function to non-annotated genes and proteins (Giron et al., 2018; Kumaran et al., 2018). A complementary approach can be achieved through gene expression analyses to examine the expression of specific genes already known to be associated with plant/herbivore interactions (Zheng & Dicke, 2008). One of the most reliable techniques for gene expression studies is quantitative reverse transcription PCR (RT-qPCR) (Prasch & Sonnewald, 2013).

The RT-qPCR approach has been used to provide insight to plant defense pathways and insect detoxification gene expression during plant/herbivore interactions in several systems (De Oliveira, Pallini & Janssen, 2019; Altuntaş, Duman & Kılıç, 2020; Dixit et al., 2020; Quais et al., 2020). For example, in fruit/pathogen interaction studies, the over-expression



Figure 1 Schematic representation of the process followed to undertake a RT-qPCR study, and the structural outline of the following sections of this paper.

Full-size  DOI: [10.7717/peerj.11762/fig-1](https://doi.org/10.7717/peerj.11762/fig-1)

of 10 phenylpropanoid genes in orange fruit infested by *Penicillium digitatum* which led to changes in the metabolite profile of the fruit (Ballester, Lafuente & González-Candelas, 2013) was determined using RT-qPCR; while in apples infested with *Penicillium expansum* RT-qPCR was used to track upregulation of defense-related genes and reactive oxygen species genes (Vilanova et al., 2014). Similarly, RT-qPCR has been used to document the over-expression of stress perception genes such as *Prosystemin* in tomato and tobacco plants in response to *Manduca sexta* larval feeding (Orozco-Cardenas, McGurl & Ryan, 1993; Gilardoni et al., 2011); and the upregulation of direct defense genes such as *CYP79B2/B3* in *Arabidopsis* and *TD* gene in tomato plant tissue against *Spodoptera exigua* larval feeding (Müller et al., 2010; Gonzales-Vigil et al., 2011). In insects, RT-qPCR has also been used to demonstrate the upregulation of known detoxification genes *Slgstel*, *Cyp321a7*, *Cyp321a9* and *Cyp6ab14* which increased *Spodoptera litura* larval resistance against plant toxic allelochemicals (Wang et al., 2015a; Wang et al., 2015b; Zou et al., 2016; Wang et al., 2017a; Wang et al., 2017b).

While a valuable counterpart to untargeted transcriptomic and proteomic studies, carrying out a RT-qPCR study from the beginning is not trivial. Before initiating such a study, it first needs to be determined if enough is already known about the system to support such an approach, if so then what are the appropriate genes for study, PCR primers need to be developed for those genes, the appropriate experiments and subsequent RT-qPCR analyses have to be undertaken, and then the results analyzed (Fig. 1). In this paper we work through the RT-qPCR developmental pipeline for a specific fruit fly/fruit system (Queensland fruit fly in tomato fruit) focusing particularly on the selection of appropriate herbivore-induced fruit defense genes and insect detoxification and sequestration genes. While doing so we identify generic issues for consideration to facilitate other RT-qPCR studies in fruit flies and their host fruit, and present a review on the metabolic pathways and associated genes known to be linked with fruit defense and insect detoxification.

MATERIALS AND METHODS

Choosing organisms for study

The frugivore: Queensland fruit fly-Bactrocera tryoni

Selection of the study organism should be driven by research priority, but ideally also the ability to extrapolate results across to related organisms and the availability of some existing genomic information. In Australia, *Bactrocera tryoni* (Froggatt) is a highly polyphagous horticultural pest, attacking most fleshy vegetables and fruit crops (Clarke et al., 2011), and so there is a strong local need for research on this organism. While locally important,

B. tryoni can also serve as a suitable model species for other tephritids as the biology of related species is quite similar (Clarke, 2019). Published transcriptomes of *B. tryoni* are available (Gilchrist et al., 2014; Kumaran et al., 2018) and there is a close similarity of the genetics on this fly and congeneric species. For example, according to the National Centre for Biotechnology Information (NCBI) database, putative *B. tryoni* detoxification pathway genes have an above 90% identity match with *Bactrocera dorsalis* (Hendel), one of the world's most destructive agricultural pests (Qin et al., 2018).

The fruit: tomato-*Solanum lycopersicum*

We chose tomato, *Solanum lycopersicum*, as our model fruit for a number of reasons which should be considered when thinking about what fruit type to use. Firstly, tomato has an accessible, fully sequenced genome (Tomato-Genome-Consortium, 2012) with numerous related genetic and genomic resources available through the National Center for Biotechnology Information (NCBI) database. Its genome is also relatively small (950 Mb) and, conveniently for genomic research, is a diploid species (Gerszberg et al., 2015). Tomato can be grown under many different cultivation conditions (from fully-controlled environments to open-field) with a relatively short life-cycle and has well documented and accessible cultivar variation. These attributes make it an already well-established model system for the study of plant/pathogen and plant/herbivore molecular interactions (Rodriguez-Saona et al., 2010; Kawazu et al., 2012). Finally, for our work, it was already known that different tomato cultivars and ripening stages have significant phenotypic effects on *B. tryoni* offspring performance (Balagawi et al., 2005; Roohigohar, Prentis & Clarke, 2020), and we hypothesized that the difference in performance has a molecular basis.

Strategy in choosing genes of interest

The process for selecting genes of interest is summarized in Fig. 2. Gene selection is a sequential process that involves a combination of literature research, PCR primer design, and laboratory testing. Not all candidate genes identified through literature research may end up being selected because of bioinformatic limitations, or because the genes themselves have very low expression in preliminary trials. The process to identify *B. tryoni* detoxification genes and tomato defense genes follow.

***Bactrocera tryoni* detoxification genes**

Larvae of tephritids such as *B. tryoni*, must stay in a single fruit to complete development (Fitt, 1984); thus, they may utilise specific molecular mechanisms to detoxify fruit toxic secondary compounds. Understanding which molecular pathways fruit fly larvae use to survive in fruit is an essential component when studying frugivore-fruit molecular interactions. In the absence of prior studies on *B. tryoni* larval detoxification genes, we selected target genes using four steps (Fig. 2): (i) a comprehensive literature review on insect detoxification mechanisms against plant allelochemicals and chemical pesticides; (ii) an exhaustive review of any similar studies in other tephritids; (iii) searching for detoxification genes in the *B. tryoni* functional annotation database (Kumaran et al., 2018); and (iv) checking nucleotide and peptide sequences in NCBI-BLAST database,




Figure 2 Schematic representation of the workflow used in the present study to choose inducible defense-related genes in tomato fruit and detoxification-related genes in *Bactrocera tryoni*.

Full-size [DOI: 10.7717/peerj.11762/fig-2](https://doi.org/10.7717/peerj.11762/fig-2)

tblastn, blastx, blastp, smartblast, and Universal Protein Resource/Uniprot to check for homologous proteins, protein domains and genes with >80% identity match from species within the tribe Dacini, such as *Bactrocera dorsalis*. Insects mostly utilize the same enzymes for detoxification of plant allelochemicals and insecticides (Dai et al., 2019), hence searching for genetic information from both plant allelochemical and insecticide studies is appropriate.

The genes identified through this process were associated with different phases of the insect chemical-detoxification process, which occurs for both the detoxification of plant secondary chemicals and pesticides (Heidel-Fischer & Vogel, 2015; Heckel, 2018). During Phase I, genes/enzymes such as P450 monooxygenases (P450s) and carboxylesterases (COEs) are involved in oxidation, hydrolysis or reduction of toxic compounds; subsequently, Phase II involves the conjugation of the modified toxins with hydrophilic groups such as glutathiones, sulphate and sugars by glutathione S-transferases (GSTs) and UDP-glucosyltransferase (UGT) to enhance the polarity of the molecules and so help excretion; while finally, in Phase III, ATP-binding cassette transporters export the conjugated toxins out of the cell (Donkor et al., 2019). Each phase of the detoxification process is associated with major gene families (Fig. 3). The following section describes each of selected gene families, and then provides a list of the individual selected genes for *B. tryoni* larvae. Not all potential genes identified through initial literature searching progressed to the selection stage. Listing all discarded genes is space prohibitive, but for illustrative purposes a selection of the excluded genes, and why they were discarded, are shown as a Data S1.

PHASE I

Cytochrome P450 (P450s). Cytochrome P450 monooxygenases (cytochrome P450s) are a large, complex, and highly conserved gene family of heme-thiolate proteins that encode P450 enzymes (Feyereisen, 2006). The P450s contribute to the catalysis of numerous oxidative reactions during endogenous and exogenous metabolism (Li & Liu, 2019).




Figure 3 The general process of chemical detoxification by insects. Phase I and Phase II involve metabolizing enzymes altering toxic chemicals, while Phase III involves transport and export of conjugated metabolites and elimination of toxins pre-biotransformation. The major gene families associated with each phase are followed: CYP450, Cytochrome P450; EST, Carboxylesterase; GST, Glutathione S-transferase; UGT, UDP-Glucosyltransferase; MDR, Multidrug Resistance Protein; MRP, Multidrug Resistance Associated Protein; ABC, ATP Binding Cassette (Berenbaum & Johnson, 2015; Saha, 2016).

Full-size DOI: 10.7717/peerj.11762/fig-3

Numerous studies report the important role of P450 genes in the metabolism of xenobiotics (George, Rao & Rahangadale, 2019) and plant allelochemicals (Hazzouri et al., 2020).

In Lepidoptera, the members of the P450 CYP6 subfamily play a crucial role in detoxifying a variety of plant toxic secondary compounds (Li, Berenbaum & Schuler, 2001). In Hemiptera, over-expression of the *CYP6CY3* gene in *Myzus persica* helped to detoxify nicotine from the tobacco plant (Bass et al., 2013). In *B. dorsalis*, high expression of *CYPA41* and *CYP6EK1* in the larval and adult malpighian tubules suggest their potential role in detoxification of pesticides (Huang et al., 2012). Amongst 12 P450 genes, *CYP6D9*, *CYP12C2*, and *CYP314A1* were upregulated in *B. dorsalis* following insect exposure to malathion and beta-cypermethrin; while *CYP4E9* expression was upregulated in response to abamectin and beta-cypermethrin exposure (Huang et al., 2013). Insecticide resistance linked to higher expression of P450 genes and their related enzymes has been proposed for both *B. dorsalis* (Jing et al., 2020) and *B. oleae* (Pavlidis et al., 2013).

For *B. tryoni* larvae, 20 genes were selected from the cytochrome P450 gene family: CP6A9, CP313, CP134, CP4D8, CP6G1, C12E1, CP6T1A, CP6T1B, C12C1, C12B1, C12B2, CP304A, C304B, CP306, C6A14, C4AC2, CP4S3, CP132, CP316 and CP6G2.

PHASE II

Carboxylesterase (CarEs). Carboxylesterase (CarEs) are a multigene superfamily ubiquitous in almost all organisms (Marshall *et al.*, 2003). CarEs are involved in hydrolyzing a broad range of ester-containing xenobiotics such as drugs, environmental toxicants, and insecticides (Feng, Li & Liu, 2018). Studies on insect CarEs are mainly focused on their role in metabolizing insecticides and differential expression of CarE genes has been associated with insecticide resistance in different number of insects (Farnsworth *et al.*, 2010; Wang *et al.*, 2019).

CarE genes are associated with the development of malathion resistance in *B. dorsalis* (Wang *et al.*, 2015a; Wang *et al.*, 2015b; Wang, 2016), with the functional role of the esterase B1 (*BdB1*) gene strongly confirmed (Wang *et al.*, 2017a; Wang *et al.*, 2017b). In *B. oleae*, 15 CarE genes were identified as being involved in the metabolism of plant phytotoxins and insecticides (Pavlidis *et al.*, 2013). Two genes from the CarEs superfamily in *B. tryoni* larvae were selected: *ESTF* and *EST1*.

Glutathione S-transferase (GST). Glutathione S-transferases (GSTs) are another multigene family, present in most organisms, which are associated with detoxification (Hayes, Flanagan & Jowsey, 2005). In insects, GSTs have a diversity of functions such as participation in olfaction, oxidative stress responses, and the development and bioactivation of ecdysteroids and hormones (Enya *et al.*, 2015; Zhao *et al.*, 2020), but they are mainly associated with detoxification of endogenous and xenobiotic compounds (Enayati, Ranson & Hemingway, 2005; Che-Mendoza, Penilla & Rodríguez, 2009). The upregulation of GSTs and insecticide resistance is well documented (Shi *et al.*, 2012; Zhang *et al.*, 2019), as is the association between GSTs and insect detoxification of plant allelochemicals (Mittapalli, Neal & Shukle, 2007; Huang *et al.*, 2011).

In *B. oleae*, 33 GSTs are involved with the metabolism of xenobiotics, such as chemical insecticides and plant phytotoxins (Pavlidis *et al.*, 2013). GSTs activities were significantly higher in malathion and λ -cyhalothrin treated *B. zonata* (Yaqoob *et al.*, 2013); while overexpression of the GST gene *BdGSTd9* has been identified as a component of malathion resistance in *B. dorsalis* (Meng *et al.*, 2020).

Three genes from the GST superfamily identified for *B. tryoni* larvae were *GSTD1*, *GSTT1* and *GSTT7*.

PHASE III

ATP binding cassette (ABC) transporters. ATP binding cassette (ABC) transporters are one of the largest transporter gene families across the metazoans (Xiao *et al.*, 2018). ABCs are found in almost all organisms, where they typically have a role in the ATP-dependent transport of various substrates across biological membranes (Broehan *et al.*, 2013). Most ABC transporter genes encode membrane-bound proteins which transport a wide range of molecules, such as amino acids, peptides, sugars, vitamins, sterols, lipids, hormones,

endogenous metabolites, inorganics and xenobiotics, across membranes (Dean, Hamon & Chimini, 2001). Studies of the physiological functions of ABC transporters in arthropods are limited to only a few species (Xiao et al., 2018), typically the “model” species such as *D. melanogaster*, *B. mori*, *Anopheles gambiae*, *Apis mellifera*, and *Tribolium castaneum* (Roth et al., 2003; Liu et al., 2011; Broehan et al., 2013), but large numbers of genes are known to be involved in ABC transporter pathway (Rösner & Merzendorfer, 2020).

The role of ABC transporters in efflux pumps, facilitating cellular excretion of insecticides or metabolites, strengthens a hypothesis for their playing a role in insecticide resistance in insects (Rösner & Merzendorfer, 2020). In a recent study of *S. litura* resistance to pyrethroid, the *ABC5* gene was significantly upregulated and showed a strong correlation with insecticide resistance (Xu et al., 2020). In *B. dorsalis*, ABC transporter genes might play roles in the insecticide resistance, with several *bdABC* genes significantly upregulated after treatment of *B. dorsalis* with malathion, abamectin, and beta-cypermethrin (Xiao et al., 2018). In *B. oleae*, 18 ABC transporter genes were reported for their possible roles in handling xenobiotics, such as plant phytotoxins and insecticides (Pavliidi et al., 2013).

Five genes from the ABC transporters family were selected from *B. tryoni* larvae: *ABCG1*, *ABCA3*, *SUR*, *L259* and *MDR49*.

Selection of tomato defense genes

Genes associated with induced-defense responses in tomato were mostly selected based on previous studies of *Solanaceae*–insect/pathogen molecular interactions; either in plant vegetative tissue or fruit. While defense genes in tomato are much better known than putative detoxification genes in *B. tryoni*, a literature review on tomato/herbivore and tomato/pathogen molecular interactions was still needed to ensure an appropriate selection of genes from across different defense pathways (Fig. 4).

Plant induced defense responses towards herbivores and pathogens first requires recognition systems, such as receptor-like kinase (RLK) and lectin receptor-like kinase (LecRLK), that can perceive herbivore-associated elicitors (HAEs), herbivore-associated molecular patterns (HAMPs), damage-associated molecular patterns (DAMPs) and pathogen-associated molecular patterns (PAMPs) (Santamaria et al., 2013). This recognition triggers the plant’s cell defense responses which are started by ion fluctuation across the membrane and production of reactive oxygen species (ROS), and then continued by mitogen-activated protein kinase (MAPK) cascades phosphorylation and responses, lipoxygenase (LOX) pathway or GABA signalling pathway stimulation (Nejat & Mantri, 2017). After specific phytohormonal crosstalk among a plant’s essential defense-related phytohormones, which include salicylic acid (SA), jasmonic acid (JA) and ethylene (ET) as the core components of plant immune system (Tsuda & Katagiri, 2010), plant defense genes and enzymes such as polyphenol oxidase (PPOs), and protease inhibitors (PIs) are expressed and accumulate in damaged and undamaged plant tissue (Liu, 2018). In the next section, a brief description of each defense pathway and the related gene families are provided before the introduction of individual genes.




Figure 4 Plant inducible defense responses against arthropod and pathogen stressors. At the stress perception stage, DAMPs = damage-associated molecular patterns; HAEs = herbivore-associated elicitors; HAMPs = herbivore-associated molecular patterns; and PAMPs = pathogen-associated molecular patterns. The major defense pathways and defense gene families are as follow: RLK, Receptor-Like Kinase; LecRLK, Lectin Receptor-Like Kinase; ROS, Reactive Oxygen Species; MAPK, Mitogen-Activated Protein Kinase; LOX, Lipoxygenase; GABA, Gamma-Aminobutyric Acid; PPO, Polyphenol Oxidase; R genes, Resistance genes; PI, Protease Inhibitor (Santamaria et al., 2013; Santamaria et al., 2018).

Full-size DOI: 10.7717/peerj.11762/fig-4

Plant perception

Receptor-Like kinase. A plant's perception systems allow them to detect physical injury and pest chemical elicitors through the use of specific receptors, such as receptor-like kinases (RLKs) (Santamaria et al., 2018). RLKs are composed of a transmembrane region, an intracellular kinase domain, and an ectodomain that potentially contributes to ligand binding (Macho & Zipfel, 2014). Plant perception of phytophagous arthropod attack through RLKs has been predominantly investigated in lepidopteran and aphid attack on vegetative tissue (Gouhier-Darimont et al., 2013; Gouhier-Darimont et al., 2019), and pathogen attack to both vegetative (Hashemi et al., 2020) and fruit tissues (Haile et al., 2019).

The target genes associated with RLKs in tomato were PEPR1/2 Ortholog Receptor-Like Kinase1 (*PORK1*) and Lectin receptor kinase1 (*LecRK1*).

The *PORK1* gene (also known as the Tomato Protein Kinase1b (*TPK1b*) interacting protein, (Xu et al., 2018)) has biological functions in wound-systemin signalling and systemin-mediated plant responses to both fungal infestation and insect attack (Liu et al., 2013; Klauser et al., 2015), where systemin is a polypeptide hormone unique to, but common within the Solanaceae (Ryan & Pearce, 2003). *PORK1* is a key determinant of systemin responses in tomato with an important role in tomato plant resistance to *B. cinerea* fungi and *M. sexta* larvae (Xu et al., 2018). *LecRK1* gene activity is important in suppressing insect-mediated inhibition of jasmonic acid-induced defense responses in *Nicotiana attenuata* during herbivory by *M. sexta* larvae (Gilarioni et al., 2011); while conversely suppressing the expression of *LecRK1* in *N. attenuata* increased *M. sexta* folivory (Bonaventure, 2012).

Plant defense signalling transduction

D-mannose/L-galactose pathway. The imposition of excessive biotic and abiotic stress to plants can increase the amount of Reactive Oxygen Species (ROS) and cause critical damage to plant cells (Hasanuzzaman *et al.*, 2020). To regulate ROS in cells, plants have developed enzymatic and non-enzymatic antioxidative defense systems in different cell parts (Ishikawa & Shigeoka, 2008). Among these, ascorbate (AsA) is the most abundant water-soluble antioxidant with multiple functions in metabolism, electron transport, control of the cell cycle, and the response of plants to pathogens and biotic stress (Davey *et al.*, 2000; Smirnoff, 2000). Four pathways exist for AsA biosynthesis, with the D-mannose/L-galactose pathway the dominant (Lorence *et al.*, 2004). The L-galactose pathway is regulated by a number of genes which express differentially during the oxidative stress response of a plant against insects and pathogens (Urzica *et al.*, 2012; Lianthanzauva *et al.*, 2020).

We selected the GDP-L-galactose gene (*GGP2*) as our target gene for the L-galactose pathway.

GGP2 plays an important role in tomato defense responses against abiotic stress and pathogenic infection (Yang *et al.*, 2017); while conversely, deficiency in levels of *GGP2* leads to increased stress susceptibility of tomato plants (Alegre *et al.*, 2020).

Mitogen-activated protein kinase pathway. In all eukaryotic cells, the mitogen-activated protein kinase (MAPK) cascade is one of the major defense pathways involving the transduction of extracellular stimuli into intracellular responses (Zhang & Klessig, 2001). MAPK activation can facilitate signal translocation to the nucleus where, through phosphorylation and activation of transcription factors, gene expression is modulated (Neill *et al.*, 2002). MAPKs are involved in plant signal transduction in response to stress signals from pathogens, drought, cold, wounding, O₃, ROS, and hormone stimuli (Moon *et al.*, 2003; Mittler *et al.*, 2004). Studies have shown the role of MAPK signaling in plant defense against herbivorous insects (Kandath *et al.*, 2007; Wu *et al.*, 2007; Wu & Baldwin, 2010), and in fruit response against pathogens and other stresses (Blanco-Ulate *et al.*, 2013; Zhang *et al.*, 2020a; Zhang *et al.*, 2020b). The target genes associated with MAPKs signalling pathway in tomato are *LeMPK1*, *LeMPK2* and *LeMPK3*.

In tomato, *MPK1*, *MPK2*, and *MPK3* genes have been shown to play an essential role in the wound response signalling pathway and increased plant resistance against *M. sexta* larval herbivory (Kandath *et al.*, 2007). Conversely, inhibition of tomato *MPK1*, *MPK2*, and *MPK3* genes suppressed tomato fruit defense signaling pathways and increased fruit susceptibility to *B. cinerea* infestation (Zheng *et al.*, 2015). In a broader study on tomato plants, *LeMPK1*, *LeMPK2* and *LeMPK3* genes were activated in response to stress caused by the wound-signalling peptide systemin, oligosaccharides elicitors, and fungal toxin fusaric acid (Holley *et al.*, 2003; Higgins *et al.*, 2007).

Lipoxygenase (LOX). The LOX genes play important roles during seed germination, plant growth, and in stress response (Porta & Rocha-Sosa, 2002). LOX catalyzes the initial reaction in the biosynthesis and metabolism of jasmonic acid by inserting molecular oxygen into position 13 of α -linolenic acid (α -LeA) (Christensen *et al.*, 2013). Jasmonic acid (JA) itself

plays an important role in mediating anti-herbivore defense responses in plants (Howe & Jander, 2008). The LOX family genes have been comprehensively studied in different plant species, including tomato (Mariutto et al., 2011), due to their functions in various physiological and molecular events (Viswanath et al., 2020), including their key role in plant defense response against herbivores and pathogens (Park et al., 2010; Vellosillo et al., 2013).

The target genes associated with LOX defense pathway are *LOXB* and *LOXD*.

In tomato plant, the expression level of the chloroplast-targeted LOX gene, *LOXD*, is rapidly induced by leaf wounding (Heitz, Bergey & Ryan, 1997), whereas the *LOXB* gene is expressed only in seeds or fruits (Ferrie et al., 1994). Antisense suppression of tomato *LOXB* caused JA production deficiency in transgenic tomato fruit (Kausch et al., 2012). Similarly, suppression of *LOXD* severely compromised tomato resistance to *H. armigera* and *B. cinerea* (Diaz, Ten Have & Van Kan, 2002; Flors et al., 2007; Shen et al., 2014).

The GABA signalling pathway

Gamma-Aminobutyric Acid (GABA) is a non-protein, four-carbon amino-acid that occurs naturally in microorganisms, plants, and animals which has various metabolic and physiological functions (Ramos-Ruiz, Martinez & Knauf-Beiter, 2019). These functions include acting as an endogenous signalling molecule in the regulation of plant growth and development (Renault et al., 2011), and being an important component in the regulation of carbon/nitrogen metabolism (Bouche & Fromm, 2004). One of the main roles of GABA accumulation in plants is to increase plant resistance to insect herbivory (Bown & Shelp, 2016). GABA, synthesized from Glutamate decarboxylase (GAD) (Shelp, Mullen & Waller, 2012), is a jasmonic-independent pathway induced rapidly after the wounding of plant tissue and cell disruption by feeding insects (Scholz et al., 2015). GAD activation and GABA accumulation due to disruption of cell structure contribute to constitutive and induced direct-defenses against invertebrates (Bown & Shelp, 2016).

The target gene from the GABA pathway is *LeGAD2*.

Overexpression of Glutamate decarboxylase 2 (*LeGAD2*) gene in transgenic tobacco plants reduced feeding by tobacco budworm larvae (MacGregor et al., 2003). The same study supported the hypothesis that mechanically-induced GABA accumulation contributes a resistance mechanism against invertebrate pests, but this may be dependent on changes in the level of gene expression of proteinase inhibitors or other defense products (Kessler & Baldwin, 2002). In tomato plant, silencing of the *GAD2* gene increased the susceptibility of the plant to bacterial (*Ralstonia solanacearum*) infestation (Wang et al., 2019).

The phenylpropanoid pathway

Phenylpropanoids are a large class of plant secondary metabolites that are widely distributed in the plant kingdom (Deng & Lu, 2017). Phenylpropanoids mainly include phenolic acids, stilbenes, coumarins, monolignols, and flavonoids (Vogt, 2010; Liu, Osbourn & Ma, 2015). These metabolites have crucial roles in plant development by acting as essential components of cell walls, protectants against UV radiation, phytoalexins against herbivores and pathogens, and floral pigments to mediate plant–pollinator interactions

(Seal et al., 2004; López-Ráez et al., 2010; Mandal, Chakraborty & Dey, 2010; De Oliveira et al., 2015). Many phenylpropanoid compounds are induced after wounding or herbivore feeding (Bernards & Båstrup-Spohr, 2008) and participate in the establishment of plant resistance (Vogt, 2010).

The target gene from the phenylpropanoids pathway in tomato tissue is *CCoAOMT*.

The caffeoyl-CoAO methyltransferase gene, *CCoAOMT*, was recorded to be involved in production of coumarin and lignin in plant tissue during plant-pathogen interactions (Do et al., 2007; Kai et al., 2008). In tomato fruit, *CCoAOMT* was found to contribute in the biosynthesis of aromatic compounds and lignin in response to pathogen attack and wounding (Miao et al., 2008).

Plant induced defense genes (Anti-nutritional activity)

Plant Polyphenol Oxidase genes (PPOs) are distributed widely in different plant tissues and their discoloration effects in damaged and diseased plant tissue have been known for many years (Tran, Taylor & Constabel, 2012). There is strong evidence of constitutive and induced expression patterns of these genes associated with plant defense against pathogens and insects (Thipyapong et al., 2004; Bhonwong et al., 2009). In tomato plant, the PPO gene family consists of seven members: PPO A, Á, B, C, D, E and F (Newman et al., 1993). These PPO gene members are differentially expressed in vegetative and reproductive tissues of tomato in response to biotic and abiotic stressors (Thipyapong, Joel & Steffens, 1997; Thipyapong, Hunt & Steffens, 2004).

The target genes from the PPO family in tomato tissue are *SIPPO1-2*.

The *SIPPO1-2* genes (are also known as *ppo1-2*, *PPO1-2* and *slPPO1-2*) are the *S. lycopersicum* polyphenol oxidase genes (Kampatsikas, Bijelic & Rompel, 2019). *PPO* gene activity is associated with tomato resistance against phloem-feeding and leaf-chewing insects and also pathogens (Ryan, Gregory & Tingey, 1982; Stout et al., 1998). *PPO1* gene overexpression increased tomato plant resistance against *S. litura* larvae (Thipyapong et al., 2004; Mahanil et al., 2008). Both *PPO1* and *PPO2* genes were highly expressed in tomato leaves infested by *Alternaria solani* fungi (Salim et al., 2011).

Proteinase inhibitor

Plant Protease Inhibitors (PIs) are small proteins that are predominantly present in plant storage tissues, but they have been also found in aerial plant parts (Rehman et al., 2017). Plant PIs are classified as serine proteinase inhibitors, alpha-amylase/trypsin inhibitors, potato type I and type II proteinase inhibitors, serpins, and squash inhibitors (Birk, 2003; Damle et al., 2005). In plant vegetative tissue PIs are induced by insect wounding of plant tissue and play a substantial role in inhibiting folivory (Telang et al., 2009; Chen et al., 2014). Protease inhibitors I and II are also well-known markers of JA mediated defense response in tomato plants and have an anti-nutritive role to feeding herbivores by decreasing the digestibility of dietary protein (Farmer & Ryan, 1992; Felton, 2005). The proteinaceous alpha-amylase inhibitors are accumulated in plant tissues in which they can act as defensive proteins against an insect-herbivore's digestive alpha-amylases (Franco et al., 2002).

The selected genes, which are responsible for the activity of proteinase inhibitor and alpha-amylase inhibitor, are *PII* and *CEVI57* (*PI-II*), and *a-AIs1*.

Solanum lycopersicum wound-induced serine-type proteinase inhibitor I and II (*PII* and *CEVI57*, or *PI-II*), exist in many *Solanaceae* (Bryant *et al.*, 1976; Pearce, Johnson & Ryan, 1993; Fan *et al.*, 2020). Both proteinase inhibitor I and II genes are upregulated in response to mechanical wounding and pathogen attack in tomato plant (Xu *et al.*, 2001; Hamza *et al.*, 2018; Zhang *et al.*, 2020a; Zhang *et al.*, 2020b). Expression of potato inhibitor-II (*Pin-II*) gene in tobacco plants decreased *M. sexta* larval growth (Johnson *et al.*, 1989); while silencing the *PI* gene in transgenic potato plants increased *Leptinotarsa decemlineata* and *Spodoptera exigua* larval weight (Ortego *et al.*, 2001). The expression of the *PIN2* (proteinase inhibitor II) in both mutant and wild tomato plants was influenced by *Helicoverpa zea* feeding (Tian *et al.*, 2014). Alpha-amylase inhibitor 1 (*a-AIs1*) in tomato negatively impacts a feeding herbivore's digestive enzymes (Da Lage, 2018). The alpha-amylase inhibitor level was significantly upregulated in damaged leaves of *Amaranthus* by *M. sexta* larvae in comparison to control leaves (Sánchez-Hernández *et al.*, 2004).

Plant resistance *R* genes

Resistance (*R*) genes are responsible for the plant's innate immune system (Dangl & Jones, 2001). Most *R* genes encode proteins characterized by the existence of a central nucleotide-binding site (NBS), leucine-rich repeats (LRRs), and a variable amino-terminal domain (Takken, Albrecht & Tameling, 2006). The amino-terminal domain determines signalling specificity, while the LRRs are mainly involved in recognition (Martin, Bogdanove & Sessa, 2003). These proteins are distributed across most plant taxa, with the main function being to detect infection by specific pathogens and pests in plant tissue (Chisholm *et al.*, 2006).

The selected gene from this group introduced was *Mi-1.1*.

The NBS-LRR gene, *Mi-1*, is involved in tomato plant resistance against three root-knot nematodes species, potato aphids, tomato powdery mildew and whiteflies (Vos *et al.*, 1998; Nombela, Williamson & Muñiz, 2003; Seifi *et al.*, 2011).

Primer design

The PCR primers for genes were designed using the Primer-BLAST (NCBI) online tool which combines BLAST with global alignment algorithm to ensure full primer-target alignment while being sensitive enough to detect targets with a noticeable number of mismatches to primers (Ye *et al.*, 2012). The following criteria were considered when designing primer pairs: (i) the annealing T_m (melting temperature) should be minimum 58 °C and maximum 62 °C; (ii) the PCR product size must be between 120 to 250 bp; (iii) maximum Poly-x should be 3. For both tomato and *B. tryoni* larvae, mRNA FASTA sequences were used as a PCR template. Each designed primer was tested by inputting to the Primer-BLAST and checking the output gene.

RESULTS

Primer check

RNA extraction and cDNA library synthesis

Snap frozen tissue of tomato and *B. tryoni* larvae were homogenized by Qiagen TissueLyser II (Retsch) in TRIzol reagent. RNA was extracted from infested tomato and larvae separately using the Isolate II RNA Mini Bioline Kit with a subsequent DNase treatment using the Turbo DNA-free kit. The quality and quantity of total RNA were checked by running samples on 1.5% denaturing agarose gel and to ensure DNA was absent a Nanodrop was used. The SensiFAST cDNA synthesis kit (BIO-65053) was used to synthesize a cDNA library by adding 15 μ l of extracted RNA, 4 μ l of TransAmp buffer and 1 μ l of reverse transcriptase enzyme. The master mix was placed in the thermal cycler and the cycling conditions were those provided with the kit.

Primer check by qPCR analysis

The PCR primer pairs designed for tomato and *B. tryoni* were also tested in qPCR reactions with cDNA of tomato fruit and *B. tryoni* larvae as the experimental samples, respectively. As negative controls, we used No Template Control (NTC) and no-primer control reactions with two technical replicates. qPCR was performed using the SensiFast SYBR No-ROX Kit (BIO-98020). For testing each PCR primer pair, 10 μ l of SensiFast SYBR, 0.8 μ l of each forward and reverse primers, 0.5 μ l cDNA and 7.9 μ l H₂O were used with the final volume of 20 μ l. We used LightCycler[®]96 Instrument (Roche) by adjusting two steps cycling and melting: 1 cycle (polymerase activation) in 95 °C for 2 min and 40 cycles in 95 °C in 5 s for denaturation and 60–65 °C in 15–30 s for annealing/extension. The cycle quantification of each target gene were checked (Table 1) and genes with a high cycle threshold (>34) were removed from further analysis. Genes with high NTC cycle threshold (>33) were acceptable for inclusion in the further study. The final PCR primer pairs that were selected in this study are shown in Table 2.

Primer consistency in experimental samples

To demonstrate the C_q consistency of the PCR primer pairs, the qPCR results from three replicates of our subsequent study have been presented in Table 3. The results were obtained from phenotypic and molecular studies to identify tomato fruit induced defense responses against *B. tryoni* larvae. The experiment was conducted under semi-natural conditions (glasshouse) while tomatoes were still on the plant. Fourty fruit from each of the two different cultivars and two different ripening stages were inoculated with 40 *B. tryoni* neonate larvae. After inoculation, half of the fruit were picked immediately and kept in the same condition as unpicked fruit. Inoculated fruit were then dissected at two different time points (48 hr and 120 hr) to reflect the two different larval stages under normal developmental conditions. Surviving *B. tryoni* larvae and infested tomato tissue from each of the fourty replicates were transferred to 2.00 ml microtubes separately and then snap frozen using liquid nitrogen and kept at –80 °C until required for RNA extraction.

Here in Table 3, the cycle threshold of three replicates from unpicked and picked treatments (tomato tissue and surviving larvae) at 48 hr timepoint shows the primer

Table 1 Mean qPCR primer check results of primer pairs with and without *Bactrocera tryoni* larval tissue cDNA or tomato fruit tissue cDNA. Two technical replications were carried out and the cycle quantification analysis was done in LightCycler® 96, version 1.1.0.1320, Roche. Genes with NTC Cq mean above 33 were acceptable for inclusion in the further study.

Gene name	Experimental Sample		No Template Control (NTC)	
	Cq mean	Cq error (SD)	Cq mean	Cq error
<i>B. tryoni</i>				
<i>GSTD1</i>	21.04	0.01	–	–
<i>GSTT1</i>	18.51	0.00	–	–
<i>GSTT7</i>	27.01	0.16	–	–
<i>ESTF</i>	24.79	0.08	–	–
<i>EST1</i>	29.84	0.18	–	–
<i>SUR</i>	28.12	0.19	–	–
<i>ABCG1</i>	25.76	0.07	–	–
<i>ABCA3</i>	23.00	0.01	–	–
<i>L259</i>	25.36	0.02	37.54	0.00
<i>MDR49</i>	24.97	0.04	–	–
<i>CP6A9</i>	27.98	0.01	39.28	0.00
<i>CP313</i>	24.39	0.01	–	–
<i>CP134</i>	24.19	0.05	–	–
<i>CP4D8</i>	27.80	0.23	37.91	0.00
<i>CP6G1</i>	25.07	0.03	–	–
<i>C12E1</i>	21.44	0.02	–	–
<i>CP6T1A</i>	28.24	0.01	37.16	0.00
<i>C12C1</i>	24.28	0.01	37.00	0.23
<i>CP6T1B</i>	28.16	0.11	32.96	1.29
<i>C12B2</i>	22.29	0.03	37.78	0.98
<i>C12B1</i>	26.79	0.01	–	–
<i>CP306</i>	27.16	0.02	–	–
<i>CP304A</i>	27.74	0.11	–	–
<i>C6A14</i>	23.34	0.05	37.83	0.91
<i>C4AC2</i>	27.24	0.01	30.82	0.08
<i>CP4S3</i>	23.99	0.05	38.13	1.02
<i>CP132</i>	24.01	0.02	–	–
<i>CP316</i>	28.08	0.03	36.24	0.08
<i>CP304B</i>	26.93	0.02	–	–
<i>CP6G2</i>	26.66	0.08	38.62	0.00
Tomato fruit				
<i>PORK1</i>	25.22	0.08	37.34	0.00
<i>SIPO1</i>	33.57	1.10	–	–
<i>SIPO2</i>	33.10	0.01	–	–
<i>LeRK1</i>	21.90	0.02	–	–
<i>PIIF</i>	33.43	0.04	–	–

(continued on next page)

Table 1 (continued)

Gene name	Experimental Sample		No Template Control (NTC)	
	Cq mean	Cq error (SD)	Cq mean	Cq error
<i>CEVI57</i>	18.88	0.22	–	–
<i>LeMPk1</i>	21.97	0.09	–	–
<i>LeMPK2</i>	20.21	0.18	–	–
<i>LeMPK3</i>	18.44	0.01	38.22	0.00
<i>GGP2</i>	19.37	0.20	–	–
<i>Mi_1.1</i>	26.15	0.00	35.83	0.92
<i>LOXB</i>	16.23	0.06	36.95	0.00
<i>LOXD</i>	17.41	0.00	35.32	0.74
<i>CCoAOMT</i>	18.77	0.11	–	–
<i>LeGAD2</i>	20.08	0.06	–	–
<i>a-AIs1</i>	30.23	0.30	37.36	0.00

consistency in candidate genes under the experimental conditions. The amount of tissue in each of the three replicates included the tissue collected in 2.00 ml microtubes for tomato and 20–25 larvae for *B. tryoni*.

CONCLUSION

Through a combined worked-example and literature review, this paper has identified genes known to be associated with induced-defense against herbivores and pathogens in tomatoes, and genes putatively associated with detoxification in *B. tryoni* based on their known action in other insect herbivore systems. Applied to *B. tryoni* larvae and tomato fruit harvested under different experimental conditions, the genes selected have been shown to respond based on the predictable patterns from the literature (S Roohigohar, AR Clarke, PJ Prentis, 2021, unpublished data). Of 30 selected genes for *B. tryoni* larvae, two genes (*C4AC2* and *C12C1*) were excluded from our study due to PCR primer failure (Table 1) or high Cq in most of replicates. In tomato, one gene (*CEVI57*) from 16 selected genes was excluded due to no Cq in most of replicates. The PCR primers designed are specific for *B. tryoni* and tomato, but the approach followed is directly transferable to other systems so long as there is already at least some genomic resources, at a minimum an annotated transcriptome.

The gene selection process for *B. tryoni* larvae developed here is novel in insect frugivory research. In contrast, the much more straight-forward gene selection process for tomato fruit (Fig. 2) shows the advantage of having expanded functional genomic studies which are now common in plant pathology. Plant protection entomologists are clearly still lagging with respect to their plant pathology colleagues in this field.

Untargeted molecular approaches, such as comparative transcriptomics, provide important insights into the overall changes in gene expression associated with two or more states, such as larvae growing in different fruit types (Corrado et al., 2012). However, more quantitative candidate gene studies, such as RT-qPCR approaches, are also needed if the intent of the research is to create resistant fruit genotypes. For a fruit fly/fruit system,

Table 2 A list of genes and their PCR primers developed for studying the fruit induced-defense/frugivorous insect-detoxification interactions occurring between *Bactrocera tryoni* larvae feeding in tomato fruit.

<i>Bactrocera tryoni</i> detoxification pathways genes and primers		
Gene symbol	Forward sequence 5'–3'	Reverse sequence 5'–3'
<i>GSTD1</i>	GCCGATTTACCACGTATGC	GCGTGTATCGCTGAAACGTC
<i>GSTT1</i>	TTAGCACCATAGACGTGGCG	TGG GCAATACTGCGGAACCT
<i>GSTT7</i>	TGGCCGGTGATCAGTTGAAA	GCTGATCGACCATAGCACGA
<i>ESTF</i>	AGCTAAACCTTCCACCACGG	CACCCATTGCAAAGCCAGAC
<i>EST1</i>	CGCTGTTTACGCATTCTCTG	AGCGGACGCATACTCATAGC
<i>SUR</i>	TTGCTCAAGGCAAAGCGAAC	CATCGTCATCCGTCTGCTCA
<i>ABCG1</i>	TTCTTTGTGCGGTGCTACGCT	ATGGGCGTTCCAAGCCATAA
<i>ABCA3</i>	GGAATAGCGATTGCGGGTA	CGTTCTTCCATGTGATGCG
<i>L259</i>	CAGGAGCCAGCACGTAAGA	GGTCCAATGACGGCCACTAA
<i>MDR49</i>	TGAGGCAACCTCGGCTTTAG	CCGAGCGCATAAGTTCAACG
<i>CP6A9</i>	GTATCGCTTGCAACTCGCTG	CGCACGATGCGCATAAAGAA
<i>CP313</i>	AACACTTCAAACCGGAGGCA	CTCCAGCTGACACAACGGAT
<i>CP134</i>	AGGGCATTTCGATTGGCAGA	TCACCCGCATCGTTTCGTTA
<i>CP4D8</i>	ATTTACTCGCACGCCATCCA	CGGCACACTGGGATAGAGAC
<i>CP6G1</i>	TGGACGAAGTGTGCGCTTA	GGATCGAAAAGTGTCCGGGTT
<i>C12E1</i>	ATGTGGACTTGGAGAACGCA	TCCATTTCCCGAATGGCAGT
<i>CP6T1A</i>	TGCATAATCATGCGCTGCTG	GTCTCCAGCTTACCGCCAAT
<i>CP6T1B</i>	CGCGCACATCTTTACTCAGC	GCCAGTAACAAGAAAGCGGC
<i>C12B2</i>	CAGCTTTCGGATGTTGCGAG	ACCGGCCAGATGGTTTCATT
<i>C12B1</i>	TACGCACACTGCCGAAAGAT	TTCCGGACAAGCACTCTCAC
<i>CP306</i>	CCTGCTCGCGCTATTAGTCA	TTCAAGAATTCCTCCGACC
<i>CP304</i>	AGCGTCGTGCTGACGATTAT	GTATGCCCATTCGCGTGTTC
<i>C6A14</i>	ACACTGCGGAAATACACGGT	CGAAACGATCGGGTTCAGGA
<i>CP4S3</i>	AAGCGCTGAAGGTAAGTGCAT	AAGTGTGCGACTTCTTCGCGT
<i>CP132</i>	AGCACACCTCTTCAATCCCG	CTGCGATCTCAGCATAACGC
<i>CP316</i>	AATCGGTTCCGGTGCAGAAGT	ATGATCTGCGCTGTGTAGCA
<i>CP304</i>	TGAGGTCGTAGGTAGAGGGC	GCTCCGTGTCTACCAATGCT
<i>CP6G2</i>	CGCGCTGTGTTCAAGTTCAG	CGCAGAAACTCGGTAGAGGT
Tomato defensive pathways genes and primers		
<i>PORK1</i>	AGACCCTCAATGAAAGAGGTA	GGTGGAGCTAGAAGTGAGACA
<i>sLPO1</i>	GTGGACAGGATGTGGAACGA	CTTCTTGGTGTCCAGGCAGT
<i>sLPO2</i>	AGTTGTTGCCCTCCTGTACC	CCCTCATTCGACTCGTAGCC
<i>LecRK1</i>	CTTTGCAGGCATCGTGCTTT	GCGCAAAGGTGAAGGGATTG
<i>PIIF</i>	TGGTGTACCAACAAAGCTTGC	GCATTTGTACAACAAAGCCCA
<i>LeMPK1</i>	GATGGTTCCGTTCCGCAAAC	GAACCTGCCACCATGGCTTA
<i>LeMPK2</i>	GCGCTTGCTCATCTTACCT	AATCCAACAGCAAACGAGCG
<i>LeMPK3</i>	CGCCCTTACGAAGGGAGTTT	ACTTTAGCCCACGGAGAAGC
<i>GGP2</i>	CCTCCACTTCCAGGCGTATT	GCATCAGACAAATCACGGGC
<i>Mi-1.1</i>	AAAGCTCACCAGTGGATCGG	CCATGCACGAAGGTCGAAAC

(continued on next page)

Table 2 (continued)

Bactrocera tryoni detoxification pathways genes and primers

Gene symbol	Forward sequence 5'–3'	Reverse sequence 5'–3'
<i>LOXB</i>	GCGTTTAAGGCTTTGTGCGA	GTAGGCCTTGACCATCCGTT
<i>LOXD</i>	GCAGATCGCTAAAGCACACG	GCGCTTAAGTGCCTATGTGC
<i>CCoAOMT</i>	ACCAAATGATTGACGACGGC	TCCGTTCCAAAGGGTGTGT
<i>LeGAD2</i>	TGAGCCCTGAGAAAGCTGTG	GGAGTGTCCCACCCTGTTC
<i>a-AIs1</i>	AAGTGCCTCACCAACACCAT	CAGAATTCGTCGCGGATGGA

Table 3 Mean qPCR primer check results of primer pairs for selected putative detoxification genes in *Bactrocera tryoni* and induced-defence mechanism genes in tomato fruit. For *B. tryoni* larvae, *RS10B*, *RK18A*, *RT15*, *RT14* genes and for tomato *FPPS1*, *ID11* genes used as housekeeping genes (internal control). The qPCR template was cDNA for *B. tryoni* larvae or tomato tissue. The Cq means calculated from two technical replications and the cycle quantification analysis was done in LightCycler® 96, version 1.1.0.1320, Roche.

Gene name	Unpicked status			Picked status		
	Cq mean			Cq mean		
	Rep 1	Rep 2	Rep 3	Rep 1	Rep 2	Rep 3
<i>B. tryoni</i>						
<i>GSTD1</i>	19.66	20.07	19.19	20.23	20.38	19.96
<i>GSTT1</i>	17.10	17.71	17.45	17.93	17.79	18.86
<i>GST7</i>	25.64	25.34	25.00	25.67	25.90	25.05
<i>ESTF</i>	23.82	24.10	23.22	23.91	23.81	24.21
<i>EST1</i>	28.63	28.39	28.25	28.38	28.42	28.26
<i>SUR</i>	27.10	26.81	26.05	27.36	27.17	27.34
<i>ABCG1</i>	24.46	24.71	24.84	25.01	25.18	25.68
<i>ABCA3</i>	22.69	22.82	21.74	22.27	22.78	22.39
<i>L259</i>	24.81	25.07	24.70	24.51	24.64	24.53
<i>MDR49</i>	24.06	25.17	22.37	24.15	24.16	23.29
<i>CP6A9</i>	26.98	27.33	26.84	26.80	27.08	26.53
<i>CP313</i>	22.66	23.67	21.76	22.99	23.79	22.27
<i>CP134</i>	27.25	27.19	24.71	26.77	26.95	25.07
<i>CP4D8</i>	28.46	29.37	29.54	27.91	29.18	26.57
<i>CP6G1</i>	27.03	27.09	26.98	27.75	27.71	27.08
<i>C12E1</i>	23.12	22.76	21.52	22.60	22.77	21.69
<i>CP6T1A</i>	28.71	29.11	28.43	28.34	29.32	27.43
<i>CP6T1B</i>	28.72	28.86	28.57	28.01	29.08	27.09
<i>C12B2</i>	22.96	22.24	21.03	22.05	22.15	22.03
<i>C12B1</i>	26.86	27.17	25.89	26.92	26.88	26.00
<i>CP306</i>	25.65	25.84	25.30	25.13	25.76	25.93
<i>CP304A</i>	28.68	28.49	29.74	28.05	28.93	27.42
<i>C6A14</i>	24.44	23.56	21.97	23.83	23.47	22.85
<i>CP4S3</i>	24.70	23.96	23.80	24.11	24.91	25.08

(continued on next page)

Table 3 (continued)

Gene name	Unpicked status			Picked status		
	Cq mean			Cq mean		
	Rep 1	Rep 2	Rep 3	Rep 1	Rep 2	Rep 3
<i>CP132</i>	23.03	22.59	21.98	22.71	22.30	22.00
<i>CP316</i>	27.65	27.64	26.09	27.34	26.97	26.74
<i>CP304B</i>	27.68	28.01	28.03	27.32	28.55	27.09
<i>CP6G2</i>	27.43	27.60	26.89	27.06	27.91	26.84
HK genes						
<i>RS10B</i>	14.60	14.49	14.12	14.42	14.87	14.84
<i>RK18A</i>	13.92	13.76	13.48	13.94	14.11	13.91
<i>RT15</i>	19.59	19.39	19.41	19.25	19.56	20.28
<i>RT14</i>	19.46	19.06	18.77	19.26	19.37	19.50
Tomato fruit						
<i>PORK1</i>	24.56	25.10	25.4	25.31	26.13	27.01
<i>SIPO1</i>	30.63	29.33	30	31.92	30.81	30.04
<i>SIPO2</i>	30.55	29.86	28.87	30.18	31.53	30.52
<i>LeRK1</i>	20.90	20.94	21.58	21.34	22.14	22.70
<i>PII</i>	21.33	20.74	18.89	21.01	21.48	29.22
<i>LeMPK1</i>	21.73	21.48	21.47	21.91	22.41	24.42
<i>LeMPK2</i>	19.44	19.72	19.72	20.11	20.10	21.10
<i>LeMPK3</i>	18.94	18.56	18.74	18.99	19.41	20.29
<i>GGP2</i>	19.01	19.48	19.07	19.33	19.38	21.01
<i>Mi_1.1</i>	26.53	26.02	27.06	27.74	27.61	29.83
<i>LOXB</i>	15.56	17.04	16.88	15.29	14.83	14.41
<i>LOXD</i>	17.63	17.55	18.79	18.18	20.06	20.71
<i>CCoAOMT</i>	19.10	19.96	19.84	19.64	20.21	21.62
<i>LeGAD2</i>	19.37	19.89	19.49	20.06	20.27	23.93
<i>a-AIs1</i>	24.60	29.13	25.71	29.20	32.32	30.03
HK genes						
<i>FPFS1</i>	18.22	19.70	18.46	18.40	18.78	19.63
<i>IDI1</i>	18.00	18.60	18.17	18.39	18.04	18.70

knowing when and where (i.e., on what cellular or metabolic pathway) larvae are most challenged by plant defenses, and similarly when and how the fruit are challenging the larvae, is fundamental to any manipulation of the system.

ACKNOWLEDGEMENTS

The authors would like to thank Ms Thelma Peek and Mrs Linda Clarke (Queensland Department of Agriculture and Fisheries) for kindly supplying the *B. tryoni* used in experiments. Lab space was provided by Molecular Genomics Research Facility and technical support was provided by Vincent Chand and Victoria Coyne at Queensland University of Technology, Brisbane, Australia. Thanks to QUT colleagues Francesca Strutt, Chloe van der Burg and Hayden Smith for their guidance in RNA isolation, designing

PCR primer pairs and RT-qPCR techniques. Similarly, we thank Kumaran Nagalingam for suggestions in choosing detoxification genes in *B. tryoni*.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

Anthony R. Clarke and the QUT fruit fly lab are funded by the Australian Research Council through its Discovery Project (DP180101915) and Industrial Transformation Training Centre (IC150100026) schemes. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Australian Research Council through its Discovery Project: DP180101915.
Industrial Transformation Training Centre: IC150100026.

Competing Interests

Peter J. Prentis is an Academic Editor for PeerJ.

Author Contributions

- Shirin Roohigohar conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Anthony R. Clarke and Peter J. Prentis conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The raw data is available at Zenodo: Shirin Roohigohar. (2020). Gene selection for studying frugivore-plant interactions: a review and an example using Queensland fruit fly in tomato. <http://doi.org/10.5281/zenodo.4267156>.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.11762#supplemental-information>.

REFERENCES

- Alegre ML, Steelheart C, Baldet P, Rothan C, Just D, Okabe Y, Ezura H, Smirnoff N, Geogoff Grozeff GE, Bartoli CG. 2020. Deficiency of GDP-1-galactose phosphorylase, an enzyme required for ascorbic acid synthesis, reduces tomato fruit yield. *Planta* 251:54 DOI 10.1007/s00425-020-03345-x.
- Alkan N, Friedlander G, Ment D, Prusky D, Fluhr R. 2015. Simultaneous transcriptome analysis of *Colletotrichum gloeosporioides* and tomato fruit pathosystem reveals novel

- fungal pathogenicity and fruit defense strategies. *New Phytologist* **205**:801–815 DOI [10.1111/nph.13087](https://doi.org/10.1111/nph.13087).
- Altuntaş H, Duman E, Kılıç G. 2020.** Juglone induced oxidative and genotoxic stress in the model insect *Galleria mellonella* L. (Pyralidae: Lepidoptera). *International Journal of Tropical Insect Science* **40**:611–619 DOI [10.1007/s42690-020-00107-w](https://doi.org/10.1007/s42690-020-00107-w).
- Aluja M, Arredondo J, Diaz-Fleischer F, Birke A, Rull J, Niogret J, Epsky N. 2014.** Susceptibility of 15 Mango (Sapindales: Anacardiaceae) cultivars to the attack by *Anastrepha ludens* and *Anastrepha obliqua* (Diptera: Tephritidae) and the role of underdeveloped fruit as pest reservoirs: Management implications. *Journal of Economic Entomology* **107**:375–388 DOI [10.1603/EC13045](https://doi.org/10.1603/EC13045).
- Aluja M, Díaz-Fleischer F, Arredondo J. 2004.** Non-host status of *Persea americana* cultivar Hass to *Anastrepha ludens*, *A. obliqua*, *A. serpentina*, and *A. striata* (Diptera: Tephritidae). *Journal of Economic Entomology* **97**:293–309 DOI [10.1093/jee/97.2.293](https://doi.org/10.1093/jee/97.2.293).
- Aluja M, Mangan RL. 2008.** Fruit fly (Diptera: Tephritidae) host status determination: critical conceptual, methodological, and regulatory considerations. *Annual Review of Entomology* **53**:473–502 DOI [10.1146/annurev.ento.53.103106.093350](https://doi.org/10.1146/annurev.ento.53.103106.093350).
- Baba VY, Constantino LV, Ivamoto ST, Moreira AFP, Madeira TB, Nixdorf SL, Rodrigues R, Gonçalves LSA. 2019.** *Capsicum-Colletotrichum* interaction: identification of resistance sources and quantification of secondary metabolites in unripe and ripe fruits in response to anthracnose infection. *Scientia Horticulturae* **246**:469–477 DOI [10.1016/j.scienta.2018.11.011](https://doi.org/10.1016/j.scienta.2018.11.011).
- Balagawi S, Vijaysegaran S, Drew RA, Raghu SJ. 2005.** Influence of fruit traits on oviposition preference and offspring performance of *Bactrocera tryoni* (Froggatt) (Diptera: Tephritidae) on three tomato (*Lycopersicon lycopersicum*) cultivars. *Australian Journal of Entomology* **44**:97–103 DOI [10.1111/j.1440-6055.2005.00459](https://doi.org/10.1111/j.1440-6055.2005.00459).
- Ballester AR, Teresa Lafuente M, González-Candelas L. 2013.** Citrus phenylpropanoids and defense against pathogens. Part II: gene expression and metabolite accumulation in the response of fruits to *Penicillium digitatum* infection. *Food Chemistry* **136**:285–291 DOI [10.1016/j.foodchem.2012.08.006](https://doi.org/10.1016/j.foodchem.2012.08.006).
- Bass C, Zimmer CT, Riveron JM, Wilding CS, Wondji CS, Kausmann M, Field LM, Williamson MS, Nauen R. 2013.** Gene amplification and microsatellite polymorphism underlie a recent insect host shift. *Proceedings of the National Academy of Sciences of the United States of America* **110**:19460–19465 DOI [10.1073/pnas.1314122110](https://doi.org/10.1073/pnas.1314122110).
- Berenbaum MR, Johnson RM. 2015.** Xenobiotic detoxification pathways in honey bees. *Current Opinion in Insect Science* **10**:51–58 DOI [10.1016/j.cois.2015.03.005](https://doi.org/10.1016/j.cois.2015.03.005).
- Bernards MA, Båstrup-Spohr L. 2008.** Phenylpropanoid metabolism induced by wounding and insect herbivory. In: Schaller A, ed. *Induced plant resistance to herbivory*. Dordrecht: Springer, 189–211 DOI [10.1007/978-1-4020-8182-8_9](https://doi.org/10.1007/978-1-4020-8182-8_9).
- Dugé de Bernonville T, Carqueijeiro I, Lanoue A, Lafontaine F, Sánchez Bel P, Liesecke F, Musset K, Oudin A, Glévarec G, Pichon O, Besseau S, Clastre M, St-Pierre B, Flors V, Maury S, Huguet E, O'Connor SE, Courdavault V. 2017.** Folivory elicits a strong defense reaction in *Catharanthus roseus*: metabolomic and transcriptomic

- analyses reveal distinct local and systemic responses. *Scientific Reports* 7:40453 DOI 10.1038/srep40453.
- Bhonwong A, Stout MJ, Attajarusit J, Tantasawat P. 2009.** Defensive role of tomato polyphenol oxidases against cotton bollworm (*Helicoverpa armigera*) and beet armyworm (*Spodoptera exigua*). *Journal of Chemical Ecology* 35:28–38 DOI 10.1007/s10886-008-9571-7.
- Birk Y. 2003.** *Plant protease inhibitors: significance in nutrition, plant protection, cancer prevention and genetic engineering*. Berlin: Springer.
- Blanco-Ulate B, Vincenti E, Powell A, Cantu D. 2013.** Tomato transcriptome and mutant analyses suggest a role for plant stress hormones in the interaction between fruit and *Botrytis cinerea*. *Frontiers in Plant Science* 4:142 DOI 10.3389/fpls.2013.00142.
- Bonaventure G. 2012.** Perception of insect feeding by plants. *Plant Biology* 14:872–880 DOI 10.1111/j.1438-8677.2012.00650.x.
- Bouche N, Fromm H. 2004.** GABA in plants: just a metabolite? *Trends in Plant Science* 9:110–115 DOI 10.1016/j.tplants.2004.01.006.
- Bown AW, Shelp BJ. 2016.** Plant GABA: not just a metabolite. *Trends in Plant Science* 21:811–813 DOI 10.1016/j.tplants.2016.08.001.
- Broehan G, Kroeger T, Lorenzen M, Merzendorfer H. 2013.** Functional analysis of the ATP-binding cassette (ABC) transporter gene family of *Tribolium castaneum*. *BMC Genomics* 14:6 DOI 10.1186/1471-2164-14-6.
- Bryant J, Green TR, Gurusaddaiah T, Ryan CA. 1976.** Proteinase inhibitor II from potatoes: isolation and characterization of its protomer components. *Biochemistry* 15:3418–3424 DOI 10.1021/bi00661a004.
- Che-Mendoza A, Penilla RP, Rodriguez DA. 2009.** Insecticide resistance and glutathione S-transferases in mosquitoes. *African Journal of Biotechnology* 8:1386–1397.
- Chen PJ, Senthilkumar R, Jane WN, He Y, Tian Z, Yeh KW. 2014.** Transplastomic *Nicotiana benthamiana* plants expressing multiple defense genes encoding protease inhibitors and chitinase display broad-spectrum resistance against insects, pathogens and abiotic stresses. *Plant Biotechnology Journal* 12:503–515 DOI 10.1111/pbi.12157.
- Chisholm ST, Coaker G, Day B, Staskawicz BJ. 2006.** Host-microbe interactions: shaping the evolution of the plant immune response. *Cell* 124:803–814 DOI 10.1016/j.cell.2006.02.008.
- Choudhary JS, Dhakar MK, Mukherjee D, Das MB, Singh AK, Bhatt BP. 2018.** Influence of some fruit traits of mango, *Mangifera indica* L. varieties against maggot development and infestation of mango fruit fly, *Bactrocera dorsalis* (Hendel). *Journal of Entomology and Zoology Studies* 6:2621–2627.
- Christensen SA, Nemchenko A, Borrego E, Murray I, Sobhy IS, Bosak L, De Blasio S, Erb M, Robert CA, Vaughn KA. 2013.** The maize lipoxygenase, Zm LOX 10, mediates green leaf volatile, jasmonate and herbivore-induced plant volatile production for defense against insect attack. *The Plant Journal* 74:59–73 DOI 10.1111/tpj.12101.
- Clarke AR. 2019.** *Biology and management of bactrocera and related fruit flies*. Wallingford: CAB International.

- Clarke AR, Powell KS, Weldon CW, Taylor PW. 2011. The ecology of *Bactrocera tryoni* (Diptera: Tephritidae): what do we know to assist pest management? *Annals of Applied Biology* 158:26–54 DOI 10.1111/j.1744-7348.2010.00448.
- Corrado G, Alagna F, Rocco M, Renzone G, Varricchio P, Coppola V, Coppola M, Garonna A, Baldoni L, Scaloni A, Rao R. 2012. Molecular interactions between the olive and the fruit fly *Bactrocera oleae*. *BMC Plant Biology* 12:86 DOI 10.1186/1471-2229-12-86.
- Da Lage JL. 2018. The amylases of insects. *International Journal of Insect Science* 10:1–14.
- Dai L, Gao H, Ye J, Fu D, Sun Y, Chen H. 2019. Isolation of *CarE* genes from the Chinese white pine beetle *Dendroctonus armandi* (Curculionidae: Scolytinae) and their response to host chemical defense. *Pest Management Science* 75:986–997 DOI 10.1002/ps.5205.
- Damle MS, Giri AP, Sainani MN, Gupta VS. 2005. Higher accumulation of proteinase inhibitors in flowers than leaves and fruits as a possible basis for differential feeding preference of *Helicoverpa armigera* on tomato (*Lycopersicon esculentum* Mill, Cv. Dhanashree). *Phytochemistry* 66:2659–2667 DOI 10.1016/j.phytochem.2005.09.006.
- Dangl JL, Jones JDG. 2001. Plant pathogens and integrated defense responses to infection. *Nature* 411:826–833 DOI 10.1038/35081161.
- Davey MW, Montagu Mv, Inze D, Sanmartin M, Kanellis A, Smirnoff N, Benzie IJJ, Strain JJ, Favell D, Fletcher F. 2000. Plant L-ascorbic acid: chemistry, function, metabolism, bioavailability and effects of processing. *Journal of the Science of Food and Agriculture* 80:825–860 DOI 10.1002/(SICI)1097-0010(20000515)80:7<825::AID-JSFA598>3.0.CO;2-6.
- De Oliveira DM, Finger-Teixeira A, Rodrigues Mota T, Salvador VH, Moreira-Vilar FC, Correa Molinari HB, Dantas dos Santos W. 2015. Ferulic acid: a key component in grass lignocellulose recalcitrance to hydrolysis. *Plant Biotechnology Journal* 13:1224–1232 DOI 10.1111/pbi.12292.
- De Oliveira EF, Pallini A, Janssen A. 2019. Herbivore performance and plant defense after sequential attacks by inducing and suppressing herbivores. *Insect Science* 26:108–118 DOI 10.1111/1744-7917.12499.
- Dean M, Hamon Y, Chimini G. 2001. The human ATP-binding cassette (ABC) transporter superfamily. *Journal of Lipid Research* 42:1007–1017 DOI 10.1016/S0022-2275(20)31588-1.
- Deng Y, Lu S. 2017. Biosynthesis and regulation of phenylpropanoids in plants. *Critical Reviews in Plant Sciences* 36:257–290 DOI 10.1080/07352689.2017.1402852.
- Díaz J, Ten Have A, Van Kan JA. 2002. The role of ethylene and wound signaling in resistance of tomato to *Botrytis cinerea*. *Plant Physiology* 129:1341–1351 DOI 10.1104/pp.001453.
- Díaz-Fleischer F, Aluja M. 2003. Clutch size in frugivorous insects as a function of host firmness: the case of the tephritid fly *Anastrepha ludens*. *Ecological Entomology* 28:268–277 DOI 10.1046/j.1365-2311.2003.00517.
- Dixit G, Srivastava A, Rai KM, Dubey RS, Srivastava R, Verma PC. 2020. Distinct defensive activity of phenolics and phenylpropanoid pathway genes in different

- cotton varieties toward chewing pests. *Plant Signaling & Behavior* **15**:1747689
DOI 10.1080/15592324.2020.1747689.
- Do CT, Pollet B, Thévenin J, Sibout R, Denoue D, Barrièr Y, Lapierre C, Jouanin L. 2007.** Both caffeoyl Coenzyme A 3-O-methyltransferase 1 and caffeic acid O-methyltransferase 1 are involved in redundant functions for lignin, flavonoids and sinapoyl malate biosynthesis in Arabidopsis. *Planta* **226**:1117–1129
DOI 10.1007/s00425-007-0558-3.
- Donkor D, Mirzahosseini Z, Bede J, Bauce E, Despland E. 2019.** Detoxification of host plant phenolic aglycones by the spruce budworm. *PLOS ONE* **145**:e0208288
DOI 10.1371/journal.pone.0208288.
- Enayati AA, Ranson H, Hemingway J. 2005.** Insect glutathione transferases and insecticide resistance. *Insect Molecular Biology* **14**:3–8
DOI 10.1111/j.1365-2583.2004.00529.x.
- Enya S, Daimon T, Igarashi F, Kataoka H, Uchibori M, Sezutsu H, Shinoda T, Niwa R. 2015.** The silkworm glutathione S-transferase gene *noppera-bo* is required for ecdysteroid biosynthesis and larval development. *Insect Biochemistry and Molecular Biology* **61**:1–7 DOI 10.1016/j.ibmb.2015.04.001.
- Fan Y, Yang W, Yan Q, Chen C, Li J. 2020.** Genome-wide identification and expression Analysis of the protease inhibitor gene families in tomato. *Genes* **11**:1
DOI 10.3390/genes11010001.
- Farmer EE, Ryan CA. 1992.** Octadecanoid precursors of jasmonic acid activate the synthesis of wound-inducible proteinase inhibitors. *The Plant Cell* **4**:129–134
DOI 10.1105/tpc.4.2.129.
- Farnsworth CA, Teese MG, Yuan G, Li Y, Scott C, Zhang X, Wu Y, Russell RJ, Oakeshott JG. 2010.** Esterase-based metabolic resistance to insecticides in heliothine and spodopteran pests. *Journal of Pesticide Science* **35**:275–289
DOI 10.1584/jpestics.R10-13.
- Felton GW. 2005.** Indigestion is a plant's best defense. *Proceedings of the National Academy of Sciences of the United States of America* **102**:18771–18772
DOI 10.1073/pnas.0509895102.
- Feng X, Li M, Liu N. 2018.** Carboxylesterase genes in pyrethroid resistant house flies, *Musca domestica*. *Insect Biochemistry and Molecular Biology* **92**:30–39
DOI 10.1016/j.ibmb.2017.11.007.
- Ferrie BJ, Beaudoin N, Burkhardt W, Bowsher CG, Rothstein SJ. 1994.** The cloning of two tomato lipoxygenase genes and their differential expression during fruit ripening. *Plant Physiology* **106**:109–118 DOI 10.1104/pp.106.1.109.
- Feyereisen R. 2006.** Evolution of insect P450. *Biochemical Society Transactions* **34**:1252–1255 DOI 10.1042/bst0341252.
- Fitt GP. 1984.** Oviposition behaviour of two tephritid fruit flies, *Dacus tryoni* and *Dacus jarvisi*, as influenced by the presence of larvae in the host fruit. *Oecologia* **62**:37–46
DOI 10.1007/BF00377370.
- Flors V, Leyva MdO, Vicedo B, Finiti I, Real MD, García-Agustín P, Bennett AB, González-Bosch C. 2007.** Absence of the endo- β -1, 4-glucanases Cel1 and Cel2

- reduces susceptibility to *Botrytis cinerea* in tomato. *The Plant Journal* **52**:1027–1040 DOI [10.1111/j.1365-313X.2007.03299.x](https://doi.org/10.1111/j.1365-313X.2007.03299.x).
- Franco OL, Rigden DJ, Melo FR, Grossi-de Sá MF. 2002.** Plant α -amylase inhibitors and their interaction with insect α -amylases: Structure, function and potential for crop protection. *European Journal of Biochemistry* **269**:397–412 DOI [10.1046/j.0014-2956.2001.02656.x](https://doi.org/10.1046/j.0014-2956.2001.02656.x).
- George A, Rao C, Rahangadale SJ. 2019.** Current status of insecticide resistance in *Aphis gossypii* and *Aphis spiraecola* (Hemiptera: Aphididae) under central Indian conditions in citrus. *Cogent Biology* **5**:1660494 DOI [10.1080/23312025.2019.1660494](https://doi.org/10.1080/23312025.2019.1660494).
- Gerszberg A, Hnatuszko-Konka K, Kowalczyk T, Kononowicz AK. 2015.** Tomato (*Solanum lycopersicum* L.) in the service of biotechnology. *Plant Cell, Tissue and Organ Culture* **120**:881–902 DOI [10.1007/s11240-014-0664-4](https://doi.org/10.1007/s11240-014-0664-4).
- Gilardoni PA, Hettenhausen C, Baldwin IT, Bonaventure G. 2011.** Nicotiana attenuata LECTIN RECEPTOR KINASE1 suppresses the insect-mediated inhibition of induced defense responses during *Manduca sexta* herbivory. *The Plant Cell* **23**:3512–3532 DOI [10.1105/tpc.111.088229](https://doi.org/10.1105/tpc.111.088229).
- Gilchrist AS, Shearman DC, Frommer M, Raphael KA, Deshpande NP, Wilkins MR, Sherwin WB, Sved JA. 2014.** The draft genome of the pest tephritid fruit fly *Bactrocera tryoni*: resources for the genomic analysis of hybridising species. *BMC Genomics* **15**:1153 DOI [10.1186/1471-2164-15-1153](https://doi.org/10.1186/1471-2164-15-1153).
- Giron D, Dubreuil G, Bennett A, Dedeine F, Dicke M, Dyer LA, Erb M, Harris MO, Huguet E, Kaloshian I. 2018.** Promises and challenges in insect–plant interactions. *Entomologia Experimentalis et Applicata* **166**:319–343 DOI [10.1111/eea.12679](https://doi.org/10.1111/eea.12679).
- Gloss AD, Abbot P, Whiteman NK. 2019.** How interactions with plant chemicals shape insect genomes. *Current Opinion in Insect Science* **36**:149–156 DOI [10.1016/j.cois.2019.09.005](https://doi.org/10.1016/j.cois.2019.09.005).
- Gonzales-Vigil E, Bianchetti CM, Phillips GN, Howe GA. 2011.** Adaptive evolution of threonine deaminase in plant defense against insect herbivores. *Proceedings of the National Academy of Sciences of the United States of America* **108**:5897–5902 DOI [10.1073/pnas.1016157108](https://doi.org/10.1073/pnas.1016157108).
- Gouhier-Darimont C, Schmiesing A, Bonnet C, Lassueur S, Reymond P. 2013.** Signalling of *Arabidopsis thaliana* response to *Pieris brassicae* eggs shares similarities with PAMP-triggered immunity. *Journal of Experimental Botany* **64**:665–674 DOI [10.1093/jxb/ers362](https://doi.org/10.1093/jxb/ers362).
- Gouhier-Darimont C, Stahl E, Glauser G, Reymond P. 2019.** The *Arabidopsis* lectin receptor kinase LecRK-I.8 is involved in insect egg perception. *Frontiers in Plant Science* **10**:623 DOI [10.3389/fpls.2019.00623](https://doi.org/10.3389/fpls.2019.00623).
- Grasso F, Coppola M, Carbone F, Baldoni L, Alagna F, Perrotta G, Pérez-Pulido AJ, Garonna A, Facella P, Daddiego L. 2017.** The transcriptional response to the olive fruit fly (*Bactrocera oleae*) reveals extended differences between tolerant and susceptible olive (*Olea europaea* L.) varieties. *PLOS ONE* **12**:e0183050 DOI [10.1371/journal.pone.0183050](https://doi.org/10.1371/journal.pone.0183050).

- Hafsi A, Facon B, Ravigné V, Chiroleu F, Quilici S, Chermiti B, Duyck P-F. 2016. Host plant range of a fruit fly community (Diptera: Tephritidae): does fruit composition influence larval performance? *BMC Ecology* 16:40 DOI 10.1186/s12898-016-0094-8.
- Haile ZM, Guzman EGNagpala-De, Moretto M, Sonogo P, Engelen K, Zoli L, Moser C, Baraldi E. 2019. Transcriptome profiles of strawberry (*Fragaria vesca*) fruit interacting with *Botrytis cinerea* at different ripening stages. *Frontiers in Plant Science* 10:1131 DOI 10.3389/fpls.2019.01131.
- Hamza R, Pérez-Hedo M, Urbaneja A, Rambla JL, Granell A, Gaddour K, Beltrán JP, Cañas LA. 2018. Expression of two barley proteinase inhibitors in tomato promotes endogenous defensive response and enhances resistance to *Tuta absoluta*. *BMC Plant Biology* 18:24 DOI 10.1186/s12870-018-1240-6.
- Hasanuzzaman M, Bhuyan M, Zulfiqar F, Raza A, Mohsin SM, Mahmud JA, Fujita M, Fotopoulos V. 2020. Reactive oxygen species and antioxidant defense in plants under abiotic stress: revisiting the crucial role of a universal defense regulator. *Antioxidants* 9:681 DOI 10.3390/antiox9080681.
- Hashemi L, Golparvar AR, Nasr-Esfahani M, Golabadi M. 2020. Expression analysis of defense-related genes in cucumber (*Cucumis sativus* L.) against *Phytophthora melonis*. *Molecular Biology Reports* 47:4933–4944 DOI 10.1007/s11033-020-05520-5.
- Hayes JD, Flanagan JU, Jowsey IR. 2005. Glutathione transferases. *Annual Review of Pharmacology and Toxicology* 45:51–88 DOI 10.1146/annurev.pharmtox.45.120403.095857.
- Hazzouri KM, Sudalaimuthuasari N, Kundu B, Nelson D, Al-Deeb MA, Le Mansour A, Spencer JJ, Desplan C, Amiri KMA. 2020. The genome of pest *Rhynchophorus ferrugineus* reveals gene families important at the plant-beetle interface. *Communications Biology* 3:323 DOI 10.1038/s42003-020-1060-8.
- Heckel DG. 2018. Insect detoxification and sequestration strategies. *Annual Plant Reviews Online* 47:77–114 DOI 10.1002/9781119312994.apr0507.
- Heidel-Fischer HM, Vogel H. 2015. Molecular mechanisms of insect adaptation to plant secondary compounds. *Current Opinion in Insect Science* 8:8–14 DOI 10.1016/j.cois.2015.02.004.
- Heitz T, Bergey DR, Ryan CA. 1997. A gene encoding a chloroplast-targeted lipoxygenase in tomato leaves is transiently induced by wounding, systemin, and methyl jasmonate. *Plant Physiology* 114:1085–1093 DOI 10.1104/pp.114.3.1085.
- Higgins R, Lockwood T, Holley S, Yalamanchili R, Stratmann JW. 2007. Changes in extracellular pH are neither required nor sufficient for activation of mitogen-activated protein kinases (MAPKs) in response to systemin and fusicoccin in tomato. *Planta* 225:1535–1546 DOI 10.1007/s00425-006-0440-8.
- Holley SR, Yalamanchili RD, Moura DS, Ryan CA, Stratmann JW. 2003. Convergence of signaling pathways induced by systemin, oligosaccharide elicitors, and ultraviolet-B radiation at the level of mitogen-activated protein kinases in *Lycopersicon peruvianum* suspension-cultured cells. *Plant Physiology* 132:1728–1738 DOI 10.1104/pp.103.024414.

- Howe GA, Jander G. 2008.** Plant immunity to insect herbivores. *Annual Review of Plant Biology* **59**:41–66 DOI [10.1146/annurev.arplant.59.032607.092825](https://doi.org/10.1146/annurev.arplant.59.032607.092825).
- Huang Y, Jiang H-B, Shen G-M, Dou W, Wang J-J. 2012.** Molecular characterizations of two cytochrome P450 genes encoding and *CYP6EK1* from the oriental fruit fly, *Bactrocera dorsalis* (Diptera: Tephritidae). *Archives of Insect Biochemistry and Physiology* **79**:31–46 DOI [10.1002/arch.21003](https://doi.org/10.1002/arch.21003).
- Huang Y, Shen G-M, Jiang H-B, Jiang X-Z, Dou W, Wang J-J. 2013.** Multiple P450 genes: Identification, tissue-specific expression and their responses to insecticide treatments in the oriental fruit fly, *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae). *Pesticide Biochemistry and Physiology* **106**:1–7 DOI [10.1016/j.pestbp.2013.03.001](https://doi.org/10.1016/j.pestbp.2013.03.001).
- Huang Y, Xu Z, Lin X, Feng Q, Zheng S. 2011.** Structure and expression of glutathione S-transferase genes from the midgut of the Common cutworm, *Spodoptera litura* (Noctuidae) and their response to xenobiotic compounds and bacteria. *Journal of Insect Physiology* **57**:1033–1044 DOI [10.1016/j.jinsphys.2011.05.001](https://doi.org/10.1016/j.jinsphys.2011.05.001).
- Ishikawa T, Shigeoka S. 2008.** Recent advances in ascorbate biosynthesis and the physiological significance of ascorbate peroxidase in photosynthesizing organisms. *Bioscience, Biotechnology, and Biochemistry* **72**:1143–1154 DOI [10.1271/bbb.80062](https://doi.org/10.1271/bbb.80062).
- Jing TX, Wang DF, Ma YP, Zeng LL, Meng LW, Zhang Q, Dou W, Wang JJ. 2020.** Genome-wide and expression-profiling analyses of the cytochrome P450 genes in *Bactrocera dorsalis* (Hendel) and screening of candidate P450 genes associated with malathion resistance. *Pest Management Science* **76**:2932–2943 DOI [10.1002/ps.5891](https://doi.org/10.1002/ps.5891).
- Johnson R, Narvaez J, An G, Ryan C. 1989.** Expression of proteinase inhibitors I and II in transgenic tobacco plants: effects on natural defense against *Manduca sexta* larvae. *Proceedings of the National Academy of Sciences of the United States of America* **86**:9871–9875 DOI [10.1073/pnas.86.24.9871](https://doi.org/10.1073/pnas.86.24.9871).
- Kai K, Mizutani M, Kawamura N, Yamamoto R, Tamai M, Yamaguchi H, Sakata K, Shimizu BI. 2008.** Scopoletin is biosynthesized via *ortho*-hydroxylation of feruloyl CoA by a 2-oxoglutarate-dependent dioxygenase in *Arabidopsis thaliana*. *The Plant Journal* **55**:989–999 DOI [10.1111/j.1365-313X.2008.03568.x](https://doi.org/10.1111/j.1365-313X.2008.03568.x).
- Kampatsikas I, Bijelic A, Rompel A. 2019.** Biochemical and structural characterization of tomato polyphenol oxidases provide novel insights into their substrate specificity. *Scientific Reports* **9**:1–13 DOI [10.1038/s41598-019-39687-0](https://doi.org/10.1038/s41598-019-39687-0).
- Kandath PK, Ranf S, Pancholi SS, Jayanty S, Walla MD, Miller W, Howe GA, Lincoln DE, Stratmann JW. 2007.** Tomato MAPKs *LeMPK1*, *LeMPK2*, and *LeMPK3* function in the systemin-mediated defense response against herbivorous insects. *Proceedings of the National Academy of Sciences of the United States of America* **104**:12205–12210 DOI [10.1073/pnas.0700344104](https://doi.org/10.1073/pnas.0700344104).
- Kausch KD, Sobolev AP, Goyal RK, Fatima T, Laila-Beevi R, Saftner RA, Handa AK, Mattoo AK. 2012.** Methyl jasmonate deficiency alters cellular metabolome, including the aminome of tomato (*Solanum lycopersicum* L.) fruit. *Amino Acids* **42**:843–856 DOI [10.1007/s00726-011-1000-5](https://doi.org/10.1007/s00726-011-1000-5).

- Kawazu K, Mochizuki A, Sato Y, Sugeno W, Murata M, Seo S, Mitsuhara I. 2012.** Different expression profiles of jasmonic acid and salicylic acid inducible genes in the tomato plant against herbivores with various feeding modes. *Arthropod-Plant Interactions* **6**:221–230 DOI [10.1007/s11829-011-9174-z](https://doi.org/10.1007/s11829-011-9174-z).
- Kessler A, Baldwin IT. 2002.** Plant responses to insect herbivory: the emerging molecular analysis. *Annual Review of Plant Biology* **53**:299–328 DOI [10.1146/annurev.arplant.53.100301.135207](https://doi.org/10.1146/annurev.arplant.53.100301.135207).
- Klauser D, Desurmont GA, Glauser G, Vallat A, Flury P, Boller T, Turlings TC, Bartels SJ. 2015.** The *Arabidopsis* Pep-PEPR system is induced by herbivore feeding and contributes to JA-mediated plant defense against herbivory. *Journal of Experimental Botany* **66**:5327–5336 DOI [10.1093/jxb/erv250](https://doi.org/10.1093/jxb/erv250).
- Kumar K, Gambhir G, Dass A, Tripathi AK, Singh A, Jha AK, Yadava P, Choudhary M, Rakshit S. 2020.** Genetically modified crops: current status and future prospects. *Planta* **251**:1–27 DOI [10.1007/s00425-020-03372-8](https://doi.org/10.1007/s00425-020-03372-8).
- Kumaran N, Van der Burg CA, Qin Y, Cameron SL, Clarke AR, Prentis PJ. 2018.** Plant-mediated female transcriptomic changes post-mating in a tephritid fruit fly, *Bactrocera tryoni*. *Genome Biology and Evolution* **10**:94–107 DOI [10.1093/gbe/evx257](https://doi.org/10.1093/gbe/evx257).
- Li W, Berenbaum M, Schuler M. 2001.** Molecular analysis of multiple *CYP6B* genes from polyphagous *Papilio* species. *Insect Biochemistry and Molecular Biology* **31**:999–1011 DOI [10.1016/S0965-1748\(01\)00048-0](https://doi.org/10.1016/S0965-1748(01)00048-0).
- Li T, Liu N. 2019.** Role of the G-Protein-Coupled receptor signaling pathway in insecticide resistance. *International Journal of Molecular Sciences* **20**:4300 DOI [10.3390/ijms20174300](https://doi.org/10.3390/ijms20174300).
- Lianthanzauva D, Koramutla MK, Subramanian S, Rohit C, Ramcharan B. 2020.** Comparative transcriptomics revealed differential regulation of defense related genes in *Brassica juncea* leading to successful and unsuccessful infestation by aphid species. *Scientific Reports* **10**:10583 DOI [10.1038/s41598-020-66217-0](https://doi.org/10.1038/s41598-020-66217-0).
- Liu J. 2018.** Temporal and stage-specific variation in mite-induced responses of tomato plants.. Doctoral dissertation, University of Amsterdam, Netherlands Available at <https://pure.uva.nl/ws/files/28145400/1.pdf>.
- Liu J, Osbourn A, Ma P. 2015.** MYB transcription factors as regulators of phenylpropanoid metabolism in plants. *Molecular Plant* **8**:689–708 DOI [10.1016/j.molp.2015.03.012](https://doi.org/10.1016/j.molp.2015.03.012).
- Liu Z, Wu Y, Yang F, Zhang Y, Chen S, Xie Q, Tian X, Zhou J-M. 2013.** BIK1 interacts with PEPRs to mediate ethylene-induced immunity. *Proceedings of the National Academy of Sciences of the United States of America* **110**:6205–6210 DOI [10.1073/pnas.1215543110](https://doi.org/10.1073/pnas.1215543110).
- Liu S, Zhou S, Tian L, Guo E, Luan Y, Zhang J, Li S. 2011.** Genome-wide identification and characterization of ATP-binding cassette transporters in the silkworm, *Bombyx mori*. *BMC Genomics* **12**:491 DOI [10.4238/2011.October.25.9](https://doi.org/10.4238/2011.October.25.9).
- López-Ráez JA, Flors V, García JM, Pozo MJ. 2010.** AM symbiosis alters phenolic acid content in tomato roots. *Plant Signaling & Behavior* **5**:1138–1140 DOI [10.4161/psb.5.9.12659](https://doi.org/10.4161/psb.5.9.12659).

- Lorence A, Chevone BI, Mendes P, Nessler CL. 2004.** Myo-inositol oxygenase offers a possible entry point into plant ascorbate biosynthesis. *Plant Physiology* 134:1200–1205 DOI 10.1104/pp.103.033936.
- MacGregor KB, Shelp BJ, Peiris S, Bown AW. 2003.** Overexpression of glutamate decarboxylase in transgenic tobacco plants deters feeding by phytophagous insect larvae. *Journal of Chemical Ecology* 29:2177–2182 DOI 10.1023/A:1025650914947.
- Macho AP, Zipfel C. 2014.** Plant PRRs and the activation of innate immune signaling. *Molecular Cell* 54:263–272 DOI 10.1016/j.molcel.2014.03.028.
- Mahanil S, Attajarusit J, Stout MJ, Thipyapong P. 2008.** Overexpression of tomato polyphenol oxidase increases resistance to common cutworm. *Plant Science* 174:456–466 DOI 10.1016/j.plantsci.2008.01.006.
- Mandal SM, Chakraborty D, Dey S. 2010.** Phenolic acids act as signaling molecules in plant-microbe symbioses. *Plant Signaling & Behavior* 5:359–368 DOI 10.4161/psb.5.4.10871.
- Mariutto M, Duby F, Adam A, Bureau C, Fauconnier M-L, Ongena M, Thonart P, Dommes J. 2011.** The elicitation of a systemic resistance by *Pseudomonas putida* BTP1 in tomato involves the stimulation of two lipoxygenase isoforms. *BMC Plant Biology* 11:1–15 DOI 10.1186/1471-2229-11-29.
- Marshall SD, Putterill JJ, Plummer KM, Newcomb RD. 2003.** The carboxylesterase gene family from *Arabidopsis thaliana*. *Journal of Molecular Evolution* 57:487–500 DOI 10.1007/s00239-003-2492-8.
- Martin GB, Bogdanove AJ, Sessa G. 2003.** Understanding the functions of plant disease resistance proteins. *Annual Review of Plant Biology* 54:23–61 DOI 10.1146/annurev.arplant.54.031902.135035.
- Medjkouh L, Costa A, Tamendjari A, Bekdouche F, Bouarroudj K, Oliveira MBPP. 2018.** Susceptibility of eight Algerian olive cultivars to *Bactrocera oleae* infestation –a pomological and nutritional quality perspective. *Phytoparasitica* 46:595–605 DOI 10.1007/s12600-018-0697-z.
- Meng L-W, Peng M-L, Chen M-L, Yuan G-R, Zheng L-S, Bai W-J, Smagghe G, Wang J-J. 2020.** A glutathione S-transferase (*BdGSTd9*) participates in malathion resistance via directly depleting malathion and its toxic oxide malaoxon in *Bactrocera dorsalis* (Hendel). *Pest Management Science* 76:2557–2568 DOI 10.1002/ps.5810.
- Miao L, Shou S, Zhu Z, Jiang F, Zai W, Yang Y. 2008.** Isolation of a novel tomato caffeoyl CoA 3-O-methyltransferase gene following infection with the bacterium *Ralstonia solanacearum*. *Journal of Phytopathology* 156:588–596 DOI 10.1111/j.1439-0434.2008.01406.x.
- Mittapalli O, Neal JJ, Shukle RH. 2007.** Tissue and life stage specificity of glutathione S-transferase expression in the Hessian fly, *Mayetiola destructor*: implications for resistance to host allelochemicals. *Journal of Insect Science* 7:20 DOI 10.1673/031.007.2001.
- Mittler R, Vanderauwera S, Gollery M, Van Breusegem F. 2004.** Reactive oxygen gene network of plants. *Trends in Plant Science* 9:490–498 DOI 10.1016/j.tplants.2004.08.009.

- Moon H, Lee B, Choi G, Shin D, Prasad DT, Lee O, Kwak S-S, Kim DH, Nam J, Bahk J. 2003.** NDP kinase 2 interacts with two oxidative stress-activated MAPKs to regulate cellular redox state and enhances multiple stress tolerance in transgenic plants. *Proceedings of the National Academy of Sciences of the United States of America* **100**:358–363 DOI [10.1073/pnas.252641899](https://doi.org/10.1073/pnas.252641899).
- Müller R, De Vos M, Sun JY, Sønderby IE, Halkier BA, Wittstock U, Jander G. 2010.** Differential effects of indole and aliphatic glucosinolates on lepidopteran herbivores. *Journal of Chemical Ecology* **36**:905–913 DOI [10.1007/s10886-010-9825-z](https://doi.org/10.1007/s10886-010-9825-z).
- National Center for Biotechnology Information. 1988.** Available at <https://www.ncbi.nlm.nih.gov/> (accessed on 18 June 2020).
- Neill SJ, Desikan R, Clarke A, Hurst RD, Hancock JT. 2002.** Hydrogen peroxide and nitric oxide as signalling molecules in plants. *Journal of Experimental Botany* **53**:1237–1247 DOI [10.1093/jexbot/53.372.1237](https://doi.org/10.1093/jexbot/53.372.1237).
- Nejat N, Mantri N. 2017.** Plant immune system: crosstalk between responses to biotic and abiotic stresses the missing link in understanding plant defense. *Molecular Biology* **23**:1–16 DOI [10.21775/cimb.023.001](https://doi.org/10.21775/cimb.023.001).
- Newman SM, Eannetta NT, Yu H, Prince JP, De Vicente MC, Tanksley SD, Steffens JC. 1993.** Organisation of the tomato polyphenol oxidase gene family. *Plant Molecular Biology* **21**:1035–1051 DOI [10.1007/BF00023601](https://doi.org/10.1007/BF00023601).
- Nombela G, Williamson VM, Muñiz M. 2003.** The root-knot nematode resistance gene *Mi-1.2* of tomato is responsible for resistance against the whitefly *Bemisia tabaci*. *Molecular Plant-Microbe Interactions* **16**:645–649 DOI [10.1094/MPMI.2003.16.7.645](https://doi.org/10.1094/MPMI.2003.16.7.645).
- Nunes MZ, Boff M, Santos Rdos, Franco C, Wille P, Rosa Jda. 2015.** Damage and development of *Anastrepha fraterculus* (Diptera: Tephritidae) in fruits of two pear cultivars. *Agrociencia Uruguay* **19**:42–48.
- Orozco-Cardenas M, McGurl B, Ryan CA. 1993.** Expression of an antisense prosystemin gene in tomato plants reduces resistance toward *Manduca sexta* larvae. *Proceedings of the National Academy of Sciences of the United States of America* **90**:8273–8276 DOI [10.1073/pnas.90.17.8273](https://doi.org/10.1073/pnas.90.17.8273).
- Ortego F, Novillo C, Sanchez-Serrano J, Castañera P. 2001.** Physiological response of Colorado potato beetle and beet armyworm larvae to depletion of wound-inducible proteinase inhibitors in transgenic potato plants. *Journal of Insect Physiology* **47**:1291–1300 DOI [10.1016/S0022-1910\(01\)00118-4](https://doi.org/10.1016/S0022-1910(01)00118-4).
- Papachristos D, Kimbaris A, Papadopoulou N, Polissiou M. 2009.** Toxicity of citrus essential oils against *Ceratitis capitata* (Diptera: Tephritidae) larvae. *Annals of Applied Biology* **155**:381–389 DOI [10.1111/j.1744-7348.2009.00350.x](https://doi.org/10.1111/j.1744-7348.2009.00350.x).
- Papachristos DP, Papadopoulou NT, Nanos GD. 2008.** Survival and development of immature stages of the Mediterranean fruit fly (Diptera: Tephritidae) in citrus fruit. *Journal of Economic Entomology* **101**:866–872 DOI [10.1093/jee/101.3.866](https://doi.org/10.1093/jee/101.3.866).
- Park Y-S, Kunze S, Ni X, Feussner I, Kolomiets MV. 2010.** Comparative molecular and biochemical characterization of segmentally duplicated 9-lipoxygenase genes *ZmLOX4* and *ZmLOX5* of maize. *Planta* **231**:1425–1437 DOI [10.1007/s00425-010-1143-8](https://doi.org/10.1007/s00425-010-1143-8).

- Pavliidi N, Dermauw W, Rombauts S, Chrisargiris A, Van Leeuwen T, Vontas J. 2013. Analysis of the olive fruit fly *Bactrocera oleae* transcriptome and phylogenetic classification of the major detoxification gene families. *PLOS ONE* 8:e66533 DOI 10.1371/journal.pone.0066533.
- Pearce G, Johnson S, Ryan CA. 1993. Purification and characterization from tobacco (*Nicotiana tabacum*) leaves of six small, wound-inducible, proteinase iso-inhibitors of the potato inhibitor II family. *Plant Physiology* 102:639–644 DOI 10.1104/pp.102.2.639.
- Porta H, Rocha-Sosa M. 2002. Plant lipoxygenases. Physiological and molecular features. *Plant Physiology* 130:15–21 DOI 10.1104/pp.010787.
- Prasch CM, Sonnewald U. 2013. Simultaneous application of heat, drought, and virus to *Arabidopsis* plants reveals significant shifts in signaling networks. *Plant Physiology* 162:1849–1866 DOI 10.1104/pp.113.221044.
- Qin Y-j, Krosch MN, Schutze MK, Zhang Y, Wang X-x, Prabhakar CS, Susanto A, Hee AKW, Ekesi S, Badji K, Khan M, Wu J-j, Wang Q-l, Yan G, Zhu L-h, Zhao Z-h, Liu L-j, Clarke AR, Li Z-h. 2018. Population structure of a global agricultural invasive pest, *Bactrocera dorsalis* (Diptera: Tephritidae). *Evolutionary Applications* 11:1990–2003 DOI 10.1111/eva.12701.
- Qin Y, Paini DR, Wang C, Fang Y, Li Z. 2015. Global establishment risk of economically important fruit fly species (Tephritidae). *PLOS ONE* 10:e0116424 DOI 10.1371/journal.pone.0116424.
- Quais MK, Munawar A, Ansari NA, Zhou W-W, Zhu Z-R. 2020. Interactions between brown planthopper (*Nilaparvata lugens*) and salinity stressed rice (*Oryza sativa*) plant are cultivar-specific. *Scientific Reports* 10:1–14 DOI 10.1038/s41598-020-64925-1.
- Ramos-Ruiz R, Martinez F, Knauf-Beiter G. 2019. The effects of GABA in plants. *Cogent Food & Agriculture* 5:1670553 DOI 10.1080/23311932.2019.1670553.
- Rao S, Nandineni MR. 2017. Genome sequencing and comparative genomics reveal a repertoire of putative pathogenicity genes in chilli anthracnose fungus *Colletotrichum truncatum*. *PLOS ONE* 12:e0183567 DOI 10.1371/journal.pone.0183567.
- Rattanapun W, Amornsak W, Clarke AR. 2009. *Bactrocera dorsalis* preference for and performance on two mango varieties at three stages of ripeness. *Entomologia Experimentalis et Applicata* 131:243–253 DOI 10.1007/s10529-017-2298-1.
- Rehman S, Aziz E, Akhtar W, Ilyas M, Mahmood T. 2017. Structural and functional characteristics of plant proteinase inhibitor-II (PI-II) family. *Biotechnology Letters* 39:647–666 DOI 10.1007/s10529-017-2298-1.
- Renault H, Amrani AEI, Palanivelu R, Updegraff EP, Yu A, Renou JP, Preuss D, Bouchereau A, Deleu C. 2011. GABA accumulation causes cell elongation defects and a decrease in expression of genes encoding secreted and cell wall-related proteins in *Arabidopsis thaliana*. *Plant and Cell Physiology* 52:894–908 DOI 10.1093/pcp/pcr041.
- Ricroch A. 2019. Global developments of genome editing in agriculture. *Transgenic Research* 28:45–52 DOI 10.1007/s11248-019-00133-6.

- Rodriguez-Saona CR, Musser RO, Vogel H, Hum-Musser SM, Thaler JS. 2010.** Molecular, biochemical, and organismal analyses of tomato plants simultaneously attacked by herbivores from two feeding guilds. *Journal of Chemical Ecology* **36**:1043–1057 DOI [10.1007/s10886-010-9854-7](https://doi.org/10.1007/s10886-010-9854-7).
- Roohigohar S, Prentis PJ, Clarke AR. 2020.** Effect of tomato fruit cultivar and ripening stage on *Bactrocera tryoni* (Froggatt) egg and larval survival. *Journal of Applied Entomology* **144**:797–805 DOI [10.1111/jen.12813](https://doi.org/10.1111/jen.12813).
- Rösner J, Merzendorfer H. 2020.** Transcriptional plasticity of different ABC transporter genes from *Tribolium castaneum* contributes to diflubenzuron resistance. *Insect Biochemistry and Molecular Biology* **116**:103282 DOI [10.1016/j.ibmb.2019.103282](https://doi.org/10.1016/j.ibmb.2019.103282).
- Roth CW, Holm I, Graille M, Dehoux P, Rzhetsky A, Wincker P, Weissenbach J, Brey PT. 2003.** Identification of the *Anopheles gambiae* ATP-binding cassette transporter superfamily genes. *Molecules and Cells* **15**:150–158.
- Ryan JD, Gregory P, Tingey WM. 1982.** Phenolic oxidase activities in glandular trichomes of *Solanum berthaultii*. *Phytochemistry* **21**:1885–1887 DOI [10.1016/0031-9422\(82\)83008-2](https://doi.org/10.1016/0031-9422(82)83008-2).
- Ryan CA, Pearce G. 2003.** Systemins: a functionally defined family of peptide signals that regulate defensive genes in *Solanaceae* species. *Proceedings of the National Academy of Sciences of the United States of America* **100**:14577–14580 DOI [10.1073/pnas.1934788100](https://doi.org/10.1073/pnas.1934788100).
- Saha D. 2016.** Biochemical insecticide resistance in tea pests. In: Trdan S, ed. *Insecticides resistance*. Croatia: InTech, 347–390 DOI [10.5772/61949](https://doi.org/10.5772/61949).
- Salim AP, Saminaidu K, Marimuthu M, Perumal Y, Rethinasamy V, Palanisami JR, Vadivel K. 2011.** Defense responses in tomato landrace and wild genotypes to early blight pathogen *Alternaria solani* infection and accumulation of pathogenesis-related proteins. *Archives of Phytopathology and Plant Protection* **44**:1147–1164 DOI [10.1080/03235408.2010.482763](https://doi.org/10.1080/03235408.2010.482763).
- Sánchez-Hernández C, Martínez-Gallardo N, Guerrero-Rangel A, Valdés-Rodríguez S, Délano-Frier J. 2004.** Trypsin and α -amylase inhibitors are differentially induced in leaves of amaranth (*Amaranthus hypochondriacus*) in response to biotic and abiotic stress. *Physiologia Plantarum* **122**:254–264 DOI [10.1111/j.0031-9317.2004.00398.x](https://doi.org/10.1111/j.0031-9317.2004.00398.x).
- Santamaria ME, Arnaiz A, Gonzalez-Melendi P, Martinez M, Diaz I. 2018.** Plant perception and short-term responses to phytophagous insects and mites. *International Journal of Molecular Sciences* **19**:1356 DOI [10.3390/ijms19051356](https://doi.org/10.3390/ijms19051356).
- Santamaria ME, Martínez M, Cambra I, Grbic V, Diaz I. 2013.** Understanding plant defense responses against herbivore attacks: an essential first step towards the development of sustainable resistance against pests. *Transgenic Research* **22**:697–708 DOI [10.1007/s11248-013-9725-4](https://doi.org/10.1007/s11248-013-9725-4).
- Sarwar M. 2015.** Genetic control tactic against fruit flies (Diptera: Tephritidae) insect to escape destruction of perishable horticulture crops. *International Journal of Animal Biology* **1**:209–214.
- Scholz SS, Reichelt M, Mekonnen DW, Ludewig F, Mithöfer A. 2015.** Insect herbivory-elicited GABA accumulation in plants is a wound-induced, direct, systemic,

- and jasmonate-independent defense response. *Frontiers in Plant Science* **6**:1128 DOI [10.3389/fpls.2015.01128](https://doi.org/10.3389/fpls.2015.01128).
- Seal AN, Pratley JE, Haig T, An M. 2004. Identification and quantitation of compounds in a series of allelopathic and non-allelopathic rice root exudates. *Journal of Chemical Ecology* **30**:1647–1662.
- Seifi A, Kaloshian I, Vossen J, Che D, Bhattarai KK, Fan J, Naher Z, Goverse A, Tjallingii WF, Lindhout P. 2011. Linked, if not the same, Mi-1 homologues confer resistance to tomato powdery mildew and root-knot nematodes. *Molecular Plant-Microbe Interactions* **24**:441–450 DOI [10.1094/MPMI-06-10-0145](https://doi.org/10.1094/MPMI-06-10-0145).
- Shelp BJ, Mullen RT, Waller JC. 2012. Compartmentation of GABA metabolism raises intriguing questions. *Trends in Plant Science* **17**:57–59 DOI [10.1016/j.tplants.2011.12.006](https://doi.org/10.1016/j.tplants.2011.12.006).
- Shen J, Tieman D, Jones JB, Taylor MG, Schmelz E, Huffaker A, Bies D, Chen K, Klee HJ. 2014. A 13-lipoxygenase, TomloxC, is essential for synthesis of C5 flavour volatiles in tomato. *Journal of Experimental Botany* **65**:419–428 DOI [10.1093/jxb/ert382](https://doi.org/10.1093/jxb/ert382).
- Shi H, Pei L, Gu S, Zhu S, Wang Y, Zhang Y, Li B. 2012. Glutathione S-transferase (GST) genes in the red flour beetle, *Tribolium castaneum*, and comparative analysis with five additional insects. *Genomics* **100**:327–335 DOI [10.1016/j.ygeno.2012.07.010](https://doi.org/10.1016/j.ygeno.2012.07.010).
- Smirnoff N. 2000. Ascorbic acid: metabolism and functions of a multi-faceted molecule. *Current Opinion in Plant Biology* **3**:229–235 DOI [10.1016/S1369-5266\(00\)80070-9](https://doi.org/10.1016/S1369-5266(00)80070-9).
- Stout MJ, Workman KV, Bostock RM, Duffey SS. 1998. Stimulation and attenuation of induced resistance by elicitors and inhibitors of chemical induction in tomato (*Lycopersicon esculentum*) foliage. *Entomologia Experimentalis et Applicata* **86**:267–279 DOI [10.1046/j.1570-7458.1998.00289.x](https://doi.org/10.1046/j.1570-7458.1998.00289.x).
- Subramanyam S, Nemacheck JA, Hargarten AM, Sardesai N, Schemerhorn BJ, Williams CE. 2019. Multiple molecular defense strategies in *Brachypodium distachyon* surmount Hessian fly (*Mayetiola destructor*) larvae-induced susceptibility for plant survival. *Scientific Reports* **6**:2596 DOI [10.1038/s41598-019-39615-2](https://doi.org/10.1038/s41598-019-39615-2).
- Takken FLW, Albrecht M, Tameling WIL. 2006. Resistance proteins: molecular switches of plant defense. *Current Opinion in Plant Biology* **9**:383–390 DOI [10.1016/j.pbi.2006.05.009](https://doi.org/10.1016/j.pbi.2006.05.009).
- Telang MA, Giri AP, Pyati PS, Gupta VS, Tegeder M, Franceschi VR. 2009. Winged bean chymotrypsin inhibitors retard growth of *Helicoverpa armigera*. *Gene* **431**:80–85 DOI [10.1016/j.gene.2008.10.026](https://doi.org/10.1016/j.gene.2008.10.026).
- Thipyapong P, Hunt MD, Steffens JC. 2004. Antisense downregulation of polyphenol oxidase results in enhanced disease susceptibility. *Planta* **220**:105–117 DOI [10.1007/s00425-004-1330-6](https://doi.org/10.1007/s00425-004-1330-6).
- Thipyapong P, Joel DM, Steffens JC. 1997. Differential expression and turnover of the tomato polyphenol oxidase gene family during vegetative and reproductive development. *Plant Physiology* **113**:707–718 DOI [10.1104/pp.113.3.707](https://doi.org/10.1104/pp.113.3.707).
- Thipyapong P, Mahanil S, Bhonwong A, Attajarusit J, Stout M, Steffens J. 2004. Increasing resistance of tomato to lepidopteran insects by overexpression of

- polyphenol oxidase. In: *IX international symposium on the processing tomato*. vol. 724. 29–38.
- Tian D, Peiffer M, De Moraes CM, Felton GW. 2014.** Roles of ethylene and jasmonic acid in systemic induced defense in tomato (*Solanum lycopersicum*) against *Helicoverpa zea*. *Planta* **239**:577–589 DOI [10.1007/s00425-013-1997-7](https://doi.org/10.1007/s00425-013-1997-7).
- Tomato-Genome-Consortium xx. 2012.** The tomato genome sequence provides insights into fleshy fruit evolution. *Nature* **485**:635–641 DOI [10.1038/nature11119](https://doi.org/10.1038/nature11119).
- Tran LT, Taylor JS, Constabel CP. 2012.** The polyphenol oxidase gene family in land plants: Lineage-specific duplication and expansion. *BMC Genomics* **13**:1–12 DOI [10.1186/1471-2164-13-395](https://doi.org/10.1186/1471-2164-13-395).
- Tsuda K, Katagiri F. 2010.** Comparing signaling mechanisms engaged in pattern-triggered and effector-triggered immunity. *Current Opinion in Plant Biology* **13**:459–465 DOI [10.1016/j.pbi.2010.04.006](https://doi.org/10.1016/j.pbi.2010.04.006).
- Urzica EI, Adler LN, Page MD, Linster CL, Arbing MA, Casero D, Pellegrini M, Merchant SS, Clarke SG. 2012.** Impact of oxidative stress on ascorbate biosynthesis in *Chlamydomonas* via regulation of the VTC2 gene encoding a GDP-L-galactose phosphorylase. *Journal of Biological Chemistry* **287**:14234–14245 DOI [10.1074/jbc.M112.341982](https://doi.org/10.1074/jbc.M112.341982).
- Vellosillo T, Aguilera V, Marcos R, Bartsch M, Vicente J, Cascón T, Hamberg M, Castresana C. 2013.** Defense activated by 9-lipoxygenase-derived oxylipins requires specific mitochondrial proteins. *Plant Physiology* **161**:617–627 DOI [10.1104/pp.112.207514](https://doi.org/10.1104/pp.112.207514).
- Vilanova L, Wisniewski M, Norelli J, Viñas I, Torres R, Usall J, Phillips J, Droby S, Teixidó N. 2014.** Transcriptomic profiling of apple in response to inoculation with a pathogen (*Penicillium expansum*) and a non-pathogen (*Penicillium digitatum*). *Plant Molecular and Biology Reporter* **32**:566–583 DOI [10.1007/s11105-013-0676-y](https://doi.org/10.1007/s11105-013-0676-y).
- Viswanath KK, Varakumar P, Pamuru RR, Basha SJ, Mehta S, Rao AD. 2020.** Plant lipoxygenases and their role in plant physiology. *Journal of Plant Biology* **1**:1–13 DOI [10.1007/s12374-020-09241-x](https://doi.org/10.1007/s12374-020-09241-x).
- Vogt T. 2010.** Phenylpropanoid biosynthesis. *Molecular Plant* **3**:2–20 DOI [10.1093/mp/ssp106](https://doi.org/10.1093/mp/ssp106).
- Vos P, Simons G, Jesse T, Wijbrandi J, Heinen L, Hogers R, Frijters A, Groenendijk J, Diergaarde P, Reijans M, Fierens-Onstenk J, Md Both, Peleman J, Liharska T, Hontelez J, Zabeau M. 1998.** The tomato *Mi-1* gene confers resistance to both root-knot nematodes and potato aphids. *Nature Biotechnology* **16**:1365–1369 DOI [10.1038/4350](https://doi.org/10.1038/4350).
- Wang L-L. 2016.** Functional characterization of an-esterase gene involving malathion detoxification in *Bactrocera dorsalis* (Hendel). *Pesticide Biochemistry and Physiology* **130**:44–51 DOI [10.1016/j.pestbp.2015.12.001](https://doi.org/10.1016/j.pestbp.2015.12.001).
- Wang R-L, He Y-N, Staehelin C, Liu S-W, Su Y-J, Zhang J-E. 2017b.** Identification of two cytochrome monooxygenase P450 genes, *CYP321A7* and *CYP321A9*, from the tobacco cutworm moth (*Spodoptera litura*) and their expression in response

- to plant allelochemicals. *International Journal of Molecular Sciences* **18**:2278 DOI [10.3390/ijms18112278](https://doi.org/10.3390/ijms18112278).
- Wang L-L, Huang Y, Lu XP, Jiang XZ, Smaghe G, Feng ZJ, Yuan GR, Wei D, Wang JJ. 2015a.** Overexpression of two α -esterase genes mediates metabolic resistance to malathion in the oriental fruit fly, *Bactrocera dorsalis* (Hendel). *Insect Molecular Biology* **24**:467–479 DOI [10.1111/imb.12173](https://doi.org/10.1111/imb.12173).
- Wang G, Kong J, Cui D, Zhao H, Niu Y, Xu M, Jiang G, Zhao Y, Wang W. 2019.** Resistance against *Ralstonia solanacearum* in tomato depends on the methionine cycle and the γ -aminobutyric acid metabolic pathway. *The Plant Journal* **97**:1032–1047 DOI [10.1111/tpj.14175](https://doi.org/10.1111/tpj.14175).
- Wang L-L, Lu X-P, Smaghe G, Meng L-W, Wang J-J. 2017a.** Functional characterization of *BdBI*, a well-conserved carboxylesterase among tephritid fruit flies associated with malathion resistance in *Bactrocera dorsalis* (Hendel). *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* **200**:1–8 DOI [10.1016/j.cbpc.2017.07.001](https://doi.org/10.1016/j.cbpc.2017.07.001).
- Wang R-L, Staehelin C, Xia Q-Q, Su Y-J, Zeng R-S. 2015b.** Identification and characterization of *CYP9A40* from the tobacco cutworm moth (*Spodoptera litura*), a cytochrome P450 gene induced by plant allelochemicals and insecticides. *International Journal of Molecular Sciences* **16**:22606–22620 DOI [10.3390/ijms160922606](https://doi.org/10.3390/ijms160922606).
- Wu J, Baldwin IT. 2010.** New insights into plant responses to the attack from insect herbivores. *Annual Review of Genetics* **44**:1–24 DOI [10.1146/annurev-genet-102209-163500](https://doi.org/10.1146/annurev-genet-102209-163500).
- Wu J, Hettenhausen C, Meldau S, Baldwin IT. 2007.** Herbivory rapidly activates MAPK signaling in attacked and unattacked leaf regions but not between leaves of *Nicotiana attenuata*. *The Plant Cell* **19**:1096–1122 DOI [10.1105/tpc.106.049353](https://doi.org/10.1105/tpc.106.049353).
- Xiao L-F, Zhang W, Jing T-X, Zhang M-Y, Miao Z-Q, Wei D-D, Yuan G-R, Wang J-J. 2018.** Genome-wide identification, phylogenetic analysis, and expression profiles of ATP-binding cassette transporter genes in the oriental fruit fly, *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae). *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics* **25**:1–8 DOI [10.1016/j.cbd.2017.10.001](https://doi.org/10.1016/j.cbd.2017.10.001).
- Xu L, Mei Y, Liu R, Chen X, Li D, Wang C. 2020.** Transcriptome analysis of *Spodoptera litura* reveals the molecular mechanism to pyrethroids resistance. *Pesticide Biochemistry and Physiology* **169**:104649 DOI [10.1016/j.pestbp.2020.104649](https://doi.org/10.1016/j.pestbp.2020.104649).
- Xu S, Liao C-J, Jaiswal N, Lee S, Yun D-J, Lee SY, Garvey M, Kaplan I, Mengiste T. 2018.** Tomato *PEPR1* ORTHOLOG RECEPTOR-LIKE KINASE1 regulates responses to systemin, necrotrophic fungi, and insect herbivory. *The Plant Cell* **30**:2214–2229 DOI [10.1105/tpc.17.00908](https://doi.org/10.1105/tpc.17.00908).
- Xu Z-F, Qi W-Q, Ouyang X-Z, Yeung E, Chye M-L. 2001.** A proteinase inhibitor II of *Solanum americanum* is expressed in phloem. *Plant Molecular Biology* **47**:727–738 DOI [10.1023/A:1013623628857](https://doi.org/10.1023/A:1013623628857).
- Yang D-Y, Li M, Ma N-N, Yang X-H, Meng Q-W. 2017.** Tomato *SlGGP-LIKE* gene participates in plant responses to chilling stress and pathogenic infection. *Plant Physiology and Biochemistry* **112**:218–226 DOI [10.1016/j.plaphy.2017.01.006](https://doi.org/10.1016/j.plaphy.2017.01.006).

- Yaqoob R, Tahir H, Khan S, Naseem S. 2013.** Insecticide resistance in *Bactocera zonata* (Diptera: Tephritidae) in district, Sargodha, Pakistan. *Biochemistry and Pharmacology* 2:114 DOI [10.4172/2167-0501.1000114](https://doi.org/10.4172/2167-0501.1000114).
- Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. 2012.** Primer-BLAST: A tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics* 13:134 DOI [10.1186/1471-2105-13-134](https://doi.org/10.1186/1471-2105-13-134).
- Zhang G, Jia S, Yan Z, Wang Y, Zhao F, Sun Y. 2020a.** A strawberry mitogen-activated protein kinase gene, *FaMAPK19*, is involved in disease resistance against *Botrytis cinerea*. *Scientia Horticulturae* 265:109259 DOI [10.1016/j.scienta.2020.109259](https://doi.org/10.1016/j.scienta.2020.109259).
- Zhang S, Klessig DF. 2001.** MAPK cascades in plant defense signaling. *Trends in Plant Science* 1385:520–527 DOI [10.1016/S1360-1385\(01\)02103-3](https://doi.org/10.1016/S1360-1385(01)02103-3).
- Zhang Y, Ma M, Han Y, Wang L, Liu Z, Guo H, Fang J. 2019.** Transcript-level analysis of detoxification gene mutation-mediated chlorpyrifos resistance in *Laodelphax striatellus* (Hemiptera: Delphacidae). *Journal of Economic Entomology* 112:1285–1291 DOI [10.1093/jee/toy411](https://doi.org/10.1093/jee/toy411).
- Zhang M-Z, Sun C-H, Liu Y, Feng H-Q, Chang H-W, Cao S-N, Li G-H, Yang S, Hou J, Zhu-Salzman K, Zhang H, Qin G-M. 2020b.** Transcriptome analysis and functional validation reveal a novel gene, *BcCGF1*, that enhances fungal virulence by promoting infection-related development and host penetration. *Molecular Plant Pathology* 21:834–853 DOI [10.1111/mpp.12934](https://doi.org/10.1111/mpp.12934).
- Zhao YJ, Wang ZQ, Zhu JY, Liu NY. 2020.** Identification and characterization of detoxification genes in two cerambycid beetles, *Rhaphuma horsfieldi* and *Xylotrechus quadripes* (Coleoptera: Cerambycidae: Clytini). *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* 243-244:110431 DOI [10.1016/j.cbpb.2020.110431](https://doi.org/10.1016/j.cbpb.2020.110431).
- Zheng S-J, Dicke M. 2008.** Ecological genomics of plant-insect interactions: from gene to community. *Plant Physiology* 146:812–817 DOI [10.1104/pp.107.111542](https://doi.org/10.1104/pp.107.111542).
- Zheng Y, Yang Y, Liu C, Chen L, Sheng J, Shen L. 2015.** Inhibition of *SIMP1*, *SIMP2*, and *SIMP3* disrupts defense signaling pathways and enhances tomato fruit susceptibility to *Botrytis cinerea*. *Journal of Agricultural and Food Chemistry* 63:5509–5517 DOI [10.1021/acs.jafc.5b00437](https://doi.org/10.1021/acs.jafc.5b00437).
- Zou X, Xu Z, Zou H, Liu J, Chen S, Feng Q, Zheng S. 2016.** Glutathione S-transferase *SIGSTE1* in *Spodoptera litura* may be involved in feeding adaption of host plants. *Insect Biochemistry and Molecular Biology* 70:32–43 DOI [10.1016/j.ibmb.2015.10.005](https://doi.org/10.1016/j.ibmb.2015.10.005).