

Reeling them in: Taxonomy of marine annelids used as bait by anglers in the Western Cape Province, South Africa

Carol Simon ^{Corresp., 1}, Jyothi Kara ^{1, 2}, Alheit Du Toit ¹, Hendré van Rensburg ¹, Caveshlin Naidoo ¹, Conrad Matthee ¹

¹ Department of Botany and Zoology, University of Stellenbosch, Stellenbosch, Western Cape, South Africa

² Research and Exhibitions, Iziko Museums of South Africa, Cape Town, Western Cape, South Africa

Corresponding Author: Carol Simon

Email address: csimon@sun.ac.za

Background. Common names are frequently used inconsistently for marine annelid species used as bait in the peer-reviewed literature, field guides and legislative material. The taxonomy of many such species based on morphology only also ignores cryptic divergences not yet detected. Such inconsistencies hamper effective management of marine annelids, especially as fishing for recreation and subsistence is increasing. This study investigates the scale of the problem by studying the use and names of bait marine annelids in the Western Cape Province of South Africa. **Methods.** Fifteen recreational and six subsistence fishers at 12 popular fishing sites in the Western Cape Province donated 194 worms which they identified by common name. Worms were assigned scientific names according to a standard identification key for polychaetes from South Africa, and mitochondrial cytochrome oxidase I (COI) amplified and sequenced. **Results.** This study identified 11 nominal species known by 10 common names, in the families Siphonosomatidae, Arenicolidae, Sabellaridae, Lumbrineridae, Eunicidae, Onuphidae and Nereididae. Cryptic diversity was investigated through employing mitochondrial COI sequences and these data will facilitate future identifications among widely distributed species. Several species (*Siphonosoma dayi*, *Abarenicola gilchristi*, *Scoletoma* species, *Marphysa corallina*, *Lysidice natalensis*, *Heptaceras quinque-dens*, *Perinereis latipalpa*) are reported as bait for the first time, and while the names blood- and moonshineworms were consistently applied to members of Arenicolidae and Onuphidae, respectively, coralworm was applied to members of Sabellaridae and Nereididae. Analysis of COI sequences supported morphological investigations that revealed the presence of two taxonomic units each for specimens initially identified as *Gunnarea gaimardi* and *Scoletoma tetraura* according to identification keys. Similarly, sequences for *Scoletoma* species and *Lysidice natalensis* generated in this study do not match those from specimens in China and India, respectively. Further research is required to resolve the species complexes detected and also to refine the use of names by fishermen over a wider geographic range.

Reeling them in: Taxonomy of marine annelids used as bait by anglers in the Western Cape Province, South Africa

Simon, C.A.^{*1}, Kara, J.^{1#}, du Toit, A.¹, van Rensburg, H.¹, Naidoo, C.¹, Matthee, C.A.¹

¹ Department of Botany and Zoology, University of Stellenbosch, Stellenbosch, South Africa

Current Address: Research and Exhibitions Department, Iziko Museums of South Africa, Cape Town, South Africa

*Corresponding Author: csimon@sun.ac.za

CAS: 0000-0001-9613-5222

CAM: 0000-0002-6836-069X

JK: 0000-0003-2961-1713

HVR: 0000-0001-7441-9929

Abstract

Background. Common names are frequently used inconsistently for marine annelid species used as bait in the peer-reviewed literature, field guides and legislative material. The taxonomy of many such species based on morphology only also ignores cryptic divergences not yet detected. Such inconsistencies hamper effective management of marine annelids, especially as fishing for recreation and subsistence is increasing. This study investigates the scale of the problem by studying the use and names of bait marine annelids in the Western Cape Province of South Africa.

Methods. Fifteen recreational and six subsistence fishers at 12 popular fishing sites in the Western Cape Province donated 194 worms which they identified by common name. Worms were assigned scientific names according to a standard identification key for polychaetes from South Africa, and mitochondrial cytochrome oxidase I (COI) amplified and sequenced.

Results. This study identified 11 nominal species known by 10 common names, in the families Siphonosomatidae, Arenicolidae, Sabellaridae, Lumbrineridae, Eunicidae, Onuphidae and Nereididae. Cryptic diversity was investigated through employing mitochondrial COI sequences and these data will facilitate future identifications among widely distributed species. Several species (*Siphonosoma dayi*, *Abarenicola gilchristi*, *Scoletoma* species, *Marphysa corallina*, *Lysidice natalensis*, *Heptaceras quinquedens*, *Perinereis latipalpa*) are reported as bait for the first time, and while the names blood- and moonshineworms were consistently applied to members of Arenicolidae and Onuphidae, respectively, coralworm was applied to members of Sabellaridae and Nereididae. Analysis of COI sequences supported morphological investigations that revealed the presence of two taxonomic units each for specimens initially identified as *Gunnarea gaimardi* and *Scoletoma tetraura* according to identification keys. Similarly, sequences for *Scoletoma* species and *Lysidice natalensis* generated in this study do not match those from specimens in China and India, respectively. Further research is required to resolve the species complexes detected and also to refine the use of names by fishermen over a wider geographic range.

Keywords: COI mtDNA, common name, pseudocryptic species, taxonomy, bloodworm,
coralworm, moonshineworm, musselworm, polychaetes, puddingworm, wonderworm

Introduction

In South Africa, shore-based marine fishing is an important recreational activity and part of the livelihood for many subsistence fishermen and has shown a steady increase over the last decades (McGrath et al., 1997; Sowman et al., 2014; Saayman et al., 2017). There is a close link between shore fishing and bait collecting (MacKenzie, 2005), so an increase in fishing intensity will certainly correlate with an increase in harvesting of natural stocks of bait species (Nel & Branch, 2014 cf. Hodgson, Allanson & Cretchley, 2000; Napier, Turpie & Clark, 2009; Simon et al., 2019a). However, in a recent assessment of the impacts of recreational and subsistence fishing in marine ecosystems in South Africa, impacts of bait collecting received just a passing mention (Majiedt et al., 2019). This supports Watson et al. (2017) who suggested that despite their wide use, marine annelids (i.e., polychaete worms) and probably many other bait species are universally a poorly managed resource.

A wide variety of marine invertebrates are used as bait by South African fishermen (MacKenzie 2005; Branch et al., 2016; DAFF, 2017). However, live marine annelids (indigenous or imported) are not sold in bait shops and are instead collected by subsistence and recreational fishermen who should possess appropriate permits (DAFF, 2017). Collection is controlled by taxon-specific daily limits (DAFF, 2017), but these restrictions have remained almost unchanged for decades (cf. Gaigher, 1979; van Herwerden et al., 1989 and DAFF, 2017). Furthermore, although nearly 2.5 million worms are harvested annually (Turpie, Heydenrych & Lamberth, 2003), biological information to inform management strategies is limited (Gaigher, 1979; van Herwerden, 1989; Lewis, 2005; Simon et al. 2020) while restrictions on collection also do not accommodate the different bait collecting habits by recreational and subsistence fishermen (Simon et al. 2019a). Knowing which species are being utilised is an important step towards improving management of a resource as many bait species, including those that may be morphologically very similar, may have different life history traits and habitat requirements (Hutchings & Lavesque, 2021), which may influence the vulnerability of species to exploitation. The Marine Recreational Activity Information Brochure issued by the Department of Agriculture, Forestry and Fisheries

in South Africa (now the Department of Forestry, Fisheries and Environment; DAFF 2017) identifies bait worms generically as seaworms, polychaetes and flatworms, and by various common names. The only taxa identified by genus are *Arenicola* Lamarck, 1801, *Nereis* Linnaeus, 1758, *Pseudonereis* Kinberg, 1865b and *Gunnarea* Johannson, 1927. As no images are included in the brochure, it is unclear what the worms listed by common name are. However, the popular *Two Oceans: A guide to the Marine Life of southern Africa* (Branch et al., 2016) provides images and common and scientific names for some baitworms: bloodworm (*Arenicola loveni* Kinberg, 1866), musselworm (*Pseudonereis podocirra* (Schmarda, 1861) as *P. variegata* (Grube & Kröyer, 1858)), wonderworm (*Eunice aphroditois* (Pallas, 1788)), Cape reef worm (*Gunnarea gaimardi* (Quatrefages, 1848), as *G. capensis* (Schmarda, 1861) in earlier editions), and the estuarine wonderworm (*Marphysa haemasoma* Quatrefages, 1866 as *M. elityeni* Lewis & Karageorgopolous, 2008, see Kara et al. 2020). The species names for bloodworm, musselworm and Cape reef worm (also known as coralworm in Branch et al., 2016) correspond with those provided in the Government Gazette No. 39790 (Marine Living Resources Act 2014). The latter source, however, uses different names for *E. aphroditois* (Bobbit or errant worm), *Arabella iricolor* (Montagu, 1804) (moonshineworm) and *M. haemasoma* (wonderworms and listed as *M. sanguinea* (Montagu, 1813)). The situation is further complicated by reports of bait worms in other sources; for example, *Diopatra* Audouin & Milne-Edwards, 1833 species have been called case worm (Day, 1974), moonshineworm (Napier, Turpie & Clark, 2009; van Rensburg, Matthee & Simon, 2020), estuarine wonderworm (Smith & Smith, 2012; Allanson et al., 2016) and coralworm (Fielding, 2007, Fielding personal communication), while *E. aphroditois* has also been called coralworm (Wooldridge & Coetzee, 1998). Thus, management of utilised worms may be hampered by confusion around the identities of the species that are harvested, and a lack of consensus in the names used among fishermen, scientists and managers active in South Africa.

The confusion around the use of common names is further complicated by recent taxonomic research which emphasised how poor our understanding of the biodiversity of South African marine annelids, including some used as bait, is. For example, *P. podocirra* and *M. haemasoma*

were removed from synonymy with apparently globally widespread *P. variegata* and *M. sanguinea*, respectively, so both are in fact indigenous to South Africa (Lewis & Karageorgopoulos, 2008; Kara, Macdonald & Simon, 2018; 2020). By contrast, the *Diopatra* species used as bait in two estuaries on the south and southeast coasts of the country (van der Westhuizen & Marais, 1977; Fielding, 2007; Napier, Turpie & Clark, 2009; Simon et al., 2019a), was only recently identified as *D. aciculata* Knox & Cameron, 1971 (van Rensburg, Matthee & Simon, 2020). This species was originally described in Australia (Knox & Cameron, 1971) and is probably alien in South Africa (Elgetany et al., 2020; van Rensburg, Matthee & Simon, 2020). At least two other bait species, *A. iricolor* and *E. aphroditois*, are also apparently globally widespread with type localities geographically distant from South Africa (see Day, 1967), and may therefore either be misidentified indigenous, or unacknowledged alien species. Some species that are harvested (e.g., *P. podocirra*, *E. aphroditois*, *A. iricolor*, *G. gaimardi*) are also widespread within South Africa (Day, 1967; Branch et al., 2016). The ranges of these species, which have planktonic larvae, span multiple known phylogeographic barriers to gene flow in the region. It is thus likely that complexes of genetically distinct but morphologically identical or similar lineages exist (i.e., cryptic or pseudocryptic species, respectively), each with discrete distributions. This was shown for species previously identified as *Pseudopolydora antennata* Claparède, 1869 from temperate and subtropical regions of the country (Simon, Sato-Okoshi & Abe, 2019b), emphasising the need for thorough taxonomic studies of seemingly widespread species.

This study builds on taxonomic information gathered to date, and explores the use of common names and the nomenclature of marine annelid worms used as bait in the Western Cape Province where fishing is particularly popular (Madjiet et al., 2019), and where harvesting of worms is high (Turpie, Heydenrych & Lamberth, 2003). Furthermore, the province spans two vicariant barriers to gene flow at Cape Point and Cape Agulhas (Teske et al., 2011, Fig. 1), and this may also split species into different taxonomic units. The specific aims of the study are to: 1) identify and provide updated descriptions of the annelid species collected as bait by recreational and subsistence fishermen in the Western Cape Province of South Africa; 2) collate

the common names used by the fishermen towards developing consensus for improved management; and 3) generate mtCOI sequences to explore the existence of species complexes locally and globally and facilitate identifications.

Methodology

Sample sites and collection

Sampling was conducted at 12 popular beach and estuarine fishing locations in the Western Cape Province, South Africa (Fig. 1), from June 2016 to May 2017. Collectively, these sites included sandy (Saldanha Bay, Muizenberg, Strand, Betty's Bay, Pearly Beach, Struisbaai, Witsand, Knysna) and or rocky (Velddrif, Melkbosstrand, Kommetjie, Betty's Bay, Hermanus, Witsand) habitats, which would influence the presence and absence of species collected. Most worms were donated by bait collectors who all gave their prior consent to participate in the project. Involvement by most recreational fishermen was confirmed prior to sampling via fishing mailing lists or word of mouth. Some additional recreational and all subsistence fishermen were approached on site. After the aims of the study were described to participants and verbal consent received (ethical clearance number: SU-HSD-001609 from Stellenbosch University), worms were collected according to the permitted methods (DAFF 2017), under permit RES2017-27 issued to CAS by the Department of Forestry, Fisheries and Environment. Additional samples of arenicolids (bloodworm) were collected by the authors using the same techniques (see Simon et al., 2020). The common names used by the bait collectors were noted for all worms. All sampling was conducted during low tide, with specific collection methods for the different taxa included in the systematic accounts below. In some instances, fishermen were only willing to donate a small piece of the worm that was sufficient for genetic analysis.

Specimen identification and processing

Samples were relaxed in an isotonic solution of 7% MgCl₂ in tap water, measured and photographed. A section of each specimen from the mid-body or posterior was placed in 96% ethanol for molecular analysis. The rest of the specimen was fixed in 4% formalin in seawater for at least 2 days, washed in distilled water and stored in 70% ethanol. Samples were

examined on Leica DM1000 light and MZ75 dissecting microscopes, and photographed with a Leica EC3 camera attachment, or on Leica DM750 light and M80 dissecting microscopes and photographed with an Olympus Targus TG5 attached to the microscope eyepieces. Where necessary, images were stacked in Helicon Focus Version 7.6.4 and processed in Photoshop Version C6. Specimens were identified using Day (1967; 1974), and where necessary, more recent literature appropriate to individual taxa. All specimens were deposited at IZIKO South African Museum (Table 1).

DNA extraction, amplification and sequencing

Approximately 25 mg of tissue was used either from mid-section or posterior end to extract DNA using the Zymo Quick DNA™ MiniPrep Plus kit (Zymo-Spin™) according to manufacturer's protocol. The universal primer pair: LCO1490 and HCO2198 (Folmer et al., 1994) was used to amplify a fragment of the cytochrome oxidase subunit 1 gene for all species. The following PCR thermal conditions were used: 94°C for 3 minutes; 34 cycles with 94°C for 45 seconds, 42°C for 1 minute and 72°C for 1 minute and a final extension at 72°C for 7 minutes (Bleidorn et al., 2005). The amplified PCR products were visualised on a 1% agarose gel using 3 µl of PCR product and 5 µl of Quick-Load Purple 100bp DNA ladder (New England BioLabs Inc.), following Simon et al. (2020). All PCR products were sequenced using Sanger sequencing at the Central Analytical Facility at Stellenbosch University. All newly generated sequences were uploaded on GenBank (Table 1).

Molecular analysis

Sequences were edited in BioEdit Version 7.2.6 (Hall, 1999) and aligned using ClustalW with default parameters in MEGA X (Kumar et al., 2018). Neighbour joining trees were constructed in the same program, per family. Nodal support was obtained using 10 000 bootstrap replicates using the maximum composite likelihood method, with uniform rates and pairwise deletion.

Results

Worms were donated by 15 recreational and six subsistence fishers, with two additional fishers who were not categorised (Table 1). In total, these fishers donated 194 specimens belonging to seven families and 11 nominal species: *Siphonosoma dayi* Stephen, 1942, *Abarenicola gilchristi* Wells, 1963, *Arenicola loveni*, *Gunnarea gaimardi*, *Scoletoma* cf. *tetraura*, *Marphysa* cf. *corallina*, *M. haemasoma*, *Lysidice natalensis* Kinberg, 1865, *Heptaceras quinquedens* (Day, 1951), *Perinereis latipalpa* (Schmarda, 1861) and *Pseudonereis podocirra* (Table 1). Together, these species were referred to by 10 common names (Table 1). Sequences could not be generated for *Heptaceras quinquedens* and *Perinereis latipala* even after multiple attempts, with the remaining nine species representing 11 genetically distinct species, including two species each of *Gunnarea* and *Scoletoma* (Table 1).

Taxonomic account

Order: Sipuncula Stephen, 1964

Family: Siphonosomatidae Kawauchi, Sharma & Giribet, 2012

Genus: *Siphonosoma* Spengel, 1912

Species: *Siphonosoma dayi* Stephen, 1942

Figs 2 & 3

Siphonosoma dayi Stephen, 1942: 246 – 247, Pl. XI, Figs 1 & 2; Day 1974: 49

Common name: Sandworm.

Material examined: Knysna: 34°03'56.0"S 23°02'57.4"E, 2 specimens, MB-A090313 and MB-A090318, 27 January 2017, coll. A. N. du Toit, mid-intertidal sandflats in estuary.

Description: Trunk length 198 and 230 mm, introvert of former 17 mm. In life body light to dark pink, colour retained after fixation (Fig. 2A), internally pearlescent pink (Fig. 2E – H). Skin covered with oval shaped papillae in longitudinal rows, following contours of circular muscle, appear white after fixation. Introvert has terminal mouth ringed with short tentacles (Fig. 2B); papillae chitinised, tubular, scale-like and with dark edges arranged in rows on circular muscle

bands (Fig. 2B – D); larger and more numerous in anterior end (Fig. 2C) than posterior (Fig. 2D). Longitudinal muscle-layer divided into 21 or 22 bands (Fig. 2E, G, black arrows), anastomosing anteriorly to form single sheet in region of introvert (Fig. 2E, black arrowhead). Four retractor muscles; dorsal pair attached to body wall anteriorly, ventral pair attached more posteriorly (Fig. 2E, G, H white arrowheads). Two branches of spindle muscle insert close to dorsal retractor muscles (Fig. 2H, black arrowhead). One pair of nephridia (Fig. 2E, F).

Remarks: New specimens match the original description by Stephen (1942). Although only two specimens were collected and sequenced, *S. dayi* (Fig. 3) forms a well-supported clade which is independent from other known species within the genus.

Collection method: Hand digging and pumping.

Type locality: Knysna, Western Cape Province, South Africa.

Known distribution in South Africa: Knysna (Day, 1974).

Ecology: In sand in low to mid intertidal in estuary.

Subclass: Sedentaria Lamarck, 1818

Infraclass: Scolecida Rouse & Fauchald, 1997

Family: Arenicolidae Johnston, 1835

Genus: *Arenicola* Lamarck, 1801

Species: *Arenicola loveni* Kinberg, 1866

Figs 4 – 6A & B

Arenicola loveni Kinberg, 1866: 355; Ashworth 1911: 2 – 17, Figs 1 – 3; Wells 1962: 348, Pl. 2 & 4; Day 1967: 610, Fig. 29.1 f – k; Day 1974: 62, Fig. 54; Branch et al. 2016: 72, Fig. 27.9

Common name: Bloodworm.

Material examined: Betty's Bay: 34°22'39.6"S 18°51'21.6"E, 3 specimens, MB-A090220 – MB-A090222, 10 February 2017, mid-intertidal, sandy beach, coll. A. du Toit. Knysna: 34°03'28.6"S

23°02'30.9"E, 3 specimens, MB-A090231 – MB-A090233, 27 January 2017, 34°03'54.3"S
 23°03'03.7"E, 2 specimens MB-A090234 – MB-A090235, 28 January 2017, 2 specimens,
 34°03'54.3"S 23°03'03.7"E, 29 January 2017, MB-A090236 – MB-A090237, mid-intertidal sandy
 beach, coll. A. du Toit. Muizenberg, 34°06'18.7"S 18°28'47.4"E, 1 specimen, MB-A090230, 13
 March 2017, coll. A. du Toit, 34°06'27.6"S 18°28'22.3"E, 1 specimen, MB-A090227, 2 specimens,
 34°06'18.7"S 18°28'47.4"E, MB-A090228 – MB-A090229, 25 February 2017, 34°06'27.6"S
 18°28'22.3"E, 1 specimen, MB-A090374, 25 February 2017, coll. A. du Toit and C. Naidoo; low
 intertidal in surf zone, sandy beach. Pearly Beach: 34°39'33"S 19°29'27.43.6"E, 3 specimens
 MB-A090246 – MB-A090248, 12 February 2017, coll. A. du Toit and H. van Rensburg, low-
 intertidal, sandy beach. Saldanha Bay: 33°00'26.9"S 17°56'46.3"E, 7 specimens, MB-A090257 –
 MB-A090263, 27 May 2017; 32°59'49.3"S 17°57'58.3"E, 3 specimens, MB-A090264 – MB-
 A090266, 27 May 2017, 33°00'26.9"S 17°56'46.3"E, 1 specimen, MB-A090375, 27 May 2017,
 coll. C. Naidoo, low intertidal, sandy beach. Struisbaai: 34°47'41.1"S 20°02'57.6"E, 1 specimen,
 MB-A090238, 12 February 2017; 3 specimens, MB-A090239 – MB-A090241, 10 April 2017, 4
 specimens, MB-A090242 – MB-A090245, coll. C. Naidoo, A. du Toit and H. van Rensburg, mid to
 low intertidal, sandy beach. Witsand: 34°23'59.9"S 20°49'47.5"E, 7 specimens, MB-A090250 –
 MB-A090256, coll. C. Naidoo, low intertidal, sandy beach.

Description: Live specimens up to 580 mm, including tail. Fixed specimens up to 296 mm long
 (excluding achaetous tail), 19.2 mm wide at chaetiger 1. In life, body colour variable; pink to
 brown, dark brown to black; usually darker in anterior, becoming lighter from branchial region
 posteriorly (Fig. 4A, B, D), colour retained when fixed. Epidermis tessellated to chaetiger 5 or 6,
 papillated from chaetae 6 or 7 onward, including achaetous tail. Chaetigerous annuli
 prominent, number of annuli between first 4 chaetigers 2-3-4, thereafter 4 (Fig. 4D).
 Anterior region consists of trilobed, non-retractable prostomium with nuchal groove on each
 side (Fig. 4E, arrows). One achaetous segment with 2 annuli (Fig. 4C). Proboscis eversible;
 covered with papillae, no pigment (Fig. 4C, E – H). Papillae on proximal section large and
 triangular (Fig. 4C, E, F, H). Papillae in median section more densely packed, small and nipple-
 shaped, becoming larger and more conical distally (Fig. 4C, F, G). One pair of long septal

pouches that reach back to at least third diaphragm (Fig. 5A, B). One pair of conical oesophageal caecae (Fig. 5A). Thorax with 19 chaetigers. Notopodia rounded triangles, retractable lobes in oval torus (Fig. 5E). Notochaetae capillaries in two rows, anterior row shorter than posterior; with lateral toothed-crests and spinulose lamina (Fig. 6A). Neuropodia oval bearing single row of unidentate hooks (Fig. 6B), sometimes with faint denticle. Neuropodia long, approach midline of venter in branchiate region. Branchiae on chaetigers 7 – 19 (13 pairs), highly vascularised, highly branched, arborescent (tree-shaped) (Fig. 5F). On chaetiger 7 branchiae vestigial; 2 – 10 short gill stems, palmar membrane sometimes inconspicuous (Fig. 5E). Up to 22 main gill stems on branchiae on chaetigers 8 to 18, usually fewer on chaetiger 19. Palmar membrane fuse lower third of gill stems (Fig. 5F), sometimes papillated (Fig. 5G). Five pairs of nephridia on chaetigers 5 – 9; nephridiopores hooded, partially hooded (Fig. 5C, D) or unhooded, posterior to dorsal end of neuropodium. Tail achaetous, papillated, anus terminal.

Remarks: Specimens examined here conform to descriptions by Ashworth (1911) and Wells (1962) which included type material, but maximum size is larger. However, oval depressions seen by Ashworth (1911) ventral to some notopodia were not observed. The colour variants of *A. loveni* from all sites form a well-supported clade (Fig. 7) which is exemplified by the fact that those illustrated in Fig. 4A, B and D are represented by an identical sequence (MK 922158). This clade includes two subclades, representing specimens collected on the west and south coasts, respectively. The structure seen here was previously reported in Simon et al. (2020), where nuclear data confirmed that these west and south coast clades represent a single species. The separation between these clades is demarcated by Cape Point, a location known to present a barrier to gene flow (Teske et al. 2011; Simon et al. 2020).

Type locality: Durban, KwaZulu-Natal, South Africa.

Collection method: By pump or digging with hand or trowel and hooking out with a wire. In Muizenburg collected from within surfzone.

Known distribution in South Africa: Saldanha Bay (Western Cape Province) to Durban (Kwa-Zulu Natal) (Day 1967).

Ecology: In sand in low to mid intertidal on sheltered sandy shores and estuaries.

Genus: *Abarenicola* Wells, 1959

Species: *Abarenicola gilchristi* Wells, 1963

Figs 6C & D & 8

Abarenicola gilchristi Wells, 1963: 147 – 149, Fig. 6c, Pl. 2 & 5; Day 1967: 611 – 612, Fig. 29.2

Arenicola assimilis var. *affinis* Ashworth 1911: 18, Figs 4 & 5 (in part); Day 1955: 427

Common name: Bloodworm, bakkiewurm.

Material examined: Betty's Bay: 34°22'S 18°51'E, 4 specimens (incomplete) (MB-A090223 – MB-A090226), 3 June 2016, mid-intertidal, sand, coll. E. Newman. Pearly Beach: 34°39'48.4"S 19°29'17.2"E, 1 specimen (MB-A090249), 10 April 2017, low-intertidal, sand, coll. A. du Toit and C. Naidoo.

Description: Up to 89 mm long (excluding achaetous tail), 11 mm wide at chaetiger 1. In life, body orange-pink (Fig. 8A), light to dark pink when fixed (Fig. 8B). Epidermis tessellated to chaetiger 4 or middle of chaetiger 5, papillated thereafter. Chaetigerous annuli of first three chaetigers prominent, number of annuli between first four chaetigers 2-2 (3 in one specimen)-4, thereafter 4 (Fig. 8B). Anterior region consists of trilobed, non-retractable prostomium and one achaetous segment (Fig. 8B, D). Nuchal groove on each side (Fig. 8D). Proboscis eversible; covered in papillae, no pigment (Fig. 8C). Papillae on proximal section sparsely distributed, prominent, irregular in size, rounded (Fig. 8H). Papillae in median section densely packed, small, rounded, skin folded (Fig. 8G). Papillae of distal section densely packed, conical (Fig. 8F). Oesophageal caecae with one elongate and 11 to 20 smaller caecae on either side of mid-line (Fig. 8E), elongate double to more than triple length of short caecae.

Thorax with 19 chaetigers. Notopodia rounded triangles, retractable lobes in oval torus. Notochaetae spinulose capillaries (Fig. 6C) in single row. Neuropodia oval bearing single row of unidentate, finely serrated, hooks (Fig. 6D). Neuropodia short, do not approach midline of venter. Branchiae on chaetigers 8 – 19 (12 pairs) (Fig. 8A). Branchiae highly vascularised, large, up to 19 main gill stems; highly branched, arborescent (tree-shaped), with lateral branches and gill filaments off each stem (Fig. 8I). Palmar membrane fuse lower third to half of gill stems (Fig. 8I). Five pairs of nephridia on chaetigers 5 – 9; nephridiopores unhooded, hooded, and partially hooded (Fig. 8J – L), posterior to dorsal end of neuropodium. Tail achaetous, papillate, anus terminal.

Remarks: Specimens examined here conform to description by Wells (1963) and Day (1967), but are smaller. *Abarenicola gilchristi* formed part of a distinct lineage in a well-supported clade (Fig. 7) also comprising *Abarenicola brevior* (Wells, 1963) and *A. wellsi* Darbyshire, 2017.

Collection method: By hand or digging with trowel.

Type locality: Buffelsbaai, Cape Peninsula, Eastern Cape province, South Africa.

Known distribution in South Africa: Lambert's Bay to Walker Bay. Presence in Pearly Beach. extends known distribution (Day, 1967) eastwards by only a few kilometres. Namibia: Luderitz. Report in Tamil Nadu, India (Thilagavathi et al., 2013) must be treated with caution.

Ecology: In sand in mid to low intertidal on sheltered shores.

Order: Sabellida Levinsen, 1883

Family: Sabellariidae Johnston, 1865

Genus: *Gunnarea* Johannson, 1927

Species: *Gunnarea gaimardi* (Quatrefages, 1848)

Fig. 9

?*Pallasia gaimardi* Quatrefages, 1848a: 24, 1866: 322, Pl. 13. Figs 17 & 18

?*Hermella capensis* Schmarda, 1861: 23, Pl. 23. Fig. 171. ?*Sabellaria capensis* McIntosh 1885: 418, Pl. 25A Figs 24 & 25, Pl. 26A Figs 11 712

Gunnarea capensis Day 1967: Fig. 33.2.d-i (NOT Schmarda, 1861), *in partum*

Gunnarea gaimardi Kirtley 1994: Fig. 3.1.2.a – e, *in partum*

Gunnarea gaimardi Branch et al. 2016: 73, Fig. 28.3

Common name: Coralworm, Cape reef worm, polworm.

Material examined: Velddrif: 32°46'08.8"S 18°08'44.2"E, 10 specimens (incomplete), MB-A090356 – MB-A090358, MB-A090360, MB-A090364, MB-A090367 – MB-A090371, 26 May 2017, sand reefs in the mid-intertidal rock pools, coll. A. N. du Toit. Bettys Bay: 34°22'39.6"S 18°51'21.6"E, 5 specimens (incomplete), MB-A090336, MB-A090337, MB-A090339 – MB-A090441), 3 June 2016, reefs in the lower intertidal zone, coll. E. Newman. Hermanus: 34°24'41.1"S 19°16'44.8"E, 8 specimens (incomplete), MB-A090341 – MB-A090348), 11 February 2017, low to mid intertidal, coll. A. N. du Toit and H. van Rensburg.

Description: Body a maximum of 110 mm in length; body colour opaque white and cream with irregular dark brown spots when fixed (Fig. 9A – C). Opercular crown and opercular stalk completely fused (Fig. 9B). Two rows of golden outer and inner paleae, arranged in two concentric rows (Fig. 9B). Approximately 44 – 48 outer paleae and 35 – 46 inner paleae. Outer paleae geniculate, obtuse in shape with a single weak tooth on the antero-lateral margin (Fig. 9D1, 3). Inner paleae geniculate with elongate, wedge-shaped peaks with sharp tips (Fig. 9D2), arranged toward the midline of the crown with no overlap in paleae (Fig. 9B – C). Anterior margin of crown with 49 – 73 conical papillae (Fig. 9B). Pair of ciliated palps in front of the mouth (Fig. 9E). Buccal lips present, with upper, lower and lateral lips (Fig. 9E). Tentacular filaments compound and branched (Fig. 9C). U-shaped building organ on the thorax (Fig. 9E); neurochaetae consists of capillaries with bipinnate blade margins (Fig. 9F); Parathorax consist of three chaetigers; notochaetae lanceolate interspersed with capillaries (Fig. 9H); neurochaetae alternating lanceolate chaetae of two lengths (Fig. 9G); neurochaetae thinner than notochaetae. Abdomen with pairs of branchiae on each segment; neuropodial lobes

reduced on abdominal chaetigers , surrounded by tori; uncini with five teeth (Fig. 9J);
neurochaetae verticillate chaetae (Fig. 9I); ventral cirri conical with tapering ends, becoming
digitiform with rounded ends, spanning the neuropodial lobe.

Remarks: Specimens collected from all western sites (Veldrif, Betty's Bay and Hermanus)
conformed to the general descriptions according to Day (1967) and Kirtley (1994). Nonetheless,
differences in the morphology of the outer paleae were observed. Day (1967) described two
incurving teeth present on the outer paleae, but this differs from what was observed in
specimens collected in the present study: one tooth on the antero-lateral margin of the outer
paleae. Additionally, Day (1967), described the inner paleae as completely concealing the
"fleshy disk" or opercular disk, however, this was not observed for our specimens, instead the
opercular disk was visible in the mid-section where paleae did not overlap, which was similar to
Kirtley (1994). Nonetheless, all other characters observed for our specimens were similar to
specimens as described by Kirtley (1994) and Day (1967), suggesting that they most likely
represent *Gunnarea gaimardi* sensu stricto. Molecular analyses (Fig. 10) will be discussed under
Gunnarea sp. 1.

Collection method: Breaking off pieces of reef by hand or narrow blade to remove worms from
tubes.

Type locality: Cape of Good Hope, Western Cape Province, South Africa.

Known distribution in South Africa: the nominal species has been reported from KwaZulu-Natal
on the east coast to the west coast of the Western Cape Province; Namibia: Walvis Bay to
Luderitz (Day, 1967), but this needs to be revised.

Ecology: Species form extensive reefs by building sandy tubes on rocks in the low to mid
intertidal of exposed shores.

Species: *Gunnarea* sp. 1

Fig. 11

Material examined: Witsand: 34°23'31.9"S 20°51'50.1"E, 2 specimens (incomplete), MB-A090293, MB-A090294, 30 April 2017, low to mid intertidal, coll. A. N. du Toit.

Description: Body maximum of 43 mm (MB-A090293) in length (MB-A090294 = 34 mm), when fixed body colour opaque white with black pigmentation throughout (Fig. 11A, B). Opercular crown and opercular stalk completely fused (Fig. 11B). Two rows of golden inner and outer paleae (Fig. 11B). Approximately 34 – 38 inner paleae and 42 – 43 outer paleae. Outer paleae geniculate with a single tooth on the antero-dorsal margin (Fig. 11G2, 3), inner paleae geniculate with elongate, wedge-shaped peaks with a sharp tip (Fig. 11G1), and orientated toward the midline, with both rows overlapping and concealing the opercular disk (Fig. 11B). Anterior margin of opercular crown with 50 conical papillae (Fig. 11B). Pair of ciliated palps in front of mouth (Fig. 11E). Buccal lips present with upper, lower and lateral lips (Fig. 11E). Tentacular filaments compound and branched (Fig. 11E). U-shaped building organ as part of thorax (Fig. 11E); neurochaetae capillaries with bipinnate blade margins (Fig. 11C). Parathorax of three chaetigers; notochaetae alternating lanceolate and capillary chaetae (Fig. 11H), neurochaetae lanceolate chaetae of two lengths (Fig. 11I), neurochaetae thinner than notochaetae. Abdomen with a pair of branchiae on each segment; reduced neuropodial lobes surrounded by tori, uncini with seven teeth (Fig. 11F); neurochaetae verticillate (Fig. 11D). Ventral cirri conical with tapering ends, becoming digitiform with rounded ends spanning the neuropodial lobes.

Remarks: Specimens collected from Witsand (southern site) conformed to the general description by Day (1967), including having paleae that completely conceal the “fleshy disk” or opercular disk. Specimens from the southern site generally resemble *Gunnarea gaimardi* (from western sites) in having a single tooth on the antero-lateral margin of the outer paleae. Nonetheless several differences were observed. Firstly, western site specimens were longer (max. of 110 mm), whereas southern specimens were a maximum of 43 mm. The most distinct feature between these two morpho-groups was the shape, orientation and arrangement of paleae on the opercular crown. The peaks of the outer and inner paleae are longer in

specimens from the southern site compared to that observed in specimens from the western sites; the angle of inclination between the handle and peaks of the inner paleae is larger in western specimens than southern specimens; the outer paleae blades are wider and shorter in specimens from southern sites compared to the longer, thinner blades observed in western specimens. The inner paleae in western specimens do not overlap at the midpoint of the opercular disk, thereby exposing the disk, whereas in southern specimens the paleae overlap, completely concealing the disk. Additionally, the abdominal uncini of western specimens have five teeth, which is two less than that observed for southern specimens. Lastly, western specimens have more opercular papillae than southern specimens when comparing similar sized animals; 73, length 45 mm and 50, length 43 mm, respectively. These differences noted between specimens collected from the southern site and *G. gaimardi* from western sites indicate that they are indeed separate species and that specimens from the southern site (Witsand) most likely represents a new undescribed species of the genus. These morphological differences are supported by the molecular analysis which recovered two well supported clades (Fig. 10) and a genetic distance of 6% (± 0.02), thus confirming their separation as independent species. The first clade, designated *G. gaimardi*, included specimens from Velddrif, Betty's Bay and Hermanus (western group) and the second, designated *Gunnarea* sp. 1, included only the specimens from Witsand (Fig. 10). Morphological differences together with the genetic separation of the clades indicate the presence of two species in what has, till now, been considered a monospecific genus (Capa, Hutchings & Peart 2012). Preliminary observations of *Gunnarea* sp. from Port Shepstone in KwaZulu-Natal suggest that they conform to the description of *Gunnarea* sp. 1 and studies are underway to confirm this.

Collection method: Breaking off pieces of reef by hand or narrow blade to remove worms from tubes.

Known distribution in South Africa: Witsand, Western Cape Province, South Africa.

Ecology: Species form extensive reefs by building sandy tubes on rocks in the low to mid intertidal of exposed shores.

Order: Eunicida Dales, 1962

Family: Lumbrineridae Schmarda, 1861

Genus: *Scoletoma* Blainville, 1828

Species: *Scoletoma* sp. 1

Fig. 12

?*Lumbrinereis tetraurus* Day 1953: 435

?*Lumbrineris tetraura* Day 1967: 437, 439, Fig. 17.16 U – W, Branch et al. 2016: 70, Fig. 26.10

Common name: Puddingworm.

Material examined: Betty's Bay: 34°22'S 18°51'E, 1 specimen (incomplete), MB-A090332, 3 June 2016, sandy sediment, coll. E. Newman.

Description: more than 145 mm; L10 = 8.1 mm, W10 = 3.4 mm (Fig. 12A). Prostomium conical, peristomium with two rings, second slightly shorter than first (Fig. 12A). No eyes. Prechaetal lobes truncate throughout, postchaetal lobe longer and bluntly triangular, becoming longer towards posterior end (Fig. 12C – E). Winged capillary chaetae from chaetiger 1 to approximately chaetiger 57. Long-headed simple multidentate hooded hooks (about 0.2 mm long) from approximately chaetiger 4, shortening posteriorly (Fig. 12F, G), after about chaetiger 35, head becomes even shorter with flared hood (Fig. 12H), appearing white. Aciculae yellow. Dental formula: MI = 1 + 1, MII = 5 + 5, MIII = 2 + 2, MIV = 1 + 1 (Fig. 12B), MV free, lateral to MIV and MIII.

Remarks: The specimen conforms to the general description of *S. tetraura* according to Schmarda (1861) and Day (1967), and no characters could be identified to distinguish the specimen collected here from the description of specimens from Chile. However, the specimen differs morphologically and genetically from others collected in this study that also conform to the description of *S. tetraura* by Day (1967), and genetically from specimens identified as *S.*

tetraura in China (Fig. 13; discussed under *Scoletoma* sp. 2 below). The species is therefore identified here as *Scoletoma* sp. 1.

Collection method: Collected with a small trowel from sediment.

Apparent distribution in South Africa: Known only from a single site. It closely resembles *Scoletoma tetraura* which has been reported from Namibia to KwaZulu-Natal.

Ecology: Burrows into sand in rock pools.

Species: *Scoletoma* sp. 2

Fig. 14

?*Lumbrinereis tetraurus* Day 1953: 435

?*Lumbrineris tetraura* Day 1967: 437, 439, Fig. 17.16 u – w, Branch et al. 2016: 70, Fig. 26.10

Common name: Puddingworm.

Material examined: Hermanus, Kammabaai: 34°24'41.1"S 19°16'44.8"E, 6 specimens (incomplete), MB-A090349 – MB-A090354, 11 February 2017, from rock pools in low to mid intertidal, coll. A. N. du Toit and H van Rensburg.

Description: up to more than 300 mm; L10 = 6.8 to 9.8 mm, W10 = 1.9 to 3.4 mm. Prostomium conical, peristomium with two rings, second slightly shorter than first (Fig. 14A). No eyes. Prechaetal lobes truncate, short and rounded throughout, postchaetal lobe longer and bluntly triangular in anterior chaetigers, becoming digitiform and longer towards posterior end (Fig. 14C – E). Winged capillary chaetae from chaetiger 1 to approximately chaetiger 56 to 70. Long-headed simple multidentate hooded hooks (about 0.15mm long) from approximately chaetiger 4, shortening posteriorly (Fig. 14F, G), after about chaetiger 25, but usually after about chaetiger 30 to 35, head becomes even shorter with flared hood (Fig. 14H), appearing white (Fig. 14E). Aciculae yellow. Dental formula (variation): MI = 1 + 1, MII = 5 (6) + 5, MIII = 2 (1) + 1 (2), MIV = 1 + 1 (Fig. 14B), MV free, lateral to MIV and MIII.

Remarks: All six specimens conform to the general description of *S. tetraura* according to Schmarda (1861) and Day (1967), and no characters could be identified to distinguish the specimens collected here from the description of specimens from Chile. However, this species differs from the specimen from Betty's Bay. In *Scoletoma* sp. 2 from Hermanus, the long-headed simple hooded hooks are about 25% shorter than those of *Scoletoma* sp. 1 from Betty's Bay, and post-chaetal lobes are about 30% longer in the posterior. Furthermore, the segments of *Scoletoma* sp. 2 appear to be longer than those of *Scoletoma* sp. 1; in specimens that are similarly wide, specimens of the former are 1.5 to 1.8 mm longer for the first 10 chaetigers than in the latter. Finally, specimens of the two species were collected from different habitats. Further research is needed to determine which, if any, refers to the species recorded previously by Day (1967) as *S. tetraura*.

The morphological separation is supported by molecular analyses (Fig. 13) that retrieved two well-supported operational taxonomic units, *Scoletoma* sp. 1 (from Betty's Bay) and *Scoletoma* sp. 2 (from Hermanus). The two *Scoletoma* species from South Africa form part of a weakly supported clade together with *Scoletoma fragilis* (O.F. Müller, 1776), *Lumbrineris aberrans* Day, 1963, *Lumbrineris erecta* Moore, 1904, *Lumbrineris japonica* Marenzeller, 1879, and *Lumbrineris perkinsi* Carrera-Parra, 2001 which is separate from *S. tetraura* from China. The separation of *Scoletoma* spp 1 and 2 from South Africa and *S. tetraura* from China in two different clades with high support suggests that they are independent species. However, without sequences from the species' type locality in Chile, it is impossible to determine whether the specimens found in China and South Africa all represent new species or whether one of them is an alien. Specimens from the extended global distribution of *S. tetraura* need to be examined, as there are likely more species within this complex. Additionally, *S. tetraura* and *S. fragilis* were previously considered members of *Lumbrineris*, so the other *Lumbrineris* species in the clade should be revised to determine whether they are also in the genus *Scoletoma*, or whether this genus is paraphyletic.

Collection method: Samples from Hermanus collected among broken pieces of *Gunnarea* tubes.

Apparent distribution in South Africa: Known only from a single site. It closely resembles

Scoletoma tetraura, which has been reported from Namibia to KwaZulu-Natal.

Ecology: Burrows into sand among *Gunnarea* tubes.

Family: Eunicidae Berthold, 1827

Genus: *Marphysa* Quatrefages, 1866

Species: *Marphysa* cf. *corallina*

Fig. 15

Marphysa corallina Day 1967: 400, Fig. 17.7 F – J; Branch et al. 2016: 70, Fig 26.7

Common name: Wonderworm.

Material examined: Witsand: 34°23'31.9"S 20°51'50.1"E, 5 specimens, (incomplete) MB-

A090276 – MB-A090280, 30 April 2017, under rocks in rock pools in mid-intertidal, coll. A. N. du Toit.

Description: Body length more than 120 mm; L10 = 8 – 11 mm, W10 = 0.4 – 0.5 mm. In live specimens, body colour medium to dark brown in anterior becoming light brown in posterior; iridescent throughout (Fig. 15A). Prostomium bilobed, lobes frontally rounded; sulcus deep. Prostomial appendages semi-circular with white tapering tips (Fig. 15A); pair of palps extend to second peristomial ring; pair of lateral antennae reaching second segment and one median antenna extending to third segment (Fig. 15A). Black reniform eye spots below pair of lateral antennae (Fig. 15A, black arrows). Four pairs of maxillary plates and one maxilla; MI = 1 + 1, MII = 3 + 3, MIII = 5 + 0, MIV = 4 + 6, MV = 1 + 1 (Fig. 15F). Branchiae pectinate, from chaetiger 35 – 47 onwards present as a single filament, reaching up to five to seven filaments in middle chaetigers (Fig. 15B). Dorsal cirri digitiform in anterior, middle and posterior chaetigers. Ventral cirri conical in anterior chaetigers and reduces to an oval swelling with a rounded tip in posterior chaetigers. Aciculae blunt with dark brown tips and ends and black shafts (Fig. 15B, E);

3 per fascicle in anterior segments, reducing to 2 and then 1 in middle segments; subacicular hooks, light brown tips with black shafts, present from 40th chaetiger with bidentate tips and guards (Fig. 15B, E). Limbate capillaries present in supracicular fascicle throughout (Fig. 15D, E). Pectinate chaetae present in supracicular fascicle; isodont broad blades and fine teeth (Fig. 15E). Compound falcigers, bidentate tips, short blades with guards, present in subacicular fascicle (Fig. 15C).

Remarks: Specimens collected in this study conform to the general description according to Day (1967). Unfortunately, the original description of *M. corallina* (Kinberg, 1865a) was poor, with no illustrations against which to compare the specimens collected in this study. However, since the type locality of *M. corallina* is in Hawaii and the species has a global disjunct distribution, it is probable that the specimens collected here are really an incorrectly identified indigenous species. We therefore take the more conservative route and refer to the species collected in South Africa as *M. cf. corallina*. All specimens collected during this study were incomplete, missing their posterior ends, so characters such as anodont chaetae, the number of branchial filaments and the number of aciculae in the posterior regions were not documented and thus could not be commented on.

All sequences generated clustered with *M. corallina* from KwaZulu-Natal (KT823410) (Kara, 2015), with high bootstrap support, indicating that it is a single species (Fig. 16). Further investigation is underway to confirm the taxonomic status of *M. corallina* in South Africa.

Collection method: By hand from sediment under rocks.

Known distribution in South Africa: Mabibi in northern KwaZulu-Natal to Mgazana in the Eastern Cape Province, Witsand in Western Cape Province (Day, 1967; current study).

Apparent distribution globally: Mozambique, New Zealand, Red Sea, Australia, Marshall Islands, Lakshadweep Island and Juluit Atoll (Day, 1967; Read & Fauchald, 2021).

Ecology: Occupies burrows in sediment under rocks in the mid-intertidal zone.

Species: *Marphysa haemasoma* Quatrefages, 1866

Fig. 17

Marphysa haemasoma Quatrefages, 1866: 334 – 334, Figs. 4B, 6 & 7; Grube 1870: 299

Marphysa sanguinea Day 1967: 396, fig. 17.5 U – Y (NOT Montagu, 1815)

Marphysa elityeni Lewis & Karageorgopoulos 2008: 280 – 281, Figs. 1 & 2; Branch et al. 2016: 69, Fig. 2.5

Marphysa haemasoma Kara et al. 2020: 16 – 21, Figs 4B, 6 & 7

Common name: Wonderworm, bloukoppies. Listed as estuarine wonderworm in Branch et al. (2016).

Material examined: Knysna: 34°02'17.5"S 23°02'23.4"E, 2 specimens (incomplete), MB-

A090326, MB-A090328), 29 January 2017, coll. A. N. du Toit. Betty's Bay: 34°22'S 18°51'E, 5

specimens (incomplete), MB-A090331, MB-A090333 – MB-A090335, MB-A090338), 3 June

2016, digging with a trowel in mid-intertidal rock pools, coll. E. Newman. Strand: 34°07'03.2"S

18°49'29.4"E, 2 specimens, MB-A090271, MB-A090315), 13 January 2017, digging with trowel

in gravel under rocks in the mid-intertidal, coll. A. N. du Toit. Soetwater: 34°09'33.0"S

18°19'40.7"E, 5 specimens (incomplete specimens), MB-A090272 – MB-A090275, MB-A090317,

10 March 2017, under rocks in mid-intertidal rock pools, coll. A. N. du Toit. Melkbosstrand:

33°43'40.3"S 18°26'17.6"E, 4 specimens (incomplete), MB-A090267 – MB-A090270), 26

February 2017, under rocks in mid-intertidal rocky reef, coll. A. N. du Toit and C. Naidoo.

Description: Body length more than 470mm. In life body colour variable: dark brown/red

anterior with white iridescent spots for about 7 chaetigers (Fig. 17A), becoming medium brown

in middle and darker towards the posterior. Specimens from Knysna and Betty's Bay with blue

colouration in anterior for about 6 chaetigers (Fig. 17B, white arrow), becoming light brown in

middle to posterior. Body iridescent in all specimens. Prostomium bilobed, lobes frontally

rounded, sulcus deep (Fig. 17A, B). Prostomial appendages in semi-circle with a brown band just

before the tapering ends in live specimens; pair of palps reaching first peristomial ring, pair of

lateral antennae extending to second segment and one median antenna reaching first chaetiger (Fig. 17A, B). Pair of eyes under the lateral antennae. Four pairs of maxillary plates and a maxilla (variation); MI = 1 + 1, MII = 3 (4) + 4, MIII = 5 + 0, MIV = 3 + 5, MV = 1 + 1 (Fig. 17C). Branchiae pectinate, present from chaetiger 26 onwards as two filaments, reaching a maximum of 8 filaments in middle, reducing to a single filament in middle to posterior, absent in posterior end near pygidium. Acicula black (Fig. 17E) throughout, five per fascicle in anterior chaetigers, reducing to three in posterior; subacicular hooks not observed. Simple capillaries and pectinate chaetae present in supracicular fascicle. Four types of pectinate chaetae; isodonts with fine teeth and symmetrical blades (Fig. 17F) in anterior segments and anodonts with medium and coarse teeth (Fig. 17D, G) in middle to posterior chaetigers. Compound spinigers with short and long blades present in subacicular fascicle throughout.

Remarks: Specimens collected here conformed to the description by Kara et al. (2020), except for those collected from Knysna and Betty's Bay which have a blue anterior (approximately first six chaetigers), becoming light brown in the middle to posterior end. Phylogenetic analysis recovered a single well-supported clade that comprised all specimens from Knysna, Betty's Bay, Strand, Kommetjie (Soetwater) and Melkbosstrand, indicating that the colour morphs are a single species (Fig. 16). Lewis & Karageorgopolous (2008) observed colour variation in specimens which included iridescent blues and greens for the reproductive segments along the length of the body, from chaetigers 70 – 80. This does not conform to the colour morphs found in the specimens in the present study in which the colour was observed in the anterior regions. Nonetheless, the colour on the remaining parts of the body, "medium brown in the middle and darker toward the posterior", conform to that reported by Lewis & Karageorgopolous (2008).

The use of two species of *Marphysa* in the Western Cape Province supports recent research showing that globally, multiple species of this genus, especially members of the *M. sanguinea* complex, are used as bait, even within regions (see review by Hutchings & Lavesque, 2021). Although the current study showed that different colour morphs represent a single species,

further research is needed to determine whether individuals occupying different habitats, as described by Day (1967) and Lewis & Karageorgopoulos (2008), are also a single species.

Collection method: By hand from sediment under boulders in boulder fields.

Type locality: Cape of Good Hope, Western Cape, South Africa.

Known distribution in South Africa: Langebaan Lagoon on the west coast to Port Elizabeth on the south coast (Day, 1967; Kara et al., 2020).

Ecology: Occupies burrows in sediment typically grey/black medium to coarse grains and rich in sulphur. In Knysna, specimens were found in sandier sediments.

Genus: *Lysidice* Lamarck, 1818

Species: *Lysidice natalensis* Kinberg, 1865

Fig. 18

Lysidice natalensis Kinberg, 1865: 566; Hartman 1948: 84, 85, Pl. XI Figs 1 – 2; Day 1951: 40; Day 1953: 435; Day 1960: p336; Day 1967: 401, Fig. 17.7 K – R; Branch et al. 2016: 70, Fig. 26.9

Lysidice atra Schmarda, 1861

Lysidice capensis Grube, 1868: 12, Fig. 4; Day 1934: 53

Common name: Musselworm. Listed as three-antennaed worm in Branch et al. (2016).

Material examined: Witsand: 34°23'31.9"S 20°51'50.1"E, 11 specimens (2 complete), MB-

A090281 – MB-A090289, MB-A090291, MB-A090292, 30 April 2017, from under rocks, in rock pools in mid-intertidal zone, coll. A. du Toit.

Description: Complete specimens 62 and 63 mm long for 126 and 156 chaetigers. L10 5.28 – 9.8 mm, W10 1.84 – 4.5 mm. Colour reddish-brown with white spots, both extending into middle of prostomium and antennae, margin of prostomium and tips and base of antennae white (Fig. 18A). Prostomium bilobed, antennae tapered, lateral antennae shorter than prostomium, median antenna slightly longer, proximal part brown, tips white (Fig. 18A). Mandibles thick; MI

1 + 1; MII 3 + 3; MIII 2-3 + 0; MIV 2-3 +4-7; MV 1 + 1. Parapodia with slender dorsal cirri (Fig. 18B), becoming shorter and thinner from chaetiger 22 to 38 onwards (Fig. 18E, H). Ventral cirrus bluntly triangular (Fig. 18B), getting shorter posteriorly (Fig. 18E), nipple-shaped in posteriormost chaetigers (Fig. 18H). Post-chaetal lobe truncate (Fig. 18B), getting shorter posteriorly (Fig. 18E), inconspicuous in posteriormost chaetigers (Fig. 18H). Superior chaetae limbate capillaries and comb chaetae of two sizes (Fig. 18F). Inferior compound chaetae with short blades, bidentate, teeth usually of similar sizes (Fig. 18C, D), but proximal tooth may be thicker and or longer. Acicula black with blunt tips, one in anterior chaetigers, two in middle and posterior (Fig. 18B, E, H); bidentate acicula hook with small hood from chaetiger 25 – 28 onwards (Fig. 18E, H), teeth may be worn, giving unidentate appearance (Fig. 18G).

Remarks: Original description by Kinberg (1865a) is poor, but this material is later described by Hartman (1948). Specimens collected here generally match this latter description, and those by Day (1951, 1953, 1967), although the posterior ventral cirrus is more prominent than described by Day (1967). The wide distribution within South Africa is suggestive of multiple species and may be further reflected by the two species that Day (1967) synonymised with *L. natalensis* without explanation. It is therefore possible that *L. capensis* and *L. atra*, both originally described from the temperate Western Cape Province in Kalk Bay and the Cape of Good Hope, respectively, are not *L. natalensis* which was first described from Durban in the subtropical KwaZulu-Natal. Additionally, Day (1967) provides no explanation for why *L. atra*, which was described four years before *L. natalensis* and therefore claims priority, was synonymised with the latter. More specimens from throughout the distribution range and any available type material need to be examined to resolve the taxonomy of this species. The description of *L. natalensis* from Pakistan by Mustaqim (2000) is not very detailed, and the only differences from samples examined here are differently shaped post-chaetal lobes. All specimens from Witsand form a well-supported clade that is not reciprocally monophyletic with *L. natalensis* from India (Fig. 16; Sigamani et al., 2020). Identity of the species in Pakistan is also doubtful.

Collection method: By hand.

Type locality: Durban, KwaZulu-Natal, South Africa.

Known distribution in South Africa: From Namibia to northern KwaZulu-Natal (Day 1967).

Ecology: Habitat variable; in the current study specimens were collected from under rocks in rock pools, Day (1934) reported them from muddy sand.

Family: Onuphidae Kinberg, 1865

Genus: *Heptaceras* Ehlers, 1868

Species: *Heptaceras quinquedens* (Day, 1951)

Fig. 19

Onuphis quinquedens Day, 1951: 40 – 42, Fig. 6A – H; Day 1967: 422, Fig. 17.13A – E; Fauchald 1982: 100, Fig. 28B

Heptaceras quinquedens Paxton 1986: 58 – 60, Fig. 36I, J

Common name: moonshineworm.

Material examined: Pearly Beach: 34°40'00.5"S 19°29'42.7"E, 5 specimens (incomplete), MB-A090432 – MB-A090436, 23 January 2017, coll. H. van Rensburg & A. du Toit. Strand beach: 34°06'37.6"S 18°49'14.6"E, 1 specimen (incomplete), MB-A090442, 13 January 2017, coll. H. van Rensburg and A. du Toit. Struisbaai Main Beach: 34°47'32.3"S 20°02'54.8"E, 15 specimens (incomplete), MB-A090421 – MB-A090431, MB-A090437 – MB-A090440, 27 January 2017, coll. H. van Rensburg, A. du Toit and C. Naidoo.

Description: Large species reaching 350mm in length and 6mm width at 10th chaetiger. Anterior section rounded, becoming dorso-ventrally flattened and ventrally convex from chaetiger 3 – 6 onward (Fig. 19D). In life, prostomium and peristomium white (Fig. 19B), rest of body pale, white-brown ventrally and more reddish-brown dorsally (Fig. 19E), becoming paler towards median and posterior sections, dorsum covered with small red-brown spots, more prominent towards anterior (Fig. 19B). Irregularly spaced red-brown or black dots on ceratophoral rings with a single white patch within final elongated distal ring (Fig. 19B). All colouring disappears

after preservation (Fig. 19A, C, D). Iridescent shine observed over entire body in live and preserved specimens (Fig. 19A – D).

Prostomium with frontal extension forming palpohores for frontal palps (Fig. 19C). Lateral antennae reaching chaetiger 4 – 7 on posterior part of prostomium, shorter median antenna reaching chaetiger 2 – 4 placed anterior to lateral antennae. Proximal ceratophoral rings wide, covering most of prostomium (Fig. 19A). Ceratophores with 15 – 30 rings on median antennae and 20 – 48 rings on lateral antennae, each terminating in an elongated distal ring.

Ceratophores at least as long as styles but up to twice the length of styles which taper distally (Fig. 19B, D). Peristomium as long as, or longer than, prostomium with deep mid-dorsal notch on the dorsal margin, flanking an elevated prostomial ridge (Fig. 19A). Peristomial cirri as long as peristomium, slender and tapering, situated distally on peristomium on either side of the mid-dorsal notch, curving laterally (Fig. 19A, B).

Parapodia mounted marginally, anterior three pairs projecting anteriorly, slightly elongated (Fig. 19B, D) and modified with four or five hooded bi- or tridentate pseudo-compound falcigers (Fig. 19G), remaining parapodia directed dorsally. Dorsal cirri simple tapering filament anteriorly with small basal process towards posterior end (Fig. 19H), shorter than branchiae (Fig. 19D, F). Ventral cirri subulate on anterior five chaetigers changing to pad-like globular form (Fig. 19D). Pectinate chaetae from chaetiger 6 – 8 with 22 – 28 teeth (Fig. 19J). Superior limbate chaetae from chaetiger 1. Branchiae start as simple tapered filaments on chaetiger 1 (Fig. 19D), become pectinate on chaetiger 8 – 10 with maximum of 7 – 12 filaments per branchia (Fig. 19F), continuing throughout rest of body (Fig. 19E). Hooded bidentate acicular chaetae appear from 10th chaetiger to the end of the body (Fig. 19I).

Remarks: The specimens examined here match earlier descriptions (Day, 1951; Day, 1967; Fauchald, 1982), but this is the first observation of tridentate falcigers in the modified parapodia, although tridentate falcigers are known to occur within the genus (Paxton, 1986). The third tooth is small (Fig. 19G) and not always present so can easily be overlooked. According to Fauchald (1982) the median antenna is longer than the posterior lateral ones in the holotype (reaching chaetiger three vs. two) but in all of the material examined here, the

posterior lateral antennae were longer than the median antenna, conforming to the description by Paxton (1986). The iridescent shine seen on the body of *H. quinquedens* is similar to that of *Diopatra aciculata* (van Rensburg, Matthee & Simon, 2020) and may be why fishermen commonly refer to both species as moonshineworms.

Collection method: “prawn pumps” during low tide.

Type locality: Umpangazi, KwaZulu-Natal, South Africa.

Known distribution in South Africa: Western Cape Province to KwaZulu-Natal (Day, 1967).

Apparent distribution globally: report in India (Sigamani et al., 2020) needs to be confirmed.

Ecology: They build temporary tubes in the intertidal of sandy beaches, but do not build conspicuous chimneys.

Order: Phyllodocida Dales, 1962

Family: Nereididae Blainville, 1818

Genus: *Perinereis* Kinberg, 1865b

Species: *Perinereis latipalpa* (Schmarda, 1861)

Fig. 20

Nereis (Nereis) latipalpa Schmarda, 1861: 104 – 105, txt-fig. A, B, Ka & b, Pl. 31,244

Neanthes latipalpa Kinberg, 1865b: 171; von Marenzeller 1888: 6 – 7, Fig. 2

Neanthes latipalpa typica Willey, 1904: 260 – 261, Pl. 13, Fig. 9, Pl. 14, Fig. 1 – 2, 2a & b

Perinereis nuntia vallata Day 1967: 334, Fig. 14.12 P – S; Branch et al. 2016: 67, Fig. 25.4 (NOT Grube & Kroyer in Grube 1858).

Perinereis namibia Wilson & Glasby, 1993: 265 – 266, Fig. 10A – K.

Perinereis latipalpa Villalobos-Guerrero 2019: 474 – 483, Figs 3 – 7.

Common name: Coralworm.

Material examined: Kommetjie: 34°08'34.5"S 18°19'20.4"E, 3 specimens (complete), MB-A090297 – MB-A090299), 10 March 2017, under rocks in the mid-intertidal zone, coll. A. N. du Toit.

Description: Body up to 170 mm. Live specimens, body colour dark green in anterior region, light brown in the middle, to a pale yellow in the posterior. Red blotchy pigment in the middle of each segment, prominent from chaetiger 7 – 10 onwards (Fig. 20A, B, black arrows). Rectangular palpophores with rounded palpostyles. Two antennae, slender with tapering ends (Fig. 20A, D). Two pairs of black eyes in a trapezoidal arrangement (Fig. 20A). Maxillary ring with conical paragnaths (Fig. 20C), Area I = 1 – 2, Area II (variation) = 4(9) + 6(10), Area III = 11 – 17 in an oval patch, Area IV = 8(33) + 16(32), spoon shaped patch. Oral ring with conical paragnaths (Fig. 20D, E), Area V = 1, Area VI = 8(10) + 9(12) in a long arc, Area VII – VIII = 34 – 58 cones in two irregular rows. Dorsal and ventral cirri present throughout. Notochaetae, homogomph spinigers with serrated blades, first 3 teeth at the base of the blade larger, becoming smaller and uniform till the tip (Fig. 20F, hoS). Neurochaetae, homogomph, heterogomph spinigers with serrated blades, uniform teeth (Fig. 20E, heS) and heterogomph falcigers with medium sized blades, finely serrated (Fig. 20G).

Remarks: Specimens collected in this study conformed to the recent redescription in Villalobos-Guerrero (2019). However, variation in body size and paragnath arrangement was noted; total length of paratype is 127 mm and paragnath arrangement, Area III = 9, Area IV = 18-23, Area VII – VIII = 53.

Collection method: From under rocks in the mid-intertidal zone.

Type locality: Table Bay, Cape of Good Hope, South Africa.

Known distribution in South Africa: Hondeklip Bay on the west coast to Port St Johns on the east coast; Namibia: extending north to Luderitz Bay; Mozambique (Day, 1967). However, records in Mozambique have not been confirmed and require further investigation.

Genus: *Pseudonereis* Kinberg, 1865b

Species: *Pseudonereis podocirra* (Schmarda, 1861)

Fig. 21

Mastigonereis podocirra Schmarda, 1861: 108, Fig. 217.

Nereis (Nereilepas) stimpsonis Grube, 1866: 176.

Pseudonereis variegata Day 1967: 331, Fig. 14.12A – F (NOT Grube & Kröyer, 1858); Branch et al., 2016: 66, Fig. 25.1

Pseudonereis podocirra Kara, Macdonald & Simon 2018: 1286 – 1291, Figs 2 – 4

Common name: Musselworm.

Material examined: Velddrif: 34°08'34.5"S 18°19'20.4"E, 9 specimens (incomplete), MB-A090355, MB-A090359, MB-A090361 – MB-A090363, MB-A090365, MB-A090366, MB-A090372, MB-A090373), 26 May 2017, from rock pools in the mid-intertidal, coll. A. N. du Toit. Betty's Bay: 34°22'39.6"S 18°51'21.6"E, 3 specimens (incomplete), MB-A090302, MB-A090304, MB-A090305, 10 February 2017, from under mussel beds in the mid-intertidal mussel belt, coll. A. N. du Toit. Hermanus: 34°24'41.1"S 19°16'44.8"E, 6 specimens (incomplete), MB-A090306 – MB-A090310, MB-A090443, 11 February 2017, from under mussels in the mid-intertidal mussel belt, coll. A.N. du Toit and H. van Rensburg.

Description: Body length up to more than 140 mm. Colour variable: greenish-brown, greyish-brown and medium brown (Fig. 21A, B) with white pigmented spots around 4 eyes on prostomium. Black pigmented spots along midpoint of segment boundaries from chaetiger 13 (Fig. 21B). A mix of different types of paragnaths; conical, shield-shaped and p-bars; arranged in distinct areas on pharynx. Area I = 1 conical, Area II = 15 – 17 conical in a wedge shape, Area III = 22 conical in three-four rows, Area IV = 27 – 32 conical and p-bars in a closely spaced arc shape, Area V = 1 conical, Area VI = large shield-shaped bars and Area VII – VIII = 40 conical and p-bars alternating in 2 – 4 rows (Fig. 21C). Oral ring (Fig. 21C), AVI-V-AVI pattern, v-shaped: ridges of AVI sub-medially separated producing parallel furrows. Notopodial ligule enlarged and

elongated from chaetiger 13 to posterior (Fig. 21F). Dorsal and ventral cirri present (Fig. 21F). Homogomph spinigers with finely serrated blades (Fig. 21E) and heterogomph falcigers (Fig. 21D) with concaved and finely serrated blades.

Remarks: Specimens collected in the study conformed to the redescription in Kara, Macdonald & Simon (2018), except for body length which was larger, measuring up to a maximum of 140 mm. Molecular analyses (Fig. 22) recovered a single monophyletic group with strong maximum likelihood support, indicating a single genetically similar population, further supporting Kara, Macdonald & Simon (2018). Synonymy of *P. podocirra* with *P. variegata* was recently reversed (Kara, Macdonald & Simon 2018), but it is not known whether *P. variegata* in KwaZulu-Natal in South Africa, Namibia and Mozambique, as reported by Day (1967), are a single species.

Collection method: Breaking off mussels by hand from the mussel bed, or by pouring household bleach over the bed (ADT; pers. obs.). Collection of nereidid species is no longer permitted (DAFF 2017).

Type locality: Cape of Good Hope, Western Cape, South Africa.

Known distribution in South Africa: Lamberts Bay to Kidds Beach (Kara, Macdonald & Simon, 2018), possibly extending up the east coast to KwaZulu-Natal and Mozambique and up the west coast to Namibia (Day, 1967).

Ecology: In low intertidal among mussel beds and abandoned *Gunnarea* tubes and barnacle shells.

Discussion

This study found that more marine annelid taxa are utilised in South Africa as bait than what has previously been reported. In addition to the widely reported and investigated bait species (*Arenicola loveni*, *Gunnarea gaimardi*, *Marphysa haemasoma* and *Pseudonereis podocirra*; e.g., van Herwerden, 1989; Lewis 2005; Sowman, 2006; Lewis & Karageorgopoulos, 2008; Branch et al., 2016), several taxa were recorded for the first time (*Abarenicola gilchristi*, *Gunnarea* sp. 1, *Heptaceras quinquedens*, *Lysidice natalensis*, *Marphysa* cf. *corallina*, *Perinereis latipalpa*,

Scoletoma spp 1 and 2). This is also the first published report of *Siphonosoma dayi* being used, even though there have been anecdotal reports of fishermen collecting sandworm in Knysna since at least 2009 (M.K.S. Smith, South African national Parks, Knysna, pers. comm.). By contrast, *Arabella iricolor* and *Eunice aphroditois* (or species matching their general descriptions), which are listed as bait in legislation and field guides (Marine Living Resources Act, 2014; Branch et al., 2016), were not collected in this study. This suggests that more species are used in the province than collected by us, possibly because these species did not occur at the sites sampled, and or that identifications of these species being used were incorrect. For example, one of the authors never found *E. aphroditois* in the Western Cape Province even after extensive sampling in apparently appropriate substrate, although she did find *Eunice* species in the subtidal in KwaZulu-Natal on the east coast (JK, personal observations). Furthermore, *A. iricolor* is superficially similar to lumbrinerid species, and it is possible that both taxa are collected, or that these species were confused in the records for bait collecting. Finally, it is not possible to determine whether species that were collected at single sites (*L. natalensis* and *P. latipalpa*) are targeted more widely, or were misidentified since both were called by names more widely used for other species.

Including *Diopatra aciculata* collected in Knysna in a parallel study (van Rensburg, Matthee & Simon, 2020), 14 species were identified in the Western Cape Province by ten common names, excluding Afrikaans translations. For species collected multiple times and from different locations, individual common names were sometimes applied to more than one species. Species of the same family or genus were often known by a single common name; for example, arenicolids (*Arenicola loveni* and *A. gilchristi*) are bloodworm, onuphids (*D. aciculata* and *H. quinquedens*) are moonshineworms, *Scoletoma* species are puddingworms and *Marphysa* species are wonderworms. For the arenicolids and onuphids this is true even when the species show clear morphological or environmental differences which may have been noted by fishermen, as evidenced by fishermen in Pearly Beach who distinguished between bloodworm (*A. loveni*) and the bakkiewurm (*A. gilchristi*). This was the first time that a second arenicolid is reported as bait, even though DAFF (2017) acknowledges that more than one species may be

used when they specify that bloodworm are “All species of the genus *Arenicola*”, although this is inaccurate as only one species of *Arenicola* has been recorded locally. Individual species were sometimes called by multiple common names that were not translations of the same thing. For example, *M. haemasoma* was identified as wonderworm, bloodworm or bloukoppie (this is Afrikaans for ‘blue head’, referring to the blue anterior of worms from Knysna and Betty’s Bay); *G. gaimardi* was identified as coralworm and polwurm (‘pol’ is Afrikaans for a tuft, tussock or clump of grass, and may here refer to the clumps of tubes formed by the worms); *P. podocirra* was identified as musselworm and coralworm, while *D. aciculata* was also called the pypiewurm (this is Afrikaans for ‘pipe worm’, undoubtedly alluding to the chimneys that extend from the mouths of the tubes) by bait collectors in Port Elizabeth (van Rensburg, Pers. Obs.). It is also apparent that individual common names were sometimes applied to species from different families, such as coralworm (*G. gaimardi*, *P. latiplapa*, *P. podocirra*) and musselworm (*P. podocirra*, *L. natalensis*).

For the most part, subsistence and recreational fishermen used the same names (e.g., for arenicolids, sabellarids, onuphids and *Marphysa* species). Variations in use of names may suggest unfamiliarity with bait worms among some subsistence fishermen, such as bloodworm for *M. haemasoma* in Melkbosstrand, or differences in the use of names depending on geographic region and or type of fishermen, such as coralworm for nereidids at Kommetjie and Velddrif. Interestingly, none of the fishermen used the names from Branch et al. (2016) for *M. haemasoma* (estuarine wonderworm which distinguishes it from *E. aphroditois*, the wonderworm), *G. gaimardi* (Cape reef-worm), *L. natalensis* (three-antennaed worm) or *Scoletoma* species (*S. tetraura* false earthworm). Finally, several common names that appear in DAFF (2017), such as rock, shingle, or pot worms, were not used for any of the species collected in this study. The results of this study confirm that common names are sometimes applied in an inconsistent manner by managers and bait collectors. These differences may be maintained through the transfer of knowledge, across generations of bait collectors, of the identification of worms by morphology and ecological patterns. However, it is possible that the application of

common names has changed (e.g., the name moonshineworm applied to onuphids and not *A. iricola* (Marine Living Resources Act, 2014)).

The genetic data confirmed the presence of complexes of morphologically similar species within South Africa and globally. Day (1967) reported *Gunnarea gaimardi* and *S. tetraura* from Namibia to northern KwaZulu-Natal. Given that this range spans the cold Namaqua, warm Agulhas, and subtropical Natal ecoregions (Sink et al., 2018) and barriers to gene flow at Cape Point, Cape Agulhas, Algoa Bay and Wild Coast (Teske et al., 2011), it is not surprising that these nominal species each included two genetically distinct species with geographic and habitat separation, respectively. This may also apply to *L. natalensis* that has a similar distribution (Day, 1967; Branch et al., 2016). Even though all specimens identified here as *Gunnarea* (including *Gunnarea* sp. and *G. gaimardi*) and the *Scoletoma* species from Hermanus and Betty's Bay matched the descriptions of the nominal species provided in Day (1967; *G. gaimardi* and *S. tetraura*, respectively), the two genetic groups identified in each could be easily distinguished after thorough morphological examination. This supports Hutchings & Kupriyanova (2018) who suggested that many descriptions contained in Day (1967), especially of species described before the 1900s such as the two species under discussion, are too generic to enable accurate identification. Similarly, sequences of *L. natalensis* and *Scoletoma* species 1 and 2 generated in this study do not match those generated for *L. natalensis* and *S. tetraura* collected in India and China, respectively (Zhou et al., 2010; Sigamani et al., 2020; Unpublished data: Chen et al., 2017; Yao et al., 2017; Xing & Zang 2020), indicating the presence of complexes of species that may be morphologically similar but genetically distinct, from different locations around the world. Sigamani et al. (2020) used Day (1967) to identify their samples which also included *H. quinquedens*, originally described from South Africa; unfortunately, we were unable to obtain sequences for the samples that we gathered to test whether the specimens from the two countries are conspecific. However, our results again support Hutchings & Kupriyanova (2018) who warned that using Day (1967) to identify polychaetes outside of southern Africa may erroneously inflate the distribution ranges of polychaete species.

Resolving the identities of marine annelids used as bait has several important management implications. This is exemplified by the recent discovery that moonshineworm collected in Swartkops and Knysna estuaries is *D. aciculata*, a species originally described in Australia and is probably an alien in South Africa (Elgetany et al., 2020; Van Rensburg et al., 2020). Thus, the focus of management of this species must change from conserving populations to preventing further population growth and spread (Van Rensburg, 2019; Van Rensburg et al. 2020). This could be done by permitting increased removal by bait collectors, but preliminary investigations suggest that this is unfeasible (van Rensburg, 2019) and that alternative management strategies need to be explored. Knowing the identity of the worms used may also have important implications for the movement of bait species between sites where worms are collected and where fish are caught, since unused bait that can regenerate is frequently discarded in the latter (M.K.S. Smith, South African National Parks, Knysna). This is especially important if the species is alien (as *D. aciculata*), or if species thought to be locally widespread are multiple species with restricted distributions (as may be the case for *Gunnarea*, *Scoletoma* and *Lysidice* species).

The disjunction between the common names used by collectors and managers is especially problematic when considering the worms that should not be collected. The most recent brochure issued by DAFF (2017) states that Cape reef worm (specified as *Gunnarea*), cannot be collected, but that coralworm can. Since collectors contributing to this study all called *Gunnarea* coralworm, and because it is unlikely that many would know the genus name, bait collectors could collect this species not knowing that they are breaking the law (or use it as a defence if they do). The prohibition on collection of *Gunnarea* and musselworms (identified as *Nereis* and *Pseudonereis* by DAFF (2017)) is related to the structural damage caused to reefs and mussel beds during collection (van Herwerden, 1989), although this is not clearly articulated in the information brochure. It may therefore be more effective to specify the prohibition of taxa based on the habitats that they occupy, and not just name.

This study was limited by several constraints. Firstly, the geographical coverage was restricted relative to the total coastline of the province; the fishing sites were selected according to where participants could be recruited in advance (because bait collecting is time consuming and needs to coincide with low tides which further limited sampling opportunities, we contacted a core of the participants via fishing mailing lists to ensure success in collection) while we also avoided sites that were potentially unsafe, such as Strandfontein and Monwabisi beaches along the northern shores of False Bay. Because of this sampling strategy, there was a bias towards recreational fishermen because subsistence fishermen could not be contacted in advance. Instead, subsistence fishermen were approached on an *ad hoc* basis if they were active at the preselected sampling sites. Additionally, many subsistence fishermen were unwilling to donate bait to the project because bait collecting is so time consuming. We were also reluctant to offer compensation to fishermen because the sale of worms is prohibited by law (Marine Living Resources Act, 2014). Consequently, our understanding of the use of common names is still incomplete because species reportedly used as bait, but not found, could not be addressed in this study. This is further exacerbated when fishermen from different fishing sectors and who speak different languages (e.g., English, Afrikaans, isiXhosa) use different names.

In conclusion, the current study has confirmed that more polychaete species are currently used as bait than previously reported. Furthermore, the inconsistent application of common names across taxa and among users, including for the more popular and widespread species, may hamper effective management. The detection of pseudocryptic species complexes among some bait species may have further implications for the management of these taxa as individual species should form separate management units, especially if they are spatially separated. Finally, diversity of marine annelids in general, and bait species in particular, has been underestimated in South Africa, and the global distribution of some has been overestimated. Research to clarify the taxonomy of the members of the pseudocryptic species complexes identified here, i.e., *Scoletoma* species 1 and 2 and *Gunnarea* sp. 1, and the use of polychaetes and common names across a wider geographic range is ongoing.

Acknowledgements

The authors thank Amy Williams for assisting with processing samples, Teresa Derbyshire for commenting on descriptions of the arenicolids, Breyten van der Merwe, Celeste du Toit, the many fishermen who were so generous with their time and expertise, and two anonymous reviewers whose feedback greatly improved the quality of the manuscript.

Additional information and declarations

Author contributions

CAS: Conceived and funded project, processed samples and wrote the manuscript
 JK: Processed and analysed samples and contributed to writing the manuscript
 HvR: Collected and processed samples and contributed to writing the manuscript
 AdT and CN: Collected and processed samples
 CAM: Co-supervised student authors, and participated in analysis of samples and editing final document.

References

- Ashworth JH. 1911.** The Arenicolidae of South Africa, including an account of *Arenicola loveni* Kinberg. *Annals of the South African Museum* **11(1)**:1–27, plate I
- Allanson BR, Human, LRD, Claassens L. 2016.** Observations on the distribution and abundance of a green tide along an intertidal shore, Knysna Estuary. *South African Journal of Botany* **107**:49–54
- Audouin JV, Milne Edwards H. 1832.** [Part 1] Classification des Annélides et description de celles qui habitent les côtes de la France. *Annales des Sciences Naturelles, Paris*. **1(27)**:337–447.
- Audouin JV, Milne Edwards H. 1833.** [Part 2.] Classification des Annélides et description de celles qui habitent les côtes de la France. *Annales des Sciences Naturelles, Paris*. **1(28)**:187–247.
- Berthold AA. 1827.** Naturliche Familien des Thierreichs aus dem Franzosischen. Mit Anmerkungen und Zusätzen. Landes-Industrie Comptoirs, Weimar, x + 606 pp.
- Blainville H M D de. 1818.** Mémoire sur la classe des Sétipodes, partie des Vers à sang rouge de M. Cuvier, et des Annélides de M. de Lamarck. *Bulletin des Sciences, par la Société Philomatique de Paris*. 1818, **3**:78–85

- 1083 **Blainville H. M. D de. 1828.** Mollusques, Vers et Zoophytes [entries in VEA-VERS, volume
1084 **57**]. In: *Dictionnaire des Sciences naturelles, dans lequel on traite methodiquement des differens*
1085 *etres de la nature, consideres soit en eux-memes. d'apres l'etat actuel de nos connais sciences,*
1086 *soit relativement a l'utilite qu'en peuvent retirer la medicine, l'agriculture, le commerce et les*
1087 *arts. Suive d'une biographie des plus celebres naturalistes.* vol. 57 [Tome LVII. Vea - Vers] F.G.
1088 Levrault, Strasbourg & Paris.
- 1089 **Bleidorn C, Vogt L, Bartolomaeus T. 2005.** Molecular phylogeny of lugworms (Annelida,
1090 Arenicolidae) inferred from three genes. *Molecular Phylogenetics and Evolution* **34**:673–679.
- 1091 **Branch GM, Griffiths CL, Branch ML, Beckley LE. 2016.** *Two Oceans: A Guide to the Marine Life*
1092 *of Southern Africa.* Cape Town: Penguin Random House South Africa. 456 pp.
- 1093 **Capa M, Hutchings P, Peart R. 2012.** Systematic revision of Sabellariidae (Polychaeta) and their
1094 relationships with other polychaetes using morphological and DNA sequence data. *Zoological*
1095 *Journal of the Linnean Society* **164**:245–284.
- 1096 **Carrera-Parra LF. 2001.** Lumbrineridae (Annelida: Polychaeta) from the Grand Caribbean region
1097 with the description of six new species. *Journal of the Marine Biological Association of the*
1098 *United Kingdom* **81**(4):599–621
- 1099 **Dales RP. 1962.** The polychaete stomodeum and the inter-relationships of the families of
1100 Polychaeta. *Proceedings of the Zoological Society* **139**:389–428.
- 1101 **Darbyshire T. 2017.** A re-evaluation of the *Abarenicola assimilis* group with a new species from
1102 the Falkland Islands and key to species. *Journal of the Marine Biological Association of the*
1103 *United Kingdom* **97**(5):897–910 doi:10.1017/S0025315417000741
- 1104 **Day JH. 1934.** On a collection of South African Polychaeta with a catalogue of the species
1105 recorded from South Africa, Angola, Mozambique and Madagascar. *Journal of the Linnean*
1106 *Society, Zoology* **39**:15–82
- 1107 **Day JH. 1951.** The polychaete fauna of South Africa. Part 1. The intertidal and estuarine
1108 Polychaeta of Natal and Mozambique. *Annals of the Natal Museum* **12**(1):1–67
- 1109 **Day JH. 1953.** The polychaete fauna of South Africa. Part 2: Errant species from Cape shores
1110 and estuaries. *Annals of the Natal Museum* **12**(3):397–441
- 1111 **Day JH. 1960.** The polychaete fauna of South Africa. Part 6: Sedentary species dredged off Cape
1112 coasts with a few records from inshore. *Annals of the South African Museum* **45**:261–373.

- 1113 **Day JH. 1963.** The polychaete fauna of South Africa. Part 8: New species and records from grab
1114 samples and dredgings. *Bulletin of the British Museum (Natural History), Series Zoology*. **10(7)**:
1115 381–445
- 1116 **Day JH. 1967.** A monograph on the Polychaeta of Southern Africa. *British Museum of Natural*
1117 *History Publications* **646**:1–878.
- 1118 **Day JH. 1974.** A guide to marine life on South African shores. 2nd Edition. A.A. Balkema, Cape
1119 Town/Rotterdam. 300 pp.
- 1120 **(DAFF) Department of Agriculture, Forestry and Fisheries. 2017.** Marine Recreational Activity
1121 Information Brochure. [http:// www.daff.gov.za/daffweb3/Branches/Fisheries-](http://www.daff.gov.za/daffweb3/Branches/Fisheries-Management/Marine-Resource-Management/Recreational-Fishing)
1122 [Management/Marine-Resource-Management/Recreational-Fishing](http://www.daff.gov.za/daffweb3/Branches/Fisheries-Management/Marine-Resource-Management/Recreational-Fishing)
- 1123 **Ehlers E. 1868.** Die Borstenwürmer (Annelida Chaetopoda) nach systematiechen und
1124 anatomischen Untersuchungen dargestellt. Engelmann, Leipzig, pp 269–748
- 1125 **Elgetany AH, van Rensburg H, Hektoen M, Matthee CA, Budaeva N, Simon CA, Struck TH.**
1126 **2020.** Species delineation in the speciation grey zone – the case of *Diopatra* (Annelida,
1127 Onuphidae). *Zoologica Scripta* **49(4)**: 516–534, DOI: 10.1111/zsc.12421
- 1128 **Fauchald K. 1982.** Revision of *Onuphis*, *Nothria*, and *Paradiopatra* (Polychaeta:Onuphidae)
1129 based upon type material. *Smithsonian Contributions to Zoology*. **356**:1–120.
- 1130 **Fauvel P. 1950.** Contribution a la Faune des Annelides polychetes du Sénégal. *Bulletin de*
1131 *l'Institut Francais d'Afrique Noire* **12(2)**:335–394.
- 1132 **Fielding PJ. 2007.** The subsistence bait fishery in the Swartkops Estuary, Port Elizabeth, Eastern
1133 Cape. Analysis of data collected by monitors December 2005–June 2006. Prepared for
1134 Environmental Evaluation Unit, University of Cape Town.
- 1135 **Fisher WK, MacGinitie GE. 1928.** A new echiuroid worm from California. *Annals and Magazine*
1136 *of Natural History* (Series 10) **1(2)**:199–204.
- 1137 **Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994.** DNA primers for amplification of
1138 mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular*
1139 *Marine Biology and Biotechnology* **3(5)**:294–299.
- 1140 **Gaigher CM. 1979.** Aspects of the population dynamics and ecology of the bloodworm
1141 (*Arenicola loveni* Kingerg). MSc thesis. University of Cape Town, South Africa, 113 pp.

- 1142 **Gallardo VA. 1968.** Polychaeta from the Bay of Nha Trang, *South Viet Nam. Naga*
1143 *Report* **4(3)**:35–279.
- 1144 **Grube AE. 1856 (pub. 1857).** Annulata Örstediana. Enumeratio Annulorum, quae in itinere
1145 per Indiam occidentalem et Americam centralem annis 1845-1848 suscepto legit cl. A.S. Örsted,
1146 adjectis speciebus nonnullis a cl. H. Kröyero in itinere ad Americam meridionalem collectis.
1147 *Videnskabelige Meddelelser fra Dansk naturhistorisk Forening i Köbenhavn* [Part 2]:44–62.
- 1148 **Grube A-E. 1857.** Annulata Örstediana. Enumeratio Annulorum, quae in itinere per Indiam
1149 occidentalem et Americam centralem annis 1845-1848 suscepto legit cl. A. S. Örsted, adjectis
1150 speciebus nonnullis a cl. H. Kröyero in itinere ad Americam meridionalem collectis.
1151 (Fortsættelse). *Vidensk. Meddel. Fra. D. naturhist. Foren, Kobenhavn* [Part 2]:158–166.
- 1152 **Grube A-E. 1866.** Beschreibungen neuer von der Novara-Expedition mitgebrachter Anneliden
1153 und einer neuen Landplanarie. *Verhandlungen der kaiserlich-königlichen zoologisch-*
1154 *botanischen Gesellschaft in Wien* **16**:173–184
- 1155 **Grube A-E. 1867.** Anneliden. *Reise der Österreichischen Fregatte Novara um die Erde in den*
1156 *Jahren 1857, 1858, 1859 unter den befehlen des Commodore B. von Wüllerstorff-Urbair.*
1157 *Zoologischer Theil.* **2(3.2)**:1–46, plates I-IV.
- 1158 **Grube AE. 1878.** Annulata Semperiana. Beiträge zur Kenntniss der Annelidenfauna der
1159 Philippinen nach den von Herrn Prof. Semper mitgebrachten Sammlungen. *Mémoires*
1160 *l'Académie Impériale des Sciences de St.- Pétersbourg.* (série 7) **25(8)**:1–300.
- 1161 **Hall TA. 1999.** BioEdit: a user-friendly biological sequence alignment editor and analysis
1162 program for Windows 95/98/NT. *Nucleic Acids Symposium Series* **41**: 95–98
- 1163 **Hartman O. 1948.** The marine annelids erected by Kinberg with notes on some other types in
1164 the Swedish State Museum. *Arkiv för Zoologi Stockholm* **42(1)**:1–137
- 1165 **Hodgson AN, Allanson BR, Cretchley R. 2000.** The exploitation of *Upogebia africana* (Crustacea:
1166 Thalassinidae) for bait in the Knysna Estuary. *Transactions of the Royal Society of South Africa*
1167 **55(2)**:197–204.
- 1168 **Hutchings P, Kupriyanova E. 2018.** Cosmopolitan polychaetes – fact or fiction? Personal and
1169 historical perspectives. *Invertebrate Systematics* **32**:1–9.
- 1170 **Hutchings P, Lavesque N. 2021.** I know who you are, but do others know? Why correct
1171 scientific names are so important for the biological sciences. *Zoosymposia* **19**:151–163.

- 1172 **Johansson KE. 1927.** Beiträge zur Kenntnis der Polychaeten-Familien Hermellidae Sabellidae
1173 und Serpulidae. [inaugural-dissertation]. *Zoologiska Bidrag från Uppsala* **11**:1–184.
- 1174 **Johnston G. 1835.** Illustrations in British Zoology (45 *Lernaea uncinata*, 46 *Arenicola*
1175 *ecaudata*). *Magazine of Natural History and Journal of Zoology, Botany, Mineralogy, Geology*
1176 *and Meteorology*. **8**:565–569, figs. 53–54. (x-1835)
- 1177 **Johnston G. 1865.** A catalogue of the British non-parasitical worms in the collection of the
1178 British Museum. British Museum. London. 365 pp.
- 1179 **Kara J. 2015.** A phylogeny of South African east coast intertidal rocky-shore Polychaete worms
1180 (Annelida) and the genetic structure and demographic history of an example, *Marphysa*
1181 *corallina*. MSc thesis. University of KwaZulu-Natal, South Africa.
- 1182 **Kara J, Macdonald AHH, Simon CA. 2018.** Integrative taxonomic methods reveal an incorrect
1183 synonymisation of the South African *Pseudonereis podocirra* (Schmarda) as the widespread
1184 *Pseudonereis variegata* (Grube) from Chile. *Invertebrate Systematics* **32**:1282–1297.
- 1185 **Kara J, Molina-Acevedo IC, Zanol J, Simon C, Idris I. 2020.** Morphological and molecular
1186 systematic review of *Marphysa* Quatrefages, 1865 (Annelida: Eunicidae) species from South
1187 Africa. *PeerJ* **8**:e10076 <http://doi.org/10.7717/peerj.10076>
- 1188 **Kawauchi GY, Sharma PP, Giribet G. 2012.** Sipunculan phylogeny based on six genes, with a
1189 new classification and the descriptions of two new families. *Zoologica Scripta* **41**(2):186–210.
- 1190 **Kinberg JGH. 1865a.** Annulata nova. *Öfversigt af Königlich Vetenskapsakademiens*
1191 *förhandlingar, Stockholm* **21**(10):559–574.
- 1192 **Kinberg JGH. 1865b.** Annulata nova. [Continuatio.]. *Öfversigt af Königlich*
1193 *Vetenskapsakademiens förhandlingar, Stockholm* **22**(2):167–179.
- 1194 **Kinberg JGH. 1866.** Annulata nova. [Continuatio.]. *Öfversigt af Königlich Vetenskapsakademiens*
1195 *förhandlingar, Stockholm*. **23**(9):337–357.
- 1196 **Kirtley DW. 1994.** A review and taxonomic revision of the family Sabellariidae Johnston, 1865
1197 (Annelida; Polychaeta). *Sabecon Press Science Series* **1**:1–223.
- 1198 **Knox GA, Cameron DB. 1971.** Port Phillip Bay Survey 1957–1963, Victoria, Australia. Part 2:
1199 Polychaeta. *Memoirs of the National Museum of Victoria* **32**:21–42.
- 1200 **Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018.** MEGA X: Molecular Evolutionary Genetics
1201 Analysis across computing platforms. *Molecular Biology and Evolution* **35**:1547–1549.

- 1202 **Lamarck JB. 1801.** System of animals without vertebrae, or a general table of classes, orders
1203 and genera of these animals; Showing their essential characters and their distribution, after
1204 consideration of their natural relationships and their organization, and following the
1205 arrangement established in the galleries of the Museum of Natural History, among their
1206 preserved remains; Preceded by the opening speech of the Course of Zoology, given in the
1207 National Museum of Natural History year 8 of the Republic. Published by the author and
1208 Deterville, Paris: viii + 432 pp.
- 1209 **Lamarck JB. 1818.** Histoire naturelle des Animaux sans Vertèbres, présentant les caractères
1210 généraux et particuliers de ces animaux, leur distribution, leurs classes, leurs familles, leurs
1211 genres, et la citation des principales espèces qui s'y rapportent; precedes d'une Introduction
1212 offrant la détermination des caractères essentiels de l'Animal, sa distinction du végétal et
1213 des autres corps naturels, enfin, l'Exposition des Principes fondamentaux de la Zoologie. *Paris,*
1214 *Deterville. 5:612 pp.*
- 1215 **Lewis C. 2005.** Aspects of the reproductive biology of the South African polychaete, *Arenicola*
1216 *loveni loveni* (Kinberg 1866). *Invertebrate Reproduction and Development* **48(1–3):147–160.**
- 1217 **Lewis C, Karageorgopoulos P. 2008.** A new species of *Marphysa* (Eunicidae) from the western
1218 Cape of South Africa. *Journal of the Marine Biological Association UK* **88:277–287**
- 1219 **Linnaeus C. 1758.** Systema Naturae per regna tria naturae, secundum classes, ordines, genera,
1220 species, cum characteribus, differentiis, synonymis, locis. *Editio decima, reformata [10th revised*
1221 *edition]. Laurentius Salvius Holmiae* **1:824 pp**
- 1222 **MacKenzie BL. 2005.** An assessment of the shore baitfishery in the Eastern Cape. MSc thesis.
1223 Rhodes University, South Africa, 115 pp
- 1224 **Majiedt PA, Holness S, Sink KJ, Reed J, Franken M, van der Bank MG, Harris LR, Adams L,**
1225 **Perschke M, Miza SA, Currie J, Dunga LV. 2019. Chapter 4: Pressures on Marine Biodiversity.**
1226 **In: Sink KJ, van der Bank MG, Majiedt PA, Harris LR, Atkinson LJ, Kirkman SP, Karenzi N (eds).**
1227 **2019. South African National Biodiversity Assessment 2018 Technical Report Volume 4: Marine**
1228 **Realm.** South African National Biodiversity Institute, Pretoria. South Africa.
1229 <http://hdl.handle.net/20.500.12143/6372>
- 1230 **Marine Living Resources Act of 2014.** Government Gazette, Volume 609, No. 39790. Available
1231 at www.gpwonline.co.za

- 1232 **McGrath MD, Horner CCM, Brouwer SL, Lamberth SJ, Mann BQ, Sauer WHH, Erasmus C. 1997.**
1233 An economic valuation of the South African linefishery. *South African Journal of Marine Science*
1234 **18(1):**203–211.
- 1235 **McIntosh WC. 1885.** Report on the Annelida Polychaeta collected by H.M.S. Challenger during
1236 the years 1873 – 76. *Report on the Scientific Results of the voyage of H.M.S. Challenger*
1237 *(Zoology)* **12:**1-554.
- 1238 **Marenzeller E von. 1879.** Südjapanische Anneliden. I. (Amphinomea, Aphroditea, Lycoridea,
1239 Phyllodocea, Hesionea, Syllidea, Eunicea, Glycera, Sternaspidea, Chaetopterea, Cirratulea,
1240 Amphictenea.). *Denkschriften der Kaiserlichen Akademie der Wissenschaften, Mathematisch-*
1241 *naturwissenschaftliche Classe, Wien.* **41(2):**109–154, plates I–VI.
- 1242 **Montagu G. 1804.** Description of several marine animals found on the south coast of
1243 Devonshire. *Transactions of the Linnean Society, London* **7:**61–85, pl. 6–7.
- 1244 **Montagu G. 1813.** Descriptions of several new or rare animals, principally marine, discovered
1245 on the South coast of Devonshire. *Transactions of the Linnean Society of London* **11(1):**1–26.
- 1246 **Moore JP. 1904.** New Polychaeta from California. *Proceedings of the Academy of Natural*
1247 *Sciences of Philadelphia* **56:**484–503, plates XXXVII–XXXVIII.
- 1248 **Moore JP. 1911.** The polychaetous annelids dredged by the U.S.S. "Albatross" off the coast of
1249 Southern California in 1904. III. Euprosynidae to Goniadidae. *Proceedings of the Academy of*
1250 *Natural Sciences of Philadelphia.* **63:**234–318
- 1251 **Müller OF. 1776.** *Zoologiæ Danicæ Prodrömus, seu Animalium Daniæ et Norvegiæ indigenarum*
1252 *characteres, nomina, et synonyma imprimis popularium.* Havniæ [Copenhagen]:Hallageri. xxxii +
1253 274 pp
- 1254 **Napier V R, Turpie JK, Clark BM. 2009.** Value and management of the subsistence fishery at
1255 Knysna Estuary, South Africa. *African Journal of Marine Science* **31(3):**297–310.
- 1256 **Nel P, Branch GM. 2014.** The effect of bait collectors on stocks of *Callichirus kraussi* and
1257 *Upogebia* species in Langebaan Lagoon. *African Zoology* **49(1):**30–43.
- 1258 **Onwuteaka J. 2016.** Salinity induced longitudinal zonation of polychaete fauna on the Bonny
1259 River Estuary. *Annual Research & Review in Biology* **10(2):**1–14.
- 1260 **Pallas PS. 1788.** Marina varia nova et rariora. *Nova Acta Academiae Scientiarum Imperialis*
1261 *Petropolitanae* **2:**229–249, plates 5–7.

- 1262 **Paxton H. 1986.** Generic revision and relationships of the family Onuphidae (Annelida:
1263 Polychaeta). *Records of the Australian Museum* **38(1)**:1–74.
- 1264 **Pazoki S, Rahimian H, Struck TH, Katouzian AR, Kupriyanova EK. 2020.** A new species of the
1265 *Spirobranchus kraussii*-complex (Annelida, Serpulidae) from the Persian Gulf and Gulf of Oman.
1266 *Zootaxa* **4748(3)**:401–430.
- 1267 **Quatrefages A de. 1848.** Études sur les types inférieurs de l'embranchements des annelés.
1268 Mémoire sur la famille des Hermelliens (*Hermellea* Nob.). *Annales des Sciences Naturelles*,
1269 *Paris. Série 3. Zoologie* **10**:5–58, plate 2.
- 1270 **Quatrefages A de. 1866.** Histoire naturelle des Annelés marins et d'eau douce. Annélides et
1271 Géphyriens. *Librarie Encyclopédique de Roret. Paris.* **1**:1–588
- 1272 **Read G, Fauchald K. (Ed.) 2021.** World Polychaeta database. *Marphysa corallina* (Kinberg,
1273 1865). Accessed at:
1274 <http://www.marinespecies.org/polychaeta/aphia.php?p=taxdetails&id=209831> on 2021-03-30
- 1275 **Saayman M, Saayman A, Zeelie E, Potts W, Mann BQ, Weyl O, Van der Merwe P, Wood A,**
1276 **Raemeakers S, Cowley P, Pledger J, Bova C, Scholtz M. 2017.** Economic significance of
1277 recreational angling in South Africa. Potchefstroom: Tourism Research in Economic Environs &
1278 Society.
- 1279 **Schmarda LK. 1861.** Neue Wirbellose Thiere: Beobachtet und Gesammelt auf einer Reise um
1280 die Erdr 1853 bis 1857. In *Turbellarien, Rotatorien und Anneliden. Leipzig, Verlag von Wilhelm*
1281 *Engelmann.* Erster Band, Zweite Hälfte, 164 pp.
- 1282 **Sowman M. 2006.** Subsistence and small-scale fisheries in South Africa: A ten-year review.
1283 *Marine Policy* **30**:60–73.
- 1284 **Sowman M, Sunde J, Raemaekers S, Schultz O. 2014.** Fishing for equality: Policy for poverty
1285 alleviation for South Africa’s small-scale fisheries. *Marine Policy* **46**:31–42.
- 1286 **Simon CA, du Toit AN, Smith MKS, Claassens L, Smith F & Smith P. 2019a.** Bait collecting by
1287 subsistence and recreational fishers in Knysna Estuary may impact management and
1288 conservation, *African Zoology* **54(2)**:91–103, DOI: 10.1080/15627020.2019.1608862
- 1289 **Simon CA, Sato-Okoshi W, Abe H. 2019b.** Hidden diversity within the cosmopolitan species
1290 *Pseudopolydora antennata* (Claparède, 1869) (Spionidae: Annelida). *Marine Biodiversity* **49**:25–
1291 42.

- 1292 **Simon CA, Kara J, Naidoo C, Matthee CA. 2020.** Genetic structure of bloodworm, *Arenicola*
1293 *loveni* (Annelida; Arenicolidae) suggests risk of local extinction in the face of overexploitation is
1294 lower than expected. *African Zoology* **55(2)**:175–183,
1295 <https://doi.org/10.1080/15627020.2020.1723440>
- 1296 **Smith F, Smith P. 2012.** *A brief field guide to the common shore creatures of the Knysna*
1297 *estuary*. The Knysna Basin Project, Knysna, South Africa. ISBN 978099217590-0
- 1298 **Spengel JW. 1912.** Einige Organisationsverhältnisse von Sipunculusarten und ihre Bedeutung
1299 für die Systematik dieser Tiere. *Deutsch. Zool. Ges.* **22**:261–272.
- 1300 **Stair JB. 1847.** An account of *Palolo*, a sea-worm eaten in the Navigator Islands, with a
1301 description by JE Gray. *Proceedings of the Zoological Society of London*, **15**:17–18.
- 1302 **Stephen AC. 1942.** The South African intertidal zone and its relation to ocean currents: Notes
1303 on the intertidal sipunculids of Cape Province and Natal. *Annals of the Natal Museum*
1304 **10(2)**:245–256
- 1305 **Stephen AC. 1964.** A revision of the classification of the phylum Sipuncula. *Annals and*
1306 *Magazine of Natural History* **7**:457–462.
- 1307 **Teske PR, Von der Heyden S, McQuaid CD, Barker NP. 2011.** A review of marine
1308 phylogeography in southern Africa. *South African Journal of Science* **107(5/6)**:1–11.
- 1309 **Turpie JK, Heydenrych BJ, Lamberth SJ. 2003.** Economic value of terrestrial and marine
1310 biodiversity in the Cape Floristic Region: implications for defining effective and socially optimal
1311 conservation strategies. *Biological Conservation* **112**:233–251.
- 1312 **van Der Westhuizen HC, Marais JFK. 1977.** Stomach content analyses of *Pomadasyys*
1313 *commersonni* from the Swartkops Estuary (Pisces: Pomadasyidae). *Zoologica Africana* **12**:500–
1314 504.
- 1315 **van Herwerden L. 1989.** Collection of musselworms *Pseudonereis variegata* for bait – a
1316 legislative anachronism. *South African Journal of Marine Science* **8**:363–366.
- 1317 **van Herwerden L, Griffiths CL, Bally R, Blaine M, du Plessis C.. 1989.** Patterns of shore
1318 utilization in a metropolitan area: the Cape Peninsula, South Africa. *Ocean & Shoreline*
1319 *Management* **12**: 331–346.
- 1320 **van Rensburg H. 2019.** Taxonomy and Distribution of Moonshine Worms (*Diopatra* sp.) in
1321 Knysna Estuary. Unpublished MSc thesis, Stellenbosch University, South Africa.

- 1322 **van Rensburg H, Matthee CA, Simon CA. 2020.** Moonshineworms (*Diopatra aciculata*:
1323 Onuphidae, Annelida) in the Knysna Estuary, South Africa; taxonomy and distribution. *Journal of*
1324 *the Marine Biological Association of the United Kingdom* **100(6)**:897–907.
1325 <https://doi.org/10.1017/S0025315420000740>
- 1326 **Villalobos-Guerrero TF. 2019.** Redescription of two overlooked species of the *Perinereis nuntia*
1327 complex and morphological delimitation of *P. nuntia* (Savigny in Lamarck, 1818) from the Red
1328 Sea (Annelida, Nereididae). *Zoosystema* **41(24)**:465–496.
- 1329 **Von Marenzeller E. 1888.** Polychäten der Angra Pequena-Bucht. *Zoologische Jahrbücher,*
1330 *Abteilung für Systematik, Geographie und Biologie der Tiere* **3(1)**:1–24.
- 1331 **Watson GJ, Murray JM, Schaefer, Bonner A. 2017.** Bait worms: a valuable and important
1332 fishery with implications for fisheries and conservation management. *Fish and Fisheries*
1333 **18(2)**:374–388
- 1334 **Wells G P. 1959.** The genera of Arenicolidae (Polychaeta). *Proceedings of the Zoological Society*
1335 *of London.* **133(2)**:301–314
- 1336 **Wells G P. 1963.** The lugworms of the southern cold temperate zone (Arenicolidae, Polychaeta).
1337 *Proceedings of the Zoological Society of London* **140(1)**:121–159.
- 1338 **Willey A. 1904.** Littoral Polychaeta from the Cape of Good Hope. *Transactions of the Linnean*
1339 *Society of London, Series 2, Zoology* **9(6)**:255–268.
- 1340 **Wilson RS, Glasby CJ. 1993.** A revision of the *Perinereis nuntia* species group (Polychaeta:
1341 Nereididae). *Records of the Australian Museum* **45(3)**:253–277.
- 1342 **Wooldridge TH, Coetzee PS. 1998.** Marine invertebrates, Chapter 7, pp 117-134. In: Lubke R &
1343 de Moor I, ed. *Field guide to the Eastern and Southern cape coasts.* University of Cape Town
1344 Press, Cape Town.
- 1345 **Zhou H, Zhang Z, Chen H, Sun R, Wang H, Huo L, Pan H. 2010.** Integrating a DNA barcoding
1346 project with an ecological survey: a case study on temperate intertidal polychaetes
1347 communities in Qingdao, China. *Chinese journal of Oceanology and Limnology* **28(4)**:899–910.

Table 1(on next page)

Museum and GenBank accession numbers with location and collector details

Table 1: Museum and GenBank accession numbers with location and collector details.

GenBank accession numbers may be repeated when haplotypes are shared among different individuals. Samples were received from contributing fishermen and processed by Alheit du Toit (AdT), Caveslin Naidoo (CN), Carol Simon (CS), and Hendré van Rensburg (HvR). NS - no sequences. # Sequences were previously published in Simon et al. (2020).

Table 1: Baitworm species from Western Cape, South Africa, found in this study, including common names, locations, collector details. GenBank accession numbers may be repeated when haplotypes are shared among different individuals. Samples were received from contributing fishermen and processed by Alheit du Toit (AdT), Caveshlin Naidoo (CN), Carol Simon (CS), Ethan Newman (EN) and Hendré van Rensburg (HvR). NS - no sequences. # Sequences were previously published in Simon et al. (2020).

Species name	Common name according to fisherman [§]	Location	Fisherman's name	Type of Bait Collector	Collector and sample processor	GenBank Accession Number (COI) (Number of individuals)	Museum Accession Number (number of individuals)
<i>Siphonosoma dayi</i>	Sand worm	Knysna Estuary	Gerrie Barnard	Recreational	AdT	MW598440	MB-A090313
	Sand worm	Knysna Estuary	Gerrie Barnard	Recreational	AdT	MW598441	MB-A090318
<i>Abarenicola gilchristi</i>	Bloodworm	Betty's Bay	Ethan Newman	Recreational	EN & CS	NS	MB-A090223 - MB-A090226 (4)
	Bakkiewurm	Pearly Beach	Frans	Recreational	CN & AdT	MW595992	MB-A090249
	Bakkiewurm	Pearly Beach	Frans	Recreational	CN & AdT	MW595993	DNA only
	Bakkiewurm	Pearly Beach	Frans	Recreational	CN & AdT	MW595994	DNA only
	Bakkiewurm	Pearly Beach	Frans	Recreational	CN & AdT	MW595995	DNA only
<i>Arenicola loveni</i> [#]	Bloodworm	Betty's Bay	Morne & Victor	Recreational	AdT	MK922184	MB-A090220
	Bloodworm	Betty's Bay	Morne & Victor	Recreational	AdT	MK922185	MB-A090221
	Bloodworm	Betty's Bay	Morne & Victor	Recreational	AdT	MK922163	MB-A090222
	Blood worm	Knysna Estuary	Gerrie Barnard	Recreational	AdT	MK922157	MB-A090231
	Blood worm	Knysna Estuary	Gerrie Barnard	Recreational	AdT	MK922158	MB-A090232
	Blood worm	Knysna Estuary	Gerrie Barnard	Recreational	AdT	MK922159	MB-A090233
	Blood worm	Knysna Estuary	Dewald Kamp	Recreational	AdT	MK922160	MB-A090234
	Blood worm	Knysna Estuary	Dewald Kamp	Recreational	AdT	MK922161	MB-A090235
	Blood worm	Knysna Estuary	Albert Kapp	Recreational	AdT	MK922158	MB-A090236, MB-A090237
	Bloodworm	Muizenberg	Anonymous	Recreational	AdT & CN	MK922158	MB-A090227, MB-A090229
	Bloodworm	Muizenberg	Anonymous	Recreational	AdT & CN	MK922164	MB-A090228
	Bloodworm	Muizenberg	Anonymous	Recreational	AdT	NS	MB-A090230
	Bloodworm	Muizenberg	Anonymous	Recreational	AdT & CN	NS	MB-A090374
	Bloodworm	Pearly Beach	Ferdi Joubert	Recreational	AdT & HvR	MK922163	MB-A090246, MB-A090247

	Bloodworm	Pearly Beach	Ferdi Joubert	Recreational	AdT & HvR	MK922183	MB-A090248
	Bloodworm	Saldanha Bay	Anonymous	Unspecified	CN	MK922165	MB-A090257
	Bloodworm	Saldanha Bay	Anonymous	Unspecified	CN	MK922166	MB-A090258, MB-A090264
	Bloodworm	Saldanha Bay	Anonymous	Unspecified	CN	MK922167	MB-A090259
	Bloodworm	Saldanha Bay	Anonymous	Unspecified	CN	MK922168	MB-A090260
	Bloodworm	Saldanha Bay	Anonymous	Unspecified	CN	MK922169	MB-A090261
	Bloodworm	Saldanha Bay	Anonymous	Unspecified	CN	MK922170	MB-A090262
	Bloodworm	Saldanha Bay	Anonymous	Unspecified	CN	MK922171	MB-A090263, MB-A090266
	Bloodworm	Saldanha Bay	Anonymous	Unspecified	CN	MK922172	MB-A090265
	Bloodworm	Saldanha Bay	Anonymous	Unspecified	CN	NS	MB-A090375
	Bloodworm	Struisbaai	Gert Kotze	Recreational	CN, AdT & HvR	MK922163	MB-A090238, MB-A090242
	Bloodworm	Struisbaai	Gert Kotze	Recreational	CN & AdT	MK922173	MB-A090239
	Bloodworm	Struisbaai	Gert Kotze	Recreational	CN & AdT	MK922174	MB-A090240
	Bloodworm	Struisbaai	Gert Kotze	Recreational	CN & AdT	MK922158	MB-A090241
	Bloodworm	Struisbaai	Gert Kotze	Recreational	CN	MK922175	MB-A090243
	Bloodworm	Struisbaai	Gert Kotze	Recreational	CN	MK922176	MB-A090244
							MB-A090245, MB-A090250, MB-A090251, MB-A090254, MB-A090255
	Bloodworm	Struisbaai	Gert Kotze	Recreational	CN	MK922158	
	Bloodworm	Witsand	Paul	Recreational	CN	MK922178	MB-A090252
	Bloodworm	Witsand	Paul	Recreational	CN	MK922179	MB-A090253
	Bloodworm	Witsand	Paul	Recreational	CN	MK922157	MB-A090256
<i>Gunnarea gaimardi</i>	Coral worm	Betty's Bay	Morne & Victor	Recreational	AdT	MN045177	DNA only
	Coral worm	Betty's Bay	Morne & Victor	Recreational	AdT	MN045178	DNA only
	Coral worm	Betty's Bay	Morne & Victor	Recreational	AdT	MN045179	DNA only
							MB-A090336, MB-A090337, MB-A090339
	Coral worm	Betty's Bay	Ethan Newman	Recreational	CS	MN045177	
	Coral worm	Betty's Bay	Ethan Newman	Recreational	CS	MN045181	MB-A090340
	Coral worm	Betty's Bay	Ethan Newman	Recreational	CS	MN045180	MB-A090441

<i>Gunnarea</i> sp.1	Polwurm	Hermanus	Hein Engelbrecht	Recreational	AdT & HvR	MN045177	MB-A090341, MB-A090342, MB-A090344, MB-A090345, MB-A090347, MB-A090348
	Polwurm	Hermanus	Hein Engelbrecht	Recreational	AdT & HvR	NS	MB-A090343
	Polwurm	Hermanus	Hein Engelbrecht	Recreational	AdT & HvR	MN045182	MB-A090346
	Coral worm	Velddrif	Anonymous	Subsistence	AdT	MN045177	MB-A090356 - MB-A090358, MB-A090364, MB-A090367 - MB-A090371 (9)
	Coral worm	Velddrif	Anonymous	Subsistence	AdT	MN045179	MB-A090360
	Coral worm	Witsand	Paul	Recreational	AdT	MN045184	MB-A090293
	Coral worm	Witsand	Paul	Recreational	AdT	MN045183	MB-A090294
<i>Scoletoma</i> cf. <i>tetraura</i> sp. 1 (Betty's Bay)	Pudding worm	Betty's Bay	Ethan Newman	Recreational	CS	MN419154	MB-A090332
<i>Scoletoma</i> cf. <i>tetraura</i> sp. 2 (Hermanus)	Pudding worm	Hermanus	Hein Engelbrecht	Recreational	AdT & HvR	NS	MB-A090349
	Pudding worm	Hermanus	Hein Engelbrecht	Recreational	AdT & HvR	MN419157	MB-A090350
	Pudding worm	Hermanus	Hein Engelbrecht	Recreational	AdT & HvR	NS	MB-A090351
	Pudding worm	Hermanus	Hein Engelbrecht	Recreational	AdT & HvR	NS	MB-A090352
	Pudding worm	Hermanus	Hein Engelbrecht	Recreational	AdT & HvR	MN419156	MB-A090353
	Pudding worm	Hermanus	Hein Engelbrecht	Recreational	AdT & HvR	MN419155	MB-A090354
<i>Marphysa</i> <i>corallina</i>	Wonderworm	Witsand	Paul	Recreational	AdT	MN067881	MB-A090276 - MB-A090278, MB-A090280 (4)
	Wonderworm	Witsand	Paul	Recreational	AdT	MN067882	MB-A090279
<i>Marphysa</i> <i>haemasoma</i>	Wonderworm	Betty's Bay	Ethan Newman	Recreational	CS	NS	MB-A090331
	Wonderworm	Betty's Bay	Ethan Newman	Recreational	CS	MN067877	MB-A090333, MB-A090335, MB-A090338 (3)

	Wonderworm	Betty's Bay	Ethan Newman	Recreational	CS	NS	MB-A090334
	Wonderworm	Knysna Estuary	Anonymous	Recreational	AdT	MN067879 (3)	DNA only
	Wonderworm	Knysna Estuary	Anonymous	Recreational	AdT	MN067878 (2)	DNA only MB-A090326, MB-A090328
	Bloukoppie	Knysna Estuary Melkbos	Anonymous	Subsistence	AdT	MN067878	(2)
	Bloodworm	Strand Melkbos	Lucas	Subsistence	AdT & CN	MN067877 (2)	DNA only MB-A090267 - MB-A090270
	Bloodworm	Strand Soetwater	Lucas	Subsistence	AdT & CN	MN067877	(4)
	Wonderworm	Kommetjie Soetwater	Altus	Subsistence	AdT	MN067877	DNA only
	Wonderworm	Kommetjie Soetwater	Altus	Subsistence	AdT	NS	MB-A090272
	Wonderworm	Kommetjie Soetwater	Altus	Subsistence	AdT	MN067877	MB-A090273 - MB-A090275, MB-A090317 (4)
	Wonderworm	Strand	Marnus	Subsistence	AdT & HvR	MN067880	DNA only MB-A090271, MB-A090315
	Wonderworm	Strand	Marnus	Subsistence	AdT & HvR	MN067880	(2)
<i>Lysidice natalensis</i>	Musselworm	Witsand	Paul	Recreational	AdT	MN419162	MB-A090281
	Musselworm	Witsand	Paul	Recreational	AdT	MN419168	MB-A090282 MB-A090283, MB-A090285
	Musselworm	Witsand	Paul	Recreational	AdT	MN419165	(2)
	Musselworm	Witsand	Paul	Recreational	AdT	MN419164	MB-A090284
	Musselworm	Witsand	Paul	Recreational	AdT	MN419160	MB-A090286
	Musselworm	Witsand	Paul	Recreational	AdT	MN419161	MB-A090287
	Musselworm	Witsand	Paul	Recreational	AdT	MN419158	MB-A090288
	Musselworm	Witsand	Paul	Recreational	AdT	MN419159	MB-A090289
	Musselworm	Witsand	Paul	Recreational	AdT	MN419167	MB-A090291
	Musselworm	Witsand	Paul	Recreational	AdT	MN419163	MB-A090292
<i>Heptaceras quinuedens</i>	Moonshineworm	Pearly Beach	Ferdi Joubert Hermann Schuch & Charlie Friess	Recreational	AdT & HvR	NS	MB-A090432 - MB-A090436 (5)
	Moonshineworm	Strand		Recreational	AdT & HvR	NS	MB-A090442

	Moonshineworm	Struisbaai	Gert Kotze	Recreational	CN, AdT & HvR	NS	MB-A090421 - MB-A090431, MB-A090437 - MB-A090440 (15)
<i>Perinereis latipalpa</i>	Coral worm	Kommetjie	Mario	Subsistence	AdT	NS	MB-A090297 - MB-A090299 (3)
<i>Pseudonereis podocirra</i>	Musselworm	Betty's Bay	Morne & Victor	Recreational	AdT	MN067871	MB-A090302, MB-A090305 (2)
	Musselworm	Betty's Bay	Morne & Victor Hein	Recreational	AdT	MN067870	MB-A090304
	Musselworm	Hermanus	Engelbrecht Hein	Recreational	AdT & HvR	MN067872	MB-A090306
	Musselworm	Hermanus	Engelbrecht Hein	Recreational	AdT & HvR	MN067873	MB-A090307
	Musselworm	Hermanus	Engelbrecht Hein	Recreational	AdT & HvR	MN067871	MB-A090308, MB-A090309, MB-A090443
	Musselworm	Hermanus	Engelbrecht	Recreational	AdT & HvR	MN067872	MB-A090310
	Coral worm	Velddrif	Anonymous	Subsistence	AdT	MN067874	MB-A090355, MB-A090362, MB-A090363, MB-A090365 (4)
	Coral worm	Velddrif	Anonymous	Subsistence	AdT	MN067871	MB-A090359, MB-A090361 (2)
	Coral worm	Velddrif	Anonymous	Subsistence	AdT	MN067872	MB-A090366
	Coral worm	Velddrif	Anonymous	Subsistence	AdT	MN067875	MB-A090372
	Coral worm	Velddrif	Anonymous	Subsistence	AdT	MN067876	MB-A090373

\$ The English names are listed, although fishermen frequently use Afrikaans translations: bloodworm (bloedwurm), Coral worm (koraalwurm), mussel worm (mosselwurm), moonshine worm (maanskynwurm), pudding worm (poedingwurm), wonderworm (wonderwurm). English names were never used for polwurm or bakkiewurm.

Figure 1

Map of South Africa and the Western Cape province with sample sites.

Figure 1. Sample sites in the Western Cape Province, South Africa: Velddrif, Saldanha Bay, Melkbosstrand, Kommetjie (Soetwater), Muizenberg, Strand, Betty's Bay, Hermanus, Pearly Beach, Struisbaai, Witsand, Knysna, with the two main barriers to gene flow in the Western Cape Province, and three main ecoregions along the South African coast. (Map created in <http://www.simplenmappr.net/> and edited in Photoshop Version C6.)




Figure 2

Morphology of *Siphonosoma dayi*

Figure 2. *Siphonosoma dayi* Stephen, 1942. A) Fixed specimen, in two pieces, B) Everted introvert with tentacles, C) Scales on anterior of introvert, D) scales on posterior of introvert, E) Anterior, internal structure showing insertions of introvert muscles (white arrowheads), bands of longitudinal muscles (black arrows), anastomosed sheet of muscle in anterior (black arrowhead) and rectum (*), F) Pair of nephridia (N) and broken rectum (*) with insertion of anus (white arrowhead), G) Magnification of digestive system showing insertions of introvert muscles (arrowheads) and bands of longitudinal muscle (black arrow), H) Close-up of insertion of dorsal introvert muscle (white arrowhead) and spindle muscle (black arrowhead). Scale Bars: A = 10 mm; B, H = 2.5 mm; C = 0.5mm, D = 1 mm; E, F, G = 5 mm; AB - D = MB A090318; E - H = MB A090313





Figure 3

Neighbour Joining tree of *Siphonosoma* species

Figure 3. Neighbour Joining tree using mitochondrial sequences of various *Siphonosoma* Spengel, 1912 species, including *S. dayi* Stephen, 1942 from Knysna. * Indicates bootstrap support greater than or equal to 80%. Sequences obtained in this study are highlighted in grey. *Urechis caupo* Fisher & MacGinite, 1928 was used as the outgroup. Scale bar represents number of substitutions per site.



—

0.050

Figure 4

External morphology of *Arenicola loveni*

Figure 4. *Arenicola loveni* Kinberg, 1866. A) Almost uniformly dark specimen from Muizenberg, dorsal view, B) Dark specimen with distinctly lighter tail from Struisbaai, lateral view, C) Close-up of proboscis of specimen in (B) showing annuli (white arrowheads), D) Light brown specimen with distinctly lighter branchial and tail region from Muizenberg, showing annuli in anterior chaetigers (white arrowheads), dorsal view, E) Prostomium and partially everted proboscis, dorsal view, arrows show nuchal grooves, F) Proboscis showing papillae in different regions, dorsal view, G) Papillae of distal part of proboscis, dorsal view, H) Papillae of proximal part of proboscis, dorsal view. Scale bars: A, B, D = 2 cm, C, E, F = 5 mm, G, H = 2.5 mm. A = MB-A090229, B = MB-A090241, D = MB-A090227, E - H: MB-A090259




Figure 5

Morphology of *Arenicola loveni*

Figure 5. *Arenicola loveni* Kinberg, 1866. A) Dorsal view of digestive system with septal pouches (arrowhead) and single pair of oesophageal caecae (arrow), B) Close up of anterior digestive system (ventral view) and septal pouches (arrowhead) and partially everted proboscis, C) Hooded nephridiopore, D) Partially hooded nephridiopore, E) Chaetiger 7 with vestigial branchia (arrowhead), F) Fully formed branchiae on chaetiger 14, G) Close up of palmar membrane showing papillated surface. Scale bars: A, B, = 10 mm, C, G = 1.5mm, D - F = 2.5mm, A: MB-A090252, B, D: MB-A090250, C, E - G: MB-A090259.




Figure 6

Arenicolid chaetae

Figure 6. Arenicolid chaetae. A) Notochaetae and B) Neuropodial hooks of *Arenicola loveni* Kinberg, 1866, C) Notochaetae and D) Neuropodial hooks of *Abarenicola gilchristi* Wells, 1963. Scale Bars: A - D = 0.1mm. A, B = MB-A090261, C, D = MB-A090225




Figure 7

Neighbour joining tree of Arenicolidae

Figure 7. Neighbour Joining tree using mitochondrial sequences belonging to various *Arenicola Lamarck, 1801* and *Abarenicola Wells, 1959* species, including *A. loveni* Kinberg, 1866 and *A. gilchristi* Wells, 1963 from South Africa. * Indicates bootstrap support greater than or equal to 80%. Areas highlighted in grey represent sequences generated in this study. *Maldanidae* sp. was used as outgroup. Scale bar represents substitutions per site.




Figure 8

Morphology of *Abarenicola gilchristi*

Figure 8. *Abarenicola gilchristi* Wells, 1963 A) Live specimen, B) Dorso-lateral view of head, showing annulations on chaetigers 1 to 2 (arrowheads), C) Proboscis showing papillations in different regions, D) Dorsal view of head showing prostomium and partially everted proboscis, arrows show nuchal grooves, E) Digestive caecae; one large pair and multiple smaller pairs, F) Papillae of distal part of proboscis, G) Papillae of median part of proboscis, H) Papillae of proximal part of proboscis, I) Branchia on chaetiger 9, J) Unhooded nephridiopore, K) Hooded nephridiopore, L) Partially hooded nephridiopore. Scale bars: A = 2 mm, B, C, D, E = 5mm; I = 2mm, F - H = 2mm, J - L = 0.5mm; A, J: MB-A090223, B, I, K, L = MB-A090224; C - H = MB-A090226.




Figure 9

Morphology of *Gunnarea gaimardi*

Figure 9. *Gunnarea gaimardi* (Quatrefages, 1848). A) Live specimen from Betty's Bay, B) Dorsal view of crown showing the inner paleae (ip), outer paleae (op), opercular disk (od), papillae (cp), C) Right side view of crown showing the tentacular filaments (tf), D) Paleae, 1 & 3: outer geniculate paleae with tooth, 2: inner geniculate paleae, E) Ventral view of anterior region showing palps (p), median ridge (mr), and U-shaped building organ (bo), F) Bipinnate capillaries, neurochaetae, G) Lanceolate chaetae of two lengths, neurochaetae, H) Lanceolate and capillaries, notochaetae, I) Verticillate chaetae, neurochaetae, J) Uncini. Scale bars: A & C = 5 mm, B & E = 2mm, D = 0.5mm, F, I, J = 50µm, G & H = 0.2 mm. B, F - J = MB-A090337, c = MB-A090343, d = MB-A090371.




Figure 10

Neighbour joining tree of *Gunnarea*

Figure 11. Neighbour Joining tree of mitochondrial sequences of various species from family Sabellariidae Johnston, 1865 including *Gunnarea capensis* (Schmarda, 1861). * Indicates bootstrap support greater than 80%. Areas in grey represent sequences generated in this study. *Spirobranchus sinuspersicus* Pazoki, Rahimian, Struck, Katouzian & Kupriyanova, 2020 was used to root the tree. Scale bar represents substitutions per site.




Figure 11

Morphology of *Gunnarea* species

Figure 10. *Gunnarea* Johansson, 1927 species collected from Witsand. A) Live specimen, B) Crown showing the inner paleae (ip) and outer paleae (op), C) Neurochaetae of first thoracic chaetiger, D) Abdominal neurochaetae, E) Anterior region showing palps (p), Buccal lip (upper and lateral sides) and tentacular filaments (tf), F) Posterior uncinus, G) Palaea 1- inner geniculate paleae, 2 & 3 - outer geniculate paleae with tooth, H) Lanceolate notochaetae, I) Neurochaetae. Scale bars: A= 5mm, B & E = 1mm, C, D, F, G = 0.5mm, H-I = 0.2mm. A, C, F - I = MB-A090293, B, E = MB-A090294




Figure 12

Morphology of *Scoletoma* species from Betty's Bay

Figure 12. *Scoletoma* species 1 from Betty's Bay. A) Dorsal anterior, B) MIII and MIV of jaws, ventral view, C) Chaetiger 5 showing pre-chaetal (Prcl) and post chaetal (Pcl) lobes, anterior view, D) Chaetiger 31 showing prechaetal and post chaetal lobes, anterior view, E) Posterior chaetiger showing prechaetal and post chaetal lobes, anterior view, F) Long-headed multidentate hooded hooks on chaetiger 5, G) Long-headed multidentate hooded hook on chaetiger 31, H) Short-headed multidentate hook with flared hood from posterior chaetiger. Scale bars: A = 5mm, B - E = 0.5mm, F - H = 0.2mm. A - H = MB-A090332




Figure 13

Neighbour joining tree of Lumbrineridae

Figure 14. Neighbour joining tree of mitochondrial sequences of various species in Lumbrineridae, including *Scoletoma tetraura* (Schmarda, 1861) from China. * Indicates bootstrap support greater than 80%. Areas highlighted in grey represent sequences generated in this study; *Scoletoma* species sp. 1 and sp. 2. Area outlined with grey dashed line represents *S. tetraura* from China. *Arabella iricolor* (Montagu, 1804) and *A. semimaculata* (Moore, 1911) were used as outgroups. Scale bar indicates substitutions per site.




Figure 14

Morphology of *Scoletoma* species from Hermanus

Figure 13. *Scoletoma* species 2 from Hermanus. A) Dorsal anterior, B) MIII and MIV of jaws, ventral view, C) Chaetiger 3 showing pre-chaetal (Prcl) and post chaetal (Pcl) lobes anterior view, D) Chaetiger 30 showing pre- and post chaetal lobes, anterior view, E) Posterior chaetiger showing pre- and post chaetal lobes, anterior view, F) Long-headed multidentate hooded hooks on chaetiger 5, G) Long-headed multidentate hooded hook on chaetiger 31, H) Short-headed multidentate hook with flared hood from posterior chaetiger. Scale bars: A = 5mm, B - E = 0.5mm, F - H = 0.2mm. A - H = MB-A090353




Figure 15

Morphology of *Marphysa corallina*

Figure 15. *Marphysa cf corallina* A) Anterior region (dorsal), live specimen, B) Posterior chaetiger with branchiae, C) Compound bidentate falcigers with guards , D) Limbate capillaries, E) Chaetae, acicula and subacicular hooks, F) Dorsal view of Maxillary apparatus Scale bars: A = 5mm, B = 0.5mm, C & D = 50µm, E = 0.2mm, F = 2mm. A, C & D = MB-A090276, E & F = MB-A090280. Figure 16. Neighbour Joining tree of various species belonging to family Eunicidae Berthold, 1827, including *Marphysa* Quatrefages, 1866 and *Lysidice* Lamarck, 1818 from South Africa. * Indicates bootstrap support greater than or equal to 80%. Grey highlighted areas indicate sequences generated in this study. Red branch represents a questionable sequence labelled as *Lysidice natalensis* Kinberg, 1865 from India. *Palola viridis* Gray in Stail, 1847, *Leodice rubra* Grube, 1856 and *Hyalinocea* sp. were used as outgroups. Scale bar represented substitutions per site.




Figure 16

Neighbour Joining tree of Eunicidae

Figure 16. Neighbour Joining tree of various species belonging to family Eunicidae, including *Marphya* and *Lysidice* from South Africa. * Indicates bootstrap support greater than 80%. Grey highlighted areas indicate sequences generated in this study. Red branch represents a questionable sequence labelled as *Lysidice natalensis* from India. *Palola viridis*, *Eunice rubra* and *Hyalinocea* sp. were used as outgroups. Scale bar represented substitutions per site.




Figure 17

Morphology of *Marphysa haemasoma*

Figure 17. *Marphysa haemasoma* Quatrefages, 1866. A) Dorsal anterior, live specimen B) Dorsal anterior of live specimen, arrow showing blue colouration C) Dorsal view of maxillary apparatus, D) Pectinate anodont chaetae E) Middle chaetiger with branchia, F) Pectinate isodont chaetae, G) Pectinate anodont chaetae. Scale bars: A & B= 5mm, C = 2mm, D, F & G = 50µm, E = 0.25mm. A = MB-A090326, B = MB-A090328, C = MB-A090274, D, F - G = MB-A090273.




Figure 18

Morphology of *Lysidice natalensis*.

Figure 18. *Lysidice natalensis* Kinberg, 1865. A) Dorsal of head, live specimen, B) Chaetiger 5, anterior view, C) Compound falciger of chaetiger 5, D) Compound falciger of chaetiger 28, E) Chaetiger 28, anterior view F) Limbate and comb (white arrowheads) chaetae of chaetiger 28, G) Acicula hook of chaetiger 28, H) Posterior chaetiger. Scale bars: A = 1mm, B, E, H = 0.5mm, C, D, F, G = 0.05mm. A - H = MB-A090291.




Figure 19

Morphology of *Heptaceras quinquedens*

Figure 19. *Heptaceras quinquedens* (Day, 1951). A) Dorsal anterior of preserved specimen showing peristomial notch flanking prostomial ridge, laterally curving peristomial cirri and iridescent shine that remains after preservation, B) Dorsal anterior of live specimen, insert shows freckled spots on anterior dorsum, C) Ventral anterior of preserved specimen, D) Lateral anterior view of preserved specimen showing cylindrical shape of modified parapodia and progression of ventral cirri from subulate to globular form, E) Dorsal view of live specimen from chaetiger 11 – 92 showing fading of colouration from anterior to middle of body, F) Lateral view of live mid-section, G) Bidentate and tridentate falcigers, with minor third tooth encircled, H) Dorsal cirri from chaetiger 82 with small basal process encircled, I) Bidentate acicular chaetae, J) Pectinate chaetae. DC = Dorsal cirri, BR = Branchiae, VCS = Ventral cirri subulate form, VCG = Ventral cirri globular form, C5 = Chaetiger five. Scale bars: A = 1mm; B, C, D, F = 5mm; E = 10mm; G, I = 0.1mm; J = 5µm. A, D, H = MB-A090434; B, E, F = MB-A090442; C, G, J = MB-A090424.




Figure 20

Morphology of *Platynereis latipalpa*

Figure 20. *Perineseis latipalpa* (Schmarda, 1861). A) Anterior region (dorsal), B) Middle region showing red colouration on segments (dorsal), C) Everted pharynx showing Areas 1, 2, 3, 4 (antero-ventral), D) Everted pharynx showing Areas V and VI (dorsal), E) Everted pharynx showing Areas VII-VIII (ventro-lateral), F) Heterogomph spinigers (heS) and homogomph spinigers (hoS) with enlarged teeth at base, G) Heterogomph falcigers with finely serrated blade. Scale bars: A & B, D = 5mm, C & E = 2mm, F - G = 0.1 mm.




Figure 21

Morphology of *Pseudonereis podocirra*

Figure 21. *Pseudonereis podocirra* (Schmarda, 1861). A) Anterior region (dorsal), B) Posterior region, (dorsal), C) Paragnaths on pharynx, Areas I - VIII, D) Compound falciger with serrated blade, E) Compound spiniger with serrated blade, F) 30th chaetiger (dorsal). Scale bars: A - C = 2mm, D & E = 0.1mm, F = 1mm.




Figure 22

Neighbour joining tree of *Pseudonereis*

Figure 22. Neighbour Joining tree of various species belonging to *Pseudonereis* Kinberg, 1865b. * Indicates bootstrap support greater than or equal to 80%. Area highlighted in grey indicates sequences generated in the present study. *Perinereis aibuhitensis* (Grube, 1878) was used as an outgroup. Scale bar represents substitutions per site.

