

# Non-linear genetic diversity and notable population differentiation caused by low gene flow of Bermudagrass [*Cynodon dactylon* (L.) Pers.] along longitude gradients

Jing-Xue Zhang<sup>1,2</sup>, Miaoli Wang<sup>2</sup>, Jibiao Fan<sup>1</sup>, Zhi-Peng Guo<sup>2</sup>, Yongzhuo Guan<sup>2</sup>, Gen Qu<sup>2</sup>, Chuan-Jie Zhang<sup>1</sup>, Yu-Xia Guo<sup>Corresp., 2</sup>, Xuebing Yan<sup>Corresp. 1</sup>

<sup>1</sup> College of Animal Science and Technology, Yangzhou University, Yangzhou, Jiangsu, China

<sup>2</sup> College of Animal and Veterinary Science, Henan Agricultural University, Zhengzhou, Henan, China

Corresponding Authors: Yu-Xia Guo, Xuebing Yan

Email address: guoyuxia@henau.edu.cn, yxbjzz@163.com

**Background.** Environmental variation related to ecological habitat is the main driver of plant adaptive divergence. Longitude plays an important role in the formation of plant population structure, indicating that environmental differentiation can significantly shape population structure.

**Methods.** Genetic diversity and population genetic structure were estimated using 105 expressed sequence tag-derived simple sequence repeat (EST-SSR) loci. A total of 249 *C. dactylon* (L.) Pers. (common bermudagrass) individuals were sampled from 13 geographic sites along the longitude (105°57'34" -119°27'06" E).

**Results.** There was no obvious linear trend of intra-population genetic diversity along longitude and the intra-population genetic diversity was not related to climate in this study. Low gene flow ( $Nm = 0.7701$ ) meant a rich genetic differentiation among populations of *C. dactylon* along longitude gradients. Significantly positive Mantel correlation ( $r = 0.438$ ,  $P = 0.001$ ) was found between genetic distance and geographical interval while no significant partial Mantel correlation after controlling the effect of mean annual precipitation, which indicated geographic distance correlated with mean annual precipitation affect genetic distance. The genetic diversity of *C. dactylon* with higher ploidy level was higher than that with lower ploidy level and groups of individuals with higher ploidy level were separated further away by genetic distance from the lower ploidy levels. Understanding the different genetic bases of local adaptation comparatively between latitude and longitude is one of the core findings in the adaptive evolution of plants.

# Non-linear Genetic Diversity and Notable Population Differentiation Caused by Low Gene Flow of Bermudagrass [*Cynodon dactylon* (L.) Pers.] along Longitude Gradients

Jing-Xue Zhang<sup>1,2</sup>, Miao-Li Wang<sup>2</sup>, Ji-Biao Fan<sup>1</sup>, Zhi-Peng Guo<sup>2</sup>, Yong-Zhuo Guan<sup>2</sup>, Gen Qu<sup>2</sup>, Chuan-Jie Zhang<sup>1</sup>, Yu-Xia Guo<sup>2\*</sup> and Xue-Bing Yan<sup>1\*</sup>

<sup>1</sup>. College of Animal Science and Technology, Yangzhou University, Yangzhou, Jiangsu, China

<sup>2</sup>. College of Animal and Veterinary Science, Henan Agricultural University, Zhengzhou, Henan, China

Corresponding Author:

Yu-Xia Guo<sup>2</sup> and Xue-Bing Yan<sup>1</sup>

63 Nongye Road, Zhengzhou, Henan, 450002, China

NO 48, Wenhui east road, Yangzhou, Jiangsu, 225000, China

Email address: guoyuxia@henau.edu.cn and yxbbjzz@163.com

## Abstract

**Background.** Environmental variation related to ecological habitat is the main driver of plant adaptive divergence. Longitude plays an important role in the formation of plant population structure, indicating that environmental differentiation can significantly shape population structure.

**Methods.** Genetic diversity and population genetic structure were estimated using 105 expressed sequence tag-derived simple sequence repeat (EST-SSR) loci. A total of 249 *C. dactylon* (L.) Pers. (common bermudagrass) individuals were sampled from 13 geographic sites along the longitude (105°57'34" -119°27'06" E).

**Results.** There was no obvious linear trend of intra-population genetic diversity along longitude and the intra-population genetic diversity was not related to climate in this study. Low gene flow ( $N_m = 0.7701$ ) meant a rich genetic differentiation among populations of *C. dactylon* along longitude gradients. Significantly positive Mantel correlation ( $r = 0.438$ ,  $P = 0.001$ ) was found between genetic distance and geographical interval while no significant partial Mantel correlation after controlling the effect of mean annual precipitation, which indicated geographic distance

correlated with mean annual precipitation affect genetic distance. The genetic diversity of *C. dactylon* with higher ploidy level was higher than that with lower ploidy level and groups of individuals with higher ploidy level were separated further away by genetic distance from the lower ploidy levels. Understanding the different genetic bases of local adaptation comparatively between latitude and longitude is one of the core findings in the adaptive evolution of plants.

**Keywords:** *Cynodon dactylon*; Landscape genetics; Genetic diversity; Longitude; Polyploidy.

## Introduction

Plants will undergo adaptive genomic and phenotypic changes in response to environmental heterogeneity on a spatial scale (Yang et al., 2017; Rellstab et al., 2015). Identification of adaptive genetic diversity and population structure of various populations of different environments is of great significance for improving our understanding of the evolutionary process and exploring the adaptive potential of plants (Li et al., 2017). Landscape genetics is helpful for assessing the ability to produce and transmit adaptive genes (Jordan et al., 2017). Genetic differentiation and landscape genetic structure are the results of evolutionary processes including gene migration, random genetic drift and natural selection related to environmental variation due to latitude and longitude. However, gene flow among different populations can effectively prevent genetic difference caused by genetic drift (Schoville et al., 2012; Rundle and Nosil, 2005). Further study of the spatial patterns of genetic diversity is valuable in estimating evolutionary process and predicting the ability of populations to respond to climate change. Studying the genetic structure of more widely distributed plant populations would provide insights into genetic differentiation and evolution. Morphological plasticity and stress tolerances of utilized germplasm should match the environmental characteristics of a specific geographical region (Casler et al., 2004; Boe and Casler, 2005). Population distributions in a large scale could be shaped by regional climate and geographic factors such as latitude and longitude (Peters et al., 2012; Sánchez-gonzález and López-mata, 2005). Longitude is correlated with some environmental variables, and environmental factors (like temperature and precipitation) also play important roles in shaping these assemblages along the longitudes. Longitude was the driving factor behind the distribution of some study species, and the distribution of *Carex physodes* was controlled by longitude (Zhang et al., 2016). Decreasing MAT (mean annual temperature) and MAP (mean annual precipitation), along with

increasing altitude, appear from the east to west along the longitude. Genetic isolation along longitude gradients caused by mountainous environments has been found in other widely distributed Europe-Asian plants (Bartha et al., 2015). Plants in eastern and western Greenland have been genetically different because the Greenlandic ice cap blocks gene flow (Eidesen et al., 2013; Alsos et al., 2009).

Bermudagrass (*Cynodon dactylon* (L.) Pers.) is a genus of the tribe Chlorideae (Poaceae), which inhabits most countries and islands, usually in between about 45°N and 45°S and penetrates to approximately latitude 53°N in Europe (Burton, 1947; Casler et al., 2003). The warm season grass is widely used for turf and pasture because of its drought and heat tolerance and low maintenance requirements (Harlan, 1970; Beard, 1972). *C. dactylon* is an enormously variable and cosmopolitan warm-season turfgrass used for lawns, parks, and sports fields (Shi et al., 2013), which justifies new and innovative approaches toward evaluating and understanding the genetic relationships among different populations at different longitudes. Some studies have been used to analyze genetic diversity and population structure of bermudagrass because of its efficient dispersal and establishment abilities, rapid population growth, and phenotypic plasticity (Linder et al., 2018; Zheng et al., 2017). Identifying the correlation between simple sequence repeat (SSR) markers and turf-quality-related traits among different bermudagrass populations can efficiently help to select high turf quality for bermudagrass. Different ploidy levels including diploid, triploid, tetraploid, pentaploid and hexaploid have been reported for *C. dactylon* (Harlan et al., 1970; Wu and Taliaferro, 2009). Polyploidy can improve competitive ability to adapt to new environments, which is vital for the worldwide distribution of grasses (Te Beest et al., 2012; Linder and Barker, 2014). There have been multiple polyploidy events during plant evolution (Wendel, 2015), so evolutionary significance of polyploidy on wild plant populations is being studied. Polyploidy has been the subject of many studies for the genetic and genomic consequences of WGDs (whole-genome duplication) (Soltis et al., 2016).

The lack of genomic information makes genotyping difficult, but molecular markers enhance the study of plant genetic diversity (Govindaraj et al., 2015). SSRs are characterized by high reproducibility, high polymorphism, co-dominant inheritance, and relative abundance across the entire genome (Vasemägi et al., 2005), and are better selective markers for genetic population analysis of some orphan species (Carneiro Vieira et al., 2016; Hodel et al., 2016). EST-SSRs may reveal some different genetic patterns, because selective processes may affect their polymorphism

(Defaveri et al., 2013). These molecular markers have great potential in genetic diversity and population genetic structure evaluation. Genetic diversity and population structure of 690 *C. dactylon* accessions have been analyzed using expressed sequence tag-simple sequence repeats (EST-SSRs), which is a crucial step of studying genetic evolution (Vlachou et al., 1997; Jewell et al., 2010). Although *C. dactylon* has important research value, the understanding of its population genetic structure and geographic variation is limited. Genetic diversity and genetic structure of *C. dactylon* along a latitudinal gradient across China and the relationship between genetic diversity and ploidy levels were also studied in our previous study (Zhang et al., 2019). Sampling includes maximum geographical variation from flooded east to arid west along the longitude to quantify the relative importance of longitude for adaptive genetic divergence of different bermudagrass populations. The genetic diversity of *C. dactylon* along longitude gradients may be correlated to complex geographic conditions and its biological characteristics including perennial growth, wind-pollination, polyploidy, sexual and asexual reproduction. Different environment factors along longitude gradients can also promote genetic variation and complex genetic structure of *C. dactylon*. Therefore, this study has four objectives (1) quantify population genetic diversity of 249 individuals from 13 populations ranging from different longitudes using EST-SSRs; (2) estimate the spatial population genetic structure and gene flow along longitude; (3) test the relationship between ploidy levels and genetic diversity; (4) compare genetic pattern between longitude and latitude. Studying genetic variation of the wild *C. dactylon* along longitude gradients can promote broadening the genetic resources, understanding the genetic process of adaptation to environmental variation along with longitude and providing some valuable information for the better utilization of wild germplasms of different ploidy levels in breeding.

## Materials & Methods

### Plant materials and DNA extraction

Plant materials consisting of 249 *C. dactylon* individuals were sampled at 13 different geographic origins locating 105°57'34" - 119°27'06" E, China (Fig. 1). Mean annual precipitation and temperature at the collection sites were provided by the China Meteorological Administration (Table 1). Total genomic DNA was extracted using a Zymo Research Plant DNA Extraction Kit (Xinmai jie, Zhengzhou, China) and the quality and concentration of DNA were checked by electrophoresis on 1% agarose gels.

# **EST-SSR amplification**

The 20 EST-SSR primer pairs from previous research (Zhang et al., 2019) produced clear and polymorphic bands and were subsequently used to evaluate genetic diversity for this study. 25 µL PCR amplifications reaction solution contains 1 µL of template DNA, 2.5 µL 10× PCR buffer, 2 µL of 25 mM MgCl<sub>2</sub>, 0.5 µL of 10 mM deoxy nucleoside triphosphate (dNTP), 0.5 µL each of forward and reverse primers, 0.2 µL of Taq DNA polymerase, and 17.8 µL of nuclease-free water.

The amplifications were performed with the following PCR protocol used in the previous research (Zhang et al., 2019): First 1 cycle of 3 min at 95°C, followed by 10 cycles at 95°C, 60°C and 72°C respectively for 30 s, finally terminating with 1 cycle of 6 min at 72°C. Next, the samples were subjected to 1 cycle at 95 °C for 3 min, then 20 cycles of 30 s at 95°C, 55°C, and 72°C respectively, and finally 1 cycle at 72 °C for 6 min. PCR amplified fragments labelled with one fluorescent tag (6-FAM (6-carboxyfluorescein) (blue) and HEX (green)) were detected on a 3730XL Genetic Analyzer (ABI, USA) with the size standard GS500 LIZ. Capillary electrophoresis analysis was made by using GeneMapper Software Version 4.0 (Applied Biosystems, Foster City, CA, USA).

# **Analysis of genetic diversity**

Genetic diversity for all populations and ploidy levels was calculated using POPGENE V. 1.32 Software (Yeh et al., 1997). These indicators included Nei's gene diversity index (He), Shannon diversity index (I) and polymorphic information content (PIC) (Smith et al., 1997; Yeh et al., 1999). He was generated according to genetic distance to analysis genetic diversity. I was computed based on the obtained allele frequencies, and PIC reflected the degree of variation of microsatellite DNA. Ploidy level of each sampled individual was determined by using a flow cytometer (Cube8, Partec, Germany) based on the method outlined by Zhang et al. (2020). Linear association between ploidy levels and longitude was investigated under a regression analysis using the software package SPSS13.0 for Windows (SPSS Inc. Chicago, USA) considering correlation coefficient values and significance of variables in the model. All individuals of the same ploidy level were treated as a population to compare the genetic diversity among different ploidy levels. Significant differences in genetic diversity parameters among different populations and ploidy levels were evaluated under a one-way analysis by Duncan's test using SAS 9.3 software (Institute Inc., Cary, NC, USA) (Duncan, 1955). We divided all the individuals into four groups by longitude

from east to west to calculate the genetic diversity parameters for the four groups with four populations within one group as described above.

### **Analysis of genetic relationship among all populations of *C. dactylon***

Distance matrices of intra-population genetic diversity indicators, environmental distance matrices, genetic distance matrices and geographical distance matrices based on the dataset for 13 populations were calculated using GenAlEx 6.1 (Peakall and Smouse, 2006). Mantel correlation tests (Mantel, 1967) between intra-population genetic diversity indexes and environmental factors (temperature and precipitation) were conducted to examine the effects of environments on the intra-population genetic diversity. Mantel correlation coefficient between genetic distances and longitude distances for *C. dactylon* at different spatial scales was also investigated using NTSYS pc version 2.0 (Khan et al., 2002) and the probabilities for the significance were assessed using Bonferroni inequality adjustment. Partial Mantel test was conducted in R (ver. 4.0.5; R Core Development Team 2011) to estimate correlation between genetic distances and geographic distance, while controlling for mean annual precipitation. Analyses of molecular variance (AMOVA) and population differentiation index ( $F_{ST}$ ) were estimated using ARLEQUIN v3.5 (Excoffier and Lischer, 2010). The significance of  $F_{ST}$  was determined by random resampling of the genotypic data through 1000 permutations. Principal coordinates (PCoA) analysis of genetic variations at different longitudes was performed using GenAlEx Version 6.1. With the help of clustering in NTSYSpc-v.2.1 software, UPGMA (unweighted pair-group method with arithmetic means) was used for cluster analysis to generate a dendrogram for genetic variation along longitude gradients (Rohlf, 2000).

### **Population structure analysis**

The population structure of the 249 individuals was constructed using a Bayesian clustering method implemented in STRUCTURE 2.3 (Hubisz et al., 2009) as previously described in Zhang et al. (2019). The model-based approach implemented was used to subdivide the individuals into different subgroups. Specifically, twenty independent runs were performed for each simulated value of K ranging from 2 to 13. For each run, 10,000 iterations were set as the burn-in time following 100,000 iterations of Markov Chain Monte Carlo (MCMC). Outputs were processed with CLUMPP V1.1.2 (Jakobsson and Rosenberg, 2007) and STRUCTURE barplots were displayed using DISTRUCT 1.1 (Rosenberg, 2004). The best K value was determined from the

modal value of the run with the delta K. Gene flow was calculated using POPGENE V. 1.32 Software (Yeh et al., 1997) based on the formula of  $N_m = 0.5(1 - G_{st})/G_{st}$ .

Mismatch distribution analysis (MDA; Rogers and Harpending, 1992) was used to investigate whether *C. dactylon* had undergone recent population expansion. Goodness-of-fit was tested with the sum of squared deviations (SSD) between observed and expected mismatch distributions (Librado and Rozas, 2009) and Harpending's raggedness index (Harpending, 1994) (HRag), using 1000 parametric bootstrap replicates. The formula  $t = \tau/2u$  was used to estimate the time since expansion began for the expanding population. The value  $u$  was calculated as  $u = \mu kg$ , where  $\mu$  is the substitution rate ( $6.5 \times 10^{-9}$  synonymous substitutions per synonymous site per year (Gaut et al., 1996)),  $k$  is the average sequence length used for analysis (105 bp) and  $g$  is the generation time in years (1 year was used as an approximation for  $g$  according to age of first reproduction of *C. dactylon*). Fu's FS (Fu, 1997) and Tajima's D (Tajima, 1989) neutrality test were used to infer demographic history of each population. Statistical significance was estimated by performing 1000 random permutations. Demographic analyses were performed in ARLEQUIN.

## Results

### Genetic diversity and differentiation along longitude gradient

The genetic diversity parameters including the PIC value,  $H_e$  and  $I$  were calculated for EST-SSR markers in 13 populations (Table 2). The PIC values ranged from a high of 0.68 and a low of 0.40. Highest  $H_e$  value of 0.19 and lowest of 0.06 were calculated. The value of  $I$  ranged from 0.11 to 0.34 respectively for EST-SSR markers. Non-significant difference in genetic diversity parameters among groups (group 1 consisted of populations 1-3; group 2 consisted of populations 4-6; group 3 consisted of populations 7-9; group 4 consisted of populations 10-13) was observed along longitude gradients ( $H_e$  among populations:  $F = 0.027$ ,  $P = 0.974$ ). However, populations 4, 5, 6 and 8 at mid-longitude and populations 1, 12, 13 at lowest and highest longitude had higher genetic diversity than other populations according to  $I$  and  $H_e$  (Fig. 2). In general, there was also no significant correlation ( $P > 0.05$ ) between intra-population genetic diversity and climate (mean annual temperature and precipitation), suggesting that temperature and precipitation showed little influence on intra-population genetic diversity along longitudes.

AMOVA results indicated that 38.23% of the genetic variation was attributed to difference among populations. About 61.77% of the genetic variation showed within population, which

accounted for a greater part of the whole genetic variation (Table 3). Mantel's test revealed significantly positive correlation between genetic distance and longitudinal distance ( $r = 0.276$ ,  $P = 0.022$ ). Environmental measures along longitude gradients (geographical distance ( $r = 0.438$ ,  $P = 0.001$ ); mean annual precipitation ( $r = 0.305$ ,  $P = 0.033$ )) were correlated with genetic distance, while mean annual temperature was not ( $r = 0.006$ ,  $P = 0.418$ ). However, a partial Mantel test accounting for mean annual precipitation was used to estimate that genetic distance was not independently correlated with geographic distance ( $P = 0.146$ ). Gene flow ( $Nm$ ) among these *C. dactylon* populations along longitude gradients was 0.7701 ( $< 1$ ), which mainly reflected low communication frequency of gene among populations.

### Genetic separation by UPGMA dendrogram and principal coordinate analysis

A UPGMA dendrogram dependent on Nei's genetic distance showed that 13 different populations were grouped into three distinct clusters (Fig. 3). The first cluster consisted of populations from Lianyungang and Tancheng that located at high-longitude in the east of China near the Yellow Ocean. Most populations along longitude gradient assigned to the second cluster, so a certain degree of kinship was identified among populations of *C. dactylon* located at different regions along longitude gradient. Populations 7, 9, 10 clustered together and population 3 was close to population 5. The third mixture cluster contained plants coming exclusively from interval locations along the longitude, such as Shanxian (4), Zhengzhou (6), Sanmenxia (8), and Baoji (12). In addition, each of four populations in the mixture cluster had higher intra-population genetic diversity. The Principal coordinate analysis (PCoA) (Fig. 4) revealed a similar pattern as inferred with UPGMA dendrogram. The first, second and third principal coordinates respectively explained 31.57%, 17.78% and 11.90%, altogether 61.25% of the variability. At axis one, 13 populations could be divided into two parts with one of cluster 2 and others of clusters 1, 3. At axis two, cluster 1 consisting of populations 1 and 2 could be clearly separated from cluster 3. Additionally, populations of cluster 1 were clearly far from the other populations. Populations 3, 5, 7, 9 and 10 clustered together in the cluster 2. Populations 4, 6, 8 and 12 with relatively higher genetic diversity formed the cluster 3.

### Population genetic structure of *C. dactylon*

Bayesian analysis of the population structure of *C. dactylon* was used to conduct separate calculation from  $K = 2$  to  $K = 20$ . The results clearly demonstrated that the appropriate  $K$  values were obtained when the 13 populations were subdivided into 4 or 8 groups (Fig. S1). The results

of 13 populations from  $K = 2-13$  strongly indicated different clusters in populations of *C. dactylon* (Fig. 5; Fig. S2). When  $K=4$ , population 1 and 2 clustered together (blue) and the cluster with green consisted of populations 4, 6, 8, 12. Populations 5, 7, 10, 13 were differentiated from populations 3, 9, 11 compared to  $K=3$ . And when  $K=8$ , one group included populations 3, 5, 7, 9, 10, and another group consisted of the populations 8 and 12. Each of the other six groups consisted of only one population respectively, which indicated a rich genetic differentiation along longitude gradients. Admixture among populations of medium longitude (populations 5, 6, 8, 9) could be observed from the structure of *C. dactylon* populations ( $K = 2$  to 13), the clustering of populations at low and high longitude were relatively stable except for the populations 1, 12, 13.

### **Mismatch analysis and neutral test of *C. dactylon***

The mismatch distribution for each population of *C. dactylon* (Table 4) suggested populations 1-11, 13 possibly underwent recent population expansion. The result was also proved by significantly negative values of Fu's  $F_S$  among the 13 populations of *C. dactylon*. Significant SSD values and non-significantly positive Tajima's  $D$  values of the population 12 might suggest relatively stable population size in Baoji. In addition, the demographic expansion time of each population was calculated based on the value of  $\tau$ .

### **Genetic variation and differentiation among different ploidy levels**

Genetic diversity ( $H_e$  and  $I$ ) results of *C. dactylon* with five ploidy levels showed that triploid had significantly lower genetic diversity than pentaploid ( $P = 0.019$  for  $H_e$ ;  $P = 0.008$  for  $I$ ) and hexaploid ( $P = 0.049$  for  $H_e$ ;  $P = 0.020$  for  $I$ ) (Table S1). The genetic diversity of the *C. dactylon* individuals with higher ploidy level was significantly higher than that of individuals with lower ploidy level except for diploid. Significant genetic diversity exists among different ploidy level, but there is no significant difference in genetic diversity between pentaploid and hexaploid based on trend lines of PIC ( $P > 0.05$ ). Diploid and triploid respectively had significantly lower genetic diversity than other polyploidy. All the individuals of *C. dactylon* at different ploidy levels fell into three major clusters based on UPGMA. The first and second clusters contained diploid and triploid individuals, respectively. Tetraploid, pentaploid, and hexaploid individuals formed the third cluster (Fig. 6). The genetic distance between two groups among different ploidy levels increased with the increase of ploidy level. In addition, there was no obvious growth trend of the distribution of ploidy levels, and regression analysis indicated that there was no significant linear association between ploidy level and longitude ( $r = 0.071$ ;  $P > 0.05$  – Fig. 7).

# Discussion

## Genetic diversity and structure among populations along longitude gradients

Identifying relative roles of spatial and environmental factors in shaping patterns of population genetic differentiation is of interest for understanding landscape genetics. Each of 13 populations of *C. dactylon* along longitude gradients had a remarkable genetic variation within population based on the study. Genetic diversity can be improved by sexual propagation, and clonal propagation by rhizome and stolon can promote stable inheritance not being easily affected by genetic drift (Loveless and Hamrick, 1984). Intra-population genetic diversity had no obvious linear relationship with longitude in this study. In addition, mean annual temperature and precipitation at different longitudes had no significant influence on the intra-population genetic diversity. However, according to our previously published paper, higher within-population genetic diversity appeared at lower latitude with higher temperature ( $> 25^{\circ}\text{C}$ ), while relatively low temperature at higher latitude might have little effect on the intra-population genetic diversity (Zhang et al., 2019). So higher temperature ( $> 25^{\circ}\text{C}$ ) may be associated with higher genetic variation of *C. dactylon*. From sampling regions, there was a little higher genetic diversity of populations at mid-longitudes (4, 5, 6, 8) than other populations at low- and high-longitudes, because outcrossing and self-incompatibility of *C. dactylon* could produce low communication among different populations of the east-west transitional locations. Genetic differences among distant geographical sites with unique alleles can increase genetic diversity (Brochmann et al., 2003; Westergaard et al., 2011; Parducci et al., 2012; Eidesen et al., 2013).

Landscape heterogeneity may lead to the formation of local adaptive populations, and different genotypes may be selected to adapt to different environments (Felsenstein, 1976; Rieseberg and Willis, 2007; Anderson et al., 2011). Environmental distance matrices were also calculated to estimate the influences of environmental distance matrices (IBE) on the genetic distance. No significant correlation was observed between genetic distance and geographic distance after controlling the effect of mean annual precipitation, which might be due to a correlation between geographic distance and mean annual precipitation. *P. arundinacea* (Phalaris) also exhibits a certain broad-scale population genetic structure in east-to-west gradients (Perdereau et al., 2017). The adaptive differentiation among populations is influenced by the natural selection of heterogeneous landscape and gene flow (Dionne et al., 2008; Poelchau and Hamrick, 2012). Low

east-to-west inter-population gene flow ( $N_m = 0.7701$ ) indicated that there was little adaptive selective pressure mediated evolutionary dynamics among different populations along longitude gradients (Casler et al., 2007). In addition, these factors to hinder further communication among populations include the transmission mechanism of pollen and seeds of species and the barrier caused by environmental conditions. In our study, restricted seed dispersal may exist among populations at different longitudes exhibiting significant IBD, and mountains in the Henan–Shanxi region may hinder further genetic communication among populations as natural barriers. Alpine plants from several mountain ranges in Europe usually have a strong patterns of longitudinal differentiation (Ronikier et al., 2008; 2012). Geographical isolation along an ecological gradient or environmental isolation can lead to strong genetic differentiation for widespread plants (Hoskin et al., 2005).

The result of AMOVA indicated 61.77% of the genetic variation was attributed to differences within populations and  $F_{ST}$  was 0.3823. Higher  $F_{ST}$  along with increasing distance along longitude than latitude gradients may indicate local genetic differentiation among populations along longitude gradients (Edillo et al., 2009). Both spatial and environmental factors interweave in the formation of population differentiation pattern. In our case, low gene flow among neighboring populations may show the isolation-by-distance pattern among two neighboring populations. Several reproductive barriers in different environments within the same geographical area can block gene flow (Lowry, 2012; Fishman et al., 2013). Interestingly, some populations separated by 200 km in distance are close in genetic distance. A probable explanation is that human-mediated introduction allow contact among these populations. The mismatch distribution test and neutrality test both indicated expansion within population along longitude gradients, with an estimated time around 12.5 Ma (under the demographic expansion model), which may be related to forests shrink and grasslands spread during Miocene time. The dramatic expansion of C4 grasslands in India is a part of increasing tropical C4 savannah vegetation worldwide in the late Miocene, which has been variously ascribed to uplift of the Tibetan plateau (Polissar et al., 2015). In contrast, the population in Baoji with non-significant SSD and HRag values was interpreted as reflecting past demographic stability.

The low number of admixed genotypes at mid-longitude can be explained by the geographical distribution of the collection sites. Natural or human-mediated transmission may have increased genetic diversity and affected the genetic structure of several grass species (Balfourier et al., 2000;

McGrath et al., 2007; Petit et al., 2003; Taberlet et al., 1998). Hybridization at population boundaries and gene migration appear to promote homogenization. Populations at low and high-longitude may evolve much earlier than populations from sites of mid-longitude based on the result. Therefore, populations of *C. dactylon* at mid-longitude may evolve from complex introgressions between populations at low and high-longitude with close genetic affiliation. Historic changes in landscape configuration, spatial isolation and related gene flow changes may gradually lead to intraspecific genetic variation between eastern and western populations of *C. dactylon* in different geographical areas. The admixture among populations determines their potential to adapt to new environmental conditions (Lee, 2002).

### **Relationship between ploidy levels and genetic diversity**

The higher values of  $H_e$ ,  $I$  and  $PIC$  were observed in populations with pentaploid and hexaploid than triploid. Polyploid retains most of the population's gene pool in the form of fixed heterozygosity (Brochmann et al., 2004). Hybridization of polyploid progeny from different parents of the same species can increase the genetic diversity. Multiple introductions from different origins and recombination between different genomes led to genetic mixture. Moreover, groups of individuals with the same ploidy level were separated by genetic distance along with the increasing ploidy levels. The higher genetic distance between two different ploidy levels along with the increasing ploidy levels was observed in the study, which indicated that each chromosome replication process might gradually increase the genetic dissimilarity. Diploid plants can participate in the formation of many polyploid (Ramsey and Schemske, 2002; Otto, 2007), and polyploid are often separated from their progenitors to produce new genotypes (Čertner et al., 2017; Alsayied et al., 2015; Sweigart et al., 2008). Different chromosome numbers could lead to reproductive segregation or sterility, which limited gene flow between hybrids and their parents (Tate et al., 2005). Polyploidy not only produce direct genetic redundancy, but also promote emergence of new gene functions and drive diversification and evolution (Alix et al., 2017). The hybrid genome composition of two related heterotetraploid species was analyzed using molecular cytogenetics to elucidate the role of heteropolyploid in the realization of *Hordeum* diversity (Cuadrado et al., 2017). From the facts above, polyploidization can lead to rich genetic variation and diversification for probable speciation of *C. dactylon*.

### **Comparison of genetic pattern between longitude and latitude**

No significant difference exists in genetic diversity parameters within-population along longitude gradients, while higher within-population genetic diversity appears at low latitude with higher temperature along latitudinal gradients ( $> 25^{\circ}\text{C}$ ) (Zhang et al., 2019). The result showed higher temperature might promote higher within-population genetic variation of *C. dactylon*. The barrier of hybridization of rice mainly focused on the environmental variation, just as temperature influenced cellularization of the endosperm to promote hybrid (Folsom et al., 2014). Higher temperatures may have affected the genetic similarity of *Culex pipiens* populations sampled early and later in the breeding season (Edillo et al., 2009).

A relatively low gene flow ( $N_m = 0.7701$ ) means a rich genetic differentiation along longitude gradients ( $N_m < 1$ ) (Wright, 1965). In our previous study on the genetic diversity and genetic structure of *C. dactylon* along a latitudinal gradient, low genetic differentiation and notable admixture structure existed among populations along latitudes (Zhang et al., 2019). Low genetic differentiation caused by different latitudes may be attributed to the wind-pollination outcrossing system and vegetative materials through human activities, which indicated that the latitude interval could not be an obstacle to genetic exchange. In addition, stronger admixture could be observed along with the latitude in another study (Zhang et al., 2019) than that along with longitude in this study, which indicated relatively lower gene flow among populations of *C. dactylon* occurred along the longitude gradients. North-south direction gene flow has existed between the eastern *Lemus chinense* populations and the western ones (Long et al., 2019). The eastern Eurasian populations of some species with global distribution pattern are a little different from the western Eurasian populations, which occurs between late Miocene and Pleistocene (Tabata et al., 2018; Hardouin et al., 2015; Rodrigues et al., 2014). East Asia was less extensively glaciated and had a relatively mild glacial climate. The Qinghai-Tibetan Plateau uplifted rapidly in Pleistocene and the Asian summer monsoon intensified, so the formation of the inland arid belt of Eurasia resulted in long-term ecological isolation (Clark and Mix, 2002; López-Pujol et al., 2011). A probable explanation is that the genetic differentiation along longitude gradients was little and the gene flow was frequent before the uplift of Tibet Plateau, but the situation reversed after that because populations were ecologically isolated. Therefore making a comparison between latitude and longitude, a stronger gene flow from the north-south than east-west direction is characteristic of genetic structure of *C. dactylon* in China. Distinct genetic divergence of *C. dactylon* presented between the lower and the higher ploidy level along both longitude and latitude gradients. In

addition, widespread polyploidization may cause significantly increasing genetic diversity with the ploidy level.

## Conclusions

A large-scale population genetic survey of genetic diversity of *C. dactylon* has been conducted to evaluate genetic variation and population structure along a longitude gradient. Landscape heterogeneity and geographical isolation along several mountain ranges can lead to formation of locally adapted populations with reduced gene flow between habitats. Geographic distance correlated with precipitation may influenced genetic diversity among populations along longitude gradients. A higher level of genetic differentiation among populations of *C. dactylon* along longitude gradients than those along latitudinal gradients indicated a stronger gene flow from the north-south than east-west direction. Decreased genetic diversity is estimated for individuals with lower ploidy level compared to higher ploidy level, and relatively rich genetic differentiation exists among different ploidy levels. Identifying genetic diversity along longitude gradients in contrast with latitudinal gradients provides important genetic information and improves the ability to protect biodiversity.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was supported by National Natural Science Foundation of China (no. 31702165).

### Competing interests

The authors declare that they have no competing interests.

### Authors Contributions

- Jing-Xue Zhang conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Xue-Bing Yan and Yu-Xia Guo conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Miao-Li Wang, Biao-Ji Fan, Zhi-Peng Guo, Yong-Zhuo Guan, Gen Qu performed the experiments and approved the final draft. Chuan-Jie Zhang checked linguistic accuracy and approved the final draft.

# References

- Alix K, Gérard PR, Schwarzacher T, Heslop-Harrison JS. 2017. Polyploidy and interspecific hybridization: partners for adaptation, speciation and evolution in plants. *Annals of Botany* 120(2):183–194.
- Alsayied N, Fernández JA, Schwarzacher T, Heslop-Harrison JS. 2015. Diversity and relationships of *Crocus sativus* and its relatives analysed by inter-retroelement amplified polymorphism (IRAP). *Annals of Botany* 116:359–368.
- Alsos IG, Torbjorn, A, Normand S, Brochmann C. 2009. Past and future range shifts and loss of diversity in dwarf willow (*Salix herbacea* L.) inferred from genetics, fossils and modelling. *Global Ecology and Biogeography* 18:223–239.
- Anderson JT, Willis JH, Mitchell-Olds T. 2011. Evolutionary genetics of plant adaptation. *Trends in Genetics* 27:258–266.
- Balfourier F, Imbert C, Charmet G. 2000. Evidence for phylogeographic structure in *Lolium* species related to the spread of agriculture in Europe. A cpDNA study. *Theoretical and Applied Genetics* 101:131–138.
- Bartha L, Sramkó G., Volkova PA, Surina B, Ivanov AL, Banciu HL. 2015. Patterns of plastid DNA differentiation in *Erythronium* (Liliaceae) are consistent with allopatric lineage divergence in Europe across longitude and latitude. *Plant Systematics and Evolution* 301(6):1747–1758.
- Beard JB. 1972. Turfgrass: Science and Culture. Prentice-Hall, Englewood Cliffs.
- Boe A, Casler MD. 2005. Hierarchical analysis of switchgrass morphology. *Crop Science*. 45:2465–2472.
- Brochmann C, Brysting AK, Alsos IG, Borgen L, Grundt HH, Scheen AC, Elven R. 2004. Polyploidy in arctic plants. *Biological Journal of the Linnean Society* 82:521–536.
- Brochmann C, Gabrielsen TM, Nordal I, Landvik JY, Elven R. 2003. Glacial survival or tabula rasa? The history of North Atlantic biota revisited. *Taxon* 52:417–450.
- Burton GW. 1947. Breeding bermudagrass for the Southeastern United States. *Agronomy Journal* 39:8.
- Carneiro Vieira ML, Santini L, Diniz AL, de Freitas Munhoz C. 2016. Microsatellite markers: what they mean and why they are so useful. *Genetics and Molecular Biology* 39:312–28.
- Casler MD, Duncan RR. 2003. Turfgrass biology, genetics and breeding. John Wiley & Sons.
- Casler MD, Vogel KP, Taliferro CM, Wynia RL. 2004. Latitudinal adaptation of switchgrass populations. *Crop Science* 44:293–303.
- Casler MD, Vogel KP, Taliaferro CM, Ehlke NJ, Berdahl JD, Brummer EC, Kallenbach RL, West CP, Mitchell RB. 2007. Latitudinal and longitudinal adaptation of switchgrass populations. *Crop Science* 47(6):2249–2260.
- Čertner M, Fenclová E, Kúr P, Kolář F, Koutecký P, Krahulcová A, Suda J. 2017. Evolutionary dynamics of mixed-ploidy populations in an annual herb: dispersal, local persistence and recurrent origins of

473 polyploids. *Annals of Botany* 120:303–315.

474 Clark PU, Mix AC. 2002. Ice sheets and sea level of the Last Glacial Maximum. *Quaternary Science*  
475 *Reviews* 21:1–7.

476 Cuadrado A, de Bustos A, Jouve N. 2017. On the allopolyploid origin and genome structure of the closely related  
477 species *Hordeum secalinum* and *Hordeum capense* inferred by molecular karyotyping. *Annals of Botany*  
478 120:245–255.

479 Defaveri J, Viitaniemi H, Leder E, Merila J. 2013. Characterizing genic and nongenic molecular markers:  
480 Comparison of microsatellites and SNPs. *Molecular Ecology Resources* 13:377–392.

481 Dionne M, Caron F, Dodson JJ, Bernatchez L. 2008. Landscape genetics and hierarchical genetic structure in  
482 Atlantic salmon: the interaction of gene flow and local adaptation. *Molecular Ecology* 17(10):2382–96.

483 Duncan DB. 1955. Multiple Range and Multiple F Tests. *Biometrics* 11:1–42.

484 Edillo F, Kiszewski A, Manjourides J, Pagano M, Hutchinson M, Kyle A, Arias J, Gaines DN, Lampman R,  
485 Novak R, Foppa IM, Lubelczyk C, Smith R, Moncayo A, Spielman A. 2009). Effects of latitude and  
486 longitude on the population structure of *Culex pipiens* sl, vectors of West Nile virus in North America.  
487 *American Journal of Tropical Medicine & Hygiene* 81(5):842–848.

488 Eidesen PB, Ehrich D, Bakkestuen V, Alsos IG, Gilg O, Taberlet P, Brochmann C. 2013. Genetic roadmap of  
489 the Arctic: Plant dispersal highways, traffic barriers and capitals of diversity. *New Phytologist* 200:898–910.

490 Excoffier L, Lischer HEL. 2010. Arlequin suite ver 3.5: a new series of programs to perform population genetics  
491 analyses under Linux and windows. *Molecular Ecology Resources* 10:564–567.

492 Felsenstein J. 1976. The theoretical population genetics of variable selection and migration. *Annual Review of*  
493 *Genetics* 10:253–280.

494 Fishman L, Stathos A, Beardsley PM, Williams CF, Hill JP. 2013. Chromosomal rearrangements and the  
495 genetics of reproductive barriers in *Mimulus* (monkey flowers). *Evolution* 67:2547–2560.

496 Folsom JJ, Begcy K, Hao X, Wang D, Walia H. 2014. Rice fertilization-independent endosperm1 regulates seed  
497 size under heat stress by controlling early endosperm development. *Plant Physiology* 165(1):238–248.

498 Fu YX. 1997. Statistical tests of neutrality of mutations against population growth, hitchhiking and background  
499 selection. *Genetics* 147(2): 915–925.

500 Gaut BS, Morton BR, McCaig BC, Clegg MT. 1996. Substitution rate comparisons between grasses and palms:  
501 synonymous rate differences at the nuclear gene *Adh* parallel rate differences at the plastid gene *rbcl*.  
502 *Proceedings National Academy of Sciences of the United States of America* 93:10274–10279.

503 Govindaraj M, Vetriventhan M, Srinivasan M. 2015. Importance of genetic diversity assessment in crop plants  
504 and its recent advances: an overview of its analytical perspectives. *Genetic Research International* 215:14.

505 Hardouin EA, Orth A, Teschke M, Darvish J, Tautz D, Bonhomme F. 2015. Eurasian house mouse (*Mus*  
506 *musculus* L.) differentiation at microsatellite loci identifies the Iranian plateau as a phylogeographic  
507 hotspot. *BMC Evolutionary Biology* 15:26.

- Harlan JR. 1970. *Cynodon* species and their value for grazing or hay. *Herbage Abstracts* 40: 233–238.
- Harlan JR, de Wet JMJ, Rawal KM, Felder MR, Richardson WL. 1970. Cytogenetic studies in *Cynodon* L.C. Rich. (Gramineae). *Crop Science* 10:288–291.
- Harpending HC. 1994. Signature of ancient population growth in a low-resolution mitochondrial DNA mismatch distribution. *Human Biology* 66(4): 591–600.
- Hodel RG, Segovia-Salcedo MC, Landis JB, Crowl AA, Sun M, Liu X, Gitzendanner MA, Douglas NA, Germain-Aubrey C, Chen SC, Soltis DE, Soltis PS. 2016. The report of my death was an exaggeration: a review for researchers using microsatellites in the 21st century. *Applications in Plant Sciences* 4:17.
- Hoskin CJ, Higgie M, McDonald KR, Moritz C. 2005. Reinforcement drives rapid allopatric speciation. *Nature* 437:1353–1356.
- Hubisz MJ, Falush D, Stephens M, Pritchard JK. 2009. Inferring weak population structure with the assistance of sample group information. *Molecular Ecology Resources* 9:1322–1332.
- Jakobsson M, Rosenberg NA. 2007. CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* 23:1801–1806.
- Jewell MC, Frere CH, Prentis PJ, Lambrides CJ, Godwin ID. 2010. Characterisation and multiplexing of EST-SSR primers in *Cynodon* (Poaceae) species. *American Journal of Botany* 97:e99–e101.
- Jordan R, Hoffmann AA, Dillon SK, Prober SM. 2017. Evidence of genomic adaptation to climate in *Eucalyptus microcarpa*: implications for adaptive potential to projected climate change. *Molecular Ecology* 26(21):6002–6020.
- Khan A, Nawaz M, Khan S, Steele R. 2002. Detection and characterization of erythromycin-resistant methylase genes in Gram-positive bacteria isolated from poultry litter. *Applied Microbiology and Biotechnology* 59:377–381.
- Lee CE. 2002. Evolutionary genetics of invasive species. *Trends in Ecology and Evolution* 17: 386–391.
- Li Y, Zhang XX, Mao RL, Yang J, Miao CY, Li Z, Qiu XY. 2017. Ten years of landscape genomics: challenges and opportunities. *Front Plant Science* 8:2136.
- Librado P, Rozas J. 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25(11): 1451–1452.
- Linder HP, Barker NP. 2014. Does polyploidy facilitate long-distance dispersal? *Botany-Botanique* 113:1175–1183.
- Linder HP, Lehmann CER, Archibald S, Osborne CP, Richardson DM. 2018. Global grass (Poaceae) success underpinned by traits facilitating colonization, persistence and habitat transformation. *Biological Reviews* 93:1125–1144.
- Long X, Weng Y, Liu S, Hao Z, Sheng Y, Guan L, Shi J, Chen J. 2019. Genetic diversity and differentiation of relict plant *Liriodendron* populations based on 29 novel EST-SSR markers. *Forests* 10(4):334.
- López-Pujol J, Zhang FM, Sun HQ, Ying TS, Ge S. 2011. Centres of plant endemism in China: Places for survival or for speciation? *Journal of Biogeography* 38:1267–1280.

- Loveless MD, Hamrick JL. 1984. Ecological determinants of genetic structure in plant populations. *Annual Review of Ecology Evolution and Systematics* 15: 65–95.
- Lowry DB. 2012. Ecotypes and the controversy over stages in the formation of new species. *Biological Journal of the Linnean Society* 106:241–257.
- Mantel N. 1967. The detection of disease clustering and a generalized regression approach. *Cancer Research* 27(2): 209–220.
- McGrath S, Hodkinson TR, Barth S. 2007. Extremely high cytoplasmic diversity in natural and breeding populations of *Lolium* (Poaceae). *Heredity* 9:531–544.
- Otto SP. 2007. The evolutionary consequences of polyploidy. *Cell* 131(3):452–462.
- Parducci L, Jørgensen T, Tollefsrud MM, Elverland E, Alm T, Fontana SL, Alsos IG, Willerslev E. 2012. Glacial survival of boreal trees in northern Scandinavia. *Science* 335(6072):1083–1086.
- Peakall R, Smouse P. 2006. GENALEX 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6:288–295.
- Perdereau A, Klaas M, Barth S, Hodkinson TR. 2017. Plastid genome sequencing reveals biogeographical structure and extensive population genetic variation in wild populations of *Phalaris arundinacea* L. in north-western Europe. *Gcb Bioenergy* 9(1):46–56.
- Peters DPC, Yao J, Sala OE, Anderson JP. 2012. Directional climate change and potential reversal of desertification in arid and semiarid ecosystems. *Global Change Biology* 18:151–63.
- Petit RJ, Aguinalalde I, de Beaulieu JL, Bittkau C, Brewer S, Cheddadi R, Ennos R, Fineschi S, Grivet D, Lascoux M, Mohanty A, Mu G, Demesure-musch B, Palme A, Martı J, Rendell S, Vendramin G. 2003. Glacial refugia: Hotspots but not melting pots of genetic diversity. *Science* 300:1563–1565.
- Poelchau MF, Hamrick JL. 2012. Differential effects of landscape-level environmental features on genetic structure in three codistributed tree species in Central America. *Molecular Ecology* 21(20):4970–4982.
- Polissar PJ, Uno KT, Phelps SR, Pensky JL. 2015. The role of hydroclimate in the late-Miocene expansion of C 4 grasses in India. *AGUFM* 2015: PP21B-2245.
- Ramsey J, Schemske DW. 2007. Neopolyploidy in flowering plants. *Annual Review of Ecology Evolution and Systematics* 2002(33):589–639.
- Rellstab C, Gugerli F, Eckert AJ, Hancock AM, Holderegger R. 2015. A practical guide to environmental association analysis in landscape genomics. *Molecular Ecology* 24(17):4348–4370.
- Rogers AR, Harpending HC. 1992. Population growth makes waves in the distribution of pairwise genetic differences. *Molecular Biology and Evolution* 9: 552–569.
- Ronikier M, Cieřlak E, Korbecka G. 2008. High genetic differentiation in the alpine plant *Campanula alpina* Jacq. (Campanulaceae): evidence for glacial survival in several Carpathian regions and long-term isolation between the Carpathians and the Alps. *Molecular Ecology* 17:1763–1775.
- Ronikier M, Schneeweiss GM, Schönswetter P. 2012. The extreme disjunction between Beringia and Europe in

- Ranunculus glacialis* s.l. (Ranunculaceae) does not coincide with the deepest genetic split—a story of the importance of temperate mountain ranges in arctic–alpine phylogeography. *Molecular Ecology* 21:5561–5578.
- Rieseberg LH, Willis JH. 2007. Plant speciation. *Science* 317:910–914.
- Rodrigues ASB, Silva SE, Marabuto E, Silva DN, Wilson MR, Thompson V, Yurtsever S, Halkka A, Borges PAV, Quartau JA, Paulo OS, Seabra SG. 2014. New mitochondrial and nuclear evidences support recent demographic expansion and an atypical phylogeographic pattern in the spittlebug *Philaenus spumarius* (Hemiptera, Aphrophoridae). *PLoS ONE* 9:e98375.
- Rohlf FJ, Rohlf FJ, Rohlf F, Rohlf FJ. 2000. NTSYS-pc numerical taxonomy and multivariate analysis system version 2.1. Owner manual.
- Rosenberg NA. 2004. DISTRUCT: a program for the graphical display of population structure. *Molecular Ecology Notes* 4:137–138.
- Rundle HD, Nosil P. 2005. Ecological speciation. *Ecology Letters* 8(3):336–352.
- Sánchez-gonzález A, López-mata L. 2005. Plant species richness and diversity along an altitudinal gradient in the Sierra Nevada, Mexico. *Divers Distribution* 11:567–75.
- Schoville S, Bonin A, Francois O, Lobreaux S, Melodelima C, Manel S. 2012. Adaptive genetic variation on the landscape: methods and cases. *Annual Review of Ecology Evolution and Systematics* 43:23–43.
- Shi H, Ye T, Chan Z. 2013. Exogenous application of hydrogen sulfide donor sodium hydrosulfide enhanced multiple abiotic stress tolerance in bermuda grass (*Cynodon dactylon* (L). Pers.). *Plant Physiology Biochemistry* 71: 226–234.
- Smith JSC, Chin ECL, Shu H, Smith OS, Wall SJ, Senior ML, Mitchell SE, Kresovich S, Ziegler J. 1997. An evaluation of the utility of SSR loci as molecular markers in maize (*Zea mays* L.): comparisons with data from RFLPs and pedigree. *Theoretical and Applied Genetics* 95:163–173.
- Soltis DE, Visger CJ, Marchant DB, Soltis PS. 2016. Polyploidy: pitfalls and paths to a paradigm. *American Journal of Botany* 103:1146–1166.
- Sweigart AL, Martin NH, Willis JH. 2008. Patterns of nucleotide variation and reproductive isolation between a *Mimulus* allotetraploid and its progenitor species. *Molecular Ecology* 17:2089–2100.
- Tabata R, Kawaguchi F, Sasazaki S, Yamamoto Y, Bakhtin M, Kazymbet P, Meldevekob A, Suleimenov MZ, Nishibori M, Mannen H. 2018. The Eurasian Steppe is an important goat propagation route: A phylogeographic analysis using mitochondrial DNA and Y-chromosome sequences of Kazakhstani goats. *Animal Science Journal* 90:317–322.
- Taberlet P, Fumagalli L, Wust-Saucy AG, Cosson JF. 1998. Comparative phylogeography and postglacial colonization routes in Europe. *Molecular Ecology* 7:453–464.
- Tajima F. 1989. Statistical-method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123(3): 585–595.

- Tate JA, Soltis DE, Soltis PS. 2005. CHAPTER 7–Polyploidy in Plants. *The Evolution of the Genome* 2005:371–426.
- Te Beest M, Le Roux JJ, Richardson DM, Brysting AK, Suda J, Kubešová M, Pyšek P. 2012. The more the better? The role of polyploidy in facilitating plant invasions. *Annals of Botany* 109:19–45.
- Vasemägi A, Nilsson J, Primmer CR. 2005. Expressed sequence tag-linked microsatellites as a source of gene-associated polymorphisms for detecting signatures of divergent selection in atlantic salmon (*Salmo salar* L.). *Molecular Biology and Evolution* 22:1067–1076.
- Vlachou D, Konsolaki M, Tolias PP, Kafatos FC, Komitopoulou K. 1997. Conserved synteny, amplification and tissue specificity but sequence divergence and altered temporal regulation. *Genetics* 147:1829–1842.
- Wendel JF. 2015. The wondrous cycles of polyploidy in plants. *American Journal of Botany* 102:1753–1756.
- Westergaard KB, Alsos IG, Popp M, Engelskjøn T, Flatberg KI, Brochmann C. 2011. Glacial survival may matter after all: Nunatak signatures in the rare European populations of two west-arctic species. *Molecular Ecology* 20:376–393.
- Wright S. 1965. The interpretation of population structure by F-Statistics with special regard to systems of mating. *Evolution* 19:395.
- Wu Y, Taliaferro CM. 2009. Bermuda grass. In: Genetic resources, chromosome engineering and crop improvement. CRC Press, New York 5:289.
- Yang J, Miao CY, Mao RL, Li Y. 2017. Landscape population genomics of forsythia (*Forsythia suspensa*) reveal that ecological habitats determine the adaptive evolution of species. *Front in Plant Science* 8:481.
- Yeh FC, Yang RC, Boyle T, Ye ZH, Mao JX. 1997. POPGENE, the user friendly shareware for population genetic analysis. Molecular Biology and Biotechnology Center, University of Alberta: Edmonton, AB, Canada.
- Yeh FC, Young RC, Boyle T. 1999. Microsoft Window-based Freeware for Population Genetic Analysis (POPGENE, Ver. 1.31). <ftp://ftp.microsoft.com/softlib/mslfiles/hppl.exe>
- Zhang JX, Wang ML, Guo ZP, Guan YZ, Guo YX, Yan XB. 2018. Variations in morphological traits of bermudagrass and relationship with soil and climate along latitudinal gradients. *Hereditas* 155(1):31.
- Zhang JX, Wang ML, Guo ZP, Guan YZ, Liu JY, Yan XB, Guo YX. 2019. Genetic diversity and population structure of bermudagrass [*Cynodon dactylon* (L.) Pers.] along latitudinal gradients and the relationship with polyploidy level. *Diversity* 11(8):135.
- Zhang JX, Wang ML, Guo ZP, Guan YZ, Guo YX, Yan XB. 2020. Variation in ploidy level and genome size of *Cynodon dactylon* (L.) Pers. along a latitudinal gradient. *Folia Geobotanica* 2020:1–12.
- Zhang R, Liu T, Zhang JL, Sun QM. 2016. Spatial and environmental determinants of plant species diversity in a temperate desert. *Journal of Plant Ecology* 9(2):124–131.
- Zheng Y, Xu S, Liu J, Zhao Y, Liu J. 2017. Genetic diversity and population structure of Chinese natural bermudagrass [*Cynodon dactylon* (L.) Pers.] germplasm based on SRAP markers. *PLoS ONE* 12(5):

e0177508.

# Figure legends

Fig. 1 Locations of sampled populations of *C. dactylon*. The blue dots represent the sampling sites along longitude gradients.

Fig. 2 Summary statistics of within-population genetic variation of 13 *C. dactylon* populations from eastern to western. He: Nei's gene diversity index; I: Shannon diversity index.

Fig. 3 The UPGMA (unweight pair-group method with arithmetic means) dendrogram generated cluster analysis based on EST-SSR data of 13 *C. dactylon* populations. The *C. dactylon* populations in this study were grouped into three clusters (blue, red, green).

Fig. 4 Principal component analysis of the 13 populations based on the genetic distance derived from EST-SSR markers. Populations in this study were grouped into three clusters (black circles).

Fig. 5 Population structure analysis of 249 *C. dactylon* individuals sampled from 13 populations using the model-based program STRUCTURE as follows: **A**-K = 4 and **B**-K = 8. Each color shows one Bayesian ancestry group (K) and the length of the colored segment represents the estimated membership proportions in different Bayesian ancestry groups (K).


Fig. 6 The UPGMA (unweight pair-group method with arithmetic means) dendrogram of bermudagrass groups with different ploidy level.

Fig. 7 Regression analysis between ploidy level of bermudagrass (*C. dactylon*) and longitude.

# Figure 1

Locations of sampled populations of *C. dactylon*.


Fig. 1 Locations of sampled populations of *C. dactylon*. The blue dot represents the sampling sites along longitudinal gradient.



# Figure 2

Summary statistics of within-population genetic variation of 13 *C. dactylon* populations from eastern to western.


Fig. 2 Summary statistics of within-population genetic variation of 13 *C. dactylon* populations from eastern to western. He: Nei's gene diversity index; I: Shannon diversity index.



# Figure 3

The UPGMA (unweight pair-group method with arithmetic means) dendrogram generated cluster analysis based on EST-SSR data of 13 *C. dactylon* populations.


Fig. 3 The UPGMA (unweight pair-group method with arithmetic means) dendrogram generated cluster analysis based on EST-SSR data of 13 *C. dactylon* populations. The *C. dactylon* populations in this study were grouped into three clusters (blue, red, green).



# Figure 4

Principal component analysis of the 13 populations based on the genetic distance derived from EST-SSR markers.


Fig. 4 Principal component analysis of the 13 populations based on the genetic distance derived from EST-SSR markers. Populations in this study were grouped into three clusters (black circles).



# Figure 5

Population structure analysis of 249 *C. dactylon* plants sampled from 13 populations using the model-based program STRUCTUREas follows: A-K = 4 and B-K = 8.


Fig. 5 Population structure analysis of 249 *C. dactylon* plants sampled from 13 populations using the model-based program STRUCTUREas follows: **A**-K = 4 and **B**-K = 8. Each color shows one Bayesian ancestry group (K) and the length of the colored segment represents the estimated membership proportions in different Bayesian ancestry groups (K).



# Figure 6

The UPGMA (unweight pair-group method with arithmetic means) dendrogram of bermudagrass groups with different ploidy level.


Fig. 6 The UPGMA (unweight pair-group method with arithmetic means) dendrogram of bermudagrass groups with different ploidy level.



# Figure 7

Regression analysis between ploidy level of bermudagrass (*C. dactylon*) and longitude.

Fig.7 Regression analysis between ploidy level of bermudagrass (*C. dactylon*) and longitude.



**Table 1**(on next page)

Information on geographical sites and climate of *C. dactylon* populations collected from different longitudes in China.

Table 1 Information on geographical sites and climate of *C. dactylon* populations collected from different longitudes in China.

Code	Region	Habitat	Longitude (E)	Latitude (N)	Altitude (m)	MAT (°C)	MAP (mm)
1	Lianyungang	Island	119°27'06"	34°46'09"	50	14.5	883.9
2	Tancheng	Roadside	118°16'08"	34°38'37"	30	13.8	832.9
3	Zaozhuang	Roadside	117°49'20"	34°38'48"	89	14.4	820.3
4	Shanxian	Roadside	116°09'11"	34°46'31"	30	14.2	621.4
5	Lankao	Roadside	114°44'55"	34°49'32"	60	14.3	631.1
6	Zhengzhou	Swamp	113°38'20"	34°54'04"	90	14.7	640.8
7	Luoyang	Roadside	112°19'30"	34°43'20"	210	14.4	637.2
8	Sanmenxia	Roadside	111°03'49"	34°42'29"	340	14	558.1
9	Tongguan	Roadside	110°13'18"	34°33'41"	540	13.1	602.9
10	Jingyang	Roadside	108°50'07"	34°32'32"	410	13.5	504.1
11	Fufeng	Roadside	107°52'41"	34°20'35"	570	12.8	569.9
12	Baoji	Ditch side	107°41'03"	34°21'54"	630	13.5	645.9
13	Tianshui	Roadside	105°57'34"	34°32'43"	1050	11.4	500.7

MAT, mean annual temperature; MAR, mean annual precipitation

4

**Table 2**(on next page)

Within-population genetic diversity of 13 *C. dactylon* populations at different longitudes.

1 Table 2. Within-population genetic diversity of 13 *C. dactylon* populations at different  
2 longitudes.

Pop	N	He	I	PIC
1	18	0.1939	0.2895	0.55
2	20	0.1367	0.2231	0.56
3	20	0.1157	0.1872	0.47
4	20	0.1837	0.2769	0.57
5	20	0.1843	0.2878	0.68
6	18	0.1859	0.279	0.53
7	19	0.0619	0.1131	0.40
8	16	0.1809	0.271	0.52
9	20	0.0984	0.1617	0.43
10	20	0.1338	0.2142	0.51
11	20	0.1603	0.2493	0.57
12	18	0.231	0.3408	0.62
13	20	0.1818	0.2779	0.57
Total	249	0.2587	0.4029	1.00

3 He, Nei's gene diversity index; I, Shannon's diversity index; PIC, Polymorphic information  
4 content

**Table 3**(on next page)

Analysis of molecular variance (AMOVA) of based on 249 individuals from the 13 populations of bermudagrass.

1 Table 3. Analysis of molecular variance (AMOVA) of bermudagrass of different population.

Source of variation	df	Sum of squares	Variance components	Percentage of variation
Among populations	12	1336.521	5.364	38.23%
Within populations	236	2045.455	8.667	61.77%
Total	248	3381.976	14.031	

2 Note: df, Degree of freedom

3

4

5

6

# **Table 4**(on next page)

Results of the mismatch distribution analysis and neutrality tests of *C. dactylon* of each population along longitudinal gradients.

Results of the mismatch distribution analysis and neutrality tests of *C. dactylon* of each population along longitudinal gradients. SSD, sum of squared deviation under expansion model; HRI, raggedness; \*Significant ( $P \leq 0.05$ ); \*\*highly significant ( $P \leq 0.01$ ).

1 Table 4 Results of the mismatch distribution analysis and neutrality tests of *C. dactylon* of each population along longitude gradients.

2

Populations	Tau ( $\tau$ )	Time since expansion began(t, Ma)	SSD	P-value	HRI	P-value	Fu's FS	P-value	Tajima's D	P-value
1	29.500	21.612	0.006	0.960	0.013	0.820	-5.704	0.017*	1.152	0.916
2	0.000	0.000	0.013	1.000	0.017	0.940	-9.177	0.000**	-0.370	0.377
3	2.000	1.465	0.008	0.870	0.011	0.810	-10.399	0.001**	-0.299	0.435
4	16.000	11.722	0.005	0.390	0.009	0.610	-7.288	0.002**	0.813	0.838
5	24.500	17.949	0.010	0.650	0.020	0.280	-7.270	0.007**	0.072	0.559
6	21.200	15.531	0.008	0.540	0.011	0.880	-5.904	0.013*	1.116	0.888
7	0.400	0.293	0.012	0.700	0.019	0.800	-14.636	0.000**	-1.731	0.033*
8	30.400	22.271	0.023	0.340	0.020	0.670	-4.805	0.017*	0.944	0.881
9	2.600	1.905	0.011	0.390	0.009	0.830	-11.688	0.000**	-0.570	0.301
10	8.700	6.374	0.008	0.540	0.011	0.680	-9.330	0.002**	-0.116	0.495
11	21.500	15.751	0.004	0.900	0.007	0.960	-8.116	0.004**	0.194	0.619
12	35.300	25.861	0.021	0.000	0.020	0.500	-4.927	0.020*	1.493	0.940
13	28.900	21.172	0.009	0.780	0.016	0.560	-7.349	0.002**	0.762	0.840
<b>Mean</b>	17.003	12.457	0.011	0.620	0.014	0.718	-8.199	0.007**	0.266	0.625

3 SSD, sum of squared deviation under expansion model; HRI, raggedness; \*Significant ( $P \leq 0.05$ ); \*\*highly significant ( $P \leq 0.01$ ).