

Who's who in *Magelona*? Phylogenetic hypotheses under Magelonidae Cunningham & Ramage, 1888 (Annelida: Polychaeta) (#61394)

1

First submission

Guidance from your Editor

Please submit by **4 Jun 2021** for the benefit of the authors (and your \$200 publishing discount) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

18 Figure file(s)

3 Table file(s)

1 Raw data file(s)



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Who's who in *Magelona*? Phylogenetic hypotheses under Magelonidae Cunningham & Ramage, 1888 (Annelida: Polychaeta)

Kate Mortimer^{Corresp., 1}, Kirk Fitzhugh², Ana Claudia Brasil³, Paulo Lana⁴

¹ Natural Sciences, Amgueddfa Cymru – National Museum Wales, Cardiff, Wales, United Kingdom

² Natural History Museum of Los Angeles County, Los Angeles, California, United States of America

³ Departamento de Biologia Animal, Instituto de Ciências Biológicas e da Saúde, Universidade Federal Rural do Rio de Janeiro, Seropédica, Rio de Janeiro, Brazil

⁴ Centro de Estudos do Mar, Universidade Federal do Paraná, Pontal do Sul, Paraná, Brazil

Corresponding Author: Kate Mortimer

Email address: Katie.Mortimer@museumwales.ac.uk

Known as shovel head worms, members of Magelonidae comprise a group of polychaetes readily recognised by the uniquely shaped, dorso-ventrally flattened prostomium and paired ventro-laterally inserted papillated palps. The present study is the first published account of inferences of phylogenetic hypotheses within Magelonidae. Members of 72 species of *Magelona* and two species of *Octomagelona* were included, with outgroups including members of one species of Chaetopteridae and four of Spionidae. The phylogenetic inferences were performed to causally account for 176 characters distributed among 79 subjects, and produced 2,417,600 cladograms, each with 404 steps. A formal definition of Magelonidae is provided, represented by a composite phylogenetic hypothesis explaining seven synapomorphies: shovel-shaped prostomium, prostomial ridges, absence of nuchal organs, ventral insertion of palps and their papillation, presence of a burrowing organ, and unique body regionation. *Octomagelona* is synonymised with *Magelona* due to the latter being paraphyletic relative to the former. The consequence is that Magelonidae is monotypic, such that *Magelona* cannot be formally defined as associated with any phylogenetic hypothesis. As such, the name is an empty placeholder, recognised only because of the binomial name requirement mandated by the International Code of Zoological Nomenclature.

Who's who in *Magelona*? Phylogenetic hypotheses under Magelonidae Cunningham & Ramage, 1888 (Annelida: Polychaeta)

Kate Mortimer¹, Kirk Fitzhugh², Ana Claudia Brasil³ and Paulo Lana⁴

¹ Amgueddfa Cymru – National Museum Wales, Cardiff, CF10 3NP, Wales, United Kingdom

² Natural History Museum of Los Angeles County, Los Angeles, California, United States of
America

³ Departamento de Biologia Animal, Instituto de Ciências Biológicas e da Saúde, Universidade
Federal Rural do Rio de Janeiro, 23890-000 Seropédica, RJ, Brazil

⁴ Centro de Estudos do Mar, Universidade Federal do Paraná, 83255-000 Pontal do Sul, Paraná,
Brazil

Corresponding author:

Kate Mortimer¹

Amgueddfa Cymru – National Museum Wales, Cardiff, CF10 3NP, Wales, United Kingdom

Email address: Katie.Mortimer@museumwales.ac.uk

ABSTRACT

Known as shovel head worms, members of Magelonidae comprise a group of polychaetes readily recognised by the uniquely shaped, dorso-ventrally flattened prostomium and paired ventro-laterally inserted papillated palps. The present study is the first published account of inferences of phylogenetic hypotheses within Magelonidae. Members of 72 species of *Magelona* and two species of *Octomagelona* were included, with outgroups including members of one species of Chaetopteridae and four of Spionidae. The phylogenetic inferences were performed to causally account for 176 characters distributed among 79 subjects, and produced 2,417,600 cladograms, each with 404 steps. A formal definition of Magelonidae is provided, represented by a composite phylogenetic hypothesis explaining seven synapomorphies: shovel-shaped prostomium, prostomial ridges, absence of nuchal organs, ventral insertion of palps and their papillation, presence of a burrowing organ, and unique body regionation. *Octomagelona* is synonymised with *Magelona* due to the latter being paraphyletic relative to the former. The consequence is that Magelonidae is monotypic, such that *Magelona* cannot be formally defined as associated with any phylogenetic hypothesis. As such, the name is an empty placeholder, recognised only because of the binomial name requirement mandated by the International Code of Zoological Nomenclature.

Subjects Evolutionary Hypotheses, Invertebrates, Philosophy of Science, Systematics, Zoology

Keywords Abductive reasoning, Cladograms, Phylogenetic Hypothesis, Systematics, Zoological Nomenclature

INTRODUCTION

Felix qui potuit rerum cognoscere causas – Virgil, *Georgics*, Vol. 1, Books I–II

The Magelonidae, known commonly as the shovel head worms, gain their name from their uniquely flattened, spade-shaped prostomia used in burrowing through soft sediments. They comprise a relatively small family of marine annelids, with members distributed among 72 species, although many more individuals to which new species hypotheses will refer are likely. Magelonids are generally found at depths of less than 100 m, although members of deeper water species at depths of 1000–4000 m have been recorded (Hartman, 1971; Aguirrezabalaga et al., 2001). They predominately burrow through sands and muds, although members of several tubicolous species are known (Mills & Mortimer, 2019; Mortimer, 2019). Individuals feed using two slender palps (Mortimer & Mackie, 2014), which are unique amongst polychaetes in being papillated and ventrally inserted.

Magelonids were given the rank of family by Cunningham & Ramage (1888) but their unusual morphology has often led to difficulties in relating them to other annelid groups. Johnston (1865), puzzled by their peculiar external form, placed them at the end of his catalogue, under the family Maeadae. McIntosh, who published extensively on magelonid morphology (McIntosh, 1878, 1879, 1911), suggested similarities with spioniforms such as members of *Prionospio* Malmgren, 1867, and *Heterospio* Ehlers, 1874, but also with the chaetopterid *Spiochaetopterus* Sars, 1853. He additionally noted that the way the ‘proboscis’ (cf. **Character descriptions** below regarding terminology) operated and the structure of the “snout and circulatory organs” are features *sui generis*. However, the placement of magelonids with spioniform polychaetes has continued until semi-recently. The latest studies have proposed placement alongside the Chaetopteridae, Sipuncula, and Oweniidae (Struck et al., 2015; Weigert

et al., 2014; Helm et al., 2018), or as sister taxon of all other annelids along with the Oweniidae (Palaeoannelida) (Weigert et al., 2016). These phylogenetic considerations will be discussed more fully later in this paper.

Whilst the systematic position of the family may have received some attention, phylogenetic relationships within have received far less. Perhaps the relatively uniform bodies of magelonids have been a contributing factor. Currently the family contains two genera: the type genus *Magelona* F. Müller, 1858, and the monotypic *Octomagelona* Aguirrezabalaga, Ceberio & Fiege, 2001, differentiated by the number of thoracic chaetigers. All previously introduced generic names for the group have been synonymised (*Maea* Johnston, 1865, *Rhynophylla* Carrington, 1865, *Meredithia* Hernández-Alcántara & Solís-Weiss, 2000). Brasil (2003) inferred phylogenetic hypotheses among members of the family based on external morphological characters, confirming the monophyletic status of the group. Her analysis concluded that both *Octomagelona* and *Meredithia* were paraphyletic, but no further conclusions could be made. The present study aims to expand the unpublished analysis performed by Brasil (2003) to include additional taxa and characters.

MATERIAL AND METHODS

Methodological considerations

What is and is not contained in the present study is based on several interrelated principles that initially might not appear consistent with some of today's thinking regarding biological systematics. These principles are not of our making, but rather are firmly established perspectives within philosophy of science, and familiar to many fields of science beyond systematics. The last several decades have witnessed systematics engaging in a growing

tendency toward developing insular views that are at odds with basic tenets of logic, reasoning, and scientific inquiry. For these reasons, we regard it as **imperative** to present justifications for what is and is not included in this study from a methodological perspective. The disinterested reader can ignore this section without damage, but the information provided does offer a useful summary for those more inclined toward critical thinking in systematics.

The goal of scientific inquiry

As systematics is a field of science and subfield of evolutionary biology, it should cohere with the acknowledged goal of scientific inquiry. That goal is to not only describe phenomena but also pursue causal understanding of what is encountered (Hanson, 1958; Hempel, 1965; Rescher, 1970; Popper, 1983, 1992; Salmon, 1984b; Van Fraassen, 1990; Strahler, 1992; Mahner & Bunge, 1997; Hausman, 1998; Thagard, 2004; Nola & Sankey, 2007; de Regt et al., 2009; Hoyningen-Huene, 2013; Potochnik, 2017, 2020; Currie, 2018; Anjum & Mumford, 2018). In the context of systematics that pursuit includes describing the characteristics of organisms, in the form of differentially shared characters, as well as exploring possible past causal events that account for those characters, either as matters of proximate and/or ultimate causes *sensu* Mayr (1961, 1993; Fitzhugh, 2012, 2016a). As aptly expressed by Uller & Laland (2019: 1), “Scientific inference typically relies on establishing causation. This is also the case in evolutionary biology, a discipline charged with providing historical accounts of the properties of living things, as well as an understanding of the processes that explain the origin of those properties.”

Whilst it is fashionable to speak of ‘*the* phylogeny’ of a group of organisms, this is something of a misnomer. Inferring phylogenetic hypotheses from a set of observed character data only achieves the objective of explaining those data, with implicit acknowledgement that no

phylogenetic inference can exhaustively account for all potentially observable characters of organisms. Phrases of the form ‘*the* phylogeny of *X*’ incorrectly connote that a final solution is being offered, when in fact such would be impossible. Emphasising the term phylogeny has the undesirable consequence of suggesting that this is the objective of systematics, thus overshadowing the pursuit of causes, which is the real aim of scientific inquiry. The same critique applies to the commonly used phrase ‘molecular phylogeny.’ There can be no ‘phylogeny’ of molecules, just as there can be no separate phylogenies for any arbitrary subdivisions of classes of characters that are in need of being explained.

The causal objective in systematics is the arena within which taxa are considered. Taxa are best regarded not as either class constructs, mere groupings, things, entities, or ontological individuals, but rather as the variety of explanatory hypotheses routinely and purposely inferred. These hypotheses include species and phylogenetic hypotheses, albeit there are other classes of explanatory hypotheses that in their own right deserve to be called taxa (Fig. 1; cf. Hennig, 1966: fig. 6) given that they as well are inferred causal accounts (Fitzhugh, 2012, 2016a; [Fitzhugh](#), 2005b, 2006a, 2008a, 2009, 2013, 2015). Consider for instance the phylogenetic and specific ‘relations’ shown in Fig. 1. Each clearly illustrates that individuals along the top of the diagram are what exist in the present, and lines connecting to lower individuals indicate past tokogenetic events. This diagram also implies past, albeit quite vague, causes such as novel character origin and fixation events among individuals in populations, as well as population splitting events. To say Fig. 1 shows several ‘lineages’ is just an imprecise way of referring to the totality of past causal events that account for what are observed of the organisms in the present. A further consequence is that taxa are neither described, discovered, nor delimited; they are inferred as reactions to what we observe of the properties of organisms. What *are* described are individuals

at specific moments in their life history (Fitzhugh, 2012, 2016a, 2008a, 2009, 2013, 2015) (operational examples include Fitzhugh, 2010b; ~~Fitzhugh et al., 2015~~; Nogueira et al., 2010, 2013, 2017, 2018); what Hennig (1966) termed semaphoronts (Fig. 1; cf. Hennig, 1966: fig. 6). The interplay between the descriptive and the explanatory will become apparent later (cf. *Phylogenetic inference = abduction*).

The present study seeks to infer taxa in the form of phylogenetic hypotheses, which are subsumed under the more inclusive (composite) phylogenetic hypothesis formally called Magelonidae. We acknowledge that species hypotheses have been previously and separately inferred and are not the focus of this paper. Whilst this distinction between phylogenetic and specific hypotheses has been the typical and inferentially appropriate approach in systematics, exceptions can be found among some publications (e.g., Nygren et al., 2018; Shimabukuro et al., 2019; Radashevsky et al., 2020), in which species hypotheses are simultaneously inferred with phylogenetic hypotheses that only causally account for sequence data, after which morphological characters are incorrectly introduced in a post hoc manner. Notwithstanding the fact that the requirement of total evidence (RTE; Fitzhugh, 2006b; see *Sequence data and explanatory hypotheses*, below) is violated, such inferences have questionable merits for the fact that explaining shared nucleotides or amino acids requires, at a minimum, discriminating between causes such as genetic drift, and selection via downward causation (see *Sequence data and explanatory hypotheses*, below). The nature of phylogenetics algorithms is such that they are agnostic as to detailed causal parameters, requiring that one must give specific consideration to drift or selection, which can determine whether or not sequence data can be directly explained via phylogenetic inferences. This is an issue that has been largely ignored in discussions of

phylogenetic inference (Fitzhugh, 2016b). Once again, the objective of inferring these different classes of hypotheses is to pursue causal understanding of the properties of organisms.

Characters and observation statements

Scientific inquiry is in reaction to the objects and events we perceive in the universe. Conceptualising and then communicating our perceptions require accurate conveyance of observation statements. Systematics has had a long tradition of speaking of organismal observations in terms of characters and states (Sokal & Sneath, 1963; Sneath & Sokal, 1973). Fitzhugh (2006c) remarked on the fact that the character/state distinction as well as other, similar perspectives in systematics are not accurate representations of observation statements, especially compared to the established views in epistemology, where observation statements are presented as relations between subjects and predicates (Strawson, 1971; Alston, 1993; Audi, 1998). Consider for instance, the statement, ‘These chaetae are bidentate.’ The terms ‘chaetae’ and ‘bidentate’ do not refer to character and state, respectively. That one can observe objects they call chaetae is because of the properties or characters of those objects. Instead of character and state, the observation statement refers to the *subject* ‘chaetae,’ and the *predicate* ‘bidentate’ is applied to that subject, reflecting a particular property or character perceived of the object as subject (Hanson, 1958; Mahner & Bunge, 1997; Strawson, 1971; Gracia, 1988; Armstrong, 1997).

Whilst observation statements of organisms are often of intrinsic properties, there also are relational or extrinsic characters (Findlay, 1936; Sider, 1996; Armstrong, 1997; Francescotti, 2014; Allen, 2016). A common example in systematics involves sequence data, e.g., ‘Individuals to which species hypothesis *X-us x-us* refer have [subject] nucleotide G at [predicate] position 546.’ Being in a particular position is relative to other objects. The character/state distinction

ignores this relational aspect, leading to nonsensical characterisations, such as, “...if nucleotide A is observed to occur at position 139 in a sequence, ‘position 139’ is the character [sic] and ‘A’ is the state assigned to that character” (Swofford et al., 1996: 412). Obviously, a position cannot be a character of an object since the position is dependent upon other objects.

The traditional emphasis on ‘character coding’ leading to compilations of observations in the form of the data matrix has treated columns as ‘characters’ and each cell as a ‘state.’ Fitzhugh (2006c: fig. 1) pointed out, however, that if observations are to be accurately implied by entries in a matrix then each cell represents a complete observation statement of subject-predicate relations, such that each column denotes each subject (Fig. 2A). This also accommodates relational characters (Fitzhugh, 2016b). But with the emphasis on causal inquiry in systematics, a data matrix is not only a codified representation of observation statements. The matrix must also imply the why-questions that inquiry seeks to answer in the form of explanatory hypotheses known as taxa. This necessitates knowing the formal structure of those questions and how they, like observation statements, are implied by the matrix. This will be addressed next.

Basis for phylogenetic inference: why-questions

The inferences of explanatory hypotheses vis-à-vis taxa do not appear *ex nihilo*. As a significant role of scientific inquiry is the pursuit of causal understanding, there is a vital conceptual link between our observations and explanatory hypotheses. That link exists through the implicit or explicit why-questions that are prompted by observations, such that the hypotheses we infer are intended as answers to those questions. The formal structure of why-questions has relevance to systematics for the fact that those questions are implicitly present in a character data matrix provided to a computer algorithm that infers phylogenetic hypotheses (Fitzhugh, 2006c, 2016b).

Whilst we tend to think of why-questions as having the form, ‘Why (is) q (the case)?,’ such a structure is an incomplete representation. The form of why-questions is instead what is often referred to as contrastive, ‘Why (is) q (the case) in contrast to p ?’ (Salmon, 1984b; Sober, 1984, 1986, 1994; Salmon, 1989; Van Fraassen, 1990; Lipton, 2004; Fitzhugh, 2006a, 2006b, 2006c; 2016b; Lavelle et al., 2013). A contrastive why-question contains observation statement q , the *fact* to be explained, and p , the *foil*, which is usually a condition ordinarily expected or previously explained. Asking a why-question often occurs because we are faced with a situation that is unexpected or surprising (the *fact*), and it is that element of surprise, against what is expected (the *foil*), that leads to the desire for an explanation. The fact/foil distinction associated with implied why-questions can be found in the data matrices used to infer phylogenetic hypotheses, when ‘outgroup’ (= *foil*) and ‘ingroup’ (= *fact*) taxa are designated (Fitzhugh, 2006c, 2016b) (Fig. 2B).

It is often acknowledged (Bromberger, 1966; Sober, 1986, 1988; Barnes, 1994; Marwick, 1999; Sintonen, 2004; Schurz, 2005) that why-questions include the presupposition that fact and foil statements must be true, otherwise there is no basis for asking such questions. This also follows from the common-sense notion that we assume the truth of our observation statements; otherwise, scientific inquiry would not be feasible. This assumption has special implications for explaining sequence data when rates of substitution are involved, which will be addressed later (see *Sequence data and explanatory hypotheses*). Explaining a fact entails also explaining the foil, and while those explanations are due to separate causal events, they should be due to the same type of cause (Sober, 1986; Barnes, 1994; see also Cleland, 2001, 2002, 2009, 2011, 2013; Turner, 2007; Jeffares, 2008; ~~Tucker, 2004, 2011~~). Designations of outgroup and ingroup accommodate this requirement (Fitzhugh, 2006c).

226 *Forms of reasoning*

227 In speaking of inferring phylogenetic hypotheses, or any other class of taxon for that matter, it is
 228 necessary to acknowledge the type of reasoning used to produce those hypotheses as well as the
 229 reasoning available for empirically testing them. This provides a clear basis for establishing
 230 which, if any, of the currently available phylogenetic inference procedures are logically and
 231 scientifically sound.

232 Three forms of reasoning are often recognised: abduction, deduction, and induction.
 233 Among formal treatments of logic, abduction, if acknowledged at all, is typically subsumed
 234 under induction (e.g., Kneale & Kneale, 1964; Salmon, 1967, 1984a; Copi & Cohen, 1998),
 235 since the purview of logic is to identify and discriminate logically valid (deductive: true premises
 236 give a true conclusion) from invalid or fallacious (inductive: true premises do not guarantee true
 237 conclusions) arguments. For purposes of explicating the processes of scientific inquiry, however,
 238 it is appropriate to regard abduction as distinct from induction. Operationally each form of
 239 reasoning plays a respective role in different stages of inquiry, where those stages include at a
 240 minimum the inferences of theories and hypotheses, and the subsequent empirical testing of
 241 those propositions. The stage of inquiry most associated with systematics is abduction (Peirce,
 242 1878, 1931, 1932, 1933a, 1933b, 1934, 1935, 1958a, 1958b; Hanson, 1958; Achinstein, 1970;
 243 Fann, 1970; Reilly, 1970; Curd, 1980; Nickles, 1980; Thagard, 1988; Josephson & Josephson,
 244 1994; Baker, 1996; Hacking, 2001; Magnani, 2001, 2009, 2017; Psillos, 2002, 2007, 2011;
 245 Godfrey-Smith, 2003; Norton, 2003; Walton, 2004; Gabbay & Woods, 2005; Aliseda, 2006;
 246 Schurz, 2008; Park, 2017), i.e., the inferences of hypotheses as the means of causally accounting
 247 for the various differentially shared characters observed among organisms (Fitzhugh, 2005a,
 248 2005b, 2006a, 2006b, 2008a, 2008b, 2008c, 2009, 2010a, 2013, 2014, 2015, 2016a, 2016b,

2016c, 2016d, 2021). Subsequent empirical assessments of explanatory hypotheses rely on deducing predictions of expected consequences – potential test evidence – if the causal claims in hypotheses are true. The act of testing, which involves determining manifestations of test evidence, is inductive. The operational relations between ab-, de-, and induction received extensive attention in the 19th and early 20th centuries by the polymath, Charles Sanders Peirce (1835–1914) (Peirce, 1931, 1932, 1933a, 1933b, 1934, 1935, 1958a, 1958b), but full appreciation of Peirce’s ideas did not become realised until the second half of the 20th century (Hanson, 1958; Fann, 1970; Reilly, 1970; Thagard, 1988; Josephson & Josephson, 1994; Magnani, 2001, 2009, 2017; Psillos, 2002, 2011; Walton, 2004; Gabbay & Woods, 2005; Aliseda, 2006; Schurz, 2008; Park, 2017). As noted by Peirce (1932: 2.106),

Abduction... is merely preparatory. It is the first step of scientific reasoning, as induction is the concluding step. Abduction makes its start from the facts, without, at the outset, having any particular [hypothesis] in view, though it is motivated by the feeling that a [hypothesis] is needed to explain the surprising facts. Induction makes its start from a hypothesis which seems to recommend itself, without at the outset having any particular facts in view, though it feels the need of facts to support the [hypothesis]. Abduction seeks a [hypothesis].

Induction seeks for facts.

The ‘facts’ Peirce refers to in relation to abduction are the various characteristics of organisms in need of explanation, whereas the ‘facts’ associated with induction are test evidence. Discussions of relations between these stages of inquiry in systematics can be found in Fitzhugh (2005a, 2006a, 2006b, 2008b, 2010a, 2012, 2013, 2016a, 2016c, 2016d).

Whilst systematics in recent decades, especially beginning with the school of thought called ‘cladistics,’ has laid great emphasis on the testing of hypotheses, usually at the expense of conflating testing with hypothesis inference (reviewed by Fitzhugh, 2016c, 2016d). Actual testing by way of induction of the variety of hypotheses implied by cladograms is rarely performed. Instead, the tendency is to merely replace results of previous abductions with new abductions as new observed effects become available, which is neither an act of testing nor a basis for claiming that previous hypotheses have been ‘defeated’ or overturned. Associated with confusing hypothesis inference with testing has been the misconception that character data used to infer hypotheses also provide evidential support for those hypotheses, leading to false claims that phylogenetic hypotheses are ‘robust’ or ‘strongly supported’ by the characters used to infer those hypotheses. We will address this misunderstanding later (see *Explanatory hypotheses and the myth of evidential support*).

In the context of phylogenetic hypotheses, examples of each of the stages of inquiry can be represented by the following abbreviated forms, where the premises lie above the double (non-deductive) or single (deductive) line, and the conclusion(s) allowed by those premises below the line(s):

(1) Abduction – inferring hypotheses as answers to why-questions:

- background knowledge, b
- theory(ies) t , such as ‘common ancestry’
- observed effects, as differentially shared characters, e

- explanatory hypotheses, e.g., cladograms, h

(2) Deduction – predictions of consequences given the truth of hypotheses:

- background knowledge, b

- theory(ies) t relevant to the observed effects
 - specific causal conditions presented in explanatory hypothesis via **(1)**
 - proposed conditions needed to perform test
-
- observed effects e , originally prompting h [cf. **(1)**]
 - *predicted test evidence, i.e., effects related as closely as possible with the specific causal conditions of the hypothesis*

(3) Induction – hypothesis testing:

- background knowledge, b
 - theory(ies) relevant to observed effects, t
 - test conditions performed
 - confirming/disconfirming evidence, e_2 [observations of predicted test evidence in **(2)**, or alternative observations]
-
- h is confirmed/disconfirmed.

A useful way to think of these inferences is as reactions to particular questions. The form of each question determines what type of inference is used to produce a conclusion that serves as an answer. Respective questions and inferences are shown in Table 1.

Several points of logic need to be highlighted regarding the three types of reasoning, as they are relevant to inferences of phylogenetic hypotheses (in fact all taxa, for that matter). For any inference to be deemed *factually correct*, premises must be assumed to be true. A deduction is *valid* only if true premises guarantee a true conclusion, which is established by particular rules of logic that can be found in standard logic textbooks. Because of those rules, the content of a deductive conclusion is already present in the premises, such that information in the conclusion

cannot go beyond what is already stated in the premises (Salmon, 1984a). Such reasoning is said to be non-ampliative. By its very nature, deduction cannot introduce new ideas. All non-deductive reasoning is ampliative, thus does not satisfy the criterion of logical validity. This makes abduction a fundamentally important mode of reasoning in everyday life as well as all of science. Abduction is the only type of reasoning from which new ideas are conceived. With true premises, abductive and inductive reasoning do not guarantee true conclusions. The content of those conclusions will extend beyond what is provided in the premises. Abductive conclusions can be regarded as merely plausible: “By plausibility, I mean the degree to which a theory [or hypothesis] ought to recommend itself to our belief independently of any kind of [test] evidence other than our instinct urging us to regard it favorably” (Peirce, 1958b: 8.223). Inductive conclusions, on the other hand, are probable given that they are determined by available test evidence.

Phylogenetic inference = abduction

Representing phylogenetic inference in a more complete form compared to what is shown in (1), the following example of abduction is somewhat closer to the actual abductive structure leading to phylogenetic hypotheses. The basis for this abduction would be to answer the why-question, ‘Why do semaphoronts to which specific hypotheses *x-us* and *y-us* refer have ventrolateral margins with appendages in contrast to smooth as seen among individuals to which other species hypotheses (*a-us*, *b-us*, etc.) refer?’ (Fitzhugh, 2006c, 2012, 2013, 2015, 2016a, 2016b):

(4) • *Background knowledge, b*

• *Phylogenetic theory*: If character $x(0)$ exists among individuals of a reproductively isolated, gonochoristic or cross-fertilising hermaphroditic population and character $x(1)$ originates by mechanisms $a, b, c \dots n$, and becomes fixed within the population by

mechanisms $d, e, f \dots n$ (= ancestral species hypothesis), followed by event(s) $g, h, i \dots n$, wherein the population is divided into two or more reproductively isolated populations, then individuals to which descendant species hypotheses refer would exhibit $x(1)$.

- *Observations (effects)*: Individuals to which specific hypotheses x -us and y -us refer have ventrolateral margins with appendages in contrast to smooth as seen among individuals to which other species hypotheses (a -us, b -us, etc.) refer.

- *Causal conditions (phylogenetic hypothesis X -us)*: Ventrolateral margin appendages originated by some unspecified mechanism(s) within a reproductively isolated population with smooth ventrolateral margins, and the appendage condition became fixed in the population by some unspecified mechanism(s) (= ancestral species hypothesis), followed by an unspecified event(s) that resulted in two or more reproductively isolated populations.

Background knowledge b , as auxiliary theories and hypotheses, is generally not listed as a premise, but instead is established knowledge accepted as true that is needed for the other premises in an inference. Background knowledge is specified here for reference later when considering abduction in relation to sequence data. The major premise, *Phylogenetic theory*, is intentionally vague regarding causal specifics. This reflects the near-complete absence of causal information that goes into the development of phylogenetic algorithms as well as the lack of explanatory content in phylogenetic hypotheses implied by cladograms (Fitzhugh, 2012, 2013, 2016b). The theory as presented here is intentionally limited to gonochoristic or cross-fertilising hermaphroditic organisms (Fitzhugh, 2013) since population-splitting events (‘speciation’) are not relevant to asexually reproducing, obligate parthenogenetic, or self-fertilising hermaphroditic

organisms. The theory states a common cause, which is consistent with the necessary presupposition that observation statements are true, mentioned earlier; if we assume the truth of our observations, then we should pursue explanations of those observations in a manner that maximises the truth element as much as possible in those explanatory accounts. The minor premise in this example, *Observations (effects)*, presents only one set of observation statements, as subject-predicate relations, i.e., ventrolateral body margins. This premise can be expanded to include other relevant characters to be explained, as is typical of the standard character data matrix. A data matrix implies both why-questions as well as the totality of observation statements comprising this second premise (Fitzhugh, 2006c, 2016b). The conclusion, *Causal conditions*, is the result of applying the *Phylogenetic theory* to *Observations* of effects. Such conclusions, often represented by cladograms, serve as answers to the why-questions. The limited causal details provided in the conclusion is a consequence of what little is offered in the theory in terms of causal particulars. For this reason, cladograms are largely devoid of explanatory information. As part of the ampliative nature of abduction, the premises can lead to a single conclusion or multiple, mutually exclusive conclusions. Any valid deduction, on the other hand, being non-ampliative, will only produce a single conclusion.

Whilst we often refer to a cladogram as a phylogenetic ‘hypothesis,’ they are in fact composite constructs, implying a minimum of three classes of hypotheses: previously inferred species hypotheses, hypotheses of character origin/fixation among members of ancestral populations, and population splitting events (‘speciation’) (Fig. 3). A fourth, ad hoc class of hypothesis is typically invoked, called homoplasy (Lankester, 1870).

To manifest abductive inferences in a phylogenetics computer algorithm that emulate what is presented in (4), the inferential form would be akin to the following (cf. Josephson & Josephson, 1994),

- (5)
- d is a collection of (presumed true) observation statements of differentially-shared characters
 - hypotheses, implied by cladograms $X_1, X_2 \dots X_n$, are possible composite explanations of d
 - select subset of cladograms with score S_n (minimal ‘tree length’ = minimise ad hoc hypotheses of homoplasy)

-
-
- cladograms with score S_n are most plausible, mutually exclusive explanations of d .

The algorithm functions in a manner that produces conclusions *as if* the common cause *Phylogenetic theory* in (4) were operative. Notice as well that the first premise in (5) contains the assumption that observation statements regarding shared characters are true; the necessary presupposition of truth associated with implied why-questions in data matrices, if one accepts that the intent of these inferences is to explain observations. The phrase in the conclusion, ‘cladograms with score S_n are most plausible,’ is intentional. Plausibility indicates hypotheses are worthy of further consideration:

By Plausible, I mean a theory [or hypothesis] that has not yet been subjected to any test, although more or less surprising phenomena have occurred which it would explain if it were true, is in itself of such a character as to recommend it for further examination or, if it be *highly* plausible, justify us in seriously inclining

toward belief in it, as long as the phenomena be inexplicable otherwise (Peirce, 1932: 2.662, emphasis original; Kapitan, 1997: 481). Epistemically, a plausible hypothesis is far weaker than a hypothesis that has been successfully tested with the introduction of test evidence, i.e., (2), (3); a distinction that is too often ignored in the systematics literature because of the general misunderstanding of the meaning of evidential support (see *Phylogenetic hypotheses and the myth of evidential support*).

At first sight, the algorithmic representation of abduction in (5) might appear to be what has often been called ‘parsimony analysis.’ The difficulty with this perspective is that parsimony is not a form of reasoning that produces explanatory hypotheses. That responsibility resides with abduction. Parsimony is a comparative measure, where for instance of two hypotheses that predict the same results, the hypothesis that requires fewer assumptions is the simpler of the two. The third premise in (5) invokes this principle. That intended action of selection, based on minimising ad hoc hypotheses of homoplasy, is implemented for the sole purpose of serving as a surrogate for the application of the *Phylogenetic theory* in (4). This is not tantamount to a parsimony analysis or method. It is an instance of abductive reasoning.

It was noted earlier that abduction is the only type of reasoning that produces explanatory hypotheses; it is the only form of reasoning that introduces new ideas. The depiction of abduction in (4), or its algorithmic form in (5), is not only at odds with ‘parsimony’ being an inference procedure but also contrary to claims in systematics that so-called ‘likelihood’ or ‘Bayesian’ methods can infer explanatory hypotheses. With his introduction of the likelihood principle, Fisher (1934; see also Hacking, 1965, 2001, Edwards, 1972) was clear that it is only relevant to considerations of test evidence *subsequent* to inferring hypotheses. Evidence e in representations of likelihood, e.g., $p(h | e) = l(h | e)$, refer to *test evidence* [cf. (2), (3)].

Probability p quantifies anticipation of an outcome of testing, whereas likelihood l quantifies trust in the hypothesis as a consequence of testing. Characters used to abductively infer phylogenetic hypotheses [cf. (1), (4), (5)] do not constitute test evidence, and likelihood does not provide the means to infer explanatory hypotheses (*contra* e.g., Felsenstein, 1981, 2004; Huelsenbeck & Crandall, 1997; ~~Swofford et al., 1996~~; Haber, 2011).

The intent of Bayes Theorem is to quantify changes in belief in hypotheses following the introduction of test evidence (Salmon, 1967; Howson & Urbach, 1993; Hacking, 2001; *contra* e.g., Huelsenbeck & Ronquist, 2001; Huelsenbeck et al., 2001; Archibald et al., 2003; Felsenstein, 2004; Ronquist et al., 2009). Bayesianism is inductive *sensu* [3]; it has no relevance to abduction. The attempt to force Bayes' Theorem into phylogenetic inference (e.g., Huelsenbeck et al., 2001: fig. 1) is represented as,

$$p(\text{tree} \mid \text{characters}) = \frac{p(\text{characters} \mid \text{tree}) \cdot p(\text{tree})}{p(\text{characters})}.$$

This is not a meaningful implementation of the Theorem (Fitzhugh, 2010a, 2012, 2016a, 2016c, 2016d). The supposed prior probability, $p(\text{tree})$, refers to all possible 'tree topologies' for taxa considered. Such topologies are nothing but branching diagrams devoid of the empirical content typical of cladograms that imply the several classes of causal events discussed earlier (Fig. 2). Tree diagrams disconnected from characters explained by those diagrams cannot be assigned prior probabilities. More problematic is the use of differentially shared characters as 'evidence' to determine posterior probabilities. Character data cannot serve as test evidence for the hypotheses intended to explain those data (see *Phylogenetic hypotheses and the myth of evidential support*). As with 'parsimony' and 'likelihood' methods, the 'Bayesian' approach in systematics fails to live up to its name.

In sum, neither ‘parsimony,’ ‘likelihood,’ nor ‘Bayesianism’ are valid forms of reasoning or methods for inferring explanatory hypotheses. Unfortunately, wider recognition of the fundamental importance of abductive reasoning did not take hold among scientists until Hanson (1958) pointed out its significance in his *Patterns of Discovery*, and later acknowledgment of its importance in artificial intelligence (Thagard, 1988) and science in general were embraced (Fann, 1970; Reilly, 1970; Thagard, 1988; Josephson & Josephson, 1994; Magnani, 2001, 2009, 2017; Psillos, 2002, 2011; Walton, 2004; Gabbay & Woods, 2005; Klien hans et al., 2005; Aliseda, 2006; Schurz, 2008; Klien hans et al., 2010; Park, 2017; Anjum & Mumford, 2018). In the absence of awareness about abduction, systematics since the 1960’s was left with an inferential void that was filled with incorrect applications under the false dichotomy of parsimony versus induction vis-à-vis statistics, where the latter led to incorrectly applying likelihood and Bayesianism. What is actually being performed in phylogenetic inference under these labels is abductive reasoning, albeit in a most peculiar and not entirely philosophically or scientifically acceptable manner.

Regarding abductive inferences of phylogenetic hypotheses to explain sequence data, there are two significant issues to consider. One issue will be discussed here, while the other will be addressed later (see *Sequence data and explanatory hypotheses*). The treatment of sequence data has often been seen as requiring the assumption of rates of substitution as part of the inferences of phylogenetic hypotheses, along with ‘branch lengths’ (i.e., number of substitutions or character changes that have occurred ‘along’ the branches of phylogenetic trees) (Swofford et al., 1996; Felsenstein, 2004; Schmidt & von Haeseler, 2009). Once we acknowledge that trees are only graphic devices for implying the various classes of explanatory hypotheses accounting for observed characters (Fig. 2), it becomes apparent that those trees cannot have properties like

lengths of branches. The need to consider substitution rates has warranted the development of so-called ‘model-based’ approaches (Jukes & Cantor, 1969; Felsenstein, 1978, 2004; Farris, 1983; Swofford et al., 1996; Sullivan & Joyce, 2005; Rindal & Brower, 2011; Kapli et al., 2020), under the guises of ‘likelihood’ and ‘Bayesian,’ versus the ‘model-free’ method of ‘parsimony analysis.’ It should be apparent by now that this distinction, just like the methods, is without merit. If we accept that phylogenetic inferences are abductive, then the inclusion of substitution rates within abduction would have the following form (Fitzhugh, 2021):

(6) *Background knowledge* (partim): Observation statements assumed true, per alignments of sequence data

Theories: (a) Rates of sequence substitution, applied to sequence data;

(b) Common cause *Phylogenetic theory* of common ancestry, cf. (4), per

Background knowledge

Observation statements: Represented in data matrix as why-questions



Phylogenetic hypotheses: Diagrammatically implied by graphic devices called cladograms.

Notice that this inference relies on theories that are contradictory. The theory of rates of sequence substitution is at odds with the *Background knowledge* of true observation statements of aligned sequences, and by extension also at odds with the implied why-questions one would ask for which the abduction seeks answers using a common-cause theory. Correcting for this contradiction would necessitate removing the theory of substitution rates as a premise in the abduction and placing it within the *Background knowledge*, such that originally aligned sequence data would be modified to take substitution rates into consideration. By doing this, revised

observation statements could be presented, which re-establish assumptions that observation statements are true. The *Background knowledge* in (6) would then have the following revised form (Fitzhugh, 2021),

(7) *Background knowledge, revised:*

- (a) observation statements of differentially shared characters, are initially assumed true, per alignments of sequence data;
- (b) rates of sequence substitution taken into consideration;
- (c) per (b), the assumption of truth of observation statements in (a) is not necessarily correct;
- (d) per (b) and (c), observation statements are revised such that apparent shared nucleotides are renamed as *different* observation statements where applicable to re-establish true statements.

With the revised *Background knowledge* in (7), the ensuing abductive inference would then have the form shown in (4), which again provides answers to why-questions by way of common cause; there is no need to consider substitution rates within the inference. The inherent difficulty of accomplishing step (7)(d), however, will likely preclude even proceeding forward with causally accounting for those sequence data given that there are no empirical criteria for discerning identical nucleotides or amino acids as not being identical. This is a matter that has been overlooked in systematics. The alternative solution is to simply accept the truth of shared characters obtained through alignment, in which case abduction would again be as shown in (4) and neither substitution rates nor branch lengths would be considered.

We have emphasised that cladograms or phylogenetic trees satisfy the goal of scientific inquiry if they are interpreted as composite explanatory hypotheses (Fig. 3). Whilst the

explanatory nature of cladograms is sometimes explicitly acknowledged (e.g., Farris, 1983; Swofford et al., 1996; Schmidt & von Haeseler, 2009; Kapli et al., 2020), often the greater focus of phylogenetic studies is on just obtaining ‘trees,’ with little to no causal considerations given (e.g., Felsenstein, 1981, 2004; Sullivan & Joyce, 2005; Kapli et al., 2020). Consequences of this emphasis on trees as opposed to explanations includes an erroneous method called character mapping (Fitzhugh, 2014). Mapping most often involves obtaining phylogenetic trees using sequence data, then ‘optimising’ other characters, usually morphological, onto these diagrams, from which are determined vague evolutionary conclusions regarding the mapped characters. This approach has become popular among polychaete phylogenetic studies (e.g., Struck et al., 2011; Borda et al., 2012; Glasby et al., 2012; Goto et al., 2013; Weigert et al., 2014; Aguado et al., 2015; Andrade et al., 2015; Struck et al., 2015; Goto, 2016; Weigert & Bleidorn, 2016; Kobayashi et al., 2018; Nygren et al., 2018; Langeneck et al., 2019; Shimabukuro et al., 2019; Radashevsky et al., 2020; San Martín et al., 2020; Tilic et al., 2020). Much credence has been given to mapping, albeit without foundation. For instance, Kapli et al. (2020) state that,

Mapping heritable character states (phenotypic or genotypic) onto a tree is the basis of different evolutionary analyses: it allows us, for example, to make inferences [sic] about character homology and also to gain insights into character loss and convergent evolution. Whereas trees were initially based to a great extent on morphological characters, biological molecules – nucleic acids and proteins – provide a far more powerful [sic] and plentiful source of information for reconstructing trees.

Mapping is flawed for a simple reason. That phylogenetic inference is abductive means inferring hypotheses, implied by a cladogram, for one set of characters produces conclusions relevant only

to those characters. Any subsequent ‘mapping’ of additional characters on that cladogram is not a legitimate form of inference, abductive or otherwise. The premises of a ‘mapping inference’ make this apparent:

- (8)
- tree topology, (*a-us (b-us (c-us (d-us, e-us))))*)
 - observed effects, as differentially shared characters, *e*
-
- ???

As a premise, a tree topology is an empirically empty statement since it would have been previously inferred to account for characters other than those in the second premise. Attempting to apply that first premise to the second is pointless since this cannot lead to any interpretable conclusion – certainly not explanatory hypotheses. As such it is impossible to speak of explanations of mapped characters. Like the past and present uncritical advocacy of phylogenetic methods called ‘parsimony,’ ‘likelihood,’ and ‘Bayesian,’ the popularity of character mapping is a further testament to the lack of consideration of formal reasoning in systematics.

Phylogenetic hypotheses and the myth of evidential support

The present study does not assess the veracity of either the Magelonidae or less inclusive phylogenetic hypotheses in terms of evidential ‘support,’ nor are there attempts to use popular methods such as the bootstrap or Bremer analysis to assert that hypotheses are either ‘strongly supported’ or ‘robust.’ Such perspectives have become all too common in systematics research, including those on annelids (e.g., Struck et al., 2011; Borda et al., 2012; Weigert et al., 2014; Aguado et al., 2015; Andrade et al., 2015; Struck et al., 2015; Weigert et al., 2016; Gonzalez et al., 2018; Nygren et al., 2018; Shimabukuro et al., 2019; San Martín et al., 2020; Stiller et al., 2020; Tilic et al., 2020). Reasons for not speaking of support in the present study follow from the basics of reasoning described above. More detailed treatments of misconceptions regarding

evidence of, and support for phylogenetic hypotheses can be found in Fitzhugh (2012, 2016b, 2016c).

First, to correctly speak of *evidence* means to refer to the premises that allow for a certain conclusion (Longino, 1979; Salmon, 1984a; Achinstein, 2001). Whilst that evidence is said to ‘support’ a given conclusion, it is only inductive, i.e., test evidence or support (cf. [3]) that matters in the pursuit of empirical evaluation, whether in matters of everyday life or during scientific inquiry. An emphasis on test evidence certainly permeates all of evolutionary biology and by extension it is understandable that it would be of concern in systematics. But this requires being clear about (1) what is meant by the test evidence, (2) if that evidence is being used in the proper context, and (3) whether or not bootstrap and Bremer ‘support’ values are meaningfully interpretable in relation to phylogenetic hypotheses.

Recall that the examples of ab-, de-, and induction in (1)–(3) draw clear distinctions between premises and conclusions derived from those premises. Those premises constitute the *evidence for* the respective conclusions [see also (4)–(6)]. Beyond basic matters of logic, we need to be cognisant of the way evidence is interpreted in the separate stages of inquiry. Evidence in relation to abduction is unremarkable in the sense that conclusions must be as they are because of the premises [cf. (1), (4), (5)] (Fitzhugh, 2010a, 2012, 2016a, 2016c, 2016d). As well, abductive conclusions cannot be changed or ‘defeated’ with the introduction of additional effects to be explained. Adding those effects as premises only results in the inference of a new set of hypotheses that have no relevance to previously inferred hypotheses.

A related misconception is that statistical consistency applies to phylogenetic inference (e.g., Felsenstein, 1978, 2004; Swofford et al., 1996; Heath et al., 2008; Assis, 2014; Brower, 2018). When correctly interpreted, consistency is the view that as more and more *test evidence* is

introduced in support of a hypothesis, the closer one is supposedly getting to a true proposition. The mistaken interpretation of consistency in systematics has led to the popular yet erroneous view that sequence data are beneficial for discerning ‘phylogenies’ because of the vast increase in characters that are available (e.g., Goto, 2016; Tilic et al., 2020). For example, in relation to phylogenetic inference, Felsenstein (2004: 107, 121, respectively) states,

“An estimator is consistent if, as the amount of data gets larger and larger (approaching infinity), the estimator converges to the true value of the parameter with probability 1...”

“The inconsistency of parsimony [sic] has been the strongest challenge to its use. It becomes difficult to argue that parsimony methods have logical and philosophical priority, if one accepts that consistency is a highly desirable property.”

Contrary to claims that consistency is relevant to phylogenetic inference, it has been acknowledged since the early 20th century that continued additions of effects to abductive inferences provides no indication that a true, as opposed to plausible, set of explanatory hypotheses has been attained (Peirce, 1902, 1932: 2.774–777; Rescher, 1978; Fitzhugh, 2012, 2016b). As new effects in need of explanation are sequentially added to abductive premises, each subsequent conclusion will only offer a new set of hypotheses that have no relevance to previously inferred hypotheses. The emphasis on consistency in systematics is yet another consequence of abduction not being recognised in lieu of a misplaced statistical (inductive) mindset.

Deduction serves to conclude predictions of potential test evidence [cf. (2)] that might be later sought during the process of testing. Much like premises-as-evidence in abduction, the

premises in deduction do not warrant any special attention in terms of ‘support.’ Where evidence is of importance is during the act of testing, i.e., induction [cf. (3)]. For instance, a conclusion that test evidence supports or confirms a hypothesis can be subsequently revised if additional support through further testing is obtained, or additional test evidence might eventually lead to a conclusion of disconfirmation. As noted by Lipton (2005), the process of testing puts hypotheses at risk of being disconfirmed since there is no guarantee predicted test evidence, via (2) and (3), will be found. Obtaining contrary evidence is always a possibility. It is test evidence that matters when speaking of support. Abductive ‘support’ lacks that qualification, and character data are not test evidence.

Because an emphasis on causality has been largely wanting in recent decades in relation to inferences of taxa-as-explanatory-hypotheses, references to support for phylogenetic hypotheses has centered on support [sic] for ‘groups’ or ‘clades’ within cladograms (Fitzhugh, 2012). Such claims of support are misleading for two reasons. First, as mentioned above, ‘support’ for any abductive conclusion is trivial since the conclusion could not be otherwise given the premises. In other words, unlike test evidence associated with induction, observations of characters do not present any risk to the hypotheses explaining those characters. To assert, for example, that characters ‘support’ some phylogenetic hypothesis verges on circularity: the conclusion(s) is/are determined by the premises and the premises support the conclusion(s). Second, our concern is not for groups, but rather various explanatory hypotheses. As all phylogenetic hypotheses are composite (Fig. 3), an emphasis on groups is both artificial and denies consideration of the variety of explanatory accounts implied by cladograms. Support in relation to groups is epistemically meaningless.

There is a third difficulty, faced by the two popular ‘support’ measures for phylogenetic hypotheses [sic] (Efron, 1979; Efron & Gong, 1983; Felsenstein, 1985, 2004; Bremer, 1988, 1994; Efron & Tibshirani, 1993; Davis, 1995; Efron et al., 1996; Holmes, 2003; Soltis & Soltis, 2003; Fitzhugh, 2006a, 2012, 2016c; Lemoine et al., 2018). Neither method provides epistemically meaningful values regarding evidential support, contrary to the popularity of their use, or claims that they provide a basis to say phylogenetic hypotheses are ‘robust.’ The bootstrap was originally developed to test statistical hypotheses through a process of random resampling from an original set of random samples taken from a population. The intent is to test the hypothesis that a statistical parameter, inferred from the original samples, has the hypothesised value, without having to perform additional sampling from the population. It is important to note that the bootstrap approach is designed to test statistical, *not* explanatory hypotheses. Phylogenetic hypotheses are not statistical constructs, as clearly shown in (4) and (5). Like explanatory hypotheses, however, hypotheses of statistical estimates are inferred by way of abduction. Such estimates are generalisations that account for observed instances as representative of the population from which the sample was drawn. For example,

- (8)
- Some balls in this bag are red
 - 25% of balls in this random sample are red
-
- *Hypothesis*: 25% of all balls in this bag are red.

Testing the hypothesis would proceed by predicting what should be observed if additional random samples are taken. Such an inference could not be deductive, but rather what Peirce (1932: 2.268) called a ‘statistical deduction,’

- (9)
- *Hypothesis to be tested*: 25% of all balls in this bag are red
 - A random sample of balls will be taken from the bag
-

- *Predicted test evidence*: 25% of balls in the sample will be red.

Notice that the prediction of test evidence in **(9)** is different from what is shown for the explanatory hypothesis in **(2)**. Statistical hypotheses are tested by seeking the same class of data used to originally infer the hypotheses, whereas testing explanatory hypotheses requires test evidence that is different from the effects originally used to infer the hypotheses (Fitzhugh, 2016c, 2016d).

As suggested by Felsenstein (1985, 2004), the bootstrap method applied to phylogenetic hypotheses involves randomly sampling characters from an original character data matrix to create contrived data sets of the same size from which new cladograms are produced. Keep in mind that these cladograms, as ‘bootstrap replicates,’ are *not* explanatory hypotheses; they are merely branching diagrams derived from artificial data sets. Yet, it is claimed that the more often ‘clades’ (actually only branching patterns) are obtained among these ‘replicates’ that match the original groupings indicates higher ‘support’ for those clades. This claim is, however, entirely incorrect. The ‘bootstrap replicates’ have no epistemic relation to the original phylogenetic hypothesis(es) under consideration, such that any measure of ‘support’ would be impossible to ascertain. Along with the statistical, i.e., inductive, mindset that was introduced into phylogenetic inference in the 1970’s, the bootstrap method applied to phylogenetic hypotheses was largely an exercise in filling a perceived methodological void. Had the abductive nature of systematics been recognised early on, and character data not confused with test evidence, the state of the field might have avoided so many scientifically questionable schemes. The only support relevant to cladograms is valid test evidence produced in the process of testing the various classes of hypotheses within a (composite) phylogenetic hypothesis [Fig. 3; cf. **(2)**, **(3)**]. But as noted

earlier, such testing virtually never occurs, making the need for talk of support both gratuitous and positively deceptive.

Just as the bootstrap provides results that cannot be interpreted as support for phylogenetic hypotheses, the Bremer Index produces values that are equally impossible to defend. Like the bootstrap, Bremer Index is claimed to determine support for groups, as opposed to any particular hypotheses implied by cladograms. This support is based on the extent to which groups or ‘clades’ are present in cladograms of ever-increasing ‘length’ beyond the original minimum-length tree. The greater the tree length with a particular group still present is supposed to represent greater ‘support’ for or resiliency by that group. The problem is that trees of greater length can only be interpreted as sets of explanatory hypotheses that have no epistemic relation to the originally inferred hypotheses. The consequence is that values provided by the Bremer Index offer no indication of support. Once again, in the absence of actually testing hypotheses, as indicated in (2)–(3), no evidential test support is possible for hypotheses.

Finally, an oblique yet equally specious approach to garnering support comes from comparisons of cladograms inferred from different sets of characters, sometimes referred to as taxonomic congruence (Fitzhugh, 2014, 2016c). The idea is that if tree topologies are identical, or nearly so, this justifies a sense of greater confidence or belief in those topologies (not the implied explanatory hypotheses). Just as was noted earlier with regard to the error of character mapping, comparisons of cladograms are an inferentially indefensible approach. Cladograms inferred to explain separate sets of characters are distinct compilations of explanatory constructs that have no epistemic relevance to each other. A popular, somewhat related approach is to speak of comparisons of past and present cladogram topologies inferred from updated compilations of observations, remarking on the degree of similarities between groups in those topologies. The

same basic criticism against taxonomic congruence applies here. For example, previous abductive inferences using data set D_1 give composite phylogenetic hypothesis H_1 . With subsequent characters added, D_{1+2} , a new composite phylogenetic hypothesis, H_2 , is inferred. It might seem reasonable to speak of topological similarities and differences between H_1 and H_2 but this is a mistake. Again, H_1 and H_2 are distinct sets of explanatory hypotheses from different sets of abductive premises. Just as there are no meaningful comparisons to be made between the actual explanatory hypotheses implied by the different cladograms (cf. Fig. 2), there are no relevant conclusions to be made regarding topological similarity or difference. In the absence of recognising the nature of abduction or associated intent of those inferences, focusing on phylogenetic tree comparisons offers no scientifically meaningful evaluation.

Sequence data and explanatory hypotheses

The pursuit of sequence data was not given consideration in this study. The reasoning behind this decision will be outlined in this subsection. A more complete account is presented in Fitzhugh (2016b; see also Nogueira et al., 2017: 683–684, Lovell & Fitzhugh, 2020: 270 and Fitzhugh, 2021). Contrary to a claim made by a reviewer of an earlier draft of this paper, what is presented in this section should not be interpreted as the assertion that we ‘do not believe in sequence data.’ Rather, our intent is to point out inherent and significant difficulties that generally preclude causally accounting for differentially shared nucleotides or amino acids via inferences of phylogenetic hypotheses.

The line of argument developed in this more inclusive section – establishing the basis for systematics – follows from the fact that the goal of scientific inquiry is to continually acquire causal understanding of phenomena, in the form of differentially shared characters of organisms, that we encounter as well as describe. The pursuit of that understanding begins with reactions to

those effects, in the form of implicit or explicit why-questions, and subsequently inferring answers to those questions by way of abductive reasoning. As discussed earlier, abduction produces hypotheses that posit possible past causal events/conditions that account for observed effects, as shown in (1), (4), and (5). And, almost all actions in systematics are centered around abduction as opposed to testing hypotheses via induction (Fitzhugh, 2005a, 2006a, 2006b, 2008a, 2009, 2010a, 2012, 2013, 2014, 2015, 2016a, 2016b, 2016c, 2016d).

We pointed out earlier that composite phylogenetic hypotheses provide almost no causal specifics among the four classes of hypotheses implied by cladograms (Fig. 3). This is a consequence of the fact that phylogenetics computer algorithms emulate a vague form of abduction [e.g., (4), (5)]. Consider again what the *Phylogenetic theory* in (4) offers regarding character origin/fixation:

character $x(1)$ originates by [unstated] mechanisms $a, b, c \dots n$, and becomes fixed within the population by [unstated] mechanisms $d, e, f \dots n$ (= ancestral species hypothesis), followed by [unstated] event(s) $g, h, i \dots n$, wherein the population is divided into two or more reproductively isolated populations.

The cause of character fixation in an ancestral population could at a minimum be assumed to be natural selection or genetic drift. In the absence of explicit causal specifics being presented in the *Phylogenetic theory*, it would have to be assumed that either selection or drift are reasonable causal alternatives. This equivalence is plausible for explaining phenotypic characters but cannot be applied to sequence data. For selection to operate, whether purifying or directional, variation among heritable traits within a population must be identified that directly result in differential fitness among individuals. Nucleotides and amino acids lack emergent properties that could *directly* manifest fitness differences (for important nuances, cf. Linquist et al., 2020). Whilst

there are tests that can indicate whether selection has occurred in relation to sequence data, such as the McDonald–Kreitman test (McDonald & Kreitman, 1991; Sawyer, 1994; Hey, 1999; Hurst, 2002; Biswas & Akey, 2006; Zhang & Yu, 2006; Koonin, 2012; Petrov, 2014), these tests cannot discriminate at what organizational level selection might have been directed. It is at higher organisational levels that phenotypic characters can result in fitness variation. If selection is a causal factor, it comes into play due to phenotypic effects at these higher levels. Alternatively, since there is no *direct* selection for sequence data, phenotypic characters have the potential to lead to differential fitness and intergenerational selection that can influence the occurrences of those sequence data that produce the selected characters. The result is a form of *indirect* selection, termed downward causation by Campbell (1974: 180),

“Where natural selection operates through life and death at a higher level of organisation, the laws of the higher-level selective system determine in part the distribution of lower-level events and substances.”

Since the 1970’s the importance of downward causation has become increasingly recognised (e.g., Vrba & Eldredge, 1984; Salthe, 1985; Lloyd, 1988; Auletta et al., 2008; Ellis, 2008, 2012, 2013; Ellis et al., 2011; Jaeger & Calkins, 2011; Laland et al., 2011; Martínez & Moya, 2011; Davies, 2012; Okasha, 2012; Walker et al., 2012; Griffiths & Stotz, 2013; Martínez & Esposito, 2014; Walker, 2014; Mundy, 2016; Callier, 2018; Pouyet et al., 2018; Salas, 2019; Yu et al., 2020), but its relevance to how sequence data are explained via phylogenetic hypotheses has only recently been considered (Fitzhugh, 2016b).

If there can be no direct selection for sequence data, the issue then becomes one of deciding to explain individual nucleotides or amino acids either directly via drift or indirectly through downward causation. If the latter, it is first necessary to associate those sequence data

with the higher-level phenotypic characters to which selection is the hypothesised causal condition. Those associated sequence data would then be excluded from phylogenetic inferences since they would already be explained in conjunction with higher-level characters (Fitzhugh, 2016b). On the other hand, it might be argued that one could avoid the issue altogether by assuming all sequence data should be explained by drift. This would be unrealistic as it necessitates that selection at higher organisational levels never occurs. In the absence of being able to discriminate between drift versus selection by downward causation, the only sensible option is to forgo altogether attempts to causally account for shared nucleotides or amino acids by way of phylogenetic inference. Otherwise, to explain shared nucleotides or amino acids with phylogenetic hypotheses inferred under an entirely agnostic perspective [i.e., the *Phylogenetic theory* in (4)] would be epistemically and scientifically unwarranted.

The inherent limitations for inferring explanatory hypotheses for sequence data have associated relevant consequences for mapping morphological characters on cladograms only inferred for sequence data. As discussed earlier (*Phylogenetic inference = abduction*; see also (Fitzhugh, 2014), mapping is an epistemically unfounded approach to explaining characters. This problem is compounded by the fact that not discriminating between genetic drift and natural selection as possible causal factors when explaining sequence data yet again precludes rational acceptance of phylogenetic hypotheses accounting for mapped characters.

Finally, it might appear at first sight that the exclusion of sequence data on either the basis of downward causation or inability to distinguish drift from selection is at odds with the requirement of total evidence (RTE; cf. Fitzhugh, 2006b, for a discussion of the RTE in relation to systematics). Such is not the case. The RTE is a recognised maxim for all non-deductive reasoning (the requirement is automatically satisfied in deduction), wherein rational acceptance

of a conclusion is based on considering all relevant evidence that can affect support (either negative or positive) for that conclusion (Carnap, 1950; Barker, 1957; Hempel, 1962, 1965, 1966, 2001; Salmon, 1967, 1984a, 1984b, 1989, 1998; Sober, 1975; Fetzer, 1993; Fetzer & Almeder, 1993): “All that the requirement of total evidence says is that one’s confidence in a hypothesis must be proportional to the support that that hypothesis receives from one’s evidence...” (Neta, 2008: 91). Nearly all of the philosophical literature on the RTE has focused on test evidence in relation to induction [cf. (3)], which is understandable given that, for instance, to exclude known test evidence that could enhance or compromise acceptance of a theory or hypothesis would be less than rational.

Whilst abductive reasoning has almost never figured into discussions of the RTE, the non-deductive nature of abduction necessitates its attention (Fitzhugh, 2006b). Consider the following example. For members of three species, *a-us*, *b-us*, and *c-us*, there are characters distributed among subjects 1–3 (cf. Fitzhugh, 2006c):

Subjects:	1	2	3
Outgroup	0	0	0
<i>a-us</i>	0	1	1
<i>b-us</i>	1	0	1
<i>c-us</i>	1	1	0.

If the characters among members of *a-us*, *b-us*, and *c-us* are explained separately, there would be three sets of explanatory hypotheses (outgroup is excluded), represented here in parenthetical form:

Subject 1: (*a-us* (*b-us*, *c-us*))
 Subject 2: (*b-us* (*a-us*, *c-us*))

Subject 3: (c-us (a-us, b-us)).

The three sets of hypotheses are contradictory, and while each is supported by abductive evidence [cf. (1)], choosing among them based on the information provided would be problematic. This is the sort of situation the RTE is intended to address, as well as those instances that involve other degrees of data partitioning in relation to inferring phylogenetic hypotheses and character mapping. The nature of the abductive inferences involves explaining features of semaphoronts, such that the observed characters are part of the integrated network that makes up those individuals. The various phylogenetic theories applied to those characters [cf. (1)] assume that past causes operated on individuals, not distinct characters. Thus, explaining the characters under each subject would be relevant to explaining the other characters. The plausibility for any of the phylogenetic hypotheses would be based on an abductive inference where all the characters are treated as part of the premises.

Related to the topic highlighted in this subsection, it would not be a violation of the RTE to exclude from phylogenetic inferences those sequence data for which no empirical basis is available for discriminating drift from selection-via-downward-causation for sequence data. The difficulty faced is not a matter that falls under the purview of the RTE, but rather an inability to engage in abductive reasoning for a particular class of characters.

Sequence data are not more objective than other classes of observations

Consider the following statements,

“Molecular data are more objective and subject to considerably more rigor than morphological data. DNA sequence contains four easily identified and mutually exclusive character states→ Morphological and embryological character definitions and scoring of character states are far more subjective, and most

characters have been repeatedly used without critical evaluation, calling into question the utility of morphological cladistic studies that span Metazoa.” (Halanych, 2004: 230).

“As we move forward with the deep-animal tree [sic], we should not employ morphology and developmental data to reconstruct the tree. There is too much historical baggage and subjectivity with these data. We should reconstruct the tree with molecular data and then use that tree to independently interpret the morphology and development in light of that tree.” (Halanych, 2016: 325).

“We present the view that rigorous and critical anatomical studies of fewer morphological characters, in the context of molecular phylogenies, is a more fruitful approach to integrating the strengths of morphological data with those of sequence data. We argue that a main constraint of morphology-based phylogenetic inference concerns the limited number of unambiguous characters available for analysis in a transformational framework” (Scotland et al., 2003: 539).

Taken at face value, each statement appears damning of all or most observations of features of organisms other than sequence data in relation to inferring phylogenetic hypotheses. We have seen throughout this section, however, that the disconnect between principles of scientific inquiry and recent views on phylogenetic inference have led to perspectives that cannot be defended. This outcome extends to the notion that sequence data are somehow ‘more objective,’ ‘more fruitful,’ or more reliable than other observations. Asserting that one class of observations is more effective than other classes presumes that the objective of phylogenetic inference is to obtain ‘phylogenies’ or ‘trees,’ which is at odds with pursuing explanations of all relevant

observations. If the objective of systematics is causal explanation, thus consistent with all of scientific inquiry, then the pursuit of ‘phylogenies’ or ‘trees’ is not only contrary to that objective but has led to erroneous inclinations in addition to the largely contrived objective/subjective dichotomy. These erroneous pursuits include, but are not limited to (1) inferring phylogenetic hypotheses for separate classes of characters, especially sequence data versus morphology, and drawing comparisons between tree ‘topologies’ (cf. *Phylogenetic hypotheses and the myth of evidential support*), and (2) inferring phylogenetic hypotheses for sequence data, then ‘mapping’ morphological characters on the ‘tree’ and making evolutionary claims regarding the latter characters (cf. *Phylogenetic inference = abduction*). To address the claim that sequence data are more objective or less ambiguous requires that we first consider what is meant by scientific objectivity. Then there are several additional issues in the above quotes that need to be addressed.

As might be expected, objectivity has multiple meanings. Useful overviews can be found in Daston & Galison (2007), Gaukroger (2012), Reiss & Sprenger (2017), and Wilson (2017). Common interpretations of objectivity range from the view that judgements should be (a) free of prejudice or bias, (b) free of assumptions, and/or (c) an accurate representation of, or ‘faithfulness’ to facts (Gaukroger, 2012). Each meaning can be applied to everyday life as well as within fields of science. Regarding freedom from prejudice or bias, this is to speak of limiting levels of distortion beyond what is perceived. Freedom from assumptions would be impossible to accomplish given the theory-laden nature of observation statements and other types of propositions: “Every instance of scientific inquiry, every study, rests on a vast submerged set of political, moral, and ultimately metaphysical assumptions” (Wilson, 2017). Scientific objectivity has often been interpreted along the lines of ‘faithfulness to facts’ (Daston & Galison, 2007;

Reiss & Sprenger, 2017), where we attempt to minimise arbitrary judgements; we want to answer questions on the basis of the most appropriate evidence we have available. Objectivity occurs in degrees – there is no absolute objectivity, which is why objectivity cannot be equated with truth, which is absolute. Ultimately, “While we can strive for objectivity, inquiry is inherently subjective” (Spencer, 2020).

Reiss and Sprenger (2017) distinguish ‘product’ and ‘process’ objectivity. Product objectivity regards the products of scientific inferences, such as theories, laws, and observation statements, as accurate representations of the external world. Process objectivity refers to processes and methods that are not dependent on social or ethical values, or the individual biases of the scientist. It is product objectivity that is of concern when it comes to assessing the quotes given above, claiming sequence data are in some way more objective or less ambiguous than other classes of observations.

There is nothing in the conception of product objectivity that would support the claim that observation statements of morphological features are less objective than sequence data, much less that such a contrived distinction warrants the exclusion of an entire class of observations from the goal of systematics. Indeed, that biologists hold different interpretations of what they perceive of the properties of organisms is not tantamount to proffering less objective observation statements. Those statements are themselves products of abductive reasoning (Peirce, 1935; Hoffman, 1995; Burton, 2000; Magnani, 2009; Anjum & Mumford, 2018), constrained by one’s background knowledge, and like all hypotheses, are open to empirical evaluation. If anything, it can be argued that sequence data are less than objective relative to other classes of characters given the fact that attempted phylogenetic inferences explaining differentially shared nucleotides or amino acids typically operate under the assumption that rates

of substitution must be imposed within abductive inferences (cf. *Phylogenetic inference* = *abduction*). That assumption carries with it the implication that observation statements of shared characters must be treated as potentially false, hence the need to interject substitution rates within inferences, as indicated by the all too common emphasis on branch lengths. Considering the rule for valid reasoning – that we should assume the truth of premises, discussed earlier – the treatment of sequence data with ‘likelihood’ and ‘Bayesian’ methods pales against any criterion of objectivity.

Rather than focusing on the false argument from objectivity as a basis to eliminate observations of morphological characters in need of explanation, systematists should embrace as much as possible all relevant observations per the goal of scientific inquiry discussed earlier and the requirement of total evidence (Fitzhugh, 2006b). Two notable exceptions to considering explaining differentially shared characters by way of phylogenetic hypotheses come from conditions discussed earlier: (1) why-questions addressing some characters might not be amenable to being answered by phylogenetic hypotheses, but instead one of the other classes of explanatory hypotheses applied in systematics (Fig. 1); (2) epistemic limitations to explaining sequence data because of an inability to discern explanations due to genetic drift versus selection by way of downward causation (cf. *Sequence data and explanatory hypotheses*). Pursuing causality is the intent – not producing ‘trees’ or ‘phylogenies,’ which are typically devoid of causal considerations in relation to sequence data.

Outgroup and ingroup taxa

Outgroup and ingroup taxa considered in this study include the following:

outgroups – Chaetopteridae: *Phyllochaetopterus limicolus* (Hartman, 1960)

Spionidae: *Spio filicornis* (Müller, 1776)

933 *Prionospio ehlersi* Fauvel, 1928

934 *P. lighti* Maciolek, 1985

935 *Laonice cirrata* (Sars, 1851)

936 ingroup – *Magelona* (72 species)

937 *Octomagelona* (2 species)

938 The choice of outgroups ideally would follow from more inclusive phylogenetic studies from
 939 which the immediate sister groups to Magelonidae have been ascertained. In this regard, the
 940 merits of recent phylogenetic hypotheses are questionable. In her review of the Magelonidae,
 941 Mortimer (2019) summarised past studies of phylogenetic relationships of magelonids to other
 942 groups within Polychaeta. As would be expected for a group as moderately disparate as
 943 magelonids, earlier conclusions were that they exhibit close relations with other spioniforms.
 944 Recent phylogenetic studies based exclusively on morphological (Capa et al., 2012; Chen et al.,
 945 2020) or sequence characters (Weigert et al., 2014; Helm et al., 2018) have, however, allied
 946 Magelonidae with Oweniidae, while Weigert et al.'s (2016) use of mitochondrial sequences had
 947 Magelonidae and Chaetopteridae as sister taxa. Each approach has aspects that call into question
 948 the respective conclusions. For instance, Capa et al. (2012; see also Capa et al., 2019) cautiously
 949 suggested a Magelonidae-Oweniidae clade based on the absence of nuchal organs, fusion of pro-
 950 and peristomium, presence of a ventral buccal organ, and monociliated cells. The absence of
 951 nuchal organs is not exclusive to magelonids and oweniids (Purschke, 1997, 2005). Mortimer
 952 and Mackie (2014) and Mortimer (2019) have shown that pro- and peristomium are not entirely
 953 fused, and the magelonid 'burrowing organ' is not homologous to the oweniid buccal organ
 954 [Mortimer, 2019; see also **Character descriptions**, *Burrowing organ* (subject 14), below].
 955 Monociliated cells have only been recorded from among members of *Magelona mirabilis*

(Bartolomaeus, 1995, N.B. fig. 1A in this publication is not *M. mirabilis*; Patrick et al., 2019). As part of their description of a fossil magelonid, Chen et al. (2020: fig. 8a; see also Parry et al., 2016) presented phylogenetic hypotheses also suggesting Oweniidae and Magelonidae are exclusive sister taxa, but their inferences are based on an overly broad taxon coverage and the compilation of characters is inadequate to address phylogenetic relationships of magelonids to other polychaetes. As discussed in the previous section (**Methodological considerations – Sequence data and explanatory hypotheses**), the inherent problems associated with attempting to explain shared nucleotides or amino acids undermine the plausibility of phylogenetic hypotheses inferred from those premises. With the emphasis on an extensive treatment of magelonid morphological characters in this study, the greatest number of characters is shared with members of Spionidae, making them an appropriate outgroup. Members of several genera and species were included to accommodate variation within Spionidae. An additional spioniform representative from the Chaetopteridae, *Phyllochaetopterus limicolus*, was also included.

Ideally, all currently recognised magelonid taxa would be included in inferences of the phylogenetic hypotheses. This, however, relies on several factors. Initially, taxa were chosen based on access to specimens, preferably type material, some of which had been observed previously during descriptions and re-descriptions by the present authors. Specimens were borrowed from the following institutions:

BMNH – Natural History Museum, London

MB – Museu Nacional de História Natural, Lisboa

MNCN – Museo Nacional de Ciencias Naturales, Madrid

MNHM – Muséum National d'Histoire Naturelle, Paris

NHMG – Gothenburg Museum of Natural History

979 **NHMLAC** – Natural History Museum of Los Angeles County
 980 **NMW** – Amgueddfa Cymru – National Museum Wales
 981 **PMBC** – Phuket Marine Biological Center
 982 **SMF** – Senckenberg Research Institute and Natural History Museum, Frankfurt
 983 **USNM** – Smithsonian National Museum of Natural History
 984 **ZMBN** – University Museum of Bergen
 985 **ZMHB** – Museum für Naturkunde der Humboldt-Universität zu Berlin
 986 **ZMUC** – Zoological Museum, University of Copenhagen
 987 **ZUEC** – Museu de Zoologia, Universidade de Campinas

988 Where specimens were not available for observation, the remaining taxa were chosen
 989 based on the level of information and detail in the original descriptions. It is only comparatively
 990 recently that the importance of illustrating and fully describing all thoracic chaetigers has been
 991 realized for the treatment of taxa for example, and further characters defined. Thus, many of the
 992 earliest descriptions lack much of the detail required for the current analysis. All ingroup taxa
 993 included in the analyses are listed in Table 2, including which specimens were observed and the
 994 institution from which they were borrowed. Table 2 also indicates whether observations were
 995 based solely on published material.

996 **Character descriptions**

997 A total of 176 characters distributed among 79 subjects (Table 3; Table S1) were used to infer
 998 phylogenetic hypotheses. Descriptions of characters are presented here.

999 *Characters of the anterior end: prostomium (subjects 1–9)*

The form of the magelonid prostomium (**subject 1**) is considered a synapomorphy of the group, the name ‘shovel head worms’ coined due to its peculiar shape. The dorso-ventrally flattened, shovel-shaped prostomium [character 1(1), Figs. 4 and 5] is a feature not observed among members of the outgroups or any other polychaete group. Nuchal organs (**subject 2**) have not been recorded among adult members of any magelonid species [character 2(0)] but are present among all members of the outgroups [character 2(1)].

The distal margin of the magelonid prostomium (**subject 3**) can vary from smooth [character 3(0), Figs. 4A and 4F], crenulate [character 3(1), Figs. 4B and 4C] or medially indented [character 3(2), Fig. 4D]. Uebelacker & Jones (1984) stated that the presence of a crenulated prostomial margin is an excellent character for separation of species, although the degree of crenulation present shows great intraspecific variability. The crenulations may differ in number and form, for example, the distinct, triangular crenulations of *Magelona* sp. L of Uebelacker & Jones (1984) or *M. crenulifrons* (Fig. 4B), in comparison to the numerous minute crenulations observed for *M. longicornis* or *M. wilsoni* (Fig. 4C). The crenulations among members of some species are difficult to discern and this is perhaps the reason they have been previously overlooked in some descriptions and erroneously reported as absent in others. The prostomial distal margin (**subject 4**) may additionally vary in shape, from triangular [character 4(0), Fig. 4E], to rounded [character 4(1), Figs. 4A and 4F] or straight [character 4(2), Figs. 4G, 4H, 5C and 5E]. As subjects 3 and 4 relate directly to the shovel-shaped prostomium, characteristic of magelonids, it does not apply to any members of the outgroups.

Horns are the distal projections of the prostomium (**subject 5**) that occur among members of some species of magelonids [character 5(1)], such as *Magelona pacifica* (Fig. 5A), or *M. montera* (Fig. 5B). Uebelacker & Jones (1984) drew attention to this character for the initial

separation of magelonid species. When present, horns can vary in size/shape (**subject 6**), from rudimentary [character 6(0), Figs. 4G, 4H, 5C and 5E], such as those observed among members of *M. variolamellata* and *M. allenii* (often described as a squared anterior margin), or conspicuous structures as seen among members of species such as *M. posterelongata* [character 6(1), Fig. 5D]. In the latter case, these are often very distinct from the distal prostomial margin. Prostomial horns do not occur in the outgroups, thus subjects 5 and 6 are not applicable.

Several authors have identified dimensions of the prostomium (**subject 7**) as an important feature in the separation of magelonid species (Hartman, 1944; Wilson, 1958, 1959; Jones, 1963, 1971, 1978; Uebelacker & Jones, 1984; Bolívar & Lana, 1986; Blake, 1996; Mortimer, 2019). According to Uebelacker & Jones (1984), the general shape and relative dimensions are fairly constant within each species, something which the current authors are in agreement. Consequently, the ratio of width to length was used to recognise three prostomial characters: prostomium longer than wide [character 7(0), Figs. 4A, 4F, 5B and 5D], as wide as long [character 7(1), Figs. 4B, 4D and 4G], and wider than long [character 7(2), Figs. 4C, 4H and 5E]. Members of three species within the outgroups (*Spio filicornis*, *Prionospio lighti* and *P. ehlersi*) are considered to possess prostomia which are longer than wide [character 7(0)], whilst members of *Phyllochaetopterus limicolus* and *Laonice cirrata* have prostomia which are wider than long [character 7(2)].

Magelonid prostomia carry paired ‘muscular’, dorsal longitudinal ridges (**subject 8**), which extend from the base of the prostomium towards the distal tip [character 8(1)]. Jones (1968) stated that their “inner surfaces, particularly dorsally and ventrally, are provided with longitudinal muscles,” describing them as hollow, cylindroid structures, which he presumed were fluid filled. The number of prostomial ridges (**subject 9**) varies from one pair [character 9(0),

Fig. 4H], such as members of *Magelona equilamellae* or two pairs [character 9(1), Figs. 4A–C, 4E and 5B], as observed among members of *M. pacifica* (Fig. 5A). The ridges may vary in size and distinctiveness, the outer pair among members of some species being more difficult to discern, e.g., *M. allenii* (Fig. 4G), to the distinct and transversely ridged structures seen in, for example, members of *M. wilsoni* (Fig. 4C). Prostomial ridges are not present among members of the outgroups [character 8(0)] and thus subject 9 is not applicable.

Characters of the palps (subjects 10–13)

Magelonids possess two long palps, considered to be peristomial, arising from the larval prototroch (Wilson, 1982). There has been much discussion about palp origin (**subject 10**), since they are ventrally inserted [character 10(1), Figs. 4F and 5G], rather than dorsally [character 10(0)], the latter of which is seen among members of all other polychaete groups with palps, including the outgroups. In addition, the palps do not have the characteristic feeding groove seen among members of other taxa such as spionids and *Myriowenia*. However, an inconspicuous ventral line devoid of papillae is generally present. According to Orrhage (1966), palps among members of the Spionidae and Magelonidae are homologous because they are innervated by nerves from the same part of the brain. More recently Beckers et al. (2019) stated that two nerves innervate the palps of members of the Magelonidae, Spionidae, Chaetopteridae and Oweniidae, indicating the purported plesiomorphic condition in Annelida.

Magelonid palp surfaces (**subject 11**) are uniquely papillated, with ampulliform to digitiform papillae arranged in rows along the longitudinal axis [character 11(1), Figs. 4H, 5B and 5F–H]. Members of the outgroups, along with all other polychaetes possess non-papillated palps [character 11(0)]. Although the number of rows of papillae varies among members of magelonid species, it is relatively stable among members of any particular species. As papillae

are involved in obtaining food (Jones, 1968; Fauchald & Jumars, 1979; Mortimer & Mackie, 2014), variations in number of papillae may reflect different feeding strategies. In general, the number of rows of papillae are fewer distally in comparison to proximally (Jones, 1963, 1971, 1978). The number of rows of papillae in the proximal region of the palp (**subject 12**) were coded as three characters: two rows of papillae [character 12(0)], 4–8 rows [character 12(1)], and 10–14 rows [character 12(2)]. The number of rows of papillae at the distal end (**subject 13**) were coded for 2 [character 13(0)], 4 [character 13(1)], or more than 4 rows [character 13(2)]. Where a range was observed among members of a species, the maximum number of rows was the value used. As with palps in other polychaete groups, those of magelonids are easily lost upon collection. Many descriptions do not include information about them, whilst others do not detail differences in number of papillae along the palp length. Many authors have undervalued the importance of palp characteristics in the descriptions of members of species. These two factors present the largest problem in coding these subjects. It should be noted that the number of rows of papillae may vary in animals with regenerating palps, particularly at the distal tips, thus affecting the ranges observed. Subjects 12 and 13 are not applicable to members of the outgroups since papillated palps are unique to magelonids.

Burrowing organ (subject 14)

Magelonids have long been described to possess a ventral eversible proboscis. A heart-shaped sac (when fully everted; oval when only partially everted), carrying longitudinal ridges (Mortimer, 2019: fig. 4.2.4). However, recent studies show that this structure plays no part in feeding (and is not connected to the buccal region) but functions in burrowing only (Mortimer, 2019). The term ‘burrowing organ’ should be applied to this structure. The burrowing organ is

present in members of all known magelonid species [character 14(1)] but not present in any members of the outgroups [character 14(0)].

Characters of body regionation (subjects 15–16)

The body of magelonids (**subject 15**) is divided into an anterior region comprising the head and ‘thorax’ (the latter of which is characterised by the presence of capillary chaetae), and an ‘abdomen’ comprising many segments with hooded hooks [character 15(1)] (Figs. 6A and 11A). As an aside, the terms thorax and abdomen are ill-defined when applied to polychaetes overall, other than to vaguely indicate anterior-posterior distinctions along the body. Unfortunately, the characters associated with these regions are not uniformly applied across all of Polychaeta.

The body of members of the outgroups do not have this regionation [character 15(0)]. Behind the head region (**subject 16**), the thorax carries an achaetous first segment, followed by either eight [character 16(0), *Octomagelona*; Fig. 6] or nine [character 16(1), *Magelona*; Figs. 11, 15 and 16] chaetigers (Fig. 11A). Subject 16 does not apply to members of the outgroups.

Characters of thoracic parapodia (subjects 17–30)

Unique terminology introduced and later modified by Jones (1963, 1971, 1978) has been applied to the family, particularly related to parapodial structures (Fig. 7). Brasil (2003: fig. 4) provided a comprehensive review of this terminology and problems related to inconsistencies in applying terms. Consequently, several authors have suggested its abandonment (Rouse, 2001; Mortimer & Mackie, 2003). Magelonid chaetigers possess biramous parapodia with lamellae in both the noto- and the neuropodia. Lamellae are generally postchaetal, flattened structures that range from filiform to foliaceous in shape. They encircle the chaetal bundle, attaching to a low, triangular or rounded prechaetal lamella, almost cuff-like (Fig. 8A). However, among members of some

1114 species, the lamellae are somewhat subchaetal in position, underneath the chaetal bundle and
 1115 appearing somewhat U-shaped when viewed laterally (Figs. 8B, 8D). All members of all species
 1116 in the ingroup and outgroups have postchaetal noto- and neuropodial lamellae in
 1117 thoracic/anterior chaetigers (**subjects 17** and **19**, respectively) except for *Phyllochaetopterus*
 1118 *limicolus*, in which they are absent [characters 17(0) and 19(0)].

1119



1120

1121 The development of notopodial (**subject 18**) and neuropodial lamellae (**subject 20**) along
 1122 the anterior chaetigers may be similar in all chaetigers [characters 18(0) and 20(0), respectively],
 1123 such as among members of *Magelona papillicornis*, and *M. minuta* (Fig. 7A), or vary along the
 1124 thorax [characters 18(1) and 20(1), respectively]. Subjects 18 and 20 are considered inapplicable
 1125 to members of all species in the outgroups. The variation in degree of development can be easily
 1126 observed among members of *M. obockensis* (Fig. 7C), and *M. conversa* (Fig. 7D), in both the
 1127 noto- and neuropodia, whilst among members of *Octomagelona bizkaiensis* it is evident only in
 1128 the neuropodia (Fig. 6E). The lamellae of chaetigers eight and nine often vary in comparison to
 1129 those of the first seven chaetigers, e.g., *M. crenulifrons*, and *M. johnstoni* (Fig. 9I). Thus,
 1130 characters of the lamellae of the first seven chaetigers were coded separately (**subjects 21–30**) to
 1131 those for chaetiger eight (**subjects 34–40**), and chaetiger nine (**subjects 41–49**).

1132 The notopodial lamellae in chaetigers 1–7 (**subject 21**) may be postchaetal [character
 1133 21(0); noted also for members of the outgroups, except *Phyllochaetopterus limicolus*, for which
 1134 the subject is not applicable] or subchaetal [character 21(1)], whilst neuropodial lamellae
 1135 (**subject 26**) may be postchaetal [character 26(0); observed for members of the outgroups, except

P. limicolus, for which the subject is not applicable], subchaetal [character 26(1)] or prechaetal [character 26(2)]. Prechaetal neuropodial lamellae, such as are observed among members of *Magelona conversa* and *M. mirabilis* (Figs. 7D and 8C), are less frequent than both post- (Fig. 7A) or subchaetal (Fig. 9E). The position of the thoracic neuropodial lamellae (**subject 27**) may be consistent throughout the thorax [character 27(0)], or may vary [character 27(1)], e.g., being prechaetal on chaetiger 1, occurring in a ventral position in the mid-thorax, and distinctly postchaetal in the posterior thorax (Figs. 8F and 8G). This subject was not considered applicable to members of the outgroups.

Thoracic notopodial **lamellar** shape (**subject 22**) may be filiform, having a long, tapering structure [character 22(0), Figs. 9A, 9B and 9G] or more foliaceous [character 22(1), Figs. 9C–F]. The margins of the latter (**subject 23**) may be smooth [character 23(0)], crenulate [character 23(1), e.g., *M. montera*, Figs. 9D and 9E] or ~~bilobe~~ [character 23(2), e.g., *M. obockensis*, Fig. 9F]. All members of the outgroups are considered to have foliaceous [character 22(1)], smooth-edged [character 23(0)] anterior notopodia, except for members of *Phyllochaetopterus limicolus* for which subjects 22 and 23 are not applicable. Above the notopodial lamellae and in a slightly prechaetal position, superior dorsal lobes (**subjects 24, 25, 37, 45**) (dorsal medial lobes of Jones) may be present (Figs. 7C, 7D, 8D, 8E and 9C–F) or absent [characters 24(0), 37(0), 45(0)]. These structures are generally smaller than the notopodial lamellae and somewhat cirriform to digitiform in shape. However, among members of some species they may be larger and distinctly foliaceous. When present, superior dorsal lobes may be present on all thoracic chaetigers [characters 25(1), 37(1), 45(1)], on chaetigers 1–8 only [characters 25(1), 37(1), 45(0)], or only occurring on a small number of thoracic chaetigers [often starting in the mid-thoracic region, e.g., *M. johnstoni* or *M. conversa*, character 25(0), Fig. 7D]. Superior dorsal lobes are absent

among all members of the outgroups on chaetigers 1–9 (thus subject 25 is not applicable), and subjects 24, 25, 37 and 45 are considered inapplicable for *Phyllochaetopterus limicolus*.

The shape of thoracic neuropodial lamellae (**subject 28**) may be filiform [character 28(0), Figs. 7C, 8F and 8G] or foliaceous [character 28(1), Fig. 9C, also seen in members of the outgroups, excluding *Phyllochaetopterus limicolus*, for which the subject is not applicable]. The distal ends (**subject 29**) of which may be pointed [character 29(0)], as among members of *M. equilamellae* (Figs. 8F and 8G), or distally expanded and scoop-shaped [character 29(1)] as observed among members of *M. cincta* (Figs 9H). The latter subject is not applicable to members of the outgroups. The lengths of noto- and neuropodial lamellae (**subject 30**) may be equivalent [character 30(0), Fig. 9G], the notopodial lamellae may be longer [character 30(1), Figs. 9D–F], or the neuropodial lamellae may be longer [character 30(2), Fig. 7D], e.g., chaetiger 8 of *M. conversa*. The thoracic notopodial lamellae among members of the outgroups, *Prionospio lighti* and *P. ehlersi* are longer than neuropodial lamellae [character 30(1)], whilst for members of *Spio filicornis* and *Laonice cirrata* they are equivalent [character 30(0)]. Subject 30 is not applicable to *P. limicolus*.

Characters of the chaetae from chaetigers 1–8 (subjects 31–33)

Capillary chaetae (**subject 31**) occur only in the thorax of magelonids (Figs. 6, 11, 15 and 16), but many authors have undervalued the differences between them, describing them simply as limbate capillary chaetae (Hartman, 1944, 1961, 1965; Nateewathana & Hylleberg, 1991). Jones (1977) was one of the few authors who referred to the presence of unilimbate [character 31(0), Fig. 10C] and bilimbate capillary chaetae [character 31(1)] in descriptions of members of species. For some individuals, both unilimbate and bilimbate chaetae have been recorded, however, the predominate form in each case was coded in the current analysis. Bolívar & Lana

(1986) and Blake (1996) additionally noted some members of species possess bilimbate chaetae (**subject 32**) with an irregular blade [character 32(1)], such as *Magelona variolamellata* (Fig. 10E), and Brasil (2003) additionally noted it to occur among members of species such as *M. riojai* (Fig. 10D), *M. pacifica* and *M. pitelkai*. Members of the outgroups possess unilimbate capillary chaetae [character 31(0)], whilst subject 32 is not applicable. Additional to the limbations of capillary chaetae, the relative lengths of noto- and neurochaetae may vary (**subject 33**). Either being equivalent in length [character 33(0)] e.g., *M. nonatoi*, longer in the notopodia [character 33(1)], e.g., *M. sacculata*, or longer in the neuropodia [character 33(2)], e.g., *M. variolamellata*. For members of *Phyllochaetopterus limicolus*, *Spio filicornis* and *Prionospio lighti*, chaetae are longer in the notopodia [character 33(1)], whilst for members of *Laonice cirrata* and *P. ehlersi* they are equivalent in length [character 33(0)]. In some members of magelonid species chaetae of all thoracic chaetigers are similar, however, in others the chaetae of the ninth chaetiger are modified. For this reason, the chaetae were coded collectively for the first eight chaetigers, and separately for chaetiger nine (see below).

Characters of parapodia of chaetiger 8 (subjects 34–40)

As described above, development of lamellae in the magelonid thorax may be similar on all chaetigers or vary between them. For members of species in which they vary, the parapodia of chaetigers 8 and 9 in particular may show larger differences in comparison to preceding chaetigers. However, the same characters that were coded for the first seven chaetigers can be applied to the posterior-most thoracic chaetigers. The notopodial lamellae (**subjects 34–36**) of chaetiger 8 may be postchaetal [character 34(0)] or subchaetal [character 34(1)] in position relative to the chaetal bundle, filiform [character 35(0)] or foliaceous [character 35(1)] in shape, with smooth [character 36(0)], crenulate [character 36(1)] or bilobed [character 36(2)] margins.

Subjects 34–36 are considered not applicable to members of *Phyllochaetopterus limicolus*. However, remaining members of the outgroups possess postchaetal [character 34(0)], foliaceous [character 35(1)] parapodia with smooth margins [character 36(0)]. Superior dorsal lobes (**subject 37**) may be absent [character 37(0)] or present [character 37(1)]. The neuropodial lamellae (**subjects 38–40**) of chaetiger 8 may be postchaetal [character 38(0)], subchaetal [character 38(1)] or prechaetal [character 38(2)] in position relative to the chaetal bundle, and may be filiform [character 40(0)] or foliaceous [character 40(1)] in shape. However, an additional postchaetal expansion (**subject 39**), often triangular in shape, as is seen among members of *Magelona montera* (Fig. 9D) is sometimes observed in the neuropodia of chaetiger 8 (and 9, see below) [character 39(1)]. Subjects 38–40 are considered not applicable to members of *Phyllochaetopterus limicolus*. Members of *Spio filicornis*, *Prionospio lighti*, *P. ehlersi* and *Laonice cirrata* have postchaetal [character 38(0)], foliaceous [character 40(1)] neuropodia in chaetiger 8, without an additional postchaetal expansion [character 39(0)].

Characters of the parapodia of chaetiger 9 (subjects 41–49)

For chaetiger 9, observed characters were coded in the same way as for chaetiger 8. The notopodial lamellae of chaetiger 9 may be postchaetal [character 41(0)], or subchaetal [character 41(1)] in position in comparison to the notochaetae, and may be filiform [character 43(0)] or foliaceous [character 43(1)] in shape, with smooth [character 44(0)], crenulate [character 44(1)] or bilobe [character 44(2)] margins. Members of the outgroups possess postchaetal [character 41(0)] notopodia chaetiger 9 which are foliaceous [character 43(1)], with smooth margins [character 44(0)]. However, subjects 41, 43 and 44 are not applicable to *Phyllochaetopterus limicolus*. Superior dorsal lobes may be present [character 45(1)] or absent [character 45(0)].

The neuropodial lamellae of chaetiger 9 may be postchaetal [character 46(0)], subchaetal [character 46(1)] or prechaetal [character 46(2)] in position in comparison to the chaetal bundle, and filiform [character 48(0)] or foliaceous [character 48(1)] in shape. As with chaetiger 8, an additional postchaetal expansion may be absent [character 49(0)] or present [character 49(1)]. Members of the outgroups possess foliaceous [character 48(1)], postchaetal [character 46(0)] lamellae without an additional postchaetal expansion [character 49(0)], except for *P. limicolus* for which these subjects are not applicable.

The parapodia in chaetiger 9 among members of some magelonid species may be markedly different in comparison to preceding chaetigers. For example, the lamellae in chaetiger 9 among members of *Magelona johnstoni* (Fig. 9I) are much broader and lower [characters 42(0) and 47(0)] in comparison to the elongate lamellae of chaetigers 1–8 in both the noto- and neuropodia, and the elongate lamellae of chaetiger 9 amongst members of other species, e.g., *M. equilamellae* [characters 42(1) and 47(1), Fig. 7B]. For this reason, additional subjects were added to accommodate the height of lamellae in both rami of chaetiger 9 (**subjects 42 and 47**). As members of *Octomagelona* species only possess eight thoracic chaetigers these subjects were coded as inapplicable for these taxa and additionally for members of *Phyllochaetopterus limicolus*. Members of *Spio filicornis* possess broad and low parapodia in chaetiger 9 [characters 42(0) and 47(0)], whilst for members of *Prionospio lighti*, *P. ehlersi* and *Laonice cirrata* they are more elongate [characters 42(1) and 47(1)].

Characters of the chaetae of chaetiger 9 (subjects 50–51)

The chaetae of chaetiger 9 in magelonids may be similar to or vary from those of chaetigers 1–8 (**subjects 50 and 51**). They may be the same length [character 50(0), as in the outgroups], shorter than [character 50 (1)], or longer than those of chaetigers 1–8 [character 50(2)]. The

distal ends of chaetiger 9 chaetae may be gently tapered as is seen in the preceding chaetigers [character 51(0), Figs. 8F and 8G] (observed also for members of the outgroups), or have distinctly mammiform, expanded, mucronate tips [character 51(1), Figs. 7C, 7D, 9I and 10F–H] such as among members of *Magelona johnstoni* and *M. mirabilis*. In members of other species, the distal ends are pennoned [character 51(2), Fig. 10I], such as *M. pitelkai*, *M. hobsonae*, and *M. hartmanae*.

Characters of the thoracic region (subjects 52–53)

Whilst many authors have noted the often-marked difference between thoracic and abdominal regions in magelonids (Figs. 8B and 15A), other characteristics of the thoracic region have been largely overlooked. Mortimer (2019) noted that among members of certain magelonid species, the thoracic interparapodial margins (**subject 52**) may be characteristically bulbous and rounded [character 52 (1), Fig. 11A], as seen among members of *Magelona allenii*, whilst among members of others, such as *M. johnstoni*, they are straight [character 52(0), Fig. 11B]. Members of the outgroups possess straight thoracic interparapodial margins [character 52(0)].

Uebelacker & Jones (1984) first coined the phrase ‘oblique lateral slits’ (**subject 53**) in relation to members of their undescribed species *Magelona* sp. I from the Gulf of Mexico. Later, Bolívar and Lana (1986) described ‘sulcos latero-dorsais’ (dorso-lateral slits) and ‘sulcos transversais’ (transverse slits) for members of *M. variolamellata* (Fig. 11C). Brasil (2003) used the term dorsal furrows/grooves, highlighting them to be present also among members of *M. polydentata* [character 53(1)], whilst herein, the term dorso-lateral grooves is utilised. Brasil (2003) stated that they vary in number, position, and location, and they were noted to be absent in the majority of members of magelonid species [character 53(0)] and are additionally absent among members of the outgroups.

Dimensions of post-chaetiger 9 segments (subject 54)

Uebelacker & Jones (1984) described posterior segments that are much longer than wide [character 54(1), Fig. 12A] among members of an undescribed species, *Magelona* sp. H, from the Gulf of Mexico. Later, *M. posterelongata* was the first species with individuals formally described with this feature. Many authors have ignored this characteristic, a problem compounded by the difficulty in collecting entire specimens. A further problem in coding this character is that abdominal chaetigers may vary in proportion along the length of the posterior. It is believed, however, that the majority of magelonid species possess abdominal chaetigers equal in length and width, or only marginally longer than their width; a characteristic shared with members of the outgroups [character 54(0)].

Characters of the parapodia of post-chaetiger nine segments (subjects 55–59)

The abdominal region of magelonids consists of many chaetigers carrying biramous parapodia. In contrast to the thorax, the abdominal lamellae of the notopodia and neuropodia are generally symmetrical and of similar size and shape in both rami. The abdominal region starts at chaetiger 9 for members of *Octomagelona* (Fig. 6) and chaetiger 10 for members of *Magelona* (Fig. 11A) and is marked by a change in chaetal type, from capillary chaetae to hooded hooks. Abdominal lamellar shape (**subject 55**) varies among members of species, from triangular to rounded. They may be basally constricted [those with rounded lamellae, character 55(1), Figs. 12C, 12H], or without a basal constriction, being broad based [character 55(0), Fig. 12B]. The latter situation being generally observed among members of species possessing more triangular lamellae, tapering to pointed tips (Fig. 12D). Subject 55 is not applicable to members of *Phyllochaetopterus limicolus*, however all other members of the outgroups have lamellae without a basal constriction [character 55(0)].

At the inner margins of chaetal rows, triangular to digitiform processes (**subject 56**) may be present [character 56(1), Figs. 12C, 12F, 12H, 12I]. Jones termed these structures dorsal medial lobes (DML) for notopodia and ventral medial lobes (VML) for neuropodia. Uebelacker and Jones (1984) believed that they may occur universally throughout the family, but this is a viewpoint not shared by the current authors. They are absent [character 56(0)] in members of the outgroups, however, this subject is considered not applicable to *Phyllochaetopterus limicolus*. In members of magelonid species, the size of the medial lobes in the abdomen vary widely, from minute and difficult to discern, to long and conspicuous structures, e.g., among members of *Magelona montera*. For comparison, the medial lobes were coded depending on their height relative to abdominal hooded hooks (**subject 57**): character 57(0) for members of species in which they are smaller than the hooks and character 57(1) when they are larger than the hooks. The latter subject is not applicable to the outgroups.

The abdominal lamellae may extend behind chaetal rows (**subject 58**). This postchaetal expansion (previously termed the interlamella) is often more conspicuous in the anterior abdomen and when present [character 58(1), Figs. 12C, 12E and 12H] it may be rounded to triangular. Among members of species in which it is absent [character 58(0), Figs. 12B, 13D and 14B], the abdominal hooded hooks arise from a distinct ridge (Figs. 12B and 13E). Postchaetal expansions are absent among members of the outgroups, whilst in members of *Phyllochaetopterus limicolus* the subject is considered inapplicable.

As stated above, abdominal lamellae (**subject 59**) in magelonids are generally symmetrical in size and shape [character 59(0), Figs. 12D and 12H] as in members of the outgroups. However, in some magelonids, such as members of *Magelona alleni*, the lamellae are

sub-equal between the two rami [character 59(1), Fig. 12G]. This subject is inapplicable to members of *Phyllochaetopterus limicolus*.

Characters of lateral abdominal pouches (subjects 60–65)

Lateral pouches (**subject 60**) are present among members of some species of magelonids, located laterally between the parapodia of certain abdominal chaetigers [character 60(1)]. Their presence [character 60(1)] or absence [character 60(0)] is often a key diagnostic feature, although their function is currently unknown (Jones, 1968; Mortimer & Mackie, 2014). Pouches vary in morphology, direction of opening, number, and location. However, descriptions of these structures have generally been limited, often only indicated as being present or absent. Fiege et al. (2000) described two types of lateral pouch: Σ -shaped occurring in the anterior abdomen, generally paired, and opening anteriorly, whilst C-shaped pouches were noted to occur on median and posterior abdominal chaetigers, opening posteriorly and occurring in pairs or singly. Mortimer (2010) suggested these terms should be abandoned to enable variations to be better understood, and subsequently, Mortimer & Mackie (2014) and Mortimer (2019) detailed further pouch morphologies. Following the latter authors, lateral pouches when present, were coded based on six subjects: arrangement of pouches (**subject 61**), direction of pouch openings (**subject 62**), posteriorly open lateral pouch arrangement (**subject 63**), lateral pouch distribution (**subject 64**), and margins of posteriorly open pouches (**subject 65**). Firstly, pouches are either all paired [character 61(0), Fig. 13C] or unpaired [character 61(1), Fig. 13B], or both types are present [character 61(2), Fig. 13A]. Secondly, both anteriorly and posteriorly opening pouches are present [character 62(1), Figs. 13A, 13E and 13G], e.g., members of *Magelona johnstoni*, or only posteriorly open pouches are present [character 62(0), Figs. 13B and 13C]. Posteriorly open pouches may occur on alternate segments [character 63(1), Fig. 13B], or on consecutive

segments [character 63(0), Fig. 13C], or alternatively, they may be present on both consecutive and alternating chaetigers at varying points along the abdomen [character 63(2)], e.g., members of *M. pulchella*. Fiege et al. (2000) described the first occurrence of lateral pouches on the body as an important distinguishing character between members of species and is relatively easy to observe. In members of some species, lateral pouches are present throughout most of the abdomen [character 64(0)], whilst in others they are present only on median and posterior chaetigers [character 64(1)], and in others restricted to the posterior abdomen only [character 64(2)] (Mortimer & Mackie, 2014). Whilst the margins of most posteriorly open pouches are smooth [character 65(0)], in members of some species, such as *M. pacifica*, they are medially slit [character 65(1), Fig. 13H]. One of the biggest problems in coding these characters is the lack of information known about the posterior regions of many magelonids, whilst some pouches are present throughout most of the abdomen and are therefore likely to be recorded even in posteriorly incomplete specimens. For members of species in which pouches only occur in the extreme posterior region, pouches are likely to be recorded as absent if individuals are described from anterior fragments only. Lateral pouches are present [character 60(1)] in members of outgroups, *Laonice cirrata* and *Prionospio ehlersi*. These are paired [character 61(0)], posteriorly open [character 62(0)] and with smooth margins [character 65(0)]. They occur on consecutive segments [character 63(0)] on median and posterior chaetigers [character 64(1)]. Lateral pouches are absent [character 60(0)] in the remaining members of the outgroups.

Characters of post-chaetiger 9 chaetae (subjects 66–74)

Magelonids possess hooded hooks in the abdominal region (**subject 66**), located between the lamellae and medial lobes (when present). They are present [character 66(1)] in members of all magelonid taxa and members of the outgroup except *Phyllochaetopterus limicolus* [character

66(0)]. In magelonids, hooded hooks generally occur in one row per ramus, but recent studies suggest that they can occur in two rows, at least in the medial part of the rami (Mortimer et al., 2020) (Fig. 14B for members of *M. equilamellae*). Although, this may prove to be an important characteristic in future studies, due to lack of information for the majority of species this was not considered in this study.

The dentition of abdominal hooks (**subject 67**) may be bidentate [character 67(0), e.g., members of *Magelona minuta* and *M. papillicornis*, Figs. 14C and 14D] with one secondary tooth above the main fang, tridentate [character 67(1), Fig. 14E], with two secondary teeth above, or polydentate [character 67(2)], having three or more secondary teeth above the main fang. Quadridentate, pentadentate, and hexodont hooks (Figs. 14F–H) have been recorded among Magelonidae, with some polydentate individuals carrying more than one type of hook. Where members of species have been recorded with more than one, the dentition which predominates was coded. For example, members of *M. minuta* are recorded to possess the odd sporadic tridentate hook (Mills & Mortimer, 2018), but carry predominately bidentate hooks. Members of the outgroup *Spio filicornis* possess bidentate [character 67(0)] hooded hooks, whilst members of *Prionospio lighti*, *P. ehlersi* and *Laonice cirrata* possess polydentate hooded hooks [character 67(2)]. This subject is not applicable to members of *Phyllochaetopterus limicolus*, which have uncini.

Jones (1971) was the first author to draw attention to the fact that in members of some species the hook at the base of the lamellae (**subject 68**) may be smaller than the others [character 68(1), Figs. 12E and 14I]. Later it has been suggested that not only their size but also their appearance may indicate groups of closely related species (*Magelona pitelkai*, *M. hobsonae*, *M. hartmanae*, *M. dakini*, *M. filiformis*, *M. capensis*). However, Brasil (2003), upon

re-examination of the holotypes of the first four species, observed no differences in shape, only differences in size. Among members of other magelonid species, the hook at the base of the lamellae is of equivalent size to the rest [character 68(0), Fig. 13D]. The subject is not applicable to members of the outgroups.

Uebelacker & Jones (1984) first reported the presence of enlarged chaetae (**subject 69**) in the abdominal region of members of four undescribed species from the Gulf of Mexico; *Magelona* sp. C, possessing an enlarged recurved hook with a minute apical tooth in each ramus; *Magelona* sp. D, with enlarged hooded apical spines (Fig. 12F); *Magelona* sp. E, possessing large bidentate hooks akin to ‘ordinary’ hooks; and lastly *Magelona* sp. H, with large unidentate recurved spines. Hernández-Alcántara & Solís-Weiss (2000) formally described members of two of these species (*Magelona* spp. D and H) as *Meredithia spinifera* and *M. uebelackerae*, respectively, erecting the new genus based on this distinctive feature. The authors stated that “although this character is the only one that differentiates this group from the other species described in this monogeneric group, it is sufficiently distinctive to make it a valid generic level character.” However, Mortimer & Mackie (2003) felt that stronger evidence for recognising other genera within the family was needed, preferring to follow Uebelacker & Jones (1984) by including species with enlarged abdominal chaetae in *Magelona*, and synonymising *Meredithia* with *Magelona*. Two additional species with enlarged abdominal chaetae have since been described: *Magelona magnahamata* and *Magelona falcifera*. For the current analysis, the presence [character 69(1)] and absence [character 69(0)] of enlarged chaetae in the abdomen was noted as well as the morphology of the distal ends (**subject 70**), i.e., recurved [character 70(0), Fig. 14K], spine-like [character 70(1)], or an enlarged normal hook [character 70(2)]. Enlarged

hooks are not present in members of the outgroups [character 69(0)], therefore subject 70 is not applicable (together with subject 69 for *Phyllochaetopterus limicolus*).

The number of hooded hooks per ramus in abdominal chaetigers (**subject 71**) is variable and characters distinguished as eight or more hooks per ramus [character 71(0), e.g., members of *Magelona equilamellae*, Fig. 14B, and members of outgroups *Prionospio lighti*, *P. ehlersi* and *Laonice cirrata*], and less than eight [character 71(1), Fig. 14J], such as among members of *M. papillicornis* and members of outgroup *Spio filicornis*. This separation in number was based on the examinations of specimens from a number of different species and assessing several chaetigers of the same specimen. As variations in the number of hooks have been observed along the length of the abdomen (e.g., members of *M. papillicornis* have a higher number of hooks in the mid-abdominal region), the number of hooks was counted in the anterior abdomen. Subject 71 is not applicable for members of *Phyllochaetopterus limicolus*. Hooks may be arranged in two formations (**subject 72**): occurring in two groups per ramus with main fangs arranged face-to-face (vis-à-vis) [character 72(0), Figs. 12C, 12G, 12H, 14B and 14J], or in unidirectional rows (vis-à-dos) with main fangs pointing in one direction [character 72(1), Fig. 13D]. Vis-à-vis orientation is more common among members of the Magelonidae, whilst vis-à-dos is present for all members of the outgroups (subject 72 is not applicable to *P. limicolus*).

Neuropodia with sabre chaetae (**subject 73**) are present [character 73(1)] among members of the spionid outgroups. Such chaetae are absent [character 73(0)] among members of *Phyllochaetopterus limicolus* and all ingroup species.

Internal support chaetae, or aciculae (**subject 74**), have been reported among members of several magelonid species. These curved and slender chaetae support the bases of the lamellae, and it is possible that they are more prevalent among members of species with larger abdominal

lamellae. The proximal ends of the aciculae may overlap in the junction between the noto- and neuropodia (Figs. 12C and 12H). Notes on their presence [character 74(1)] or absence [character 74(0)], as well as their description, are poorly reported in the literature. Jones (1978) stated that all magelonid abdominal chaetigers are supported by these structures, although this is not an observation supported by the current authors. Aciculae are generally more conspicuous in larger specimens, whilst slide preparations of parapodia may be necessary for smaller individuals. The relative obscurity of aciculae may be a factor in the limited inclusion in descriptions of specimens. Aciculae are absent in all members of the outgroups [character 74(0)].

Characters of body pigmentation (subjects 75–76)

Among members of some species pigmentation in the posterior thorax (**subject 75**) is evident even in fixed material [character 75(1)], often deep reddish to brown in colour. Very little is known about this character beyond presence or absence [character 75(0)]. When pigment is present, there are two patterns of distribution (**subject 76**). Among members of some species, such as *Magelona alleni* and *M. equilamellae*, the pigment forms a distinct band [character 76(0), Figs. 15A and 15C], often from chaetigers 4/5 to 8; among members of other species only light, dispersed pigmentation in posterior chaetigers is noted [character 76(1)], e.g., members of *M. symmetrica*. Moreover, members of *Magelona* sp. 5 from West Africa are noted to have distinct stripy pigmentation over much of the body (Fig. 15B). Pigmentation on the palps of members of some species, such as *M. mirabilis*, has been noted (see Mills & Mortimer, 2019: fig. 4.2.6). However, insufficient information is currently available to enable coding of this character. Relevant pigmentation is absent in members of the outgroups [character 75(0)].

Characters of granular bodies (subjects 77–78)

The surfaces of segments can have granular bodies (**subject 77**) appearing as spots, distributed in a variety of patterns on both the thorax and abdomen, and remain upon fixation and preservation of material [character 77(1), Figs. 6A, 9H and 15C]. Jones (1971) felt that this characteristic had no systematic significance. However, patterns appear to be species specific, particularly those of the thoracic region. The latter may appear as transverse stripes (Fig. 16A) or distinct smaller circular regions adjacent to the parapodia (Fig. 16B). Abdominal interparapodial areas commonly have granular bodies occurring in patches (**subject 78**) [character 78(1), Figs. 16C and 16D] among members of Magelonidae and may cover a large part of the lateral region between parapodia of adjacent chaetigers. Both thoracic and abdominal bodies may stain with dyes such as Rose Bengal, or be highlighted by methyl green, in which the granules appear white against the contrasting stain (Figs. 16B and 16C). Whilst staining patterns can be extremely useful in separating members of species (Figs. 11D, 11E and 16A), it was not included in the current study. However, it may prove useful in the future, as more methyl green patterns are described. Granular bodies and abdominal interparapodial areas are absent in members of the outgroups [character 77(0), character 78(0)].

Tube construction (subject 79)

In general, magelonids are relatively motile, burrowing more or less continually through sediments (Jumars et al., 2015), without constructing tubes [character 79(0)]. However, members of several species are known to build distinct multi-layered tubes covered in sand [character 79(1), Fig. 16E]. Mills & Mortimer (2019) investigated the permanency of these tube-lined burrows among members of the European species, *Magelona alleni*. Members of some species, such as *M. filiformis*, have been described living in fragile tubes of a secretion to which sand grains adhere. These are not considered as tubes *per se*, but are believed to be the worm's

response to removal from its habitat. Members of the outgroups, *Spio filicornis* and *Phyllochaetopterus limicolus*, are known to build distinct tubes [character 79(1)], whilst the situation is unknown for members of *Prionospio lighti*, *P. ehlersi* and *Laonice cirrata*.

Characters not included, but of possible future consideration

Prostomial features

Magelonid prostomia may carry distinct markings on either side of the dorsal muscular ridges (Figs. 4A, 4C–E, 5A, 5B and 5E), the pattern of which is generally species specific, whilst members of other species show no obvious markings (Figs. 4G and 4H). Although this can be an important diagnostic feature, this was not included in the current analysis due to difficulties in adequately describing this character.

Thoracic chaetae

Mills & Mortimer (2018) highlighted differences in the number of thoracic chaetae among members of different species, and additionally observed variations in number of chaetae on different chaetigers of the same animal for members of several magelonid species. They noted that members of some species, such as *Magelona cincta*, have distinctly splayed chaetae in the posterior thorax when compared to members of other species. Whilst these three characteristics may prove to be useful, limited information is currently available, prohibiting their use in the current study.

Features of the thoracic region

More recently, several additional characteristics of the thoracic region have been noted. Mortimer & Mackie (2009) and Mortimer (2019) described distinct V- and X-shaped ventral

markings in the mid-thoracic region of members of species such as *Magelona crenulifrons* and *M. pulchella* (Figs. 11D and 11E). Among members of other magelonid species, thoracic ventral pads/swellings have been recorded, such as those observed among members of *M. mirabilis* (Mortimer & Mackie, 2014) or *M. obockensis* (Mortimer, 2010). These swellings may be oval to reniform in shape (Figs. 11F and 11G). However, occurrence of these two features in members of the majority of species is unknown.

Although, the constriction at the thorax/abdomen junction can be marked and may vary between species, the extent to which it is constricted can be greatly influenced by preservation and fixation and was not considered herein. Despite this, however, it is a character worthy of further study, and should be detailed more fully in future descriptions. Brasil (2003) indicated that members of *Octomagelona* present a constriction despite lacking a 9th ‘thoracic’ chaetiger, suggesting that the character may not simply indicate a transition between the thorax and the abdomen.

Whilst average length and the average number of chaetigers appear to be species specific, the lack of information for members of many species prohibits this as a character in the analysis at the present time. Many individuals of magelonid species are described from anterior fragments only. The breadth of individuals, however, may prove an important character. Members of species like *Magelona minuta*, true to their name, rarely attain widths greater than 0.5 mm, being somewhat long and slender. Members of species such as *M. alleni* are distinctly broad and stout, often measuring over 1 mm in width (Mortimer et al., 2021). Additionally, Mills & Mortimer (2019) indicate that stouter magelonid species generally are shorter in comparison to width and have a fewer number of chaetigers. This is something which warrants further investigation.

Abdominal hooded hooks

Mills & Mortimer (2018) highlighted differences in the angle between the main fang and secondary teeth in abdominal hooded hooks of members of several magelonid species and further suggested characteristics such as the width and roundedness of the main fang, and the angle between it and the axis of the hook shaft, as noted for example by Jones (1963, 1977), may prove useful in future studies.

Pygidium

Magelonids possess a pair of digitiform pygidial cirri on either side of the pygidium (Figs. 16F and 16G). Uebelacker & Jones (1984) noted the presence of three pygidial cirri on individuals of an undescribed species, *Magelona* sp. B, from the Gulf of Mexico, however, it is believed that the third cirrus may actually represent the elongated, rounded tip of the pygidium itself. As this is the only magelonid that has been described with this feature, the number of pygidial cirri was not included in this study. Mills & Mortimer (2019) noted that pygidial cirri of members of *M. alleni* were more truncate and triangular (Fig. 16H) in comparison to other magelonids, however, this is something that warrants further investigation.

Whilst Rouse (2001) stated that the magelonid anus is terminal, early illustrations from McIntosh (1878) of members of *Magelona mirabilis* (possibly *M. johnstoni*; see Fiege et al., 2000) clearly show the anus in a distinctly ventral position. Unfortunately, for members of many species the pygidium is unknown, or the amount of information provided by authors has been relatively limited. When included in descriptions, the pygidium is often drawn from a dorsal perspective, thus conclusions about the position of the anus are often difficult to make. Mills & Mortimer (2019) investigated the situation among members of five European species, concluding that the anus was ventral in members of four species, whilst in the latter, *M. alleni*, it was distinctly terminal in position (Fig. 16H). Subsequently, Mortimer et al. (2021) have noted that

members of *Magelona* sp. 5 from West Africa also have a terminal anus. Whilst this may prove to be a valuable character in subsequent studies, the lack of information among individuals of many species precludes inclusion in the current study.

Inferring phylogenetic hypotheses

Phylogenetic inferences were performed using PAUP* 4.0b10 (Swofford, 2001), with all observations weighted equally and multiple subject–predicate relations (‘multistate characters’) treated as non-additive (Table S1). The following command string was executed (Larkin et al., 2006; Fitzhugh, 2010b; Fitzhugh et al., 2015; Nogueira et al., 2010; Nogueira et al., 2013):
hsearch enforce = no start = stepwise addseq = random nreps = 100000 nchuck = 5 chuckscore = 1; hsearch enforce = no start = current chuckscore = no. As noted earlier in **Methodological considerations**, the commands used are not tantamount to what has been incorrectly called ‘parsimony analysis.’ Rather, implementation of these commands is consistent with abductive reasoning for the purpose of causally accounting for differentially shared characters by way of common causes as fully as possible. Character transformation series were examined using Mesquite 3.61 (Maddison & Maddison, 2011).

RESULTS

The phylogenetic inference using data in Table S1 produced 2,417,600 cladograms of 404 steps each. The consistency index for each is 0.243 and retention index is 0.744. The strict consensus tree for all cladograms is shown in Fig. 17. What is notable is the mix of clades and grades within the monophyletic Magelonidae. A consequence of the overall arrangements of these phylogenetic hypotheses is that there can be no generic-level phylogenetic hypotheses that can be formally named without also incurring paraphyletic taxa. The result, as will be pointed out in

the **DISCUSSION**, is that the only (composite) phylogenetic hypothesis that can be formally recognised is Magelonidae. The Magelonidae clade involves seven phylogenetic hypotheses referring to the following respective characters:

1. Shovel-shaped prostomium: absent (0) → *present* (1)
2. Nuchal organs: present (1) → *absent* (0)
8. Prostomial ridges: absent (0) → *present* (1)
10. Palp origin: dorsal (0) → *ventral* (1)
11. Palp surface: non-papillate (0) → *papillate* (1)
14. Burrowing organ: absent (0) → *present* (1)
15. Body regionation: non-magelonid (0) → *magelonid-like* (1).

The present results indicate that *Octomagelona* cannot be maintained, given that the clade is nested within *Magelona*. And while the arrangements of the phylogenetic hypotheses among all the cladograms impose severe limits that only allow for formally recognising Magelonidae, there are no phylogenetic hypotheses to which the type genus *Magelona* refer. The implications of these results will be addressed in the **DISCUSSION**.

In addition to the Magelonidae, there are 20 additional clades, indicated by letters **a** through **t** in Fig. 17, from which phylogenetic hypotheses deserve mention. None of these clades can be formally named, given the various grade groups also present among the cladograms. Whilst serving as synapomorphies for these clades, those characters that are homoplasious (including reversals and convergence) are denoted by an asterisk (*):

- Clade a** – 22. Chaetigers 1–7 notopodial lamellae shape: foliaceous (1) → *filiform* (0)*
26. Chaetigers 1–7 neuropodial lamellae position relative to ~~notochaetae~~

- 1590 postchaetal (0) → *subchaetal* (1)*
- 1591 28. Chaetigers 1–7 neuropodial lamellae shape: foliaceous (1) →
- 1592 *filiform* (0)*
- 1593 **Clade b** – 5. Prostomial horns: present (1) → *absent* (0)*
- 1594 20. Development of neuropodia along thorax: different in some chaetigers,
- 1595 (1) → *all similar* (0)*
- 1596 26. Chaetigers 1–7 neuropodial lamellae position relative to ~~notochaetae~~  notochaetae
- 1597 subchaetal (1) → *postchaetal* (0)*
- 1598 30. Chaetigers 1–7 lengths of noto- and neuropodial lamellae: notopodial
- 1599 longer (1) → *equivalent* (0)*
- 1600 **Clade c** – 4. Prostomium distal shape: straight (2) → *triangular* (0)*
- 1601 6. Shape of prostomial horns: rudimentary (1) → *distinct* (0)*
- 1602 46. Chaetiger 9 neuropodial lamellae position: postchaetal (0) →
- 1603 *subchaetal* (1)*
- 1604 **Clade d** – 67. Post-chaetiger 9 noto- and neuropodial hooded hooks dentition:
- 1605 bidentate (0) → *tridentate* (1)*
- 1606 **Clade e** – 16. Number of anterior body region chaetigers: nine (1) → *eight* (0)
- 1607 18. Development of notopodia along thorax: all similar (0) → *different in*
- 1608 *some chaetigers* (1)*
- 1609 **Clade f** – 29. Chaetigers 1–7 neuropodial filiform lamellae shape distal ends:
- 1610 pointed (0) → *distally expanded/scoop shaped* (1)
- 1611 60. Post-chaetiger 9 lateral pouches: absent (0) → *present* (1)*
- 1612 **Clade g** – 22. Chaetigers 1-7 notopodial lamellae shape: filiform (0) → *foliaceous* (1)*

- 1613 35. Chaetiger 8 notopodial lamellae shape: filiform (0) → *foliaceous* 35(1)*
- 1614 55. Post-chaetiger 9 lamellae shape: without basal constriction (0) → *with*
- 1615 *basal constriction* (1)*
- 1616 **Clade h** – 12. Number of rows of proximal palp papillae: 10–14 (2) → 4–8 (1)*
- 1617 43. Chaetiger 9 notopodial lamellae shape: filiform (0) → *foliaceous* (1)*
- 1618 75. Pigmentation in posterior thorax: present (1) → *absent* (0)*
- 1619 **Clade i** – 18. Development of notopodia along thorax: different in some chaetigers (1)
- 1620 → *all similar* (0)*
- 1621 45. Chaetiger 9 dorsal superior lobes: absent (0) → *present* (1)*
- 1622 **Clade j** – 38. Chaetiger 8 neuropodial lamellae position: subchaetal (1) → *prechaetal*
- 1623 (2)*
- 1624 60. Post-chaetiger 9 lateral pouches: absent (0) → *present* (1)*
- 1625 **Clade k** – 22. Chaetigers 1-7 notopodial lamellae shape: foliaceous (1) → *filiform* (0)*
- 1626 57. Post-chaetiger 9 dorsal medial lobes length: smaller than hooks (0) →
- 1627 longer than hooks (1)*
- 1628 68. Post-chaetiger 9 hooded hooks adjacent to notopodial subchaetal
- 1629 or neuropodial suprachaetal lamellae: same size (0) → *smaller than rest*
- 1630 (1)*
- 1631 **Clade l** – 46. Chaetiger 9 neuropodial lamellae position: subchaetal (1) → *prechaetal*
- 1632 (2)*
- 1633 **Clade m** – 4. Prostomium distal shape: straight (2) → *rounded* (1)
- 1634 5. Prostomial horns: present (1) → *absent* (0)*
- 1635 42. Chaetiger 9 notopodial lamellae height: elongate (1) → *low* (0)*

- 1636 50. Lengths of chaetiger 9 fascicles relative to chaetigers 1–8 fascicles:
 1637 same length as chaetae in chaetigers 1–8 (0) → *shorter than chaetae in*
 1638 *chaetigers 1–8* (1)
- 1639 51. Distal ends of chaetae in chaetiger 9: gently tapered, similar to
 1640 chaetigers 1–8 (0) → *mucronate* (1)
- 1641 55. Post-chaetiger 9 lamellae shape: with basal constriction (1) →
 1642 *without basal constriction* (0)*
- 1643 61. Post-chaetiger 9 lateral pouch arrangement: unpaired (1) → *both*
 1644 *paired and unpaired* (2)*
- 1645 62. Direction of lateral pouch openings: posteriorly (0) → *posteriorly*
 1646 *and anteriorly* (1)*
- 1647 **Clade n** – 13. Number of rows of distal palp papillae: four (1) → *two* (0)*
- 1648 **Clade o** – 36. Chaetiger 8 margins of foliaceous notopodial lamellae: smooth (0) →
 1649 *bilobed* (2)
- 1650 44. Chaetiger 9 notopodial foliaceous lamellae margins: smooth (0) →
 1651 *crenulate* (1)*
- 1652 **Clade p** – 72. Post-chaetiger 9 arrangement of hooded hooks: vis-à-vis (0) → *vis-à-dos*
 1653 (1)*
- 1654 **Clade q** – 52. Shape of thoracic interparapodial margins: straight (0) → *rounded*,
 1655 *bulbous* (1)*
- 1656 60. Post-chaetiger 9 lateral pouches: present (1) → *absent* (0)*
- 1657 **Clade r** – 26. Chaetigers 1–7 neuropodial subchaetal lamellae position relative to
 1658 ~~notochaetae~~ varying in position along thorax (1) → *same position*

- 1659 *along thorax* (0)*
- 1660 41. Chaetiger 9 notopodial lamellae positions: postchaetal (0) → *subchaetal*
- 1661 (1)*
- 1662 **Clade s** – 57. Post-chaetiger 9 dorsal medial lobes length: smaller than hooks (0) →
- 1663 *longer than hooks* (1)*
- 1664 **Clade t** – 7. Prostomium dimensions: longer than wide (0) → *wider than long* (2)*
- 1665 33. Chaetigers 1-8 lengths of noto- and neuropodial capillary chaetae:
- 1666 equivalent (0) → *neuropodial longer* (2)*
- 1667

1668 DISCUSSION

1669 A point mentioned in the **RESULTS** is that only one composite phylogenetic hypothesis can be

1670 formally recognised, i.e., Magelonidae. There can be no phylogenetic hypothesis(es) to which

1671 the name *Magelona* refers that is not redundant with Magelonidae. From a nomenclatural, as

1672 opposed to strictly scientific perspective, this assertion is contrary to what is required by the

1673 International Code of Zoological Nomenclature (ICZN; 1999): at least one formal name at the

1674 rank of genus must be established relative to the ranks of family and species. The present results

1675 offer another good example (see also Fitzhugh, 2008a, 2010b; Nogueira et al., 2010; Nogueira et

1676 al., 2017) of the conflict that can occur between the science of biological systematics and a

1677 mandated international nomenclature system that is not entirely aligned with the goal of that

1678 science. Compromise between these two positions currently favours the nomenclatural system,

1679 but by properly acknowledging that formal taxon names should be defined in terms of being

1680 explanatory hypotheses (cf. **Methodological considerations**), the conflict between naming and

1681 scientific practice can be somewhat mitigated. Emendations of the formal names Magelonidae,

1682 *Magelona*, and *Octomagelona* in relation to the phylogenetic hypotheses inferred in this study
1683 are presented below.

1684 **Morphological characters in future specimen descriptions**

1685 Various authors have discussed the ‘crucial morphological characters’ in differentiating members
1686 of magelonid species. Jones (1963), in his review of magelonids of the Gulf of Mexico listed six:
1687 (1) the fine structure of the hooded hooks of the posterior region (subject 67); (2) the presence or
1688 absence of prostomial horns (subject 5); (3) the presence or absence of medial lamellae in the
1689 posterior region (subject 56); (4) the presence or absence of specialised chaetae of various types
1690 on chaetiger 9 (subject 51); (5) the morphology of the anterior lateral lamellae (subjects 18–30,
1691 34–49); and (6) the relative dimensions of the prostomium (subject 7). Blake (1996) listed seven
1692 ‘principal diagnostic characters important in differentiating species,’ adding to Jones’ list the
1693 following: (7) presence or absence of dorsal median lobes on thoracic notopodia (subjects 24, 37,
1694 45, superior dorsal lobes); (8) presence and location of lateral pouches between abdominal
1695 segments (subjects 60, 64); and (9) the presence/absence and form of interlamellae on abdominal
1696 parapodia (herein termed the postchaetal expansion behind chaetal rows, subject 58). Brasil
1697 (2003: fig. 23) concluded that the position and form of noto- and neuropodial lamellae, and the
1698 division of the thorax into three different blocks of segments (i.e., A – all anterior chaetigers with
1699 identical morphology; B – first eight chaetigers being identical but possessing a ninth chaetiger
1700 having a distinct morphology; and C – in which the first seven identical chaetigers differ from
1701 chaetigers eight and nine) reflected the relationships found within clades and sub-clades within
1702 the family.

1703 Of the characters previously mentioned, the current results suggest those of particular
1704 importance to be included in future descriptions of members of species are the presence/absence

of prostomial horns (in addition to their form, i.e., distinct or rudimentary), the relative dimensions of the prostomium, the morphology of the anterior lamellae, including presence/absence of superior dorsal lobes, the presence/absence of specialised chaetae on chaetiger 9, and the presence of lateral abdominal pouches. As highlighted by Brasil (2003), the thoracic lamellae are extremely diagnostic. Great care must be taken to fully describe and illustrate all thoracic chaetigers in descriptions, taking into account the size, shape and position of thoracic lamellae and whether/how they vary along the thorax. Often variations in position of the lamellae may be subtle between thoracic parapodia, particularly that of filiform subchaetal neuropodial lamellae. It is particularly important to examine the parapodia of chaetigers 8 and 9.

In addition to the characters already highlighted, the prostomial distal shape (subject 3), the number of prostomial ridges (subject 9), the shape of thoracic interparapodial margins (subject 52), the arrangement of abdominal lateral pouches (subject 63) and the arrangement of abdominal hooded hooks (subject 72) warrant further investigation and inclusion in descriptions.

1718

1719 Systematics

1720 Magelonidae Cunningham & Ramage, 1888, emended

1721 Type genus. *Magelona* by monotypy.

1722 **Definition.** A composite phylogenetic hypothesis (Fig. 17), causally accounting for (a) presence
1723 of a shovel-shaped prostomium, (b) absence of nuchal organs, (c) presence of prostomial ridges,
1724 (d) ventral palps with (e) papillate surfaces, (f) presence of a burrowing organ, and (g)
1725 magelonid-like body regionation.

Within a reproductively isolated population of individuals in the past, the following causal events occurred: (a') the presence of a shovel-shaped prostomium [character 1(1)] originated by unspecified mechanism(s) among individuals with a rounded prostomium [character 1(0)], subsequent to which the novel character became fixed in the population by an unspecified mechanism(s); (b') the loss of nuchal organs [character 2(1)] originated by unspecified mechanism(s) among individuals with nuchal organs [character 2(0)], subsequent to which the novel character became fixed in the population by an unspecified mechanism(s); (c') the presence of prostomial ridges [character 8(1)] originated by unspecified mechanism(s) among individuals with no ridges [character 8(0)], subsequent to which the novel character became fixed in the population by an unspecified mechanism(s); (d') the presence of ventrally inserted palps [character 10(1)] originated by unspecified mechanism(s) among individuals with dorsal palps [character 10(0)], subsequent to which the novel character became fixed in the population by an unspecified mechanism(s); (e') the presence of palps with papillae [character 11(1)] originated by unspecified mechanism(s) among individuals with smooth palps [character 11(0)], subsequent to which the novel character became fixed in the population by an unspecified mechanism(s); (f') the presence of a burrowing organ [character 14(1)] originated by unspecified mechanism(s) among individuals without such an organ [character 14(0)], subsequent to which the novel character became fixed in the population by an unspecified mechanism(s); (g') the magelonid-like body [character 15(1)] originated by unspecified mechanism(s) among individuals without such body regionation [character 15(0)], subsequent to which the novel character became fixed in the population by an unspecified mechanism(s). Following character origin/fixation events (a')–(g') was a population splitting event by unspecified mechanism(s) leading to individuals to which subsequent phylogenetic and specific hypotheses refer.

Remarks. Results of the phylogenetic inferences by Brasil (2003: figs. 45 and 46; Fig. 18) acknowledged the monophyly of Magelonidae based on three of the synapomorphies also referred to in this study (cf. **RESULTS**): presence of a spade-shaped prostomium [character 1(1); phylogenetic hypothesis a' in the above **Definition**] and ventral insertion of papillated palps [characters 10(1), 11(1); phylogenetic hypotheses d' and e', respectively, in the above **Definition**]. Fauchald and Rouse (1997: 103) stated evidence for monophyly of the Magelonidae as “Palps with rounded cross-section and a subdistal expanded area covered with papillae. Prostomium shovel-shaped.” They additionally added that “chaetiger 9 often modified with unusual postchaetal lobes or different chaetae from those segments in front or behind it or both.” This does not, however, follow for all members of species within the Magelonidae.

The most basal group in the phylogenetic study by Brasil (2003) (Fig. 18: ~~clade I~~) was that containing members of five species: *Magelona californica*, *M. minuta*, *M. pettiboneae*, *M. papillicornis* and *Magelona* sp. C of Uebelacker & Jones (1984). Members of these species possess postchaetal lamellae [characters 21(0), 26(0), 34(0), 38(0), 41(0), 46(0)] of equivalent size [character 30(0)] in both rami of thoracic chaetigers. **Clade b** in the current study (Fig. 17) contains the same five species, along with members of five additional species: *M. falcifera*, *M. fauchaldi*, *M. magnahamata*, *M. phyllisae* and *M. pygmaea*. As well as the characters highlighted by Brasil (2003), individuals in this clade possess smooth [characters 23(0), 36(0), 44(0)], filiform [characters 22(0), 28(0), 35(0), 40(0), 43(0), 48(0)] lamellae without superior dorsal lobes [characters 24(0), 37(0), 45(0)]. They also have prostomia the widths approximately equivalent to their lengths [character 7(1)], with straight [character 4(2)] and smooth [character 3(0)] prostomial anterior margins, but only one pair of prostomial ridges [character 9(0)]. As the names of two species within the clade highlight, i.e., *M. minuta* and *M. pygmaea*, members of

these species rarely attain the sizes seen in members of other magelonid species and are slender and thread-like. No lateral abdominal pouches have been recorded among members of these species [character 66(0)], although as noted above (cf. **Character descriptions**) this should be accepted with caution since lateral abdominal pouches are likely underreported within the family. As noted by Mills & Mortimer (2018) for members of *M. minuta*, granular bodies [subject 77(1)] appear as distinct transverse stripes in the thoracic region, and spots or stripes of this kind are certainly reported in over half of these species. This is a character worthy of further investigation.

The analysis by Brasil (2003: fig. 47; Fig. 18) highlighted a ~~clade VIII.2~~ containing members of eight species: *Magelona mirabilis*, *M. johnstoni*, *M. riojai*, *M. crenulata*, *Magelona* sp. B of Uebelacker & Jones (1984), *M. obockensis*, *M. jonesi* and *M. sacculata*. The synapomorphies of the clade include the presence of mucronate chaetae [character 51(1)] and abdominal hooded hooks in a vis-à-dos orientation [character 72(1)], the latter of which is uncommon amongst members of Magelonidae. **Clade m** (Fig. 17) in the present study includes all the above species (except *M. jonesi* which will be discussed below) and additionally members of *M. conversa*, *M. debeerei*, *M. parochilis*, *Magelona* sp. A of Uebelacker & Jones (1984) and *M. tinae*. Members of these species possess longer than wide prostomia [character 7(0)], with rounded [character 4(1)], smooth anterior margins [character 3(0)], carrying two pairs of prostomial ridges [character 9(1)] but lacking prostomial horns [character 5(0)]. The development of lamellae along the thorax is different [characters 18(1), 20(1)], particularly the lamellae of chaetiger 9 which are low and broad [characters 42(0), 47(0)] in comparison to the elongate lamellae of preceding chaetigers. In addition to the chaetae of chaetiger 9 being mucronate, they are shorter than those occurring on chaetigers 1–8 [character 50(1)]. Lateral

abdominal pouches are always present [character 60(1)] and whilst members of several species within the clade possess both anteriorly and posteriorly open pouches [characters 62(1)], only the latter is present in members of other species [character 62(0)]. **Clade m** is further divided into clades **o** and **p** (Fig. 17), separating the species based on arrangement of abdominal hooks (**clade o** with vis-à-vis and **clade p** with vis-à-dos).

As noted above, *Magelona jonesi* in the present study is plesiomorphic to all magelonid species in **clade a** (Fig. 17). However, despite possessing tapering chaetae in chaetiger 9 [character 51(0)], similar to preceding chaetigers, in the analysis of Brasil (2003) members of *M. jonesi* occurred in ~~clade VIII.3~~  the synapomorphies of which were the presence of mucronate chaetae [character 51(1)] and abdominal hooded hooks in vis-à-dos orientation. Members of this species are in need of redescription and this may help to resolve the situation in a future study. It is perhaps the orientation of the abdominal hooded hooks which is in most need of examination. Whilst Hartmann-Schröder (1980) illustrated the abdominal hooks in a vis-à-dos orientation, she made no mention of that orientation in her description. Looking at her neuropodial figures (Hartmann-Schröder, 1980: figs. 116 and 117), several hooks are drawn facing anteriorly. It has been recently highlighted by Mortimer et al. (2020) that *M. filiformis*, a species also originally recorded as possessing vis-à-dos hooks and occurring in ~~clade VIII.3~~ of Brasil (2003: fig. 47; Fig. 18) along with *M. pacifica*, has in fact a vis-à-vis orientation. This misinterpretation of orientation can happen in members of species in which the hooks are not split equally between the two groups of a ramus, particularly if the quality of the material is poor. This warrants further clarification for *M. jonesi* and also *M. pacifica*. Unfortunately for the latter species, the abdominal hooks in the type material are mostly all broken (Mortimer et al., 2012) making it extremely difficult to discern. This may explain why these three species, *M. jonesi*, *M. filiformis*

1818 and *M. pacifica* were part of clade VIII.2 with members of species possessing mucronate chaetae
1819 in Brasil's (2003) study.

1820 **Clade k** in the current results (Fig. 17) contains members of five species: *Magelona*
1821 *dakini*, *M. filiformis*, *M. hartmanae*, *M. hobsonae* and *M. pitelkai*. They share similarities in
1822 possessing prostomia which are longer than wide [character 7(0)], with straight [character 4(2)],
1823 smooth anterior margins [character 3(0)] formed into rudimentary horns [character 6(0)], and
1824 which carry two pairs of prostomial ridges [character 9(1)]. Individuals have similar
1825 development of lamellae along the thorax in the notopodia [character 18(0)] but varying
1826 development in the neuropodia [character 20(1)]. They possess smooth edged [characters 23(0),
1827 36(0), 44(0)], filiform thoracic lamellae [characters 22(0), 28(0), 35(0), 40(0), 43(0), 48(0)]
1828 which are larger in the notopodia [character 31(1)], and basally constricted abdominal lamellae
1829 [character 55(1)]. Posteriorly open lateral abdominal pouches are reported [character 60(1)] in all
1830 but one species, *M. hobsonae*, however members of this species are only known from short
1831 anterior fragments, so this character needs verification. Members of three species, *M. hartmanae*,
1832 *M. hobsonae* and *M. pitelkai*, within this clade also possess pennoned chaetae of chaetiger 9
1833 [character 51(2)].

1834 Perhaps worth of noting is **clade n** (Fig. 17), which includes members of species
1835 possessing prostomia with distinct horns [character 6(1)], triangular anterior margins [character
1836 4(0)] and two pairs of prostomial ridges [character 9(1)]. Specimens also possess foliaceous
1837 notopodial thoracic lamellae [characters 22(1), 35(1), 43(1)] which are larger than the
1838 neuropodial lamellae [character 30(1)]. Superior dorsal lobes are present on chaetigers 1–8
1839 [characters 24(1), 37(1)]. The thoracic neuropodial lamellae are filiform [characters 28(0),
1840 40(0), 48(0)] with additional postchaetal expansions on chaetigers 8 and 9 [characters 39(1),

49(1)] (N.B. the latter needs verifying in *Magelona anuheone*). Abdominal lamellae are rounded with basal constrictions [character 55(1)]. Whilst nothing more can be concluded from **clade r** (Fig. 17), which contains numerous polytomies, characters such as prostomial width may prove to be important characters in other clades. **Clade t** contains members of four species (Fig. 17), *M. berkeleyi*, *M. lenticulata*, *M. wilsoni* and *Magelona* sp. J of Uebelacker & Jones (1984), which all have wide prostomia [character 7(2)], rounded bulbous thoracic interparapodial margins [character 52(1)], and lack abdominal lateral pouches [character 60(0)].

***Magelona* F. Müller, 1858, emended**

Meredithia Hernández-Alcántara & Solís-Weiss, 2000

Octomagelona Aguirrezabalaga, Ceberio & Fiege, 2001

Definition. There is no phylogenetic hypothesis to which this genus-rank name refers (Fig. 17). The consequence is that ICZN (International Commission on Zoological Nomenclature, 1999) Article 13.1.1, where a name must be ‘accompanied by a description or definition that states in words characters that are purported to differentiate the taxon,’ cannot be reasonably applied.

Remarks. The name *Magelona*, lacking any associated phylogenetic hypothesis, is an empty placeholder that is only recognised to satisfy the requirement that species names be binomial. If one wishes to insist on a definition for the name *Magelona* or list of characters that ‘differentiate the taxon,’ those would have to be identical to what is provided for Magelonidae.

Hernández-Alcántara & Solís-Weiss (2000) introduced the genus *Meredithia* for members of species possessing large hooded recurved spines on abdominal chaetigers [Table 3: character 69(1)], adding the two species *M. spinifera* and *M. uebelackerae*. However, the nature of the enlarged chaetae differ among members of the two species; members of *M. spinifera*

possessing enlarged spines [Table 3: character 70(1)], whilst members of *M. uebelackerae* possess unidentate, enlarged re-curved hooks [Table 3: character 70(0)]. The descriptions of specimens with enlarged abdominal chaetae, but which differ greatly in other aspects (e.g., lacking prostomial horns and in the nature of the thoracic lamellae), called the validity of the genus into question (Mortimer & Mackie, 2003). Subsequently in the phylogenetic analysis performed by Brasil (2003) it was concluded that *Meredithia* is paraphyletic, which is in agreement with the current results. Herein, members of species possessing enlarged chaetae appear in three separate clades (Fig. 17: **clades b, c and r**), and the two species originally assigned to *Meredithia* are split between **clades c and r**. *Magelona falcifera*, *M. magnahamata* and *Magelona* sp. C of Uebelacker & Jones (1984) (**clade b**) are all species with members lacking prostomial horns, whilst *M. uebelackerae*, *Magelona* sp. E of Uebelacker & Jones (1984) (**clade c**) and *M. spinifera* (**clade r**) all possess distinct prostomial horns. The present results further support the synonymisation of *Meredithia* with *Magelona*.

Results of the phylogenetic inferences by Brasil (2003: fig. 45) additionally shed doubt on the validity of *Octomagelona*, with *O. bizkaiensis* and an undescribed species from Mexico as a paraphyletic group nested within *Magelona* and closely related to species such as *M. variolamellata*, *M. equilamellae*, *Magelona* sp. I of Uebelacker & Jones (1984) and *M. polydentata*. This finding is in agreement with the current analysis, indicating that *Octomagelona* cannot be maintained (Fig. 17). Whilst members of *Octomagelona* may differ from other members of *Magelona* in possessing only eight thoracic chaetigers, they share similarities with members of some species in terms of possessing a prostomium which is wider than long [character 7(2)], with rudimentary horns [character 6(0)], but with only one pair of prostomial ridges [character 9(0)], filiform subchaetal thoracic lamellae in the noto- and neuropodia

1887 [characters 21(1), 22(0), 26(1), 28(0)], without superior dorsal lobes [characters 24(0), 37(0),
1888 45(0)] and abdominal lamellae without basal constrictions [character 55(0)].

1889

1890 ACKNOWLEDGMENTS

1891 The authors would like to thank Kimberley Mills (Amgueddfa Cymru – National Museum Wales
1892 and School of Earth and Environmental Sciences, Cardiff University), Andrew Mackie and
1893 James Turner (Amgueddfa Cymru – National Museum Wales) for provision of images used
1894 within this publication.

1895

1896 REFERENCES

- 1897 **Achinstein P. 1970.** Inference to scientific laws. In: Stuewer RH, ed. *Volume V: historical and*
1898 *philosophical perspectives of science. Minnesota studies in the philosophy of science.*
1899 Minneapolis: University of Minnesota Press 87–111.
- 1900 **Achinstein P. 2001.** *The book of evidence.* New York: Oxford University Press.
- 1901 **Aguado MT, Glasby CJ, Schroeder PC, Weigert A, Bleidorn C. 2015.** The making of a
1902 branching annelid: an analysis of complete mitochondrial genome and ribosomal data
1903 of *Ramisyllis multicaudata*. *Scientific Reports* **5**:12072.
- 1904 **Aguirrezabalaga F, Ceberio A, Fiege D. 2001.** *Octomagelona bizkaiensis* (Polychaeta:
1905 Magelonidae) a new genus and species from the Capbreton Canyon (Bay of Biscay, north-
1906 east Atlantic). *Journal of the Marine Biological Association of the United Kingdom*
1907 **81**:221–224.

- 1908 **Aliseda A. 2006.** *Abductive reasoning: logical investigations into discovery and explanation.*
- 1909 Dordrecht: Springer.
- 1910 **Allen SR. 2016.** *A critical introduction to properties.* New York: Bloomsbury Academic.
- 1911 **Alston WP. 1993.** *The reliability of sense perception.* Ithaca: Cornell University Press.
- 1912 **Andrade SCS, Novo M, Kawauchi GY, Worsaae K, Pleijel F, Giribet G, Rouse GW. 2015.**
- 1913 Articulating “Archiannelids”: phylogenomics and annelid relationships, with emphasis on
- 1914 meiofaunal taxa. *Molecular Biology and Evolution* **32**:2860–2875.
- 1915 **Anjum RL, Mumford S. 2018.** *Causation in science and the methods of scientific discovery.*
- 1916 New York: Oxford University Press.
- 1917 **Archibald JK, Mort ME, Crawford DJ. 2003.** Bayesian inference of phylogeny: a non-
- 1918 technical primer. *Taxon* **52**:187–191.
- 1919 **Armstrong DM. 1997.** *A world of states of affairs.* New York: Cambridge University Press.
- 1920 **Assis LCS. 2014.** Testing evolutionary hypotheses: from Willi Hennig to angiosperm phylogeny
- 1921 group. *Cladistics* **30**:240–242.
- 1922 **Audi R. 1998.** *Epistemology: a contemporary introduction to the theory of knowledge.* New
- 1923 York: Routledge.
- 1924 **Auletta G, Ellis GFR, Jaeger L. 2008.** Top-down causation by information control: From a
- 1925 philosophical problem to a scientific research programme. *Journal of the Royal Society*
- 1926 *Interface* **5**:1159–1172.
- 1927 **Baker VR. 1996.** Hypotheses and geomorphological reasoning. In: Rhoads BL, Thorn CE, eds.
- 1928 *The scientific nature of geomorphology: proceedings of the 27th Binghamton symposium in*
- 1929 *geomorphology held 27–29 September 1996.* New York: Wiley 57–85.
- 1930 **Barker SF. 1957.** *Induction and hypothesis.* New York: Cornell University Press.

- 1931 **Barnes E. 1994.** Why P rather than Q? The curiosities of fact and foil. *Philosophical Studies*
1932 **73**:35–53.
- 1933 **Bartolomaeus T. 1995.** Secondary monociliarity in the Annelida: monociliated epidermal cells
1934 in larvae of *Magelona mirabilis* (Magelonida). *Microfauna Marina* **103**:327–332.
- 1935 **Beckers P, Helm C, Bartolomaeus T. 2019.** The anatomy and development of the nervous
1936 system in Magelonidae (Annelida) – insights into the evolution of the annelid brain. *BMC*
1937 *Evolutionary Biology* **19**:173.
- 1938 **Biswas S, Akey JM. 2006.** Genomic insights into positive selection. *Trends in Genetics* **22**:437–
1939 446.
- 1940 **Blake JA. 1996.** Family Magelonidae Cunningham and Ramage, 1888. In: Blake JA, Hilbig B,
1941 Scott PH, eds. *Taxonomic atlas of the benthic fauna of the Santa Maria Basin and Western*
1942 *Santa Barbara Channel, volume 6: the Annelida. Part 3: Polychaeta: Orbiniidae to*
1943 *Cossuridae*. Santa Barbara: Santa Barbara Museum of Natural History 252–261.
- 1944 **Bolívar GA, Lana PC. 1986.** Magelonidae (Annelida: Polychaeta) do litoral sudeste do Brasil.
1945 *Neritica* **1**:131–147.
- 1946 **Borda E, Kudenov JD, Bienhold C, Rouse GW. 2012.** Towards a revised Amphinomidae
1947 (Annelida, Amphinomida): description and affinities of a new genus and species from the
1948 Nile Deep-sea Fan, Mediterranean Sea. *Zoologica Scripta* **41**:307–325.
- 1949 **Brasil AC dos S. 2003.** Filogenia de Magelonidae Cunningham & Ramage, 1888 (Annelida –
1950 Polychaeta) com base na morfologia externa. Doctoral dissertation. Universidade Federal
1951 do Paraná.
- 1952 **Bremer K. 1988.** The limits of amino acid sequence data in angiosperm phylogenetic
1953 reconstruction. *Evolution* **42**:795–803.

- 1954 **Bremer K. 1994.** Branch support and tree stability. *Cladistics* **10**:295–304.
- 1955 **Bromberger S. 1966.** Why-questions. In: Colodny RG, ed. *Mind and cosmos: Essays in*
1956 *contemporary science and philosophy. University of Pittsburgh series in the philosophy of*
1957 *science, volume 3.* Pittsburgh: University of Pittsburgh Press 86–111.
- 1958 **Brower, AVZ. 2018.** Statistical consistency and phylogenetic inference: a brief review.
1959 *Cladistics* **34**:562–567.
- 1960 **Burton RB. 2000.** The problem of control in abduction. *Transactions of the Charles S. Peirce*
1961 *Society* **36**:149–156.
- 1962 **Callier V. 2018.** Theorists debate how ‘neutral’ evolution really is. *Quanta*. Available from:
1963 [https://www.quantamagazine.org/neutral-theory-of-evolution-challenged-by-evidence-for-](https://www.quantamagazine.org/neutral-theory-of-evolution-challenged-by-evidence-for-dna-selection-20181108/)
1964 [dna-selection-20181108/](https://www.quantamagazine.org/neutral-theory-of-evolution-challenged-by-evidence-for-dna-selection-20181108/).
- 1965 **Campbell DT. 1974.** Downward causation in hierarchically organized biological systems. In:
1966 Ayala FJ, Dobzhansky T, eds. *Studies in the philosophy of biology: reduction and related*
1967 *problems.* Berkeley: University of California Press 179–186.
- 1968 **Capa M, Parapar J, Hutchings P. 2012.** Phylogeny of Oweniidae (Polychaeta) based on
1969 morphological data and taxonomic revision of Australian fauna. *Zoological Journal of the*
1970 *Linnean Society* **166**:236–278.
- 1971 **Capa M, Parapar J, Hutchings P. 2019.** 4.1 Oweniidae Rioja, 1917. In: Purschke G,
1972 Böggemann, M, Westheide W, eds. *Volume 1: Annelida basal groups and Pleistoannelida,*
1973 *Sedentaria I.* Boston: De Gruyter 91–112.
- 1974 **Carnap R. 1950.** *Logical foundations of probability.* Chicago: University of Chicago Press.

- 1975 **Chen H, Parry LA, Vinther J, Zha, D, Hou X, Ma X. 2020.** A Cambrian crown annelid
1976 reconciles phylogenomics and the fossil record. *Nature* Available from:
1977 <https://doi.org/10.1038/s41586-020-2384-8>.
- 1978 **Cleland CE. 2001.** Historical science, experimental science, and the scientific method. *Geology*
1979 **29**:987–990.
- 1980 **Cleland CE. 2002.** Methodological and epistemic differences between historical science and
1981 experimental science. *Philosophy of Science* **69**:474–496.
- 1982 **Cleland CE. 2009.** Philosophical issues in natural history and its historiography. In: Tucker A,
1983 ed. *A companion to the philosophy of history and historiography*. Oxford: Wiley-Blackwell
1984 44–62.
- 1985 **Cleland CE. 2011.** Prediction and explanation in historical natural science. *British Journal for*
1986 *the Philosophy of Science* **62**:551–582.
- 1987 **Cleland CE. 2013.** Common cause explanation and the search for a smoking gun. In: Baker VR,
1988 ed. *Rethinking the fabric of geology. The Geological Society Special Papers* **502**:1–9.
- 1989 **Copi IM, Cohen C. 1998.** *Introduction to logic*. Upper Saddle River: Prentice-Hall.
- 1990 **Cunningham JT, Ramage GA. 1888.** The Polychaeta Sedentaria of the Firth of Forth.
1991 *Transactions of the Royal Society of Edinburgh* **33**:635–684.
- 1992 **Curd MV. 1980.** The logic of discovery: An analysis of three approaches. In: Nickles T, ed.
1993 *Scientific discovery, logic and rationality*. Dordrecht: D. Reidel Publishing Company 201–
1994 219.
- 1995 **Currie A. 2018.** *Rock, bone, and ruin: an optimist’s guide to the historical sciences*. Cambridge:
1996 The MIT Press.
- 1997 **Daston L, Galison P. 2007.** *Objectivity*. New York: Zone Books.

- 1998 **Davies PCW. 2012.** The epigenome and top-down causation. *Interface Focus* **2**:42–48.
- 1999 **Davis JI. 1995.** A phylogenetic structure for the monocotyledons, as inferred from chloroplast
2000 DNA restriction site variation, and a comparison of measures of clade support. *Systematic*
2001 *Botany* **20**:503–527.
- 2002 **Edwards AWF. 1972.** *Likelihood*. New York: Cambridge University Press
- 2003 **Efron B. 1979.** Bootstrap methods: Another look at the jackknife. *The Annals of Statistics* **7**:1–
2004 26.
- 2005 **Efron B, Halloran E, Holmes S. 1996.** Bootstrap confidence levels for phylogenetic trees.
2006 *Proceedings of the National Academy of Science* **93**:7085–7090.
- 2007 **Efron B, Gong G. 1983.** A leisurely look at the bootstrap, the jackknife, and cross-validation.
2008 *The American Statistician* **37**:36–48.
- 2009 **Efron B, Tibshirani RJ. 1993.** *An introduction to the bootstrap*. New York: Chapman & Hall.
- 2010 **Ellis G. 2013.** Time to turn cause and effect on their heads. *New Scientist* **2930**:28–29.
- 2011 **Ellis GFR. 2008.** On the nature of causation in complex systems. *Transactions of the Royal*
2012 *Society of South Africa* **63**:69–84.
- 2013 **Ellis GFR. 2012.** Top-down causation and emergence: some comments on mechanisms.
2014 *Interface Focus* **2**:126–140.
- 2015 **Ellis GFR, Noble D, O'Connor T. 2011.** Top-down causation: An integrating theme within and
2016 across the sciences? *Interface Focus* **2**:1–3.
- 2017 **Fann KT. 1970.** *Peirce's theory of abduction*. The Hague: Martinus Nijhoff.
- 2018 **Farris JS. 1983.** The logical basis of phylogenetic analysis. In: Platnick NI, Funk VA, eds.
2019 *Advances in cladistics, volume 2: proceedings of the second meeting of the Willi Hennig*
2020 *society*. New York: Columbia University Press 7–36.

- 2021 **Fauchald K, Jumars PA. 1979.** The diet of worms: a study of polychaete feeding guilds.
2022 *Oceanography and Marine Biology Annual Review* **17**: 93–284.
- 2023 **Fauchald K, Rouse G. 1997.** Polychaete systematics: past and present. *Zoologica Scripta*
2024 **26**:71–138.
- 2025 **Felsenstein J. 1978.** Cases in which parsimony or compatibility methods will be positively
2026 misleading. *Systematic Zoology* **27**:401–410.
- 2027 **Felsenstein J. 1981.** Evolutionary trees from DNA sequences: a maximum likelihood approach.
2028 *Journal of Molecular Evolution* **17**:368–376.
- 2029 **Felsenstein J. 1985.** Confidence limits on phylogenies: an approach using the bootstrap.
2030 *Evolution* **39**:783–791.
- 2031 **Felsenstein J. 2004.** *Inferring phylogenies*. Sunderland: Sinauer Associates Inc.
- 2032 **Fetzer JH. 1993.** *Philosophy of science*. New York: Paragon House.
- 2033 **Fetzer JH, Almeder RF. 1993.** *Glossary of epistemology/philosophy of science*. New York:
2034 Paragon House.
- 2035 **Fiege D, Licher F, Mackie ASY. 2000.** A partial review of the European Magelonidae
2036 (Annelida: Polychaeta): *Magelona mirabilis* redefined and *M. johnstoni* sp. nov.
2037 distinguished. *Journal of the Marine Biological Association of the United Kingdom*
2038 **80**:215–234.
- 2039 **Findlay JN. 1936.** Relational properties. *Australasian Journal of Psychology and Philosophy*
2040 **14**:176–190.
- 2041 **Fisher RA. 1934.** *Statistical methods for research workers*. Edinburgh: Oliver & Boyd.
- 2042 **Fitzhugh K. 2005a.** Les bases philosophiques de l’inférence phylogénétique: une vue
2043 d’ensemble. *Biosystema* **24**:83–105.

- 2044 **Fitzhugh K. 2005b.** The inferential basis of species hypotheses: the solution to defining the term
2045 ‘species’. *Marine Ecology* **26**:155–165.
- 2046 **Fitzhugh K. 2006a.** The abduction of phylogenetic hypotheses. *Zootaxa* **1145**:1–110.
- 2047 **Fitzhugh K. 2006b.** The ‘requirement of total evidence’ and its role in phylogenetic systematics.
2048 *Biology and Philosophy* **21**:309–351.
- 2049 **Fitzhugh K. 2006c.** The philosophical basis of character coding for the inference of
2050 phylogenetic hypotheses. *Zoologica Scripta* **35**:261–286.
- 2051 **Fitzhugh K. 2008a.** Abductive inference: implications for ‘Linnean’ and ‘phylogenetic’
2052 approaches for representing biological systematization. *Evolutionary Biology* **35**:52–82.
- 2053 **Fitzhugh K. 2008b.** Fact, theory, test and evolution. *Zoologica Scripta* **37**:109–113.
- 2054 **Fitzhugh K. 2008c.** Clarifying the role of character loss in phylogenetic inference. *Zoologica*
2055 *Scripta* **37**:561–569.
- 2056 **Fitzhugh K. 2009.** Species as explanatory hypotheses: refinements and implications. *Acta*
2057 *Biotheoretica* **57**:201–248.
- 2058 **Fitzhugh K. 2010a.** Evidence for evolution versus evidence for intelligent design: parallel
2059 confusions. *Evolutionary Biology* **37**:68–92.
- 2060 **Fitzhugh K. 2010b.** Revised systematics of *Fabricia oregonica* Banse, 1956 (Polychaeta:
2061 Sabellidae: Fabriciinae): an example of the need for a uninominal nomenclatural system.
2062 *Zootaxa* **2647**:35–50.
- 2063 **Fitzhugh K. 2012.** The limits of understanding in biological systematics. *Zootaxa* **3435**:40–67.
- 2064 **Fitzhugh K. 2013.** Defining ‘species’, ‘biodiversity’, and ‘conservation’ by their transitive
2065 relations. In: Pavlinov IY, ed. *The species problem—Ongoing problems*. New York:
2066 InTech 93–130.

- 2067 **Fitzhugh K. 2014.** Character mapping and cladogram comparison versus the requirement of
2068 total evidence: does it matter for polychaete systematics? *Memoirs of Museum Victoria*
2069 **71**:67–78.
- 2070 **Fitzhugh K. 2015.** What are species? Or, on asking the wrong question. *The Festivus* **47**:229–
2071 239.
- 2072 **Fitzhugh K. 2016a.** Ernst Mayr, causal understanding, and systematics: an example using
2073 sabelliform polychaetes. *Invertebrate Biology* **135**:302–313.
- 2074 **Fitzhugh K. 2016b.** Sequence data, phylogenetic inference, and implications of downward
2075 causation. *Acta Biotheoretica* **64**:133–160.
- 2076 **Fitzhugh K. 2016c.** Dispelling five myths about hypothesis testing in biological systematics.
2077 *Organisms Diversity & Evolution* **16**:443–465.
- 2078 **Fitzhugh K. 2016d.** Phylogenetic hypotheses: neither testable nor falsifiable. *Evolutionary*
2079 *Biology* **43**:257–266.
- 2080 **Fitzhugh K. 2021.** Phylogenetic inference and the misplaced premise of substitution rates. *Acta*
2081 *Biotheoretica*. [OI 10.1007/s10441-021-09412-4](https://doi.org/10.1007/s10441-021-09412-4).
- 2082 **Fitzhugh K, Nogueira JMM, Carrerette O, Hutchings P. 2015.** An assessment of the status of
2083 Polycirridae genera (Annelida: Terebelliformia) and evolutionary transformation series of
2084 characters within the family. *Zoological Journal of the Linnean Society* **174**:666–701.
- 2085 **Francescotti R. 2014.** Intrinsic/extrinsic: a relational account defended. In: Francescotti RM, ed.
2086 *Companion to intrinsic properties*. Berlin: De Gruyter 175–198.
- 2087 **Gabbay DM, Woods J. 2005.** *The reach of abduction: insight and trial. A practical logic of*
2088 *cognitive systems, volume 2*. Amsterdam: Elsevier.
- 2089 **Gaukroger S. 2012.** *Objectivity: a very short introduction*. New York: Oxford University Press.

- 2090 **Glasby CJ, Schroeder PC, Aguado MT. 2012.** Branching out: a remarkable new branching
2091 syllid (Annelida) living in a *Petrosia* sponge (Porifera: Demospongiae). *Zoological Journal*
2092 *of the Linnean Society* **164**:481–497.
- 2093 **Godfrey-Smith P. 2003.** *Theory and reality: an introduction to the philosophy of science.*
2094 Chicago: University of Chicago Press.
- 2095 **Gonzalez BC, Martínez A, Borda E, Iliffe TM, Eibye-Jacobsen D, Worsaae K. 2018.**
2096 Phylogeny and systematics of Aphroditiformia. *Cladistics* **34**:225–259.
- 2097 **Goto R. 2016.** A comprehensive molecular phylogeny of spoon worms (Echiura, Annelida):
2098 Implications for morphological evolution, the origin of dwarf males, and habitat shifts.
2099 *Molecular Phylogenetics and Evolution* **99**:247–260.
- 2100 **Goto R, Okamoto T, Ishikawa H, Hamamura Y, Kato M. 2013.** Molecular phylogeny of
2101 echiuran worms (Phylum: Annelida) reveals evolutionary pattern of feeding mode and
2102 sexual dimorphism. *PLOS One* **8**:e56809.
- 2103 **Gracia JJE. 1988.** *Individuality: an essay on the foundations of metaphysics.* Albany: State
2104 University of New York Press.
- 2105 **Griffiths P, Stotz K. 2013.** *Genetics and philosophy: an introduction.* New York: Cambridge
2106 University Press.
- 2107 **Haber M. 2011.** Phylogenetic inference. In: Tucker A, ed. *A companion to the philosophy of*
2108 *history and historiography.* Malden: Wiley-Blackwell 231–242.
- 2109 **Hacking I. 1965.** *Logic of statistical inference.* New York: Cambridge University Press.
- 2110 **Hacking I. 2001.** *An introduction to probability and inductive logic.* New York: Cambridge
2111 University Press.

- 2112 **Halanych KM. 2004.** The new view of animal phylogeny. *Annual Review of Ecology and*
2113 *Evolutionary Systematics* **35**:229–256.
- 2114 **Halanych KM. 2016.** How our view of animal phylogeny was reshaped by molecular
2115 approaches: lessons learned. *Organisms Diversity & Evolution* **16**:319–328.
- 2116 **Hanson NR. 1958.** *Patterns of discovery: an inquiry into the conceptual foundations of science.*
2117 New York: Cambridge University Press.
- 2118 **Hartman O. 1944.** Polychaetous annelids from California, including two new genera and nine
2119 new species. *Allan Hancock Pacific Expeditions* **10**:239–304.
- 2120 **Hartman O. 1961.** Polychaetous annelids from California. *Allan Hancock Pacific Expeditions*
2121 **25**:1–226.
- 2122 **Hartman O. 1965.** Deep-water benthic polychaetous annelids off New England to Bermuda and
2123 other North Atlantic areas. *Occasional Papers of the Allan Hancock Foundation* **28**:1–378.
- 2124 **Hartman O. 1971.** Abyssal polychaetous annelids from the Mozambique Basin off southeast
2125 Africa, with a compendium of abyssal polychaetous annelids from world-wide areas.
2126 *Journal of the Fisheries Research Board of Canada* **28**:1407–1428.
- 2127 **Hartmann-Schröder G. 1980.** Die Polychaeten der tropischen Nordwestküste Australiens
2128 (zwischen Port Samson im Norden und Exmouth im Süden) In: Hartmann-Schröder G,
2129 Hartmann G, eds. *Zur Kenntnis des Eulitorals der australischen Küsten unter besonder*
2130 *Berücksichtigung der Polychaeten und Ostracoden. Mitteilungen aus dem Hamburgischen*
2131 *zoologischen Museum und Institut.* **77**:41–110.
- 2132 **Hausman DM. 1998.** *Causal asymmetries.* New York: Cambridge University Press.
- 2133 **Heath T A, Hedtke SM, Hillis DM. 2008.** Taxon sampling and the accuracy of phylogenetic
2134 analysis. *Journal of Systematics and Evolution* **46**:239–257.

- 2135 **Helm C, Beckers P, Bartolomaeus T, Drukewitz SH, Kourtesis I, Weigert A, Purschke G,**
2136 **Worsaae K, Struck TH, Bleidorn C. 2018.** Convergent evolution of the ladder-like
2137 ventral nerve cord in Annelida. *Frontiers in Zoology* **15**. Available from:
2138 <https://doi.org/10.1186/s12983-018-0280-y>.
- 2139 **Hempel CG. 1962.** Deductive nomological vs. statistical explanation. In: Feigl H, Maxwell G,
2140 eds. *Minnesota studies in the philosophy of science, volume 3*. Minneapolis: University of
2141 Minnesota Pres 98–169.
- 2142 **Hempel CG. 1965.** *Aspects of scientific explanation and other essays in the philosophy of*
2143 *science*. New York: The Free Press.
- 2144 **Hempel CG. 1966.** Recent problems of induction. In: Colodny RG, ed. *Mind and cosmos*.
2145 Pittsburgh: University of Pittsburgh Press 112–134.
- 2146 **Hempel CG. 2001.** *The philosophy of Carl G. Hempel: studies in science, explanation, and*
2147 *rationality*. In: Fetzer JH, ed. New York: Oxford University Press.
- 2148 **Hennig W. 1966.** *Phylogenetic systematics*. Urbana: University of Illinois Press.
- 2149 **Hernández-Alcántara P, Solís-Weiss V. 2000.** Magelonidae from the Mexican Pacific and
2150 northern Gulf of Mexico, with the description of a new genus (*Meridithia*) and four new
2151 species. In: Reish DJ, Lana P, eds. Proceedings of the 6th international polychaete
2152 conference, Curitiba, Brazil, 1998. *Bulletin of Marine Science* **67**:625–644.
- 2153 **Hey J. 1999.** The neutralist, the fly and the selectionist. *Trends in Ecology & Evolution* **14**:35–
2154 38.
- 2155 **Hoffmann M. 1995.** Problems with Peirce’s concept of abduction. *Foundations of Science*
2156 **4**:271–305.

2157 **Holmes S. 2003.** Bootstrapping phylogenetic trees: theory and methods. *Statistical Science*
2158 **18**:241–255.

2159 **Howson C, Urbach P. 1993.** *Scientific reasoning: the Bayesian approach*. Chicago: Open
2160 Court.

2161 **Hoyningen-Huene P. 2013.** *Systematicity: the nature of science*. New York: Oxford University
2162 Press.

2163 **Huelsenbeck JP, Crandall KA. 1997.** Phylogeny estimation and hypothesis testing using
2164 maximum likelihood. *Annual Review of Ecology and Systematics* **28**:437–466.

2165 **Huelsenbeck JP, Ronquist F. 2001.** MrBayes: Bayesian inference of phylogeny. *Bioinformatics*
2166 **17**:754–755.

2167 **Huelsenbeck JP, Ronquist F, Nielsen R, Bollback JP. 2001.** Bayesian inference of phylogeny
2168 and its impact on evolutionary biology. *Science* **294**:2310–2314.

2169 **Hurst LD. 2002.** The Ka/Ks ratio: diagnosing the form of sequence evolution. *Trends in*
2170 *Genetics* **18**:486–487.

2171 **International Commission on Zoological Nomenclature. 1999.** *International code of*
2172 *zoological nomenclature*. London: The International Trust for Zoological Nomenclature.

2173 **Jaeger L, Calkins ER. 2011.** Downward causation by information control in micro-organisms.
2174 *Interface Focus*; **2**:26–41.

2175 **Jeffares B. 2008.** Testing times: regularities in the historical sciences. *Studies in History and*
2176 *Philosophy of Biological and Biomedical Sciences* **39**:469–475.

2177 **Johnston GA. 1865.** *Catalogue of the British non-parasitical worms in the collection of the*
2178 *British Museum*. London: Trustees of the British Museum.

- 2179 **Jones ML. 1963.** Four new species of *Magelona* (Annelida, Polychaeta) and a redescription of
2180 *Magelona longicornis* Johnson. *American Museum Novitates* **2164**:1–31.
- 2181 **Jones ML. 1968.** On the morphology, feeding, and behavior of *Magelona* sp. *Biological Bulletin*
2182 **134**:272–297.
- 2183 **Jones ML. 1971.** *Magelona berkeleyi* n. sp. from Puget Sound (Annelida: Polychaeta) with a
2184 further redescription of *Magelona longicornis* Johnson and a consideration of recently
2185 described species of *Magelona*. *Journal of the Fisheries Research Board of Canada*
2186 **28**:1445–1454.
- 2187 **Jones ML. 1977.** A redescription of *Magelona papillicornis* F. Müller. In: Reish DJ, Fauchald
2188 K, eds. *Essays on polychaetous annelids in memory of Dr. Olga Hartman*. Los Angeles:
2189 Allan Hancock Foundation, University of Southern California 247–266.
- 2190 **Jones ML. 1978.** Three new species of *Magelona* (Annelida, Polychaeta) and a redescription of
2191 *Magelona pitelkai* Hartman. *Proceedings of the Biological Society of Washington* **91**:336–
2192 363.
- 2193 **Josephson JR, Josephson SG, eds. 1994.** *Abductive inference: computation, philosophy,*
2194 *technology*. New York: Cambridge University Press.
- 2195 **Jukes TH, Cantor CR. 1969.** Evolution of protein molecules. In: Munro HN, ed. *Mammalian*
2196 *protein metabolism*. New York: Academic Press 21–132.
- 2197 **Jumars PA, Kelly, MD, Lindsay SM. 2015.** Diet of worms emended: an update of polychaete
2198 feeding guilds. *Annual Review of Marine Science* **7**:497–520.
- 2199 **Kapitan T. 1997.** Peirce and the structure of abductive inference. In: Houser N, Roberts DD,
2200 Evra JV, eds. *Studies in the logic of Charles Sanders Peirce*. Bloomington: Indiana
2201 University Press 477–496.

- 2202 **Kapli P, Yang Z, Telford MJ. 2020.** Phylogenetic tree building in the genomic age. *Nature*
2203 *Reviews Genetics* **21**:428–444.
- 2204 **Klienhans MG, Buskes CJJ, de Regt HW. 2005.** Terra incognita: explanation and reduction in
2205 Earth science. *International Studies in the Philosophy of Science* **19**:289–317.
- 2206 **Klienhans MG, Buskes CJJ, de Regt HW. 2010.** Philosophy of Earth science. In: Allhoff F,
2207 ed. *Philosophies of the sciences: a guide*. Oxford: Wiley-Blackwell 213–236.
- 2208 **Kneale W, Kneale M. 1964.** *The development of logic*. New York: Oxford University Press.
- 2209 **Kobayashi G, Goto R, Takano T, Kojima S. 2018.** Molecular phylogeny of Maldanidae
2210 (Annelida): multiple losses of tube-capping plates and evolutionary shifts in habitat depth.
2211 *Molecular Phylogenetics and Evolution* **127**:332–344.
- 2212 **Koonin EV. 2012.** *The logic of chance: the nature and origin of biological evolution*. Upper
2213 Saddle River: FTS Press.
- 2214 **Laland KN, Sterelny K, Odling-Smee J, Hoppitt W, Uller T. 2011.** Cause and effect in
2215 biology revisited: Is Mayr’s proximate-ultimate dichotomy still useful? *Science* **334**:1512–
2216 1516.
- 2217 **Langeneck J, Barbieri M, Maltagliati F, Castelli A. 2019.** Molecular phylogeny of Paraonidae
2218 (Annelida). *Molecular Phylogenetics and Evolution* **136**:1–13.
- 2219 **Lankester ER. 1870.** II.—On the use of the term homology in modern zoology, and the
2220 distinction between homogenetic and homoplastic agreements. *Annals and Magazine of*
2221 *Natural History* **6**:34–43.
- 2222 **Larkin LL, Neff JL, Simpson BB. 2006.** Phylogeny of the *Callandrena* subgenus of *Andrena*
2223 (Hymenoptera: Andrenidae) based on mitochondrial and nuclear DNA data: polyphyly and
2224 convergent evolution. *Molecular Phylogenetics and Evolution* **38**:330–343.

- 2225 **Lavelle JS, Botterill G, Lock S. 2013.** Contrastive explanation and the many absences problem.
2226 *Synthese* **190**:3495–3510.
- 2227 **Lemoine F, Entfellner J-BD, Wilkinson E, Correia D, Felipe MD, De Oliveira T, Gascuel**
2228 **O. 2018.** Renewing Felsenstein’s phylogenetic bootstrap in the era of big data. *Nature*
2229 **556**:452–456.
- 2230 **Linguist S, Doolittle WF, Palazzo AF. 2020.** Getting clear about the F-word in genomics.
2231 *PLOS Genetics* **16**:e1008702.
- 2232 **Lipton P. 2004.** *Inference to the best explanation*. New York: Routledge.
- 2233 **Lipton P. 2005.** Testing hypotheses: prediction and prejudice. *Science* **307**:219–221.
- 2234 **Lloyd EA. 1988.** *The structure and confirmation of evolutionary theory*. Princeton: Princeton
2235 University Press.
- 2236 **Longino HE. 1979.** Evidence and hypothesis: an analysis of evidential relations. *Philosophy of*
2237 *Science* **46**:35–56.
- 2238 **Lovell LL, Fitzhugh K. 2020.** Taking a closer look: an SEM review of *Levinsenia* species
2239 (Polychaeta: Paraonidae) reported from California. *Zootaxa* **4751**:257–275.
- 2240 **Maddison WP, Maddison DR. 2011.** *Mesquite: a modular system for evolutionary analysis*.
2241 *Version 3.61*. Available from: <http://mesquiteproject.com>.
- 2242 **Magnani L. 2001.** *Abduction, reason, and science: processes of discovery and explanation*. New
2243 York: Kluwer Academic.
- 2244 **Magnani L. 2009.** *Abductive cognition: the epistemological and eco-cognitive dimensions of*
2245 *hypothetical reasoning*. Berlin: Springer-Verlag.
- 2246 **Magnani L. 2017.** *The abductive structure of scientific creativity: an essay on the ecology of*
2247 *cognition*. Berlin: Springer-Verlag.

- 2248 **Mahner M, Bunge M. 1997.** *Foundations of biophilosophy*. New York: Springer.
- 2249 **Martínez M, Esposito M. 2014.** Multilevel causation and the extended synthesis. *Biological*
- 2250 *Theory* **9**:209–220.
- 2251 **Martínez M, Moya A. 2011.** Natural selection and multi-level causation. *Philosophy, Theory,*
- 2252 *and Practice in Biology* **3**:e202.
- 2253 **Marwick P. 1999.** Interrogatives and contrasts in explanation theory. *Philosophical Studies*
- 2254 **96**:183–204.
- 2255 **Mayr E. 1961.** Cause and effect in biology. *Science* **131**:1501–1506.
- 2256 **Mayr E. 1993.** Proximate and ultimate causation. *Biology and Philosophy* **8**:95–98.
- 2257 **McDonald JH, Kreitman M. 1991.** Adaptive protein evolution at the *Adh* locus in *Drosophila*.
- 2258 *Nature* **351**:652–654.
- 2259 **McIntosh WC. 1877.** On the structure of *Magelona*. *Annals and Magazine of Natural History,*
- 2260 *Series 4* **20**:147–152.
- 2261 **McIntosh WC. 1878.** Beiträge zur Anatomie von *Magelona*. *Zeitschrift für Wissenschaftliche*
- 2262 *Zoologie* **31**:401–472.
- 2263 **McIntosh WC. 1879.** On the circulatory system of *Magelona*. *Journal of Anatomy and*
- 2264 *Physiology* **13**:331–345.
- 2265 **McIntosh WC. 1911.** On the structure of *Magelona*. *Annals and Magazine of Natural History,*
- 2266 *Series 8* **7**:417–457. [Translation of McIntosh, 1878].
- 2267 **Mills K, Mortimer K. 2018.** Redescription of *Magelona minuta* Eliason, 1962 (Annelida), with
- 2268 discussions on the validity of *Magelona filiformis minuta*. *Zootaxa* **4527**:541–559.
- 2269 **Mills K, Mortimer K. 2019.** Observations on the tubicolous annelid *Magelona allenii*
- 2270 (Magelonidae), with discussions on the relationship between morphology and behaviour of

- 2271 European magelonids. *Journal of the Marine Biological Association of the United Kingdom*
- 2272 **99**:715–727.
- 2273 **Mortimer K. 2010.** Magelonidae (Polychaeta) from the Arabian Peninsula: A review of known
- 2274 species, with notes on *Magelona tinae* from Thailand. *Zootaxa* **2628**:1–26.
- 2275 **Mortimer K. 2019.** 4.2 Magelonidae Cunningham & Ramage, 1888. In: Purschke G,
- 2276 Böggemann M, Westheide W, eds. *Volume 1: Annelida basal groups and Pleistoannelida,*
- 2277 *Sedentaria I.* Boston: De Gruyter 112–131.
- 2278 **Mortimer K, Cassà S, Martin D, Gil J. 2012.** New records and new species of Magelonidae
- 2279 (Polychaeta) from the Arabian Peninsula, with a re-description of *Magelona pacifica* and a
- 2280 discussion on the magelonid buccal region. *Zootaxa* **3331**:1–43.
- 2281 **Mortimer K, Kongsrud JA, Willassen E. 2021.** Integrative taxonomy of West African
- 2282 *Magelona* (Annelida: Magelonidae): species carrying posterior thoracic pigmentation.
- 2283 *Zoological Journal of the Linnean Society* **—**.
- 2284 **Mortimer K, Mackie ASY. 2003.** The Magelonidae (Annelida: Polychaeta) from the
- 2285 Seychelles, with the description of three new species. In: Sigvaldadóttir E, Mackie ASY,
- 2286 Helgason GV, Reish DJ, Svavarsson J, Steingrímsson SA, Gudmundsson G, eds. *Advances*
- 2287 *in polychaete research.* *Hydrobiologia* **496**:163–173.
- 2288 **Mortimer K, Mackie ASY. 2009.** Magelonidae (Polychaeta) from Hong Kong, China; with
- 2289 discussions on related species and redescrptions of three species. In: Maciolek NJ, Blake
- 2290 JA, eds. *Proceedings of the ninth international polychaete conference.* *Zoosymposia* **2**:179–
- 2291 199.

- 2292 **Mortimer K, Mackie ASY. 2014.** Morphology, feeding and behaviour of British *Magelona*
2293 (Annelida: Magelonidae), with discussions on the form and function of abdominal lateral
2294 pouches. *Memoirs of Museum Victoria* **71**:177–201.
- 2295 **Mortimer K, Mills K, Jordana E, Pinedo S, Gil J. 2020.** A further review of European
2296 Magelonidae (Annelida), including redescrptions of *Magelona equilamellae* and *Magelona*
2297 *filiformis*. *Zootaxa* **4767**:89–114.
- 2298 **Mundy NI. 2016.** Population genomics fits the bill: genetics of adaptive beak variation in
2299 Darwin’s finches. *Molecular Ecology* **25**:5265–5266.
- 2300 **Nateewathana A, Hylleberg J. 1991.** Magelonid polychaetes from Thailand, the Andaman Sea,
2301 with descriptions of eight new species. In: Petersen ME, Kirkegaard JB, eds. *Systematics,*
2302 *biology and morphology of world Polychaeta. Proceedings of the 2nd international*
2303 *polychaete conference, Copenhagen, 1986. Ophelia Supplement* **5**:169–184.
- 2304 **Neta R. 2008.** What evidence do you have? *The British Journal for the Philosophy of Science*
2305 **59**:89–119.
- 2306 **Nickles T. 1980.** Introductory essay: scientific discovery and the future of philosophy of science.
2307 In: Nickles T, ed. *Scientific discovery, logic and rationality*. Dordrecht: D. Reidel
2308 Publishing Company 1–59.
- 2309 **Nogueira JMM, Carrerette O, Hutchings P, Fitzhugh K. 2018.** Systematic review of the
2310 species of the family Telothelepodidae Nogueira, Fitzhugh & Hutchings, 2013 (Annelida:
2311 Polychaeta), with descriptions of three new species. *Marine Biology Research* **14**:217–257.
- 2312 **Nogueira JMM, Fitzhugh K, Hutchings P. 2013.** The continuing challenge of phylogenetic
2313 relationships in Terebelliformia (Annelida: Polychaeta). *Invertebrate Systematics* **27**:186–
2314 238.

- 2315 **Nogueira JMM, Fitzhugh K, Hutchings P, Carrerette O. 2017.** Phylogenetic analysis of the
2316 family Telothelepodidae Nogueira, Fitzhugh & Hutchings, 2013 (Annelida: Polychaeta).
2317 *Marine Biology Research* **13**:671–692.
- 2318 **Nogueira JMM, Fitzhugh K, Rossi MCS. 2010.** A new genus and new species of fan worms
2319 (Polychaeta: Sabellidae) from Atlantic and Pacific Oceans – the formal treatment of taxon
2320 names as explanatory hypotheses. *Zootaxa* **2603**:1–52.
- 2321 **Nola R, Sankey H. 2007.** *Theories of scientific method: an introduction*. Ithaca; McGill-Queen’s
2322 University Press.
- 2323 **Norton JD. 2003.** A material theory of induction. *Philosophy of Science*. **70**:647–670.
- 2324 **Nygren A, Parapar J, Pons J, Meißner K, Bakken T, Kongsrud JA, Oug E, Gaeva D,**
2325 **Sikorski A, Johansen RA, Hutchings PA, Lavesque N, Capa M. 2018.** A mega-cryptic
2326 species complex hidden among one of the most common annelids in the North East
2327 Atlantic. *PLOS One* **13**:e0198356.
- 2328 **Okasha S. 2012.** Emergence, hierarchy and top-down causation in evolutionary biology.
2329 *Interface Focus* **2**:49–54.
- 2330 **Orrhage L. 1966.** Über die anatomie des zentralen nervensystemes der sedentären polychaeten.
2331 *Arkiv för zoologi* **19**:99–133.
- 2332 **Park W. 2017.** *Abduction in context: the conjectural dynamics of scientific reasoning*. Berlin:
2333 Springer-Verlag.
- 2334 **Parry LA, Edgecombe GD, Eibye-Jacobsen D, Vinther J. 2016.** The impact of fossil data on
2335 annelid phylogeny inferred from discrete morphological characters. *Proceedings of the*
2336 *Royal Society B* **283**:20161378. Available from: <http://dx.doi.org/10.1098/rspb.2016.1378>.

- 2337 **Patrick B, Helm C, Bartolomaeus T. 2019.** The anatomy and development of the nervous
- 2338 system in Magelonidae (Annelida) – Insights into the evolution of the annelid brain. *BMC*
- 2339 *Evolutionary Biology* **19**:173. Available from: <https://doi.org/10.1186/s12862-019-1498-9>.
- 2340 **Peirce CS. 1878.** Illustrations of the logic of science. Sixth paper—Deduction, induction, and
- 2341 hypothesis. *Popular Science Monthly* **13**:470–482.
- 2342 **Peirce CS. 1902.** Reasoning. In: Baldwin JA, Rand B, eds. *Dictionary of philosophy and*
- 2343 *psychology: prefatory note. Text, Le-Z. Addenda: indices. I. Greek terms. II. Latin terms.*
- 2344 *III. German terms. IV. French terms. V. Italian terms.* New York: The Macmillan
- 2345 Company 426–428.
- 2346 **Peirce CS. 1931.** *Collected papers of Charles Sanders Peirce, volume 1, principles of*
- 2347 *philosophy.* In: Hartshorne C, Weiss P, Burks A, eds. Cambridge: Harvard University
- 2348 Press.
- 2349 **Peirce CS. 1932.** *Collected papers of Charles Sanders Peirce, volume 2, elements of logic.* In:
- 2350 Hartshorne C, Weiss P, Burks A. eds. Cambridge: Harvard University Press.
- 2351 **Peirce CS. 1933a.** *Collected papers of Charles Sanders Peirce, volume 3, exact logic.* In:
- 2352 Hartshorne C, Weiss P, Burks A, eds. Cambridge: Harvard University Press.
- 2353 **Peirce CS. 1933b.** *Collected papers of Charles Sanders Peirce, volume 4, the simplest*
- 2354 *mathematics.* In: Hartshorne C, Weiss P, Burks A, eds. Cambridge: Harvard University
- 2355 Press.
- 2356 **Peirce CS. 1934.** *Collected papers of Charles Sanders Peirce, volume 5, pragmatism and*
- 2357 *pragmaticism.* In: Hartshorne C, Weiss P, Burks A, eds. Cambridge: Harvard University
- 2358 Press.

- 2359 **Peirce CS. 1935.** *Collected papers of Charles Sanders Peirce, volume 6, scientific metaphysics.*
2360 In: Hartshorne C, Weiss P, Burks A. eds. Cambridge: Harvard University Press.
- 2361 **Peirce CS. 1958a.** *Collected papers of Charles Sanders Peirce, volume 7, science and*
2362 *philosophy.* In: Hartshorne C, Weiss P, Burks A, eds. Cambridge: Harvard University
2363 Press.
- 2364 **Peirce CS. 1958b.** *Collected papers of Charles Sanders Peirce, volume 8, correspondence and*
2365 *bibliography.* In: Burks A, ed. Cambridge: Harvard University Press.
- 2366 **Petrov DA. 2014.** Searching for adaptation in the genome. In: Losos JB, Baum DA, Futuyma
2367 DJ, Hoekstra HE, Lenski RE, Moore AJ, Peichel CL, Schluter D, Whitlock MC, eds. *The*
2368 *Princeton guide to evolution.* Princeton: Princeton University Press 466–474.
- 2369 **Popper KR. 1983.** *Objective knowledge: an evolutionary approach.* New York: Oxford
2370 University Press.
- 2371 **Popper KR. 1992.** *Realism and the aim of science.* New York: Routledge.
- 2372 **Potochnik A. 2017.** *Idealization and the aims of science.* Chicago: University of Chicago Press.
- 2373 **Potochnik A. 2020.** What constitutes an explanation in biology? In: Kampourakis K, Uller T,
2374 eds. *Philosophy of science for biologists.* New York: Cambridge University Press —.
- 2375 **Pouyet F, Aeschbacher S, Thiéry A, Excoffier L. 2018.** Background selection and biased gene
2376 conversion affect more than 95% of the human genome and bias demographic inferences.
2377 *eLife* 7:e36317.
- 2378 **Psillos S. 2002.** Simply the best: a case for abduction. In: Kakas AC, Sadri F, eds.
2379 *Computational logic: logic programming and beyond.* New York: Springer 605–625.
- 2380 **Psillos S. 2007.** *Philosophy of science A-Z.* Edinburgh: University Press.

- 2381 **Psillos S. 2011.** An explorer upon untrodden ground: Peirce on abduction. In: Gabbay D,
2382 Hartmann S, Woods J, eds. *The handbook of the history of logic, volume 10, inductive*
2383 *logic*. Oxford: Elsevier 117–151.
- 2384 **Purschke G. 1997.** Ultrastructure of nuchal organs in polychaetes (Annelida) – New results and
2385 review. *Acta Zoologica* **78**:123–143.
- 2386 **Purschke G. 2005.** Sense organs in polychaetes (Annelida). *Hydrobiologia* **535/536**:53–78.
- 2387 **Radashevsky VI, Malyar VV, Pankova VV, Gambi MC, Giangrande A, Keppel E, Nygren**
2388 **A, Al-Kandari M, Carlton JT. 2020.** Disentangling invasions in the sea: molecular
2389 analysis of a global polychaete species complex (Annelida: Spionidae: *Pseudopolydora*
2390 *paucibranchiata*). *Biological Invasions*. Available from: [https://doi.org/10.1007/s10530-](https://doi.org/10.1007/s10530-020-02346-x)
2391 [020-02346-x](https://doi.org/10.1007/s10530-020-02346-x).
- 2392 **de Regt HW, Leonelli S, Eigner K. 2009.** Focusing on scientific understanding. In: de Regt H,
2393 Leonelli S, Eigner K, eds. *Scientific understanding: philosophical perspectives*. Pittsburgh:
2394 University of Pittsburgh Press 1–17.
- 2395 **Reilly FE. 1970.** *Charles Peirce's theory of scientific method*. New York: Fordham University
2396 Press.
- 2397 **Reiss J, Sprenger J. 2017.** Scientific objectivity. In: *The Stanford encyclopedia of*
2398 *philosophy (Winter 2017 edition)*. Zalta EN, ed. Available from:
2399 <https://plato.stanford.edu/archives/win2017/entries/scientific-objectivity/>.
- 2400 **Rescher N. 1970.** *Scientific explanation*. New York: The Free Press.
- 2401 **Rescher N. 1978.** *Peirce's philosophy of science: critical studies in his theory of induction and*
2402 *scientific method*. Notre Dame: University of Notre Dame Press.

- 2403 **Rindal E, Brower AVZ. 2011.** Do model-based phylogenetic analyses perform better than
2404 parsimony? A test with empirical data. *Cladistics* **27**:331–334.
- 2405 **Ronquist F, van der Mark P, Huelsenbeck JP. 2009.** Bayesian phylogenetic analysis using
2406 MrBayes. In: Lemey P, Salemi M, Vandamme A-M, eds. *The phylogenetic handbook: a*
2407 *practical approach to phylogenetic analysis and hypothesis testing*. New York: Cambridge
2408 University Press 210–266.
- 2409 **Rouse GW. 2001.** *Magelona* Müller, 1858. In: Rouse GW, Pleijel F, authors. *Polychaetes*.
2410 Oxford: Oxford University Press 261–263.
- 2411 **Salas A. 2019.** The natural selection that shapes our genomes. *Forensic Science International:*
2412 *Genetics* **39**:57–60.
- 2413 **Salmon WC. 1967.** *The foundations of scientific inference*. Pittsburgh: University of Pittsburgh
2414 Press.
- 2415 **Salmon WC. 1984a.** *Logic*. Englewood Cliffs: Prentice-Hall, Inc.
- 2416 **Salmon WC. 1984b.** *Scientific explanation and the causal structure of the world*. Princeton:
2417 Princeton University Press.
- 2418 **Salmon WC. 1989.** Four decades of scientific explanation. In: Kitcher P, Salmon WC, eds.
2419 *Scientific explanation. Minnesota studies in the philosophy of science, volume XIII*.
2420 Minneapolis: University of Minnesota Press 3–219.
- 2421 **Salmon WC. 1998.** *Causality and explanation*. New York: Oxford University Press.
- 2422 **Salthe SN. 1985.** *Evolving hierarchical systems: their structure and representation*. New York:
2423 Columbia University Press.

- 2424 **San Martín G, Álvarez-Campos P, Kondo Y, Núñez J, Fernández-Álamo MA, Pleijel F,**
- 2425 **Goetz FE, Nygren A, Osborn K. 2020.** New symbiotic association in marine annelids:
- 2426 ectoparasites of comb jellies. *Zoological Journal of the Linnean Society*. 191:672–694.
- 2427 **Sawyer SA. 1994.** Inferring selection and mutation from DNA sequences: the McDonald–
- 2428 Kreitman test revisited. In: Golding B, ed. *Non-neutral evolution: theories and molecular*
- 2429 *data*. Dordrecht: Springer 77–87.
- 2430 **Schmidt HA, von Haeseler A. 2009.** Phylogenetic inference using maximum likelihood
- 2431 methods. In: Lemey P, Salemi M, Vandamme A-M, eds. *The phylogenetic handbook: a*
- 2432 *practical approach to phylogenetic analysis and hypothesis testing*. New York: Cambridge
- 2433 University Press 181–198.
- 2434 **Schurz G. 2005.** Explanations in science and the logic of why-questions: discussion of the
- 2435 Halonen–Hintikka-approach and alternative proposal. *Synthese* **143**:49–178.
- 2436 **Schurz G. 2008.** Patterns of abduction. *Synthese* **164**:201–234.
- 2437 **Scotland RW, Olmstead RG, Bennett JR. 2003.** Phylogeny reconstruction: the role of
- 2438 morphology. *Systematic Biology* **52**:539–548.
- 2439 **Shimabukuro M, Carrerette O, Alfaro-Lucas JM, Rizzo AE, Halanych, KM, Sumida PYG.**
- 2440 **2019.** Diversity, distribution and phylogeny of Hesionidae (Annelida) colonizing whale
- 2441 falls: new species of *Sirsoe* and connections between ocean basins. *Frontiers in Marine*
- 2442 *Science*. Available from: <https://doi.org/10.3389/fmars.2019.00478>.
- 2443 **Sider T. ~~Intrinsic properties~~. 1996. Philosophical Studies** **83**:1–27.
- 2444 **Sintonen M. 2004.** Reasoning to hypotheses: where do questions come? *Foundations of Science*
- 2445 **9**:249–266.

- 2446 **Sneath PHA, Sokal RR. 1973.** *Numerical taxonomy: the principles and practice of numerical*
2447 *classification*. San Francisco: WH Freeman.
- 2448 **Sober E. 1975.** *Simplicity*. New York: Oxford University Press.
- 2449 **Sober E. 1984.** *The nature of selection: evolutionary theory in philosophical focus*. Cambridge:
2450 MIT Press.
- 2451 **Sober E. 1986.** Explanatory presupposition. *Australasian Journal of Philosophy* **64**:143–149.
- 2452 **Sober E. 1988.** *Reconstructing the past: parsimony, evolution, and inference*. New York: MIT
2453 Press.
- 2454 **Sober E. 1994.** *From a biological point of view: essays in evolutionary biology*. Cambridge:
2455 Cambridge University Press.
- 2456 **Sokal RR, Sneath PHA. 1963.** *Principles of numerical taxonomy*. San Francisco: WH Freeman.
- 2457 **Soltis PS, Soltis DE. 2003.** Applying the bootstrap in phylogeny reconstruction. *Statistical*
2458 *Science* **18**:256–267.
- 2459 **Spencer AR. 2020.** *American pragmatism: an introduction*. New York: Polity.
- 2460 **Stiller J, Tilic E, Rousset V, Pleijel F, Rouse GW. 2020.** Spaghetti to a tree: a robust
2461 phylogeny for Terebelliformia (Annelida) based on transcriptomes, molecular and
2462 morphological data. *Biology* **9**:73–100.
- 2463 **Strahler AN. 1992.** *Understanding science: an introduction to concepts and issues*. Buffalo:
2464 Prometheus Books.
- 2465 **Strawson PF. 1971.** *Individuals: an essay in descriptive metaphysics*. New York: Routledge.
- 2466 **Struck TH, Golombek A, Weigert A, Franke FA, Westheide W, Purschke G, Bleidorn C,**
2467 **Halanych KM. 2015.** The evolution of annelids reveals two adaptive routes to the
2468 interstitial realm. *Current Biology* **25**:1993–1999.

- 2469 **Struck TH, Paul C, Hill N, Hartmann S, Hösel C, Kube M, Lieb B, Meyer A, Tiedemann R,**
- 2470 **Purschke G, Bleidorn C. 2011.** Phylogenomic analyses unravel annelid evolution. *Nature*
- 2471 **471:95–98.**
- 2472 **Sullivan J, Joyce P. 2005.** Model selection in phylogenetics. *Annual Review of Ecology and*
- 2473 *Systematics* **36:445–466.**
- 2474 **Swofford DL. ~~PAUP*~~. 2001.** *Phylogenetic analysis using parsimony (*and other methods).*
- 2475 *Version 4.0b10.* Sunderland: Sinauer Associates.
- 2476 **Swofford DL, Olsen GJ, Waddell PJ, Hillis DM. 1996.** Phylogenetic inference. In: Hillis DM,
- 2477 Moritz C, Mable BK, eds. *Molecular systematics.* Sunderland: Sinauer Associates 407–
- 2478 514.
- 2479 **Thagard P. 1988.** *Computational philosophy of science.* Cambridge: MIT Press.
- 2480 **Thagard P. 2004.** Rationality and science. In: Mele A, Rawlings P, eds. *Handbook of*
- 2481 *rationality.* Oxford: Oxford University Press 363–379.
- 2482 **Tilic E, Sayyari E, Stiller J, Mirarab S, Rouse GW. 2020.** More is needed – Thousands of loci
- 2483 are required to elucidate the relationships of the ‘flowers of the sea’ (Sabellida, Annelida).
- 2484 *Molecular Phylogenetics and Evolution* **151:** —.
- 2485 doi.org/10.1016/j.ympev.2020.106892.
- 2486 **Tucker A. 2004.** *Our knowledge of the past: a philosophy of historiography.* New York:
- 2487 Cambridge University Press.
- 2488 **Tucker A. 2011.** Historical science, over- and underdetermined: a study of Darwin’s inference
- 2489 of origins. *The British Journal for the Philosophy of Science* **62:805–829.**
- 2490 **Turner D. 2007.** *Making prehistory: historical science and the scientific realism debate.* New
- 2491 York: Cambridge University Press.

- 2492 **Uebelacker JM, Jones ML. 1984.** Family Magelonidae. In: Uebelacker JM, Johnson PG, eds.
2493 *Taxonomic guide to the polychaetes of the northern Gulf of Mexico. Final report to the*
2494 *Minerals Management Service, contract 14-12-001-29091.* Barry A. Vittor and Associates,
2495 Mobile, Alabama 7.1–7.29.
- 2496 **Uller T, Laland KN. 2019.** Evolutionary causation. In: Uller T, Laland KN, eds. *Evolutionary*
2497 *causation: biological and philosophical reflections.* Cambridge: The MIT Press 1–12.
- 2498 **Van Fraassen BC. 1990.** *The scientific image.* New York: Clarendon Press.
- 2499 **Vrba E, Eldredge N. 1984.** Individuals, hierarchies and processes: towards a more complete
2500 evolutionary theory. *Paleobiology* **10**:146–171.
- 2501 **Walker SI. 2014.** Top-down causation and the rise of information in the emergence of life.
2502 *Information* **5**:424–39.
- 2503 **Walker SI, Cisneros L, Davies PCW. 2012.** Evolutionary transitions and top-down causation.
2504 *Proceedings of Artificial Life* **13**:283–290.
- 2505 **Walton D. 2004.** *Abductive reasoning.* Tuscaloosa: The University of Alabama Press.
- 2506 **Weigert A, Bleidorn C. 2016.** Current status of annelid phylogeny. *Organisms Diversity &*
2507 *Evolution* **16**:345–362.
- 2508 **Weigert A, Golombek A, Gerth M, Schwarz F, Struck TH, Bleidorn C. 2016.** Evolution of
2509 mitochondrial gene order in Annelida. *Molecular Phylogenetics and Evolution* **94**:196–206.
- 2510 **Weigert A, Helm C, Meyer M, Nickel B, Arendt D, Hausdorf B, Santos SR, Halanych KM,**
2511 **Purschke G, Bleidorn C, Struck TH. 2014.** Illuminating the base of the annelid tree using
2512 transcriptomics. *Molecular Biology and Evolution* **31**:1391–401.

- 2513 **Wilson DP. 1958.** The polychaete *Magelona alleni* n. sp. and a re-assessment of *Magelona*
2514 *cincta* Ehlers. *Journal of the Marine Biological Association of the United Kingdom*
2515 **37**:617–626.
- 2516 **Wilson DP. 1959.** The polychaete *Magelona filiformis* sp. nov. and notes on other species of
2517 *Magelona*. *Journal of the Marine Biological Association of the United Kingdom*
2518 **38**:547–556.
- 2519 **Wilson DP. 1982.** The larval development of three species of *Magelona* (Polychaeta) from
2520 localities near Plymouth. *Journal of the Marine Biological Association of the United*
2521 *Kingdom* **62**:385–401.
- 2522 **Wilson WA. 2017.** The myth of scientific objectivity. Available from:
2523 <https://www.firstthings.com/article/2017/11/the-myth-of-scientific-objectivity>.
- 2524 **Yu L, Boström C, Franzenburg S, Bayer T, Dagan T, Reusch TBH. 2020.** Somatic genetic
2525 drift and multilevel selection in a clonal seagrass. *Nature Ecology & Evolution* **4**:952–962.
- 2526 **Zhang Z, Yu J. 2006.** Evaluation of six methods for estimating synonymous and
2527 nonsynonymous substitution rates. *Genomics Proteomics Bioinformatics* **4**:173–181.

Table 1(on next page)

Forms of questions leading to ab-, de-, and inductive inferences, and respective answers provided by those inferences.

1 **Table 1. Forms of questions leading to ab-, de-, and inductive inferences, and respective**
 2 **answers provided by those inferences.**

Question	Inferential Reaction	Conclusion (= answer)
‘Why is Y the case in contrast to X ?’	Abduction, cf. (1)	Explanatory hypothesis
‘What consequence(s) are expected if hypothesis h is true?’	Deduction, cf. (2)	Potential test evidence
‘Should hypothesis h be accepted as true?’	Induction, cf. (3)	Hypothesis confirmation / disconfirmation

3

4

Table 2(on next page)

Sources from which character data were obtained for this study. Museum abbreviations are explained in the text.

- 1 Table 2. Sources from which character data were obtained for this study. Museum
- 2 abbreviations are explained in the text.

Species	Material
<i>M. alexandrae</i> Magalhães, Bailey-Brock & Watling, 2018	Original description plus photos provided by the authors
<i>M. alleni</i> Wilson, 1958	Holotype: BMNH 1958.5.2.1 Paratypes: BMNH 1958.5.2.2-10
<i>M. annulata</i> Hartmann-Schröder, 1962	Original description
<i>M. anuheone</i> Magalhães, Bailey-Brock & Watling, 2018	Original description plus photos provided by the authors
<i>M. berkeleyi</i> Jones, 1971	Original description
<i>M. californica</i> Hartman, 1944	Original description
<i>M. cepiceps</i> Mortimer & Mackie, 2006	Holotype: NMW.Z.2000.020.0209 paratype: NMW.Z.2000.020.0208
<i>M. cerae</i> Hartman & Reish, 1950	Original description
<i>M. cincta</i> Ehlers, 1908	Holotype (ZMHB 4531)
<i>M. cinthya</i> Magalhães, Bailey-Brock & Watling, 2018	Original description plus photos provided by the authors
<i>M. conversa</i> Mortimer & Mackie, 2003	Holotype: NMW.Z.2000.020.0001 Paratypes: NMW.Z.2000.020.0002-0007
<i>M. cornuta</i> Wesenberg-Lund, 1949	Holotype (ZMUC-POL-969; ZMUC-POL-963)

<i>M. crenulata</i> Bolívar & Lana, 1986	Original description
<i>M. crenulifrons</i> Gallardo, 1968	Paratypes: (ZMUC-POL-1416- 1421)
<i>M. dakini</i> Jones, 1978	Original description
<i>M. debeerei</i> Clarke, Paterson, Florence & Gibbons, 2010	Original description
<i>M. equilamellae</i> Harmelin, 1964	Syntypes: (SMF 4675)
<i>M. falcifera</i> Mortimer & Mackie, 2003	Holotype: NMW.Z.2000.020.0009 Paratypes: NMW.Z.2000.020.0008, NMW.Z.2000.020.0010-0017
<i>M. fauchaldi</i> Shakouri, Mortimer & Dehani, 2017	Holotype: NMW.Z.2015.012.0002a Paratypes: NMW.Z.2015.012.0001-0002; NMW.Z.2010.037.0002-0006
<i>M. filiformis</i> Wilson, 1959	Holotype: BMNH 1959.4.2.1 Paratypes: BMNH 1959.4.2.2-10
<i>M. gemmata</i> Mortimer & Mackie, 2003	Holotype: NMW.Z.2000.020.0019 Paratypes: NMW.Z.2000.020.0018 _f , NMW.Z.2000.020.0020
<i>M. hartmanae</i> Jones, 1978	Original description
<i>M. hobsonae</i> Jones, 1978	Original description
<i>M. johnstoni</i> Fiege, Licher & Mackie, 2000	Cited material from original description from NMW collections
<i>M. jonesi</i> Hartmann-Schröder, 1980	Original description
<i>M. lenticulata</i> Gallardo, 1968	Original description

<i>M. lusitanica</i> Mortimer, Gil & Fiege, 2011	Holotype: SMF 9246/1 Paratypes: MB29-000176-000181; NMW.Z.2010.010.0001-0006; SMF 9245/1
<i>M. magnahamata</i> Aguado & San Martín, 2004	Original description
<i>M. mahensis</i> Mortimer & Mackie, 2006	Holotype: NMW.Z.2000.020.0188 Paratypes: NMW.Z.2000.020.0176-0187; NMW.Z.2000.020.0189-0193
<i>M. marianae</i> Hernández-Alcántara & Solís-Weiss, 2000	Original description
<i>M. minuta</i> Eliason, 1962	Holotype: NHMG Polych. 11491 Additional material: USNM 52510
<i>M. mirabilis</i> (Johnston, 1865)	Cited material from original description from NMW collections
<i>M. montera</i> Mortimer, Cassà, Martin & Gil, 2012	Holotype: MNHN A895
<i>M. nonatoi</i> Bolívar & Lana, 1986	Original description
<i>M. obockensis</i> Gravier, 1905	Syntypes: MNHN Type 1357
<i>M. pacifica</i> Monro, 1933	Syntypes: BMNH Type 1933.7.10.65/70
<i>M. papillicornis</i> F. Müller, 1858	Redescription [242]
<i>M. parochilis</i> Zhou & Mortimer, 2013	Holotype: ECSFRI100532 Paratypes: ECSFRI100533-535; NMW.Z.2012.033.0001

<i>M. paulolanai</i> Magalhães, Bailey-Brock & Watling, 2018	Original description plus photos provided by the authors
<i>M. pettiboneae</i> Jones, 1963	Original description
<i>M. phyllisae</i> Jones, 1963	Original description
<i>M. pitelkai</i> Hartman, 1944	Original description
<i>M. polydentata</i> Jones, 1963	Original description
<i>M. posterelongata</i> Bolívar & Lana, 1986	Original description
<i>M. pulchella</i> Mohammad, 1970	Holotype: BMNH 1969.391
<i>M. pygmaea</i> Nateewathana & Hylleberg, 1991	Paratype: PMBC 4220 Additional material: PMBC 4234; PMBC 4227; PMBC 4241
<i>M. riojai</i> Jones, 1963	Original description
<i>M. sacculata</i> Hartman, 1961	Holotype: NHMLA: LACM-AHF POLY 596
<i>M. sinbadi</i> Mortimer, Cassà, Martin & Gil, 2012	Holotype: NMW.Z.2010.037.0001
<i>M. spinifera</i> (Hernández-Alcántara & Solís-Weiss, 2000)	Original description
<i>M. symmetrica</i> Mortimer & Mackie, 2006	Holotype: NMW.Z.2000.020.0175
<i>M. tehuanaensis</i> Hernández-Alcántara & Solís-Weiss, 2000	Original description
<i>M. tinae</i> Nateewathana & Hylleberg,	Paratypes: PMBC 3180; PMBC 4251; PMBC

1991	4253; PMBC 4254
<i>M. uebelackerae</i> (Hernández-Alcántara & Solís-Weiss, 2000)	Original description
<i>M. variolamellata</i> Bolívar & Lana, 1986	Original description
<i>M. wilsoni</i> Glémarec, 1966	Paratypes: MNHN Poly Type 1415
<i>Magelona</i> sp. A	Uebelacker & Jones, 1984
<i>Magelona</i> sp. B	Uebelacker & Jones, 1984
<i>Magelona</i> sp. C	Uebelacker & Jones, 1984
<i>Magelona</i> sp. D	Uebelacker & Jones, 1984
<i>Magelona</i> sp. E	Uebelacker & Jones, 1984
<i>Magelona</i> sp. F	Uebelacker & Jones, 1984
<i>Magelona</i> sp. G	Uebelacker & Jones, 1984
<i>Magelona</i> sp. H	Uebelacker & Jones, 1984
<i>Magelona</i> sp. I	Uebelacker & Jones, 1984
<i>Magelona</i> sp. J	Uebelacker & Jones, 1984
<i>Magelona</i> sp. K	Uebelacker & Jones, 1984
<i>Magelona</i> sp. L	Uebelacker & Jones, 1984
<i>Magelona</i> sp. 4 (West Africa) Mortimer et al. (in prep)	ZMBN; AC-NMW
<i>Magelona</i> sp. 4B (West Africa) Mortimer et al. (in prep)	ZMBN; AC-NMW
<i>Magelona</i> sp. 4C (West Africa)	ZMBN; AC-NMW

Mortimer et al. (in prep)	
<i>Magelona</i> sp. 5A (West Africa)	ZMBN; AC-NMW
Mortimer et al. (in prep)	
<i>Magelona</i> sp. 5B (West Africa)	ZMBN; AC-NMW
Mortimer et al. (in prep)	
<i>O. bizkaiensis</i> Aguirrezabalaga, Ceberio & Fiege, 2001	Holotype: MNCN 16.01/6887 Paratype: SMF 10025
<i>Octomagelona</i> sp. (West Africa)	AC-NMW

Table 3(on next page)

Characters, as observation statements represented by subject-predicate relations, not ‘characters’ and ‘states’ (Fitzhugh, 2006c, 2008c), used in this study.

Table 3. Characters, as observation statements represented by subject-predicate relations, not ‘characters’ and ‘states’ (Fitzhugh, 2006c, 2008c), used in this study.

-
1. Shovel-shape prostomium: (0) absent; (1) present.
 2. Nuchal organs: (0) absent; (1) present.
 3. Prostomium distal margin: (0) smooth; (1) crenulate; (2) medially indented.
 4. Prostomium distal shape: (0) triangular; (1) rounded; (2) straight.
 5. Prostomial horns (including rudimentary): (0) absent; (1) present.
 6. Shape of prostomial horns [cf. character 5(1)]: (0) rudimentary; (1) distinct.
 7. Prostomium dimensions: (0) longer than wide; (1) as wide as long; (2) wider than long.
 8. Prostomial ridges: (0) absent; (1) present.
 9. Number of prostomial ridges [cf. character 8(1)]: (0) a pair; (1) two pairs.
 10. Palp origin: (0) dorsal; (1) ventral.
 11. Palp surface: (0) non-papillate; (1) papillate.
 12. Number of rows of proximal palp papillae [cf. character 11(1)]: (0) 2; (1) 4–8; (2) 10–14.
 13. Number of rows of distal palp papillae [cf. character 11(1)]: (0) 2; (1) 4; (2) more.
 14. Burrowing organ: (0) absent; (1) present.
 15. Body regionation: (0) non-magelonid; (1) magelonid-like.
 16. Number of anterior body region chaetigers [cf. character 15(1)]: (0) 8; (1) 9.
 17. Notopodial postchaetal lamellae: (0) absent; (1) present.
 18. Development of notopodia along thorax: (0) all similar; (1) different in some chaetigers.
 19. Neuropodial postchaetal lamellae: (0) absent; (1) present.
 20. Development of neuropodia along thorax: (0) all similar; (1) different in some chaetigers.
 21. Chaetigers 1–7 position of notopodial lamellae relative to notochaetae: (0) postchaetal; (1)

- 24 subchaetal.
- 25 22. Chaetigers 1–7 notopodial lamellae shape: (0) filiform; (1) foliaceous.
- 26 23. Chaetigers 1–7 margins of foliaceous notopodial lamellae [cf. character 22(1)]: (0) smooth;
- 27 (1) crenulate; (2) bilobed.
- 28 24. Chaetigers 1–7 dorsal superior lobes: (0) absent; (1) present.
- 29 25. Superior dorsal lobes [cf. character 24(1)]: (0) absent on some chaetigers between 1–7; (1)
- 30 present on chaetigers 1–7.
- 31 26. Chaetigers 1–7 neuropodial lamellae position relative to notochaetae (0) postchaetal;
- 32 (1) subchaetal; (2) prechaetal.
- 33 27. Chaetigers 1–7 neuropodial subchaetal lamellae position relative to notochaetae [cf.
- 34 character 26(1)]: (0) same position along thorax; (1) varying in position along thorax.
- 35 28. Chaetigers 1–7 neuropodial lamellae shape: (0) filiform; (1) foliaceous.
- 36 29. Chaetigers 1–7 neuropodial filiform lamellae shape distal ends [cf. character 28(0)]: (0)
- 37 pointed; (1) distally expanded/scoop shaped.
- 38 30. Chaetigers 1–7 lengths of noto- and neuropodial lamellae: (0) equivalent; (1) notopodial
- 39 longer; (2) neuropodial longer.
- 40 31. Chaetigers 1–8 limbations of capillary chaetae: (0) unilimbate; (1) bilimbate.
- 41 32. Chaetigers 1–8 margins of bilimbate capillary chaetae [cf. character 31(1)]: (0) smooth;
- 42 (1) irregular blade.
- 43 33. Chaetigers 1–8 lengths of noto- and neuropodial capillary chaetae: (0) equivalent; (1)
- 44 notopodial longer; (2) neuropodial longer.
- 45 34. Chaetiger 8 notopodial lamellae position relative to notochaetae: (0) postchaetal;
- 46 (1) subchaetal.

- 47 35. Chaetiger 8 notopodial lamellae shape: (0) filiform; (1) foliaceous.
- 48 36. Chaetiger 8 margins of foliaceous notopodial lamellae [cf. character 35(1)]: (0) smooth; (1)
- 49 crenulate; (2) bilobed.
- 50 37. Chaetiger 8 superior dorsal lobes: (0) absent; (1) present.
- 51 38. Chaetiger 8 neuropodial lamellae position: (0) postchaetal; (1) subchaetal; (2) prechaetal.
- 52 39. Chaetiger 8 neuropodia: (0) without additional postchaetal expansion; (1) with an additional
- 53 postchaetal expansion.
- 54 40. Chaetiger 8 neuropodial lamellae shape: (0) filiform; (1) foliaceous.
- 55 41. Chaetiger 9 notopodial lamellae positions: (0) postchaetal; (1) subchaetal.
- 56 42. Chaetiger 9 notopodial lamellae height [cf. character 41(0)]: (0) low; (1) elongate.
- 57 43. Chaetiger 9 notopodial lamellae shape: (0) filiform; (1) foliaceous.
- 58 44. Chaetiger 9 notopodial foliaceous lamellae margins [cf. 43(1)]: (0) smooth; (1) crenulate; (2)
- 59 bilobed.
- 60 45. Chaetiger 9 superior dorsal lobes: (0) absent; (1) present.
- 61 46. Chaetiger 9 neuropodial lamellae position: (0) postchaetal; (1) subchaetal; (2) prechaetal.
- 62 47. Chaetiger 9 neuropodial lamellae height [cf. character 46(0)]: (0) low; (1) elongate.
- 63 48. Chaetiger 9 neuropodial lamellae shape: (0) filiform; (1) foliaceous.
- 64 49. Chaetiger 9 neuropodia: (0) without additional postchaetal expansion; (1) with additional
- 65 postchaetal expansion.
- 66 50. Lengths of chaetiger 9 fascicles relative to chaetigers 1–8 fascicles: (0) same length as
- 67 chaetae in chaetigers 1–8; (1) shorter than chaetae in chaetigers 1–8; (2) longer than
- 68 chaetae in chaetigers 1–8.
- 69 51. Distal ends of chaetae in chaetiger 9: (0) gently tapered, similar to chaetigers 1–8; (1)

- mucronate; (2) pennoned.
52. Shape of thoracic interparapodial margins: (0) straight; (1) rounded, bulbous.
53. Dorso-lateral grooves, anterior chaetigers: (0) absent; (1) present.
54. Dimensions of post-chaetiger 9 segments: (0) as long as wide; (1) longer than wide.
55. Post-chaetiger 9 lamellae shape: (0) without basal constriction; (1) with basal constriction.
56. Post-chaetiger 9 dorsal and ventral medial lobes: (0) absent; (1) present.
57. Post-chaetiger 9 dorsal and ventral medial lobes sizes [cf. character 56(1) partim]: (0) smaller than hooks; (1) longer than hooks.
58. Post-chaetiger 9 postchaetal expansion behind chaetal rows: (0) absent; (1) present.
59. Relative sizes of post-chaetiger 9 noto- and neuropodial postchaetal lamellae: (0) same size; (1) different size.
60. Post-chaetiger 9 lateral pouches: (0) absent; (1) present.
61. Post-chaetiger 9 lateral pouch arrangement [cf. character 60(1)]: (0) paired; (1) unpaired; (2) both paired and unpaired.
62. Direction of lateral pouch openings [cf. character 60(1)]: (0) posteriorly; (1) posteriorly and anteriorly.
63. Post-chaetiger 9 posteriorly open lateral pouch arrangement [cf. character 62(0)]: (0) on consecutive segments; (1) on alternating segments; (2) on both consecutive and alternating segments.
64. Post-chaetiger 9 lateral pouches distribution [cf. character 60(1)]: (0) throughout most abdominal chaetigers; (1) median and posterior abdominal chaetigers only; (2) posterior abdomen only.
65. Margins of posteriorly-open pouches [cf. character 62(0)]: (0) smooth; (1) medially split.

- 93 66. Post-chaetiger 9 noto- and neuropodial hooded hooks: (0) absent; (1) present.
 - 94 67. Post-chaetiger 9 noto- and neuropodial hooded hooks dentition [cf. character 66(1)]: (0)
 - 95 bidentate; (1) tridentate; (2) polydentate.
 - 96 68. Post-chaetiger 9 hooded hooks adjacent to notopodial subchaetal or neuropodial suprachaetal
 - 97 lamellae [cf. character 66(1)]: (0) same size; (1) smaller than rest.
 - 98 69. Post-chaetiger 9 enlarged hooded hooks: (0) absent; (1) present.
 - 99 70. Post-chaetiger 9 enlarged hooded hooks distal ends [cf. character 69(1)]: (0)
 - 100 recurved; (1) spine-like; (2) enlarged normal hook.
 - 101 71. Post-chaetiger 9 number of hooded hooks per ramus [cf. character 66(1)]: (0) 8 or more; (1)
 - 102 less than 8.
 - 103 72. Post-chaetiger 9 arrangement of hooded hooks [cf. character 66(1)]: (0) vis-à-vis; (1) vis-à-
 - 104 dos.
 - 105 73. Neuropodial sabre chaetae: (0) absent; (1) present.
 - 106 74. Post-chaetiger 9 noto- and neuropodial internal ‘aciculae’: (0) absent; (1) present.
 - 107 75. Pigmentation in posterior thorax: (0) absent; (1) present.
 - 108 76. Pigmentation pattern in posterior thorax [cf. character 75(1): (0) discreet band; (1) dispersed,
 - 109 not forming band.
 - 110 77. Granular bodies on surfaces of segments: (0) absent; (1) present.
 - 111 78. Granular bodies within segments as abdominal interparapodial patches: (0) absent; (1)
 - 112 present.
 - 113 79. Tube construction: (0) absent; (1) present.
-

114

Classes of hypotheses in systematics. Relations between the eight classes (A–H) of explanatory hypotheses commonly inferred in biological systematics (adapted from Fitzhugh, 2013: fig. 1, 2016b: fig. 1; based on Hennig, 1966: fig. 6).




Figure 2

Observation statements, why-questions, and data matrix.

A data matrix used to infer phylogenetic hypotheses implies the why-questions to which inferences lead to answers as phylogenetic hypotheses, implied by cladograms (Fig. 1). (A) Observation statements, as subject-predicate relations, and contrastive why-questions implied by a data matrix. (B) The contrastive form of why-questions is maintained in a data matrix through the inclusion of outgroups, distinguishing foil and fact. Modified from Fitzhugh (2016b: fig. 3).

A.

		subjects				
		↑				
		1	2	3	4	5
'outgroup'	<i>a-us</i>	0	0	0	0	0
	<i>b-us</i>	1	1	0	0	1
	<i>c-us</i>	1	1	1	1	2
	<i>d-us</i>	1	1	1	1	2

Why-question:
 "Why do individuals to which species hypotheses *c-us* and *d-us* refer have character 4(1), in contrast to 4(0) as observed among members of other species?"

2(1): Subject-predicate relations among individuals to which species hypothesis *d-us* applies

B.

			1	2	3	4	5
foil:	'outgroup'	<i>a-us</i>	0	0	0	0	0
		<i>b-us</i>	1	1	0	0	1
facts:		<i>c-us</i>	1	1	1	1	2
		<i>d-us</i>	1	1	1	1	2

Figure 3

Explanatory hypotheses implied by cladograms.

Cladograms imply two general classes of explanatory hypotheses (see also Fig. 1) – specific and phylogenetic – causally accounting for differentially shared characters. Inferences of specific hypotheses are distinct from phylogenetic inferences (cf. Fitzhugh, 2005b, 2009, 2013, 2015), but both typically offer vague explanations with largely unspecified causes. Modified from Fitzhugh (2016b: fig. 2).




Figure 4

Magelonid prostomial characters.

(A) smooth distal margin of *Magelona obockensis* (BMNH:ZB.1971.54, photo by J. Turner); (B) distinctly crenulate distal margin of *M. crenulifrons* from Hong Kong (NMW.Z.2007.033.0001); (C) minutely crenulate margin of *M. wilsoni* from the Gulf of Lions (NMW.Z.2010.010.0008); (D) medially indented distal margin of *M. symmetrica* (holotype); (E) triangular distal margin of *M. simadi* (holotype); (F) rounded distal margin of *M. johnstoni* from Berwick-upon-Tweed (NMW.Z.2013.037.0008); (G–H) straight distal margins (often termed ‘rudimentary horns’) of *M. alleni* from Morocco (ZMBN, MIWA project) and *M. equilamellae* from the Ebro Delta, Catalonia. (A–E, G, H, dorsal views; F, ventral view; A, E, G, stained with methyl green; H, stained with Rose Bengal). BO = burrowing organ, IM = indented margin, IR = inner ridges, OR = outer ridge, PB = palp base, PH = prostomial horns, Pp = palp, Pr = prostomium, RH = rudimentary horns.




Figure 5

Magelonid anterior end and palp characters.

(A) prostomium and first chaetiger of *Magelona pacifica* (syntype, dorsal view); (B) prostomium and first three chaetigers of *M. montera* (holotype, dorso-lateral view, stained with methyl green; Photo by J. Turner); (C) prostomium and first three chaetigers of *M. filiformis* (paratype, BMNH 1959.4.2.6–10, dorso-lateral view); (D) prostomium of *M. posterelongata* (dorsal view); (E) prostomium and first chaetiger of an undescribed species from Nigeria (dorsal view, ZMBN, MIWA project); (F) mid-palp region of *M. johnstoni* from Berwick-upon-Tweed (NMW.Z.2013.037.0008c); (G) prostomium and anterior thorax, showing ventrally inserted papillate palps, and partially everted burrowing organ (NMW.Z.2013.037.0008b); (H) distal palp end of *M. areni* (SEM by K. Mills). Ach = achaetous first segment, BO = burrowing organ, IR = inner ridges, OR = outer ridge, PH = prostomial horns, Pp = palp, Pr = prostomium, RH = rudimentary horns. Numbers indicate chaetiger.




Figure 6

Octomagelona.

Octomagelona: (A, C-F) anterior of an undescribed species from West Africa (dorsal, ventral, antero-dorsal, postero-dorsal and lateral views respectively) (B) *Octomagelona bizkaiensis* (holotype, dorsal view). Ab = abdomen, Ach = achaetous first segment, Pr = prostomium, Th = thorax.




Figure 7

Development of magelonid thoracic parapodia.

(A) left-hand parapodia of chaetigers 1, 5, 8 and 9 of *Magelona minuta* SNM 52510, anterior views); (B) parapodia of chaetigers 1, 6, 8 and 9 of *M. equilamella* (syntype, anterior views); (C) parapodia of chaetigers 2, 3, 4 and 9 of *M. obockensis* (syntype, anterior views); (D) left-hand parapodia of chaetigers 1, 5, 8 and 9 of *M. conversa* (Iran, NMW.Z.2010.037.0007, anterior views).




Figure 8

Magelonid thoracic parapodia.

(A) thoracic postchaetal notopodial lamellae of *Magelona mirabilis* (anterior view, SEM by K. Mills); (B) chaetigers 7–10 of *M. alleni* from Morocco (ZMBN, MIWA project, lateral view, stained with methyl green); (C) thoracic prechaetal neuropodial lamellae of *M. mirabilis* (posterior view, SEM by K. Mills); (D) subchaetal notopodial lamellae of *M. riojai*; (E) left-hand parapodia of chaetiger 8 of *M. conversa* (Iran, NMW.Z.2010.037.0007, anterior view); (F) right-hand parapodia of chaetigers 2–4 of *M. equilamellae* (Ebro Delta, Catalonia, lateral view, SEM by K. Mills); (G) left-hand parapodia of chaetigers 7–9 of same specimen (SEM by K. Mills). Noto = notopodial, Neuro = neuropodia, Pre = prechaetal, Post = postchaetal, SDL = superior dorsal lobe. Numbers indicate chaetiger.











Figure 9

Magelonid thoracic parapodia.








(A) right-hand parapodia of chaetiger 4 of *Magelona fauchaldi* (paratype, NMW.Z.2015.012.0002b, antero-lateral view); (B) right-hand notopodia of *M. equilamellae* (Ebro Delta, Catalonia, anterior view; SEM by K. Mills); (C) foliaceous thoracic lamellae of *M. sp. 2* (Brasil, [16], anterior view); (D–E) parapodia of chaetigers 8 and 4 of *M. montera* respectively (holotype, anterior views); (F) right-hand parapodia of chaetiger 2 of *M. obocken* (holotype, anterior view); (G) right-hand parapodia of chaetiger 8 of *M. symmetri* (holotype, anterior view); (H) distally expanded and scoop-shaped neuropodial lamellae of chaetigers 1–3 of *Magelona* sp. cf. *M. cincta* (Iran, NMW.Z.2015.012.0004, posterolateral view, stained with methyl green); (I) notopodia of chaetigers 9 and 10 of *M. johnstoni* from Berwick-upon-Tweed (anterior view). Noto = notopodia, Neuro = neuropodia, Pre = prechaetal, Post = postchaetal, SDL = superior dorsal lobe.



Vac-High PC-Std. 10 kV x 1500 20 µm 000067



Vac-High PC-Std. 15 kV x 340 50 µm 000024



Vac-High PC-Std. 10 kV x 400 50 µm 000066

Figure 10

Magelonid thoracic chaetae.

(A) first five chaetigers of *Magelona equilamellae* (Ebro Delta, Catalonia, dorsal view; SEM by K. Mills) ~~showing thoracic capillary chaetae in noto- and neuropodia~~; (B) right-hand notopodia of chaetiger 5 (anterior view) of *M. equilamellae* (Ebro Delta, Catalonia; SEM by K. Mills); (C) unilimbate capillary chaetae of *M. minut* (SNM 52510); (D) bilimbate capillary chaetae with irregular blades from chaetiger 4 of *M. riojai*; (E) bilimbate capillary chaeta with irregular blade from chaetiger 2 of *M. variolamellata*; (F–G) mucronate chaetae from chaetiger 9 of *M. riojai*; (H) the same from *M. johnstoni* from Berwick-upon-Tweed (NMW.Z.2013.037.0011b); (I) pennoned chaetae from chaetiger 9 of *M. pitelkai*, *M. hobsonae*, *M. hartmanae* (respectively), modified from Jones (1978).




Figure 11

Magelonid thoracic region.

(A) characteristically bulbous and rounded thoracic chaetigers of *Magelona alieni* (holotype, dorsal view); (B) anterior of *M. johnstoni* (NMW.Z. 2013.037.0001, dorsal view, photo by ASYM) showing straight thoracic interparapodial margins; (C) anterior of *M. variolamellata* (dorsal view, scale 1 mm) showing oblique lateral slits; (D) anterior of *M. crenulifrons* (Hong Kong, ventral view) showing V shaped ventral marking in the mid thoracic region and methyl green staining pattern; (E) anterior of *M. pulchella* (holotype, ventral view) showing X shaped ventral marking in the mid thoracic region, stained with methyl green; (F) thoracic ventral pads/swellings of *M. mirabilis* in the mid to posterior thorax (NMW.Z.2013.037.0020, ventral view, photo by A.S.Y. Mackie); (G) posterior thorax of *M. obockensis* (syntype, ventral view) showing ventral swellings/pads. Ach = achaetous first segment, BO = burrowing organ, DS = dorsal slits, Pp = palp, Pr = prostomium, Th = thorax, V = ventral V-shaped marking, VS = ventral swellings, X = ventral X-shaped marking.




Figure 12

Magelonid abdominal chaetigers.

(A) wider than long abdominal chaetigers of *Magelona posterolongica* (lateral view, scale 1 mm); (B) right-hand parapodia of chaetigers 10 and 11 of *M. johnsoni* (NMW.Z.2013.037.0011b, anterior view), ~~showing anteriorly open pouch~~; (C) left-hand parapodia of chaetiger 10 of *M. crenulifrons* (NMW.Z.2010.037.0034b, anterior view); (D) right-hand parapodia of chaetigers 9–11 of *M. equilamellae* (Ebro Delta, Catalonia, lateral view; SEM by K. Mills); (E) abdominal parapodium of *M. sp.* 1 of Brasil (2003) (anterior view); (F) right-hand parapodia of abdominal chaetiger of *M. sp.* 2 of Uebelacker & Jones (1984) (~~modified from that publication~~); (G) left-hand parapodia of chaetiger 11 of *M. allenii* (Morocco, ZMBN, MIWA project) ~~showing sub equal lateral lamellae~~; (H) right-hand parapodia of an abdominal chaetiger of *M. sp.* 2 of Brasil (2003); (I) thoracic-abdominal junction of *M. filiformis* (paratype, BMNH 1959.4.2.6–10, ventro-lateral view). AC = acicular chaetae, AH = abdominal hooks, DF = dorsal flap, DML = dorsal medial lobe, LO = lateral organ, Neuro = neuropodia, Noto = notopodia, PE = postchaetal extension of the lamellae, PM, pouch membrane, SAH = small abdominal hook, US = unidentate spine, VF = ventral flap, VML = ventral medial lobe. Numbers indicate chaetiger.




Figure 13

Magelonid lateral abdominal pouches.

(A) schematic diagram showing a 'magelonid' with paired anteriorly open pouches in the anterior abdomen, followed by unpaired posteriorly open pouches in the mid and posterior abdomen, on alternate chaetigers; (B) schematic diagram of a 'magelonid' with unpaired posteriorly open pouches, on alternate chaetigers; (C) schematic diagram of a 'magelonid' with paired posteriorly open pouches, on consecutive chaetigers; (D) transverse section through the body and an anteriorly opening pouch situated between chaetigers 10 and 11 (posterior half of pouch and parapodia of chaetiger 11 visible) of *Magelona johnstoni* (NMW.Z.2013.037.0010c, anterior view); (E) left-hand anteriorly opening pouch between chaetigers 10 and 11 of *M. johnstoni* (NMW.Z.2013.037.0011b, lateral view); (F) the same from another specimen, ventrolateral view (NMW.Z.2013.037.0008c); (G) posteriorly open pouch of the same specimen (ventrolateral view); (H) posteriorly open pouches on consecutive segments of *M. pacifica* (syntype, lateral view), pouches showing distinct medial splits. DF = dorsal flap, DML = dorsal medial lobe, LO = lateral organ, Neuro = neuropodia, Noto = notopodia, PM = pouch membrane, PO pouch = posteriorly opening pouch, VF = ventral flap. Numbers indicate chaetiger.




Figure 14

Magelonid abdominal hooded hooks.

(A) transverse section through the body of *Magelona johnstoni* between chaetigers 10 and 11 (posterior half of pouch and parapodia of chaetiger 11 visible) (NMW.Z.2013.037.0010c, anterior view); (B) row of hooded hooks of *M. equilamellae* arranged vis-à-vis (Ebro Delta, Catalonia); (C) bidentate hooded hooks from chaetiger 29 of *M. neta* (NMW.Z.1991.075.1584a, lateral view); (D) bidentate hooded hook from *M. papillicornis* (lateral view; scale = 0.01 mm); (E) tridentate hooded hook from *M. alleni*; (F–H) quadridentate, pentadentate and hexodont hooded hooks from *M. fauchardii* (paratypes); (I) small hooded hook adjacent to the lateral lamellae of *M. sp.* 370r Brasil (2003) (anterior view); (J) bidentate hooded hooks of *M. papillicornis*, arranged vis-à-vis; (K) unidentate enlarged hook of *M. sp.* 370r Brasil [16]. (C, E–H, hoods broken. Photos B, C, E by K. Mills). Neuro = neuropodia, Noto = notopodia, SAH = small abdominal hook.











Figure 15

Magelonid pigmentation.

(A) anterior end of a live *Magelona allenii* (dorsal view) from Plymouth, showing thoracic pigment band on posterior thorax (photo by A.S.Y. Mackie); (B) anterior end (dorsal view) of *M. sp. 5* from West Africa (ZMBN, MIWA project), ~~showing distinct stripy pigmentation characteristic of the species~~; (C) pigment band of posterior thorax of *M. equilamellae* from Ebro Delta, Catalonia (photo by K. Mills). Pp = palp, Pr = prostomium.



Figure 16

Magelonid pigmentation, granular bodies, tube, pygidial cirri.

(A) anterior of *Magelona minuta* (NMW.Z.2005.014.0111, dorsal view. Photo by K. Mills) showing distinct transverse thoracic stripes; (B) anterior of *Magelona* sp. cf. *M. cincta* from Iran (NMW.Z.2015.012.0004, dorsal view) showing distinct circular regions of granular bodies adjacent to the parapodia; (C) same specimen showing abdominal interparapodial patches and granular bodies along the mid-ventral line of the abdomen (ventral view); (D) granular bodies of the dorsal surface of the thoracic region of *M. papillicornis*; (E) distinct multi-layered tube covered in sand of *M. allenii* from the Irish Sea (NMW.Z.1969.104.1094. Photo by K. Mills); (F) digitiform pygidial cirri either side of the pygidium (dorsal view); (G) posterior of *M. minuta* from the Outer Bristol Channel (NMW.Z.2003.047.5939, ventral view. Photo by K. Mills); (H) posterior of *M. allenii* (paratype, ventro-lateral view). (A–C, G, H, stained with methyl green; E, stained with Rose Bengal). A = anus, Ab = abdomen, BO = burrowing organ, GB = granular bodies, PC = pygidial cirri, Pp = palp, Pr = prostomium, Th = thorax.




Figure 17

Phylogenetic results. Strict consensus tree for 2,417,600 cladograms produced from data matrix in Table S1.

Note that *Magelona* is paraphyletic relative to *Octomagelona*. Letters **a–u** indicate clades for which some character transformation series are presented in the **Results**.




Figure 18

Phylogenetic results.

Fig 46 of the strict consensus tree for nine cladograms presented by Brasil (2003) of Magelonidae phylogenetic hypotheses. Clades referred to in that study are indicated by the Roman numerals.




Table S1. Character matrix of Magelonidae taxa and characters. Outgroups are *Phyllochaetopterus limicolus*, *Spio filicornis*,

Laonice cirrata, *Prionospio lighti*, and *P. ehlersi*. ‘?’ = unknown; ‘-’ = inapplicable; ‘*’ = polymorphism, 0/1, in *Magelona* sp. H.

	1 0	2 0	3 0	4 0	5 0	6 0	7 0
<i>Phyllochaetopterus limicolus</i>	01---20-0	0--00-0-0-	-----	0-1-----	-----0	0000-----0	-----0-000-001
<i>Spio filicornis</i>	01---00-0	0--00-1-1-	0100-0-1-0	0-10100001	0010000100	000000-000	-----10-0- 11100-001
<i>Laonice cirrata</i>	01---20-0	0--00-1-1-	0100-0-1-0	0-00100001	0110001100	000000-001	0001012-0- 01100-00?
<i>Prionospio lighti</i>	01---00-0	0--00-1-1-	0100-0-1-1	0-10100001	0110001100	000000-000	-----12-0- 01100-00?
<i>P. ehlersi</i>	01---00-0	0--00-1-1-	0100-0-1-1	0-00100001	0110001100	000000-001	0001012-0- 01100-00?
<i>Magelona alexandrae</i>	1000110111	1101111111	0111110001	10?0111110	111002101?	0001110101	000111100- 10000-110
<i>M. alleni</i>	1002101111	1221111010	10-0-10001	10010-0100	110-011000	010000-010	-----1100- 000010111
<i>M. annulata</i>	1000111111	1??1111010	1100-1?001	10011001?0	1110011010	0000111000	-----101?? 100?0-??0
<i>M. anuheone</i>	1000110111	1111111110	0101110001	10?0101100	111001100?	0001111101	0???1110- 10010-1?0
<i>M. berkeleyi</i>	1000112111	1111111111	0101111001	1120101210	0110021010	0100110100	-----1110- 000?0-?00
<i>M. californica</i>	10020-1101	1??1111010	00-0-0-000	??000-0000	010-001000	000000-000	-----1010- 10000-110
<i>M. cepiceps</i>	1002101111	1111111011	0101111001	??00101110	0110111010	0100110100	-----1100- 00000-110
<i>M. cerae</i>	10001111?1	1??1111111	00-111?001	?0200-1000	1100001000	000000-000	-----1000- 11000-110
<i>M. cincta</i>	1002101111	1111111111	10-0-11012	10?10-0000	110-001001	010000-001	101001100- 000010111
<i>M. cinthya</i>	1000110111	1101111111	0101110001	10?0101210	111002101?	0001110101	00?1?1110- 10010-110
<i>M. conversa</i>	10010-0111	1221111111	010102-001	10?0101211	0010021011	1000010001	211001000- 01010-010
<i>M. cornuta</i>	1010111111	11?1111111	0101110001	10?0101210	111002101?	0000110101	000111100- 00010-110
<i>M. crenulata</i>	10010-01?1	11?1111111	012112-001	1000121210	0011000101	1000010101	2110?1100- 10000-?10
<i>M. crenulifrons</i>	1010111111	1??1111111	0101110001	1000101210	1110021010	0000111101	101001000- 00010-110
<i>M. dakini</i>	1002100111	1101111011	00-0-11001	1000100210	0110011010	0000111101	1012?1110- 00010-?10
<i>M. debeerei</i>	10010-0111	11?1111111	0101010001	??00101211	0010000101	100000-001	?110?1100- 01000-??0
<i>M. equilamellae</i>	1002102101	1111111011	10-0-11000	10010-0000	010-001000	010000-000	-----1100- 000010111
<i>M. falcifera</i>	10020-1101	1111111010	00-0-0-000	??000-0000	010-00100?	0000010000	-----10110 10000-110
<i>M. fauchaldi</i>	10020-1101	1001111010	00-0-0-000	10000-0000	010-001000	000000-000	-----1200- 10000-110
<i>M. filiformis</i>	1002100111	1111111111	00-1110001	10000-1110	110-011010	0000111001	101201110- 00000-110
<i>M. gemmata</i>	1000110111	1101111111	0101110001	???0101110	1110021010	0000110101	000101100- 10010-110
<i>M. hartmanae</i>	1002100111	1101111111	00-112-001	10000-1200	110-021010	2000111001	1011?1110- 10000-?00
<i>M. hobsonae</i>	1002102111	1111111111	00-1110000	10010-1210	110-011010	2000111000	-----1110- 00000-?00
<i>M. johnstoni</i>	10010-0111	1111111111	0111010001	1?01101210	0010000101	1000010001	211001100- 01010-010
<i>M. jonesi</i>	1022101111	12?1111011	0100-0-1-0	1010100110	0110001100	0100110000	-----1010- 01000-?00
<i>M. lenticulata</i>	1010112111	11?1111111	010111?0?1	10211010?1	0110121010	0100110000	-----1100- 000?0-1?0
<i>M. lusitanica</i>	1000110111	1101111111	0101110001	??00101210	1110021010	0000110101	101001100- 10000-110
<i>M. magnahamata</i>	10020-1101	1001111010	00-0-0-000	???00-0000	010-00100?	0000010100	-----10110 100?0-110

Comentado [user1]: Where are the specimens used from? Where are the vouchers deposited? Please, include these data in Table 2.

<i>M. mahensis</i>	1002100111	1201111011	00-0-10001	??00-0110	010-00100?	000010-001	101101000-	00000-110
<i>M. marianae</i>	1000111111	1171111111	0111117001	1010111170	1110121010	000010-000	----1000-	00010-?10
<i>M. minuta</i>	10020-1101	1001111010	00-0-0-000	0-000-0000	010-001000	0000010000	----1000-	10000-110
<i>M. mirabilis</i>	10010-0111	1101111111	0100-2-1-1	1000100200	0010000101	1000010101	001201100-	01010-010
<i>M. montera</i>	1000110111	1101111111	0111110001	??01111110	111002101?	0000111101	000111110-	00000-110
<i>M. nonatoi</i>	1002101171	1171111011	0101111001	1000101110	0110111010	0100110000	----1110-	00000-110
<i>M. obockensis</i>	10010-0111	1111111111	0121111001	1001121210	0011021011	1000010101	211001100-	000?0-170
<i>M. pacifica</i>	1000110111	1111111111	0101110001	1110101110	1110021010	0000117101	000111000-	07000-110
<i>M. papillicornis</i>	10020-1101	1771111010	00-0-0-000	0-000-0000	010-001000	0000010000	----1010-	10000-?10
<i>M. parochilis</i>	10010-0111	1111111111	0101111001	0-11101210	0010021011	100000-001	211001100-	01000-110
<i>M. paulolanai</i>	1000111111	1101111111	0101110000	1070101270	111002101?	000?110101	007071100-	10000-110
<i>M. pettiboneae</i>	10020-1111	1101111010	00-0-0-000	0-000-0000	010-001000	0000110000	----1010-	00000-110
<i>M. phyllisae</i>	1002111101	1101111010	00-0-0-000	0-000-0000	010-001000	0000111000	----1010-	10000-?00
<i>M. piteikai</i>	1002100111	1111111111	10-1111001	11010-1210	1110021010	2000111101	1011?1110-	00010-?0?
<i>M. polydentata</i>	10020-2101	1211111011	10-0-17000	10200-0100	010-011000	011000-000	----1200-	000710111
<i>M. posterelongata</i>	1000110111	1771111011	10-0-10001	10010-0100	110-011000	0001010010	----1010-	00000-?10
<i>M. pulchella</i>	1000111111	1101111111	0101110001	??01101210	1110021010	0000110101	202001000-	00000-110
<i>M. pygmaea</i>	10020-1101	1001111010	00-0-0-000	??000-0000	010-00100?	000000-000	----1100-	10070-110
<i>M. riojai</i>	10010-0111	1111111111	1101111001	1101101211	0010000101	1000010101	101??1100-	01000-?10
<i>M. sacculata</i>	10010-0111	1701111111	0100-171-1	1011100??1	0010000101	1000010011	2110?1100-	01000-??0
<i>M. sinbadi</i>	1000110111	1101111111	0101111001	??0101210	0110021010	0100111100	----1100-	00000-110
<i>M. spinifera</i>	1010111111	1171111111	01011171-1	??00101210	0110021010	0700111070	----10711	17010-?10
<i>M. symmetrica</i>	1022102101	1211111010	10-0-0-000	??710-0000	010-00100?	010000-000	----1100-	000011111
<i>M. tehuansensis</i>	1010111111	1771111111	0101110001	1000101210	1110021010	0700110000	----1100-	10010-?10
<i>M. tinae</i>	10010-0111	1111111111	0111111001	??0121210	001102101?	1007010101	211001100-	00070-110
<i>M. uebelackerae</i>	1000110111	1771111011	00-0-10001	10100-0170	010-011010	0000110000	----10110	10000-??0
<i>M. variollamelata</i>	1002102101	1221111011	10-0-11001	11210-0100	110-011000	0110010000	----1100-	000010??1
<i>M. wilsoni</i>	1010112111	1101111011	0101111001	1020101210	0110121010	0100110000	----1100-	00070-110
<i>M. sp. 4 W Africa</i>	1002101111	??71111111	10-1110001	??010-1100	110-011010	0100010000	----1100-	00001011?
<i>M. sp. 4B-4B W Africa</i>	1002102111	??71111111	0101110001	1070101110	010-00100?	0100110100	----1100-	00001001?
<i>M. sp. 4C-4C W Africa</i>	1002102101	1211111111	0101110001	??01011100	010-00100?	0100110000	----1100-	00001111?
<i>M. sp. 5A W Africa</i>	1002101111	1211111011	10-0-11011	11710-0000	110-01101?	0100010001	101001000-	00001011?
<i>M. sp. 5B W Africa</i>	1002102111	1211111111	00-1110001	??700-1110	010-01101?	0100010000	----1000-	000011111
<i>M. sp. A</i>	10010-0111	1771111111	012112-001	1001121211	0012021011	1000010101	2110?1000-	00000-?10
<i>M. sp. B</i>	10010-0111	1771111111	012112-001	1001121210	0011000101	1000010001	2110?1100-	00000-??0
<i>M. sp. C</i>	10020-1101	1771111010	00-0-0-000	0-000-0000	010-001000	0000110000	----10110	10000-??0
<i>M. sp. D</i>	1010111111	1771111111	01011101-1	1000101210	1110021010	0700111000	----10711	10010-??0
<i>M. sp. E</i>	1010110111	1771111010	10-1110001	10000-1100	010-111010	0000110000	----10712	10000-?00
<i>M. sp. F</i>	1012101111	1771111010	00-0-17001	10000-0100	010-001000	0000010000	----1000-	00000-?00
<i>M. sp. G</i>	1000110111	1771111111	0101110001	1010101110	1110021010	0000110101	0000?1010-	00010-?10
<i>M. sp. H</i>	1000111111	1771111110	0100-10001	10000-0100	1110011000	0701111100	----1*110	00000-?00
<i>M. sp. I</i>	1002102101	1771111011	10-0-17001	10210-0170	110-011000	0110010000	----1200-	000010???
<i>M. sp. J</i>	1012112111	1771111011	0101111001	1020101210	0110121010	0100110000	----1100-	00000-?10
<i>M. sp. K</i>	1000110111	1771111111	0101111001	1000101110	1110021000	0700111100	----1100-	00000-?10
<i>M. sp. L</i>	1010111111	1771111111	0101110001	1000101210	0110011010	0700110101	0000?1100-	10010-?10

Octomagelona bizkaiensis 100210210? ???1101111 10-0-0-001 10210-0000 ----- -10000-000 -----1100- 10000-100
O. sp. W Africa 1002102101 ???1101111 10-0-0-001 ??20?00??? ----- -10?0????? ?????1??0- ??0?0-100