

# Performance in mortality prediction of SAPS 3 And MPM-III scores among adult patients admitted to the ICU of a private tertiary referral hospital in Tanzania: A retrospective cohort study

Nadeem Kassam<sup>Corresp., 1</sup>, Eric Aghan<sup>2</sup>, Samina Somji<sup>1</sup>, Omar Aziz<sup>3</sup>, James Orwa<sup>4</sup>, Salim R Surani<sup>5</sup>

<sup>1</sup> Internal medicine, Aga Khan University, Dar-es-Salaam, Tanzania

<sup>2</sup> Family medicine, Aga Khan University, Dar-es-Salaam, Tanzania

<sup>3</sup> Internal medicine, Aga Khan Hospital, Dar-es-Salaam, Tanzania

<sup>4</sup> Population health, Aga Khan University, Nairobi, Kenya

<sup>5</sup> Medicine & Pharmacy, Texas A&M University, Texas, United States

Corresponding Author: Nadeem Kassam  
Email address: nadeem.kassam@aku.edu

**Background:** Illness predictive scoring systems are significant and meaningful adjuncts of patient management in the Intensive Care Unit (ICU). They assist in predicting patient outcomes, improve clinical decision making and provide insight into the effectiveness of care and management of patients while optimizing the use of hospital resources. We evaluated mortality predictive performance of Simplified Acute Physiology Score (SAPS 3) and Mortality Probability Models (MPM<sub>0</sub>-III) and compared their performance in predicting outcome as well as identifying disease pattern and factors associated with increased mortality. **Methods:** This was a retrospective cohort study of adult patients admitted to the ICU of the Aga Khan Hospital, Dar- es- Salaam, Tanzania between August 2018 and April 2020. Demographics, clinical characteristics, outcomes, source of admission, primary admission category, length of stay and the support provided with the worst physiological data within the first hour of ICU admission were extracted. SAPS 3 and MPM<sub>0</sub>-III scores were calculated using an online web-based calculator. The performance of each model was assessed by discrimination and calibration. Discrimination between survivors and non – survivors was assessed by the area under the receiver operator characteristic curve (ROC) and calibration was estimated using the Hosmer-Lemeshow goodness-of-fit test. **Results:** A total of 331 patients were enrolled in the study with a median age of 58 years (IQR 43-71), most of whom were male (n=208, 62.8%), of African origin (n=178, 53.8%) and admitted from the emergency department (n=306, 92.4%). In- hospital mortality of critically ill patients was 16.1%. Discrimination was very good for all models, the area under the receiver-operating characteristic (ROC) curve for SAPS 3 and MPM<sub>0</sub>-III was 0.89 (95%CI:

0.844-0.935) and 0.90 (95%CI: 0.864-0.944) respectively. Calibration as calculated by Hosmer-Lemeshow goodness-of-fit test showed good calibration for SAPS 3 and MPM<sub>0</sub>-III with Chi-square values of 4.61 and 5.08 respectively and P - Value >0.05. Conclusion: Both SAPS 3 and MPM<sub>0</sub>-III performed well in our cohort of patients admitted to the intensive care unit of a private tertiary hospital. The in-hospital mortality of critically ill patients was lower compared to studies done in other intensive care units in tertiary referral hospitals within Tanzania.

# Performance in mortality prediction of SAPS 3 and MPM-III scores among adult patients admitted to the ICU of a private tertiary referral hospital in Tanzania: A retrospective cohort study

Primary Author: Nadeem Mehboob Kassam<sup>1</sup>

Co-Authors: Eric Luggar Aghan<sup>2</sup>, Samina Sadrudin Somji<sup>1</sup>, Omar Mohamed Aziz<sup>3</sup>, James Orwa Kenyatta<sup>4</sup>, Salim Ramzan Surani<sup>5</sup>

1. Department of Internal Medicine, Aga Khan University, Dar es Salaam, Tanzania
2. Department of Family Medicine, Aga Khan University, Dar es Salaam, Tanzania
3. Department of Internal Medicine, Aga Khan Hospital, Dar es Salaam, Tanzania
4. Department of Population Health, Aga Khan University, Nairobi, Kenya
5. Department of Medicine & Pharmacy, Texas A&M University, Texas, USA

Institutional address

1. Nadeem Kassam : [Nadeem.kassam@aku.edu](mailto:Nadeem.kassam@aku.edu)
2. Eric Aghan : [eric.aghan@aku.edu](mailto:eric.aghan@aku.edu)
3. Samina Somji : [samina.somji@aku.edu](mailto:samina.somji@aku.edu)
4. Omar Aziz : [omar.aziz@akhst.org](mailto:omar.aziz@akhst.org)
5. James Orwa : [james.orwa@aku.edu](mailto:james.orwa@aku.edu)
6. Salim Surani : [srsurani@hotmail.com](mailto:srsurani@hotmail.com)

Corresponding author: Nadeem Kassam  
[Nadeem.kassam@aku.edu](mailto:Nadeem.kassam@aku.edu)

# ABSTRACT

## Background

Illness predictive scoring systems are significant and meaningful adjuncts of patient management in the Intensive Care Unit (ICU). They assist in predicting patient outcomes, improve clinical decision making and provide insight into the effectiveness of care and management of patients while optimizing the use of hospital resources. We evaluated the mortality predictive performance of the Simplified Acute Physiology Score (SAPS 3) and Mortality Probability Models (MPM<sub>0</sub>-III) and compared their performance in predicting outcomes as well as identifying disease patterns and factors associated with increased mortality.

## Methods

This was a retrospective cohort study of adult patients admitted to the ICU of the Aga Khan Hospital, Dar es Salaam, Tanzania between August 2018 and April 2020. Demographics, clinical characteristics, outcomes, source of admission, primary admission category, length of stay and the support provided with the worst physiological data within the first hour of ICU admission were extracted. SAPS 3 and MPM<sub>0</sub>-III scores were calculated using an online web-based calculator. The performance of each model was assessed by discrimination and calibration. Discrimination between survivors and non-survivors was assessed by the area under the receiver operator characteristic curve (ROC) and calibration was estimated using the Hosmer-Lemeshow goodness-of-fit test.

## Results

A total of 331 patients were enrolled in the study with a median age of 58 years (IQR 43-71), most of whom were male (n=208, 62.8%), of African origin (n=178, 53.8%) and admitted from the emergency department (n=306, 92.4%). In-hospital mortality of critically ill patients was 16.1%. Discrimination was very good for all models, the area under the ROC curve for SAPS 3 and MPM<sub>0</sub>-III was 0.89 (95%CI: 0.844-0.935) and 0.90 (95%CI: 0.864-0.944) respectively. Calibration as calculated by Hosmer-Lemeshow goodness-of-fit test showed good calibration for SAPS 3 and MPM<sub>0</sub>-III with chi-square values of 4.61 and 5.08 respectively and P-Value >0.05.

## Conclusion

Both SAPS 3 and MPM<sub>0</sub>-III performed well in our cohort of patients admitted to the intensive care unit of a private tertiary hospital. The in-hospital mortality of critically ill patients was lower compared to studies done in other intensive care units in tertiary referral hospitals within Tanzania.

# BACKGROUND

The burden of critical care and ICU mortality is greatest in countries with low global national income (1). The reported ICU mortality widely varies from one setting to the other with higher rates reported in low and middle-income countries (LMICs) (1-3). As of 1<sup>st</sup> July 2020, the World Bank upgraded Tanzania's economic status from a low to lower-middle income country due to its strong economic performance over the past decade. However, the availability of intensive care units in Tanzania is very limited; none of the seven district hospitals surveyed in 2009 had an ICU. The four national referral hospitals had a total of only 38 ICU beds serving a population of 57 million (4). This is in contrast to high-income countries (HICs) which generally have between 5 to 30 ICU beds per 100 000 people (5). The availability and improvement of quality of care of critical illness in LMICs is necessary to reduce this burden and even more significant in the coming years as the population ages and prevalence of comorbidities increases (6).

Despite the use of high-cost and sophisticated devices, ICU mortality rates remain high. The burden of diseases compounded by a severe lack of resources, specialists and data makes prediction of ICU outcomes in terms of morbidity and mortality a crucial component of care across the continent. In HICs mortality prediction models are not only used to predict outcome but also as tools for quality enhancement and analytical decision making. These mortality predictive models were developed more than 25 years ago using patient characteristics. They help quantify the severity of illness, estimate the gravity of the disease, help predict outcome and assist in resource allocation (7, 8).

The three major predictive scoring systems used to predict mortality in general ICU patients are the Acute Physiologic and Chronic Health Evaluation (APACHE) scoring system, the Simplified Acute Physiologic Score (SAPS) and the Mortality Prediction Model (MPM<sub>0</sub>) (9). APACHE-IV, SAPS 3, and MPM<sub>0</sub>-III are the latest versions of the aforementioned scoring systems (10-12). When selecting a predictive scoring system for use in a given ICU, it is essential to use a model that is well-proven, established and validated contextually. APACHE-IV has long been considered more precise for predicting mortality but is perceived as burdensome and more costly especially in resource-limited settings (13). MPM<sub>0</sub>-III is considered superior in resource-limited settings since it has the lowest extraction burden among the three models and is available without cost on various medical information sites. MPM<sub>0</sub>-III has been a well-studied tool in East Africa. Nevertheless, its performance in predicting outcomes among critically ill patients admitted to ICU's in Kenya (14) and Rwanda (15) are contradictory. External validation of SAPS 3 among patients admitted to the ICU's in Austria, Brazil and Italy reported that SAPS 3 had a good ability to predict outcomes but performed poorly across all probabilities of death when compared with APACHE-IV and MPM<sub>0</sub>-III (16-18). SAPS 3 may have greater potential for international use since the

score was derived from data in more than one country (11). No study to date has assessed the performance of SAPS 3 in LMICs, especially in Sub-Saharan Africa.

These aforementioned predictive scoring systems have been compared in different studies and have produced variable results. The existence of a large number of scoring systems with contrasting performance suggests the best fit model is ICU specific. Thus, each particular ICU needs to determine which scoring system performs best in its setup; hence there was a need to carry out a comparative study in our cohort of patients to identify the best performing model. There have been no studies done in Tanzania that have compared the performance of these scoring models. The study had two main objectives: 1). To compare the performance of MPM<sub>0</sub>-III and SAPS 3 in order to identify which model best fits in the ICU of the Aga Khan Hospital, Dar es Salaam. 2). To identify disease patterns and risk factors associated with higher mortality rates among critically ill patients.

## METHODS

This was a single centre retrospective cohort study, conducted in the ICU of the Aga Khan Hospital, Dar-es-Salaam, Tanzania. The Aga Khan Hospital is the largest private hospital and the only Joint Commission International (JCI) accredited hospital in Tanzania. The ICU of the Aga Khan Hospital is a 15 bed unit which provides level III services to all kind of critically ill patients. The ICU is capable of providing mechanical ventilation, inotropic support and renal replacement therapy. The unit is divided in 3 sections – 7 adult general ICU beds (including 2 isolation rooms), 4 for cardiac patients and 4 for paediatrics. The ICU is an open-model one run by a multidisciplinary team comprising of physician of the primary specialty of care, physiotherapist and dietician led by a full-time critical care specialist. The nurse-to-patient ratio ranges between 1:1 and 1:2. All adult patients aged 18 and above admitted to the ICU were eligible for the study. Patients admitted for observation, having incomplete data and those whose duration of stay in the unit was less than an hour as well as those diagnosed with COVID-19 were excluded from the study. Admissions to the ICU are limited to those meeting a strict admitting criteria set by the hospital. A total of 747 adults patients were admitted to the ICU from August 2018 to April 2020. A sample size of 331 patients having a specific outcome (death or discharge home) was determined to be sufficient to give the study a 80% power and 95% confidence for detection of 10% difference in performance between SAPS 3 and MPM<sub>0</sub>-III. The ICU admission register was used to identify patients admitted and patient files were retrieved from the medical records. The medical file numbers were entered into a computer and computer generated random sampling was performed until the desired sample size was achieved. Patient demographics and clinical data were extracted using patient records and were

entered into a spreadsheet on Microsoft Office Excel 2010 (Redmond, WA, USA). Data was extracted by experienced junior doctors with working experience in the ICU and was independently verified by the primary author for accuracy and completeness. The reasons for admission were grouped into 11 categories: surgery, gastroenterology, neurology, endocrinology, respiratory, cardiovascular, nephrology, sepsis, oncology, hematology, obstetrics and gynecology. When multiple diagnoses were present, the leading one with the worst prognosis was selected as the main reason for admission.

Descriptive analysis of demographic characteristics were done and presented as percentages while the categorical and continuous outcome variables were analysed and presented as means and medians with interquartile ranges respectively. Categorical and continuous variables between survivors and non-survivors were compared using Pearson's chi-square test and Mann-Whitney *U* test respectively. SAPS-3 and MPM<sub>0</sub>-III were calculated using an online scoring calculator, available on [www.uptodate.com](http://www.uptodate.com). Accurate discrimination and calibration are key distinguishing features that should be met by all predictive scoring models. Discrimination of the model was assessed by the area under the receiver operating characteristic (ROC) curve. An area of 0.7–0.8 is reflected as fair, 0.8–0.9 good and > 0.9 excellent. Non-parametric Wilcoxon statistics was used to compare the area under the ROC curves (19). A Hosmer-Lemeshow goodness-of-fit test which follows chi-square distribution was used to evaluate the model fit as well as calibration of the models with a p-value of > 0.05 signifying no evidence of poor fit (19). However, all other statistical tests with a p-value of < 0.05 were considered statistically significant. Any variable with P-Value <0.05 and those considered clinically significant in explaining mortality in the ICU were considered in multivariable model. Determinants of mortality among critically ill patients were identified using binary logistic regression; odds ratio with corresponding 95% Confidence intervals (CI) and P-value were reported. All statistical analysis was done using STATA version 15. The study protocol was approved by the Ethical Research Committee (ERC) of the Aga Khan University (AKU/2020/051/fb) and individual consent of each study participant was exempted since it did not affect the rights and welfare of the patients.

## RESULTS

### General characteristics of the participants

A total of 331 patients were included in the study. Out of the 331 patients (n=278, 83.9%) survived and (n=53, 16.1%) died. Table 1 below shows general and clinical characteristic of the cohort and provides a comparison of survivors to non-survivors. The median age of the cohort was 58 years (IQR 43-71) with more than half of admitted patients being male (n=208, 62.8%) and of African origin (n=178, 53.8%).. Most of the patients were admitted to the ICU from the emergency department (n=306, 92.5%), who were

at home prior (n=318, 96.1%) with majority of them suffering from neurological disease (n=63, 19%), sepsis (n=60, 18.1%), respiratory (n=36, 10.9%) and cardiovascular (n=36, 10.9%) related conditions. Median ICU and hospital LOS were 4 (2-6) and 6 (4-10) days respectively.

When survivors and non-survivors were compared, there was a statistically significant difference (P-value <0.05) in age, length of ICU stay, admitting category and code status. Higher mortality rates were noted in the elderly aged between 75-84 years (n=15, 28.3%), those suffering from sepsis (n=13, 24.5%) and neurological diseases (n=10, 18.9%). Higher percentage of mortality was also noted in Do-Not-Resuscitate (DNR) patients (n=27, 50.9%) compared to those without limitations of care. There was no statistically significant difference between survivors and non-survivors by sex, ethnicity and prior location before ICU admission. The overall SAPS 3 score for all the patients was 42 (IQR: 32-51) of which non-survivors had a higher score 60 (IQR: 51-68) than the survivors 39 (IQR: 31-48) with p-value (<0.0001). Similarly, the median MPM<sub>0</sub>-III score in non-survivors, 5 (IQR: 4-6) was higher than survivors, 3 (IQR: 2-4) with p-value (<0.0001).

Table 2 below, illustrates the type of support patients received in the first 24 hours of ICU admission. Of the 331 patients admitted to the ICU, (n=123, 37.2%) patients received support in the first hour of ICU admission that included: mechanical ventilation, inotropes and hemodialysis. Highest mortality was noted amongst those who required inotropic support and mechanical ventilation (n=21, 39.6%) in the first hour of ICU admission.

Table 3 below highlights comorbid conditions amongst the critical ill patients admitted to the ICU. More than one comorbid condition per critically ill patient was recorded when present. The most common comorbid condition amongst our cohort was hypertension (n=174, 52.6%) and diabetes mellitus (n=107, 32.3%). When comorbid conditions were compared between survivors and non-survivors, a higher percentage of mortality with statistically significant difference (P value <0.05) was noted among those suffering with chronic kidney disease (n=14, 26.4%) and liver cirrhosis (n=7, 13.2%).

Calibration of each scoring system exhibited good performance. The goodness of fit Hosmer-Lemeshow test and p-value of each scoring system is shown in Table 4 below. The area under ROC of SAPS 3 and MPM<sub>0</sub>-III in prediction of mortality are shown below in Figure 1 below. The area under the ROC was calculated to evaluate the predictive value of the scoring systems. The area under the ROC curve for the



SAPS 3 showed a statistically significant predictive marker of mortality (AUC: 0.8892; 95%CI: 0.844-0.935). The cut-off value for SAPS 3 was 54 with the sensitivity of 72% and specificity of 90%. The MPM<sub>0</sub>-III scoring system also showed a statistically significant predictive marker for the outcome of interest (AUC: 0.904; 95%CI: 0.864-0.944). The cut-off value for MPM<sub>0</sub>-III was 4, with sensitivity of 74% and specificity of 87%. There was no statistically significant difference between the ROC curves of the two models (P-value=0.2418) (Table 5).

The overall estimated median (IQR) predicted mortality among the 331 ICU patients was 6% (2%-20%) on the SAPS 3 model and 11.5% (3.8%-27.9%) based on the MPM<sub>0</sub>-III model. The stratified analysis by survivors and non-survivors is shown in Figure 2 below. The median predicted mortality risk for survivors is lower than those of non-survivors. In the SAPS 3 model, the estimated median for survivors was 5% (IQR: 1%-11%) while for the non-survivors this was 50% (IQR: 34%-69%). Based on the MPM<sub>0</sub>-III model the median predicted mortality was 9.1% (3.1%-17%) and 68.5% (IQR: 42.7%-84.0%) for survivors and non-survivors respectively.

Multiple clinical factors were associated with increased adjusted odds of mortality. These included length of ICU stay (adjusted odds ratio [aOR], 1.462; P-value = 0.001) and those transferred from the ward (aOR, 5.341; P-value < 0.022). However, it was protective to stay longer in the hospital as the odds of mortality decreased as the length of hospitalization increased (aOR, 0.717; P-value = 0.002) (Table 6).

## DISCUSSION

To our knowledge, this is the first study to report on performance of predictive scoring models in Tanzania and more so in a private setting. Accurate discrimination and calibration are two key characteristics that should be met by all predictive scoring systems. Both SAPS 3 and MPM<sub>0</sub>-III performed well in our cohort. According to our results, a SAPS 3 score of higher than 54 can predict mortality with sensitivity of 72% and specificity of 90%. A MPM<sub>0</sub>-III score of greater than 4 can predict mortality with sensitivity of 74% and specificity of 87%.

Discrimination describes the accuracy of a given prediction. In our cohort, the discriminatory capability of both SAPS 3 (20 variables) and MPM<sub>0</sub>-III (16 variables) was good. There was no statistically

significance difference when both these models were compared, suggesting that the model with more variables was not associated with better discriminatory performance. MPM<sub>0</sub>-III has been externally validated in various ICUs in North America (12, 13, 20) and has shown to have good discrimination which was similar to our study finding. However, a study done at Aga Khan University Hospital, Nairobi, Kenya (14) and two public ICUs in Rwanda(15) showed MPM<sub>0</sub>-III to have fair discrimination amongst their cohort. This observed difference in discrimination may be due to the effect of differences in proportion of case mixes between the study settings. Similarly SAPS 3 has been externally validated in various ICUs in Italy (16), Brazil (17), Austria (18) and found to have good discriminatory capability amongst their cohort. Despite SAPS 3 having greater prospective for international generalizability there has been no published studies evaluating its performance in Sub-Saharan African ICUs. This is the first study that reports its potential for application in LMICs.

Calibration describes how the instrument performs over a wide range of predicted mortalities. Calibration is sensitive to alterations in case-mix and patient care and interventions. Despite its tendency to deteriorate over time and leading to overestimation of mortality (17), both SAPS 3 and MPM<sub>0</sub>-III were well calibrated amongst the critically ill patients admitted at our study setting. Our study findings were contrary to SAPS 3 validation studies mentioned earlier which reported poor calibration and overestimation of mortality (16-18). However, external validation studies have reported MPM<sub>0</sub>-III to have good calibration (12, 13, 20). Earlier studies mentioned that were conducted in Sub-Saharan Africa have produced contrasting results. The MPM<sub>0</sub>-III was well calibrated amongst the critical ill patients admitted to the ICU of the Aga Khan University Hospital, Nairobi (14) but showed poor calibration amongst all adult patients admitted to Rwanda's two public ICUs (15). These findings highlight the similar treatment protocols and interventions between two sister hospitals located in different geographical regions.

In this retrospective study we also aimed to identify patient demographics, disease patterns, clinical outcomes as well as factors associated with higher risk of mortality in patients admitted to the ICU of the Aga Khan Hospital, Dar -es Salaam. Based on this retrospective observational cohort, the in-hospital mortality of critically ill patients was 16.1%, which is far less than the reported mortality among all other tertiary referral hospitals in Tanzania, 41.4% (21) but slightly exceeds rates reported in western Europe and North America (1). This disparity is not surprising since the intensive care unit at our setting has access to more resources than similar units in the country and comparable in various ways to facilities in HICs. The ICU cohort studied in the four tertiary referral hospitals in Tanzania was younger (median age 34 years, IQR 21-53) compared to our study population (median age 58 years, IQR 43-71). This variation

could be due to the exclusion of patients aged less than 18 years in our study. However both the cohorts had male predominance of 57.5% and 62.8 % respectively (21). The bulk of admissions in our cohort were those suffering from neurological disease, sepsis, respiratory and cardiovascular related conditions. Mortality was highest among those admitted due to sepsis. Our results are in parallel with a large intercontinental database that emphasized the association of sepsis with high mortality rates in all countries (1). The median length of ICU stay is similar to reports from tertiary hospitals in Sub-Saharan Africa (21, 22).

Prolonged length of stay (LoS) in the ICU and patients transferred from the general ward to the ICU were factors associated with higher adjusted odds of mortality among critically ill patients. Prolonged LoS in the ICU may be attributed to development of multi- systemic complications necessitating continued organ support. There are no laws and guidelines in Tanzania with regards to withdrawal of support, hence we hypothesize that significant fraction of patients with a prolonged course of illness and with expected poor outcomes are admitted for extended intervals before succumbing to death. Our study findings are comparable to several studies done in well-equipped ICUs that concluded patients with multiple diseases and having organ dysfunction were key factors that prolong the LoS in ICU (23, 24). Contrasting results have also been published that LoS in ICU is not an independent risk factor for in-hospital mortality (25). Those patients transferred from the general ward to the ICU also had higher adjusted odds of mortality; this is not surprising since it is a mere reflection of deteriorating physiological and clinical condition. Few studies have demonstrated early transfer to the ICU for treatment to have a substantial impact on in-hospital mortality and LoS (26, 27).

We identified several limitations in our study. Firstly, this was a single center study and as such the findings may not be valid across all patient populations in Tanzania. Secondly, since our study was a retrospective design it restricted us the ability to follow up outcomes after ICU discharge and doesn't provide the same level of evidence as a prospective study design.

## CONCLUSION

In summary, this is the first and largest study to report on performance of predictive scoring models in Tanzania. Our study concluded both SAPS 3 and MPM<sub>0</sub>-III perform well among critically ill patients admitted to the ICU of the Aga Khan Hospital, Dar es Salaam, Tanzania. We found our mortality among critically ill patients to be much lower compared to other tertiary referral hospitals in Tanzania. Amongst out cohort, patients with sepsis had the highest mortality rate. Thus clinical research targeting infection prevention efforts and early implementation of targeted interventions would be important to improving

outcomes. Prolonged ICU stay and transfer from general wards to ICU were key factors of mortality. Of note, the performance of predictive scoring models tend to deteriorate over time; termed as worsening of discrimination and calibration and resulting in overestimation of mortality (17). Thus periodic updating is crucial for sustaining accuracy of these predictive models. .

## ABBREVIATIONS

- APACHE: Acute physiology and chronic health evaluation
- SAPS: Simplified acute physiology score
- MPM: Mortality probability models
- LoS : Length of Stay
- ICU :- Intensive care Unit

## DECLARATION

### Ethics approval and consent to participate

The study was approved by the Aga Khan University Ethical Research Committee (AKU-ERC, EA). The National Institute for Medical Research (NIMR) mandates the AKU-ERC to approve health research conducted by Tanzanian staff and students under the Act of Parliament No. 23 of 1979 and its amendments in 1997.

### Consent for publication

Not Applicable

### Acknowledgements

Special thanks to Dr. Kamran Hameed, head of the Internal medicine department at the Aga Khan University and hospital, Dar-es-Salaam for his continuous mentorship and guidance during my residency program.

### Author Details

1. Department of Internal Medicine, Aga Khan University, Medical College, East Africa
2. Department of Family Medicine, Aga Khan University, Medical College, East Africa
3. Department of Internal Medicine, Aga Khan Hospital, Dar-es-Salaam, Tanzania
4. Department of Population Health, Aga Khan University, Medical College, East Africa
5. Department of Medicine & Pharmacy, Texas A&M University, Texas, USA


# REFERENCE

1. Vincent JL, Marshall JC, Namendys-Silva SA, Francois B, Martin-Loeches I, Lipman J, et al. Assessment of the worldwide burden of critical illness: the intensive care over nations (ICON) audit. *Lancet Respir Med.* 2014;2(5):380-6.
2. Ilori IU, Kalu QN. Intensive care admissions and outcome at the University of Calabar Teaching Hospital, Nigeria. *J Crit Care.* 2012;27(1):105 e1-4.
3. Smith ZA, Ayele Y, McDonald P. Outcomes in critical care delivery at Jimma University Specialised Hospital, Ethiopia. *Anaesth Intensive Care.* 2013;41(3):363-8.
4. Baker T, Lugazia E, Eriksen J, Mwafongo V, Irestedt L, Konrad D. Emergency and critical care services in Tanzania: a survey of ten hospitals. *BMC Health Serv Res.* 2013;13:140.
5. Dondorp AM, Iyer SS, Schultz MJ. Critical Care in Resource-Restricted Settings. *JAMA.* 2016;315(8):753-4.
6. Adhikari NK, Fowler RA, Bhagwanjee S, Rubenfeld GD. Critical care and the global burden of critical illness in adults. *Lancet.* 2010;376(9749):1339-46.
7. Keegan MT, Gajic O, Afessa B. Severity of illness scoring systems in the intensive care unit. *Crit Care Med.* 2011;39(1):163-9.
8. Zimmerman JE, Wagner DP, Knaus WA, Williams JF, Kolakowski D, Draper EA. The use of risk predictions to identify candidates for intermediate care units. Implications for intensive care utilization and cost. *Chest.* 1995;108(2):490-9.
9. Juneja D, Singh O, Nasa P, Dang R. Comparison of newer scoring systems with the conventional scoring systems in general intensive care population. *Minerva Anesthesiol.* 2012;78(2):194-200.
10. Zimmerman JE, Kramer AA, McNair DS, Malila FM. Acute Physiology and Chronic Health Evaluation (APACHE) IV: hospital mortality assessment for today's critically ill patients. *Crit Care Med.* 2006;34(5):1297-310.
11. Metnitz PG, Moreno RP, Almeida E, Jordan B, Bauer P, Campos RA, et al. SAPS 3--From evaluation of the patient to evaluation of the intensive care unit. Part 1: Objectives, methods and cohort description. *Intensive Care Med.* 2005;31(10):1336-44.
12. Higgins TL, Teres D, Copes WS, Nathanson BH, Stark M, Kramer AA. Assessing contemporary intensive care unit outcome: an updated Mortality Probability Admission Model (MPM0-III). *Crit Care Med.* 2007;35(3):827-35.
13. Kuzniewicz MW, Vasilevskis EE, Lane R, Dean ML, Trivedi NG, Rennie DJ, et al. Variation in ICU risk-adjusted mortality: impact of methods of assessment and potential confounders. *Chest.* 2008;133(6):1319-27.
14. Lukoko LN, Kussin PS, Adam RD, Orwa J, Waweru-Siika W. Investigating SOFA, delta-SOFA and MPM-III for mortality prediction among critically ill patients at a private tertiary hospital ICU in Kenya: A retrospective cohort study. *PLoS One.* 2020;15(7):e0235809.
15. Riviello ED, Kiviri W, Fowler RA, Mueller A, Novack V, Banner-Goodspeed VM, et al. Predicting Mortality in Low-Income Country ICUs: The Rwanda Mortality Probability Model (R-MPM). *PLoS One.* 2016;11(5):e0155858.
16. Poole D, Rossi C, Anghileri A, Giardino M, Latronico N, Radrizzani D, et al. External validation of the Simplified Acute Physiology Score (SAPS) 3 in a cohort of 28,357 patients from 147 Italian intensive care units. *Intensive Care Med.* 2009;35(11):1916-24.
17. Nassar AP, Jr., Mocelin AO, Nunes AL, Giannini FP, Brauer L, Andrade FM, et al. Caution when using prognostic models: a prospective comparison of 3 recent prognostic models. *J Crit Care.* 2012;27(4):423 e1-7.
18. Metnitz B, Schaden E, Moreno R, Le Gall JR, Bauer P, Metnitz PG, et al. Austrian validation and customization of the SAPS 3 Admission Score. *Intensive Care Med.* 2009;35(4):616-22.

19. Steyerberg EW, Vickers AJ, Cook NR, Gerds T, Gonen M, Obuchowski N, et al. Assessing the performance of prediction models: a framework for traditional and novel measures. *Epidemiology*. 2010;21(1):128-38.
20. Higgins TL, Kramer AA, Nathanson BH, Copes W, Stark M, Teres D. Prospective validation of the intensive care unit admission Mortality Probability Model (MPM0-III). *Crit Care Med*. 2009;37(5):1619-23.
21. Sawe HR, Mfinanga JA, Lidenge SJ, Mpondo BC, Msangi S, Lugazia E, et al. Disease patterns and clinical outcomes of patients admitted in intensive care units of tertiary referral hospitals of Tanzania. *BMC Int Health Hum Rights*. 2014;14:26.
22. Kwizera A, Dunser M, Nakibuuka J. National intensive care unit bed capacity and ICU patient characteristics in a low income country. *BMC Res Notes*. 2012;5:475.
23. Toptas M, Sengul Samanci N, Akkoc I, Yucetas E, Cebeci E, Sen O, et al. Factors Affecting the Length of Stay in the Intensive Care Unit: Our Clinical Experience. *Biomed Res Int*. 2018;2018:9438046.
24. Moitra VK, Guerra C, Linde-Zwirble WT, Wunsch H. Relationship Between ICU Length of Stay and Long-Term Mortality for Elderly ICU Survivors. *Crit Care Med*. 2016;44(4):655-62.
25. Williams TA, Ho KM, Dobb GJ, Finn JC, Knuiman M, Webb SA, et al. Effect of length of stay in intensive care unit on hospital and long-term mortality of critically ill adult patients. *Br J Anaesth*. 2010;104(4):459-64.
26. Churpek MM, Wendlandt B, Zadavec FJ, Adhikari R, Winslow C, Edelson DP. Association between intensive care unit transfer delay and hospital mortality: A multicenter investigation. *J Hosp Med*. 2016;11(11):757-62.
27. Young MP, Gooder VJ, McBride K, James B, Fisher ES. Inpatient transfers to the intensive care unit: delays are associated with increased mortality and morbidity. *J Gen Intern Med*. 2003;18(2):77-83.


# Figure 1

Receiver operating curve for predicting mortality according to SAPS 3 and MPM<sub>0</sub> III models



# Figure 2

Median predicted mortality rates for SAPS 3 and MPM<sub>0</sub>-III





**Table 1** (on next page)

General and clinical characteristics of patients (N=331)

1 **Table 1: General and clinical characteristics of patients (N=331)**

Characteristics	All( N=331) N (%)	Survivors (n=278) n (%)	Non-survivors (n=53) n (%)	p-value
<b>Sex</b>				
Male	208(62.8)	174(62.6)	34(64.2)	0.829
Female	123(37.2)	104(37.4)	19(35.8)	
<b>Age group in years</b>				
< 45	87(26.3)	79(28.4)	8(15.1)	0.015
45-64	114(34.4)	99(35.6)	15(28.3)	
65-74	61(18.4)	51(18.4)	10(18.9)	
75-84	51(15.4)	36(12.9)	15(28.3)	
> 84	18(5.4)	13(4.7)	5(9.4)	
<b>Ethnicity</b>				
African	178(53.8)	152(54.7)	26(49.1)	0.589
Asian	136(41.1)	111(39.9)	25(47.2)	
Other	17(5.1)	15(5.4)	2(3.8)	
<b>Admitted from</b>				
Emergency	306(92.5)	257(92.5)	49(92.5)	0.800
Wards	15(4.5)	12(4.3)	3(5.7)	
Clinic	10(3.0)	9(3.2)	1(1.9)	
<b>Location before ICU admission</b>				
Home	318(96.1)	267(96.0)	51(96.2)	0.950
Hospital	13(3.9)	11(4.0)	2(3.8)	
<b>Admitting Category</b>				0.014
Surgery	38(11.5)	36(12.9)	2(3.8)	0.014
Gastroenterology	31(9.4)	25(9.0)	6(11.3)	
Neurology	63(19.0)	53(19.1)	10(18.9)	
Endocrinology	18(5.4)	17(6.1)	1(1.9)	
Respiratory	36(10.9)	35(12.6)	1(1.9)	
Cardiovascular	36(10.9)	30(10.8)	6(11.3)	
Nephrology	16(4.8)	12(4.3)	4(7.6)	
Sepsis	60(18.1)	47(16.9)	13(24.5)	
Obstetrics and Gynecology	10(2.0)	9(3.2)	1(1.9)	
Hematology	7(2.1)	5(1.4)	2(3.8)	
Oncology	16(4.8)	9(3.2)	7(13.2)	
<b>Code Status on Admission</b>				
DNR	40(12.1)	13(4.7)	27(50.9)	< 0.001
Full Code	291(87.9)	265(95.3)	26(49.1)	
<b>Age in years</b>	<b>58(43-71) †</b>	<b>55.5(41-70) †</b>	<b>70(55-78) †</b>	<b>0.0003‡</b>
<b>SAPS 3 scores</b>	<b>42(32-51) †</b>	<b>39(31-48) †</b>	<b>60(51-68) †</b>	<b>&lt; 0.0001‡</b>
<b>MPM<sub>0</sub>III scores</b>	<b>3(2-4) †</b>	<b>3(2-4) †</b>	<b>5(4-6) †</b>	<b>&lt; 0.0001‡</b>
<b>LOS ICU (days)</b>	<b>4(2-6) †</b>	<b>4(2-6) †</b>	<b>6(2-11) †</b>	<b>0.0029‡</b>
<b>LOS Hospital (days)</b>	<b>6(4-10) †</b>	<b>6(4-10) †</b>	<b>8(3-13) †</b>	<b>0.3248‡</b>

2  
3 †median (IQR); ‡: p-value for Mann-Whitney U test; LOS: Length of Stay, DNR: Do Not  
4 resuscitate. Data in median (IQR), and n (%)



# **Table 2**(on next page)

Type of support received in the first hour of ICU admission

1 **Table 2: Type of support received in the first hour of ICU admission**

Type of support	All( N=331)	Survivors (n=278)	Non-survivors (n=53)	p-value
	N (%)	n (%)	n (%)	
None	208(62.8)	196(70.5)	12(22.6)	< 0.001
Hemodialysis	13(3.9)	11(4.0)	2(3.8)	0.9498
Inotropes	33(10.0)	29(10.4)	4(7.6)	0.5206
Mechanical ventilation	26(7.9)	17(6.1)	9(17.0)	0.0070
Inotropes, Hemodialysis	5(1.5)	5(1.8)	0(0.0)	0.3252
Inotropes, mechanical ventilation	36(10.9)	15(5.4)	21(39.6)	< 0.001
Mechanical ventilation, Hemodialysis	3(0.9)	3(1.1)	0(0.0)	0.4474
Inotropes, mechanical ventilation, hemodialysis	7(2.1)	2(0.7)	5(9.4)	0.0001

2

3 *Data presented in n (%)*

**Table 3**(on next page)

Comorbid conditions among critically ill patients admitted to the ICU

**Table 3: Comorbid conditions among critically ill patients admitted to the ICU**

Comorbidity	All( N=331)	Survivors (n=278)	Non-survivors (n=53)	p-value
	N (%)	n (%)	n (%)	
Hypertension	174(52.6)	146(52.5)	28(52.8)	0.9553
Diabetes Mellitus	107(32.3)	87(31.3)	20(37.7)	0.3613
Heart Failure	53(16.0)	41(14.7)	12(22.6)	0.1510
Chronic Kidney Disease	45(13.6)	31(11.2)	14(26.4)	0.0030
HIV	21(6.3)	16(5.8)	5(9.4)	0.314
COPD	17(5.1)	15(5.4)	2(3.8)	0.6239
CAD	17(5.1)	15(5.4)	2(3.8)	0.6239
Liver Cirrhosis	16(4.8)	9(3.2)	7(13.2)	0.0022
DM & HTN	87 (26.3)	72 (25.9)	15 (28.3)	0.7157
HTN & CKD	39 (11.8)	27 (9.7)	12 (22.6)	0.0075
DM & HTN & CKD	30 (9.1)	21 (7.6)	9 (17.0)	0.0285

*Data presented in n (%)*

**Table 4**(on next page)

Goodness of fit Hosmer-Lemeshow test and p-value of each scoring model



**Table 4: Goodness of fit Hosmer-Lemeshow test and p-value of each scoring model**

Scoring Model	Chi – Square	P – Value
MPM 0- III Score	5.08	0.2791
SAPS 3	4.61	0.7980

# **Table 5**(on next page)

Area under curve and 95% confidence Intervals for the models

**Table 5: Area under curve and 95% confidence Intervals for the models**

Variable	Cut-off	AUC	LL	UL	P-value
MPM 0- III Score	4	0.904	0.864	0.944	0.2418
SAPS 3	54	0.8892	0.8440	0.935	

*LL: Lower limits; UL: Upper limits; AUC: Area under the ROC curve*

# **Table 6**(on next page)

Factors associated with increased odds of mortality among critically ill patients

1 **Table 6: Factors associated with increased odds of mortality among critically ill patients**

Characteristics	Unadjusted Odds ratio			Adjusted Odds ratio		
	OR	95%CI	P-value	OR	95%CI	P-value
<b>Age in years</b>	1.032	1.014-1.051	0.001	1.020	0.997-1.043	0.086
<b>LOS in ICU</b>	1.068	1.021-1.117	< 0.001	1.462	1.179-1.814	0.001
<b>LOS in hospital</b>	1.017	0.994-1.041	0.141	0.717	0.580-0.886	0.002
<b>Sex</b>						
Male	ref					
Female	0.935	0.507-1.723	0.829	0.870	0.399-1.893	0.725
<b>Admitted From</b>						
Emergency	ref					
Wards	1.311	0.357-4.819	0.683	5.341	1.278-22.322	0.022
Clinic	0.583	0.072-4.704	0.612	1.033	0.114-9.347	0.977
<b>Code status</b>						
DNR	ref					
Full code	0.047	0.022-0.102	< 0.001	0.052	0.021-0.129	< 0.001

2