

Global gap-analysis of amphipod barcode library

Anna Maria Jazdzewska^{Corresp., 1}, Anne Helene S. Tandberg², Tammy Horton³, Saskia Brix⁴

¹ Department of Invertebrate Zoology and Hydrobiology, Faculty of Biology and Environmental Protection, University of Lodz, Lodz, Poland

² University museum, Department of Natural History, University of Bergen, Bergen, Norway

³ National Oceanography Centre, Southampton, United Kingdom

⁴ Department for Marine Biodiversity Research (DZMB), Senckenberg am Meer, Hamburg, Germany

Corresponding Author: Anna Maria Jazdzewska
Email address: anna.jazdzewska@biol.uni.lodz.pl

In the age of global climate change and biodiversity loss there is an urgent need to provide effective and robust tools for diversity monitoring. One of the promising techniques for species identification is the use of DNA barcoding that in Metazoa utilizes the so called ‘gold-standard’ gene of cytochrome c oxidase (COI). However, the success of this method relies on the existence of trustworthy barcode libraries of the species. The Barcode of Life Data System (BOLD) aims to provide barcodes for all existing organisms, and is complemented by the Barcode Index Number (BIN) system serving as a tool for potential species recognition. Here we provide an analysis of all public COI sequences available in BOLD of the diverse and ubiquitous crustacean order Amphipoda, to identify the barcode library gaps and provide recommendations for future barcoding studies. Our gap analysis of 25702 records has shown that although 3835 BINs (indicating putative species) were recognised by BOLD, only 10 % of known amphipod species are represented by barcodes. We have identified almost equal contribution of both records (sequences) and BINs associated with freshwater and with marine realms. Three quarters of records have a complete species-level identification provided, while BINs have just 50%. Large disproportions between identification levels of BINs coming from freshwaters and the marine environment were observed, with three quarters of the former possessing a species name, and less than 40% for the latter. Moreover, the majority of BINs are represented by a very low number of sequences rendering them unreliable according to the quality control system. The geographical coverage is poor with vast areas of Africa, South America and the open ocean acting as “white gaps”. Several, of the most species rich and highly abundant families of Amphipoda (e.g. Phoxocephalidae, Ampeliscidae, Caprellidae), have very poor representation in the BOLD barcode library. As a result of our study we recommend stronger effort in identification of already recognised BINs, prioritising the studies of families that are known to be important and abundant

components of particular communities, and targeted sampling programs for taxa coming from geographical regions with the least knowledge.

Global gap-analysis of amphipod barcode library

Anna M. Jazdzewska¹, Anne Helene S. Tandberg², Tammy Horton³, Saskia Brix⁴

¹ Department of Invertebrate Zoology and Hydrobiology, Faculty of Biology and Environmental Protection, University of Lodz, Lodz, Poland, author ORCID: 0000-0003-2529-0641

² University of Bergen, University museum, Department of Natural History, Bergen, Norway, author ORCID: 0000-0003-3470-587X

³ National Oceanography Centre, Southampton, UK, author ORCID: 0000-0003-4250-1068

⁴ Senckenberg am Meer, Department for Marine Biodiversity Research (DZMB), Hamburg, Germany, author ORCID: 0000-0002-3269-8904

Corresponding Author:

Anna Jazdzewska¹
12/16 Banacha st., 90–237 Lodz, Poland
Email address: anna.jazdzewska@biol.uni.lodz.pl

Abstract

In the age of global climate change and biodiversity loss there is an urgent need to provide effective and robust tools for diversity monitoring. One of the promising techniques for species identification is the use of DNA barcoding that in Metazoa utilizes the so called ‘gold-standard’ gene of cytochrome *c* oxidase (COI). However, the success of this method relies on the existence of trustworthy barcode libraries of the species. The Barcode of Life Data System (BOLD) aims to provide barcodes for all existing organisms, and is complemented by the Barcode Index Number (BIN) system serving as a tool for potential species recognition. Here we provide an analysis of all public COI sequences available in BOLD of the diverse and ubiquitous crustacean order Amphipoda, to identify the barcode library gaps and provide recommendations for future barcoding studies. Our gap analysis of 25702 records has shown that although 3835 BINs

(indicating putative species) were recognised by BOLD, only 10 % of known amphipod species are represented by barcodes. We have identified almost equal contribution of both records (sequences) and BINs associated with freshwater and with marine realms. Three quarters of records have a complete species-level identification provided, while BINs have just 50%. Large disproportions between identification levels of BINs coming from freshwaters and the marine environment were observed, with three quarters of the former possessing a species name, and less than 40% for the latter. Moreover, the majority of BINs are represented by a very low number of sequences rendering them unreliable according to the quality control system. The geographical coverage is poor with vast areas of Africa, South America and the open ocean acting as “white gaps”. Several, of the most species rich and highly abundant families of Amphipoda (e.g. Phoxocephalidae, Ampeliscidae, Caprellidae), have very poor representation in the BOLD barcode library. As a result of our study we recommend stronger effort in identification of already recognised BINs, prioritising the studies of families that are known to be important and abundant components of particular communities, and targeted sampling programs for taxa coming from geographical regions with the least knowledge.

Keywords

DNA barcoding, Crustacea, marine, freshwater, semi-terrestrial, taxonomic identification

Introduction

Nature in the age of Anthropocene is facing numerous global changes and challenges. One of the drastic results of human associated activities is the acceleration of species extinctions, with one

million species estimated to be presently critically endangered (IPBES 2019). What is more, although the rate of species discovery grows, large numbers of species remain undescribed and it is believed many will not be recognized before they go extinct (Mora et al., 2011). This raises the challenge of efficient environmental monitoring, which is crucial for biodiversity recognition and preservation. Monitoring based on the taxonomic identification of organisms in samples is time-consuming and requires knowledge of the studied group. In the time of the taxonomic impediment (Ebach, Valdecasas & Wheeler, 2011), species identification methods offering an alternative to morphology-based methods are of great interest. Utilization of DNA-barcoding (identifying sequences of individual specimens), metabarcoding (high-throughput identification of bulk samples) and the use of environmental DNA (e-DNA, identifying DNA of taxa directly from water or soil sample, without collection of specimens) have been presented as promising methods in monitoring and ecological studies (e.g., Hajibabaei et al., 2012; Cristescu, 2014; Aylagas et al., 2018; Leese et al., 2018; Bush et al., 2019; Feio et al., 2020). The use of metabarcoding in assessing the status of ecosystems has already received the new term “Biomonitoring 2.0” (Bush et al., 2019). Such approaches require the existence of well-established barcode fragment libraries, which allow accurate recognition of organisms in the environment (Cristescu, 2014; Cowart et al., 2015; Oliveira et al., 2016; Múrria et al., 2020). Recent studies indicate that although the use of barcoding in biomonitoring has great advantages over morphological identification, the current gaps in barcode libraries may hinder their use (Weigand et al., 2019; Feio et al., 2020; Hestetun et al., 2020; Leite et al., 2020; Múrria et al., 2020).

There are two main repositories where DNA sequences are deposited: NCBI GenBank (www.ncbi.nlm.nih.gov/genbank/, Sayers et al., 2020) and Barcode of Life Data System (BOLD,

www.boldsystems.org, Ratnasingham & Hebert, 2007). In contrast to GenBank, which assembles nucleotide data of all genes, the primary aim of BOLD is to store data used for species barcoding, which in the case of Metazoa is the cytochrome *c* oxidase (COI) gene. The development of the BOLD database included the Barcode Index Number (BIN) system implementation (Ratnasingham & Hebert, 2013) that intends to help in biodiversity assessments by providing species-level taxonomic registry. Based on a molecular species delimitation method, each Molecular Operational Taxonomic Unit (MOTU) recognized by BOLD receives a unique alphanumeric code (BIN). Ideally, each BIN is associated with an accurate taxonomic (preferably species) identification and links to the voucher stored in a recognised institution. However, in practice this is not working well, and at the time of system implementation as many as 46% of BINs lacked species names (Ratnasingham & Hebert, 2013). This issue has arisen for a variety of reasons, which we investigate in this study using a particular faunal group, the Amphipoda, as a model.

The Order Amphipoda are peracarid crustaceans belonging to the class Malacostraca. They are very diverse components of aquatic environments. According to the World Amphipoda Database (WAD, Horton et al., 2020, accessed on 17-07-2020) there are 10235 accepted amphipod species, the majority of which (78%) inhabit the marine realm, around 20% are freshwater species and just 2% are terrestrial taxa (Horton et al., 2020; Väinölä et al., 2008). The discovery rate of new species has grown steadily since the first amphipod species description and has particularly accelerated in the last six decades (Horton et al., 2020) with mean number of over 100 taxa annually described since the 1960s (Coleman, 2015). If the trend from the last sixty years persists, we may expect to have ca. 8000 new species described by 2100. More conservative estimates predict that 6100 new species will be described by that date (Arfianti,

Wilson & Costello, 2018). The use of molecular methods in the studies of Amphipoda has revealed very high species diversity (e.g. Knox et al., 2012; Verheye, Backeljau & d'Udekem d'Acoz, 2016; Jazdzewska & Mamos, 2019) and revealed the existence of cryptic species complexes within widely distributed taxa (Witt, Therloff & Hebert, 2006; Mamos et al., 2014; Wysocka et al., 2014; Havermans, 2016). Amphipoda are not only a species-rich group, but they also often dominate the crustacean assemblages in which they occur (e.g., Corkum, 1989; Humphries, Davies & Mulcahy, 1996; Vinogradov, Volkov & Semenova, 1996; Jazdzewski et al., 2001; Väinölä et al., 2008; Frutos, Brandt & Sorbe, 2017; Brix et al., 2018; Havermans & Smetacek, 2018). They can be found in both the benthos and the pelagic realm, presenting a variety of states of mobility (from epibenthic clingers to fully mobile swimmers) and, as a result, possess a wide variety of feeding habits including herbivory, detritivory, necrophagy, omnivory, predation and ectoparasitism (Barnard & Karaman, 1991; Vinogradov, Volkov & Semenova, 1996; Dauby, Scailteur & De Broyer, 2001; Väinölä et al., 2008). Being diverse and abundant they are important prey items for other invertebrates and vertebrates, including fish, birds and mammals (e.g. Dalpadado et al., 2001; Dauby, Nyssen & De Broyer, 2003). Certain species of Amphipoda are used in laboratory ecotoxicological studies (Hyne & Everett, 1998; Bundschuh et al., 2011; Bundschuh et al., 2013, Major et al., 2013). Some amphipod species are well-adapted to anthropogenic environments such as artificial structures used in coastal protection or are part of fouling communities, and have shown a high invasion potential worldwide (e.g., Bij de Vaate et al., 2002; Kelly et al., 2006; Cabezas et al., 2014; Rewicz et al., 2015; Beermann et al., 2020; Sedano et al., 2020).

The combined factors of high diversity and the important role played by amphipods in the aquatic ecosystem highlights the need for accurate species identifications which are required for

biological monitoring programs. The use of DNA-barcoding may speed up the identification process, but it will only succeed if the barcode library is well-established and robust. Recent gap-analyses of the barcode libraries in aquatic European environments showed very large differences in the coverage between different taxonomic groups and geographic regions (Weigand et al., 2019; Feio et al., 2020; Hestetun et al., 2020; Leite et al., 2020). These studies used species lists restricted to particular geographic regions or chosen taxonomic groups. Basic summaries concerning the extent of amphipod data in BOLD identified problems with lack of taxonomic identification or detailed geographic information as well as contamination with human or bacterial DNA and provided recommendations to improve the data (Radulovici & Coleman, 2017; Coleman & Radulovici, 2020). However, to date there are no detailed analyses that have been conducted on a single taxon on a global scale.

In this study we have conducted a gap-analysis of the barcode library of a single crustacean order, the Amphipoda, on a global basis. In producing an up-to-date picture of the current state of knowledge, we will provide researchers with a detailed understanding of the both the strengths and the potential limitations of the use of DNA barcodes for identifications. We also propose recommendations for future initiatives that involve molecular data and produce new barcodes to fill the gaps in our knowledge of this taxon.

Material and methods

Data for the present study were retrieved from BOLD by searching the “Public Data Portal” using the keyword “Amphipoda”. A combined dataset of all records was downloaded as an .xml file on June 24th 2020.

All records of the barcoding fragment of the cytochrome c oxidase I (COI-5P in BOLD) were extracted (29016 records). This extracted dataset was used for all further analyses conducted by using various filtering options in an Excel spreadsheet. 2579 records, represented by sequences shorter than 500 bp or having more than 1% ambiguous nucleotides for which BINs were not ascribed, were removed from dataset. Continued analysis of the dataset revealed some duplicate records (1468 records, 734 cases, Supplemental file 1). These derived from data harvested by BOLD from GenBank and seemed to be associated with an update of the records GenBank. In the dataset, these records had an identical sample ID that referred to a GenBank Accession Number but with an additional ‘.1’ appended (e.g. KP713892 and KP713892.1) with an identical identification provided. The differences were often linked with more detailed geographical information in the case of one record from the pair. Only the more detailed entry was retained for continued analysis. One sequence of *Niphargus novomestanus* S. Karaman, 1952 (KR858496, BOLD:ADD1128) was removed from the dataset because it was deleted from GenBank by its submitter (“This record was removed at the submitter’s request because the source organism cannot be confirmed.” GenBank website). The resulting dataset contained 25702 records (Fig. 1, Supplemental file 2).

Each record in the dataset was then further refined by sorting into categories according to the level of taxonomic identification. The following categories were used: order, family, subfamily, genus and species. Where records were provided with a temporary species identification, i.e. they are recognised as separate morphospecies but are not determined to correspond to a known taxon - they were treated as a separate category. In the whole dataset ca. 2.5% of records (596 individuals, 145 BINs) had uncertain identification with “cf.” or “aff”. Because the majority of them (417 records, 101 BINs) were associated with five species of one genus (*Gammarus*) for

simplification all such records were treated as final species identifications. However, it is understood that the use of open nomenclature, when applied to identifications, provides an indication of the level of uncertainty, and may be intended to indicate the presence of new species or species complexes.

The data in BOLD come from wide variety of projects, some of which involve detailed taxonomic study by specialists, others are focused on monitoring or other topics in which taxonomic specialists are not involved. For the purposes of our analyses it was assumed that the identification accuracy was equal throughout the whole dataset, regardless of its origin. In several cases identification of the specimens within a single BIN varied strongly, with some records remaining at order level while others were determined to the species level. BINs aim to represent a putative species, so in the above example, the most detailed taxonomic information was applied to all records within the single BIN. Sometimes multiple (most often two) species or genus names were associated with a single BIN (87 cases). Each of these cases was checked individually. Sometimes it was an obvious misidentification of a single individual within a large group - if this was noted the misidentified record was added as an additional element to the records identified to the lowest congruent level (e.g. if the genus name matched the BIN genus, the misidentified taxon was added as an additional record identified to the genus level, if the lowest congruent level was family it was added to the family records); and the taxon identification of the majority of records was applied as correct. When it was impossible to judge which name was correct, the name of the identifier was checked and identifications carried out by taxonomists specializing in Amphipoda was prioritised over a that provided by a non-specialist study. Where this process did not give a satisfactory conclusion, the BIN was allocated an identification at a rank that was congruent for the different records. The list of taxa with

incongruent identifications together with an explanation of the final decision is presented in Supplemental file 3.

Based on the taxonomic identification of the records the associated BINs were divided into the following environmental categories:

- a) marine
- b) freshwater
- c) terrestrial.

Taxa that inhabit both marine realm and brackish environments were allocated to the marine category. Taxa from freshwater also occurring in brackish waters were allocated to the freshwater category. All representatives of the family Talitridae were treated as terrestrial taxa. Where taxonomic information was not detailed enough to provide environmental information about the particular BIN, the geographic data (coordinates and/or locality description) of the associated records were used to ascribe a particular BIN to one of the above categories. In some cases, this necessitated checking the original publication. A small number of unallocated BINs (18) and associated records (44) were removed from the dataset (Supplemental file 4).

In order to verify the correct environmental allocation of BINs, all BINs with records possessing coordinates were plotted on a map using the software QGIS2.16.1 (QGIS Development Team, 2018). Cases where incongruence between the ascribed environment and the geographic position appeared were checked individually. For those records without detailed geographic information the country of origin was taken from either BOLD or the associated publication.

In order to verify the barcode coverage within the studied group a list of BINs associated with a species name was compared with the list of accepted amphipod species names available in the World Amphipoda Database (WAD, Horton et al., 2020, accessed on 17-07-2020). A barcode

quality assessment of the species represented in BOLD, based on the grading system proposed by Oliveira et al. (2016) and slightly modified by Fontes et al. (2020) was applied. This system consists of five grades: A – consolidated concordance (>10 sequences of a single morphospecies grouped in a single BIN), B – basal concordance (same as grade A but between three and 10 sequences available in the library), C – multiple BINs (one morphospecies assigned to more than one BIN), D – insufficient data (single species is assigned to single BIN but it is represented by less than three sequences in the barcode library), E – discordant species assignment (more than one species assigned to a single BIN). Fontes et al. (2020) provide an R-based application (Barcode, Audit & Grad System – BAGS), and uses only those records possessing either coordinates or the indication of the country of origin. Since this information is very often missing in the data mined from GenBank, and our aim was to focus on all available barcode records, the assessment was carried out manually. Additionally, as a result of initial treatment of the dataset, misidentified species records or BINs with unclear species identification, were already removed, so category E (discordant species assignment, Oliveira et al., 2016; Fontes et al., 2020) was not recorded. For the purpose of the present study *Lysianassoidea incertae sedis* was treated as an additional family. The amphipod families were divided into four categories depending on the number of species in each: low species rich families (up to 10 species), moderately species rich families (from 11 to 30 species), species rich families (31-100 species), very species rich families (more than 100 species). This division allowed verification of pattern between the species richness of the family and its representation in BOLD.

Results

Of the 25702 amphipod COI records, 46.5% (11958 records) were freshwater, 43.5% (11169 records) were from the marine realm, and 9.8% (2531 records) were terrestrial taxa. Of the 3835

recognized BINs in total, 45% (1726 BINs) belonged to freshwater taxa, 50% (1920 BINs) were marine, and 4.5% (171 BINs) were from terrestrial taxa. 44 records (0.2%) and their associated 18 BINs (0.5%) could not be ascribed to the above environmental categories (Fig. 2A, B).

More than half (57.5%) of the records available in BOLD possessed coordinates, and 20% had information about the country of origin. Geographic information about the remaining 22.5% was provided only in the original publication and this information was added manually. Geographic information is more comprehensive for marine taxa, where 71% of records possessed coordinates (compared to 47% for freshwater, and 50% for terrestrial taxa). Molecular studies of freshwater Amphipoda are focused mainly in the Northern hemisphere (particularly European countries, Russia and United States) while in the Southern hemisphere, Australia, New Zealand and Argentina are well studied (Fig. 3A). There is a complete lack of records (amphipod sequences) from Brazil, equatorial America and vast areas of Africa. Similar patterns of data coverage were seen for marine amphipods, which have greater numbers of records along European, North American and East Asian coasts. In the Southern hemisphere, Australia, New Zealand and Antarctica had larger numbers of barcode records (Fig. 3B). However, vast areas of the deep sea and the Arctic Ocean remain undersampled. Terrestrial Amphipoda in Europe, North America, China, Australia and Chile were the best represented (Fig. 3C), but sampling gaps were seen in the continents of South America and Africa.

The majority of records with environmental information (69.8%, 17922 recs.) had a complete species-level identification. Of the remaining 30.2% of records, 5.6% (1433 recs.) had received temporary names (open nomenclature), 11.3% (2902 recs.) remained identified at the genus level, 0.2% (40 recs.) at subfamily, 5.0% (1285 recs.) at family, and 8.1% (2076 recs.) at the order level. Levels of identification varied according to the environment, with marine taxa

having greater proportions of taxa identified only to higher taxonomic ranks (Fig. 4A). The majority of BINs with geographic affiliation (3817) were associated with species names (55.7%, 2126 BINs). These were followed by BINs identified to the order level (13.3%, 506 BINs), generic or family level (10.7%, 407 BINs each) and those with a temporary name (9.4%, 359). BINs with only a subfamily name constituted just 0.3% (12). Greater variations between environments were seen for the BINs, with 74% (1284) of freshwater BINs having a species level identification, compared to only 39% (751) of marine BINs (Fig. 4B). More than 20% (444, 23%) of the BINs for marine taxa remained identified at the order level.

Regardless of the environmental origin, the majority of BINs were represented by a single sequence (Fig. 5). BINs represented by five or fewer sequences constituted around two thirds (67%, 114 terrestrial BINs to three quarters, 78%, 1488 marine BINs) of BINs recorded in a particular environment. Freshwater taxa had 41 BINs (2.4%) represented by more than 50 sequences, compared to 28 (1.5%) for marine taxa, and eight (4.7%) for terrestrial taxa. When only those BINs with complete species-level identifications are considered, the proportion of sequences representing a particular MOTU does not change, with freshwater taxa having 78% of BINs (1016) represented by five or fewer sequences. Almost three quarters of marine BINs (71%, 525 BINs) had five or fewer sequences in BOLD, while this proportion was 61% (56 BINs) for terrestrial taxa. Freshwater taxa had 35 BINs (3%) represented by more than 50 sequences, compared to 27 (4%) for marine taxa, and 6 (7%) for terrestrial taxa. The best represented BIN in BOLD (801 sequences) belonged to the terrestrial species *Orchestoidea tuberculata* Nicolet, 1849 (BOLD:ACQ3380), followed by the marine species *Gammarus oceanicus* Segestråle, 1947 (BOLD:AAA1262, 553 sequences), and the freshwater species *Diporeia hoyi* (S.I. Smith, 1874) (BOLD:AAA1473, 512 sequences). A further 26 BINs were

represented by more than 100 sequences, including 17 freshwater, seven marine and two terrestrial BINs (Supplemental file 5).

Out of the 3817 studied BINs, just over half (55.7%, 2126) were associated with a species-level identification, representing 1001 species. Freshwater BINs with species identification reached 1284, associated with 453 species, while 751 marine BINs were determined to 496 species. Of the 91 terrestrial BINs, 52 species were identified. Generally, a single morphological species was associated with each BIN (68%, 680 cases, 288 in freshwater, 359 marine, 33 terrestrial). 17% of the identified species were associated with two different BINs (72 freshwater, 82 marine and 14 terrestrial) (Fig. 6). There were however 19 cases when one single morphological species was represented by more than 10 BINs (17 freshwater, one marine and one terrestrial) (Supplemental file 6). The greatest number of BINs was recorded for the freshwater species *Gammarus balcanicus* Schäferna, 1923 represented by 143 BINs (45 BINs were identified as “cf.” or “aff.”) followed by another freshwater taxon *Hyalrella azteca* (Saussure, 1858) (62 BINs) and *Gammarus fossarum* Koch, 1836 (51 BINs; 19 BINs identified as “cf.” or “aff.”). Among terrestrial taxa the highest molecular variation (12 BINs) was recorded for *Morinoia japonica* (Tattersall, 1922) (present in BOLD under former generic name *Platorchestia*), while *Apohyale stebbingi* Chevreux, 1888 (with 11 BINs recognized) was the most diverse among marine species.

Of the 239 accepted families of Amphipoda (238 families and Lysianassoidea *incertae sedis*), 105 (44%) were represented by at least one species in BOLD (Table 1). The largest number of families had up to 20% of species barcoded, while only ten families had more than half of the known species barcoded (Supplemental file 7). Thirteen families lacking barcoded species had at

least one barcoded taxon identified at the genus level, a further five families had a taxon identified at the family level.

Just under ten percent (999 spp., 9.7%) of the 10330 accepted species of Amphipoda (Horton et al., 2020) had barcodes. Of the nominal species possessing barcodes almost 500 (496 spp.) are marine, 451 spp. are freshwater and 52 spp. are terrestrial taxa. The data coverage of the majority of species, no matter their environmental origin, is not sufficient for the barcodes to be trusted according to the quality control system (Table 2) (Oliveira et al., 2016; Fontes et al., 2020). Additionally, a large group of taxa is represented by multiple BINs; only 10% of species represent consolidated concordance of available barcodes.

The breakdown of amphipod families according to the assigned categories of richness and their respective representation in BOLD can be seen in Table 3. Almost every one of the very species rich families had at least one species barcoded (31 families out of 32), and 22 of 30 species rich families are represented in BOLD. For both moderately low and low species rich families 26 possessed at least one representative in BOLD constituting respectively 48% and 21% of all families each (Supplemental file 7). The mean coverage of barcodes for species in each of the above groups was around 10% with the highest observed for low species rich families (12%) and the lowest (8%) recorded for families grouping from 30 to 100 species. However, if the families without any molecular information were removed from the study these numbers considerably change. The low species rich families (1-10 spp.) had a barcode coverage at the level of 49%, moderately species rich families (11-30 spp.) reached 21% of coverage, while the rich and very rich amphipod families (more than 30 spp.) had only 9-10% of species studied.

A third of families (34) have at least one species characterized by consolidated concordance of available barcodes (category A of the quality grading system). Another third of families (38) do

not have any species in categories A or B, indicating that the species already studied represent a potential cryptic diversity or the available data are insufficient (Table 4).

Within the very species rich families, the best representation in BOLD was recorded for Niphargidae (36.5% of known species represented with a barcode), Gammaridae (31%) and Crangonyctidae (16%). Only the family Stegocephalidae did not have any representative with species level identification (although barcodes belonging to this family but identified at genus level were present). The least studied families within this group (but having at least one species barcode) were: Phoxocephalidae (1% of the species with a barcode), Dexaminidae, Liljeborgiidae and Maeridae (ca. 2% of the species with a barcode). Among species rich families 41% of the species from Pseudoniphargidae had barcodes, while the Epimeriidae and Pontogammaridae had 20% and 19%, respectively. The best represented moderately species rich families were Metacrangonyctidae, Oxycephalidae and Hyperiidae with 55%, 50% and 48% of the associated species represented with a barcode respectively. Within low species rich families four (Baikalogammaridae, Crymostygidae, Cyllopodidae and Tryphanidae) had all known species represented with barcodes, but other than Cyllopodidae (two species) the families are monotypic (Supplemental file 7).

Discussion

Extent of barcode library of Amphipoda

One of the aims of establishing the BOLD database was to store and publish barcodes, based on records uploaded by its users and supplemented by the data harvested from GenBank (Ratnasingham & Hebert, 2007). Together with the BIN system, that groups similar sequences in clusters representing putative species (Ratnasingham & Hebert, 2013), the BOLD database aids in recognising and quantifying biodiversity. The extent of data in BOLD expresses the activity of

researchers studying particular groups using molecular methods. The number of available sequences of Amphipoda in BOLD is comparatively large. At the time of download (end of June 2020) Amphipoda in BOLD were represented by almost 26 000 records (3835 BINs), and by the end of August there were more than 34 000 public sequences (3914 BINs) (BOLD accessed on 20-08-2020), indicating the great intensity of molecular studies involving this crustacean group, and that the data in BOLD are actively growing. Among other crustacean groups only Decapoda is represented by a higher number of records (64 281 records). Copepoda are represented by 18 511, Thecostraca by 15 554, Isopoda by 13 858 and Branchiopoda by 12 326 sequences. The large number of identified BINs within the Amphipoda also places this group second only to Decapoda (with 6056 BINs). Isopods and copepods are represented by 1853 and 1804 BINs, respectively, while 969 BINs were identified within Branchiopoda. Within Thecostraca only 545 BINs were identified (boldsystems.org, accessed on 20-08-2020).

Ratnasingham & Hebert (2013) when presenting the BIN system indicated that overall 12% of the sequences available in BOLD at that time were lacking a family name, 19% a genus name and 40% a species name. A comparison of these numbers with the present data on Amphipoda looks optimistic, where only 8% of sequences are without family indication, 13% without genus and 29% species identification. However, the global analysis of Ratnasingham & Hebert (2013) also identified 10% of BINs lacking family names, almost 24% lacking generic names and 46% lacking species names. These numbers are almost identical for amphipod BINs known presently (13%, 23%, 43% of BINs lacking family, genus and species information, respectively). Among all known species of Amphipoda, almost 80% of species are marine, some 20% live in freshwaters, while 2% may be considered as terrestrial (Horton et al., 2020; Väinölä et al., 2008). The above proportions are expressed neither in the number of records nor the number of

recognized BINs that are more or less evenly distributed between freshwater and marine taxa. This demonstrates that in terms of amphipod crustaceans freshwater taxa are much better studied than the marine taxa. These disproportions are even more striking when the level of identification of sequences and BINs is considered. Although the majority of data present in BOLD possess species-level identifications, marine amphipods are? less thoroughly identified. This is especially clear for marine BINs, of which only 39% had species-level identifications, while as much as one fifth are identified only as “Amphipoda”. The fact that freshwater amphipods are better studied is not surprising considering the easier access to this environment. In the case of marine fauna, obtaining samples suitable for molecular analysis can be challenging, especially when extreme habitats (polar regions, deep-sea, hydrothermal vents etc.) are considered (e.g., Riehl et al., 2014; Jazdzewska & Mamos, 2019;). Additionally, rarity is a common feature of numerous marine species (particularly in the deep-sea environment, see Kaiser et al., [2007]), where many taxa are known only from their original descriptions and type localities (Jazdzewska & Mamos, 2019). The question of how many of the BINs not associated with a species identification actually belong to already known species is also of concern. In these cases, it is highly advisable to put every effort to identify the already available material – this will relatively efficiently improve data usability. Taxa that are associated with a BIN, yet are known to be new to science are another cause for concern. This is particularly evident for marine taxa collected during recent deep-sea exploration programs (e.g., Brandt et al., 2007; Jazdzewska, 2015; Brandt et al., 2019; Brix et al., 2020). It is imperative that full scientific descriptions of new species are produced to reduce the current proliferation of ‘dark taxa’ (Page, 2016).

The geographic distribution of available amphipod sequence records shows clear sampling gaps. In particular the African continent, the northern part of South America and the Coral Triangle in Asia are complete “white spots” when freshwater and terrestrial taxa are considered. For marine species, the coasts of Africa and South America, the Coral Triangle, and large parts of the deep sea of all oceans, lack coverage. Considering the known high species diversity of these regions it will be necessary to establish targeting sampling programs before we can consider that we have adequate global coverage of the molecular diversity of the Amphipoda.

Our study shows that globally the barcoding coverage of amphipod species is only about 10%. In comparison, just over 20% of all species registered in the European Register of Marine Species (ERMS) and almost 50% of species listed in the AZTI Marine Biotic Index (AMBI) have been barcoded (Weigand et al., 2019). The percentage of barcoded European freshwater invertebrates used in environmental monitoring reaches 64.5%, and when considering only Peracarida, 24% of ERMS species, 45% of AMBI and 82% of freshwater monitored taxa have been barcoded (Weigand et al., 2019). It has to be emphasized however, that only ERMS lists all marine invertebrates from European region, while both other datasets studied by Weigand et al. (2019) consists of a subset of species from this area. More specific studies of Iberian macroinvertebrates revealed that ca. 40% of amphipod species possess barcodes (Leite et al., 2020; Múrria et al., 2020;). Hestetun et al. (2020) conducted a barcode library gap-analysis of the benthic macrofauna of one region of the North Sea, which indicated the barcode coverage varying from 42.4% to 61% (depending on the calculation method). This indicates that for smaller subset of taxa and specified geographic region it is much easier to produce good barcode coverage. It can be concluded that although Amphipoda are an actively studied taxonomic group where scientists

increasingly use molecular methods, this diverse and abundant macrofaunal taxon is still insufficiently represented in the BOLD barcode library.

Quality of amphipod barcode library

In order to provide a trusted barcode for a particular species, at least one good quality sequence associated with a species-level identification provided by taxonomic specialist is required as an absolute minimum. However, a single sequence cannot provide information about intraspecific variation, and overlooked contamination of the sample will mean the sequence cannot be validated. As such, it is advisable to provide a small number of sequences to characterise each taxon. The recently proposed barcode quality auditing system suggests providing at least three sequences to enable proper barcode evaluation (Oliveira et al., 2016; Fontes et al., 2020). Unfortunately, as we have shown in the case of Amphipoda, globally more than half of BINs are represented by only 1-2 sequences in BOLD. This low number of sequences places them in category D of the Oliveira et al. (2016) system, indicating the existing data is insufficient for use as trusted barcodes. Similar observations for a restricted amphipod dataset are made by Fontes et al. (2020).

Due to methodological differences it is impossible to make direct comparisons of our data with the results of the gap analysis of aquatic organisms in European waters (Weigand et al., 2019), but re-calculation of their data shows much improved barcode coverage. Among all freshwater invertebrates 65% of taxa barcoded are represented by more than five sequences, while this percentage rises to 77% when considering only freshwater Peracarida. This proportion of high quality datasets diminishes when marine taxa are considered; with 52% of the marine species from the AMBI list and 45% those listed in ERMS having at least five barcodes available. These numbers do not change when considering only marine Peracarida (52% and 46% of the ones

presented in AMBI and ERMS lists, respectively). Our analysis of Amphipoda shows opposite pattern with about 1/4-1/3 of BINs represented by more than five sequences but the good barcode coverage observed by Weigand et al. (2019) may be biased by the fact that they targeted the species used in water quality assessment programs. Because of their practical use such taxa receive more scientific interest and it may be assumed that their barcoding is prioritized by different institutions.

The amphipod BINs that have the largest numbers of sequences in BOLD are often the result of detailed studies of targeted species, which have produced large numbers of sequences as a secondary aim of the study. For example, 750 out of the 801 sequences in BOLD of terrestrial *Orchestoidea tuberculata* come from a single study by Brante et al. (2019); 406 records out of 411 sequences in BOLD of freshwater *Dikerogammarus haemobaphes* (Eichwald, 1841) come from Jazdzewska et al. (2020); while 232 records of 235 sequences in BOLD of marine *Caprella scaura* Templeton, 1836 come from Cabezas et al. (2014). The disproportional representation between the few species that are very thoroughly studied and the remaining majority of species that are represented only by a single, or a low number of sequences emphasises the need for more targeted sampling of less common species.

Best studied families and cryptic diversity

Almost half of the 239 known amphipod families are represented in BOLD. However, only ten of these families have more than 50% of their associated species sequenced. It is important to underline that there are 18 families in BOLD that do not have species-level identifications, but have records left at the family or genus level. A small effort to provide trusted species-level identifications for these taxa will greatly improve barcode coverage of the Amphipoda, particularly if they represent species already known to science.

458 Another concern that has arisen as part of this study relates to the format of temporary names in
 459 GenBank and BOLD, the different requirements by users for their input, and how this has
 460 changed following development of the databases. In GenBank, the incorporation of temporary
 461 names or codes is allowed (referred to as placeholder names in GenBank). In 2010, a large
 462 amount of COI data was incorporated into the BOLD database. The identifications associated
 463 with each of these imported sequences were included verbatim from GenBank. BOLD users,
 464 however, were originally able to use temporary names in the database only in private
 465 projects/dataset and when opening their data for public they were expected to provide the
 466 identification to the lowest taxonomic level possible (e.g. genus) and to provide the temporary
 467 name (e.g. incorporating “cf.” or “aff.”) as a taxonomy note (that has happened to the authors of
 468 the present paper). However, in BOLD a taxonomy note is only visible when the specimen page
 469 is open, and not in a general search. Recently, we have learnt that open nomenclature identifiers
 470 (such as ‘cf.’ and ‘aff.’) are accepted by BOLD, but it may be assumed that numerous records
 471 remain at a higher taxonomic level, with more detailed identifications available that are hidden
 472 from general searches. This discrepancy in dealing with temporary names has become apparent
 473 when analysing the whole dataset as part of this study. In particular, the inconsistent use of
 474 temporary names in these databases mean that it is very difficult to differentiate between
 475 temporary names which are being used to refer to species that are new to science, and those
 476 which have remained at a higher taxonomic level because they were simply not identified further
 477 (which could be for a variety of reasons). Molecularly well-defined temporary names for new
 478 species are likely to become more abundant and therefore critical to our knowledge of
 479 biodiversity in the coming years, and we need to ensure they are managed carefully and
 480 consistently. Recommendations for the use of open nomenclature have been proposed recently to

attempt to standardise and overcome these issues (Sigovini, Keppel & Tagliapietra, 2016; Horton et al., 2021) and it is hoped that these standard formats will be considered for use in both BOLD and GenBank.

Barcode coverage of families varies depending on the species richness. For species rich families it is around 10%, while coverage is increased for moderate and low species rich taxa. This is not surprising considering it is much easier to receive better coverage for monotypic families or those represented by only a few species. The best studied families are the ones that remain under the interest of large working groups who focus on studying specific families (e.g. Hou, Fu & Li, 2007; Hou, Sket & Li, 2014; Mamos et al., 2014; Wysocka et al., 2014; Delić et al., 2017a; Delić et al., 2017b; Fišer et al., 2017; Copilaş-Ciocianu, Sidorov & Gontcharov, 2019). It is worth noting that providing barcodes is generally more a “by-product” of other analyses than the goal per se. Another issue that should be emphasized is that species rich families are proportionally under studied. This is important because they usually do not only group many species but very often the species from these families constitute the majority of amphipods characterizing different assemblages. This is clearly shown by the Phoxocephalidae (1% of the 367 known species are barcoded), Ampeliscidae (7% of 312 of the known species barcoded) or Oedicerotidae (10% of the 246 known species barcoded), all constitute very large and important components of marine benthic communities worldwide (Brandt, 1993; Weissshappel & Svavarsson, 1998; Frutos & Sorbe, 2017; Brix et al., 2018;). Another example is provided by the Caprellidae (6% of the 443 known species are barcoded) which are an important part of many fouling communities (e.g., Ros, Vázquez-Luis & Guerra-García, 2013; Ros et al., 2013;) and where proper species identifications are crucial in the context of growing transport with their

resulting potential alien species invasions (op. cit.). The studies of these families should be prioritized in order to support marine monitoring programs based on barcode libraries.

The analysis of the amphipod BINs with a species-level identification showed that there were only a few cases where multiple names were associated with a single molecular unit. A quarter of these cases resulted from the misidentification of single individuals within a taxon. In some cases different names were associated with the description of new species (present in the database under both former and newly established name). Problems with morphological identification of cryptic species and the lack of well established diagnostic characters within closely related species may also be the reason of the presence of multiple names for single BIN.

The above problems have been recognized within *Gammarus ochridensis* Schäferna, 1926 species complex that is the group of morphologically very similar species of which two *Gammarus cryptosalemaai* Grabowski, Wysocka & Mamos, 2017 and *Gammarus cryptoparechiniformis* Grabowski, Wysocka & Mamos, 2017 are recognizable only based on molecular data (Wysocka et al., 2013; Grabowski, Wysocka & Mamos, 2017). This indicates that generally BOLD may be considered a trusted tool for species identification. Our analyses showed that in the majority of cases, a single BIN was characterising a single species, which is congruent with the results of other similar studies (Fontes et al., 2020; Leite et al., 2020). Some morphologically identified species were represented by two or even three BINs, which can indicate overlooked diversity. It has been noted however, that sometimes due to the methodology used during BIN-identification and the threshold used (2% of similarity, Ratnasingham & Hebert, 2013) some valid species may be split into two or more BINs (Lörz, Jazdzewska & Brandt, 2018; Jazdzewska & Mamos, 2019). This happens more frequently when the sample size is small and the intraspecific variation range cannot be adequately assessed. In such cases, the

use of additional genes or other data analysing methods may help to decide the proper species delineation. The present study revealed 19 morphological species that were represented by 11 or more BINs. This multi-BIN representation was much more common in freshwater environments, where 17 species with potential cryptic diversity were observed. The existence of such high cryptic diversity especially in European waters was recognized by authors of the original works (Witt, Threlhoff & Hebert, 2006; Bauzà-Ribot et al., 2011; Major et al., 2013; Mamos et al., 2014; Wysocka et al., 2014; Delić et al., 2017a; Delić et al., 2017b, Fišer et al., 2017; Tomikawa et al., 2017) most recently confirmed also by Wattier et al. (2020). A detailed study of the available barcodes and cryptic diversity of the Gammaridae and other representatives of the superfamily Gammaroidea is in preparation (Mamos pers. comm.). The large representation of freshwater taxa forming cryptic species complexes (especially in Europe) can be partly explained by the geological events that shaped the European freshwater system (Wysocka et al., 2014; Mamos et al., 2016; Wattier et al., 2020). Presence of marine cryptic or pseudo-cryptic species have also been reported (Havermans, 2016; Verheye, Backeljau & d'Udekem d'Acoz, 2016;), but the extent of molecular studies of amphipods from this realm is much smaller and as a result cryptic species may have been overlooked. A study of the marine genus *Apohyale* showed high diversification of species within the genus, and confirms that more studies are required to correctly identify species diversity and uncover cryptic diversity in marine taxa (Desiderato et al., 2019). Cases of highly diverse nominative species usually come from studies based in a single research group that was already aware of the high diversity within the taxon. There are, however, cases where multiple BINs have received the same identification but this was carried out by different authors at different times (without comparison of the material) and it is difficult to judge if the observed diversity is a result of the existence of a cryptic species or of

misidentification of the species. In such cases it is impossible to decide which of the BINs represents the known species and which are cryptic/new species that require more detailed study (Jazdzewska et al., 2018). A detailed analysis of the species represented by several BINs was not the focus of the present study, but it should be a priority for BOLD to identify such cases and inform users about the presence of possible cryptic taxa. Users of BOLD who seek to obtain identification for their own sequence should be notified that the specimen they have may belong to a group of cryptic species so that the taxonomic identification can be treated with caution. It is also highly recommended that mistakes or problematic issues that are found in the database are corrected and published e.g. the case of *Hyperrella antarctica* Bovallius, 1887/*H. dilatata* Stebbing, 1888 which was recently clarified by Havermans et al. (2019).

Conclusions and Future recommendations

We have conducted a gap-analysis of the barcode library using a single crustacean order, the Amphipoda, as a model. The high diversity and the important role played by amphipods in the aquatic ecosystem combine to highlight the need for accurate species identifications which are required for biological monitoring programs. DNA-barcoding may speed up the identification process, but success is dependent on the barcode library coverage and quality. Our gap analysis has shown that although a large number of BINs (indicating putative species) was recognized by BOLD still only 10 % of the amphipod species are represented by barcodes. Moreover, the majority of BINs is represent by a very low number of sequences that make them unreliable according to the quality control system. The geographical coverage is poor with vast areas of Africa, South America and the open ocean acting as “white gaps”, also the level of barcoding effort is skewed depending on the environment.

As such, we make the following recommendations (in order of priority), which will improve the data currently held within BOLD, and we outline steps that are needed to provide a more equal coverage of the sequence data within the Amphipoda, and thus improve the utility of the database for a variety of applications, including species identification and biomonitoring.

1. Morphological identification of the already recognised BINs (that are missing species ID) if the voucher specimens are available.

2. Analysis of the nominal species that are represented by more than one BIN, especially if identifications represented by different BINs were produced by separate working teams.

3. Prioritised barcoding of representatives from families that are known to be important and abundant components of communities; Phoxocephalidae, Ampeliscidae, Oedicerotidae, and Caprellidae should be prioritised.

4. Targeted sampling programs for taxa coming from geographical regions with the least knowledge.

5. Targeted sampling to obtain more sequences for taxa present in BOLD but represented by small numbers of sequences (especially singletons), from different parts of the species' range if possible.

6. Targeted programs to sequence type specimens stored in musea or to collect and study fresh individuals from type localities if types are unsuitable for analyses.

References

Arfianti T, Wilson S, Costello MJ. 2018. Progress in the discovery of amphipod crustaceans.

- 592 *PeerJ* 6:e5187. <https://doi.org/10.7717/peerj.5187>
- 593 **Aylagas E, Borja A, Muxika I, Rodríguez-Ezpeleta N. 2018.** Adapting metabarcoding based
- 594 benthic biomonitoring into routine marine ecological status assessment networks. *Ecological*
- 595 *Indicators* 95:194–202. <https://dx.doi.org/10.1016/j.ecolind.2018.07.044>
- 596 **Barnard JL, Karaman GS. 1991.** The families and genera of marine gammaridean Amphipoda
- 597 (except marine gammaroids). Part 1 & 2. *Records of the Australian Museum Supplement*
- 598 13:1–866.
- 599 **Bauzá-Ribot MM, Jaume D, Fornós JJ, Juan C, Pons J. 2011.** Islands beneath islands:
- 600 phylogeography of a groundwater amphipod crustacean in the Balearic archipelago. *BMC*
- 601 *Evolutionary Biology* 11(1):221. <https://doi.org/10.1186/1471-2148-11-221>
- 602 **Beermann J, Hall-Mullen AK, Havermans C, Coolen JW, Crooijmans RP, Dibbits B, Held**
- 603 **C, Desiderato A. 2020.** Ancient globetrotters—connectivity and putative native ranges of
- 604 two cosmopolitan biofouling amphipods. *PeerJ* 8:e9613 <https://doi.org/10.7717/peerj.9613>
- 605 **Bij de Vaate A, Jazdzewski K, Ketelaars HAM, Gollasch S, Van der Velde G. 2002.**
- 606 Geographical patterns in range extension of Ponto-Caspian macroinvertebrate species in
- 607 Europe. *Canadian Journal of Fisheries and Aquatic Science* 59:1159–1174.
- 608 <https://doi.org/10.1139/f02-098>
- 609 **Brandt A. 1993.** Composition, abundance, and diversity of peracarid crustaceans on a transect of
- 610 the Kolbeinsey Ridge, north of Iceland. *Polar Biology* 13:565–576.
- 611 <https://doi.org/10.1007/BF00236399>
- 612 **Brandt A, Alalykina I, Brix S, Brenke N, Błażewicz M, Golovan OA, Johannsen N, Hrinko**
- 613 **AM, Jazdzewska AM, Jeskulke K, Kamenev GM, Lavrenteva AV, Malyutina MV,**

Riehl T, Lins L. 2019. Depth zonation of deep-sea macrofauna of the Northwest Pacific.

Progress in Oceanography <https://doi.org/10.1016/j.pocean.2019.102131>

Brandt A, De Broyer C, De Mesel I, Ellingsen K, Gooday A, Hilbig B, Linse K, Thomson

MR, Tyler P. 2007. The biodiversity of the deep Southern Ocean benthos. *Philosophical*

Transactions of the Royal Society B Biological Sciences **362**:39–66.

<https://doi.org/10.1098/rstb.2006.1952>

Brante A, Guzmán-Rendón G, Barriá EM, Guillemin ML, Vera-Escalona I, Hernández

CE. 2019. Post-disturbance genetic changes: the impact of the 2010 mega-earthquake and

tsunami on Chilean sandy beach fauna. *Scientific Reports* **9(1)**:1–12.

<https://doi.org/10.1038/s41598-019-50525-1>

Brix S, Lörz A-N, Jazdzewska A, Hughes L, Tandberg AH, Pabis K, Stransky B, Krapp-

Schickel T, Sorbe J-C, Hendrycks E, Vader WJM, Frutos I, Horton T, Jazdzewski K,

Peart R, Beermann J, Coleman CO, Buhl-Mortensen L, Corbari L, Havermans C, Tato

R, Jimenez Campean A. 2018. Amphipod family distributions around Iceland. *Zookeys*

731:41–53. <https://doi.org/10.3897/zookeys.731.19854>

Brix S, Osborn KJ, Kaiser S, Truskey SB, Schnurr SM, Brenke N, Malyutina M, Martinez

Arbizu P. 2020. Adult life strategy affects distribution patterns in abyssal isopods –

implications for conservation in Pacific nodule areas. *Biogeosciences* **17**: 6163–6184.

<https://doi.org/10.5194/bg-17-6163-2020>

Bundschuh M, Zubrod JP, Klemm P, Elsaesser D, Stang C, Schulz R. 2013. Effects of peak

exposure scenarios on *Gammarus fossarum* using field relevant pesticide mixtures.

Ecotoxicological Environment Safety **95**:137–143

<https://doi.org/10.1016/j.ecoenv.2013.05.025>

- Bundschuh M, Zubrod JP, Seitz F, Stang C, Schulz R. 2011.** Ecotoxicological evaluation of three tertiary wastewater treatment techniques via meta-analysis and feeding bioassays using *Gammarus fossarum*. *Journal of hazardous materials* **192(2)**:772–778. <https://doi.org/10.1016/j.jhazmat.2011.05.079>
- Bush A, Compson ZG, Monk WA, Porter TM, Steeves R, Emilson EJ, Gagne N, Hajibabaei M, Roy M, Baird DJ. 2019.** Studying ecosystems with DNA metabarcoding: lessons from biomonitoring of aquatic macroinvertebrates. *Frontiers in Ecology and Evolution* **7** (434) <http://dx.doi.org/10.3389/fevo.2019.00434>
- Cabezas MP, Xavier R, Branco M, Santos AM, Guerra-Garcia JM. 2014.** Invasion history of *Caprella scaura* Templeton, 1836 (Amphipoda: Caprellidae) in the Iberian Peninsula: multiple introductions revealed by mitochondrial sequence data. *Biological Invasions* **16**:2221–2245. <https://doi.org/10.1007/s10530-014-0660-y>
- Coleman CO. 2015.** Taxonomy in Times of the Taxonomic Impediment – Examples from the Community of Experts on Amphipod Crustaceans. *Journal of Crustacean Biology* **35(6)**:729–740. <https://doi.org/10.1163/1937240X-00002381>
- Coleman CO, Radulovici AE. 2020.** Challenges for the future of taxonomy: talents, databases and knowledge growth. *Megataxa* **001(1)**:028–034. <https://doi.org/10.11646/megataxa.1.1.5>
- Copilaş-Ciocianu D, Sidorov D, Gontcharov A. 2019.** Adrift across tectonic plates: molecular phylogenetics supports the ancient Laurasian origin of old limnic crangonyctid amphipods. *Organisms, Diversity and Evolution* **19**:191–207. <https://doi.org/10.1007/s13127-019-00401-7>

- 658 **Corkum LD. 1989.** Patterns of benthic invertebrate assemblages in rivers of northwestern North
659 America. *Freshwater Biology* **21**:191–205. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2427.1989.tb01358.x)
660 [2427.1989.tb01358.x](https://doi.org/10.1111/j.1365-2427.1989.tb01358.x)
- 661 **Cowart DA, Pinheiro M, Mouchel O, Maguer M, Grall J, Miné J, Arnaud-Haond S. 2015.**
662 Metabarcoding Is Powerful yet Still Blind: A Comparative Analysis of Morphological and
663 Molecular Surveys of Seagrass Communities. *PLoS ONE* **10**(2):e0117562.
664 <https://doi.org/10.1371/journal.pone.0117562>
- 665 **Cristescu ME. 2014.** From barcoding single individuals to metabarcoding biological
666 communities: Towards an integrative approach to the study of global biodiversity. *Trends in*
667 *Ecology and Evolution* **29**(10):566–571. <http://dx.doi.org/10.1016/j.tree.2014.08.001>
- 668 **Dalpadado P, Borkner N, Bogstad B, Mehl S. 2001.** Distribution of *Themisto* (Amphipoda)
669 spp. in the Barents Sea and predator-prey interactions. *ICES Journal of Marine Sciences*
670 **58**:876–895. <https://doi.org/10.1006/jmsc.2001.1078>
- 671 **Dauby P, Nyssen F, De Broyer C. 2003.** Amphipods as food sources for higher trophic levels in
672 the Southern Ocean: a synthesis. In: Huiskes AHL, Gieskes WWC, Rozema J, Schorno
673 RML, van der Vies SM, Wolff WJ, editors. *Antarctic Biology in a Global Context*. Backhuys
674 Publishers, Leiden, the Netherlands, 129–134.
- 675 **Dauby P, Scailteur Y, De Broyer C. 2001.** Trophic diversity within the eastern Weddell Sea
676 community. *Hydrobiologia* **443**:69–86. <https://doi.org/10.1023/A:1017596120422>
- 677 **Delić T, Švara V, Coleman CO, Trontelj P, Fišer C. 2017a.** The giant cryptic amphipod
678 species of the subterranean genus *Niphargus* (Crustacea, Amphipoda). *Zoologica Scripta*
679 **46**(6):740–752. <https://doi.org/10.1111/zsc.12252>

Delić T, Trontelj P, Rendoš M, Fišer C. 2017b. The importance of naming cryptic species and the conservation of endemic subterranean amphipods. *Scientific Reports* **7(1)**:1–12.

<https://doi.org/10.1038/s41598-017-02938-z>

Desiderato A, Costa FO, Serejo CS, Abbiati M, Queiroga H, Vieira PE. 2019. Macaronesian islands as promoters of diversification in amphipods: The remarkable case of the family Hyalidae (Crustacea, Amphipoda). *Zoologica Scripta* **48**:359– 375.

<https://doi.org/10.1111/zsc.12339>

Ebach MC, Valdecasas AG, Wheeler QD. 2011. Impediments to taxonomy and users of taxonomy: accessibility and impact evaluation. *Cladistics* **27(5)**:550–557.

<https://doi.org/10.1111/j.1096-0031.2011.00348.x>

Feio MJ, Filipe AF, Garcia-Raventos A, Ardura A, Calapez AR, Pujante AM, Mortágua A, Múrria C, Diaz-de-Quijano D, Martins FMS, Duarte S, Sáinz Bariáin M, Cordeiro R, Rivera SF, Väisänen LOS, Fonseca A, Gonçalves V, Garcia-Vazquez E, Vieites Rodríguez D, Ivanova EA, Costa FO, Barquín J, Rojo V, Vierna J, Fais M, Suarez M, Nieminen M, Hammers-Wirtz M, Kolmakova OV, Trusova MY, Beja P, González R, Planes S, Almeida SFP. 2020. Advances in the use of molecular tools in ecological and biodiversity assessment of aquatic ecosystems. *Limnetica* **39(1)**:419–440.

<https://doi.org/10.23818/limn.39.27>

Fišer C, Konec M, Alther R., Švara V, Altermatt F. 2017. Taxonomic, phylogenetic and ecological diversity of *Niphargus* (Amphipoda: Crustacea) in the Hölloch cave system (Switzerland). *Systematics and Biodiversity* **15(3)**: 218-237

<https://doi.org/10.1080/14772000.2016.1249112>

- 702 **Fontes JT, Vieira PE, Ekrem T, Soares P, Costa FO. 2020.** BAGS: an automated Barcode,
703 Audit & Grade System for DNA barcode reference libraries. *Molecular Ecology Resources*.
704 <https://doi.org/10.1111/1755-0998.13262>
- 705 **Frutos I, Brandt A, Sorbe JC. 2017.** Deep-Sea Suprabenthic Communities: The Forgotten
706 Biodiversity. In: Rossi S, Bramanti L, Gori A, Orejas C, editors. *Marine Animal Forests*.
707 Springer International Publishing, Cham, p.475–503. [https://doi.org/10.1007/978-3-319-](https://doi.org/10.1007/978-3-319-21012-4_21)
708 [21012-4_21](https://doi.org/10.1007/978-3-319-21012-4_21)
- 709 **Frutos I, Sorbe JC. 2017.** Suprabenthic assemblages from the Capbreton area (SE Bay of Biscay):
710 faunal recovery after a canyon turbiditic disturbance. *Deep-Sea Research I* **130**:36–46.
711 <https://doi.org/10.1016/j.dsr.2017.10.007>
- 712 **Grabowski M, Wysocka A, Mamos T. 2017.** Molecular species delimitation methods provide
713 new insight into taxonomy of the endemic gammarid species flock from the ancient Lake
714 Ohrid. *Zoological Journal of the Linnean Society-London* **181**(2):272–285.
715 <https://doi.org/10.1093/zoolinnean/zlw025>
- 716 **Hajibabaei M, Spall JL, Shokralla S, van Konynenburg S. 2012.** Assessing biodiversity of a
717 freshwater benthic macroinvertebrate community through non-destructive environmental
718 barcoding of DNA from preservative ethanol. *BMC Ecology* **12**:28
719 <http://dx.doi.org/10.1186/1472-6785-12-28>
- 720 **Havermans C. 2016.** Have we so far only seen the tip of the iceberg? Exploring species
721 diversity and distribution of the giant amphipod *Eurythenes*. *Biodiversity* **17**:12–25
722 <https://doi.org/10.1080/14888386.2016.1172257>

- 723 **Havermans C, Hagen W, Zeidler W, Held C, Auel H. 2019.** A survival pack for escaping
- 724 predation in the open ocean: amphipod–pteropod associations in the Southern Ocean. *Marine*
- 725 *Biodiversity* **49(3)**:1361–1370. <https://doi.org/10.1007/s12526-018-0916-3>
- 726 **Havermans C, Smetacek V. 2018.** Bottom-up and top-down triggers of diversification: A new
- 727 look at the evolutionary ecology of scavenging amphipods in the deep sea. *Progress in*
- 728 *Oceanography* **164**:37–51. <https://doi.org/10.1016/j.pocean.2018.04.008>
- 729 **Hestetun JT, Bye-Ingebrigtsen E, Nilsson RH, Glover AG, Johansen PO, Dahlgren TG.**
- 730 **2020.** Significant taxon sampling gaps in DNA databases limit the operational use of marine
- 731 macrofauna metabarcoding. *Marine Biodiversity* **50(5)**:1–9. [https://doi.org/10.1007/s12526-](https://doi.org/10.1007/s12526-020-01093-5)
- 732 [020-01093-5](https://doi.org/10.1007/s12526-020-01093-5)
- 733 **Horton T, Lowry J, De Broyer C, Bellan-Santini D, Coleman CO, Corbari L, Costello M J,**
- 734 **Daneliya M, Dauvin J-C, Fišer C, Gasca R, Grabowski M, Guerra-García JM,**
- 735 **Hendrycks E, Hughes L, Jaume D, Jazdzewski K, Kim Y-H, King R, Krapp-Schickel T,**
- 736 **LeCroy S, Lörz A-N, Mamos T, Senna AR, Serejo C, Sket B, Souza-Filho JF, Tandberg**
- 737 **AH, Thomas J, Thurston M, Vader W, Väinölä R, Vonk R, White K, Zeidler W. 2020.**
- 738 World Amphipoda Database. Accessed at <http://www.marinespecies.org/amphipoda> on 17-
- 739 07-2020.
- 740 **Horton, T. Marsh, L., Bett, B.J., Gates, A.R., Jones, D.O.B., Benoist, N.M.A., Pfeifer, S.**
- 741 **Simon-Lledó, E. Durden, J., Vandepitte, L., Appeltans, W. 2021.** Recommendations for
- 742 the standardisation of open taxonomic nomenclature for image-based identifications.
- 743 *Frontiers in Marine Sciences, Deep-Sea Environments and Ecology*
- 744 <https://doi.org/10.3389/fmars.2021.620702>

- Hou Z, Fu J, Li S. 2007.** A molecular phylogeny of the genus *Gammarus* (Crustacea: Amphipoda) based on mitochondrial and nuclear gene sequences. *Molecular Phylogenetics and Evolution* **45**:596–611. <https://doi.org/10.1016/j.ympev.2007.06.006>
- Hou Z, Sket B, Li S. 2014.** Phylogenetic analyses of Gammaridae crustacean reveal different diversification patterns among sister lineages in the Tethyan region. *Cladistics* **30(4)**:352–365. <https://doi.org/10.1111/cla.12055>
- Humphries P, Davies PE Mulcahy ME. 1996.** Macroinvertebrate assemblages of littoral habitats in the Macquarie and Mersey rivers, Tasmania: Implications for the management of regulated rivers. *Regulated Rivers: Research & Management* **12**:99–122. doi:[10.1002/\(SICI\)1099-1646\(199601\)12:1<99::AID-RRR382>3.0.CO;2-1](https://doi.org/10.1002/(SICI)1099-1646(199601)12:1<99::AID-RRR382>3.0.CO;2-1)
- Hyne RV, Everett DA. 1998.** Application of a benthic euryhaline amphipod, *Corophium* sp., as a sediment toxicity testing organism for both freshwater and estuarine systems. *Archives of Environmental Contamination and Toxicology* **34(1)**:26–33 <https://doi.org/10.1007/s002449900282>
- IPBES. 2019.** Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. Díaz S, Settele J, Brondízio ES, Ngo HT, Guèze M, Agard J, Arneth A, Balvanera P, Brauman KA, Butchart SHM, Chan KMA, Garibaldi LA, Ichii K, Liu J, Subramanian SM, Midgley GF, Miloslavich P, Molnár Z, Obura D, Pfaff A, Polasky S, Purvis A, Razzaque J, Reyers B, Roy Chowdhury R, Shin YJ, Visseren-Hamakers IJ, Willis KJ, Zayas CN, editors. IPBES secretariat, Bonn, Germany; 2019.
- Jazdzewska A. 2015.** Kuril–Kamchatka deep sea revisited – insights into the amphipod abyssal fauna. *Deep-Sea Research II* **111**:294–300. <https://doi.org/10.1016/j.dsr2.2014.08.008>

- Jazdzewska AM, Corbari L, Driskell A, Frutos I, Havermans C, Hendrycks E, Hughes L, Lörz A-N, Stransky B, Tandberg AHS, Vader W, Brix S. 2018.** A genetic fingerprint of Amphipoda from Icelandic waters – the baseline for further biodiversity and biogeography studies. *Zookeys* **731**:55–73. <https://doi.org/10.3897/zookeys.731.19931>
- Jazdzewska AM, Mamos T. 2019.** High species richness of Northwest Pacific deep-sea amphipods revealed through DNA barcoding. *Progress in Oceanography* <https://doi.org/10.1016/j.pocean.2019.102184>
- Jazdzewska AM, Rewicz T, Mamos T, Wattier R, Bącela-Spychalska K, Grabowski M. 2020.** Cryptic diversity and mtDNA phylogeography of the invasive demon shrimp, *Dikerogammarus haemobaphes* (Eichwald, 1841), in Europe. *NeoBiota* **57**:53–86. <https://doi.org/10.3897/neobiota.57.46699>
- Jazdzewski K, De Broyer C, Pudlarz M, Zielinski D. 2001.** Seasonal fluctuations of vagile benthos in the uppermost sublittoral of a maritime Antarctic fjord. *Polar Biology* **24**:910–917. <https://doi.org/10.1007/s003000100299>
- Kaiser S, Barnes DK, Brandt A. 2007.** Slope and deep-sea abundance across scales: Southern Ocean isopods show how complex the deep sea can be. *Deep-Sea Research II* **54**:1776–1789. <https://doi.org/10.1016/j.dsr2.2007.07.006>
- Kelly DW, Muirhead JR, Heath DD, MacIsaac HJ. 2006.** Contrasting patterns in genetic diversity following multiple invasions of fresh and brackish waters. *Molecular Ecology* **15**:3641–3655. <https://doi.org/10.1111/j.1365-294X.2006.03012.x>
- Knox MA, Hogg ID, Pilditch CA. 2011.** The role of vicariance and dispersal on New Zealand's estuarine biodiversity: the case of *Paracorophium* (Crustacea: Amphipoda). *Biological*

- 790 *Journal of the Linnean Society* **103(4)**:863–874. <https://doi.org/10.1111/j.1095->
- 791 [8312.2011.01675.x](https://doi.org/10.1111/j.1095-8312.2011.01675.x)
- 792 **Knox MA, Hogg ID, Pilditch CA, Lörz AN, Hebert PDN, Steinke D. 2012.** Mitochondrial
- 793 DNA (COI) analyses reveal that amphipod diversity is associated with environmental
- 794 heterogeneity in deep-sea habitats. *Molecular Ecology* **21**:4885–4897
- 795 <https://doi.org/10.1111/j.1365-294X.2012.05729.x>
- 796 **Leese F, Bouchez A, Abarenkov K, Altermatt F, Borja A, Bruce K, Torbjörn E, Čiampor**
- 797 **F, Čiamporová Z, Zatošová Z, Costa FO, Duarte S, Elbrecht V, Fontaneto D, Franc A,**
- 798 **Geiger MF, Hering D, Kahlert M, Stroil BK, Kelly M, Keskin E, Liska I, Mergen P,**
- 799 **Meissner K, Pawlowski J, Penev L, Reyjol Y, Rotter A, Steinke D, van der Wal B,**
- 800 **Vitecek S, Zimmermann J, Weigand AM. 2018.** Why we need sustainable networks
- 801 bridging countries, disciplines, cultures and generations for aquatic biomonitoring 2.0: a
- 802 perspective derived from the DNAqua-Net COST action. *Advances in Ecological Resources*
- 803 **58**:63–99 <http://dx.doi.org/10.1016/bs.aecr.2018.01.001>
- 804 **Leite BR, Vieira PE, Teixeira MAL, Lobo-Arteaga J, Hollatz C, Borges LMS, Duarte S,**
- 805 **Troncoso JS, Costa FO. 2020.** Gap-analysis and annotated reference library for supporting
- 806 macroinvertebrate metabarcoding in Atlantic Iberia. *Regional Studies in Marine Science*
- 807 **10**:1307. <https://doi.org/10.1016/j.rsma.2020.101307>
- 808 **Lörz A-N, Jazdzewska AM, Brandt A. 2018.** A new predator connecting the abyssal with the
- 809 hadal in the Kuril-Kamchatka Trench, NW Pacific. *PeerJ* **6**:e4887.
- 810 <https://doi.org/10.7717/peerj.4887>
- 811 **Major K, Soucek DJ, Giordano R, Wetzel MJ, Soto-Adames F. 2013.** The common
- 812 ecotoxicology laboratory strain of *Hyalella azteca* is genetically distinct from most wild

strains sampled in eastern North America. *Environmental toxicology and chemistry*

32(11):2637–2647. <https://doi.org/10.1002/etc.2355>

Mamos T, Wattier R, Burzyński A, Grabowski M. 2016. The legacy of a vanished sea: a high level of diversification within a European freshwater amphipod species complex driven by 15 My of Paratethys regression. *Molecular Ecology* **3**:795–810.

<https://doi.org/10.1111/mec.13499>

Mamos T, Wattier R, Majda A, Sket B, Grabowski M. 2014. Morphological vs. molecular delineation of taxa across montane regions in Europe: the case study of *Gammarus balcanicus* Schäferna, 1922 (Crustacea: Amphipoda). *Journal of Zoological Systematics and Evolutionary Research* **52(3):237–248.** <https://doi.org/10.1111/jzs.12062>

Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B. 2011. How Many Species Are There on Earth and in the Ocean? *PLoS Biology* **9(8):e1001127.**

<https://doi.org/10.1371/journal.pbio.1001127>

Múrria C, Väisänen LO, Somma S, Wangensteen OS, Arnedo MA, Prat N. 2020. Towards an Iberian DNA barcode reference library of freshwater macroinvertebrates and fishes.

Limnetica **39(1):73–92.** <http://dx.doi.org/10.23818/limn.39.06>

Oliveira LM, Knebelsberger T, Landi M, Soares P, Raupach MJ, Costa FO. 2016.

Assembling and auditing a comprehensive DNA barcode reference library for European marine fishes. *Journal of Fish Biology* **89(6):2741–2754.** <http://dx.doi.org/10.1111/jfb.13169>

Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. 2021. The PRISMA 2020

statement: an updated guideline for reporting systematic reviews. *BMJ* **372**:n71.

<http://doi.org/10.1136/bmj.n71>

Page RDM. 2016. DNA barcoding and taxonomy: dark taxa and dark texts. *Philosophical Transactions of the Royal Society B* **371**:20150334. <https://doi.org/10.1098/rstb.2015.0334>

QGIS Development Team. 2018. QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>.

Radulovici AE, Coleman CO. 2017. Reconciling large molecular datasets, bioinformatics and taxonomy: prospects for Amphipoda. *Biodiversity Journal* **8(2)**:633–634.

Ratnasingham S, Hebert P. 2007. BOLD: The Barcode of Life Data System. *Molecular Ecology Notes* **7(3)**:355–364 <https://doi.org/10.1111/j.1471-8286.2007.01678.x>

Ratnasingham S, Hebert PDN. 2013. A DNA-based registry for all animal species: the barcode index number (BIN) system. *PLoS ONE* **8**:e66213 <https://doi.org/10.1371/journal.pone.0066213>

Rewicz T, Wattier R, Grabowski M, Rigaud T, Bącela-Spychalska K. 2015. Out of the Black Sea: Phylogeography of the Invasive Killer Shrimp *Dikerogammarus villosus* across Europe. *PLoS ONE* **10(2)**:e0118121. <https://doi.org/10.1371/journal.pone.0118121>

Riehl T, Brenke N, Brix S, Driskell A, Kaiser S, Brandt A. 2014. Field and Laboratory Methods for DNA Studies on Deep-sea Isopod Crustaceans. *Polish Polar Research* **35**:203–224. <https://doi.org/10.2478/popore-2014-0018>

Ros M, Guerra-García JM, González-Macías M, Saavedra Á, López-Fe CM. 2013. Influence of fouling communities on the establishment success of alien caprellids (Crustacea: Amphipoda) in Southern Spain. *Marine Biology Research* **9(3)**:261–273. <https://doi.org/10.1080/17451000.2012.739695>

- Ros M, Vázquez-Luis M, Guerra-García JM. 2013.** The role of marinas and recreational boating in the occurrence and distribution of exotic caprellids (Crustacea: Amphipoda) in the Western Mediterranean: Mallorca Island as a case study. *Journal of Sea Research* **83**:94–103. <https://doi.org/10.1016/j.seares.2013.04.004>
- Sayers EW, Cavanaugh M, Clark K, Ostell J, Pruitt KD, Karsch-Mizrachi I. 2020.** GenBank. *Nucleic Acids Research* **47**:D94–D93. <https://doi.org/10.1093/nar/gkz956>.
- Sedano F, Navarro-Barranco C, Guerra-García JM, Espinosa F. 2020.** From sessile to vagile: Understanding the importance of epifauna to assess the environmental impacts of coastal defence structures. *Estuarine, Coastal and Shelf Sciences* **235**:106616. <https://doi.org/10.1016/j.ecss.2020.106616>
- Sigovini M., Keppel E., Tagliapietra D. 2016.** Open nomenclature in the biodiversity era. *Methods in Ecology and Evolution* **7**:10. <https://doi.org/10.1111/2041-210X.12594>
- Tomikawa K, Kyono M, Kuribayashi K, Nakano T. 2017.** The enigmatic groundwater amphipod *Awacaris kawasawai* revisited: synonymisation of the genus *Sternomoera*, with molecular phylogenetic analyses of *Awacaris* and *Sternomoera* species (Crustacea: Amphipoda: Pontogeneiidae). *Invertebrate Systematics* **31**(2):125–140. <https://doi.org/10.1071/IS16037>
- Väinölä R, Witt JDS, Grabowski M, Bradbury JH, Jazdzewski K, Sket B. 2008.** Global diversity of amphipods (Amphipoda; Crustacea) in freshwater. *Hydrobiologia* **595**:241–255. <https://doi.org/10.1007/s10750-007-9020-6>
- Verheye ML, Backeljau T, d’Udekem d’Acoz C. 2016.** Looking beneath the tip of the iceberg: diversification of the genus *Epimeria* on the Antarctic shelf (Crustacea, Amphipoda). *Polar Biology* **39**(5):925–945. <https://doi.org/10.1007/s00300-016-1910-5>

- 882 **Vinogradov ME, Volkov AF, Semenova TN. 1996.** Hyperiid Amphipods (Amphipoda,
883 Hyperiidea) of the World Oceans. Smithsonian Institution Libraries, Washington, DC.
- 884 **Wattier, R., Mamos, T., Copilaş-Ciocianu, D., Jelić M., Ollivier A., Chaumot A., Danger M,**
885 **Felten V., Piscart C., Žganec K., Rewicz T., Wysocka A., Rigaud T., Grabowski M.**
886 **2020.** Continental-scale patterns of hyper-cryptic diversity within the freshwater model taxon
887 *Gammarus fossarum* (Crustacea, Amphipoda). *Scientific Reports* **10**:16536.
888 <https://doi.org/10.1038/s41598-020-73739-0>
- 889 **Weigand H, Beermann AJ, Čiampor F, Costa FO, Csabai Z, Duarte S, Geiger MF,**
890 **Grabowski M, Rimet F, Rulik B, Strand M, Szucsich N, Weigand AM, Willassen E,**
891 **Wyler SA, Bouchez A, Borja A, Čiamporová Z, Ferreira S, Dijkstra KB,**
892 **Eisendle U, Freyhof J, Gadawski P, Graf W, Haegerbaeumer A, van der Hoorn BB,**
893 **Japoshvili B, Keresztes L, Keskin E, Lesse F, Macher JN, Mamos T, Paz G, Pešić V,**
894 **Pfannkuchen DM, Pfannkuchen MA, Price BW, Rinkevich B, Teixeira MAL, Várбірó**
895 **G, Ekrem T. 2019.** DNA barcode reference libraries for the monitoring of aquatic biota in
896 Europe: Gap-analysis and recommendations for future work. *Science of the Total*
897 *Environment* **678**:499–524. <http://dx.doi.org/10.1016/j.scitotenv.2019.04.247>
- 898 **Weisshappel JBF, Svavarsson J. 1998.** Benthic amphipods (Crustacea: Malacostraca) in
899 Icelandic waters: diversity in relation to faunal patterns from shallow to intermediate deep
900 Arctic and North Atlantic Oceans. *Marine Biology* **131**:133–142.
- 901 **Witt JD, Threlloff DL, Hebert PD. 2006.** DNA barcoding reveals extraordinary cryptic
902 diversity in an amphipod genus: implications for desert spring conservation. *Molecular*
903 *Ecology* **15(10)**:3073–3082. <https://doi.org/10.1111/j.1365-294X.2006.02999.x>

- Wysocka A, Grabowski M, Sworobowicz L, Burzyński A, Kilikowska A, Kostoski G, Sell J. 2013.** A tale of time and depth: intralacustrine radiation in endemic *Gammarus* species flock from the ancient Lake Ohrid. *Zoological Journal of the Linnean Society-London* **167(3)**:345–359. <https://doi.org/10.1111/j.1096-3642.2012.00878.x>
- Wysocka A, Grabowski M, Sworobowicz L, Mamos T, Burzyński A, Sell J. 2014.** Origin of the Lake Ohrid gammarid species flock: ancient local phylogenetic lineage diversification. *Journal of Biogeography* **41(9)**:1758–1768 <https://doi.org/10.1111/jbi.12335>

Figure captions

Figure 1. PRISMA 2020 work-flow diagram (Page et al., 2021). Summary of the data download, identification and screening before analysis. All record removals were done by the leading author of the paper.

Figure 2. Environmental origin of the amphipod records (A) and BINs (B) in BOLD database.

Figure 3. Geographic distribution of amphipod records expressed by sequences present in BOLD (A – freshwater, B – marine, C – terrestrial). Dots indicate records with exact coordinates, for records without latitude and longitude the country of origin was checked. Background color of the country indicates this number per country.

Figure 4. Proportion of records (A) and BINs (B) with different level of identification within freshwater, marine and terrestrial amphipod taxa.

Figure 5. Number of BINs represented by given number of sequences. Upper set (A, B, C) – all BINs, lower set (D, E, F) – only BINs with complete species-level identification considered. A, D – freshwater, B, E – marine, C, F – terrestrial taxa.

Figure 6. Number of nominal species represented by given number of BINs.

926

927 **Supplemental files list**

928

929 **Supplemental file 1**

930 - File format .xlsx

931 - Title: List of the doubled records indentified in the original dataset

932 - Description: The file consists of records that appeared to be doubled in the BOLD database.

933

934 **Supplemental file 2**

935 - File format .xlsx

936 - Title: The list of records analysed (after removal of doubled records).

937 - Description: The file consists of all records that was basis of the present study. The colors of
938 the cell of recordID indicates the environment: green - marine (including brakishwater and
939 fully marine taxa), red - freshwater (including freshwater and brakishwater taxa), yellow -
940 terrestrial, blue - environment not recorded.

941

942 **Supplemental file 3**

943 - File format .xlsx

944 - Title: List of BINs possessing more than one ID variant with notes on the identification

945 - Description: The file presents BINs that have received different identifications with details of
946 the ID and comments concerning the final identification used in the study.

947

948 **Supplemental file 4**

949 - File format .xlsx

950 - Title: List of BINs for which the environment was not able to be assessed.

951 - Description: The file presents BINs for which the available data did not allow to specify the
952 environment.

953

954 **Supplemental file 5**

955 - File format .xlsx

- Title: List of BINs with the largest number of records (blue - freshwater, green - marine, yellow - terrestrial taxa).

- Description: The file presents BINs that are represented by the largest number of records. The information about environmental origin of associated species are provided.

Supplemental file 6

- File format .xlsx

- Title: Nominal species with the largest number of BINs identified.

- Description: The file presents nominal species for which the largest number of BINs has been identified. Environmental origin of species is also provided.

Supplemental file 7

- File format .xlsx

- Title: List of amphipod families with number of accepted and barcoded species as well as information of the barcoding coverage within family. Families within each category with the highest barcoding coverage indicated in bold.

Supplemental file 8

- File format .docx

- PRISMA 2020 checklist

Table 1 (on next page)

Representation of amphipod families in BOLD.

* in parentheses the number of families without barcoded species but with at least one BIN identified to the genus (g) or family (f) level.

- 1 Table 1. Representation of amphipod families in BOLD. * in parentheses the number of families
- 2 without barcoded species but with at least one BIN identified to the genus (g) or family (f) level.

| Number of families | |
|---------------------------------|------------------|
| without any barcoded species | 117 (+ 13g, 5f)* |
| with up to 10% barcoded species | 47 |
| with 11-20% barcoded species | 24 |
| with 21-50% barcoded species | 24 |
| with >50% barcoded species | 10 |

3

4

Table 2(on next page)

Number of amphipod species in each realm with indication of their barcode quality according to grading system from Fontes et al. (2020).

A – consolidated concordance, B – basal concordance, C – multiple BINs for single morphospecies, D – insufficient data; for more detailed explanation of grading system, see Material and methods section.

1 Table 2. Number of amphipod species in each realm with indication of their barcode quality
 2 according to grading system from Fontes et al. (2020). A – consolidated concordance, B – basal
 3 concordance, C – multiple BINs for single morphospecies, D – insufficient data; for more
 4 detailed explanation of grading system, see Material and methods section.

| | A | B | C | D | all species |
|------------------|-----|-----|-----|-----|-------------|
| All species | 100 | 155 | 276 | 468 | 999 |
| Freshwater spp. | 31 | 55 | 140 | 225 | 451 |
| Marine spp. | 58 | 92 | 120 | 226 | 496 |
| Terrestrial spp. | 11 | 8 | 16 | 17 | 52 |

5

6

Table 3(on next page)

Number of accepted families and species of Amphipoda (according to WAD accessed on 17-07-2020), number of families with representation in BOLD, number of species present in BOLD and mean coverage of barcodes in amphipod families represented in BOLD.

- 1 Table 3. Number of accepted families and species of Amphipoda (according to WAD accessed
- 2 on 17-07-2020), number of families with representation in BOLD, number of species present in
- 3 BOLD and mean coverage of barcodes in amphipod families represented in BOLD.

| | No. of families | No. of species | No. of families with species representation in BOLD | No. of species present in BOLD | Mean barcode coverage [%] of those families with representation in BOLD |
|---|-----------------|----------------|---|--------------------------------|---|
| Very species rich families (>100 spp.) | 33 | 7302 | 32 | 714 | 8 |
| Species rich families (31-100 spp.) | 30 | 1633 | 22 | 127 | 10 |
| Moderately species rich families (11-30 spp.) | 53 | 979 | 26 | 107 | 21 |
| Low species rich families (<10 spp.) | 123 | 416 | 26 | 51 | 49 |

4

5

Table 4(on next page)

Percent of families with species belonging to different quality grading categories (Fontes et al., 2020).

A – consolidated concordance, B – basal concordance, C – multiple BINs for single morphospecies, D – insufficient data; for more detailed explanation of grading system, see Material and methods section.

1 Table 4. Percent of families with species belonging to different quality grading categories
 2 (Fontes et al., 2020). A – consolidated concordance, B – basal concordance, C – multiple BINs
 3 for single morphospecies, D – insufficient data; for more detailed explanation of grading system,
 4 see Material and methods section.

| | % of families | | | | |
|---------------------------------------|---------------|--|---|---|--|
| | All families | Very species rich families (>100 spp.) | Species rich families (31-100 spp.) | Moderately species rich families (11-30 spp.) | Low species rich families (<10 spp.) |
| At least one sp. in the category A | 32.4 | 65.6 | 31.8 | 16 | 7.7 |
| At least one sp. in the category B | 31.4 | 28.1 | 36.4 | 28 | 34.6 |
| At least one sp. in the category C | 10.5 | 0 | 13.6 | 24 | 7.7 |
| At least one sp. in the category D | 25.7 | 6.3 | 18.2 | 32 | 50 |
| Number of families | 105 | 32 | 22 | 25 | 26 |

5

6

Figure 1

PRISMA 2020 work-flow diagram (Page et al., 2021).

Summary of the data download, identification and screening before analysis. All record removals were done by the leading author of the paper.

Figure 1. PRISMA 2020 work-flow diagram (Page et al., 2021). Summary of the data download, identification and screening before analysis. All record removals were done by the leading author of the paper.




Figure 2

Environmental origin of the amphipod records (A) and BINs (B) in BOLD database.




Figure 3

Geographic distribution of amphipod records expressed by sequences present in BOLD (A - freshwater, B - marine, C - terrestrial).

Dots indicate records with exact coordinates, for records without latitude and longitude the country of origin was checked. Background color of the country indicates this number per country.




Figure 4

Proportion of records (A) and BINs (B) with different level of identification within freshwater, marine and terrestrial amphipod taxa.




Figure 5

Number of BINs represented by given number of sequences.

Upper set (A, B, C) - all BINs, lower set (D, E, F) - only BINs with complete species-level identification considered. A, D - freshwater, B, E - marine, C, F - terrestrial taxa.




Figure 6

Number of nominal species represented by given number of BINs.

