

Multiple virulence factors regulated by AlgU contribute to the pathogenicity of *Pseudomonas savastanoi* pv. *glycinea* in soybean

Viet Tru Nguyen^{Equal first author, 1, 2}, Nanami Sakata^{Equal first author, 2}, Giyu Usuki², Takako Ishiga², Yoshiteru Hashimoto^{2, 3}, Yasuhiro Ishiga^{Corresp. 2}

¹ Western Highlands Agriculture and Forestry Science Institute, Buon Ma Thuot, Daklak, Vietnam

² Faculty of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Ibaraki, Japan

³ Microbiology Research Center for Sustainability (MiCS), University of Tsukuba, Tsukuba, Ibaraki, Japan

Corresponding Author: Yasuhiro Ishiga

Email address: ishiga.yasuhiro.km@u.tsukuba.ac.jp

Pseudomonas savastanoi pv. *glycinea* (*Psg*) causes bacterial blight of soybean. To identify candidate virulence factors, transposon-mediated mutational analysis of *Psg* was carried out. We syringe-inoculated soybean leaves with *Psg* transposon mutants and identified 28 mutants which showed reduced virulence from 1,000 mutants screened. Next, we spray-inoculated soybean leaves with these mutants and demonstrated that the *algU* mutant showed significantly reduced virulence together with reduced bacterial populations *in planta*. Expression profiles comparison between the *Psg* wild-type (WT) and *algU* mutant in HSC broth revealed that expression of coronatine (COR)-related genes (including *cmaA* and *corR*) were down-regulated in the *algU* mutant compared with *Psg* WT. Moreover, we also showed that COR production were reduced in the *algU* mutant compared with WT. We also demonstrated that *algD*, which is related to alginate biosynthesis, showed reduced expression and biofilm formation was significantly suppressed in the *algU* mutant. Furthermore, *hrpL* also showed less expression in the *algU* mutant. These results indicate that AlgU plays a critical role in promoting *Psg* pathogenesis by regulating multiple virulence factors.

Multiple virulence factors regulated by AlgU contribute to the pathogenicity of *Pseudomonas savastanoi* pv. *glycinea* in soybean

Viet Tru Nguyen^{1,2*}, Nanami Sakata^{1*}, Giyu Usuki¹, Takako Ishiga¹, Yoshiteru Hashimoto^{1,3} and Yasuhiro Ishiga¹

¹Faculty of Life and Environmental Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8572, Japan.

²Western Highlands Agriculture and Forestry Science Institute, 53 Nguyen Luong Bang Street, Buon Ma Thuot City, Daklak Province, Vietnam.

³Microbiology Research Center for Sustainability (MiCS), University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8572, Japan.

*These authors contributed equally to this work.

Corresponding Author:

Yasuhiro Ishiga¹

1-1-1 Tennodai, Tsukuba, Ibaraki 305-8572, Japan.

Email address: ishiga.yasuhiro.km@u.tsukuba.ac.jp

Abstract

Pseudomonas savastanoi pv. *glycinea* (*Psg*) causes bacterial blight of soybean. To identify candidate virulence factors, transposon-mediated mutational analysis of *Psg* was carried out. We syringe-inoculated soybean leaves with *Psg* transposon mutants and identified 28 mutants which showed reduced virulence from 1,000 mutants screened. Next, we spray-inoculated soybean leaves with these mutants and demonstrated that the *algU* mutant showed significantly reduced virulence together with reduced bacterial populations *in planta*. Expression profiles comparison between the *Psg* wild-type (WT) and *algU* mutant in HSC broth revealed that expression of coronatine (COR)-related genes (including *cmaA* and *corR*) were down-regulated in the *algU* mutant compared with *Psg* WT. Moreover, we also showed that COR production were reduced in the *algU* mutant compared with WT. We also demonstrated that *algD*, which is related to alginate biosynthesis, showed reduced expression and biofilm formation was significantly suppressed in the *algU* mutant. Furthermore, *hrpL* also showed less expression in the *algU* mutant. These results indicate that AlgU plays a critical role in promoting *Psg* pathogenesis by regulating multiple virulence factors.

Introduction

Pseudomonas savastanoi pv. *glycinea* (*Psg*) causes bacterial blight of soybean. The disease is characterized by circular necrotic lesions on leaves surrounded by a chlorotic halo (Ignjatov et al., 2007). In *P. syringae*, *P. cannabina*, and *P. savastanoi*, the phytotoxin Coronatine (COR) is important in inducing chlorosis, and contributes to bacterial growth and lesion formation (Bender et al., 1987; Budde and Ullrich, 2000; Peñaloza-Vázquez et al., 2000; Uppalapati et al., 2005; Qi et al., 2011; Sakata et al., 2021). In *Psg* PG4180, COR synthesis genes reside on a 90 kb plasmid designated p4180A (Bender et al., 1991), with a 32 kb COR gene cluster which consists of two distinct regions encoding coronafacic acid (CFA) and coronamic acid (CMA) (Bender, 1999). *Psg* produces COR not only for the biological fitness of pathogens in planta (Ullrich et al., 1993) but also *in vitro* (Hoitink and Sinden, 1970; Palmer and Bender, 1993). Many studies showed the roles of COR in chlorosis, promoting lesion formation, and suspension of both stomatal and salicylic acid (SA)-dependent defenses (Peñaloza-Vázquez et al., 2000; Kloeck et al., 2001; Zhao et al., 2003; Brooks et al., 2005; Melotto et al., 2006; Uppalapati et al., 2007). COR contributes to *P. syringae* pv. *tomato* (*Pst*) DC3000 virulence by suppressing the host defense response (Uppalapati et al., 2007). COR suppresses pathogen-associated molecular pattern (PAMP)-triggered immunity (PTI), especially stomatal-based defense in the early *Pst* DC3000 infection stage in *Arabidopsis thaliana* and tomato (Melotto et al., 2006; Ishiga et al., 2018).

Besides COR, the type three secretion system (T3SS) also plays a critical role in *P. syringae* virulence. The T3SS, is encoded by the *hrp* (hypersensitive response and pathogenicity) cluster, and transfers type three effectors (T3Es) into plant cells to suppress PTI, contribute to pathogenesis, and enhance disease symptoms and bacterial multiplication (Brooks et al., 2005; Lam et al., 2014). Furthermore, *Psg* also produces pectolytic enzymes which allow the pathogen to invade and multiply in the intercellular spaces of host tissues. This is the physiological capability of *Psg* in using plant polysaccharides and providing flexibility for its pathogen activity (Haeefelet and Lindow, 1987).

AlgU, an extracytoplasmic function (ECF) sigma factor, is also important in supporting *P. syringae* growth and disease development (Markel et al., 2016). AlgU regulates between 800 to 1,000 genes (Yu et al., 2014), and importantly contributes to virulence gene regulation as well as flagellin repression (Schreiber and Desveaux, 2011; Markel et al., 2016; Bao et al., 2020). *Pst* DC3000 AlgU is not only able to regulate gene expression associated with T3SEs and the phytotoxin COR, but also alginate biosynthesis (Ishiga et al., 2018). Moreover, AlgU (previously called AlgT) also induces transcription of the *algT-mucAB* gene cluster and the *algD* operon, which are responsible for alginate biosynthesis in *Psg* PG4180 (Schenk et al., 2006). Although AlgU has been extensively studied in several *P. syringae* pathovars, its roles in *Psg* pathogenicity have not been elucidated yet.

To identify genes related to *P. syringae* pathogenicity, researchers carried out a screen for *P. syringae* mutants with reduced virulence. *Pst* DC3000 Tn5 mutants with reduced virulence on *A. thaliana* found the crucial functions of COR in virulence (Brooks et al., 2004). Sakata et al., (2019) also screened for *P. cannabina* pv. *alisalensis* (*Pcal*) KB211 Tn5 mutants with reduced virulence on cabbage plants using a dip-inoculation method, and identified multiple virulence

factors including the T3SS, membrane transporters, transcription factors, and amino acid metabolism genes. Thus, it is tempting to speculate that each *P. syringae* pathovar has developed its own virulence factors. However, a screening study to identify *Psg* virulence factors has not been conducted previously.

To investigate *Psg* virulence factors, we constructed a bacterial mutant library based on transposon insertion in *Psg*, and screened for mutants with less or no chlorosis on soybean leaves after syringe-inoculation. We successfully identified several virulence factors including COR, T3Es, and AlgU. Expression profiles revealed that AlgU promotes virulence in host plants not only by up-regulation of COR-related gene expression. We also showed that *algU* mutant showed reduced COR production and biofilm formation compared to WT. Our results provide evidence that AlgU plays a critical role in promoting *Psg* pathogenesis.

Materials & Methods

Bacterial strains, plasmids, and growth conditions

The bacterial strains and plasmids used in this study are listed in Table 1. *Pseudomonas* strains were routinely cultured on King's B (KB; King et al., 1954) medium or mannitol-glutamate (MG; Keane et al., 1970) medium at 28°C. *Escherichia coli* (*E. coli*) cultures were grown on Luria-Bertani (LB; Sambrook et al. 1989) medium at 37°C. The bacterial cell densities at 600 nm (OD₆₀₀) were measured using a Biowave CO8000 Cell Density Meter (Funakoshi, Tokyo, Japan) as described in Sakata et al., (2021).

Bacterial *in vitro* Growth Measurements

Wild-type, the *algU* mutant, and the *algU*-complemented strain were grown at 28°C on LB medium. The bacterial suspensions were standardized to an OD₆₀₀ of 0.05 with LB, and bacterial growth was measured at OD₆₀₀ for 6, 9, and 12 h.

Plant material and inoculation procedures

Soybean plants (*Glycine max*), cultivar “Enrei”, were grown in a growth chamber at 22°C, with approximately 60% humidity, and a supplementary light intensity of 200–350 μmol/(m² sec) for a 14 h photoperiod. All soybean plants used for virulence studies were three or four-weeks-old.

Psg carrying Tn5 transposon was syringe-infiltrated into soybean leaves at an OD₆₀₀ of 0.1 (5 × 10⁷ CFU/ml) containing 0.02% Silwet L-77 (OSi Specialties Inc., Danbury, CT, USA). The disease symptoms were fully developed at 6 days post inoculation (dpi) (Fig. 1A). The mutants which showed different disease symptoms or virulence reduction in comparison to *Psg* WT were selected.

For spray inoculation, bacterial suspensions were applied to observe disease symptoms on whole soybean plants as described previously (Uppalapati et al. 2007). Plants were sprayed with a bacterial suspension at an OD₆₀₀ of 0.2 (1 × 10⁸ CFU/ml) in sterile distilled water containing 0.025% Silwet L-77 until runoff. After inoculation, plants were transferred to growth chambers at

28°C with approximately 90 to 100% humidity for 24 h in the dark before maintaining plants at approximately 70% humidity.

For syringe inoculation, bacteria were suspended at a final concentration at an OD₆₀₀ of 0.1 (5×10^7 CFU/ml), 0.01 (5×10^6 CFU/ml), and 0.001 (5×10^5 CFU/ml), and infiltrated with a 1-ml blunt syringe into leaves. The plants were then incubated at 70-80% humidity for the rest of the experimental period. Leaves were removed and photographed at 5 dpi.

To measure bacterial growth in soybean leaves after spray inoculation, individual second leaf pairs were selected at 6 dpi, weighed and surface-sterilized in 5% H₂O₂ for 3 min, and then rinsed three times with sterile water. The leaves were then homogenized, and appropriate dilutions were plated on KB medium containing the appropriate antibiotics. The bacterial colony forming units (CFU) were normalized as CFU/g using the total inoculated leaf mass. The population at 0 dpi was estimated using leaves harvested 1 h post inoculation (hpi) without surface-sterilization. For syringe-inoculation, leaf discs were harvested using a 3.5 mm-diameter cork-borer from syringe-infiltrated leaf zone. The leaves were then homogenized, and appropriate dilutions were plated on KB medium containing the appropriate antibiotics. The bacterial colony forming units (CFU) were normalized as CFU/cm² using the leaf square meters. The bacterial populations were evaluated in at least three independent experiments.

Transposon-mediated mutagenesis and identification of mutated genes

Transposon-mediated mutagenesis was carried out, as described previously (Sakata et al. 2019). Briefly, pBSLC1 (Sawada et al., 2018) carrying mini-Tn5 transposon were transferred into *Psg* to build a mutant library. We developed more than 1,000 individual *Psg* mutant lines. After the inoculation assay, we identified the mutated genes by rescuing the transposon insertion sites into an *E. coli* plasmid and sequencing (Sakata et al., 2019).

Complementation of the *algU* mutant

The *algU*-complemented strain was constructed as described in Ishiga et al., (2018). Briefly, the pDSKG vector was made from pDSK519 vector (Keen et al., 1988) by replacing kanamycin cassette to gentamycin. The *algU* and promoter region were transferred into the pDSKG vector (Keen et al., 1988) to generate pDSKG-*algU*. The pDSKG-*algU* construct was introduced into the *algU* mutant by electrophoresis to generate the complemented strain.

Real-time quantitative RT-PCR

For *Psg* gene expression profiles, data were collected as previously described in Sakata et al., (2021). Specifically, bacteria were grown in HS medium optimized for COR production (HSC; Palmer and Bender, 1993) for 3 and 48 h. Bacterial RNA was extracted using the ReliaPrep RNA Cell Miniprep System Kit (Promega, WI, USA) according to the manufacture's protocol. Two micrograms of total RNA were treated with gDNA Remover (TOYOBO, Osaka, Japan) to eliminate genomic DNA, and the DNase-treated RNA was reverse transcribed using the ReverTra Ace qPCR RT Master Mix (TOYOBO). The cDNA (1:10) was then used for RT-qPCR using the primers shown in Supplementary Table S1 with THUNDERBIRD SYBR qPCR Mix (TOYOBO)

on a Thermal Cycler Dice Real Time System (TaKaRa). *Psg outer membrane lipoprotein I (oprI)* was used to normalize gene expression.

COR quantification by HPLC

Psg WT, the *algU* mutant, and the *algU*-complemented strain were cultured in HSC for 7 days. Culture supernatant was obtained by centrifugation ($12,000 \times g$ for 5 min). Cell pellets were dried at 65°C and weighed. The 500 μl of supernatants were extracted twice with 500 μl of ethyl acetate and 25 μl of HCl, and the organic phase was transferred to a new microcentrifuge tube. The sample was dried by centrifugal evaporator at 55°C , and the dried sample was dissolved with 0.05% trifluoroacetic acid (TFA)/acetonitrile (9:1, v/v). The culture supernatant was analyzed by HPLC with a Shimadzu LC20A system equipped with a Symmetry C8 column (4.6×250 mm; Waters Corporation, MA, USA) as described previously (Sakata et al. 2021).

Hypertrophy-inducing activity assay on potato tuber tissue

Potato tubers were cut from the central tuber portion to ensure samples of high uniformity. After washing in tap water for 5 min, each disc was washed with sterile distilled water several times. Potato tuber discs were inoculated using toothpicks by placing the tip in *Psg* WT, COR-defective mutants (*cfa6* and *cmaA*), the *algU* mutant, and the *algU*-complemented strain on a KB medium plate, and then placing the toothpick on the potato tuber disc. The discs were then placed at 23°C incubator (darkness) for 5 days. Photographs were taken at 5 dpi.

Biofilm formation assay

Biofilm formation was assayed as described previously (Shao et al., 2019). Briefly, the bacterial strains were incubated overnight in LB broth and resuspended in fresh LB broth to an OD_{600} of 0.1. Bacterial suspensions (120 μl) were put into 96-well plates and incubated at 28°C for 24, 48, 72, and 96 h. The bacterial solutions were discarded and washed three times with distilled water. The biofilm forming bacteria were treated with 150 μl of 0.1% crystal violet (CV; Fujifilm, Tokyo, Japan) for 20 min without shaking. The dye was discarded and washed twice with distilled water. The plate was dried completely, subsequently the biofilm was eluted with 150 μl of 100% ethanol, and the CV were dissolved completely. Finally, the eluted biofilm sample's absorbance was measured at OD_{595} .

Results

Identification and characterization of reduced virulence mutants

To identify *Psg* virulence genes, we screened 1,000 transposon insertion mutants for reduced disease symptoms on soybean leaves using the syringe-infiltration method. Disease symptoms caused by *Psg* WT showed a small water-soaked lesion surrounded by regions of chlorosis (Fig. 1A). A total of 28 mutants showed no or less chlorosis at 6 dpi (Fig. 1B). Seventeen mutants out of 28 had transposon insertions in genes encoding COR biosynthesis-related genes (Table 1). Soybean leaves inoculated with COR biosynthesis mutants (VTD29, VTE13, VTI15, and VTR4),

the *algU* mutant (VTO41), and an unknown-function mutant (VTF6) showed no chlorosis (Fig. 1B).

Reduced disease symptoms and bacterial growth in soybean

We identified the 28 mutants which showed reduced disease symptoms compared to *Psg* WT by syringe-inoculation (Fig. 1B). We further investigated whether these mutants also showed reduced virulence via spray-inoculation, and selected 2 mutants related to COR biosynthesis (*cmaA* and *cfa6*), and others. Soybean leaves inoculated with all mutants showed significantly reduced bacterial populations than those of *Psg* WT (Fig. 2A). Among all mutants, the *algU* mutant (VTO41) showed dramatically reduced bacterial populations and disease symptoms at 6 dpi (Figs. 2A and 2B). To confirm whether the altered *algU* mutant phenotype originates from a corresponding mutation, an *algU*-complemented strain was generated. *Psg* WT and the *algU*-complemented strain showed the same bacterial population levels as well as disease symptom development in soybean (Supplementary Figs. S1A and S1B). We also confirmed that *algU* is apparently dispensable for *Psg* growth in rich LB medium, since no growth difference was observed among WT, *algU* mutant, and *algU*-complemented strain (Supplementary Fig. S2).

To further investigate the *algU* contribution to *Psg* virulence, we conducted syringe infiltration with WT, *algU* mutant, and *algU*-complemented strain. As a result, the *algU* mutant showed reduced symptoms and bacterial populations at all inoculum levels we tested (Figs. 3A–3F). Taken together, these results indicate that AlgU contributes to growth both on leaf surface and in apoplast, and to causing disease.

AlgU regulates the expression of *Psg* virulence genes in HSC medium

Pst DC3000 AlgU positively regulates virulence gene transcription (Markel et al. 2016; Ishiga et al. 2018). To investigate whether *Psg* AlgU also regulates virulence genes, we analyzed virulence gene expression profiles in HSC medium. COR biosynthesis-related genes including *cmaA* and *corR*, in the *algU* mutant showed reduced expression at 48 h after incubation (Figs. 4A and 4C). Moreover, *hrpL*, encoding HrpL (an alternative sigma factor recognizing the *hrp* box in the promoter of T3SS genes), also showed significantly less expression in the *algU* mutant at both 3 and 48 h after incubation compared to *Psg* WT (Fig. 4D). These results indicate that AlgU positively regulates COR biosynthesis-related genes and *hrpL*.

To investigate whether AlgU can coordinate gene expression involved in *Psg* motility, we determined the expression profile of *fliC* (encoding flagellin, relating to flagellar mobility). At 3 h after incubation, there was no difference in the flagellar-encoding gene expression between the *algU* mutant and *Psg* WT. However, after 48 h incubation, relative *fliC* expression was greater in the *algU* mutant compared to *Psg* WT (Fig. 4E). Additionally, *algD* expression was down regulated in the *algU* mutant (Fig. 4F), indicating that AlgU positively regulates alginate biosynthesis-related genes.

AlgU contributes to COR biosynthesis and biofilm formation in *Psg*

We demonstrated that COR biosynthesis-related genes in the *algU* mutant showed reduced expression in HSC medium (Figs. 4A–4D). To investigate whether AlgU contributes to COR production, we first conducted a hypertrophy-inducing activity test on potato tuber tissues for COR detection (Sakai et al., 1979; Völksch et al., 1989). Potato tuber tissues inoculated with *Psg* WT showed hypertrophy response, but those inoculated with COR-defective mutants (*cmaA* and *cfa6*) showed no response (Fig. 5A). The *algU* mutant-inoculated potato tuber tissues showed less hypertrophy response compared to those inoculated with *Psg* WT and the reduction was restored in the *algU*-complemented strain (Fig. 5A). Furthermore, we also quantified COR production by using HPLC. *Psg* WT produced around 90 ng/g of COR in HSC medium (Fig. 5B). However, the *algU* mutant produced only around one third as much COR compared to *Psg* WT (Fig. 5B). We also confirmed that the *algU*-complemented strain recovered COR production more than WT (Fig. 5B). Taken together, these results suggest that AlgU contributes to COR biosynthesis in *Psg*.

Since we also demonstrated that alginate biosynthesis-related genes *algD* showed reduced expression in *algU* mutant (Fig. 4F), we next investigated the biofilm formation ability in *Psg* WT and the *algU* mutant. The *algU* mutant showed a reduction in biofilm formation at 24, 48, and 72 hpi (Fig. 6). These results suggest that AlgU also contributes to biofilm formation in *Psg*.

Discussion

We attempted to identify *Psg* virulence factors that are crucial in soybean pathogenicity. We screened 1,000 *Psg* mutants by syringe-infiltration and identified 28 mutants with reduced virulence (Fig. 1B). Several important virulence factors contribute to *Psg* virulence including COR, the T3SS, and AlgU (Table 1). Sigma factor AlgU regulates not only *algD*, but also other virulence genes including *hrpL* and COR (Figs. 4A–4F). Our study provides new insights into AlgU function as a global regulatory hub for *Psg* pathogenicity by regulating the expression of multiple virulence genes.

Our screening identified that 17 out of 28 reduced virulence mutants were related to COR biosynthesis genes. These COR biosynthesis mutants were mostly disrupted by Tn5 on the *cfa* and *cma* operons (Table 1). The *cfa* and *cma* operons encode enzymes related to CFA and CMA biosynthesis, respectively, the two elements that are ligated together to form COR (Bender et al., 1999). Together, these results indicate that COR is an important *Psg* virulence factor.

The *algU* mutant showed reduced virulence in plants both spray- and syringe-inoculated (Figs. 2; Supplementary Fig. S1; Figs. 3), indicating that AlgU contributes to *Psg* multiplication both on leaf surface and in apoplast, and causing disease. Our results indicate that AlgU regulate several virulence factors. Firstly, gene expression related to COR biosynthesis, such as *cmaA* and *corR* (but not *cfl*) were suppressed in the *algU* mutant (Figs. 4A–4C). Moreover, COR production in the *algU* mutant also less than that of *Psg* WT (Figs. 5A and 5B), suggesting that AlgU contributes to COR production in *Psg*. Consistent with our results, Ishiga et al. (2018) demonstrated that gene expression related to COR biosynthesis was suppressed during *Pst* DC3000 *algU* mutant infection. Furthermore, AlgU also contributes to *Pst* DC3000 virulence by regulating COR production to overcome stomatal-based defense (Ishiga et al., 2018). Together,

these results suggest that AlgU suppresses stomatal-based defense in the early *Psg* infection stage with soybean plants. Moreover, COR contributes to virulence by overcoming apoplastic defense as well as stomatal-based defense in *Pcal* (Sakata et al., 2021). Further study on COR contribution in *Psg* virulence will be needed to understand AlgU-mediated COR regulation.

Secondary, the *algD* expression profile in the *algU* mutant was significantly reduced in comparison with *Psg* WT, indicating that AlgU is important in regulating alginate biosynthesis gene expression (Fig. 4F). This result was consistent with a previous report in *Pst* DC3000 (Ishiga et al., 2018), in which *algD* expression was significantly suppressed in an *algU* mutant. Further, alginate plays a crucial role in epiphytic fitness and survival, and contributes to *P. syringae* virulence (Yu et al., 1999). Alginate is one of the exopolysaccharides (EPSs), which are the major components of biofilms, in *Psg* (Osman et al., 1986; Sutherland, 2001). We demonstrated that biofilm formation was significantly decreased in the *algU* mutant compared with *Psg* WT (Fig. 6). *Psg* PG4180 AlgU is important in virulence and bacterial growth in host plants, but it is not dependent on alginate production (Schenk et al., 2008; Yu et al., 2014). Additionally, AlgU, but not AlgD plays a crucial role in *Pst* DC3000 virulence (Markel et al. 2016). Together, it is tempting to speculate that alginate function in virulence differs in each *P. syringae* pathovar. Thus, further study is needed to understand AlgU regulating genes involved in biofilm formation and alginate function in *Psg* virulence.

Thirdly, expression profiles also revealed *hrpL* transcripts were suppressed in the *algU* mutant compared with *Psg* WT (Fig. 4D). *Pst* DC3000 AlgU functions to regulate *hrpL* expression (Ishiga et al., 2018). Markel et al., (2016) also demonstrated that AlgU plays an important role in virulence by regulating the expression of T3Es and *hrpL*. In *P. syringae*, both the T3SS and T3Es genes are in turn encoded by the *hrp* gene cluster, while the sigma factor HrpL directly regulates both *hrc* and *hop* genes (Lam et al., 2014). Although many studies were carried out to elucidate the functions and mode of actions of the T3SS and its T3Es, AlgU regulation on the T3SS system is still unknown in *P. syringae* infection processes. Therefore, further precise characterization of AlgU-mediated T3SS regulation will be needed to understand global gene expression networks during *Psg* infection.

Lastly, *fliC* transcripts in the *algU* mutant were increased compared with those of *Psg* WT (Fig. 4E). In *Pst* DC3000, AlgU not only downregulates flagellar and chemotaxis genes *in vitro* (Markel et al., 2016), but also negatively regulates *fliC* expression during infection (Bao et al., 2020). *fliC* encodes the flagellin protein including the flg22 epitope which triggers PTI (Felix et al., 1999; Zipfel et al., 2004; Parys et al., 2021; Colaianni et al., 2021). Recent studies reported the important role of AlgU in de-flagellation during the *P. syringae*-plant interaction to reduce PTI activation, and promote bacterial fitness in its host (Bao et al., 2020). Likewise, AlgU also plays an important role in de-flagellation of *P. syringae* pv. *maculicola* ES4236, in which transposon inactivation of AlgW led to decreased AlgU activity and increased the flagella expression, as well as reduced bacterial growth *in planta* (Schreiber and Desveaux, 2011). Therefore, it is tempting to speculate that high levels of flagellin protein production in the *algU* mutant activate PTI.

Conclusions

Our findings indicate that multiple virulence factors regulated by AlgU, including COR biosynthesis, biofilm formation, and T3SS, contributes to *Psg* virulence in soybean. Our findings help to expand understanding of AlgU roles in *Psg* virulence. Further studies on AlgU regulated mechanisms will be needed to fully understand *Psg* virulence.

Acknowledgements

We thank Dr. Christina Baker for editing the manuscript. *Psg* WT were given from the NARO Genebank, Ibaraki, Japan.

References

- Bao Z, Wei HL, Ma X, Swingle B (2020) *Pseudomonas syringae* AlgU downregulates flagellin gene expression, helping evade plant immunity. *J Bacteriol* 202:e00418–19
- Bender CL (1999) Chlorosis-inducing phytotoxins produced by *Pseudomonas syringae*. *Eur J Plant Pathol* 105:1–12
- Bender CL, Young SA, Mitchell RE (1991) Conservation of plasmid DNA sequences in coronatine-producing pathovars of *Pseudomonas syringae*. *Appl Environ Microbiol* 57:993–999
- Bender CL, Srone HE, Sims JJ, Cooksey DA (1987) Reduced pathogen fitness of *Pseudomonas syringae* pv. *tomato* Tn5 mutants defective in coronatine production. *Physiol Mol Plant Pathol* 30:237–283
- Brooks DM, Bender CL, Kunkel BN (2005) The *Pseudomonas syringae* phytotoxin coronatine promotes virulence by overcoming salicylic acid-dependent defenses in *Arabidopsis thaliana*. *Mol Plant Pathol* 6:629–639
- Brooks DM, Hernández-Guzmán G, Kloek AP, Alarcón-Chaidez F, Sreedharan A, Rangaswamy V, Peñaloza-Vázquez A, Bender CL, Kunkel BN (2004) Identification and characterization of a well-defined series of coronatine biosynthetic mutants of *Pseudomonas syringae* pv. *tomato* DC3000. *Mol Plant Microbe Interact* 17:162–174
- Budde IP, Ullrich MS (2000) Interactions of *Pseudomonas syringae* pv. *glycinea* with host and nonhost plants in relation to temperature and phytotoxin synthesis. *Mol Plant Microbe Interact* 13:951–61
- Colaïanni NR, Parys K, Lee H-S, Conway JM, Kim NH, Edelbacher N, Mucyn TS, Madalinski M, Law TF, Jones CD, Belkhadir Y, Dangl JL (2021) A complex immune response to flagellin epitope variation in commensal communities. *Cell Host Microbe* 29:635–649.e9
- Felix G, Duran JD, Volko S, Boller T (1999) Plants have a sensitive perception system for the most conserved domain of bacterial flagellin. *Plant J* 18:265–276
- Haefele DM, Lindow SE (1987) Flagellar motility confers epiphytic fitness advantages upon *Pseudomonas syringae*. *Appl Environ Microbiol* 53:2528–2533

- 359 Hoitink H, and Sinden S (1970) Partial purification and properties of chlorosis inducing toxins of
360 *Pseudomonas phaseolicola* and *Pseudomonas glycinea*. *Phytopathology* 60:1236–1237
- 361 Ignjatov M, Milošević M, Nikolić Z, Vujaković M, Petrović D (2007) Characterization of
362 *Pseudomonas savastanoi* pv. *glycinea* isolates from Vojvodina. *The Polish Phytopathol*
363 *Soci* 45:43–54
- 364 Ishiga T, Ishiga Y, Betsuyaku S, Nomura N (2018) AlgU contributes to the virulence of
365 *Pseudomonas syringae* pv. *tomato* DC3000 by regulating production of the phytotoxin
366 coronatine. *J Gen Plant Pathol* 84:189–201
- 367 Keane P, Kerr A, New P (1970) Crown gall of stone fruit II. Identification and nomenclature of
368 *Agrobacterium* isolates. *Aust J Biol Sci* 23:585–596
- 369 Keen NT, Tamaki S, Kobayashi D, Trollinger D (1988) Improved broad-host-range plasmids for
370 DNA cloning in gram-negative bacteria. *Gene* 70:191–197
- 371 King EO, Ward MK, Raney DE (1954) Two simple media for the demonstration of pyocyanin
372 and fluorescein. *J Lab Clin Med* 44:301–307
- 373 Kloek AP, Verbsky ML, Sharma SB, Schoelz JE, Vogel J, Klessig DF, Kunkel BN (2001)
374 Resistance to *Pseudomonas syringae* conferred by an *Arabidopsis thaliana* coronatine-
375 insensitive (*coi1*) mutation occurs through two distinct mechanisms. *Plant J* 26:509–522
- 376 Lam HN, Chakravarthy S, Wei HL, Nguyen BH, Stodghill PV, Collmer A, Swingle BM,
377 Cartinhour SW (2014) Global analysis of the HrpL regulon in the plant pathogen
378 *Pseudomonas syringae* pv. *tomato* DC3000 reveals new regulon members with diverse
379 functions. *PLoS One* 9:e106115
- 380 Markel E, Stodghill P, Bao Z, Myers C, Swingle B (2016) AlgU controls expression of virulence
381 genes in *Pseudomonas syringae* pv. *tomato* DC3000. *J Bacteriol* 198:2330–2344
- 382 Melotto M, Underwood W, Koczan J, Nomura K, He SY (2006) Plant stomata function in innate
383 immunity against bacterial invasion. *Cell* 126:969–980
- 384 Osman SF, Fett WF, Fishman ML (1986) Exopolysaccharides of the phytopathogen
385 *Pseudomonas syringae* pv. *glycinea*. *J Bacteriol* 166:66–71
- 386 Palmer DA, Bender CL (1993) Effects of environmental and nutritional factors on production of
387 the polyketide phytotoxin coronatine by *Pseudomonas syringae* pv. *glycinea*. *Appl Environ*
388 *Microbiol* 59:1619–1626
- 389 Parys K, Colaianni NR, Lee H-S, Hohmann U, Edelbacher N, Trgovcevic A, Blahovska Z, Lee
390 D, Mechtler A, Muhari-Portik Z, Madalinski M, Schandry N, Rodríguez-Arévalo I, Becker
391 C, Sonnleitner E, Korte A, Bläsi U, Geldner N, Hothorn M, Jones CD, Dangl JL, Belkadir
392 Y (2021) Signatures of antagonistic pleiotropy in a bacterial flagellin epitope. *Cell host &*
393 *microbe*.
- 394 Peñaloza-Vázquez A, Preston GM, Collmer A, Bender CL (2000). Regulatory interactions
395 between the Hrp type III protein secretion system and coronatine biosynthesis in
396 *Pseudomonas syringae* pv. *tomato* DC3000. *Microbiology* 146:2447–2456
- 397 Qi M, Wang D, Bradley CA, Zhao Y (2011) Genome sequence analyses of *Pseudomonas*
398 *savastanoi* pv. *glycinea* and subtractive hybridization-based comparative genomics with nine
399 *Pseudomonads*. *PLoS One* 6:e16451

- Sakai R, Nishiyama K, Ichihara A, Shiraishi K, Sakamura S (1979) Studies on the mechanism of physiological activity of coronatine. *Jpn J Phytopathol* 45:645–653
- Sakata N, Ishiga T, Saito H, Nguyen VT, Ishiga Y (2019) Transposon mutagenesis reveals *Pseudomonas cannabina* pv. *alisalensis* optimizes its virulence factors for pathogenicity on different hosts. *PeerJ* 7:e7698
- Sakata N, Ishiga T, Masuo S, Hashimoto Y, Ishiga Y (2021) Coronatine contributes to *Pseudomonas cannabina* pv. *alisalensis* virulence by overcoming both stomatal and apoplastic defenses in dicot and monocot plants. *Mol Plant Microbe Interact* doi: 10.1094/MPMI-09-20-0261-R
- Sambrook J, Fritsch E, Maniatis T (1989) Molecular cloning: a laboratory manual, 2nd edition. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York
- Sawada T, Eguchi M, Asaki S, Kashiwagi R, Shimomura K, Taguchi F, Matsui H, Yamamoto M, Noutoshi Y, Toyoda K, Ichinose Y (2018) MexEF-OprN multidrug efflux pump transporter negatively controls N-acyl-homoserine lactone accumulation in *Pseudomonas syringae* pv. *tabaci* 6605. *Mol Genet Genom* 293:907–917
- Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A. (1994) Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* 145:69–73
- Schenk A, Weingart H, Ullrich MS (2008) The alternative sigma factor AlgT, but not alginate synthesis, promotes in planta multiplication of *Pseudomonas syringae* pv. *glycinea*. *Microbiology* 154:413–421
- Schenk A, Berger M, Keith LM, Bender CL, Muskhelishvili G, Ullrich MS (2006). The *algT* gene of *Pseudomonas syringae* pv. *glycinea* and new insights into the transcriptional organization of the *algT-muc* gene cluster. *J Bacteriol* 188:8013–8021
- Schreiber KJ, Desveaux D (2011) AlgW regulates multiple *Pseudomonas syringae* virulence strategies. *Mol Microbiol* 80:364–377
- Shao X, Xie Y, Zhang Y, Deng X (2019) Biofilm formation assay in *Pseudomonas syringae*. *Bio-protocol* 9:e3237–e3237
- Sutherland, I. (2001) Biofilm exopolysaccharides: a strong and sticky framework. *Microbiology* 147:3–9
- Ullrich M, Bereswill S, Volksch B, Fritsche W, Geider K (1993) Molecular characterization of field isolates of *Pseudomonas syringae* pv. *glycinea* differing in coronatine production. *Microbiology* 139:1927–1937
- Uppalapati SR, Ayoubi P, Weng H, Palmer DA, Mitchell RE, Jones W, Bender CL (2005) The phytotoxin coronatine and methyl jasmonate impact multiple phytohormone pathways in tomato. *Plant J* 42:201–217
- Uppalapati SR, Ishiga Y, Wangdi T, Kunkel BN, Anand A, Mysore KS, Bender CL (2007) The phytotoxin coronatine contributes to pathogen fitness and is required for suppression of

439 salicylic acid accumulation in tomato inoculated with *Pseudomonas syringae* pv. *tomato*
 440 DC3000. Mol Plant Microbe Interact 20:955–965

441 Völksch B, Bublitz F, Fritsche W (1989) Coronatine production by *Pseudomonas syringae*
 442 pathovars: screening method and capacity of product formation. J Basic Microbiol 29:463–
 443 468

444 Yu J, Peñaloza-Vázquez A, Chakrabarty AM, Bender CL (1999) Involvement of the
 445 exopolysaccharide alginate in the virulence and epiphytic fitness of *Pseudomonas syringae*
 446 pv. *syringae*. Mol Microbiol 33:712–720

447 Yu X, Lund SP, Greenwald JW, Records AH, Scott RA, Nettleton D, Lindow SE, Gross DC,
 448 Beattie GA (2014) Transcriptional analysis of the global regulatory networks active in
 449 *Pseudomonas syringae* during leaf colonization. mBio 5:e01683–14

450 Zhao Y, Thilmony R, Bender CL, Schaller A, He SY, Howe GA (2003) Virulence systems of
 451 *Pseudomonas syringae* pv. *tomato* promote bacterial speck disease in tomato by targeting the
 452 jasmonate signaling pathway. Plant J 36:485–499

453 Zipfel C, Robatzek S, Navarro L, Oakeley EJ, Jones JDG, Felix G, Boller T (2004) Bacterial
 454 disease resistance in Arabidopsis through flagellin perception. Nature 428:764–767

Table 1 (on next page)

Bacterial strains and plasmids used in this study

Table 1 Bacterial strains and plasmid used in this study

Bacterial strain or plasmid	Locus	Relevant characteristics	Reference or source
<i>E. coli</i> strain			
DH5 α		F- λ - ϕ 80dLacZ Δ M15 Δ (lacZYA-argF)U169 recA1 endA1 hsdR17 (rK-mK+) supE44 thi-1gyrA relA1	Takara, Kyoto, Japan
S17-1		Thi pro hsdR-hsdM + recA [chr::RP4-2-Tc::Km::Tn7]	Schäfer et al. 1994
<i>P. savastanoi</i> pv. <i>glycinea</i>			
<i>Psg</i>		<i>Psg</i> wild-type	MAFF301684
VTB8	PsgB076_27735	<i>cfa6::mTn5</i> , NaI ^r , Km ^r	This study
VTC23	PsgB076_27795	<i>cmaA::mTn5</i> , NaI ^r , Km ^r	This study
VTD16	PsgB076_27730	<i>cfa5::mTn5</i> , NaI ^r , Km ^r	This study
VTD28	PsgB076_27725	<i>cfa4::mTn5</i> , NaI ^r , Km ^r	This study
VTD29	PsgB076_27740	<i>cfa7::mTn5</i> , NaI ^r , Km ^r	This study
VTD30	PsgB076_27720	<i>cfa3::mTn5</i> , NaI ^r , Km ^r	This study
VTE13	PsgB076_27750	<i>cfa8::mTn5</i> , NaI ^r , Km ^r	This study
VTE17	PsgB076_27740	<i>cfa7::mTn5</i> , NaI ^r , Km ^r	This study
VTF3	PsgB076_27705	<i>cfa2::mTn5</i> , NaI ^r , Km ^r	This study
VTG22	PsgB076_27735	<i>cfa6::mTn5</i> , NaI ^r , Km ^r	This study
VTG29	PsgB076_27740	<i>cfa7::mTn5</i> , NaI ^r , Km ^r	This study
VTG41	PsgB076_27725	<i>cfa5::mTn5</i> , NaI ^r , Km ^r	This study
VTH39	PsgB076_27735	<i>cfa6::mTn5</i> , NaI ^r , Km ^r	This study

VTI15	PsgB076_27740	<i>cfa7::mTn5</i> , Nal ^r , Km ^r	This study
VTR4	PsgB076_27740	<i>cfa7::mTn5</i> , Nal ^r , Km ^r	This study
VTI14	PsgB076_27805	<i>cmaC::mTn5</i> , Nal ^r , Km ^r	This study
VTI22	PsgB076_27730	<i>cfa5::mTn5</i> , Nal ^r , Km ^r	This study
VTB2	PsgB076_28870	<i>Type III effector protein XopAD::mTn5</i> , Nal ^r , Km ^r	This study
VTB52	PsgB076_09885	<i>Heat shock protein 90 (Hsp90)::mTn5</i> , Nal ^r , Km ^r	This study
VTH40	PsgB076_15537	<i>MFS transporter::mTn5</i> , Nal ^r , Km ^r	This study
VTM22	PsgB076_27425	<i>Hypothetical protein::mTn5</i> , Nal ^r , Km ^r	This study
VTO15	PsgB076_01129	<i>ABC transporter permease::mTn5</i> , Nal ^r , Km ^r	This study
VTO41	PsgB076_06035	<i>Sigma factor algU::mTn5</i> , Nal ^r , Km ^r	This study
VTP20	PsgB076_26940	<i>DNA-binding protein::mTn5</i> , Nal ^r , Km ^r	This study
VTE41	PsgB076_09720	<i>Unknown</i> , Nal ^r , Km ^r	This study
VTF6	PsgB076_10315	<i>Unknown</i> , Nal ^r , Km ^r	This study
VTO37	PsgB076_19532	<i>Unknown</i> , Nal ^r , Km ^r	This study
VTI8	PsgB076_26618	<i>Unknown</i> , Nal ^r , Km ^r	This study
<i>algU</i> mutant (VTO41) + pDSKG- <i>algU</i>		<i>algU</i> mutant complemented with pDSKG- <i>algU</i> , Nal ^r , Gen ^r , Km ^r	This study
Plasmid			
pBSLC1		Transposon vector constructed by ligation of pBSL118 and pHSG396 at EcoRI site, Amp ^r , Km ^r , Cm ^r	Sawada et al. 2018
pDSK519		Broad-host-range cloning vector, Ken ^r	Keen et al. 1998
pDSKG		Broad-host-range cloning vector, Gen ^r	This study

pDSKG-*algU*

The vector containing *algU* gene inserted into
pDSK519, Gen^r

This study

Notes Amp^r ampicillin resistance, Cm^r chloramphenicol resistance, Gen^r gentamicin,
Km^r kanamycin resistance, Nal^r nalidixic acid resistance.


1

Figure 1

Disease symptoms on soybean leaves syringe-inoculated with *Pseudomonas savastanoi* pv. *glycinea* (*Psg*) wild-type (WT) and mutants.

(A) Disease symptoms on soybean leaves syringe-inoculated with 5×10^7 colony forming units (CFU)/ml of the *Psg* WT at 6 dpi. (B) Disease symptoms on soybean leaves syringe-inoculated with 5×10^7 colony forming units (CFU)/ml of the *Psg* mutants at 6 dpi. Scale bars shows 1 cm.

A



Psg WT

B




Figure 2

Bacterial populations and disease symptoms in soybean leaves.

(A) Bacterial populations in leaves spray-inoculated with *Pseudomonas savastanoi* pv.

glycinea (*Psg*) wild-type (WT) and mutants (1×10^8 colony forming units (CFU)/ml). Bacterial populations in leaves were estimated at 6 days post inoculation (dpi) by dilution plating on selective medium as described in the methods. Vertical bars indicate standard error for three independent experiments. Asterisks indicate a significant difference from WT in a *t* test (**P* < 0.05; ***P* < 0.01). (B) Disease symptoms in leaves spray-inoculated with the *Psg* WT and the *algU* mutant (1×10^8 colony forming units (CFU)/ml) at 6 dpi.




Figure 3

Disease symptoms and bacterial populations in soybean leaves after syringe inoculation.

Disease symptom and bacterial populations in leaves syringe-inoculated with *Pseudomonas syringae* pv. *glycinea* (*Psg*) wild-type (WT), the *algU* mutant, and the *algU*-complemented strain at 5×10^7 colony forming units (CFU)/ml (A, B), 5×10^6 CFU/ml (C, D), and 5×10^5 CFU/ml (E, F), respectively. Photographs were taken at 5 days post inoculation (dpi). Scale bars shows 1 cm. Bacterial populations in leaves were estimated at 5 dpi by dilution plating on selective medium as described in the methods. Vertical bars indicate standard error for three independent experiments. The different letters indicate significant statistical differences ($P < 0.05$, Turkey's HSD test).




Figure 4

Gene expression profiles involved in the virulence of *Pseudomonas savastanoi* pv. *glycinea* (Psg) wild-type (WT) and *algU* mutant in liquid HSC broth.

Psg WT and *algU* mutant were grown in HSC broth for 3 and 48 h, adjusted to an OD₆₀₀ of 0.1, and grown again in fresh HSC broth for 3 h. Gene expression was normalized using the housekeeping gene *Psg outer membrane lipoprotein I (oprI)* by real-time quantitative reverse transcription-polymerase chain reaction (RT-qPCR) with gene-specific primer sets (Supplementary Table S1). (A) *cmaA*, (B) *cfl*, (C) *corR*, (D) *hrpL*, (E) *fliC*, and (F) *algD*. Vertical bars indicate the standard error for three biological replicates. The different letters (a-c) indicate a significantly statistical difference ($P < 0.05$, Turkey's HSD test).





Figure 5

COR quantification of *Pseudomonas savastanoi* pv. *glycinea* (*Psg*) wild-type (WT), the *algU* mutant, and the *algU*-complemented strain.

(A) Observation of hypertrophy-inducing activity on potato tuber tissue inoculated with *Pseudomonas savastanoi* pv. *glycinea* (*Psg*) wild-type (WT), COR-defective mutants (*cmaA* and *cfa6*), the *algU* mutant, and the *algU*-complemented strain. Potato tuber discs were inoculated using toothpicks by placing the tips in the *Psg* WT, *cmaA*, *cfa6*, *algU* mutant, and the *algU*-complemented strain on a KB medium plate and then placing the toothpick on the potato tuber disc. Photographs were taken at 5 dpi. (B) COR quantification of *Pseudomonas savastanoi* pv. *glycinea* (*Psg*) wild-type (WT) and the *algU* mutant grown in liquid HS broth by HPLC. *Psg* WT, the *algU* mutant, and the *algU*-complemented strain were cultured in HSC broth for 7 days. HPLC analysis was conducted by a Shimadzu LC20A system equipped with a Symmetry C8 column. COR in the culture supernatant was identified, as compared with authentic COR as the standard. Vertical bars indicate the standard error for three biological replicates. Asterisks indicate a significant difference from WT in a *t* test (** $P < 0.01$). N.D. indicates not detected by HPLC.

A



B




Figure 6

Biofilm biomass measurement of *Pseudomonas savastanoi* pv. *glycinea* (Psg) wild-type (WT) and the *algU* mutant with crystal violet (CV) grown in LB broth at 28°C for 24, 48, 72, and 96 h.

The biofilm eluted sample's absorbance was measured at OD₅₉₅. Vertical bars indicate the standard error for three biological replicates. Asterisks indicate a significant difference from WT in a *t* test (* *P* < 0.05, ** *P* < 0.01).

