

The nonavian theropod quadrate I: standardized terminology and overview of the anatomy, function and ontogeny

By allowing the articulation of the mandible with the cranium, the quadrate of diapsids and most other tetrapods plays an important role morphofunctionally. In Theropoda, its morphology is particularly complex and varies importantly among different clades of nonavian theropods so that the quadrate possesses a strong taxonomic potential. Inconsistencies in the notation and terminology used in discussions of the theropod quadrate anatomy have been noticed, a number of no less than height different terms being sometimes given to a same structure. A standardization list of terms and notation for each quadrate anatomical entity is here proposed, with the goal of facilitating future descriptions of this important cranial bone. An overview of the quadrate function, pneumaticity and ontogeny in nonavian theropods is also given. The quadrate of the large majority of nonavian theropod is akinetic and the diagonally oriented sulcus of the mandibular articulation allowed both rami of the mandible to move laterally when opening the mouth in many of them. Pneumaticity of the quadrate is also present in most of tetanuran clades and the pneumatic chamber, invaded by the quadrate diverticulum of the mandibular arch pneumatic system, was connected to one or several pneumatopores on the medial, lateral, posterior, anterior or ventral sides of the quadrate. Absence of a quadrate foramen in allosauroid embryos and a poor delimitation of mandibular condyles in both embryonic and juveniles tetanurans seems to be ontogenetic features of some theropods. Finally, the numerous morphological differences existing in the quadrates of the two specimens of *Shuvuuia deserti*, interpreted by some as juvenile and adult individuals, are considered as ontogenetic, taphonomic, and perhaps also taxonomic variations.

1 **The nonavian theropod quadrate I: standardized terminology and** 2 **overview of the anatomy, function and ontogeny**

3 Christophe Hendrickx^{1,2}

4 ¹Universidade Nova de Lisboa, CICEGe, Departamento de Ciências da Terra, Faculdade de
5 Ciências e Tecnologia, Quinta da Torre, 2829-516, Caparica, Portugal.

6 ²Museu da Lourinhã, 9 Rua João Luis de Moura, 2530-158, Lourinhã, Portugal.
7 chritophe.hendrickx@hotmail.com

8 Ricardo Araújo^{2,3,4,5}

9 ²Museu da Lourinhã, 9 Rua João Luis de Moura, 2530-158, Lourinhã, Portugal.

10 ³Huffington Department of Earth Sciences, Southern Methodist University, PO Box 750395,
11 75275-0395, Dallas, Texas, USA.

12 ⁴Instituto de Plasmas e Fusão Nuclear, Instituto Superior Técnico, Universidade de Lisboa,
13 Portugal.

14 ⁵Museum für Naturkunde, Berlin, Germany.
15 rmaraujo@smu.edu

16 Octávio Mateus^{1,2}

17 ¹Universidade Nova de Lisboa, CICEGe, Departamento de Ciências da Terra, Faculdade de
18 Ciências e Tecnologia, Quinta da Torre, 2829-516, Caparica, Portugal.

19 ²Museu da Lourinhã, 9 Rua João Luis de Moura, 2530-158, Lourinhã, Portugal.
20 omateus@fct.unl.pt

ABSTRACT

By allowing the articulation of the mandible with the cranium, the quadrate of diapsids and most other tetrapods plays an important role morphofunctionally. In Theropoda, its morphology is particularly complex and varies importantly among different clades of nonavian theropods so that the quadrate possesses a strong taxonomic potential. Inconsistencies in the notation and terminology used in discussions of the theropod quadrate anatomy have been noticed, a number of no less than eight different terms being sometimes given to a same structure. A standardization list of terms and notation for each quadrate anatomical entity is here proposed, with the goal of facilitating future descriptions of this important cranial bone.

An overview of the quadrate function, pneumaticity and ontogeny in nonavian theropods is also given. The quadrate of the large majority of nonavian theropod is akinetic and the diagonally oriented sulcus of the mandibular articulation allowed both rami of the mandible to move laterally when opening the mouth in many of them. Pneumaticity of the quadrate is also present in most of tetanuran clades and the pneumatic chamber, invaded by the quadrate diverticulum of the mandibular arch pneumatic system, was connected to one or several pneumatopores on the medial, lateral, posterior, anterior or ventral sides of the quadrate. Absence of a quadrate foramen in allosauroid embryos and a poor delimitation of mandibular condyles in both embryonic and juveniles tetanurans seems to be ontogenetic features of some theropods. Finally, the numerous morphological differences existing in the quadrates of the two specimens of *Shuvuuia deserti*, interpreted by some as juvenile and adult individuals, are considered as ontogenetic, taphonomic, and perhaps also taxonomic variations.

INTRODUCTION

The quadrate (in Latin *quadratum*, meaning ‘square’) is a cranial bone of endochondral origin that articulates with the mandible in all gnathostomes except mammals, in which it evolved into the incus (Carroll 1988; Benton 2005; Brusatte 2012). In theropods, this bone plays many important functions such as a structural support for the basicranium, articulatory element with the lower jaws, attachment for several muscles, audition and hosting important nerves and vascular passages (e.g., Witmer 1990, 1997; Bakker 1998; Sedlmayr 2002; Kundrát and Janáček 2007; Holliday and Witmer 2008; Tahara and Larsson 2011; Appendix 1).

Its rather simple architecture, an elongated body bearing an anteriorly projected blade, therefore tends to vary significantly in the structure of its head, mandibular articulation, quadratojugal contact and the presence of pneumatic openings, quadrate foramen and lateral processes, among theropods with a large variability of feeding strategies (e.g., Holtz 2003; Therrien et al. 2005; Hone and Rauhut 2010; Zanno and Makovicky 2011). Such morphological variation has been recognized in avian theropods in particular, revealing the great taxonomical utility of the quadrate in this clade (e.g., Lowe 1926; Samejima and Otsuka 1987; Barbosa 1990; Elzanowski et al. 2001; Elzanowski and Stidham 2010). Likewise but to a lesser degree, the systematic potential of the quadrate bone has also been noted for nonavian theropods (Maryańska and Osmólska 1997; Currie 2006), witnessing the particular importance that should be accorded to the description of this bone in the literature on nonavian theropod anatomy. Nevertheless, the terminology and abbreviations of the quadrate anatomy has been inconsistent in nonavian theropods, several different anatomical terms for the same quadrate sub-entity being often used (Appendix 2). Although a thorough list of anatomical terms has been given by Elzanowski et al. (2001) and Elzanowski and Stidham (2010) for the avian quadrate, the terminology proposed by these authors has never been followed in the description of the nonavian theropod quadrate

hitherto. Indeed, the quadrate of birds has greatly changed in its morphology throughout the evolution of this clade and therefore displays many features absent in more primitive theropods, so that many anatomical terms coined by Elzanowski et al. (2001) and Elzanowski and Stidham (2010) cannot be applied for the nonavian theropod quadrate. Likewise, some quadrate entities such as the quadrate foramen and the lateral process observable in nonavian theropods are absent in their avian **descents** and do not appear in the list of these authors.

The present paper has two major aims. The first is to propose a standardization of the anatomical terms for the quadrate sub-units, each associated with a two to four letters abbreviation and followed by a definition, in order to facilitate future description of this bone in the literature. The second is to give a general overview of the function, pneumaticity and ontogeny of this important bone in nonavian theropods.

Institutional Abbreviations

AMNH, American Museum of Natural History, New York, U.S.A.; **BMNH**, The Natural History Museum, London, U.K.; **BYUVP**, Brigham Young University Vertebrate Paleontology, Provo, Utah, U.S.A.; **CMNH**, Carnegie Museum, Pittsburgh, Pennsylvania, U.S.A.; **FMNH**, Field Museum of Natural History, Chicago, Illinois, U.S.A.; **IGM**, Mongolian Institute of Geology, Ulaan Bataar, Mongolia; **IVPP**, Institute for Vertebrate Paleontology and Paleoanthropology, Beijing, China; **MACN**, Museo Argentino de Ciencias Naturales, Buenos Aires, Argentina; **MCF**, **PVPH**, Museo Municipal Carmen Funes, Paleontología de Vertebrados, Plaza Huincul, Argentina; **MCNA**, Museo de Ciencias Naturales y Antropológicas de Mendoza, Mendoza, Argentina; **MIWG**, Dinosaur Isle, Isle of Wight Museum Services, Sandown, U.K.; **ML**, Museu da Lourinhã, Lourinhã, Portugal; **MNHN**, Muséum national d'Histoire Naturelle, Paris, France; **MNN**, Musée National du Niger, Niamey, Niger; **MSNM**, Museo di Storia Naturale di Milano, Milan, Italy; **MUCPv**, Museo de Ciencias Naturales de la Universidad Nacional de Comahue,

90 Neuquén, Argentina; **NH**, Horniman Museum & Gardens, London, U.K.; **OUMNH**, Oxford
91 University Museum, Oxford, U.K.; **PVSJ**, Instituto y Museo de Ciencias Naturales, San Juan,
92 Argentina; **SMA**, Sauriermuseum Aathal, Aathal, Switzerland; **SMNS**, Staatliches Museum für
93 Naturkunde, Stuttgart, Germany; **RTMP**, Royal Tyrrell Museum of Palaeontology, Drumheller,
94 Alberta, Canada; **UCMP**, University of California Museum of Paleontology, Berkeley, California,
95 U.S.A.; **UC**, University of Chicago Paleontological Collection, Chicago, U.S.A.; **UMNH**, Utah
96 Museum of Natural History, Salt Lake City, Utah, U.S.A.; **USNM**, United State National
97 Museum Vertebrate Paleontology, Washington, D. C., U.S.A.; **WDIS**, Wyoming Dinamation
98 International Society, Casper, Wyoming, U.S.A.

99 PROPOSED TERMINOLOGY OF THE QUADRATE ANATOMY

100 **Figure 1**

101 **Table 1**

102 **Quadrate body (qb).** Part of the quadrate that includes the mandibular articulation, the quadrate
103 shaft, the quadrate ridge, the quadrate head, the lateral contact (quadratojugal and/or squamosal
104 contact), and the lateral process, and excludes the pterygoid flange. In posterior view, the
105 quadrate body is delimited by the lateral margin of the lateral contact and sometimes the medial
106 margin of the quadrate foramen, the ventral margin of the mandibular articulation, the dorsal
107 margin of the quadrate head, and a medial margin mostly formed by the quadrate shaft and the
108 medial fossa of the pterygoid flange. The quadrate body corresponds to the corpus ossis quadrati
109 of Baumel and Witmer (1993), and the corpus quadrati of Elzanowski et al. (2001) and
110 Elzanowski and Stidham (2010) for avian theropods.

111 **Quadrate shaft (qs).** Part of the quadrate body that excludes the lateral process and all
112 articulating surfaces (i.e., quadrate head, quadratojugal/squamosal/pterygoid contacts, and

mandibular articulation). The quadrate shaft, as called by Welles (1984), Sereno and Novas (1994), Norell et al. (2006), Sampson and Witmer (2007), Sereno et al. (2008), Carrano et al. (2011), and Brusatte et al. (2012), is also referred as the quadrate pillar by Madsen and Welles (2000), and the ascending process by Colbert (1989).

Quadrate ridge (qr). Dorsoventrally elongated column, ridge or crest located on the quadrate body and visible in posterior view. Although the quadrate ridge is present in the large majority of nonavian theropods, a description of the structure is often omitted in the literature. The quadrate ridge is referred as ‘a column’ for Welles (1984), a ‘ridge-like mediodorsal edge’ for Carr (1996), ‘a prominent rounded ridge’ for Smith et al. (2007), a ‘columnar ridge’ for (Rauhut et al. 2010) and a ‘robust ridge’ for Brusatte et al. (2012).

Quadrate ridge groove (qrg). Groove dividing the quadrate ridge in two different units at two-thirds, or more dorsally, of the quadrate body height. A quadrate ridge groove exists in some allosauroid theropods.

Quadrate head (qh). Dorsal articulation of the quadrate abutting to the cotyle of the squamosal and touching other bones of the braincase in some theropod taxa. The quadrate head, as it is called by Britt (1991), Charig and Milner (1997), Madsen and Welles (2000), Sampson and Witmer (2007), Sereno et al. (2008), Norell et al. (2009) and Brusatte et al. (2012) among others, has also been termed quadrate cotylus (Currie 2003; Coria and Currie 2006) quadrate cotyle (Currie 2003; Coria and Currie 2006), squamosal condyle (Coria and Salgado 1998), squamosal articulation (Turner et al. 2011), otic process (Maryańska and Osmólska 1997; Burnham 2004; Holliday and Witmer 2008), squamosal capitulum (Zanno 2010) and, processus oticus (Baumel and Witmer 1993; Figure 2), and caput quadrati (Elzanowski et al. 2001; Elzanowski and Stidham 2010). When the quadrate head is double, it divides into otic and squamosal capitula.

136 **Otic capitulum (otc).** Medial capitulum of the quadrate head articulating with the braincase. The
137 otic capitulum corresponds to the capitulum (condylus) oticum of Baumel and Witmer (1993;
138 Figure 2), Elzanowski et al. (2001) and Elzanowski and Stidham (2010) for avian theropods.

139 **Squamosal capitulum (sqc).** Lateral capitulum of the quadrate head articulating with the
140 squamosal. The squamosal capitulum corresponds to the capitulum (condylus) squamosum of
141 Baumel and Witmer (1993; Figure 2), Elzanowski et al. (2001) and Elzanowski and Stidham
142 (2010) for avian theropods.

143 **Intercapitular sulcus (icas).** Groove separating the ootic capitulum from the squamosal
144 capitulum on the dorsal surface of the quadrate head. The intercapitular sulcus (sensu Witmer
145 1990) corresponds to the incisura intercapitularis of Baumel and Witmer (1993; Figure 2), and the
146 vallecule intercapitularis of Elzanowski et al. (2001) and Elzanowski and Stidham (2010) for
147 avian theropods.

148 **Quadrate foramen (qf).** Aperture in the quadrate body or concavity on the lateral margin of the
149 quadrate body and delimited ventrally by the ventral quadratojugal contact and dorsally by the
150 dorsal quadratojugal contact and its ventral projection in some theropod taxa. Most authors
151 usually refer to this perforation as the quadrate foramen (e.g., Welles 1984; Sereno and Novas
152 1994; Charig and Milner 1997; Maryańska and Osmólska 1997; Currie and Carpenter 2000;
153 Coria and Currie 2006; Currie 2006; Norell et al. 2006; Zanno 2010; Choiniere et al. 2010; Foth
154 and Rauhut 2012; Brusatte et al. 2012) but it can be also called the paraquadratic foramen (e.g.,
155 Barsbold and Osmólska 1999; Kobayashi and Lü 2003; Kobayashi and Barsbold 2005), the
156 paraquadrate foramen (Sampson and Witmer 2007; Dal Sasso and Maganuco 2011), the
157 paraquadrate fenestra (Smith et al. 2007) or the quadrate fenestra (e.g., Carr 1996; Sereno et al.
158 1998; Currie 2003; Eddy and Clarke 2011). A quadrate foramen exists in all nonavian theropods
159 but Ceratosauria and Megalosauridae.

Mandibular articulation (mar). Ventral surface of the quadrate, articulating with the mandible and fitting in the glenoid fossa of the lower jaw. It includes the ectocondyle, entocondyles, sometimes a mediocondyle, and an intercondylar sulcus. Corresponds to the pars or processus mandibularis of Baumel and Witmer (1993; Figure 2), Elzanowski et al. (2001) and Elzanowski and Stidham (2010) for avian theropods. Although most authors (e.g., Currie 2006; Sampson and Witmer 2007; Rauhut et al. 2010; Brusatte et al. 2012) referred the ectocondyle and entocondyles as the lateral and medial condyles (or hemicondyles) respectively, the terms ‘ectocondyle’ and ‘entocondyle’ have been used by Welles (1984), and Madsen and Welles (2000). The condyle present in between the ecto- and entocondyles in some theropods is here coined mediocondyle.

Ectocondyle (ecc). Lateral condyle of the mandibular articulation. The ectocondyle corresponds to the condylus (mandibularis) lateralis of Baumel and Witmer (1993; Figure 2), Elzanowski et al. (2001) and Elzanowski and Stidham (2010) for avian theropods.

Entocondyle (enc). Medial condyle of the mandibular articulation. The entocondyle corresponds to the condylus (mandibularis) medialis of Baumel and Witmer (1993; Figure 2), Elzanowski et al. (2001) and Elzanowski and Stidham (2010) for avian theropods.

Mediocondyle (mdc). Median condyle of the mandibular articulation. The mediocondyle is referred as the third condyle by Clark et al. (1994) and Xu and Wu (2001), the accessory condyle by Kobayashi and Lü (2003), and the **condyles** caudalis of Baumel and Witmer (1993) and Elzanowski et al. (2001) for avian theropods.

Intercondylar sulcus (ics). Groove separating the ectocondyle from the entocondyle and **sliding** **along** the interglenoid ridge of the articular. The intercondylar sulcus, a term also used by Carrano et al. (2011), can be referred as a groove (e.g., Madsen 1976; Britt 1991; Madsen and Welles 2000; Currie 2006), swelling (Charig and Milner 1997), sulcus (e.g., Kobayashi and Lü 2003; Norell et al. 2006; Sadleir et al. 2008), trochlea (Brochu 2003; Brusatte et al. 2010), trochlear surface (Brusatte et al. 2010, 2012), and intercondylar bridge (Zanno 2010). The

intercondylar sulcus corresponds to the sulcus intercondylaris (Baumel and Witmer 1993) and the vallicula intercondylaris (Elzanowski et al. 2001; Elzanowski and Stidham 2010) of the quadrate of avian theropods.

Intercondylar notch (icn). Notch located in between the ectocondyle and entocondyle, either on the anterior or posterior margin of the mandibular articulation, and corresponding to the depressio praecondylaris of Elzanowski et al. (2001) for avian theropods.

Lateral process (lpq). Lateral or anterolateral projection of the lateral margin of the quadrate body. Also known as the dorsal wing (Welles 1984; Currie 2006), the anterolateral wing (Madsen and Welles 2000), the lateral lamina (Coria and Salgado 1998) and the lateral ramus (Sampson and Witmer 2007), this process can contact the quadratojugal and/or the squamosal and therefore either be referred to the quadratojugal ramus by Sampson and Witmer (2007) or the squamosal ramus Norell et al. (2006).

Quadratojugal contact (qjc). Contact of the quadrate with the quadratojugal on the lateral, anterolateral or, posterolateral margin of the quadrate body. The quadratojugal contact, which corresponds to the cotyla quadratojugalis of Baumel and Witmer (1993), Elzanowski et al. (2001) and Elzanowski and Stidham (2010) for avian theropods, can be divided into a ventral and a dorsal quadratojugal contact when the quadrate foramen is present and delimited by both quadrate and quadratojugal.

Ventral quadratojugal contact (vqjc). ventral contact of the quadrate with the quadratojugal. The ventral quadratojugal contact of the quadrate always receives the quadratojugal bone.

Dorsal quadratojugal contact (dqjc). dorsal contact of the quadrate with the quadratojugal. The ventral quadratojugal contact of the quadrate can either receive the quadratojugal or both quadratojugal and squamosal in some theropod taxa.

Ventral projection of the dorsal quadratojugal contact (vpdq). Small projection of the dorsal quadratojugal contact delimiting the dorsolateral margin of the quadrate foramen.

210 **Quadratojugal process (qjp).** Anterior projection of the ventral quadratojugal contact of the
 211 quadrate.

212 **Squamosal contact (sqc).** Contact on the lateral margin of the quadrate with the squamosal.

213 **Posterior fossa (pfq).** Depression or concavity situated on the posterior side of the quadrate body
 214 and dorsal to the mandibular articulation, ventral to the quadrate head and lateral to the quadrate
 215 ridge. The posterior fossa can include or exclude the quadrate foramen.

216 **Quadrate diverticulum (qdi).** Air sac invading the pneumatic chamber inside the quadrate body
 217 and communicating with other diverticula by the quadrate pneumatopores.

218 **Dorsal pneumatopore (dpne).** Pneumatic foramen or recess located on the dorsal part of the
 219 quadrate, beneath the quadrate head. The dorsal pneumatopore corresponds to the foramen
 220 pneumaticum caudomediale of Elzanowski et al. (2001) and Elzanowski and Stidham (2010) for
 221 avian theropods.

222 **Medial pneumatopore (mpne).** Pneumatic foramen or recess situated on the medial side of the
 223 quadrate, typically in the ventromedial part of the pterygoid flange. The medial pneumatopore
 224 corresponds to the foramen pneumaticum basiorbitale of Elzanowski et al. (2001) and
 225 Elzanowski and Stidham (2010) for avian theropods.

226 **Posterior pneumatopore (ppne).** Pneumatic foramen or recess on the posterior side of the
 227 quadrate body, typically at one-height of the quadrate. The posterior pneumatopore corresponds
 228 to the ‘foramen pneumaticum rostromediale’ of Elzanowski et al. (2001) and Elzanowski and
 229 Stidham (2010) for avian theropods.

230 **Ventral pneumatopore (vpne).** Pneumatic foramen or recess on the ventral side of the quadrate.
 231 The ventral pneumatopore corresponds to the foramen pneumaticum adventitium of Elzanowski
 232 et al. (2001) and Elzanowski and Stidham (2010) for avian theropods.

233 **Pterygoid flange (pfl).** Sheet-like projection attached to the quadrate body and extending
 234 anteriorly or anteromedially to contact the pterygoid bone. The pterygoid flange, a term also used

by Charig and Milner (1997), Brochu (2003), Currie (2006), Coria and Currie (2006) and (Rauhut et al. 2010), is also known as the quadrate/anterior flange (e.g., Colbert 1989; Norell et al. 2006; Brusatte et al. 2010, 2012), the pterygoid ramus (e.g., Sereno and Novas 1994; Sampson and Witmer 2007; Choiniere et al. 2010), the pterygoid wing (e.g., Welles 1984; Madsen and Welles 2000; Eddy and Clarke 2011), the pterygoid ala (e.g., Currie 2003, 2006; Sadleir et al. 2008; Dal Sasso and Maganuco 2011), the pterygoid process (Molnar 1991; Carr 1996; Sereno et al. 2008), the optic wing (Balanoff and Norell 2012), the orbital process (Clark et al. 1994; Chiappe et al. 2002), and the processus orbitalis (Baumel and Witmer 1993; Elzanowski et al. 2001; Elzanowski and Stidham 2010; Figure 2) for avian theropods.

Pterygoid contact (ptc). Contact on the medial margin of the pterygoid flange, or the quadrate body, with the pterygoid. The pterygoid contact corresponds, for avian theropods, to the facies pterygoidea in Elzanowski et al. (2001) and the facies articularis pterygoidea in Elzanowski and Stidham (2010), as well as the condylus pterygoideus, located on the quadrate body, in Baumel and Witmer (1993; Figure 2), Elzanowski et al. (2001) and Elzanowski and Stidham (2010).

Medial fossa (mfq). Depression or concavity situated ventroposteriorly on the pterygoid flange.

It is delimited by the quadrate shaft and the ventral shelf in some theropod taxa. The medial fossa corresponds to the fossa corporis quadrati of Fuchs (1954) and the fossa basiorbitalis of Elzanowski et al. (2001) and Elzanowski and Stidham (2010) for avian theropods.

Ventral shelf (vsh). a medial or medioposterior fold of the ventral margin of the pterygoid flange. The term ‘shelf’ was employed by Sereno and Novas (1994) and ‘ventral shelf’ was used by Sampson and Witmer (2007), Eddy and Clarke (2011) and Carrano et al. (2011).

Figure 2

Inter-taxic Topological Homologies

To establish comparisons between taxa with widely disparate quadrate morphology, a homology concept of the feature in question is required. Here, we will give a general account of the variability within different anatomical sub-units of the quadrate and by following the criteria summarized in Rieppel (2006).

The quadrate ridge (Figure 1, qr) is easily distinguishable in many theropod taxa such as *Aerosteon riocoloradensis* (MCNA-PV 3137) and *Proceratosaurus bradleyi* (NHM R.4860) but the limits of this structure on the quadrate body can be subtle like in *Noasaurus leali* (PVL 4061), *Majungasaurus crenatissimus* (FMNH PR 2100), and *Eustreptospondylus oxoniensis* (OUMNH J.13558). The quadrate ridge corresponds to a columnar ridge in many theropod clades like in *Dilophosaurus wetherilli* (Welles 1984), *Allosaurus* (SMA 005/02) and *Eotyrannus lengi* (MIWG, 1997.550) but can also corresponds to a thin crest as in *Tyrannosauridae* (AMNH 5027; Carr 1996; Brusatte et al. 2012). Although the ventral portion of the quadrate ridge is usually demarcated just above the medial condyle of the mandibular articulation, its dorsal termination is more variable. The dorsal termination can reach the quadrate head like in *Acrocanthosaurus atokensis* (NCSM 14345) or flatten at the mid-height of the quadrate such as in *Albertosaurus sarcophagus* (Currie 2003, figure 10B). Despite being the same structure, the quadrate ridge can be separated by a deep groove like in *Allosaurus fragilis* (AMNH 600) and *Allosaurus europaeus* (ML 415) or flare and reappear more dorsally at the second third of the quadrate such as in some derived *Spinosauridae* (pers. observ.). Likewise, the ventral portion can also be divided in two parts separated by a concavity such as in the tyrannosaurids *Albertosaurus sarcophagus*, *Daspletosaurus* sp. (Currie 2003, figure 10 and 28) and *Tyrannosaurus rex* (AMNH 5027).

The pterygoid flange (Figure 1, pfl) contacts the quadrate process of the pterygoid anteriorly or anteromedially, and sometimes other bones such as the epipterygoid in *Herrerasaurus ischigualastensis* (PVSJ 407), the basipshenoid and prootic in *Erlikosaurus andrewsi* (Clark et al. 1994), and the squamosal in *Khaan mckennai* (Balanoff and Norell 2012).

Although the pterygoid flange can be easily homologized between taxa, it may acquire subtrapezoidal, subtriangular, subrectangular and M-shaped outlines or form a large semi-oval structure. The ventral limit of the flange can reach the mandibular condyles (e.g., *Baryonyx walkeri*, *Tyrannosaurus rex*) or get attached to the quadrate body well-above the mandibular articulation (e.g., *Majungasaurus crenatissimus*). This single structure can in some instances be divided into two ridges delimited by a deep pneumatic fossa facing ventrally (e.g., *Alioramus altai*, Brusatte et al. (2012), figure 23c; *Tyrannosaurus rex* FMNH PR2081). In anterior view, the pterygoid flange can be straight and only directed anteriorly like in the carcharodontosaurid *Shaochilong maortuensis* (Brusatte et al. 2010, figure 7a), or completely recurved anteromedially and its anteroventral margin can be either straight, or fold medially and dorsally, forming the ventral shelf, such as in *Majungasaurus crenatissimus* (FMNH PR 2100), *Carnotaurus sastrei* (MACN-CH 894) and *Allosaurus fragilis* (Madsen 1976:plate 3d).

The medial fossa of the quadrate (Figure 1, mfg) can be easily recognizable between taxa as it is always situated on the pterygoid flange, usually on the ventral part of it. This fossa is posteriorly delimited by the quadrate body in nonavian theropods and sometimes by the ventral shelf of the pterygoid flange. The medial fossa can be of variable depth (deep in *Cryolophosaurus* and shallow in *Eustreptospondylus*), pneumatized (e.g., *Falcarius*), and situated in the ventralmost part of the pterygoid flange (e.g., *Tsaagan*) or at mid-height of it and just above a large pneumatic recess like in *Mapusaurus roseae* (MCF PVPH-108.102).

The posterior fossa of the quadrate (Figure 1, pfq) can be located either in between the quadrate and the quadratojugal, being centred on the lateral margin of the quadrate (e.g., *Ceratosaurus*); or in the middle of the quadrate shaft and between the quadrate ridge and the lateral margin of the quadrate (e.g., ‘*Syntarsus*’; *Majungasaurus*; *Tsaagan*). Difficulties to recognize this structure between taxa may arise because it is not always well-delimited (e.g., *Shaochilong*). The posterior fossa can either be strongly ventrodorsally elongated like in the

megalosaurid *Torvosaurus tanneri*, or form an oval concavity lateromedially wide (e.g., *Majungasaurus*). Similarly to the medial fossa, the posterior fossa can have a large pneumatic recess positioned dorsally (e.g., *Sinornithomimus*) or ventrally (e.g., *Garudimimus*) inside the fossa.

Due to the highly variable morphology of the quadrate foramen (Figure 1, qf), this structure deserves special attention. It can be completely absent (e.g., *Carnotaurus*, *Torvosaurus*), or form a very small aperture (e.g., ‘*Syntarsus*’) to a large opening (e.g., *Tsaagan*). In most nonavian theropods, the quadrate foramen is mostly delimited by the quadrate and only its lateral margin is bounded by the quadratojugal (e.g., *Sinraptor*). In some nonavian theropods, however, the medial margin of the quadrate foramen and part of the ventral and dorsal margins are formed by the quadrate, the other lateral half being delimited by the quadratojugal (e.g., *Dromaeosaurus*). Finally, in a few theropods, the foramen can be completely enclosed in the quadrate (e.g., *Aerosteon*).

The quadratojugal contact of the quadrate (Figure 1, qjc) can either be a single extensive contact or made of two contacts separated by the quadrate foramen. In the latter case, the ventral quadratojugal contact and the dorsal quadratojugal contact of the quadrate are not always clearly separated and their dorsal and ventral margins, respectively, can overlap like in the sinraptorid *Sinraptor dongi* (IVPP 10600). If the quadrate foramen is absent or located inside the quadrate, the lateral quadratojugal contact is an elongated line of variable width along the lateral margin of the quadrate. If separated by the quadrate foramen, the ventral and dorsal contacts can display a wide variety of shapes that, nevertheless, are easily recognizable inter-taxically. Both quadratojugal contacts may face laterally, anteriorly or posteriorly and their articulating surface can be smooth, irregular or deeply grooved by several radiating ridges as in *Allosaurus fragilis* (Madsen 1976). The ventral quadratojugal contact is usually D-shaped or ovoid and its anterior margin can extend far anteriorly, forming the quadratojugal process occasionally forming a large

subtriangular projection (Norell et al. 2006). The dorsal quadratojugal contact can vary from a very thin line to a broad surface and its dorsal extension can reach the dorsal condyle or terminate well beneath it. A ventral projection of this contact may be present, and such projection delimiting part of the lateral border of the quadrate can either be short, like in *Daspletosaurus* sp. (Currie 2003, figure 28A), or formed an elongated ramus, like in the therizinosaurid *Falcarius utahensis* (Zanno 2010, figure 1H) and the coelurosaur *Zuolong salleei* (Choiniere et al. 2010, figure 3B).

In some basal theropods, ceratosaurs and dromaeosaurids, the lateral process of the quadrate (Figure 1, lpq) forms a wing-like projection similar to the pterygoid flange. This process is an extension of the quadrate body laterally so that it can be difficult to delimitate and one can see the presence of such process in *Allosaurus 'jimmadseni'* (SMA 005/02), *Sinraptor dongi* (Currie 2006, figure 1D), and *Erlikosaurus andrewsi* (Clark et al. 1994, figure 7). The lateral process can also vary in shape and size, as it can be short and parabolic (e.g., *Carnotaurus*) or elongated and subtriangular (e.g., *Tsaagan*). Its ventral border can also extend to the quadrate foramen (e.g., *Bambiraptor*) or more ventrally, sometimes reaching the medial condyle of the mandibular articulation (e.g., *Ilokelesia*).

The quadrate head (Figure 1, qh) always articulates with the deep cotylus of the squamosal and contacts more rarely other bones of the braincase such as the opisthotic in oviraptorids (Maryńska and Osmólska 1997), the prootic in *Mononykus olecranus* (Perle et al. 1994; Chiappe et al. 2002) and the postorbital in *Shuvuuia deserti* (Chiappe et al. 1998, 2002). The contact of the braincase between the dorsal part of the quadrate and the opisthotic-exoccipital or the paroccipital process is also present in *Herrerasaurus ischigualastensis* (Serenó and Novas 1994), *Dilophosaurus wetherilli* (Welles 1984), *Ceratosaurus magnicornis* (Madsen and Welles 2000; Sanders and Smith 2005), tyrannosaurids (Currie 2003), *Heyuannia huangi* (Lü 2005), and perhaps *Erlikosaurus andrewsi* (Clark et al. 1994), but this contact occurs on a small medial

surface just below the quadrate head and not with the quadrate head itself. The large majority of nonavian theropods have a single headed quadrate head (Rauhut 2003; pers. observ.); however, oviraptorids (Maryńska and Osmólska 1997, figure 3B) and the alvarezsaurid *Shuvuuia deserti* (Chiappe et al. 1998) have a unique double head. A bistylic quadrate head has also been observed in the dromaeosaurid *Mahakala omnogovae* (Turner et al. 2007) but Turner et al. (2011, figure 4) later reconsidered the head of the quadrate as not being double headed. The morphology of the quadrate head is variable; it may be subtriangular in most basal theropods (Serenó and Novas 1994; UCMP 37302), oval or subcircular in megalosaurids (UC OBA1; BYUVP 9246) and allosauroids (MCNA-PV-3137; IVPP 10600; IVPP V2885.3), subquadrangular in Spinosaurinae (SMNS 58022; MSNM V6896) or conical in Oviraptoridae (Maryńska and Osmólska 1997, figure 1B). Whilst most nonavian theropods have either a convex or a flattened quadrate head, the quadrate of some allosaurids (Bakker 1998, figure 5C) and derived tyrannosaurids (FMNH PR208) can also possess a well-marked concavity on the dorsal margin of the quadrate head. Despite this variability, the quadrate head can be easily homologized inter-taxically due to the obvious location of this structure.

With the exception of the therizinosaur *Erlikosaurus andrewsi* and the ornithomimosaur *Sinornithomimus dongi* which both seem to have a unique tricondylar condition on the mandibular articulation (Clark et al. 1994; Kobayashi and Lü 2003), all other nonavian theropods have two mandibular condyles. The presence of three mandibular condyles was also noted in the alvarezsaurid *Avimimus portentosus* (Chatterjee 1995) and the dromaeosaurid *Sinornithosaurus millenii* (Xu and Wu 2001). However, Vickers-Rich et al. (2002) only found two condyles in the former and our observations confirm that the third condyle of the latter seems to be part of the much broader lateral condyle (Xu and Wu 2001, figure 4D).

The intercondylar sulcus (Figure 1, ics) varies in orientation, size and depth. It can be large, shallow and sub-perpendicular to the long axis passing through the mandibular articulation

as in *Tyrannosaurus rex* (FMNH PR2081); or narrow, deep and strongly lateromedially-oriented as in some derived spinosaurids (pers. observ.).

The shape of the mandibular articulation (Figure 1, mar) in posterior view can vary from the biconvex condition known in most theropods, to the W-shaped articulation typical of *Citipati osmolskae* (Clark et al. 2002, figure 6) or a unique convex articulation seen in some dromaeosaurids such as *Tsaagan manga* (IGM 100/1015). The intercondylar notch (Figure 1, icn) is present in *Allosaurus 'jimmadseni'* (Bakker 1998, figure 5B, C; SMA 005/02) and *Suchomimus tenerensis* (MNN GAD 502) on the posterior side of the mandibular articulation, and in *Majungasaurus crenatissimus* (FMNH PR 2100) and *Carnotaurus sastrei* (MACN-CH 894) on its anterior margin. The ectocondyle (Figure 1, ecc) and entocondyle (Figure 1, ent) are highly variable among each clade of nonavian theropods in terms of shape, size and orientation.

Pneumaticity of the quadrate can either be internal or, obviously expressed externally by pneumatopores. The establishment of inter-taxic homologies is difficult to assess, because these structures have very diverse interspecific variability. Nevertheless, as in other saurischian taxa (Schwarz et al. 2007), these pneumatic structures have phylogenetic signal. These openings can appear on different views and portions of the quadrate. The medial and posterior pneumatopores (Figure 1, ppne) usually occur in the medial and posterior fossa respectively, and their position inside the fossae is again quite variable. Pneumatopores can also be located in a pneumatic recess outside the medial fossa and just beneath it such as in the carcharodontosaurids *Mapusaurus roseae* (Coria and Currie 2006) and *Acrocanthosaurus atokensis* (Eddy and Clarke 2011). In the latter, the pneumatic aperture is divided by a septum.

OVERVIEW OF THE FUNCTION, PNEUMATICITY AND ONTOGENY

Figure 3

Function of the Quadrate

Although in all archosaurs, and all amniotes except mammals, the main function of the quadrate is the articulation of the cranium with the mandible, this bone can also play an important role in the mobility of the skull in many extant theropods. Quadrate movement inside the skull, known as streptostyly, is a fundamental property of all avian theropods, and cranial kinesis in birds, known already in the beginning of the, 19th century (Nitzsch 1816), has been extensively studied in the past sixty years (e.g., Fisher 1955; Bock 1964, 1999, 2000; Bühler 1981; Zusi 1984, 1993; Bühler et al. 1985, 1988; Chatterjee 1991, 1997; Hoese and Westneat 1996; Zweers et al. 1997; Zweers and Vanden Berge 1998; Bout and Zweers 2001; Gussekloo and Bout 2005; Meekangvan et al. 2006). Streptostyly consists of the rotation of the quadrate at its dorsal articulation against the squamosal and/or supratemporal and usually corresponds to a transversal movement, although a lateral movement of the quadrate around an anteroposteriorly directed axis occurs in some lepidosaurs taxa (Metzger 2002). Cranial kinesis in avian theropods with a streptostylic quadrate includes upward (protraction) and downward (retraction) rotation of the upper jaw relative to the braincase and three main types of kinesis are recognized relative to the position of the dorsal flexion zone of the cranium and the nature of the nasal opening in modern theropods (Bock 1964; Bühler 1981; Zusi 1984; Meekangvan et al. 2006). There are the prokinesis where the flexion occurs at the nasofrontal joint and the upper jaw thereby moves as one unit, the amphikinesis where the flexion occurs in two zones of flexibility and the upper jaw and its tip are bent upward, and finally the rhynchokinesis where the flexion happens forward from the nasofrontal joint, allowing its rostral part to be moved.

Inference of the cranial kinesis and quadrate mobility in nonavian theropods has been recently investigated by Holliday and Witmer (2008) which regard the cranium of this group of dinosaurs as partially kinetically competent, since synovial joints and protractor muscles are present, but not fully kinetic like in birds. The strong suture of the quadrate to the quadratojugal

and the immobile contact of the quadrate and the pterygoid on the medial side of the pterygoid flange in most nonavian theropods seem to indicate a very limited movement, and perhaps even the total absence of movement of this bone within the cranium. Although the synovial quadrate head joint existing in theropods, and all other archosaurs, is necessary to infer cranial kinesis, its presence in **akinetetic taxa such as crocodiles and thyreophorans** demonstrates that the synovial joint cannot be considered alone as an argument for cranial kinesis. Synovial joints have actually been interpreted as growth zones rather than articular surfaces of mobile joints based on the presence of very thin articular cartilage covering the end of this joint (Holliday and Witmer 2008). According to Holliday and Witmer (2008) “articular cartilage persists in loading environments that exert hydrostatic pressures (which result in a change in volume but not shape) but exert low shear stresses”. Indeed, one of the key centres of deformation during normal biting is the quadrate-squamosal contact, which would have experienced large shear stresses associated with torque and asymmetrical loading during biting (Rayfield 2005), and the presence of a minimal amount of cartilage between the quadrate and squamosal would therefore suggest that the synovial zone was rather a growth zone than a mobile one. A streptostylic quadrate in *Tyrannosaurus rex* (Molnar 1991, 1998), *Oviraptor philoceratops* (Smith 1992), *Heyuannia huangi* (Lü 2005) and *Dromiceiomimus breviterius* (Russell 1972) based on the saddle joint between the quadrate and squamosal only is thereby unlikely.

Nevertheless, and more convincingly, a streptostylic quadrate has also been proposed in the alvarezsaurid *Shuvuuia deserti* (Chiappe et al. 1998) **in which the quadratojugal/jugal (Dufeau (2003) suggested that the quadratojugal may be absent in *Shuvuuia deserti*), rather than being firmly sutured to the quadrate, like in other nonavian theropods, would have contacted the quadrate though a movable joint (Chiappe et al. 1998, 2002; Figure 3).** According to Chiappe et al. (1998), the absence of a latero-dorsal contact of the quadrate with the quadratojugal/jugal, as well as a ventro-lateral process of the squamosal, would have permitted to the quadrate of this

mononykine to pivot antero-posteriorly, and the upper jaw to rotate ventro-dorsally thanks to this transversal movement. Although these authors have implied the existence of a bending zone between the frontals and the nasal–preorbital bones in *S. deserti*, allowing the flexion of the snout as a single unit when the quadrate displaced forward, like in prokinetic birds, the complex contacts between the nasal, frontal and prefrontal illustrated by Sereno (2001, figure 12B) makes assessment of Chiappe et al. (1998) hypothesis quite dubious (Holliday and Witmer 2008). In addition, Holliday and Witmer (2008) noted that a maxillojugal and palatal flexion zones authorizing a true prokinesis to be present in alvarezsaurids is still not clear. Likewise, the contact between the pterygoid flange of the quadrate and the pterygoid needs also to be better documented in order to imply any specific movement of the quadrate inside the cranium of *S. deserti*.

A movable articulation between the quadrate and quadratojugal may also have been present in the oviraptorids *Heyuannia huangi* (Lü 2003) and *Nemegtia huangi* (Lü et al. 2004) in which the quadrate and quadratojugal articulation of the former corresponds to a trochlea-like structure (Lü 2003, 2005), while the quadratojugal contact of the latter is convex and fit into a quadratojugal cotyle (Lü et al. 2004).

Quadrate articulation with the mandible and orientation of the intercondylar sulcus are highly variable among nonavian theropods, therefore suggesting some variation in the movement of the rami when the jaw opened. The helical intercondylar sulcus present in many nonavian theropods, but not all of them (pers. observ.), was noticed by Bakker (1998) in primitive theropod dinosaurs, Hendrickx and Buffetaut (2008) in spinosaurids, and Molnar (1991) and (Larson 2008) in *Tyrannosaurus rex*. These authors suggested that such spiral groove of the mandibular articulation constrained the diagonal ridge of the articular glenoid fossa, which fitted into the intercondylar sulcus, to slide laterally, therefore forcing the rami of the mandible to displace

laterally when the lower jaw was depressed, enlarging the width of the larynx in order to swallow food of large size (Hendrickx and Buffetaut 2008).

In allosaurid quadrates, the marked constriction of the intercondylar sulcus, associated with the enlargement of the mandibular condyles, the backward bent of the ventral part of the quadrate, and the intercondylar notch, were interpreted by Bakker (1998) as joint-stabilization zones. According to this author, the enlargement of the articulating surface would improve the stability of the joint when the mouth was widely opened, while the posterior depression would correspond to the area for attachment of one or several ligaments within the quadrate-mandibular articulation ((Bakker 1998). An intercondylar notch has also been noticed in the abelisaurids *Carnotaurus sastrei* (MACN-CH 894) and *Majungasaurus crenatissimus* (FMNH PR 2100), and the spinosaurid *Suchomimus tenerensis* (MNN GAD 502), implying a similar mechanic of the mandibular articulation than *Allosaurus*.

Figure 4

Pneumaticity in the Quadrate

Pneumatization of the quadrate bone has long been recognized for its phylogenetic value (e.g., Gauthier 1986; Holtz 1998; Chiappe 2001; Rauhut 2003; Holtz et al. 2004; Smith et al. 2007; Benson 2010; Carrano et al. 2012; Turner et al. 2012; Novas et al. 2013; Choiniere et al. 2014). Pneumatopores within the quadrate are widespread among avetheropod clades (Gold et al. 2013; Figure 4). The presence of one or several pneumatopores has indeed been recorded in carcharodontosaurids (e.g., Coria and Currie 2006; Eddy and Clarke 2011), neovenatorids (Serenó et al. 2008), tyrannosauroids (e.g., Molnar 1991; Brochu 2003; Currie 2003; Xu et al. 2004; Brusatte et al. 2012; Gold et al. 2013), compsognathids (Currie and Chen 2001), therizinosauroids (Clark et al. 1994; Zanno 2010), oviraptorids (e.g., Maryańska and Osmólska 1997; Lü 2003; Kundrát and Janáček 2007; Balanoff and Norell 2012), *ornithomimoids* (Witmer

1997; Tahara and Larsson 2011), dromaeosaurids (Makovicky et al. 2005) and troodontids (Barsbold et al. 1987; Currie and Zhao 1993; Varricchio 1997; Xu et al. 2002; Xu and Norell 2004). An incipient development of a pneumatic recess also exists in the basal allosauroid *Sinraptor dongi* (Currie 2006), and quadrate pneumaticity therefore seems to be an avetheropod synapomorphy (Figure 4). Among nonavian avetheropods, there is however no evidence of a pneumatic quadrate in Alvarezsauroidea hitherto.

The pneumatic foramen is particularly large in some allosauroids such as *Aerosteon riocoloradensis* (Serenio et al. 2008) and *Acrocanthosaurus atokensis* (Eddy and Clarke 2011), and the therizinosaur *Falcarius utahensis* (Zanno 2010; Figure 5). It however corresponds to a small rounded or oval aperture lodged in the posterior fossa of the quadrate body in most avetheropods (Figure 5). Indeed, a posterior pneumatopore exists in the tyrannosauroid *Dilong paradoxus* (Xu et al. 2004), the compsognathid *Sinosauropteryx prima* (Currie and Chen 2001, figure 3f), the ornithomimids *Garudimimus brevipes* (the ‘paraquadrate foramen’ or ‘paraquadratic foramen’ of Kobayashi and Barsbold 2005), *Sinornithomimus dongi* (the ‘paraquadratic foramen’ of Kobayashi and Lü 2003) and *Ornithomimus edmontonicus* (Tahara and Larsson 2011), the dromaeosaurid *Buitreraptor gonzalezorum* (Makovicky et al. 2005), and the troodontids *Mei long* (Xu and Norell 2004), and *Sinovenator changii* (Xu et al. 2002). A pneumatopore can also be located in the ventral corner of the pterygoid flange, as observed in the carcharodontosaurids *Acrocanthosaurus atokensis* (Eddy and Clarke 2011), *Mapusaurus roseae* (Coria and Currie 2006), *Giganotosaurus carolinii* (MUCPv-CH-1), and the tyrannosaurid *Albertosaurus sarcophagus* (Currie 2003, figure 10B). A pneumatic opening is also present anteroventrally, within a recess on the posteroventral part of the pterygoid flange (‘funnel-like external opening on the rostral surface of the quadrate, above the condyles’ of Gold et al., 2013, p. 37) like in the therizinosauroid *Falcarius utahensis* (Zanno 2010) and the tyrannosaurids *Alioramus altai* (Brusatte et al. 2012; Gold et al. 2013), *Daspletosaurus* sp. (Currie 2003, figure

28C) and *Tyrannosaurus rex* (Brochu 2003). This ventral pneumatopore has also been reported in the basal tyrannosauroid *Dilong paradoxus* (Gold et al. 2013) but was not observed in the closely related taxa *Guanlong wucaii*, *Proceratosaurus lengi*, and *Xiongguanlong baimoensis* (Gold et al. 2013), and its presence cannot be established in *Eotyrannus lengi* (contra Gold et al. 2013; pers. obs.). More rarely, a pneumatic opening can be situated on the lateral surface of the quadrate body, as in *Aerosteon riocoloradensis* (MCNA-PV 3137), and in the anterior part of the quadrate, as in *Mapusaurus roseae* (Coria and Currie 2006), *Troodon formosus* (Currie and Zhao 1993), *Heyuannia huangi* (Lü 2005), and perhaps *Tyrannosaurus rex* (Molnar 1991).

Figure 5

Carcharodontosauridae (Coria and Currie 2006; Eddy and Clarke 2011) and Tyrannosauridae (Molnar 1991; Brochu 2003) possess several pneumatic openings which perforate different sides of the quadrate and sometimes intercommunicate (Brochu 2003). The pneumatic foramina usually enter a large pneumatic chamber within the quadrate bone such as in *Tyrannosaurus rex* (Molnar 1991; Brochu 2003), *Alioramus altai* (Gold et al. 2013), *Conchoraptor gracilis* (Kundrát and Janáček 2007) or *Ornithomimus edmontonicus* (Tahara and Larsson 2011). The neovenatorid *Aerosteon riocoloradensis* also possess a large posterior pneumatopore leading to a pneumatic chamber, as well as a shallow pneumatic recess on the lateral surface of the quadrate shaft (pers. obs.).

These pneumatopores and the pneumatic chamber associated with them are invaded by the quadrate diverticulum of the mandibular arch pneumatic system which, together with the periotic pneumatic system, forms the tympanic sinus of archosaurs (Tahara and Larsson 2011). The mandibular arch pneumatic system includes the quadrate and/or the articular diverticulum which both have their embryological origins as parts of the first pharyngeal (= mandibular) arch, like the middle ear sac itself (Witmer 1997). As nonavian theropods, the quadrate diverticulum of modern birds exhibits a large variety of morphologies, and can either pneumatize the quadrate by

entering through a single medial or anteromedial foramen or not (Witmer 1990; Tahara and Larsson 2011). In the basal theropods that do not have a pneumatic quadrate, both medial and posterior fossae of the quadrate correspond to the osteological trace of the quadrate diverticulum. In nonavian theropods with a pneumatic quadrate, the position of the quadrate diverticulum is variable such as in ornithomimids (Tahara and Larsson 2011), carcharodontosaurids and oviraptorids (pers. observ.). The quadrate diverticulum of nonavian theropods may also have communicated with other diverticula such as the squamosal diverticulum as in *Conchoraptor gracilis* (Kundrát and Janáček 2007), and the siphoneal diverticulum of the articular as in *Dilong paradoxus*, *Aerosteon riocoloradensis* and perhaps other nonavian maniraptorans (Serenó et al. 2008; Tahara and Larsson 2011). In *Tyrannosaurus rex*, however, the siphoneal diverticulum does not pass through the quadrate and the quadrate diverticulum only enters the ventral opening of the pterygoid flange, and then passes with or without the siphoneal diverticulum along the medial fossa of the pterygoid flange. Likewise, the quadrate diverticulum only pneumatizes two distinct regions of the quadrate in *Acrocanthosaurus atokensis* and *Mapusaurus roseae* (Tahara and Larsson 2011).

Figure 6

Quadrate Ontogeny

Skull ontogeny has been generally poorly studied in nonavian theropod, especially in their early stage of development (Rauhut and Fechner 2005), but the ontogeny of the quadrate bone has particularly received very little attention when compared to other cranial bones (see Carr 1999; Loewen 2010). Although the quadrate of embryonic and juvenile specimens has been reported in many nonavian theropod clades such as basal Megalosauroida (Rauhut et al. 2012), Spinosauridae (Hendrickx and Mateus 2012), basal Avetheropoda (Hendrickx and Mateus 2012), Tyrannosauridae (e.g. Bakker et al. 1988; Carr and Williamson 2010; Tsuihiji et al. 2011),

578 Compsognathidae (Dal Sasso and Maganuco 2011), alvarezsaurid (Dufeu 2003), Oviraptoridae
 579 (Norell et al. 1994, 2001; Weishampel et al. 2008) and Troodontidae (Varricchio et al. 2002), the
 580 ontogenic variations of the nonavian theropod quadrate has only been investigated by Hendrickx
 581 and Mateus (2012) hitherto. These authors reported two incomplete quadrates belonging to
 582 embryos of *Lourinhanosaurus autunesi* (Figure 6) from the Upper Jurassic of Portugal (Mateus et
 583 al. 1998; de Ricqlès et al. 2001; Mateus 2005). Comparison of these two bones with the quadrates
 584 of the closest relative of *Lourinhanosaurus*, *Sinraptor dongi* (Benson 2010; Benson et al. 2010)
 585 allowed Hendrickx and Mateus (2012) to suggest that the absence of a quadrate foramen and the
 586 poor delimitation of the two mandibular condyles and the intercondylar sulcus were most likely
 587 ontogenetical features present in the allosauroid quadrate. Hendrickx and Mateus (2012) also
 588 examined the ontogenic changes of the quadrate in spinosaurids from the Cenomanian of Eastern
 589 Morocco. Based on five quadrates belonging to juvenile, subadult and adult specimens of
 590 *Spinosaurus* sp., they were able to propose a list of ontogenetical steps that they divided into
 591 juvenile, subadult and adult stages. The ontogenetical transformation occurring in the
 592 *Spinosaurus* quadrate are the clear delimitation of the ento- and ectocondyle, the intercondylar
 593 sulcus and the quadrate head, as well as the development of a ventral projection of the dorsal
 594 quadratojugal contact and the excavation of both ventral and dorsal quadratojugal contacts,
 595 allowing a firm contact between the quadrate and quadratojugal to be present in sub-adults and
 596 adults specimens.

597 Well-preserved quadrates are present in a small and large specimens of the alvarezsaurid
 598 *Shuvuuia deserti* (Chiappe et al. 1998; Dufeu 2003). Dufeu (2003), who interprets the small
 599 skull (IGM 100-1001) and the larger one (IGM 100-977) as belonging to a juvenile and adult
 600 specimens of *Shuvuuia deserti*, respectively, comprehensively described the quadrate bone but
 601 did not investigate the ontogenetic variation occurring in the skull of *Shuvuuia deserti*. Personal
 602 examination of the two specimens allowed to observe major differences in the quadrate

morphology (Figure 3.3-3.6) that could be interpreted as ontogenetic variations. The quadrate
 body of IGM 100-1001 is extremely ventro-dorsally elongated, with a prominent and narrow
 quadrate ridge along the dorsal half of the quadrate bone. On the other hand, the quadrate body of
 the larger specimen IGM 100-977 is shorter, with a relatively shallow quadrate ridge. The
 mandibular articulation of IGM 100-1001 is lateromedially expanded and the lateral part of the
 articulation is subtriangular in outline and strongly projects laterally. The mandibular articulation
 of IGM 100-977 is particularly short, subrectangular in outline in posterior view, and lacks a
 lateral projection (Figure 3.3, 3.5). This projection may however be broken and present as an
 isolated piece of bone displaced on the ventrolateral surface of the pterygoid flange (pers. obs.).
 In IGM 100-1001, the lateral process is **ventro-dorsally** long, subtriangular in lateral view, almost
 subrectangular in posterior view, and reaches the quadrate head dorsally. The most lateral corner
 projects anteriorly to contact the postorbital, and the postorbital contact of the quadrate extends
 along the anterodorsal surface of the lateral process. The ventral most part of the lateral process
 forms a small corner dorsal to the parabolic outline of the ventral part of the quadrate body. On
 the contrary, the lateral process of IGM 100-977 only projects laterally and does not extend to the
 quadrate head dorsally. The lateral margin of the lateral process is parabolic in outline, and
 neither includes a small corner ventrally, nor an important projection anteriorly. The quadrate
 foramen of IGM 100-1001, forms a large fenestra delimited by the quadrate, jugal and,
 postorbital. If an articulation between the postorbital and the lateral corner of the lateral process
 was present in IGM 100-977 as well, the quadrate foramen of this specimen would have been
 much smaller, and delimited by the ventral half of the quadrate only. The quadrate head of the
 small specimen (IGM 100-1001) is incipiently bistylic and oriented posteromedially to contact
 the squamosal dorsally, and the braincase posteromedially (Figure 3.3). The quadrate head of
 IGM 100-977 is strongly inclined laterally and seems to include a single condyle only, although

627 Dufeu (2003) noted that the quadrate head was also divided by a very weak intercondylar sulcus
628 in the adult specimen.

629 Dufeu (2003) interpreted these differences in the two specimens of *Shuvuuia deserti* as
630 ontogenetic and taphonomic variations (Dufeu pers. comm.). According to Dufeu (2003), the
631 contact between the quadrate head and the junction of the squamosal and postorbital, which is
632 unique in IGM 100-1001 among nonavian theropods, is, for instance, interpreted as a taphonomic
633 condition where the quadrate head would be taphonomically displaced laterally. Nevertheless,
634 both left and right quadrate of IGM 100-1001 occupy the same position and share similar contact
635 with the postorbital and squamosal, and the skull suffered neither compressional nor shear
636 distortions (Dufeu 2003; pers. obs.). A taphonomic displacement of the quadrate of IGM 100-
637 1001 seems therefore unlikely, and the postorbital contact of the quadrate is an autapomorphy of
638 *Shuvuuia deserti*. Despite the fact that the right quadrate of the larger specimen IGM 100-977
639 may have been subject to important postmortem compression (as the rest of the skull which is
640 ventro-dorsally flattened), the numerous differences existing between IGM 100-1001 and IGM
641 100-977 suggests that these variations may be taphonomic, ontogenetic, but also perhaps
642 taxonomic. The ontogenetic variations occurring in the spinosaurid quadrates are subtle and only
643 concern the delimitation of the condyles and quadrate head, and the reinforcement of the quadrate
644 and quadratojugal suture, so that the general morphology of the quadrate bone does not change at
645 all. If the features differentiating IGM 100-1001 and IGM 100-977 are mostly ontogenetic, the
646 quadrate of *Shuvuuia deserti* would therefore undergo a major metamorphosis during its
647 development, including a dorsoventral shortening of the quadrate bone, rotation of the quadrate
648 head from a medial to a lateral orientation, posterior displacement of the lateral process, and
649 morphological transformation of the lateral process from a subtriangular to a parabolic process.
650 Although plausible, these ontogenetic transformations seem a bit extreme, and it is reasonable to

suggest that IGM 100-977 and IGM 100-977 may belong to two taxa, perhaps two species of *Shuvuuia*.

Quadrate pneumaticity appears early in ontogeny as it has been reported in the embryo of *Troodon formosus* (Varricchio et al. 2002) and the juvenile *Tarbosaurus baatar* (Tsuihiji et al. 2011). Although absent in the embryonic specimen of *Lourinhanosaurus autunesi*, a quadrate foramen exists in many juvenile specimens of theropods such as the hatchling *Scipionyx samniticus* (Dal Sasso and Maganuco 2011) and the early posthatchling *Sciurumimus albersdoerferi* (Rauhut et al. 2012). Although the quadrate and quadratojugal are weakly articulated to each other in immature tetanurans (Hendrickx and Mateus 2012), a fusion between the quadrate and pterygoid was already present in oviraptorid embryos (Norell et al. 2001).

CONCLUSIONS

A revised nomenclature of the quadrate bone, along with a corresponding set of abbreviations, is here proposed and provides a standard set of terms for describing this cranial bone in nonavian theropod dinosaurs. The quadrate can be divided into two regional categories, the quadrate body and the pterygoid flange, and twelve anatomical sub-units, the quadrate shaft, quadrate head, quadrate ridge, quadrate foramen, lateral process, quadratojugal contact, squamosal contact, pterygoid contact, mandibular articulation, medial fossa, and posterior fossa. Although being highly variable in shape, all of these quadrate entities, with perhaps the exception of the posterior fossa, are easily recognisable inter-taxa and a description of their morphology should be provided in the literature.

A summary of the current knowledge on the quadrate function, pneumaticity and ontogeny in nonavian theropods allows some evidence about this bone to be highlighted. The quadrate of large majority of nonavian theropods is akinetic and a streptostylic quadrate may

674 have been present in some derived coelurosaurs such as the alvarezsaurid *Shuvuuia deserti* and
675 the oviraptorid *Nemegtia huangi*. A lateral movement of the rami while the mandible was
676 depressed was permitted in many theropods such as spinosaurids thanks to a helicoidal and
677 diagonally oriented intercondylar sulcus of the mandibular articulation. Likewise, the presence of
678 an intercondylar notch constricting the intercondylar sulcus in allosaurids was interpreted to be a
679 joint-stabilization zone that would improve the stability of the mandibular articulation when the
680 mouth was widely opened.

681 A pneumatic quadrate was present in members of most nonavian avetheropod clades such
682 as allosauroids, tyrannosaurids, compsognathids, therizinosauroids, ornithomimoids, oviraptorids,
683 troodontids and dromaeosaurids, in which pneumatopores typically open in the ventral part of the
684 pterygoid flange and in the medial and lateral fossae. Although the pneumatic recess invaded by
685 the quadrate diverticulum of the mandibular arch pneumatic system was linked to only one
686 pneumatopore in most avetheropods, the presence of several pneumatic openings perforating
687 different sides of the quadrate have been recorded in Neovenatoridae, Carcharodontosauridae and
688 Tyrannosauridae.

689 A poorly delimited mandibular condyles, intercondylar sulcus and quadrate head, and a
690 quadratojugal contact with a smooth surface have been interpreted as ontogenetical features in the
691 quadrate of embryonic and juvenile basal tetanurans. The development of a quadrate foramen and
692 a ventral projection of the dorsal quadratojugal contact seem also to happen during ontogeny in
693 allosauroids and spinosaurids, respectively. On the other hand, pneumaticity and a strong suture
694 between the quadrate and quadratojugal appear early in ontogeny, in the embryonic stage of
695 coelurosaurs. Finally, based on the quadrate morphology, the two specimens assigned to
696 *Shuvuuia deserti*, considered by some as juvenile and adult specimens, may belong to two
697 distinct taxa.

ACKNOWLEDGMENTS

The quadrate of many nonavian theropods were examined at first hand in several institutions and access to the material was possible thanks to P. Sereno (Uni. Chicago), P. Makovicky (FMNH), W. Simpson (FMNH), M. Lamanna (CMNH), A. Henrici (CMNH), M. Carrano (NMNH), M. Brett-Surman (NMNH), S. Chapman (NHM), P. Barrett (NHM), P. Jeffery (OUMNH), S. Hutt (MIW), R. Allain (MNHN), R. Schoch (SMNS), H.-J. Siber (SMA), C. Dal Sasso (MSNM), A. Kramarz (MACN), F. Novas (MACN), R. Barbieri (MPCA), L. Salgado (MUCPv), J. Ignacio Canale (MUCPv-CH), R. Coria (MCF-PVPH), C. Succar (MCF-PVPH), J. Calvo (CePaLB), R. Martínez (PVSJ), C. Mehling (AMNH), M. Norell (AMNH), D. Krauze (SBU), J. Groenke (SBU), P. Brinkman (NCSM), and L. Zanno (NCSM). Photographs of theropod quadrates were kindly shared by M. Lamanna (CMNH), M. Ezcurra (MACNBR), M. Carrano (USNM), E. Buffetaut (CNRS), M. Ellison (AMNH), L. Witmer (Uni. Ohio), S. Brusatte (Uni. Edinburgh), R. Benson (Uni. Cambridge), C. Foth (BSPG), P. Currie (Uni. Alberta), J. Canale (MUCPv-CH), P. Barrett (BMNH), J. Choiniere (Uni. Witwatersrand), D. Eddy (Uni. Texas), P. Viscardi (Horniman), S. Nesbitt (Uni. Texas), Y. Kobayashi (HUM), R. Tahara (McGill Uni.), R. Pei (AMNH), C. Dal Sasso (MSNM), P. Sereno (Uni. Chicago), C. Abraczinskas (Uni. Chicago), N. Smith (Uni. Chicago), L. Zanno (FMNH), R. Tykoski (MNSD), D. Burnham (Uni. Kansas), P. Asaroff (MACNBR), R. Irmis (UMNH), V. Shneider (NCMNS), C. Brochu (Uni. Iowa), S. Lautenschlager (Uni. Bristol), M. Mortimer, K. Peyer (MNHN), R. Molnar (MNA), and the authors would like to address their sincere thanks to all of these people. A special thanks to D. Dufeu for sharing his MSc thesis on *Shuvuuia*. C.H. dedicates this paper to R. Bec.

REFERENCES

- 720 Bakker, R.T. 1998. Brontosaur killers: late Jurassic allosaurids as sabre-tooth cat analogues. *Gaia*
721 15:145–158.
- 722 Bakker, R.T., Williams, M. and Currie, P.J. 1988. *Nanotyrannus*, a new genus of pygmy
723 tyrannosaur, from the latest Cretaceous of Montana. *Hunteria* 1:1–30.
- 724 Balanoff, A.M. and Norell, M.A. 2012. Osteology of *Khaan mckennai* (Oviraptorosauria:
725 Theropoda). *Bulletin of the American Museum of Natural History* 1–77.
726 doi:10.1206/803.1.
- 727 Barbosa, A. 1990. Identification key of Iberian waders (Charadriiformes) based on the os
728 quadratum. *Miscellània Zoològica* 14:181–185.
- 729 Barsbold, R. and Osmólska, H. 1999. The skull of *Velociraptor* (Theropoda) from the Late
730 Cretaceous of Mongolia. *Acta Palaeontologica Polonica* 44:189–219.
- 731 Barsbold, R., Osmólska, H. and Kurzanov, S.M. 1987. On a new troodontid (Dinosauria,
732 Theropoda) from the Early Cretaceous of Mongolia. *Acta Palaeontologica Polonica*
733 32:121–132.
- 734 Baumel, J.J. and Witmer, L.M. 1993. Osteologia, p. 45–132. In Baumel, J.J. (ed.), *Handbook of*
735 *Avian Anatomy : Nomina Anatomica Avium*. Nuttall Ornithological Club, Cambridge.
- 736 Benson, R., Carrano, M. and Brusatte, S. 2010. A new clade of archaic large-bodied predatory
737 dinosaurs (Theropoda: Allosauroidae) that survived to the latest Mesozoic.
738 *Naturwissenschaften* 97:71–78. doi:10.1007/s00114-009-0614-x.
- 739 Benson, R.B.J. 2010. A description of *Megalosaurus bucklandii* (Dinosauria: Theropoda) from
740 the Bathonian of the UK and the relationships of Middle Jurassic theropods. *Zoological*
741 *Journal of the Linnean Society* 158:882–935.
- 742 Benton, M.J. 2005. *Vertebrate Palaeontology (Third edition)*. Wiley-Blackwell.
- 743 Bock, W.J. 1964. Kinetics of the avian skull. *Journal of Morphology* 114:1–41.
744 doi:10.1002/jmor.1051140102.

- 745 Bock, W.J. 1999. Avian cranial kinesis revisited. *Acta Ornithologica* 34:115–122.
- 746 Bock, W.J. 2000. The evolution of avian cranial kinesis, p. 1, 191–201. In Zhou, Z. and Zhang, F.
747 (eds.), *Proceedings of the 5th Symposium of the Society of Avian Paleontology and*
748 *Evolution, Beijing, 1–4 June 2000. Science Press, Beijing.*
- 749 Bonaparte, J.F., Novas, F.E. and Coria, R.A. 1990. *Carnotaurus sastrei* Bonaparte, the horned,
750 lightly built carnosaur from the Middle Cretaceous of Patagonia. *Natural History*
751 *Museum of Los Angeles County Contributions in Science* 416:1–42.
- 752 Bout, R.G. and Zweers, G.A. 2001. The role of cranial kinesis in birds. *Comparative*
753 *Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 131:197–205.
754 doi:10.1016/S1095-6433(01)00470-6.
- 755 Britt, B.B. 1991. Theropods of Dry Mesa Quarry (Morrison Formation, Late Jurassic), Colorado,
756 with emphasis on the osteology of *Torvosaurus tanneri*. *Brigham Young University*
757 *Geology Studies* 37:1–72.
- 758 Brochu, C.A. 2003. Osteology of *Tyrannosaurus rex*: Insights from a nearly complete skeleton
759 and high-resolution computed tomographic analysis of the skull. *Journal of Vertebrate*
760 *Paleontology* 22:1–138. doi:10.1080/02724634.2003.10010947.
- 761 Brusatte, S.L. 2012. *Dinosaur Paleobiology*. Wiley-Blackwell.
- 762 Brusatte, S.L., Carr, T.D. and Norell, M.A. 2012. The osteology of *Alioramus*, a gracile and long-
763 snouted tyrannosaurid (Dinosauria: Theropoda) from the Late Cretaceous of Mongolia.
764 *Bulletin of the American Museum of Natural History* 366:1–197. doi:10.1206/770.1.
- 765 Brusatte, S.L., Chure, D.J., Benson, R.B.J. and Xu, X. 2010. The osteology of *Shaochilong*
766 *maortuensis*, a carcharodontosaurid (Dinosauria: Theropoda) from the Late Cretaceous of
767 Asia. *Zootaxa* 2334:1–46.
- 768 Bühler, P. 1981. Functional anatomy of the avian jaw apparatus. *Form and function in birds*
769 2:439–468.

- 770 Bühler, P., Hecht, M.K., Ostrom, J.H., Viohl, G. and Wellnhofer, P. 1985. On the morphology of
771 the skull of Archaeopteryx. *The Beginnings of Birds* 135–140.
- 772 Bühler, P., Martin, L.D. and Witmer, L.M. 1988. Cranial kinesis in the Late Cretaceous birds
773 Hesperornis and Paraesperornis. *The Auk* 105:111–122.
- 774 Burnham, D.A. 2004. New Information on *Bambiraptor feinbergi* (Theropoda: Dromaeosauridae)
775 from the Late Cretaceous of Montana, p. 67–111. In Currie, P.J., Koppelhus, E.B., Shugar,
776 M.A., and Wright, J.L. (eds.), *Feathered Dragons: Studies on the Transition from*
777 *Dinosaurs to Birds*. Indiana University Press, Bloomington, Indiana.
- 778 Carr, T.D. 1996. Cranial osteology and craniofacial ontogeny of Tyrannosauridae: Dinosauria:
779 Theropoda from the Dinosaur Park Formation (Judith River Group, Upper Cretaceous,
780 Campanian) of Alberta. MSc. Dissertation, University of Toronto, Toronto, Ontario,
781 Canada.
- 782 Carr, T.D. 1999. Craniofacial ontogeny in Tyrannosauridae (Dinosauria, Coelurosauria). *Journal*
783 *of Vertebrate Paleontology* 19:497–520. doi:10.1080/02724634.1999.10011161.
- 784 Carr, T.D. and Williamson, T.E. 2010. *Bistahieversor sealeyi*, gen. et sp. nov., a new
785 tyrannosauroid from New Mexico and the origin of deep snouts in Tyrannosauroida.
786 *Journal of vertebrate Paleontology* 30:1–16.
- 787 Carrano, M.T., Benson, R.B.J. and Sampson, S.D. 2012. The phylogeny of Tetanurae
788 (Dinosauria: Theropoda). *Journal of Systematic Palaeontology* 10:211–300.
789 doi:10.1080/14772019.2011.630927.
- 790 Carrano, M.T., Loewen, M.A. and Sertich, J.J.W. 2011. New materials of *Masiakasaurus*
791 *knopfleri* Sampson, Carrano, and Forster, 2001, and implications for the morphology of
792 the Noasauridae (Theropoda: Ceratosauria). *Smithsonian Contributions to Paleobiology*
793 95:1–53.
- 794 Carroll, R.L. 1988. *Vertebrate Paleontology and Evolution*. W. H. Freeman and Company.

- 795 Charig, A.J. and Milner, A.C. 1997. *Baryonyx walkeri*, a fish-eating dinosaur from the Wealden of
796 Surrey. *Bulletin of the Natural History Museum* 53:11–70.
- 797 Chatterjee, S. 1991. Cranial anatomy and relationships of a new Triassic bird from Texas.
798 *Philosophical Transactions: Biological Sciences* 332:277–342.
- 799 Chatterjee, S. 1995. The Triassic bird Protoavis. *Archaeopteryx* 13:15–31.
- 800 Chatterjee, S. 1997. *The rise of birds: 225 million years of evolution*. Johns Hopkins University
801 Press Baltimore, Maryland.
- 802 Chiappe, L.M. 2001. Phylogenetic relationships among basal birds, p. 125–139. In Gauthier, J.
803 and Gall, L.F. (eds.), *New perspectives on the origin and early evolution of birds:*
804 *proceedings of the international symposium in honor of John H. Ostrom*. Yale Univ
805 Peabody Museum.
- 806 Chiappe, L.M., Norell, M.A. and Clark, J.M. 1998. The skull of a relative of the stem-group bird
807 Mononykus. *Nature* 392:275–278. doi:10.1038/32642.
- 808 Chiappe, L.M., Norell, M.A. and Clark, J.M. 2002. The Cretaceous, short-armed
809 Alvarezsauridae: *Mononykus* and its kin, p. 87–120. In Chiappe, L.M. and Witmer, L.M.
810 (eds.), *Mesozoic birds: above the heads of dinosaurs*. University of California Press.
- 811 Choiniere, J.N., Clark, J.M., Forster, C.A., Norell, M.A., Eberth, D.A., Erickson, G.M., Chu, H.
812 and Xu, X. 2014. A juvenile specimen of a new coelurosaur (Dinosauria: Theropoda)
813 from the Middle–Late Jurassic Shishugou Formation of Xinjiang, People’s Republic of
814 China. *Journal of Systematic Palaeontology* 12:177–215.
815 doi:10.1080/14772019.2013.781067.
- 816 Choiniere, J.N., Clark, J.M., Forster, C.A. and Xu, X. 2010. A basal coelurosaur (Dinosauria:
817 Theropoda) from the Late Jurassic (Oxfordian) of the Shishugou Formation in Wucuiwan,
818 People’s Republic of China. *Journal of Vertebrate Paleontology* 30:1773–1796.
819 doi:10.1080/02724634.2010.520779.

- 820 Clark, J.M., Norell, M.A. and Rowe, T. 2002. Cranial anatomy of *Citipati osmolskae* (Theropoda,
821 Oviraptorosauria), and a reinterpretation of the holotype of *Oviraptor philoceratops*.
822 *American Museum Novitates* 3364:1–24.
- 823 Clark, J.M., Perle, A. and Norell, M. 1994. The skull of *Erlicosaurus andrewsi*, a late Cretaceous
824 “Segnosaur” (Theropoda, Therizinosauridae) from Mongolia. *American Museum*
825 *Novitates* 3115:1–39.
- 826 Colbert, E.H. 1989. The Triassic dinosaur *Coelophysis*. *Museum of Northern Arizona Bulletin*
827 57:1–174.
- 828 Colbert, E.H. and Russell, D.A. 1969. The small Cretaceous dinosaur *Dromaeosaurus*. *American*
829 *Museum Novitates* 2380:1–49.
- 830 Coria, R.A. and Currie, P.J. 2006. A new carcharodontosaurid (Dinosauria, Theropoda) from the
831 Upper Cretaceous of Argentina. *Geodiversitas* 28:71–118.
- 832 Coria, R.A. and Salgado, L. 1998. A basal Abelisauria Novas, 1992 (Theropoda-Ceratosauria)
833 from the Cretaceous of Patagonia, Argentina. *Gaia* 15:89–102.
- 834 Currie, P.J. 1995. New information on the anatomy and relationships of *Dromaeosaurus*
835 *albertensis* (Dinosauria: Theropoda). *Journal of Vertebrate Paleontology* 15:576–591.
836 doi:10.1080/02724634.1995.10011250.
- 837 Currie, P.J. 2003. Cranial anatomy of tyrannosaurid dinosaurs from the Late Cretaceous of
838 Alberta, Canada. *Acta Palaeontologica Polonica* 48:191–226.
- 839 Currie, P.J. 2006. On the quadrate of *Sinraptor dongi* (Theropoda: Allosauroidae) from the Late
840 Jurassic of China. *Mesozoic and Cenozoic Vertebrates and Paleoenvironments. Tributes*
841 *to the career of Prof. Dan Grigorescu* 111–115.
- 842 Currie, P.J. and Carpenter, K. 2000. A new specimen of *Acrocanthosaurus atokensis* (Theropoda,
843 Dinosauria) from the Lower Cretaceous Antlers Formation (Lower Cretaceous, Aptian) of
844 Oklahoma, USA. *Geodiversitas* 22:207–246.

- 845 Currie, P.J. and Chen, P. 2001. Anatomy of *Sinosauropteryx prima* from Liaoning, northeastern
846 China. *Canadian Journal of Earth Sciences* 38:1705–1727. doi:10.1139/e01-050.
- 847 Currie, P.J. and Zhao, X.-J. 1993. A new troodontid (Dinosauria, Theropoda) braincase from the
848 Dinosaur Park Formation (Campanian) of Alberta. *Canadian Journal of Earth Sciences*
849 30:2231–2247. doi:10.1139/e93-194.
- 850 Dufeu, D.L. 2003. The cranial anatomy of the theropod dinosaur *Shuvuuia deserti*
851 (Coelurosauria: Alvarezsauridae), and its bearing upon coelurosaurian phylogeny. MSc.
852 Dissertation, University of Texas, Austin, Texas.
- 853 Eddy, D.R. and Clarke, J.A. 2011. New information on the cranial anatomy of *Acrocanthosaurus*
854 *atokensis* and its implications for the phylogeny of Allosauroidea (Dinosauria:
855 Theropoda). *PLoS ONE* 6:e17932. doi:10.1371/journal.pone.0017932.
- 856 Elzanowski, A., Paul, G.S. and Stidham, T.A. 2001. An avian quadrate from the Late Cretaceous
857 Lance formation of Wyoming. *Journal of Vertebrate Paleontology* 20:712–719.
- 858 Elzanowski, A. and Stidham, T.A. 2010. Morphology of the quadrate in the Eocene anseriform
859 *Presbyornis* and extant galloanserine birds. *Journal of Morphology* 271:305–323.
860 doi:10.1002/jmor.10799.
- 861 Fisher, H.I. 1955. Some aspects of the kinetics in the jaws of birds. *The Wilson Bulletin* 175–188.
- 862 Foth, C. and Rauhut, O. 2012. Macroevolutionary and morphofunctional patterns in theropod
863 skulls: a morphometric approach. *Acta Palaeontologica Polonica*.
864 doi:10.4202/app.2011.0145.
- 865 Fuchs, A. 1954. On the correlation between the skull structure and the muscles in the male
866 *Phasianus colchicus* L. IV. The attachment of the musculus protractor quadrati et
867 pterygoidei and of the musculus depressor mandibulae. *Koninklijke Nederlandse*
868 *Akademie van Wetenschappen, Proceedings, C* 57:666–672.

- 869 Gauthier, J. 1986. Saurischian monophyly and the origin of birds, p. 8, 1–55. In Padian, K. (ed.),
870 *The origin of Birds and the Evolution of Flight*. Memoirs of the California Academy of
871 Sciences, San Francisco, California.
- 872 Gilmore, C.W. 1920. Osteology of the carnivorous Dinosauria in the United State National
873 museum: with special reference to the genera *Antrodemus* (*Allosaurus*) and *Ceratosaurus*.
874 *Bulletin of the United States National Museum* 110:1–159.
- 875 Gold, M.E.L., Brusatte, S. and Norell, M. 2013. The cranial pneumatic sinuses of the
876 tyrannosaurid *Alioramus* (Dinosauria, Theropoda) and the evolution of cranial
877 pneumaticity in theropod dinosaurs. *American Museum Novitates* 3790:1–46.
- 878 Gussekloo, S.W.S. and Bout, R.G. 2005. Cranial kinesis in palaeognathous birds. *Journal of*
879 *Experimental Biology* 208:3409–3419. doi:10.1242/jeb.01768.
- 880 Hendrickx, C. and Buffetaut, E. 2008. Functional interpretation of spinosaurid quadrates
881 (Dinosauria: Theropoda) from the Mid-Cretaceous of Morocco. *56th Annual Symposium*
882 *of Vertebrate Palaeontology and Comparative Anatomy. Dublin (September 2nd-6th*
883 *2008)* 25–26.
- 884 Hendrickx, C. and Mateus, O. 2012. Ontogenetical changes in the quadrate of basal tetanurans, p.
885 20, 101–104. In Royo-Torres, R., Gascó, F., and Alcalá, L. (eds.), *10th Annual Meeting of*
886 *the European Association of Vertebrate Palaeontologists. ¡Fundamental!*
- 887 Hoese, W.J. and Westneat, M.W. 1996. Biomechanics of cranial kinesis in birds: testing linkage
888 models in the white-throated sparrow (*Zonotrichia albicollis*). *Journal of Morphology*
889 227:305–320.
- 890 Holliday, C.M. 2009. New insights into dinosaur jaw muscle anatomy. *The Anatomical Record:*
891 *Advances in Integrative Anatomy and Evolutionary Biology* 292:1246–1265.
892 doi:10.1002/ar.20982.

- 893 Holliday, C.M. and Witmer, L.M. 2008. Cranial kinesis in dinosaurs: intracranial joints,
894 protractor muscles, and their significance for cranial evolution and function in diapsids.
895 *Journal of Vertebrate Paleontology* 28:1073–1088. doi:10.1671/0272-4634-28.4.1073.
- 896 Holtz, T.R.J. 1998. A new phylogeny of the carnivorous dinosaurs. *Gaia* 15:5–61.
- 897 Holtz, T.R.J. 2003. Dinosaur Predation: Evidence and Ecomorphology, p. 325–340. In Kelley,
898 P.H., Kowalewski, M., and Hansen, T.A. (eds.), *Predator—Prey Interactions in the Fossil*
899 *Record*. Topics in Geobiology. Springer US.
- 900 Holtz, T.R.J., Molnar, R.E. and Currie, P.J. 2004. Basal Tetanurae, p. 71–110. In Weishampel,
901 D.B., Dodson, P., and Osmólska, H. (eds.), *The Dinosauria. Second Edition*. University of
902 California Press, Berkeley, California.
- 903 Hone, D.W.E. and Rauhut, O.W.M. 2010. Feeding behaviour and bone utilization by theropod
904 dinosaurs. *Lethaia* 43:232–244. doi:10.1111/j.1502-3931.2009.00187.x.
- 905 Kobayashi, Y. and Barsbold, R. 2005. Reexamination of a primitive ornithomimosaur,
906 *Garudimimus brevipes* Barsbold, 1981 (Dinosauria: Theropoda), from the Late
907 Cretaceous of Mongolia. *Canadian Journal of Earth Sciences* 42:1501–1521.
908 doi:10.1139/e05-044.
- 909 Kobayashi, Y. and Lü, J.-C. 2003. A new ornithomimid dinosaur with gregarious habits from the
910 Late Cretaceous of China. *Acta Palaeontologica Polonica* 48:235–259.
- 911 Kundrát, M. and Janáček, J. 2007. Cranial pneumatization and auditory perceptions of the
912 oviraptorid dinosaur *Conchoraptor gracilis* (Theropoda, Maniraptora) from the Late
913 Cretaceous of Mongolia. *Naturwissenschaften* 94:769–778. doi:10.1007/s00114-007-
914 0258-7.
- 915 Larson, P.L. 2008. Atlas of the skull bones of *Tyrannosaurus rex*, p. 233–243. In Larson, P.L. and
916 Carpenter, K. (eds.), *Tyrannosaurus rex, the Tyrant King*. Indiana University Press,
917 Bloomington, Indiana.

- 918 Li, D., Norell, M.A., Gao, K.Q., Smith, N.D. and Makovicky, P.J. 2010. A longirostre
919 tyrannosauroid from the Early Cretaceous of China. *Proceedings of the Royal Society B:*
920 *Biological Sciences* 277:183–190.
- 921 Loewen, M.A. 2010. Variation in the Late Jurassic theropod dinosaur *Allosaurus*: Ontogenetic,
922 functional, and taxonomic implications. Ph.D. Dissertation, The University of Utah,
923 Texas, Utah, USA.
- 924 Lowe, P.R. 1926. More notes on the quadrate as a factor in avian classification. *Ibis* 68:152–188.
925 doi:10.1111/j.1474-919X.1926.tb07563.x.
- 926 Lü, J. 2003. A new oviraptorosaurid (Theropoda: Oviraptorosauria) from the Late Cretaceous of
927 southern China. *Journal of Vertebrate Paleontology* 22:871–875. doi:10.1671/0272-
928 4634(2002)022[0871:ANOTOF]2.0.CO;2.
- 929 Lü, J. 2005. Oviraptorid dinosaurs from southern China. Ph.D. Dissertation, Southern Methodist
930 University, Dallas, Texas, USA.
- 931 Lü, J., Tomida, Y., Azunia, Y., Dong, Z. and Lee, Y.N. 2004. New Oviraptorid Dinosaur
932 (Dinosauria: Oviraptorosauria) from the Nemegt Formation of Southwestern Mongolia.
933 *Bulletin of the National Science Museum: Geology & paleontology* 30:95–130.
- 934 Madsen, J.H. 1976. *Allosaurus fragilis*: A Revised Osteology. *Utah Geological Survey Bulletin*
935 109:1–177.
- 936 Madsen, J.H. and Welles, S.P. 2000. *Ceratosaurus* (Dinosauria, Theropoda): a revised osteology.
937 *Utah Geological Survey, Miscellaneous Publication* 00-2:1–89.
- 938 Makovicky, P.J., Apesteguía, S. and Agnolín, F.L. 2005. The earliest dromaeosaurid theropod
939 from South America. *Nature* 437:1007–1011. doi:10.1038/nature03996.
- 940 Maryńska, T. and Osmólska, H. 1997. The quadrate of oviraptorid dinosaurs. *Acta*
941 *Palaeontologica Polonica* 42:361–371.

- 942 Mateus, I., Mateus, H., Antunes, M.T., Mateus, O., Taquet, P., Ribeiro, V. and Manuppella, G.
943 1998. Upper Jurassic theropod dinosaur embryos from Lourinhã (Portugal). *Memórias da*
944 *Academia das Ciências de Lisboa* 37:101–110.
- 945 Mateus, O. 2005. Dinossauros do Jurássico Superior de Portugal, com destaque para os
946 saurísquios. Ph.D. Dissertation, Universidade Nova de Lisboa, Lisbon, Portugal.
- 947 Meekangvan, P., Barhorst, A., Burton, T.D., Chatterjee, S. and Schovanec, L. 2006. Nonlinear
948 dynamical model and response of avian cranial kinesis. *Journal of Theoretical Biology*
949 240:32–47. doi:10.1016/j.jtbi.2005.08.027.
- 950 Metzger, K. 2002. Cranial kinesis in lepidosaurs: skulls in motion. *Topics in functional and*
951 *ecological vertebrate morphology. Maastricht: Shaker Publishing.* p 15–46.
- 952 Molnar, R.E. 1991. The cranial morphology of *Tyrannosaurus rex*. *Palaeontographica Abteilung*
953 *A* 217:137–176.
- 954 Molnar, R.E. 1998. Mechanical factors in the design of the skull of *Tyrannosaurus rex* (Osborn,
955 1905). *Gaia* 15:193–218.
- 956 Nitzsch, C.L. 1816. Über die bewegung des oberkiefers der vögel. *Deutsches Archiv für die*
957 *Physiologie* 2:361–380.
- 958 Norell, M.A., Clark, J.M. and Chiappe, L.M. 2001. An embryonic oviraptorid (Dinosauria:
959 Theropoda) from the Upper Cretaceous of Mongolia. *American Museum Novitates*
960 3315:1–20.
- 961 Norell, M.A., Clark, J.M., Demberelyin, D., Rhinchen, B., Chiappe, L.M., Davidson, A.R.,
962 McKenna, M.C., Altangerel, P. and Novacek, M.J. 1994. A Theropod Dinosaur Embryo
963 and the Affinities of the Flaming Cliffs Dinosaur Eggs. *Science* 266:779–782.
964 doi:10.1126/science.266.5186.779.

- 965 Norell, M.A., Clark, J.M., Turner, A.H., Makovicky, P.J., Barsbold, R. and Rowe, T. 2006. A new
966 dromaeosaurid theropod from Ukhaa Tolgod (Ömnögovi, Mongolia). *American Museum*
967 *Novitates* 3545:1–51. doi:10.1206/0003-0082(2006)3545[1:ANDTFU]2.0.CO;2.
- 968 Norell, M.A. and Hwang, S.H. 2004. A troodontid dinosaur from Ukhaa Tolgod (Late Cretaceous
969 Mongolia). *American Museum Novitates* 3446:1–9.
- 970 Norell, M.A., Makovicky, P.J., Bever, G.S., Balanoff, A.M., Clark, J.M., Barsbold, R. and Rowe,
971 T. 2009. A review of the Mongolian Cretaceous dinosaur *Saurornithoides* (Troodontidae:
972 Theropoda). *American Museum Novitates* 3654:1–63. doi:10.1206/648.1.
- 973 Novas, F.E., Agnolín, F.L., Ezcurra, M.D., Porfiri, J. and Canale, J.I. 2013. Evolution of the
974 carnivorous dinosaurs during the Cretaceous: The evidence from Patagonia. *Cretaceous*
975 *Research* 45:174–215. doi:10.1016/j.cretres.2013.04.001.
- 976 Perle, A., Chiappe, L.M. and Barsbold, R. 1994. Skeletal morphology of *Mononykus olecranus*
977 (Theropoda: Avialae) from the Late Cretaceous of Mongolia. *American Museum*
978 *Novitates* 3105:1–29.
- 979 Rauhut, O.W.M. 2003. The interrelationships and evolution of basal theropod dinosaurs. *Special*
980 *Papers in Palaeontology* 69:1–213.
- 981 Rauhut, O.W.M. and Fechner, R. 2005. Early development of the facial region in a non-avian
982 theropod dinosaur. *Proceedings of the Royal Society B: Biological Sciences* 272:1179–
983 1183. doi:10.1098/rspb.2005.3071.
- 984 Rauhut, O.W.M., Foth, C., Tischlinger, H. and Norell, M.A. 2012. Exceptionally preserved
985 juvenile megalosauroid theropod dinosaur with filamentous integument from the Late
986 Jurassic of Germany. *Proceedings of the National Academy of Sciences* 109:11746–11751.
987 doi:10.1073/pnas.1203238109.
- 988 Rauhut, O.W.M., Milner, A.C. and Moore-Fay, S. 2010. Cranial osteology and phylogenetic
989 position of the theropod dinosaur *Proceratosaurus bradleyi* (Woodward, 1910) from the

Middle Jurassic of England. *Zoological Journal of the Linnean Society* 158:155–195.

doi:10.1111/j.1096-3642.2009.00591.x.

Rayfield, E.J. 2005. Using finite-element analysis to investigate suture morphology: A case study

using large carnivorous dinosaurs. *The Anatomical Record Part A: Discoveries in*

Molecular, Cellular, and Evolutionary Biology 283A:349–365. doi:10.1002/ar.a.20168.

De Ricqlès, A., Mateus, O., Antunes, M.T. and Taquet, P. 2001. Histomorphogenesis of embryos

of Upper Jurassic theropods from Lourinhã (Portugal). *Comptes Rendus de l'Académie*

des Sciences-Series IIA-Earth and Planetary Science 332:647–656.

Rieppel, O. 2006. The merits of similarity reconsidered. *Systematics and Biodiversity* 4:137–147.

doi:10.1017/S1477200005001830.

Russell, D.A. 1972. Ostrich dinosaurs from the Late Cretaceous of Western Canada. *Canadian*

Journal of Earth Sciences 9:375–402. doi:10.1139/e72-031.

Sadleir, R., Barrett, P.M. and Powell, H.P. 2008. The anatomy and systematics of

Eustreptospondylus oxoniensis, a theropod dinosaur from the Middle Jurassic of

Oxfordshire, England. *Monograph of the Palaeontographical Society, London* 160:1–82.

Sakamoto, M. 2008. Bite force and the evolution of feeding function in birds, dinosaurs and cats.

Ph.D. Dissertation, University of Bristol, Bristol, U.K.

Samejima, M. and Otsuka, J. 1987. Observations on the Quadrate of Birds. *Japanese Journal of*

Ornithology 35:129–144.

Sampson, S.D. and Witmer, L.M. 2007. Craniofacial anatomy of *Majungasaurus crenatissimus*

(Theropoda: Abelisauridae) from the Late Cretaceous of Madagascar. *Journal of*

Vertebrate Paleontology 27:32–104. doi:10.1671/0272-

4634(2007)27[32:CAOMCT]2.0.CO;2.

Sanders, R.K. and Smith, D.K. 2005. The endocranium of the theropod dinosaur *Ceratosaurus*

studied with computed tomography. *Acta Palaeontologica Polonica* 50:601.

- 1015 Dal Sasso, C. and Maganuco, S. 2011. *Scipionyx samniticus* (Theropoda: Compsognathidae) from
- 1016 the Lower Cretaceous of Italy: osteology, ontogenetic assessment, phylogeny, soft tissue
- 1017 anatomy, taphonomy and palaeobiology. *Memorie della Società Italiana di Scienze*
- 1018 *Naturali e del Museo Civico di Storia Naturale di Milano* 37:1–281.
- 1019 Schwarz, D., Frey, E. and Meyer, C.A. 2007. Pneumaticity and soft-tissue reconstructions in the
- 1020 neck of diplodocid and dicraeosaurid sauropods. *Acta Palaeontologica Polonica* 52:167.
- 1021 Sedlmayr, J.C. 2002. Anatomy, evolution, and functional significance of cephalic vasculature in
- 1022 Archosauria. Ph.D. dissertation, Ohio University, Athens, Ohio, USA.
- 1023 Sereno, P.C. 2001. Alvarezsaurids: birds or ornithomimosaurs, p. 69–98. In Gauthier, J. and Gall,
- 1024 L.F. (eds.), *New Perspectives on the Origin and Early Evolution of Birds: Proceedings of*
- 1025 *the International Symposium in Honor of John H. Ostrom*. Yale Univ Peabody Museum.
- 1026 Sereno, P.C., Beck, A.L., Dutheil, D.B., Gado, B., Larsson, H.C.E., Lyon, G.H., Marcot, J.D.,
- 1027 Rauhut, O.W.M., Sadleir, R.W., Sidor, C.A., Varricchio, D.D., Wilson, G.P. and Wilson,
- 1028 J.A. 1998. A long-snouted predatory dinosaur from Africa and the evolution of
- 1029 spinosaurids. *Science* 282:1298–1302. doi:10.1126/science.282.5392.1298.
- 1030 Sereno, P.C., Martinez, R.N., Wilson, J.A., Varricchio, D.J., Alcober, O.A. and Larsson, H.C.E.
- 1031 2008. Evidence for avian intrathoracic air sacs in a new predatory dinosaur from
- 1032 Argentina. *PLoS ONE* 3:e3303. doi:10.1371/journal.pone.0003303.
- 1033 Sereno, P.C. and Novas, F.E. 1994. The skull and neck of the basal theropod *Herrerasaurus*
- 1034 *ischigualastensis*. *Journal of Vertebrate Paleontology* 13:451–476.
- 1035 doi:10.1080/02724634.1994.10011525.
- 1036 Smith, D. 1992. The type specimen of *Oviraptor philoceratops*, a theropod dinosaur from the
- 1037 Upper Cretaceous of Mongolia. *Neues Jahrbuch für Geologie und Paläontologie,*
- 1038 *Abhandlungen* 186:365–388.

- 1039 Smith, N.D., Makovicky, P.J., Hammer, W.R. and Currie, P.J. 2007. Osteology of
1040 *Cryolophosaurus ellioti* (Dinosauria: Theropoda) from the Early Jurassic of Antarctica
1041 and implications for early theropod evolution. *Zoological Journal of the Linnean Society*
1042 151:377–421. doi:10.1111/j.1096-3642.2007.00325.x.
- 1043 Tahara, R. and Larsson, H.C.E. 2011. Cranial pneumatic anatomy of *Ornithomimus*
1044 *edmontonicus* (Ornithomimidae: Theropoda). *Journal of Vertebrate Paleontology* 31:127–
1045 143. doi:10.1080/02724634.2011.539646.
- 1046 Therrien, F., Henderson, D.M. and Ruff, C.B. 2005. Bite me: biomechanical models of theropod
1047 mandibles and implications for feeding behavior, p. 179–237. In Carpenter, K. (ed.), *The*
1048 *Carnivorous Dinosaurs*. Indiana University Press, Bloomington, Indiana.
- 1049 Tsuihiji, T., Watabe, M., Tsogtbaatar, K., Tsubamoto, T., Barsbold, R., Suzuki, S., Lee, A.H.,
1050 Ridgely, R.C., Kawahara, Y. and Witmer, L.M. 2011. Cranial osteology of a juvenile
1051 specimen of *Tarbosaurus bataar* (Theropoda, Tyrannosauridae) from the Nemegt
1052 Formation (Upper Cretaceous) of Bugin Tsav, Mongolia. *Journal of Vertebrate*
1053 *Paleontology* 31:497–517. doi:10.1080/02724634.2011.557116.
- 1054 Turner, A.H., Hwang, S.H. and Norell, M.A. 2007. A small derived theropod from Öösh, Early
1055 Cretaceous, Baykhangor Mongolia. *American Museum Novitates* 3557:1–27.
- 1056 Turner, A.H., Makovicky, P.J. and Norell, M. 2012. A review of dromaeosaurid systematics and
1057 paravian phylogeny. *Bulletin of the American Museum of Natural History* 371:1–206.
- 1058 Turner, A.H., Pol, D. and Norell, M.A. 2011. Anatomy of *Mahakala omnogovae* (Theropoda:
1059 Dromaeosauridae), Tögrögiin Shiree, Mongolia. *American Museum Novitates* 3722:1–66.
1060 doi:10.1206/3722.2.
- 1061 Varricchio, D.J. 1997. Troodontidae, p. 749–754. In Currie, P.J. and Padian, K. (eds.),
1062 *Encyclopedia of Dinosaurs*. Academic Press, San Diego, California.

- 1063 Varricchio, D.J., Horner, J.R. and Jackson, F.D. 2002. Embryos and eggs for the Cretaceous
1064 theropod dinosaur *Troodon formosus*. *Journal of Vertebrate Paleontology* 22:564–576.
1065 doi:10.1671/0272-4634(2002)022[0564:EAEFTC]2.0.CO;2.
- 1066 Vickers-Rich, P., Chiappe, L.M. and Kurzanov, S. 2002. The enigmatic birdlike dinosaur
1067 *Avimimus portentosus*, p. 65–86. In Chiappe, L.M. and Witmer, L.M. (eds.), *Mesozoic*
1068 *birds: above the heads of dinosaurs*. University of California Press, Berkeley/Los
1069 Angeles/London.
- 1070 Weishampel, D.B., Fastovsky, D.E., Watabe, M., Varricchio, D., Jackson, F., Tsogtbaatar, K. and
1071 Barsbold, R. 2008. New oviraptorid embryos from Bugin-Tsav, Nemegt Formation
1072 (Upper Cretaceous), Mongolia, with insights into their habitat and growth. *Journal of*
1073 *Vertebrate Paleontology* 28:1110–1119. doi:10.1671/0272-4634-28.4.1110.
- 1074 Welles, S.P. 1984. *Dilophosaurus wetherilli* (Dinosauria, Theropoda). Osteology and
1075 comparisons. *Palaeontographica Abteilung A* 185:85–180.
- 1076 Witmer, L.M. 1990. The craniofacial air sac system of Mesozoic birds (Aves). *Zoological*
1077 *Journal of the Linnean Society* 100:327–378. doi:10.1111/j.1096-3642.1990.tb01865.x.
- 1078 Witmer, L.M. 1997. The evolution of the antorbital cavity of archosaurs: a study in soft-tissue
1079 reconstruction in the fossil record with an analysis of the function of pneumaticity.
1080 *Journal of Vertebrate Paleontology* 17:1–76. doi:10.1080/02724634.1997.10011027.
- 1081 Xu, X. and Norell, M.A. 2004. A new troodontid dinosaur from China with avian-like sleeping
1082 posture. *Nature* 431:838–841. doi:10.1038/nature02898.
- 1083 Xu, X., Norell, M.A., Kuang, X., Wang, X., Zhao, Q. and Jia, C. 2004. Basal tyrannosauroids
1084 from China and evidence for protofeathers in tyrannosauroids. *Nature* 431:680–684.
1085 doi:10.1038/nature02855.
- 1086 Xu, X., Norell, M.A., Wang, X., Makovicky, P.J. and Wu, X. 2002. A basal troodontid from the
1087 Early Cretaceous of China. *Nature* 415:780–784. doi:10.1038/415780a.

- 1088 Xu, X. and Wu, X.-C. 2001. Cranial morphology of *Sinornithosaurus millenii* Xu et al. 1999
1089 (Dinosauria: Theropoda: Dromaeosauridae) from the Yixian Formation of Liaoning,
1090 China. *Canadian Journal of Earth Sciences* 38:1739–1752. doi:10.1139/e01-082.
- 1091 Zanno, L.E. 2010. Osteology of *Falcarius utahensis* (Dinosauria: Theropoda): characterizing the
1092 anatomy of basal therizinosaurs. *Zoological Journal of the Linnean Society* 158:196–230.
1093 doi:10.1111/j.1096-3642.2009.00464.x.
- 1094 Zanno, L.E. and Makovicky, P.J. 2011. Herbivorous ecomorphology and specialization patterns in
1095 theropod dinosaur evolution. *Proceedings of the National Academy of Sciences* 108:232–
1096 237. doi:10.1073/pnas.1011924108.
- 1097 Zusi, R.L. 1984. A functional and evolutionary analysis of rhynchokinesis in birds. *Smithsonian*
1098 *Contributions to Zoology* 395:1–40.
- 1099 Zusi, R.L. 1993. Patterns of diversity in the avian skull. *The skull* 2:391–437.
- 1100 Zweers, G.A. and Vanden Berge, J.C. 1998. Birds at geological boundaries. *Zoology* 100:183–
1101 202.
- 1102 Zweers, G.A., Vanden Berge, J.C. and Berkhoudt, H. 1997. Evolutionary patterns of avian trophic
1103 diversification. *Zoology* 100:25–57.

FIGURE CAPTIONS AND TABLES

FIGURE 1. Anatomy of nonavian theropod quadrates. Right (1–5) quadrate of *Tsaagan mangas* (IGM 100-1015; courtesy shared by Mick Ellison © AMNH) in (1) anterior, (2) lateral, (3) posterior, (4) medial and (5) ventral views; left (6–10) and right (11) quadrates of (6) *Baryonyx walkeri* (BMNH R9951), (7) *Aerosteon riocoloradensis* (MCNA-PV-3137; courtesy shared by Martín Ezcurra), (8) an indeterminate Oviraptoridae (GIN A; Maryańska and Osmólska 1997, figure 1), (9) *Tyrannosaurus rex* (BHI 3333, cast; Larson and Carpenter, 2008), (10) *Allosaurus ‘jimmadseni’* (SMA 005/02), and (11) *Majungasaurus crenatissimus* (FMNH PR 2100, cast; courtesy shared by Lawrence Witmer) in (6–9) posterior and (10–11) ventral views.

Abbreviations: **dqjc**, dorsal quadratojugal contact; **ecc**, ectocondyle; **enc**, entocondyle; **icn**, intercondylar notch; **ics**, intercondylar sulcus; **lpq**, lateral process of the quadrate; **mar**, mandibular articulation; **mfq**, medial fossa of the quadrate; **pfl**, pterygoid flange (in green); **pfq**, posterior fossa of the quadrate; **ppne**, posterior pneumatopore; **qb**, quadrate body (in blue); **qf**, quadrate foramen (delimited by a broader line); **qh**, quadrate head; **qj**, quadratojugal; **qjp**, quadratojugal process; **qr**, quadrate ridge; **qrg**, quadrate ridge groove; **qs**, quadrate shaft (in light blue); **sqc**, squamosal contact; **vqjc**, ventral quadratojugal contact; **vpdq**, ventral projection of the dorsal quadratojugal contact; **vsh**, ventral shelf.

FIGURE 2. Avian and non-avian theropod terminology of the quadrate bone. Left quadrate of *Fregata minor* (Great Frigatebird) in medial view annotated with (1) Baumel and Witmer (1993) terminology, and (2) the here proposed terminology for the nonavian theropod quadrate (image from Witmer 1990 and Baumel and Witmer (1993; modified).

FIGURE 3. Skull comparison between (1) the bird *Corythaeola cristata* (Great Blue Turaco; NH.34.26; courtesy of the Horniman Museum & Gardens), and (2) the alvarezsaurid *Shuvuuia deserti* (IGM 100-1001; reversed) in lateral view. Comparison between the left quadrate of the juvenile? specimen of *Shuvuuia deserti* (IGM 100-1001) in (3) posterior, and (4) lateral view, and the right quadrate of the adult specimen of *Shuvuuia deserti* (IGM) in (5) posterior, and (6) posterolateral views. Abbreviations: **j**, jugal; **jc**, jugal contact; **lpq**, lateral process; **otc**, otic capitulum; **pfl**, pterygoid flange; **po**, postorbital; **poc**, postorbital contact; **qf**, quadrate foramen; **sq**, squamosal; **sqc**, squamosal capitulum. Scale bars = 2 cm (1-2), 1 cm (5-6), and 5 mm (3-4).

FIGURE 4. Morphology and position of pneumatic openings in the quadrate of nonavian Theropoda. Pneumatic fossa (1) of the right quadrate of the metriacanthosaurid *Sinraptor dongi* (IVPP 10600) in posterior view (courtesy of Philip Currie). Medial pneumatopore (2) of the right quadrate of the carcharodontosaurid *Acrocanthosaurus atokensis* (NCSM 14345) in medial view. Medial pneumatopore (3) of the left quadrate of the carcharodontosaurid *Giganotosaurus carolinii* (MUCPv CH 1) in medial view. Posterior pneumatopore (4) of the left quadrate of the neovenatorid *Aerosteon riocoloradensis* (MCNA PV 3137) in posterior view (courtesy of Martín Ezcurra). Ventral pneumatopore (5) of the right quadrate of the tyrannosaurid *Alioramus altai* (IGM 100–844) in ventral view (courtesy of Mick Ellison). Ventral pneumatopore (6) of the left quadrate of the tyrannosaurid *Tyrannosaurus rex* (FMNH PR2081; cast) in ventral view. Medial pneumatopore (7) on the right quadrate of the therizinosauroid *Falcarius utahensis* (UMNH VP 14559) in medial view (courtesy of Lindsay Zanno). Posterior pneumatopore (8) of the left quadrate of the ornithomimid *Garudimimus brevipes* (IGM 100–13) in posterior view (courtesy of Yoshitsugu Kobayashi). Posterior pneumatopore (9) of the right quadrate of the dromaeosaurid *Buitreraptor gonzalezorum* (MPCA 245) in posterior view. Scale bars = 5 cm (1–3,6), 2 cm (4,8), 1 cm (5,7), 5 mm (9).

FIGURE 5. Distribution of quadrate pneumaticity in Theropoda. Simplified cladogram of theropod dinosaurs showing the phylogenetic distribution of a pneumatic quadrate denoted in red (from Bhullar et al., 2012; Modified). Pneumaticity is present in the quadrate of Allosauroidea (*Acrocanthosaurus*, *Mapusaurus*, *Giganotosaurus*), Tyrannosauroidea (*Alioramus*, *Albertosaurus*, *Tyrannosaurus*), Compsognathidae (*Sinosauropteryx*), Ornithomimosauria (*Ornithomimus*), Therizinosauria (*Falcarius*), Oviraptorosauria (*Conchoraptor*), Troodontidae (*Sinovenator*), Dromaeosauridae (*Buitreraptor*) and many birds. There is no evidence of a pneumatic quadrate in Alvarezsauroidea hitherto. When optimized onto this simplified tree, a pneumatic quadrate is a synapomorphy of Avetheropoda.

FIGURE 6. Quadrate of *Lourinhanosaurus antunesi* embryo. Incomplete left (1–8) quadrate (ML565-150) in (1) anterior, (2) lateral, (3) posterior, (4) medial, (5) ventral, (6) dorsal, (7) posteromedial, and (8) ventromedial views (the quadrate in (7) and (8) was photographed before preparation). Abbreviations: **dqjc**, dorsal quadratojugal contact; **ecc**, ectocondyle; **enc**, entocondyle; **mfq**, medial fossa; **pfl**, pterygoid flange; **pfq**, posterior fossa; **qjp**, quadratojugal process; **qr**, quadrate ridge; **vqjc**, ventral quadratojugal contact.

TABLE 1. Standardized terminology and abbreviation of the nonavian theropod quadrate and comparison with the terminology of the avian quadrate based on Baumel and Witmer (1993), Elzanowski et al. (2001) and Elzanowski and Stidham (2010).

Nonavian theropod quadrate (q)		Avian theropod quadrate
Quadrate body	qb	Corpus quadrati
Quadrate shaft	qs	/
Quadrate ridge	qr	/
Quadrate ridge groove	qrg	/
Quadrate head	qh	Processus oticus/Pars otica/Caput quadrati
Otic capitulum	otc	Capitulum oticum
Squamosal capitulum	sqc	Capitulum squamosum
Intercapitular sulcus	icas	Incisura/Vallecula intercapitularis
Quadrate foramen	qf	/
Mandibular articulation	mar	Pars/Processus mandibularis
Ectocondyle	ecc	Condylus (mandibularis) lateralis
Entocondyle	enc	Condylus (mandibularis) medialis
Mediocondyle	mec	Condylus caudalis
Intercondylar sulcus	ics	Vallecula intercondylaris
Intercondylar notch	icn	Depressio praecondylaris
Lateral process	lpq	/
Quadratojugal contact	qjc	Cotyla quadratojugal
Ventral quadratojugal contact	vqjc	/
Dorsal quadratojugal contact	dqjc	/
Quadratojugal process	qjp	/
Ventral projection of the dorsal	vpdq	/
quadratojugal contact		
Squamosal contact	sqc	/
Posterior fossa	pfq	/
Quadrate diverticulum	qdi	/
Dorsal pneumatopore	dpne	Foramen pneumaticum caudomediale
Medial pneumatopore	mpne	Foramen pneumaticum basiorbitale
Posterior pneumatopore	ppne	Foramen pneumaticum rostromediale
Ventral pneumatopore	vpne	Foramen pneumaticum adventitium
Pterygoid flange	pfl	Processus orbitalis
Pterygoid contact	ptc	Condylus pterygoideus/Facies articularis pterygoidea
Medial fossa	mfq	Fossa basiorbitalis
Ventral shelf	vsh	/

Figure 1

Anatomy of nonavian theropod quadrates

Right (1–5) quadrate of *Tsaagan mangas* (IGM 100-1015; courtesy shared by Mick Ellison © AMNH) in (1) anterior, (2) lateral, (3) posterior, (4) medial and (5) ventral views; left (6–10) and right (11) quadrates of (6) *Baryonyx walkeri* (BMNH R9951), (7) *Aerosteon riocoloradensis* (MCNA-PV-3137; courtesy shared by Martín Ezcurra), (8) an indeterminate Oviraptoridae (GIN A; Maryńska and Osmólska 1997 :fig. 1), (9) *Tyrannosaurus rex* (BHI 3333, cast; Larson and Carpenter, 2008), (10) *Allosaurus 'jimmadseni'* (SMA 005/02), and (11) *Majungasaurus crenatissimus* (FMNH PR 2100, cast; courtesy shared by Lawrence Witmer) in (6–9) posterior and (10–11) ventral views. Abbreviations: **dqjc**, dorsal quadratojugal contact; **ecc**, ectocondyle; **enc**, entocondyle; **icn**, intercondylar notch; **ics**, intercondylar sulcus; **lpq**, lateral process of the quadrate; **mar**, mandibular articulation; **mfq**, medial fossa of the quadrate; **pfl**, pterygoid flange (in green); **pfq**, posterior fossa of the quadrate; **ppne**, posterior pneumatopore; **qb**, quadrate body (in blue); **qf**, quadrate foramen (delimited by a broader line); **qh**, quadrate head; **qj**, quadratojugal; **qjp**, quadratojugal process; **qr**, quadrate ridge; **qrg**, quadrate ridge groove; **qs**, quadrate shaft (in light blue); **sqc**, squamosal contact; **vqjc**, ventral quadratojugal contact; **vpdq**, ventral projection of the dorsal quadratojugal contact; **vsh**, ventral shelf.





Figure 2

Avian and non-avian theropod terminology of the quadrate bone

Left quadrate of *Fregata minor* (Great Frigatebird) in medial view annotated with (1) Baumel and Witmer (1993) terminology, and (2) the here proposed terminology for the nonavian theropod quadrate (image from Witmer 1990 and Baumel and Witmer (1993) ; modified).




Figure 3

Skull comparison between *Corythaeola* and *Shuvuuia*

Skull comparison between (1) the bird *Corythaeola cristata* (Great Blue Turaco; NH.34.26; courtesy of the Horniman Museum & Gardens), and (2) the alvarezsaurid *Shuvuuia deserti* (IGM 100-1001; reversed) in lateral view. Comparison between the left quadrate of the juvenile? specimen of *Shuvuuia deserti* (IGM 100-1001) in (3) posterior, and (4) lateral view, and the right quadrate of the adult specimen of *Shuvuuia deserti* (IGM) in (5) posterior, and (6) posterolateral views. Abbreviations: **j**, jugal; **jc**, jugal contact; **lpq**, lateral process; **otc**, otic capitulum; **pfl**, pterygoid flange; **po**, postorbital; **poc**, postorbital contact; **qf**, quadrate foramen; **sq**, squamosal; **sqc**, squamosal capitulum. Scale bars = 2 cm (1-2), 1 cm (5-6), and 5 mm (3-4).






Figure 4

Morphology and position of pneumatic openings in the quadrate of nonavian Theropoda

Pneumatic fossa (1) of the right quadrate of the metriacanthosaurid *Sinraptor dongi* (IVPP 10600) in posterior view (courtesy of Philip Currie). Medial pneumatopore (2) of the right quadrate of the carcharodontosaurid *Acrocanthosaurus atokensis* (NCSM 14345) in medial view. Medial pneumatopore (3) of the left quadrate of the carcharodontosaurid *Giganotosaurus carolinii* (MUCPv CH 1) in medial view. Posterior pneumatopore (4) of the left quadrate of the neovenatorid *Aerosteon riocoloradensis* (MCNA PV 3137) in posterior view (courtesy of Martín Ezcurra). Ventral pneumatopore (5) of the right quadrate of the tyrannosaurid *Alioramus altai* (IGM 100–844) in ventral view (courtesy of Mick Ellison). Ventral pneumatopore (6) of the left quadrate of the tyrannosaurid *Tyrannosaurus rex* (FMNH PR2081; cast) in ventral view. Medial pneumatopore (7) on the right quadrate of the therizinosauroid *Falcarius utahensis* (UMNH VP 14559) in medial view (courtesy of Lindsay Zanno). Posterior pneumatopore (8) of the left quadrate of the ornithomimid *Garudimimus brevipes* (IGM 100–13) in posterior view (courtesy of Yoshitsugu Kobayashi). Posterior pneumatopore (9) of the right quadrate of the dromaeosaurid *Buitreraptor gonzalezorum* (MPCA 245) in posterior view. Scale bars = 5 cm (1–3,6), 2 cm (4,8), 1 cm (5,7), 5 mm (9).





Figure 5

Distribution of quadrate pneumaticity in Theropoda

Simplified cladogram of theropod dinosaurs showing the phylogenetic distribution of a pneumatic quadrate denoted in red (from Bhullar et al., 2012; Modified). Pneumaticity is present in the quadrate of Allosauroidea (*Acrocanthosaurus*, *Mapusaurus*, *Giganotosaurus*), Tyrannosauroidea (*Alioramus*, *Albertosaurus*, *Tyrannosaurus*), Compsognathidae (*Sinosauroptryx*), Ornithomimosauria (*Ornithomimus*), Therizinosauria (*Falcarius*), Oviraptorosauria (*Conchoraptor*), Troodontidae (*Sinovenator*), Dromaeosauridae (*Buitreraptor*) and many birds. There is no evidence of a pneumatic quadrate in Alvarezsauroidea hitherto. When optimized onto this simplified tree, a pneumatic quadrate is a synapomorphy of Avetheropoda.




Figure 6

Quadrate of *Lourinhanosaurus antunesi* embryo

Incomplete left (1–8) quadrate (ML565-150) in (1) anterior, (2) lateral, (3) posterior, (4) medial, (5) ventral, (6) dorsal, (7) posteromedial, and (8) ventromedial views (the quadrate in (7) and (8) was photographed before preparation). Abbreviations: **dqjc**, dorsal quadratojugal contact; **ecc**, ectocondyle; **enc**, entocondyle; **mfq**, medial fossa; **pfl**, pterygoid flange; **pfq**, posterior fossa; **qjp**, quadratojugal process; **qr**, quadrate ridge; **vqjc**, ventral quadratojugal contact.

