

Estimating the differences in critical thermal maximum and metabolic rate of *Helicoverpa punctigera* (Wallengren) (Lepidoptera: Noctuidae) across life stages

Samuel Bawa^{Corresp., 1, 2}, Peter Gregg³, Alice Del Soccoro³, Cara Miller⁴, Nigel Andrew²

¹ Asuansi Agric. Station, Box 520, Cape Coast, Ghana, Cape Coast, Central Region, Ghana

² Zoology, Insect Ecology Laboratory, University of New England, Armidale, NSW, Australia

³ Agronomy and Soil Science, University of New England, Armidale, NSW, Australia

⁴ Science and Technology, University of New England, Armidale, NSW, Australia

Corresponding Author: Samuel Bawa
Email address: ambiasin@yahoo.com

Temperature is a crucial driver of insect activity and physiological processes throughout their life-history, and heat stress may impact life stages (larvae, pupae and adult) in different ways. Using thermolimit respirometry, we assessed the critical thermal maxima (CT_{max} - temperature at which an organism loses neuromuscular control), CO_2 emission rate ($\dot{V}CO_2$) and Q_{10} (a measure of $\dot{V}CO_2$ temperature sensitivity) of three different life stages of *Helicoverpa punctigera* (Wallengren) by increasing their temperature exposure from 25°C to 55°C at a rate of 0.25°C min⁻¹. We found that the CT_{max} of larvae (49.1°C±0.3°C) was higher than pupae (47.4°C ±0.2°C) and adults (46.9°C±0.2°C). The mean mass-specific CO_2 emission rate (ml $\dot{V}CO_2$ h⁻¹) of larvae (0.26±0.03 ml $\dot{V}CO_2$ h⁻¹) was also higher than adults (0.24±0.04 ml $\dot{V}CO_2$ h⁻¹) and pupae (0.06±0.02 ml $\dot{V}CO_2$ h⁻¹). The $Q_{10:25-35^\circ C}$ for adults (2.01±0.22) was significantly higher compared to larvae (1.40±0.06) and $Q_{10:35-45^\circ C}$ for adults (3.42±0.24) was significantly higher compared to larvae (1.95±0.08) and pupae (1.42±0.98) respectively. We have established the upper thermal tolerance of *H. punctigera*, which will lead to a better understanding of the thermal physiology of this species both in its native range, and as a pest species in agricultural systems.

Estimating the differences in critical thermal maximum and metabolic rate of *Helicoverpa punctigera* (Wallengren) (Lepidoptera: Noctuidae) across life stages

Samuel A. Bawa^{1,4*}, Peter C. Gregg², Alice P. Del Socorro², Cara Miller³, Nigel R. Andrew¹

¹ Insect Ecology Lab, Natural History Museum, School of Environmental and Rural Science, University of New England, Armidale, NSW 2351, Australia.

² Agronomy, School of Environmental and Rural Science, University of New England, Armidale, NSW 2351, Australia.

³ School of Science and Technology, University of New England, Armidale, NSW 2351, Australia

⁴ Asuansi Agric. Station, Box 520, Cape Coast, Ghana

Corresponding author: Samuel A. Bawa

Email address: abawa@myune.edu.au/ambiasin@yahoo.com

Postal address: Insect Ecology Lab, Natural History Museum, Building W77, University of New England, Armidale, NSW 2351, Australia.

Abstract

Temperature is a crucial driver of insect activity and physiological processes throughout their life-history, and heat stress may impact life stages (larvae, pupae and adult) in different ways. Using thermolimit respirometry, we assessed the critical thermal maxima (CT_{max} - temperature at which an organism loses neuromuscular control), CO_2 emission rate ($\dot{V}CO_2$) and Q_{10} (a measure of $\dot{V}CO_2$ temperature sensitivity) of three different life stages of *Helicoverpa punctigera* (Wallengren) by increasing their temperature exposure from 25°C to 55°C at a rate of 0.25°C min⁻¹. We found that the CT_{max} of larvae (49.1°C±0.3°C) was higher than pupae (47.4°C ±0.2°C) and adults (46.9°C±0.2°C). The mean mass-specific CO_2 emission rate (ml $\dot{V}CO_2$ h⁻¹) of larvae (0.26±0.03 ml $\dot{V}CO_2$ h⁻¹) was also higher than adults (0.24±0.04 ml $\dot{V}CO_2$ h⁻¹) and pupae (0.06±0.02 ml $\dot{V}CO_2$ h⁻¹). The $Q_{10:25-35^\circ C}$ for adults (2.01±0.22) was significantly higher compared to larvae (1.40±0.06) and $Q_{10:35-45^\circ C}$ for adults (3.42±0.24) was significantly higher compared to larvae (1.95±0.08) and pupae (1.42±0.98) respectively. We have established the upper thermal tolerance of *H. punctigera*, which will lead to a better understanding of the thermal physiology of this species both in its native range, and as a pest species in agricultural systems.

Key words: *Helicoverpa punctigera*, life stage, critical thermal maxima, metabolic rate, thermolimit respirometry

Introduction

Insects have a specific range of temperatures within which they can survive, grow and reproduce (Andrew et al., 2011; Huang et al., 2015; DeVries et al., 2016), and have their distribution strongly defined by a thermal envelope (Gullan & Cranston, 2014; Li et al., 2019). Assessing the impact of extreme temperature exposures on insect distribution is vital to understanding insect ecology and the current and future effects of climate change (Andrew et al., 2011; Betini et al., 2013). Within a rapidly changing climate, thermal limits are an essential part of assessing thermal physiology of insects, as they are the critical endpoints that an organism can survive or loses muscular control (Lighton & Turner, 2004; Andrew et al., 2016), quantified in terms of critical thermal maximum (CT_{max}) and critical thermal minimum (CT_{min}) (Kingsolver et al., 2018).

Thermal limit measurements like CT_{max} and CT_{min} are commonly used to understand and predict species distributions among populations (Shik et al., 2019) and responses to environmental changes (Angilletta et al., 2003; Angilletta, 2009; Kingsolver et al., 2018). CT_{max} is often aligned to the distribution of species, as it is a biologically significant and ecologically relevant thermal performance trait (Sunday et al., 2014; Shik et al., 2019). However, the CT_{max} of most insect groups is highly plastic, so can change depending on an organism's pre-exposure conditions, making it difficult to forecast their response to a rapidly changing climate (Terblanche et al., 2005; Martínez et al., 2016; Kellermann et al., 2017).

CT_{max} has been shown to differ across life stages in some insects e.g. kelp fly (*Paractora dreuxi*), Sirex wasp (*Sirex noctilio*) and mealworm (*Tenebrio molitor*) (Marais et al., 2009; Vorhees &

Bradley, 2012; Li et al., 2019), and can be influenced by body mass and surface area of the insect (Lighton & Turner, 2004). CT_{max} values depend substantially on how they are measured (Lighton & Turner, 2004; Shik et al., 2019); it is thus essential to determine a precise scale of temperature exposure for insects (Terblanche et al., 2007). Temperature ramping rates can influence the CT_{max} of an insect (Lighton & Turner, 2004; Terblanche et al., 2007). A slow temperature ramp rate of an organism leads to increased risk of heat shock, which is absent under the natural conditions (Lutterschmidt and Hutchison, 1997). The other process that can occur during a slow ramp is short-term physiological adjustments (e.g. induction of heat shock proteins) that protect cells from heat damage and might increase CT_{max} (Kingsolver, 2009; Shik et al., 2019). Fast temperature ramp leads to delay in thermal equilibrium between the air and the insect's body. Because of this, under a fast temperature ramp, the inner temperature of the insect will be lower than the recorded air temperatures leading to an overestimation of their CT_{max} (Lighton & Turner, 2004; Andrew et al., 2013; Agudelo-Cantero et al., 2019).

Intriguingly, only few studies have looked at thermal variation across different life stages (Le Lann et al., 2011). The few studies focused on a single unit of fitness in one environment (Le Lann et al., 2011). Thermal plasticity has been identified in many insect taxa e.g. ants *Iridomyrmex purpureus*, *Drosophila* species and parasitic wasps (*Aphidius rhopalosiphi*) (Chown & Terblanche, 2006; Le Lann et al., 2011; Andrew et al., 2013; Shik et al., 2019), but not all (Terblanche et al., 2010).

Insect herbivores that are found on the ground and plant surfaces in warmer and drier environments are exposed to substantial microclimate variability including seasonal and daily temperature variations at the soil and plant surface (Hodgson, 1991; Andrew et al., 2016). How organisms respond to the variable microclimates depends on the thermal physiology e.g. CT_{max} (Somero,

2010; Shik et al., 2019). The physiology of insects can be modified by the microclimate they experience (Andrew et al., 2013; Andrew et al., 2016). Predicting the future in changing climates is made difficult by our limited understanding of the eco-physiological mechanisms and evolutionary process underlying thermal phenotypic plasticity (Shik et al., 2019).

Helicoverpa punctigera is native to Australia and widely distributed across the continent (Gregg et al., 2018). It is a notable pest on crops like cotton, oilseeds, legumes and vegetables (Gregg et al., 2019). The adult can travel a long distance and can adapt locally to different environmental conditions including inland Australia (Gregg et al., 2018, 2019). Female adult *H. punctigera* lay eggs on leaves, flower buds and developing fruits of the host plant (Zalucki et al., 1986). The neonate (newly hatched) usually eat soft leaves; as they mature, they move about other feeding sites like flowers or flower buds, pods, fruits and seeds (Gregg et al., 2018). The last larval instar burrows into the soil and pupates. Inland habitats are characterised by extreme climatic conditions like very hot temperatures as high as 50°C during summer (Bureau of Meteorology, 2018). The ability of this insect to survive the extreme summer temperatures in the inland depends on their thermal tolerances and physiological sensitivities e.g., CT_{max} and biochemical shifts that are temperature-dependent (Gaitán-Espitia et al., 2014). Female adults of *H. punctigera* are responsible for host selection and dispersal, whilst feeding activities of the larvae cause economic loss to the host plants, and a complete reorganisation of the body occurs within the pupal stage, understanding the metabolic rates among the different life stages are essential.

Energy is the crucial currency of life-history traits of animals including insects (Terblanche & Chown, 2007; Desforges et al., 2019). Metabolism combines organism's energy supply and fitness-related activities (Shah et al., 2021). Measurement of metabolic rate is one way of measuring the energy cost related with different life-functions like growth, development,

movement among others (Shah et al., 2021). In insects, metabolic rates can influence not only fitness in individuals but also geographic distributions and abundances (Shah et al., 2021). Therefore, estimating the metabolic rates of organisms will assist in identifying the cost of living and performance in a particular environment (Terblanche et al., 2005; Terblanche & Chown, 2007), and can help predict geographic variations in response to global warming (Shah et al., 2021).

Metabolic rate can respond to changes in ambient temperature, activity and body mass; however, age and ontogeny, sex, feeding status, season and time of day, among others, can also exert substantial influence (Terblanche & Chown, 2007; Harrison, 2009; Altermatt, 2010). For example, in the Australian field cricket (*Teleogryllus oceanicus*) and thynnine wasp (*Zaspilothynnus nigripes*), males have a higher metabolic rate than females (Kolluru et al., 2004; Tomlinson & Phillips, 2015). Specific dynamic action is the amount of energy expenditure above the basal metabolic rate (RMR) due to the ingestion and digestion of food for use as energy or conversion to a storage form (Secor, 2009). One major metabolic variable is the temporary rise in metabolic rate following feeding status or swallowing food (Karasov & Martinez del Rio, 2007; McCue et al., 2012). This feeding status-associated rise in metabolic rate is a result of food breakdown and processing in the gut of the organism (Karasov & Martinez del Rio, 2007). High temperatures generally produce high metabolic rate, however the total energy assigned for specific dynamic actions is not fully dependent on temperature (McCue et al., 2016a; Wang et al., 2003).

The temperature sensitivity (Q_{10}), defined as the rate ratio of a given process taking place at different temperatures, is used to measure the degree of temperature dependence of a given biological process (Davidson et al., 2006; Mundim et al., 2020). Q_{10} is an important parameter in predicting the effects of temperature on carbon dioxide release (Mundim et al., 2020).

Previous studies on *Helicoverpa* species (eg. *H. armigera* and *H. punctigera*) focused on the effect of temperature on development rate, survivorship and reproduction (Room, 1983; Qayyum & Zalucki, 1987; Mironidis et al., 2008; Zalucki et al., 2005; Mironidis, 2014) but not on thermal physiology (CT_{max} and metabolic rate) of different life stages. For example, *H. punctigera* are distinctive in terms of their life stage variation and habitat (Gregg et al., 2018, 2019), just as other insects with stage-based niche changes. Across life stages, an individual will be exposed to a range of temperatures (Agudelo-Cantero et al., 2019) with different opportunities to escape extreme exposure. In the field, the larvae and adults of *H. punctigera* live on plant surfaces while the pupae lives in the soil. Based on this life stage and habitat variations, their tolerance to extreme temperatures may significantly vary among the different life stages. Our study is the first of *H. punctigera* thermal physiology across three life-stages – larvae, pupae and adult.

We assessed the thermal physiology (CT_{max} and metabolic rate) of three different life stages — late instar larvae, pupae, and adult — and sex of *H. punctigera* during temperature ramping using thermolimit respirometry, which employs a flow-through respirometry, with $\dot{V}CO_2$ as a key measure. The technique has an infrared activity detector to monitor moving activity (Lighton & Turner, 2004; Vorhees & Bradley, 2012). One important advantage of this technique is that CT_{max} can be identified directly from respiratory or activity data. In addition, it allows for CT_{max} to be determined without disturbing the organism (Vorhees & Bradley, 2012). The technique, in addition, allows for the measurement of $\dot{V}CO_2$ when exposed to biologically relevant temperature points.

The specific questions we asked are:

(i) what are the differences in $\dot{V}CO_2$ and CT_{max} across different life stages of *H. punctigera* considering the different field conditions each stage experiences? We predict that, larvae and adult of *H. punctigera*, which are exposed to the many microclimatic conditions on host plants surface, would have a higher metabolic rate and CT_{max} than pupae, because, the soil the pupae is burrowed in is more thermally stable than the leaves.

(ii) what are the metabolic rates of the life stages at five different temperatures at 5°C bins: 25°C, 30°C, 35°C, 40°C and 45°C? Here, we predict that because exposure temperature has an overwhelming effect on the metabolic rate of insects, the metabolic rate among the life stages would be lower at 25°C, 30°C and 35°C compared with 40°C and 45°C.

(iii) does sex influence the metabolic rate and CT_{max} of adult and pupae of *H. punctigera*? We predict that because of the differences in activity (e.g., host selection and dispersal in the case of adults) between male and female, sex would influence metabolic rate and CT_{max} of *H. punctigera* pupae and adult.

Materials and methods

Insect Culture and Experimental Animals

A laboratory insect culture was established in April 2018 from pupae provided by Tamworth Agricultural Institute, New South Wales Department of Primary Industries. After emergence, the adults were put into cylindrical framed mating cages containing 5% sugar solution in dental wicks held in 50 ml plastic containers. We left female moths to mate and lay fertile eggs for 72 hours following protocols outlined by Gregg et al. (2016). Eggs were collected by daily scrolling (by way of folding) of the paper towel which acts as the mating cage wall, followed by cutting the

towel containing the eggs into pieces measuring approximately 7cm x 21cm x 11cm. Eggs were cleaned by brief immersion in 0.2% sodium hypochlorite solution for disease prevention, brushed carefully for one to two minutes and then rinsed two times with distilled water onto filter paper placed in a Büchner funnel. The cleaned eggs were put in plastic eggcups coated with artificial diet and kept at 25°C. Five days after hatching, the larvae were transferred into 35ml plastic cups ventilated with small holes and containing 10-15 ml of soybean-based artificial diet (Greene et al., 1976; Teakle, 1991). The insects were reared in the Agronomy insectary maintained at controlled conditions of 70-75% relative humidity, 14h: 10h light: dark period and 25°C, University of New England, NSW, Australia. For a continuous supply of *H. punctigera* individuals at appropriate life-stages, two cultures were maintained at different developmental stages. Before and after each experiment, the larvae, pupae or adult moths used were weighed on an electronic balance (Mettler Toledo XP 404S, Greifensee, Switzerland) with an adjusted accuracy of 0.1 mg. We used fifth instar larvae in our experiment because they were available and relatively easy to manipulate without damaging their cuticle. The average live body mass of the insects used pre-experiment was 189.09 ± 15.58 mg for adults, 348.26 ± 57.57 mg for larvae and 332.38 ± 18.39 mg for pupae. One-week-old pupae were sexed into male and female under a microscope using the genital scars (Kirkpatrick, 1961) and used in pupal assays. Finally, virgin adults that were one to two days post-emergence were used for the experiment: these adults were not fed. Because of the polyphagous nature of the insect, larvae would always have a bite of a meal or would always have eaten in the field, so all measures of metabolic rate include the specific dynamic action (roughly, the cost of digestion) of the larvae. Larvae were fed with the same amount of food and quality ad-libitum, until it was time to use for the experiment.

Experimental setup

We used standard thermolimit respirometry protocols (Lighton & Turner, 2004; Terblanche et al., 2007; Boardman & Terblanche, 2015; Andrew et al., 2016; Ghaedi & Andrew, 2016). In brief, using a HiBlow HP40 air pump, room air was pushed into sodalime to remove CO₂ and Drierite to remove moisture. Air was set to flow at 180 ml min⁻¹ using a mass flow controller (Sable MFC-2, Sable Systems) and flow control valve (Side-Trak 840L; Sierra Instruments Inc., Monterey, CA, USA) through Bev-a-line tubing to a 30ml glass cuvette. Carbon dioxide and water in the air that was flowing before and after the glass cuvette was analysed by a two-channel calibrated infrared Li-7000 analyser (Li-Cor, Lincoln, NE, USA). The scrubbed airstream flowed across the experimental organism in the 30ml glass cuvette, which was held underwater in a water bath (Grant, GP200-R4), programmed (resolution $\pm 0.1^{\circ}\text{C}$) with LABWISE software. Cuvettes were placed into two plastic bags to stop any water entering the cuvette. We warmed insects from 25°C to 55°C, at 0.25°C min⁻¹ (Andrew et al., 2016), over 120 minutes in the 30ml glass cuvette. The outgoing air from the cuvette then enters the infrared Li-cor analyser again: the CO₂ concentration difference of the air before and after it flowed through the cuvette, at one-second intervals is then recorded. A type-T thermocouple attached to a data logger (PicoTech TC-08 data logger with resolution $\pm 0.025^{\circ}\text{C}$, accuracy $\pm 0.5^{\circ}\text{C}$) was used to record the temperature in the glass cuvette. The two data recordings (Licor and data logger) were combined using Microsoft excel, aligned using the computer time stamp. An infrared activity detector (AD-1; Sable Systems, Las Vegas, NV, USA) recorded the movement of the test insect (see trace in supplementary).

Data Extraction and measurement of CO₂ emission rate (metabolic rate) and CT_{max}

209 Using Expedata (Sable Systems International, Las Vegas, NV, USA) version 1.9.2 analysis
 210 software, data on CO₂ emission rate was extracted and measured from 25°C to 55°C (Boardman
 211 et al., 2015; Andrew et al., 2016). We drift-corrected our recordings to baseline recordings from
 212 an empty 30ml glass cuvette made ten min before and after trial, and converted to ml CO₂ h⁻¹
 213 (Andrew et al., 2016; Ghaedi & Andrew, 2016). We controlled chamber switching by recording
 214 from the cuvette by manually adding/removing the insect from the same cuvette as the baseline.
 215 For each replicate insect, $\dot{V}CO_2$ (n = 10 larvae, 10 pupae, 10 adults) were extracted at these
 216 temperature bins (25, 30, 35°C, 40°C, 45°C and CT_{max} ±0.25°C). The overall $\dot{V}CO_2$ (in ml CO₂
 217 h⁻¹) was calculated by extracting the area under the curve (integral of CO₂ ml h⁻¹ vs hours) in total
 218 over the ramping period from 25°C to CT_{max} by transforming ppm concentration of CO₂ to CO₂
 219 fraction and then multiplied by the flow rate. This area was equal to the volume of CO₂ produced
 220 by each replicate insect in the 30ml glass cuvette (Stevens et al., 2010; Andrew et al., 2016). The
 221 $\dot{V}CO_2$ for each replicate insect at each temperature point was extracted and multiplied by 1,000 to
 222 give $\mu\text{l CO}_2 \text{ h}^{-1}$ per insect per hour. CT_{max} (n = 10 larvae, 10 pupae, 10 adults) was determined by
 223 identifying the point at which the spiracular activity ceases i.e. the inflection point in the absolute
 224 difference sum (ADS) residuals (Lighton & Turner, 2004; Vorhees & Bradley, 2012; Andrew et
 225 al., 2016; Ghaedi & Andrew 2016). Because of differences in activity among the life stages, to
 226 standardise our analysis, we used $\dot{V}CO_2$ data to determine CT_{max} value not activity. Other studies
 227 showed that $\dot{V}CO_2$ and activities measurements produced similar estimates of CT_{max} in insects
 228 (Lighton & Turner, 2004; Vorhees & Bradley, 2012). Metabolic rates at 25°C, 35°C and 45°C for
 229 Q₁₀, were calculated as the average $\dot{V}CO_2$ over the five-minute period at each temperature e.g. start
 230 at 25°C and go to 26.25°C. The inflection point for activity ADS residuals where motor activity

ceases is the activity CT_{max} . Lighton & Turner (2004) defined ADS as the cumulative sum of the absolute difference between adjacent sampling points.

Data analysis

We considered our measures of $\dot{V}CO_2$ as a proxy for metabolic rate (Terblanche et al., 2010). Statistical analyses were carried out in SAS[®] 9.4 software (SAS Institute Inc., 2013) using the GLIMMIX procedure. CT_{max} and total metabolic rate differences between the life stages were determined using generalised linear models (GLM). Differences in metabolic rate for different temperatures were assessed using generalised linear mixed models (GLMM) with individuals as random factors to balance for the repeated measurement. A one-way analysis of variance (ANOVA) was used to compare the overall $\dot{V}CO_2$ and CT_{max} among the life stages. The Q_{10} (25–35 °C) and Q_{10} (35–45 °C) for the life stages were also recorded similar to the above described procedure. Data were reported as means plus or minus standard error (mean \pm SE). In all the models CT_{max} and $\dot{V}CO_2$ of individuals were the response variables with life stage as the explanatory variables. Data was checked for normality and homogeneity of variance. The factors in the model include life stage, sex and body mass as a covariate for metabolic rate. Because sex was not significant for CT_{max} and metabolic rate, data on sex for those parameters were combined in the subsequent analysis. Mass was removed from the final model for CT_{max} since it was not significant. Where there were significant differences, post-hoc analysis was performed using least-square means (LSmeans) and the differences compared, $\alpha=0.05$ (Tukey-Kramer Adjustment). Mass was log-transformed before analysis. T-test was used to compare treatments under two treatments.

Results

Metabolic rate (overall $\dot{V}CO_2$ (ml h⁻¹) across life stage of *H. punctigera*

There was a significant effect of life stages on overall $\dot{V}CO_2$ (ANOVA, DF=2, F=31.52, $p<0.0001$). Pupae (n=10) had an overall $\dot{V}CO_2$ of 41% lower than adults (n=10) and 50% lower than larvae (n=10) ($p<0.0001$, and $p<0.0001$, respectively; Figure 1). The $\dot{V}CO_2$ of larvae was 9% higher than adults, but not statistically significant ($p=0.20$; Figure 1).

Upper thermal tolerance (CT_{max}) across life stages of *H.punctigera*

There was a significant life stage effect on CT_{max} (ANOVA, DF=2, F=43, $p<0.0001$). Adult (n=10) and pupae (n=10) had a similar CT_{max} ($p=0.12$; Figure 2). However, larvae (n=10) had a CT_{max} on average 2.2°C higher than adults and 1.8°C higher than pupae ($p<0.0001$; $p<0.0001$, respectively; Figure 2). Activity CT_{max} and CO_2 CT_{max} for adults were similar and not significant; unlike larvae, which had significantly higher CO_2 CT_{max} (Table 1).

$\dot{V}CO_2$ of *H. punctigera* life stages at different temperature points

Temperature effect on mean $\dot{V}CO_2$ was high in adults and larvae compared to pupae (Table 2) at the different temperature points. The mean $\dot{V}CO_2$ of pupae was significantly different from that of adult and larvae at all temperature points except at 25°C (Table 2). The mean $\dot{V}CO_2$ of each of the life stages doubled at both 40°C and 45°C (Table 2) before CT_{max} . Although the mean $\dot{V}CO_2$ between adults and larvae was not significantly different at 40°C, adults had a 0.01 μl h⁻¹ higher mean $\dot{V}CO_2$ than larvae (Table 2). At equilibrium (25°C) and 30°C-35°C larvae had 0.19 μl h⁻¹, 0.13 μl h⁻¹ and 0.16 μl h⁻¹ higher than adult (Table 2) respectively. The Q_{10} 25–35 °C values for larvae and adults were lower compared to Q_{10} 35–45°C – 28% lower for larvae (Q_{10} :25–35 °C

1.40±0.06 to Q_{10} 35–45 °C 1.95±0.14), 41% lower for adults (2.01±0.22 to 3.42±0.24) but 32% higher for pupae (1.74± 0.21 to 1.42±0.98) (Figure 3). Among life stages the Q_{10} : 25–35 °C for adults was significantly higher ($p= 0.04$) compared with larvae, but not pupae ($p=0.53$; Figure 3). The Q_{10} : 25–35 °C was not significant between larvae and pupae ($p=0.35$; Figure 3). The Q_{10} :35–45 °C values were not statistically different among larvae and pupae ($p=0.07$; Figure 3) but significant among larvae and adults as well as pupae and adults ($p=0.00$, $p=0.00$; Figure 3) respectively.

The influence of sex on upper thermal limit (CT_{max}) and metabolic rates $\dot{V}CO_2$ (ml h⁻¹) of adult moth and pupae

There was no sex effect (ANOVA, $DF=3$, $F=0.15$, $p=0.92$; Figure 4) on the CT_{max} of adults (males: 46.92°C ± 0.29°C and females: 46.72°C ±0.29°C) and pupae (males: 47.47°C ± 0.29°C and females: 47.32°C ± 0.29°C). For metabolic rate assessments, there was no significant difference between female and male adults as well as between female and male pupae ($t= -0.57$, $p=0.93$; Figure 5) and ($t=1.38$, $p=0.52$; Figure 5) respectively.

Discussion

Here, we measured the upper thermal tolerance limits (CT_{max}) and metabolic rates at different temperature exposures across three life stages larvae, pupae, and adults and between sexes of *H. punctigera* using thermolimit respirometry. These life stages experience different temperatures because of the different habitats they naturally occupy, enabling an ecologically relevant investigation of their metabolic rates at different temperatures. We found that CT_{max} differed among the life stages of the insect with larvae exhibiting significantly higher CT_{max} compared with

pupae and adults (Figure 2). Similar to the findings presented here, studies by other authors have shown that CT_{max} differs among life stages of insects including *Paractora dreuxi*, *Sirex noctilio* and *Tenebrio molitor* (Marais et al., 2009; Vorhees & Bradley, 2012; Li et al., 2019). The higher CT_{max} of the larvae compared with adults however, contradicts the findings of Vorhees & Bradley, (2012) and Li et al., (2019) in *T. molitor* and *S. noctilio* respectively, where adults had a higher CT_{max} than larvae, although, the insects are of different taxa. The differences in the CT_{max} between the life stages suggest differences in upper thermal tolerance in this species with larvae having the highest thermal tolerance to high temperature (Marais et al., 2009).

The CT_{max} differences among the life stages could be caused by several different factors such as interactions of the microclimatic conditions experienced by the life stages and the level to which they compensate their physiology to avoid harsh or extreme temperatures (Vorhees & Bradley, 2012). However, this would not be the case here, as all insects were kept at a constant 25°C, so a more evolutionarily relevant mechanism must be in train here. The results imply that, larvae, pupae and adults have the ability in the wild to adapt and cope differently.

The significantly higher CT_{max} exhibited by larvae is not surprising because larvae in their natural environment are generally more exposed to extreme/hot temperatures, just above the soil surface and while feeding on plants, and, less able to behaviourally adapt to extreme periods. Hence the need to modify their physiological response to such extreme periods of heat conditions for survival (Andrew et al., 2013a; Andrew et al., 2013b; Andrew et al., 2016). Even though the insects used in the study were laboratory-reared, they were most likely to exhibit similar characteristics like those in the field, because there is little genetic variability among *H. punctigera* population (Daly et al., 1985). The limitation of our study is that; the low genetic diversity in the wild in the insect does not imply that laboratory populations will not be adapted after many generations. We

acknowledge that the climatic variability experienced in the wild could further influence in these thermal tolerance traits, but is unlikely to affect the clear life-stage differences.

The decline in the upper thermal limit between larvae, pupae and adults in that order (Figure 2) suggests that the insects upper thermal limit could be influenced by ontogeny as observed in *Drosophila melanogaster* (Bowler & Terblanche, 2008), although we did not test for age effects in a single life stage. This upper thermal limit decline across the life stages however, does not always follow a logical order as in some insect's e.g. *Drosophila buzzatii*; heat tolerance was high in pupae followed by eggs and larvae (as cited in Vorhees & Bradley, 2012). Therefore, caution must be made against the generalisation concerning the effects of ontogeny on heat resistance (Vorhees & Bradley, 2012), although we should expect ontogenetic variation (Bowler & Terblanche, 2008). One would have expected in part, no significant difference in the CT_{max} between larvae and adults because of the several factors like the microclimate conditions they experienced and the extent to which they can use their physiology to avoid the extreme temperatures. The lower CT_{max} of the adults compared with the larvae may be because the adults have a higher ability to avoid extreme temperatures by flying away from areas of extreme temperatures compared to larvae. The results suggest that there are ecological reasons for different thermal preferences: larvae are more sedentary so it is harder for them to move away from hotter temperatures they may be exposed to; adults are more mobile, so can move to preferred temperatures refuges in the evening so have a lower thermal threshold. Because of the ectothermic and regional heterothermic nature of insects, high temperatures could speed up growth and development of insects (Andrew et al., 2011; DeVries et al., 2016). The higher CT_{max} of the larvae could assist them to grow faster in warmer conditions like summer if food, and most importantly water, are not limiting factors.

The CT_{max} exhibited by the life stages is significantly greater than the highest recorded maximum daily mean ambient temperature in Tamworth region where the population originated from (40.9°C) (NicheMapR). However, the microclimatic conditions that *H. punctigera* could be exposed to may be much higher than this, particularly during the middle of the day (Andrew et al., 2013), and can put the insects under physiological stress that can change their interactions under changing climate. The higher CT_{max} exhibited by the insect could be the reason for which it is able to occupy a wide range of climatic conditions including inland of Australia, and spread in all states of Australia (Gregg et al., 2018, 2019).

Our results show that upper thermal limits were not significantly different between males and females. The similar thermal limits demonstrated by males and females indicates that sex has minimal influence on CT_{max} (Figure 4), similar to the findings in bumble bees (*Bombus terrestris*) (Oyen et al., 2016); suggesting that *H. punctigera* sex ratios will not change with adult exposure to climatic extremes.

Insects as other animals require energy to forage, survive, grow and reproduce (Terblanche et al., 2004; White et al., 2019). The processes that cause variation in metabolic rate apart from ambient temperature, body mass is also an important factor that contributes to metabolic rate of insects (White et al., 2019). Insects show a range of thermoregulatory capacities: ectothermic, regional heterothermic, poikilothermic and endothermic (Heinrich, 1993; Gullan & Cranston, 2014) their activity and physiological processes are primarily influenced by ambient temperature (Nguyen et al., 2014). Our results revealed $\dot{V}CO_2$ varied among the life stages at the different temperature points. The differences in the mean $\dot{V}CO_2$ among the life stages suggests that the metabolic demands of the life stages differ. Several factors may account for the differences in the metabolic

rates among the life stages; including ontogeny, upper thermal tolerance, activity, body mass and feeding (Terblanche et al., 2005).

Fat body tissues play crucial roles in the life of insects, as they are involved in multiple metabolic functions like energy storage and utilisation in reply to the energy call or request of the insect (Arrese & Soulages, 2010). Muscle tissue usually has a higher metabolic rate than fat body tissue.

Muscle tissues are active and burns calories even at rest unlike fat body for storing excess energy.

The significantly higher $\dot{V}CO_2$ of larvae and adult compared with pupae could be due to low amount of fat body tissues in the larvae and high muscle tissues in the case of adult. Another reason for the high metabolic rate may be due to the high activities of larvae and adult compared with pupae, similar to that reported by DeVries et al. (2013) for the bed bug, *Cimex lectularius*. Size also affects the metabolic rate of an organism (Terblanche et al., 2004; Quinlan et al., 2006; DeVries et al., 2013), and was similar in our case, as size was significant (Figure 8 in supplementary material). Pupae, which has a higher CT_{max} than adults, had a comparative lower $\dot{V}CO_2$. The lower $\dot{V}CO_2$ may be an indication of lower stress at the higher temperatures and or maybe due to more fat tissues in pupae. It is plausible that it may be an adaption strategy for survival to dry conditions (Terblanche et al., 2004; Quinlan et al., 2006). The high amount of fat tissues in the pupae because of lipid reserves needed to provide energy during the extended non-feeding periods could be the reason for the low metabolism (Arrese & Soulages, 2010). Because pupae do not have a source of free water underground, they must conserve water by reducing respiration (Table 3 in supplementary material) (Harrison, 2009).

As the effect of temperature on metabolic rates of insects is widely reported (Terblanche et al., 2004, 2007; Hill et al., 2016), only a few studies looked at that at different temperature points for different life stages for *H. punctigera*. Our results on metabolic rate between life stages considering

the different temperatures (Table 2) and Q_{10} (Figure 3) indicates that there was a correlation between temperature rise and $\dot{V}CO_2$ release on the life stages during the thermolimit respiratory (Table 2). The Q_{10} values for all the life stages except pupae almost doubled from Q_{10} : 25–35 °C to Q_{10} : 35– 45 °C, with adults having the highest thermal sensitivity. This increase in metabolic rate as temperature increases is consistent with other studies (Terblanche et al., 2005; Terblanche & Chown 2007; Andrew et al., 2016). However, the negative effects associated with increasing temperatures on metabolic efficiency could be reduced on adults, as they are able to compensate for short-term extreme microclimate temperatures by flying away. Although, increase in temperatures result in high metabolic rate, the total energy assigned for certain actions, as specific dynamic actions are not dependent on temperature (McCue et al., 2016a; Wang et al., 2003). It is possible that as insects enlarge their geographic radius or scale, they will face abiotic and biotic stress.

For better comparison among life stages, we measured $\dot{V}CO_2$ at CT_{max} , and in all cases, $\dot{V}CO_2$ among life stages was higher. Even though the $\dot{V}CO_2$ at CT_{max} increased in all the life stages, adults $\dot{V}CO_2$ were higher than larvae and pupae. Harrison (2009) reported that tracheated insects have the ability to carry or move gases by opening their spiracles. Short periods of spiracles closure follow a continuous carbon dioxide exchange (Harrison, 2009). The higher metabolic rates due to higher stress levels may be due to a decrease between spiracle opening and closing. Therefore, the higher $\dot{V}CO_2$ of adults at CT_{max} may be due to their high activity rate and a decrease between its spiracle opening and closing compared to larvae and pupae (Harrison, 2009; Boardman et al., 2015). The lower metabolic rate of pupae at CT_{max} compared to larvae and adults could not be explained beyond a physiological adaption strategy to reduce physiological cost by reducing respiration and

conserve water at this stage (Harrison, 2009), similar to what Merkey et al. (2011) reported in *Drosophila melanogaster* pupae.

The lower $\dot{V}CO_2$ of adults compared to larvae at 25°C could suggest that adults have a lower resting metabolic rate than larvae at that point. The higher resting metabolic rate of larvae could be the result of modification of the energy allocation between competing functions such as maintenance for continuous development. Additionally, the lower $\dot{V}CO_2$ of adults may be as result of the occurrence of discontinuous gas exchange in the adults at 25°C. Discontinuous gas exchange is associated with low metabolic rates. In contrast, age, as proven to affect metabolic rates in some insects like *Glossina pallidipes* (Terblanche et al., 2004) did not apply in our case, as larvae had a lower $\dot{V}CO_2$. The high $\dot{V}CO_2$ produced by the larvae could imply a high-energy utilisation (Terblanche et al., 2004) for development. This high-energy utilisation can provide a deep understanding into the life histories of the life stage across a variety of environments. Therefore, this high-energy utilisation may translate to an increase in food consumption and more considerable plant damage (Deutsch et al., 2018). The metabolic rate of pupae and adults was not influenced by sex, suggesting that the cost of living of the dispersing adults and pupae of both sexes are similar. Similar work by Terblanche et al. (2004) found out that metabolic rate did not differ between males and females of *Glossina pallidipes* and *G. morsitans*.

Conclusion

The present study provides vital information on how the different life stages of *H. punctigera* will physiologically perform in a changing thermal environment. It also provides data for predictions about metabolic rates of *H. punctigera* among life stages. The study revealed that CT_{max} of larvae was significantly higher than those of pupae and adults, and the metabolic rate of pupae was

significantly different from those of larvae and adults. Also, sex did not influence CT_{max} and metabolic rate. The results suggest that the performance and survival of the different life stages of *H. punctigera* could differ. This indicates that life-stage needs to be taken into account when assessing responses to a rapidly changing change, as ontogeny influences vulnerability and exposure. This finding can be used to predict the potential spread of the insect to other new environments and can indirectly help in management planning.

Acknowledgements

Sarah Hill assisted with the setup, calibration, maintenance and protocols of the thermolimit respirometry system. Zac Hemmings assisted in the data extraction, analysis and initial interpretations of the raw-data outputs. We thank Bianca Boss-Bishop, Sarah Hill, James O’Hanlon, Zac Hemmings, Alva Rebecka Curtsdotter and Behnaz Ghaedi for their comments on earlier versions of the manuscript. Also NSW DPI for providing the insects for the culture.

Declarations of interest: Nigel R. Andrew is an Academic editor of PeerJ

Funding

University of New England (UNE) provided funds as scholarship to Samuel Abukari Bawa as well as for equipment and consumables used in the study. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

All authors conceived and designed the experiment. The first author (Samuel Abukari Bawa) performed the experiment. The data was analysed by Samuel Abukari Bawa and Cara Miller. Samuel Abukari Bawa wrote the paper, prepared figures and tables. Nigel R. Andrew, Peter C. Gregg and Alice P. Del Socorro reviewed the drafts of the paper and contributed to writing the paper.

References

- Altermatt, F. 2010.** Climatic warming increases voltinism in European butterflies and moths. *Proceedings of the Royal Society of London B: Biological Sciences*, **277**(1685), 1281-1287.
- Arrese, E. L., & Soulages, J. L. 2010.** Insect fat body: energy, metabolism, and regulation. *Annual review of entomology*, **55**, 207-225
- Andrew, N. R., Ghaedi, B., & Groenewald, B. 2016.** The role of nest surface temperatures and the brain in influencing ant metabolic rates. *Journal of thermal biology*, **60**, 132-139.
- Andrew, N. R., Hart, R. A., Jung, M.-P., Hemmings, Z., & Terblanche, J. S. 2013.** Can temperate insects take the heat? A case study of the physiological and behavioural responses in a common ant, *Iridomyrmex purpureus* (Formicidae), with potential climate change. *Journal of Insect Physiology*, **59**(9), 870-880.
- Andrew, N.R. 2013.** Population dynamics of insects: impacts of a changing climate. *The Balance of Nature and Human Impact* (ed. K. Rohde), pp. 311-323. Cambridge University Press.
- Andrew, N. R., & Terblanche, J. S. 2013.** The response of insects to climate change. *Climate of Change: Living in a Warmer World*. Auckland: David Bateman Ltd, 38-50.

- 481 **Andrew, N. R., Hart, R. A., & Terblanche, J. S. 2011.** Limited plasticity of low temperature
482 tolerance in an Australian cantharid beetle *Chauliognathus lugubris*. *Physiological Entomology*,
483 **36**(4), 385-391.
- 484 **Andrew, N. R., & Hill, S. J. 2017.** Effect of Climate Change on Insect Pest Management.
485 *Environmental Pest Management: Challenges for Agronomists, Ecologists, Economists and*
486 *Policymakers*, 197.
- 487 **Angilletta Jr, M. J., & Angilletta, M. J. 2009.** Thermal adaptation: a theoretical and empirical
488 synthesis: Oxford University Press.
- 489 **Angilletta, J., Michael J, & Dunham, A. E. 2003.** The temperature-size rule in ectotherms:
490 simple evolutionary explanations may not be general. *The American Naturalist*, **162**(3), 332-342.
- 491 **Angilletta Jr, M. J., Wilson, R. S., Navas, C. A., & James, R. S. 2003.** Tradeoffs and the
492 evolution of thermal reaction norms. *Trends in Ecology & Evolution*, **18**(5), 234-240.
- 493 **Agudelo-Cantero, G. A., & Navas, C. A. 2019.** Interactive effects of experimental heating rates,
494 ontogeny and body mass on the upper thermal limits of anuran larvae. *Journal of thermal biology*,
495 **82**, 43-51.
- 496 **Arnold, P. A., Nicotra, A. B., & Kruuk, L. E. 2019.** Sparse evidence for selection on phenotypic
497 plasticity in response to temperature. *Philosophical Transactions of the Royal Society B*,
498 **374**(1768), 20180185.
- 499 **Bale, J. 2010.** Implications of cold-tolerance for pest management. Low temperature biology of
500 insects, 342-373.

- 501 **Bale, J. S., Masters, G. J., Hodkinson, I. D., Awmack, C., Bezemer, T. M., Brown, V. K.,**
502 **Farrar, J. 2002.** Herbivory in global climate change research: direct effects of rising temperature
503 on insect herbivores. *Global Change Biology*, **8**(1), 1-16.
- 504 **Betini, G. S., Griswold, C. K., & Norris, D. R. 2013.** Carry-over effects, sequential density
505 dependence and the dynamics of populations in a seasonal environment. Proceedings of the Royal
506 Society of London B: *Biological Sciences*, **280**(1759), 20130110.
- 507 **Betz A. & Andrew N. R. 2020.** Influence of non-lethal doses of natural insecticides Spinetoram
508 and Azadirachtin on *Helicoverpa punctigera* (native budworm, Lepidoptera: Noctuidae) under
509 laboratory conditions. *Front. Inv. Physiol.*
- 510 **Boardman, L., & Terblanche, J. S. 2015.** Oxygen safety margins set thermal limits in an insect
511 model system. *Journal of Experimental Biology*, **218**(11), 1677-1685.
- 512 **Bowler, K. and Terblanche, J. S. 2008.** Insect thermotolerance: what is the role of ontogeny,
513 ageing and senescence? *Biol. Rev.* **83**, 339-355.
- 514 **Chown, S.L., Terblanche, J.S. 2006.** Physiological diversity in insects: ecological and
515 evolutionary contexts. *Advances in insect physiology* **33**, 50-152.
- 516 **Daly, J. C., & Gregg, P. 1985.** Genetic variation in *Heliothis* in Australia: species identification
517 and gene flow in the Two-pest species *H. armigera* (Hübner) and *H. punctigera* Wallengren
518 (Lepidoptera: Noctuidae). *Bulletin of Entomological Research*, **75**(1), 169-184.
- 519 **Duncan, F. D., Krasnov, B., & McMaster, M. 2002.** Novel case of a tenebrionid beetle using
520 discontinuous gas exchange cycle when dehydrated. *Physiological Entomology*, **27**(1), 79-83.

Davidson, E. A., Janssens, I. A., & Luo, Y. (2006). On the variability of respiration in terrestrial ecosystems: moving beyond Q10. *Global Change Biology*, **12**(2), 154-164.

Dawson, T. P., Jackson, S. T., House, J. I., Prentice, I. C., & Mace, G. M. 2011. Beyond predictions: biodiversity conservation in a changing climate. *Science*, **332**(6025), 53-58.

Deutsch, C. A., Tewksbury, J. J., Tigchelaar, M., Battisti, D. S., Merrill, S. C., Huey, R. B., & Naylor, R. L. 2018. Increase in crop losses to insect pests in a warming climate. *Science*, **361**(6405), 916-919.

DeVries, Z. C., Kells, S. A., & Appel, A. G. 2013. Standard metabolic rate of the bed bug, *Cimex lectularius*: effects of temperature, mass, and life stage. *Journal of Insect Physiology*, **59**(11), 1133-1139.

DeVries, Z. C., Kells, S. A., & Appel, A. G. 2016. Estimating the critical thermal maximum (CT_{max}) of bed bugs, *Cimex lectularius*: comparing thermolimit respirometry with traditional visual methods. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, **197**, 52-57.

Diffenbaugh, N. S., Krupke, C. H., White, M. A., & Alexander, C. E. 2008. Global warming presents new challenges for maize pest management. *Environmental Research Letters*, **3**(4), 044007.

Dylan, B. 2006. Metabolic rate variation in *Glossina pallidipes* (Diptera: Glossinidae): gender, ageing and repeatability. Variability among individuals and populations: implications for arthropod physiology, 40

541 **Desforbes, J.-P., Marques, G. M., Beumer, L. T., Chimienti, M., Blake, J., Rowell, J. E., van**
542 **Beest, F. M. 2019.** Quantification of the full lifecycle bioenergetics of a large mammal in the high
543 Arctic. *Ecological Modelling*, **401**, 27-39

544 **Gaitán-Espitia, J. D., Bacigalupe, L. D., Opitz, T., Lagos, N. A., Timmermann, T., &**
545 **Lardies, M. A. 2014.** Geographic variation in thermal physiological performance of the intertidal
546 crab *Petrolisthes violaceus* along a latitudinal gradient. *Journal of Experimental Biology*, jeb.
547 108217

548 **Ghaedi, B., & Andrew, N. R. 2016.** The physiological consequences of varied heat exposure
549 events in adult *Myzus persicae*: a single prolonged exposure compared to repeated shorter
550 exposures. *PeerJ*, **4**, e2290.

551 **Greene, G., Leppla, N., & Dickerson, W. 1976.** Velvetbean caterpillar: a rearing procedure and
552 artificial medium. *Journal of Economic Entomology*, **69**(4), 487-488.

553 **Gregg, P. C., Del Socorro, A. P., Le Mottee, K., Tann, C. R., Fitt, G. P., & Zalucki, M. P.**
554 **2019.** Host plants and habitats of *Helicoverpa punctigera* and *H. armigera* (Hübner) (Lepidoptera:
555 Noctuidae) in inland Australia. *Austral Entomology*, **58**(3), 547-560.

556 **Gregg, P.C., Del Socorro, A.P., Le Mottee, K., Tann, C.R., Fitt, G.P., and Zalucki, M.P. 2018.**
557 Host plants and habitats of *Helicoverpa punctigera* and *H. armigera* (Hübner) (Lepidoptera:
558 Noctuidae) in inland Australia. *Austral Entomology*.

559 **Gregg, P. C., Henderson, G. S., Del Socorro, A. P., Le Mottee, K., & Birchall, C. 2016.**
560 Polyphagy in an uncertain environment: *Helicoverpa punctigera* in inland Australia. *Austral*
561 *Ecology*, **41**(7), 819-828. doi:10.1111/aec.12374

562 **Gullan P. J. & Cranston P. S. 2014.** *The Insects: An Outline of Entomology*. Wiley Blackwell
563 Publishing, Chichester, UK.

564 **Harris, R. M. B., Beaumont, L. J., Vance, T. R., Tozer, C. R., Remenyi, T. A., Perkins**
565 **Karasov, W. H., & del Rio, C. M. 2007.** Physiological ecology: how animals process energy,
566 nutrients, and toxins: Princeton University Press.

567 **Kirkpatrick, S. E., Bowman, D. M. J. S. 2018.** Biological responses to the press and pulse of
568 climate trends and extreme events. *Nature Climate Change*, **8**(7), 579-587. doi:10.1038/s41558-
569 018-0187-9

570 **Harrison, J. 2009.** "Respiratory System," in *Encyclopedia of Insects*, ed. R.C. Vincent
571 Resh. 2 ed (Amsterdam: Elsevier), 1168.

572 **Hazell, S. P., Neve, B. P., Groutides, C., Douglas, A. E., Blackburn, T. M., & Bale, J. S. 2010.**
573 Hyperthermic aphids: insights into behaviour and mortality. *J Insect Physiol*, **56**(2), 123-131.
574 doi:10.1016/j.jinsphys.2009.08.022

575 **Hodgson, C. 1991.** Dispersal of *Apterous aphids* (Homoptera: Aphididae) from their host plant
576 and its significance. *Bulletin of Entomological Research*, **81**(4), 417-427.

577 **Heinrich, Bernd 1993.** The hot-blooded insects: Strategies and mechanisms of thermoregulation,
578 *Cambridge, Massachusetts: Harvard University Press, p. 601*, ISBN 978-0-674-40838-8

579 **Huang, J., & Li, J. 2015.** Effects of climate change on overwintering pupae of the cotton
580 bollworm, *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae). *Int J Biometeorol*, **59**(7),
581 863-876. doi:10.1007/s00484-014-0903-8

582 **Hill, M. P., Bertelsmeier, C., Clusella-Trullas, S., Garnas, J., Robertson, M. P., &**
583 **Terblanche, J. 2016.** Predicted decrease in global climate suitability masks regional complexity
584 of invasive fruit fly species response to climate change. *Biological Invasions*, **18**(4), 1105-1119.

585 **IPCC. 2019.** IPCC Special Report on Climate Change, Desertification, Land Degradation,
586 Sustainable Land Management, Food Security, and Greenhouse gas fluxes in Terrestrial
587 Ecosystems. Summary for Policymakers.

588 **Kirkpatrick, T. 1961.** Comparative morphological studies of *Heliothis* species (Lepidoptera:
589 Noctuidae) in Queensland. *Queensland journal of agricultural science*, **18**(2).

590 **Kolluru, G. R., Chappell, M. A., & Zuk, M. 2004.** Sex differences in metabolic rates in field
591 crickets and their dipteran parasitoids. *Journal of Comparative Physiology B*, **174**(8), 641-648.

592 **Kingsolver, J. G. 2009.** The Well-Tempered Biologist: (American Society of Naturalists
593 Presidential Address). *The American Naturalist*, **174**(6), 755-768

594 **Kingsolver, J. G., Arthur Woods, H., Buckley, L. B., Potter, K. A., MacLean, H. J., &**
595 **Higgins, J. K. 2011.** Complex life cycles and the responses of insects to climate change. In: Oxford
596 University Press.

597 **Kellermann, V., van Heerwaarden, B., & Sgrò, C. M. 2017.** How important is thermal history?
598 Evidence for lasting effects of developmental temperature on upper thermal limits in *Drosophila*
599 *melanogaster*. *Proceedings of the Royal Society B: Biological Sciences*, **284**(1855), 20170447.

600 **Kingsolver, J. G., & Umbanhowar, J. 2018.** The analysis and interpretation of critical
601 temperatures. *Journal of Experimental Biology*, **221**(12), jeb167858.

- 602 **Lutterschmidt, W.I., and Hutchison, V.H. 1997.** The critical thermal maximum: 509 history and
603 critique. *Canadian Journal of Zoology* **75**(10), 1561-1574.
- 604 **Lighton, J. R., & Turner, R. J. 2004.** Thermolimit respirometry: an objective assessment of
605 critical thermal maxima in two sympatric desert harvester ants, *Pogonomyrmex rugosus* and *P.*
606 *californicus*. *Journal of Experimental Biology*, **207**(11), 1903-1913.
- 607 **Le Lann, C., Wardziak, T., Van Baaren, J., & van Alphen, J. J. 2011.** Thermal plasticity of
608 metabolic rates linked to life-history traits and foraging behaviour in a parasitic wasp. *Functional*
609 *Ecology*, **25**(3), 641-651
- 610 **Lignot, J., LeMaho, Y., & McCue, M. 2012.** Comparative Physiology of Fasting, Starvation, and
611 Food Limitation.
- 612 **Li, C., Wang, L., Li, J., Gao, C., Luo, Y., & Ren, L. 2019.** Thermal survival limits of larvae
613 and adults of *Sirex noctilio* (Hymenoptera: Siricidae) in China. *PLoS One*, **14**(6).
- 614 **Marais, E., Klok, C. J., Terblanche, J. S., & Chown, S. L. 2005.** Insect gas exchange patterns:
615 a phylogenetic perspective. *Journal of Experimental Biology*, **208**(23), 4495-4507.
- 616 **Marais, E., Terblanche, J. S., & Chown, S. L. 2009.** Life stage-related differences in hardening
617 and acclimation of thermal tolerance traits in the kelp fly, *Paractora dreuxi* (Diptera,
618 Helcomyzidae). *Journal of Insect Physiology*, **55**(4), 336-343.
- 619 **Martínez, J. D., Cadena, C. D., & Torres, M. 2016.** Critical thermal limits of *Poecilia caucana*
620 (Steindachner, 1880) (Cyprinodontiformes: Poeciliidae). *Neotropical Ichthyology*, **14**(1)

- 621 **McCue, M. D., Boardman, L., Clusella-Trullas, S., Kleynhans, E., & Terblanche, J. S. 2016.**
622 The speed and metabolic cost of digesting a blood meal depends on temperature in a major disease
623 vector. *Journal of Experimental Biology*, **219**(12), 1893-1902.
- 624 **Merkey, A. B., Wong, C. K., Hoshizaki, D. K., & Gibbs, A. G. 2011.** Energetics of
625 metamorphosis in *Drosophila melanogaster*. *Journal of Insect Physiology*, **57**(10), 1437-1445.
- 626 **Mironidis, G., & Savopoulou-Soultani, M. 2008.** Development, survivorship, and reproduction
627 of *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) under constant and alternating
628 temperatures. *Environmental Entomology*, **37**(1), 16-28.
- 629 **Mironidis, G. K. 2014.** Development, survivorship and reproduction of *Helicoverpa armigera*
630 (Hübner) (Lepidoptera: Noctuidae) under fluctuating temperatures. *Bulletin of Entomological*
631 *Research*, **104**(6), 751-764.
- 632 **Moyano, M., Candebat, C., Ruhbaum, Y., Álvarez-Fernández, S., Claireaux, G.,**
633 **Zambonino-Infante, J.-L., & Peck, M. A. 2017.** Effects of warming rate, acclimation
634 temperature and ontogeny on the critical thermal maximum of temperate marine fish larvae. *PLoS*
635 *One*, **12**(7), e0179928.
- 636 **Montejo-Kovacevich, G., Martin, S. H., Meier, J. I., Bacquet, C. N., Monllor, M., Jiggins, C.**
637 **D., & Nadeau, N. J. 2020.** Microclimate buffering and thermal tolerance across elevations in a
638 tropical butterfly. *Journal of Experimental Biology*, **223**(8).
- 639 **Mundim, K. C., Baraldi, S., Machado, H. G., & Vieira, F. M. 2020.** Temperature coefficient
640 (Q10) and its applications in biological systems: Beyond the Arrhenius theory. *Ecological*
641 *Modelling*, 431, 109127.

- 642 **Nguyen, C., Bahar, M. H., Baker, G., & Andrew, N. R. 2014.** Thermal tolerance limits of
643 diamondback moth in ramping and plunging assays. *PLoS One*, **9**(1), e87535.
- 644 **Nyamukondiwa, C., & Terblanche, J. S. 2010.** Within-generation variation of critical thermal
645 limits in adult Mediterranean and Natal fruit flies *Ceratitis capitata* and *Ceratitis rosa*: thermal
646 history affects short-term responses to temperature. *Physiological Entomology*, **35**(3), 255-264.
- 647 **Nyamukondiwa, C., Weldon, C. W., Chown, S. L., le Roux, P. C., & Terblanche, J. S. 2013.**
648 Thermal biology, population fluctuations and implications of temperature extremes for the
649 management of two globally significant insect pests. *Journal of Insect Physiology*, **59**(12), 1199-
650 1211.
- 651 **O'Neill, K. M., Kemp, W. P., & Johnson, K. A. 1990.** Behavioural thermoregulation in three
652 species of robber flies (Diptera, Asilidae: Efferia). *Animal Behaviour*, **39**(1), 181-191.
- 653 **Oyen, K. J., Giri, S., & Dillon, M. E. 2016.** Altitudinal variation in bumblebee (*Bombus*) critical
654 thermal limits. *Journal of thermal biology*, **59**, 52-57.
- 655 **Qayyum, A., & Zalucki, M. 1987.** Effects of high temperature on survival of eggs of *Heliothis*
656 *armigera* (Hübner) and *H. punctigera* Wallengren (Lepidoptera: Noctuidae). *Austral Entomology*,
657 **26**(4), 295-298.
- 658 **Quinlan, M. C., & Gibbs, A. G. 2006.** Discontinuous gas exchange in insects. *Respiratory*
659 *physiology & neurobiology*, **154**(1-2), 18-29.
- 660 **Room, P. 1983.** Calculations of temperature-driven development by *Heliothis spp.* (Lepidoptera:
661 Noctuidae) in the Namoi Valley, New South Wales. *Journal of the Australian Entomological*
662 *Society*, **22**(3), 211-215.

663 Secor, S. M. (2009). Specific dynamic action: a review of the postprandial metabolic response.
 664 *Journal of Comparative Physiology B*, 179(1), 1-56.

665 **Somero, G. 2010.** The physiology of climate change: how potentials for acclimatization and
 666 genetic adaptation will determine ‘winners’ and ‘losers’. *Journal of Experimental Biology*, **213**(6),
 667 912-920.

668 **Stevens, M. M., Jackson, S., Bester, S.A., Terblanche, J.S., & Chown, S.L. 2010.** Oxygen
 669 limitation and thermal tolerance in two terrestrial arthropod species. *Journal of Experimental*
 670 *Biology* (13), 2209-2218.

671 **SAS Institute Inc., 2013.** SAS/STAT 13.1 User's Guide. SAS Institute Inc., Cary, NC.

672 **Sunday, J. M., Bates, A. E., Kearney, M. R., Colwell, R. K., Dulvy, N. K., Longino, J. T., &**
 673 **Huey, R. B. 2014.** Thermal-safety margins and the necessity of thermoregulatory behavior
 674 across latitude and elevation. *Proceedings of the National Academy of Sciences*, 201316145.

675 **Shik, J. Z., Arnan, X., Oms, C. S., Cerdá, X., & Boulay, R. 2019.** Evidence for locally
 676 adaptive metabolic rates among ant populations along an elevational gradient. *Journal of animal*
 677 *ecology*, **88**(8), 1240-1249

678 **Shah, A. A., Woods, H. A., Havird, J. C., Encalada, A. C., Flecker, A. S., Funk, W. C., &**
 679 **Thomas, S. A. 2021.** Temperature dependence of metabolic rate in tropical and temperate aquatic
 680 insects: Support for the Climate Variability Hypothesis in mayflies but not stoneflies. *Global*
 681 *Change Biology*, 27(2), 297-311.

682 **Teakle, R. 1991.** Laboratory culture of *Heliothis* species and identification of disease. In *Heliothis:*
 683 *research methods and prospects* (pp. 22-29): *Springer*.

684 **Terblanche, J., Klok, C., & Chown, S. 2004.** Metabolic rate variation in *Glossina pallidipes*
685 (Diptera: Glossinidae): gender, ageing and repeatability. *Journal of Insect Physiology*, **50**(5), 419-
686 428.

687 **Terblanche, J. S., & Chown, S. L. 2007.** The effects of temperature, body mass and feeding on
688 metabolic rate in the tsetse fly *Glossina morsitans centralis*. *Physiological Entomology*, **32**(2),
689 175-180.

690 **Terblanche, J. S., Deere, J. A., Clusella-Trullas, S., Janion, C., & Chown, S. L. 2007.** Critical
691 thermal limits depend on methodological context. *Proceedings of the Royal Society of London B:*
692 *Biological Sciences*, **274** (1628), 2935-2943.

693 **Terblanche, J.S., Clusella-Trullas, S., Chown, S.L. 2010.** Phenotypic plasticity of gas exchange
694 pattern and water loss in *Scarabaeus spretus* (Coleoptera: Scarabaeidae): deconstructing the basis
695 for metabolic rate variation. *Journal of Experimental Biology* **213**, 2940-2949.

696 **Terblanche, J. S., Klok, C. J., & Chown, S. L. 2005.** Temperature-dependence of metabolic rate
697 in *Glossina morsitans morsitans* (Diptera, Glossinidae) does not vary with gender, age, feeding,
698 pregnancy or acclimation. *Journal of Insect Physiology*, **51**(8), 861-870.

699 **Terblanche, J. S., Sinclair, B. J., Klok, C. J., McFarlane, M. L., & Chown, S. L. 2005.** The
700 effects of acclimation on thermal tolerance, desiccation resistance and metabolic rate in *Chirodica*
701 *chalconota* (Coleoptera: Chrysomelidae). *Journal of Insect Physiology*, **51**(9), 1013-1023.

702 **Terblanche, J., Klok, C., & Chown, S. 2004.** Metabolic rate variation in *Glossina pallidipes*
703 (Diptera: Glossinidae): gender, ageing and repeatability. *Journal of Insect Physiology*, **50**(5), 419-
704 428.

705 **Thienel, M., Canals, M., Bozinovic, F., & Veloso, C. 2015.** The effects of temperature on the
 706 gas exchange cycle in *Agathemera crassa*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, **183**, 126-130.

707

708 **The R Foundation** for Statistical Computing Platform (C) **2018** (version 3.5.2.)

709 **Tomlinson, S., & Phillips, R. D. 2015.** Differences in metabolic rate and evaporative water loss
 710 associated with sexual dimorphism in thynnine wasps. *Journal of Insect Physiology*, **78**, 62-68.

711 **Vorhees, A. S., & Bradley, T. J. 2012.** Differences in critical thermal maxima and mortality
 712 across life stages of the mealworm beetle *Tenebrio molitor*. *Journal of Experimental Biology*, **215**
 713 (13), 2319-2326.

714 **Wang, T., Zaar, M., Arvedsen, S., Vedel-Smith, C., & Overgaard, J. 2002.** Effects of
 715 temperature on the metabolic response to feeding in *Python molurus*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, **133**(3), 519-527.

716

717 **Whitman, D. W., & Agrawal, A. A. 2009.** What is phenotypic plasticity and why is it important.
 718 Phenotypic plasticity of insects: Mechanisms and consequences, 1-63.

719 www.bom.gov.au/state-of-the-climate (2018).

720 **White, C. R., Marshall, D. J., Alton, L. A., Arnold, P. A., Beaman, J. E., Bywater, C. L.,**
 721 **Pirtle, E. 2019.** The origin and maintenance of metabolic allometry in animals. *Nature Ecology &*
 722 *Evolution*, **3**(4), 598-603.

723 **Zalucki, M., Daglish, G., Firempong, S., & Twine, P. 1986.** The biology and ecology of
 724 *Heliothis-armigera* (Hübner) and *Heliothis-punctigera* Wallengren (Lepidoptera: Noctuidae) in
 725 Australia-What do we know. *Australian Journal of Zoology*, **34**(6), 779-814.

726 **Zalucki, M. P., & Furlong, M. J. 2005.** Forecasting *Helicoverpa* populations in Australia: a
 727 comparison of regression based models and a bioclimatic-based modelling approach. *Insect*
 728 *Science*, **12**(1), 45-56.

Figure 1

Box plot showing the Total mean $\dot{V}\text{CO}_2$ ml h⁻¹ among life stages (larvae, pupae and adult) of *H. punctigera* over complete ramping period of about 120mins from 25°C to 55°C following thermolimit respirometry

The dark line is the median in each group




Figure 1: Metabolic rate $\dot{V}\text{CO}_2$ (ml h^{-1}) boxplot of *H. punctigera* life stages (larvae, pupae and adult; $n = 10$ for all stages) over the complete ramping period of about 120mins from 25°C to CT_{max} following thermolimit respirometry. Significant differences ($p > 0.05$) indicated by the letter above each boxplot. The box represents the lower and upper quartiles with the dark line representing the mean. The bars represents the minimum and maximum whiskers, the dot represents an outlier.

Figure 2

Box plot showing the Mean CT_{max} identified as the inflection point in the absolute difference sum (ADS) residuals among life stages (larvae, pupae and adult) of *H. punctigera* following thermolimit respirometry.

The dark line is the median of the data in each group




Figure 2: Upper thermal limit (CT_{max}) identified as the inflection point in the absolute difference sum (ADS) residuals among life stages (larvae, pupae and adult; $n = 10$ for all stages) of *H. punctigera* following thermolimit respirometry. Box plots with the same letter are not significantly different ($p > 0.05$). The box represents the lower and upper quartiles with the dark line representing the mean. The bars represent the minimum and maximum whiskers, the dot represents an outlier.

Figure 3

Box plots comparisons of *H. punctigera* life stages (larvae, pupae and adult; n = 10 for all stages) Q₁₀ (25–35 °C) and Q₁₀ (35–45 °C).

The box represents the lower and upper quartiles with the dark line representing the mean.

The bars represents the minimum and maximum whiskers, the dots represent outliers.




Figure 3: Box plots comparisons of *H. punctigera* life stages (adult, larvae and pupae; $n = 10$ for all stages) Q_{10} (25–35 °C) and Q_{10} (35–45 °C). The box represents the lower and upper quartiles with the dark line representing the mean. The bars represents the minimum and maximum whiskers, the dots represent outliers.

Figure 4

Box plots of upper thermal limit (CT_{max}) identified as the inflection point in the absolute difference sum (ADS) residuals among Life Stage (pupae and adult) and sex (male – M and female – F) of *H. punctigera* following thermolimit respirom

The box represents the lower and upper quartiles with the dark line representing the mean. The bars represent the minimum and maximum whiskers, the dots represent outliers.




Figure 4: Box plots of upper thermal limit (CT_{max}) identified as the inflection point in the absolute difference sum (ADS) residuals among Life Stage (pupae and adult) and sex (male – M and female – F) of *H. punctigera* following thermolimit respirometry ; $n = 5$ for all sexes. The box represents the lower and upper quartiles with the dark line representing the mean. The bars represent the minimum and maximum whiskers, the dots represent outliers.

Figure 5

Box plots of $\dot{V}CO_2$ ml h⁻¹ among Life Stage (pupae and adult) and sex (male – M and female – F) of *H. punctigera* over the complete temperature ramping period of about 140mins from 25°C to CT_{max} following thermo

The box represents the lower and upper quartiles with the dark line representing the median. The bars represents the minimum and maximum whiskers, the dots represent outliers.




Figure 5: Box plots of $\dot{V}\text{CO}_2$ ml h^{-1} among Life Stage (pupae and adult) and sex (male – M and female – F) of *H. punctigera* over the complete temperature ramping period of about 140mins from 25°C to CT_{max} following thermolimit respirometry. n=five in all cases; n = five for all sexes. The box represents the lower and upper quartiles with the dark line representing the median. The bars represents the minimum and maximum whiskers, the dots represent outliers.

Table 1(on next page)

Anova summary information for Activity CT_{max} , CO_2 CT_{max} and mean CT_{max} between two life stages (Larvae and Adults) of *H. punctigera* following thermolimit respirometry

1

2 Table 1: Anova summary information for Activity CT_{max} , CO_2 CT_{max} and mean CT_{max} between two life stages (Larvae
3 and Adults) of *H. punctigera* following thermolimit respirometry. n=10 for each life stage

Stage	DF	Sum Sq.	Mean Sq.	F	p	Mean CT_{max}	
						Life Stage	
						Larvae	Adults
Activity	1	8.3	8.28	0.21	0.65	44.1±1.3	43.1±2.5
CT_{max}							
CO_2 CT_{max}	1	133.1	133.08	4.27	0.05	49.0±0.3	43.5±2.4

4

Table 2 (on next page)

Mean \pm s.e. $\dot{V}CO_2$ ($\mu l CO_2 h^{-1}$) over different temperature points among life stages (adult, larvae, and pupae).

Values in bold indicate significant differences. Letters indicate significant differences among life stages. The 95% CL is 95% confidence limit

- 1 Table 2: Mean \pm s.e. $\dot{V}\text{CO}_2$ ($\mu\text{l CO}_2 \text{ h}^{-1}$) over different temperature points among life stages (adult, larvae, and pupae). Values in bold indicate significant differences.
- 2 Letters indicate significant differences among life stages. The 95% CL is 95% confidence limit

Life	Mean $\dot{V}\text{CO}_2$ ($\mu\text{l CO}_2 \text{ h}^{-1}$)											
Stage	25°C	95% CL	30°C	95% CL	35°C	95% CL	40°C	95% CL	45°C	95% CL	CT _{max}	95% CL
Larvae	0.40\pm0.04a	0.31-0.50	0.44\pm0.04a	0.43-0.62	0.53\pm0.04a	0.57-0.76	0.70\pm0.06a	0.54-0.81	1.03\pm0.22a	0.83-1.78	1.03\pm0.14a	0.90-1.50
Pupae	0.08\pm0.03b	-0.03-0.11	0.11\pm0.03b	0.02-0.15	0.14\pm0.03b	0.08-0.22	0.19\pm0.04b	-0.01-0.16	0.24\pm0.16b	-0.09-0.60	0.23\pm0.10b	-0.05-0.37
Adult	0.21\pm0.04b	0.16-0.36	0.31\pm0.04c	0.21-0.41	0.37\pm0.04c	0.22-0.43	0.71\pm0.06a	0.71-1.01	1.47\pm0.24c	0.91-1.92	1.23\pm0.15a	1.01-1.65

3

4

5

6

