

A comparison of two gluteus maximus EMG maximum voluntary isometric contraction positions

Bret Contreras, Andrew D Vigotsky, Brad J Schoenfeld, Chris Beardsley, John B Cronin

Background: The purpose of this study was to compare the peak electromyography (EMG) of the most commonly-used positions in the literature, the prone bent-leg (90°) hip extension against manual resistance applied to the distal thigh (PRONE), to a novel position, the standing glute squeeze (SQUEEZE).

Methods: Surface EMG electrodes were placed on the upper and lower gluteus maximus of thirteen recreationally active females (age = 28.9 years; height = 164 cm; body mass = 58.2 kg), before three maximum voluntary isometric contraction (MVIC) trials for each position were obtained in a randomized, counterbalanced fashion.

Results: No significant ($p \leq 0.05$) differences were observed between PRONE (upper: 91.94%; lower: 94.52%) and SQUEEZE (upper: 92.04%; lower: 85.12%) for both the upper and lower gluteus maximus. Neither the PRONE nor SQUEEZE was more effective between all subjects.

Conclusions: In agreement with other studies, no single testing position is ideal for every participant. Therefore, it is recommended that investigators employ multiple MVIC positions, when possible, to ensure accuracy. Future research should investigate a variety of gluteus maximus MVIC positions in heterogeneous samples.

A comparison of two gluteus maximus EMG maximum voluntary isometric contraction positions

Running Head: GMax MVIC

Bret Contreras, MA ¹

Andrew D. Vigotsky ²

Brad J. Schoenfeld, PhD ³

Chris Beardsley ⁴

John Cronin, PhD ^{1, 5}

¹ Auckland University of Technology, Sport Performance Research Institute New Zealand

² Kinesiology Program, Arizona State University, Phoenix, AZ

³ Department of Health Sciences, CUNY Lehman College, Bronx, NY

⁴ Strength and Conditioning Research Limited, London, UK

⁵ School of Exercise, Biomedical and Health Science, Edith Cowan University, Perth, Australia

INTRODUCTION

Maximum voluntary isometric contractions (MVIC) are often used to normalize electromyography (EMG) signals. It is important to employ an MVIC position that elicits the highest activation in order to increase the validity of EMG studies and decrease incidents of abnormally high normalized mean and peak EMG data. In order for accurate comparisons to be made between studies, it is also important for researchers to standardize MVIC positions, or at least use positions that elicit similar magnitudes of EMG activity. A number of MVIC positions have been used in the literature to assess the gluteus maximus, including the Biering-Sorenson position (Cambridge et al. 2012; McGill et al. 2009), the prone straight leg hip extension position (Barton et al. 2014; Worrell et al. 2001), the prone bent leg position (Jakobsen et al. 2013; Youdas et al. 2013), the prone straight leg position with 70° of hip flexion (Simenz et al. 2012), and the standing bent leg position (Boudreau et al. 2009). The most commonly used position, however, is the prone bent-leg (90°) hip extension with manual resistance applied to the distal thigh (PRONE) (Choi et al. 2014; Emami et al. 2014; Hislop et al. 2013; Kang et al. 2013; Kendall et al. 1993; Oh et al. 2007).

A recent study by Simenz et al. (2012) that used a prone gluteus maximus MVIC position in 70° of hip flexion, demonstrates the importance of standardizing MVIC positions across studies. Researchers have shown that lower gluteus maximus amplitude is elicited at higher degrees of hip flexion and reaches a maximum EMG amplitude at end range hip extension (Worrell et al. 2001). By employing an MVIC position that renders significantly lower EMG activity than those values that are truly maximal, the normalized data of Simenz et al. (2012) are most likely overestimated. For example, if the work of Worrell et al. (2001) is extrapolated, the MVIC

position used by Simenz et al. (2012) would only elicit approximately 80% of true MVIC, translating into 25% greater mean and peak values when compared to the true MVIC position. The data reported by Simenz et al. (2012) therefore cannot be used for comparison with exercises in other studies that utilized alternative MVIC positions with smaller hip flexion angles, as the data would have overestimated how effectively the gluteus maximus was activated. Therefore, it is apparent that researchers should only compare EMG data that utilize positions that render similar values.

Since Worrell et al. (2001) found that full hip extension elicited the greatest amount of gluteus maximus EMG activity, and this finding is corroborated by earlier work from Wheatley & Jahnke (1951) and Fischer & Houtz (1968), it is postulated that the most appropriate gluteus maximus MVIC position is at full hip extension, or hip hyperextension. PRONE is currently the recommended position in several texts on muscle testing (Hislop et al. 2013; Kendall et al. 1993), although to the authors' knowledge, this position has not been compared to others in the literature. In order to correct for individual variation, some researchers have employed multiple MVIC positions. For example, McGill et al. (2009) used both the Biering-Sorenson and PRONE positions; whichever position elicited the greatest activity was used for normalization purposes. The authors, however, are unaware of any existing research that quantitatively compares gluteus maximus MVIC positions.

The gluteus maximus muscle appears to be segmented into at least two subdivisions, which may display different EMG activity in response to certain muscle actions. McAndrew et al. (2006) used a laser-based mechanomyographic (MMG) technique to measure the mean contraction time

in six subdivisions of the gluteus maximus, both in the sagittal plane (superior, middle, inferior) and in the frontal plane (medial and lateral). The superior region displayed the longest contraction time followed by the middle region and then the inferior region. On the basis of these findings, McAndrew et al. (2006) suggested that the superior region may contain more slow twitch fibers and be more involved in postural tasks compared to the inferior region, while the inferior region may contain more fast twitch fibers and be more involved in dynamic tasks. This is further substantiated by the work of Lyons et al. (1983) and Karlsson & Jonsson (1965), who found differences between upper and lower gluteus maximus EMG during functional movement; for example, load acceptance during stair ambulation better targets the lower gluteus maximus (Lyons et al. 1983), while hip abduction better targets the upper gluteus maximus (Karlsson & Jonsson 1965).

Pilot data from our lab showed that some subjects were able to elicit greater EMG activity during a standing glute squeeze (SQUEEZE) when compared to PRONE, and this was especially true for the upper gluteus maximus. Given this observation and the findings articulated in previous paragraphs, the purpose of this investigation was to compare upper and lower gluteus maximus EMG activity in PRONE versus SQUEEZE. Based on our pilot data, it was hypothesized that SQUEEZE would elicit greater upper gluteus maximus EMG activity, while PRONE would elicit greater lower gluteus maximus EMG activity.

METHODS

Subjects

Thirteen healthy women (age = 28.9 ± 5.1 years; height = 164 ± 6.3 cm; body mass = 58.2 ± 6.4 kg) with 7.0 ± 5.8 years of resistance training experience participated in this study. Inclusion criteria required subjects to be between 20 to 40 years of age and have at least 3 years of consistent resistance training experience. All subjects were healthy and free of any musculoskeletal or neuromuscular injuries, pain, or illnesses. Subjects completed an Informed Consent form. Subjects were advised to refrain from training their lower body for 72 hours prior to testing. The study was approved by the Auckland University of Technology Ethics Committee (AUTEC Reference number 13/375).

Procedures

Subjects first performed a 10-minute general warm-up consisting of various dynamic stretches for the lower body musculature. Following warm-up, subjects practiced each testing position several times, until they felt comfortable with the technique. Subjects were asked to wear appropriate clothing for access to the EMG electrode placement sites. Before placing the electrodes on the skin, excess hair was removed with a razor, and skin was cleaned and abraded using an alcohol swab. After preparation, self-adhesive disposable silver/silver chloride pre-gelled dual snap surface bipolar electrodes (Noraxon Product #272, Noraxon USA Inc, Scottsdale, AZ) with a diameter of 1 centimeter (cm) and an inter-electrode distance of 2 cm were attached in parallel to the fibers of the right upper and lower gluteus maximus. More specifically, “[upper gluteus maximus] electrodes were placed two finger’s width above the line just under the spina iliaca posterior superior and the trochanter major; [lower gluteus maximus]

electrodes were set below the same line” (Fujisawa et al. 2014) (Figure 1, Figure 2). After the electrodes were secured, a quality check was performed to ensure EMG signal validity.

Following electrode placement, subjects completed three trials of PRONE then SQUEEZE, or vice versa. The PRONE position was performed by having the subject lie prone on the bench, flexing their knee to 90°, and was told to extending her hip while manual resistance was applied to the distal thigh (Figure 1). The SQUEEZE position was performed by having the subject stand with her feet slightly wider than shoulder width apart and hips slightly externally rotated. The subject was instructed to squeeze her glutei and focus on externally rotating and extending her hips (Figure 2). For example, if a subject was randomized to complete PRONE first, her testing order would be PRONE, SQUEEZE, rest, PRONE, SQUEEZE, rest, PRONE, SQUEEZE. Each contraction phase lasted 5 seconds, and each rest phase lasted 3 minutes. Randomization was counterbalanced so that half the subjects performed PRONE first and the other half performed SQUEEZE first. In all MVIC positions, subjects were instructed to contract the gluteus maximus “as hard as possible.”



128

129 **Figure 1.** Prone bent-leg hip extension position. Force is applied to the distal thigh while the

130 subject produces a hip extension moment.



Figure 2. Standing glute squeeze position. The subject squeezes her glutei, which generates hip extension and external rotation moments.

Raw EMG signals were collected at 2000 Hz by a Myotrace 400 EMG unit (Noraxon USA Inc, Scottsdale, AZ). Data was sent in real time to a computer via Bluetooth and recorded and

analyzed by MyoResearch 3.6 Clinical Applications software (Noraxon USA, Inc., Scottsdale, AZ). A 10-500 Hz bandpass filter was applied to EMG data. Signals of all MVIC trials were full-wave rectified and smoothed with a root mean square (RMS) algorithm with a 100 ms window. Maximal peak EMG values over a 1000 ms window, or the 1000 ms window with the greatest average EMG amplitude within the 5 second contraction, were then used to normalize peak EMG signals obtained during each MVIC trial (Vera-Garcia et al. 2010).

Statistical Analysis

Paired samples *t*-tests were performed after checking normality using Shapiro-Wilk test in Stata 13 (StataCorp LP, College Station, TX). Alpha was set *a priori* at 0.05 for significance. Effect sizes (ES) were calculated by Cohen's *d* using the formula $d = \frac{M_d}{s_d}$, where M_d is the mean of differences and s_d is the standard deviation of differences (Becker 1988; Morris 2007; Smith & Beretvas 2009). This method is slightly different than the traditional method of calculating Cohen's *d*, as it calculates the within-subject effect-size rather than group or between-subject effect size. ES were defined as small (0.20-0.49), moderate (0.50-0.79), and large (≥ 0.80) (Cohen 1988). Confidence intervals (95% CI) for each ES were also calculated.

RESULTS

The normalized peak EMG for the different exercises and gluteus maximus sections can be observed in Table 1. In terms of the upper gluteus maximus comparison, no significant differences were observed in the peak EMG for both exercises (ES = 0.005; 95% CI = -0.599 – 0.609; $t(12) = 0.018$; $p = 0.986$). With regards to the lower gluteus maximus, a small ES was

observed (0.412; 95% CI = -0.193 – 0.102) between the two positions in favor of the PRONE position; however, this outcome may have been due to chance alone ($t(12) = -1.485$; $p = 0.164$).

Table 1. Group mean \pm SD of normalized peak EMG amplitudes.

	PRONE	SQUEEZE
Upper gluteus maximus	91.94 \pm 11.64	92.04 \pm 11.30
Lower gluteus maximus	94.52 \pm 13.59	85.12 \pm 12.64

DISCUSSION

The purpose of this investigation was to compare a novel gluteus maximus MVIC position, SQUEEZE, to the current gold standard, PRONE. We have failed to reject the null hypotheses, as there were no statistically significant differences between the two positions tested (Table 1). However, despite no statistically significant differences, the peak EMG values for the lower gluteus maximus were approximately 9% higher for the PRONE compared to the SQUEEZE. Consequently, if the SQUEEZE test were used for normalization, it would render approximately 10% higher mean and peak EMG values compared to the PRONE test. Therefore, although not statistically significant, the findings could be considered practically meaningful. Furthermore, these data show a large amount of individual variation (Table 2), which has been previously described by McGill (1990) and Vera-Garcia et al. (2010) for other muscles.

Table 2. Number of subjects (percentage of subjects (%)) which each MVIC technique resulted in the greatest peak EMG amplitude.

	PRONE	SQUEEZE
Upper gluteus maximus	7 (53.85)	6 (46.15)
Lower gluteus maximus	10 (76.92)	3 (23.08)

There are several kinematic and kinetic differences between PRONE and SQUEEZE, any of which may have affected our results, either individually or in combination. During PRONE, the knee is bent to 90°, whereas during SQUEEZE, the knees are fully extended. Previous research has shown that gluteus maximus EMG activity during hip extension is greater with the knees flexed than when extended, presumably resulting from a greater reliance upon the gluteus maximus for hip extension due to decreased hamstrings length (Kwon & Lee 2013). On the other hand, extended knees allow for greater hip extension range of motion compared to flexed knees, thereby shortening the gluteal fibers to a greater extent (Van Dillen et al. 2000) and leading to a greater amount of gluteus maximus EMG activity (Worrell et al. 2001). In addition, PRONE involved primarily hip hyperextension since the pelvis was fixed, whereas SQUEEZE appeared to involve a combination of hip extension and posterior pelvic tilt. Although posterior pelvic tilt mimics hip extension (Neumann 2010), it is unclear how each of these kinematic variables might affect gluteus maximus EMG activity individually. To our knowledge, no study to date has investigated gluteus maximus EMG activity with varying combinations of hip extension and posterior pelvic tilt during MVIC actions. Moreover, PRONE is an open kinetic chain maneuver with the torso stabilized onto a bench, whereas SQUEEZE is a closed kinetic chain maneuver performed in a standing position. Stensdotter et al. (2003) investigated the EMG activity of the quadriceps muscle group during open kinetic chain and closed kinetic chain positions during MVIC actions and reported significant differences in EMG amplitude. The rectus femoris

displayed greater EMG activity during open kinetic chain maneuvers while the vastus medialis displayed greater EMG activity during closed kinetic chain maneuvers. It is therefore hard to predict whether the gluteus maximus would inherently display greater or lesser EMG activity during either open or closed kinetic chain maneuvers. Finally, PRONE required manual resistance, whereas SQUEEZE relied upon anatomical structures surrounding the hip to provide resistance against hip extension. Whether this factor has any effect on EMG activity recorded in a muscle is unclear, as the authors are unaware of any previous investigations into the effect of squeezing a muscle whereby range of motion is limited by anatomical structures on EMG activity rather than against external resistance.

This investigation was subject to several important limitations. Firstly, although we observed what may have been a practically important difference between the MVIC positions, this difference was not found to be statistically significant, which suggests that our initial estimates for the appropriate sample size may have been too small. Secondly, there were several kinematic differences between the two positions that were explored (PRONE and SQUEEZE), including different pelvic, hip, and knee joint angles. There were also kinetic differences between the two positions, in that PRONE was an open kinetic chain maneuver and SQUEEZE was a closed kinetic chain maneuver. Moreover, PRONE used external resistance and SQUEEZE utilized oppositional torques produced by internal, anatomical structures. These multiple differences make it difficult to assess whether our results arose from a combination of biomechanical factors acting in opposing directions, heterogeneity, or genuinely no difference between the conditions. Thirdly, we only compared two MVIC positions, and it is feasible that other positions might result in superior or inferior levels of EMG activity. Fourthly, we only investigated two

subdivisions of the gluteus maximus muscle and there are indications that there may be others, from proximal-to-distal, medial-to-lateral, and superficial-to deep. Furthermore, our statistical analysis was not designed to assess whether there was a difference between the EMG activity of the upper and lower gluteus maximus during either MVIC position and therefore this remains uncertain.

Conclusions

Although these data are inconclusive as to which position is superior, they do provide insight as to the complexity of MVIC positions for the gluteus maximus. More specifically, due to the large individual variations (Table 2), it is recommended that multiple MVIC positions be utilized to ensure that the greatest possible EMG amplitude be the divisor during normalization. These recommendations are well in line with other studies, which have utilized or recommended multiple MVIC positions (McGill et al. 2009; Vera-Garcia et al. 2010). Future research should use heterogeneous samples, such as athletic males, and also test more positions, such as the Biering-Sorenson position, quadruped hip extension position, and top hip thrust position (Contreras et al. 2011), each with manual resistance, along with the tall kneeling position.

REFERENCES

- Barton CJ, Kennedy A, Twycross-Lewis R, Woledge R, Malliaras P, and Morrissey D. 2014. Gluteal muscle activation during the isometric phase of squatting exercises with and without a Swiss ball. *Physical Therapy in Sport* 15:39-46.
- Becker BJ. 1988. Synthesizing standardized mean - change measures. *British Journal of Mathematical and Statistical Psychology* 41:257-278.
- Boudreau SN, Dwyer MK, Mattacola CG, Lattermann C, Uhl TL, and McKeon JM. 2009. Hip-muscle activation during the lunge, single-leg squat, and step-up-and-over exercises. *Journal of Sport Rehabilitation* 18:91-103.
- Cambridge ED, Sidorkewicz N, Ikeda DM, and McGill SM. 2012. Progressive hip rehabilitation: the effects of resistance band placement on gluteal activation during two common exercises. *Clinical Biomechanics (Bristol, Avon)* 27:719-724.
- Choi SA, Cynn HS, Yi CH, Kwon OY, Yoon TL, Choi WJ, and Lee JH. 2014. Isometric hip abduction using a Thera-Band alters gluteus maximus muscle activity and the anterior pelvic tilt angle during bridging exercise. *Journal of Electromyography and Kinesiology*.
- Cohen J. 1988. *Statistical power analysis for the behavioral sciences*: Routledge Academic.
- Contreras B, Cronin J, and Schoenfeld B. 2011. Barbell hip thrust. *Strength & Conditioning Journal* 33:58-61.
- Emami M, Arab AM, and Ghamkhar L. 2014. The activity pattern of the lumbo-pelvic muscles during prone hip extension in athletes with and without hamstring strain injury. *International Journal of Sports Physical Therapy* 9:312-319.
- Fischer FJ, and Houtz SJ. 1968. Evaluation of the function of the gluteus maximus muscle. An electromyographic study. *American Journal of Physical Medicine* 47:182-191.
- Fujisawa H, Suzuki H, Yamaguchi E, Yoshiki H, Wada Y, and Watanabe A. 2014. Hip Muscle Activity during Isometric Contraction of Hip Abduction. *Journal of Physical Therapy Science* 26:187-190.
- Hermens HJ, Freriks B, Merletti R, Stegeman D, Blok J, Rau G, Disselhorst-Klug C, and Hägg G. 1999. European recommendations for surface electromyography. *Roessingh Research and Development, Enschede*.
- Hislop H, Avers D, and Brown M. 2013. *Daniels and Worthingham's Muscle Testing: Techniques of Manual Examination and Performance Testing*: Elsevier Health Sciences.
- Jakobsen MD, Sundstrup E, Andersen CH, Aagaard P, and Andersen LL. 2013. Muscle activity during leg strengthening exercise using free weights and elastic resistance: effects of ballistic vs controlled contractions. *Human movement science* 32:65-78.
- Kang SY, Jeon HS, Kwon O, Cynn HS, and Choi B. 2013. Activation of the gluteus maximus and hamstring muscles during prone hip extension with knee flexion in three hip abduction positions. *Manual Therapy* 18:303-307.
- Karlsson E, and Jonsson B. 1965. Function of the Gluteus Maximus Muscle. An Electromyographic Study. *Acta Morphologica Neerlandico-Scandinavica* 6:161-169.
- Kendall FP, McCreary EK, Provance PG, Rodgers MM, and Romani W. 1993. Muscles, testing and function: with posture and pain.

- Kwon YJ, and Lee HO. 2013. How different knee flexion angles influence the hip extensor in the prone position. *Journal of Physical Therapy Science* 25:1295-1297.
- Lyons K, Perry J, Gronley JK, Barnes L, and Antonelli D. 1983. Timing and relative intensity of hip extensor and abductor muscle action during level and stair ambulation. An EMG study. *Physical Therapy* 63:1597-1605.
- McAndrew D, Gorelick M, and Brown J. 2006. Muscles within muscles: a mechanomyographic analysis of muscle segment contractile properties within human gluteus maximus. *Journal of Musculoskeletal Research* 10:23-35.
- McGill SM. 1990. Electromyographic activity of the abdominal and low back musculature during the generation of isometric and dynamic axial trunk torque: implications for lumbar mechanics. *Journal of Orthopaedic Research* 9:91-103.
- McGill SM, McDermott A, and Fenwick CM. 2009. Comparison of different strongman events: trunk muscle activation and lumbar spine motion, load, and stiffness. *Journal of strength and conditioning research / National Strength & Conditioning Association* 23:1148-1161.
- Morris SB. 2007. Estimating effect sizes from the pretest-posttest-control group designs. *Organizational Research Methods*.
- Neumann DA. 2010. Kinesiology of the hip: a focus on muscular actions. *Journal of Orthopaedic and Sports Physical Therapy* 40:82-94.
- Oh JS, Cynn HS, Won JH, Kwon OY, and Yi CH. 2007. Effects of performing an abdominal drawing-in maneuver during prone hip extension exercises on hip and back extensor muscle activity and amount of anterior pelvic tilt. *Journal of Orthopaedic and Sports Physical Therapy* 37:320-324.
- Simenz CJ, Garceau LR, Lutsch BN, Suchomel TJ, and Ebben WP. 2012. Electromyographical analysis of lower extremity muscle activation during variations of the loaded step-up exercise. *Journal of Strength and Conditioning Research* 26:3398-3405.
- Smith LJW, and Beretvas SN. 2009. Estimation of the Standardized Mean Difference for Repeated Measures Designs. *Journal of Modern Applied Statistical Methods* 8:27.
- Stensdotter AK, Hodges PW, Mellor R, Sundelin G, and Hager-Ross C. 2003. Quadriceps activation in closed and in open kinetic chain exercise. *Medicine and Science in Sports and Exercise* 35:2043-2047.
- Van Dillen LR, McDonnell MK, Fleming DA, and Sahrmann SA. 2000. Effect of knee and hip position on hip extension range of motion in individuals with and without low back pain. *Journal of Orthopaedic and Sports Physical Therapy* 30:307-316.
- Vera-Garcia FJ, Moreside JM, and McGill SM. 2010. MVC techniques to normalize trunk muscle EMG in healthy women. *Journal of Electromyography and Kinesiology* 20:10-16.
- Wheatley MD, and Jahnke WD. 1951. Electromyographic study of the superficial thigh and hip muscles in normal individuals. *Archives of Physical Medicine and Rehabilitation* 32:508-515.
- Worrell TW, Karst G, Adamczyk D, Moore R, Stanley C, Steimel B, and Steimel S. 2001. Influence of joint position on electromyographic and torque generation during maximal voluntary isometric contractions of the hamstrings and gluteus maximus muscles. *Journal of Orthopaedic and Sports Physical Therapy* 31:730-740.

327 Youdas JW, Foley BM, Kruger BL, Mangus JM, Tortorelli AM, Madson TJ, and Hollman JH.
 328 2013. Electromyographic analysis of trunk and hip muscles during resisted lateral
 329 band walking. *Physiotherapy Theory and Practice* 29:113-123.
 330