

Exploring the nitrogen reservoir of biodegradable household garbage and its potential in replacing synthetic nitrogen fertilizers in China

Lan Wang^{1,2}, Tianyu Qin^{1,2}, Jianshe Zhao³, Yicheng Zhang^{1,2}, Zhiyuan Wu^{1,2}, Xiaohui Cui^{1,2}, Gaifang Zhou^{1,2}, Caihong Li¹, Liyue Guo^{Corresp., 1}, Gaoming Jiang^{Corresp. 1, 2}

¹ State Key Laboratory of Vegetation and Environment Change, Institute of Botany, the Chinese 17 Academy of Sciences, Beijing, China

² College of Resources and Environment, University of Chinese Academy of Sciences, Beijing, China

³ Henan Zhongyuan Organic Agriculture Research Institute Co., Ltd., Zhengzhou, China

Corresponding Authors: Liyue Guo, Gaoming Jiang

Email address: guoliyue6789@126.com, jgm@ibcas.ac.cn

Biodegradable household garbage contains a large amount of nitrogen, which could be used as organic fertilizer to produce organic food and significantly reduce synthetic nitrogen fertilizers. There is limited information on how large the nitrogen reservoir of biodegradable household garbage is in a certain country or region. Here we took China as a case, analyzed the amount of biodegradable household garbage resources and their nitrogen reservoirs. It was noted that the biodegradable household garbage mainly included food waste, waste paper and wood chips, with the amount being 31.56, 29.55, and $6.45 \times 10^6 \text{ t} \cdot \text{a}^{-1}$, respectively. Accordingly, the nitrogen reservoirs were 65.31×10^4 , 6.80×10^4 , and $3.81 \times 10^4 \text{ t} \cdot \text{a}^{-1}$ in China. The nitrogen reservoir of food waste accounted for 86% of the total nitrogen reservoir of biodegradable household garbage, which was equivalent to 11% of the amount of actual absorption for synthetic nitrogen fertilizers ($6.20 \times 10^6 \text{ t} \cdot \text{a}^{-1}$) by agriculture plants in China. Our findings provided a scientific basis for the classification and utilization of biodegradable household garbage.

Exploring the nitrogen reservoir of biodegradable household garbage and its potential in replacing synthetic nitrogen fertilizers in China

Lan Wang^{1,2}, Tianyu Qin^{1,2}, Jianshe Zhao³, Yicheng Zhang^{1,2}, Zhiyuan Wu^{1,2}, Xiaohui Cui^{1,2}, Gaifang Zhou^{1,2}, Caihong Li¹, Liyue Guo¹*, Gaoming Jiang^{1,2}*

(1. State Key Laboratory of Vegetation and Environment Change, Institute of Botany, the Chinese Academy of Sciences, Beijing, 100093, China
2. College of Resources and Environment, University of Chinese Academy of Sciences, Beijing, 100049, China
3. Henan Zhongyuan Organic Agriculture Research Institute Co., Ltd., Zhengzhou, 450043, China)

*** Corresponding author:**

Gaoming Jiang, Tel.: +8610-6283-6286, fax: +8610-8259-6146. E-mail address: jgm@ibcas.ac.cn

Liyue Guo, Tel.: +8610-6283-6555, fax: +8610-8259-6146. E-mail address: guoliyue@ibcas.ac.cn

Abstract

Biodegradable household garbage contains a large amount of nitrogen, which could be used as organic fertilizer to produce organic food and significantly reduce synthetic nitrogen fertilizers. There is limited information on how large the nitrogen reservoir of biodegradable household garbage is in a certain country or region. Here we took China as a case, analyzed the amount of biodegradable household garbage resources and their nitrogen reservoirs. It was noted that the biodegradable household garbage mainly included food waste, waste paper and wood chips, with

the amount being 31.56, 29.55, and $6.45 \times 10^6 \text{ t} \cdot \text{a}^{-1}$, respectively. Accordingly, the nitrogen reservoirs were 65.31×10^4 , 6.80×10^4 , and $3.81 \times 10^4 \text{ t} \cdot \text{a}^{-1}$ in China. The nitrogen reservoir of food waste accounted for 86% of the total nitrogen reservoir of biodegradable household garbage, which was equivalent to 11% of the amount of actual absorption for synthetic nitrogen fertilizers ($6.20 \times 10^6 \text{ t} \cdot \text{a}^{-1}$) by agriculture plants in China. Our findings provided a scientific basis for the classification and utilization of biodegradable household garbage.

Keywords: Household garbage; Food waste; Biomass nitrogen reservoir; Agricultural sustainable development

1. Introduction

Modern agriculture is mainly characterized with large utilization of synthetic chemical substances such as fertilizers, pesticides, and herbicides. Although the crop yields have been increased by chemical substances, the long-term application of those chemicals has caused serious adverse effects on soil, air, water, food, even human health. Soil acidification (Guo et al. 2010; Sutton et al. 2011), greenhouse gas emission (Parihar et al. 2018), water eutrophication (Hansen et al. 2019) and pesticide residues (Urso and Gilbertson 2018) in agricultural system have been frequently reported. As a result, the quality of food, the nutritional contents (Yu et al. 2018), and the incomes of agriculture have been largely decreased (Stuart and Houser 2018). To a certain extent, cheaper food lead to more food waste. For instance, today, food mixed with garbage become the main components of modern urban and rural household garbages.

In the world, a large amount of household garbage is produced every day, which causes serious environmental pollution, and cost a lot of lands and money to handle (Du et al. 2018).

According to the statistics of the World Bank, the amount of household garbage generated worldwide reached to $2.01 \times 10^9 \text{ t a}^{-1}$ in 2016. In China, $4 \times 10^8 \text{ t a}^{-1}$ household garbages (fresh weight) have been produced, while the disposal rate was far behind its rate of generation (Wu et al. 2018).

The accumulation of household garbage has resulted in increasing environmental pollution, affecting the lives of residents and harming human health (Hiramatsu et al. 2009; Rao and Rathod 2019). Actually, the household garbage has an abundance of nutrients, including organic matter (39.05%), nitrogen (1.02%), phosphorus (0.50%) and potassium (1.42%) (Han et al. 2019). Nitrogen, an essential nutrient for plants growth, is the key element in the agricultural ecosystem (Sharma and Bali 2018), particularly in poor soils which need more exogenous nitrogen as an essential nutrient (Zeng et al. 2020). Globally, $1.50\text{-}2.00 \times 10^8 \text{ t a}^{-1}$ mineral nitrogen is required to produce grains, feed animals and be used as industrial products (Aulakh et al. 2017). Nowadays, the demand for nitrogen is mainly met by applying synthesized nitrogen fertilizers (Lu and Tian 2017). If the biodegradable household garbage was utilized as organic fertilizer, we would reduce the application amount of synthesized nitrogen at a certain level.

The household garbage comes from both urban and rural waste. Urban household garbage, known as municipal solid waste, includes food waste, recyclables, hazardous waste and others. However, the largest proportion in municipal solid waste is food waste, accounting to 61.2% of the total (Gu et al. 2017). The main components of rural household garbage in China include inert waste, food waste, glass and paper (Han et al. 2019). Regardless of urban or rural household garbage, biodegradable food waste accounts for a large proportion (Gallipoli et al. 2010), with

the main components being carbohydrate polymers (starch, cellulose, hemicellulose), proteins, organic acids, lignin, lipids, etc. Those biological materials can be decomposed into reducing sugars, free amino acids, phosphates and nitrates under the action of microbial hydrolysis which could be absorbed by agriculture plants (Xing et al. 2019). In addition, food waste has low content of salts and heavy metals, which might be directly used as organic fertilizer (Xiong 2015).

Among 94 countries in the world, China takes a leading position in the amount of food waste, followed by the United States and South Korea (Zhang et al. 2018). However, intensive researches have just focused on the technology innovation, management methods and bioenergy utilization potentials of food waste, in some countries such as the United States (Breunig et al. 2017), South Korea (Seo et al. 2019), and Ireland (Dennehy et al. 2017) , Turkey (Guven et al. 2019), *etc.* In many European countries, there were more reports on food waste composting technology (Ghinea et al. 2019), and food waste collection methods (Cecchi and Cavinato 2019). Although the UK had the most detailed, directly measured household food waste data, only the nutrients and energy content had been calculated (Cooper et al. 2018). Some investigators from Italy (Stathopoulos 2017; Giordano et al. 2019) and Canada (von Massow et al. 2019; Heidari et al. 2020) analyzed the main behaviors and attitudes based on the amount of food waste. Nevertheless, there was no statistics on the overall nitrogen reservoir. It is still not clear the resources and nitrogen reservoir of the biodegradable components of household garbage in the developing countries like China. The potential of replacing chemical fertilizers has been seldom reported, so the relevant scientific research is urgently needed.

The scientific hypothesis of this paper was that the biodegradable household garbage contained a large amount of nitrogen resources that could significantly replace the synthetic nitrogen fertilizers to develop organic agriculture. The treatment of biodegradable household garbage and the development of organic agriculture could be perfectly unified with the help of farmland. This study tried to use the statistical data to reveal: 1) The biodegradable components and resources in urban and rural household garbage together with the nitrogen reservoir in China; 2) The potential of biodegradable household garbage in replacing chemical fertilizers. This study might provide scientific basis for both the utilization of household garbage resources and the health development of organic agriculture.

2. Materials & Methods

2.1 Data Sources

The biodegradable household garbage was divided into urban and rural one. According to the composition and characteristics, the biodegradable garbage was further divided into three categories, *i.e.*, food waste, waste paper and wood chips (Li et al. 2019). Data were collected as previously described in Cui date (Cui *et al.* 2021). Specifically the original data mainly came from: China Statistical Yearbook (2011-2020) and China Statistical Yearbook on Urban and Rural Construction (2011-2020). The rest of related information was obtained by searching “household garbage”, “biodegradable household garbage”, and “food waste” through “Web of Science” and “Chinese National Knowledge Infrastructure (CNKI)”. Because of the special statistical system and the situation of China, all data reported in this paper were included except Hong Kong, Macau and Taiwan.

2.2 Food waste resources and the nitrogen reservoir

The total amount of food waste resources contained urban and rural one. The former was calculated by multiplying urban household garbage by the proportion of urban food waste in urban household garbage. The latter was calculated by multiplying the rural population by the average daily waste generation per capita, the number of days per year and the ratio of rural food waste to rural household garbage. The nitrogen reservoir of food waste was calculated by multiplying the total amount of food waste by its average nitrogen content.

$$TF = (UF + RF) \times (1 - w1) = (UG \times a1 + RG \times b1) \times (1 - w1) = (UG \times a1 + RP \times DG \times D \times b1) \times (1 - w1) \quad (1)$$

$$TN1 = TF \times \alpha1 \quad (2)$$

Where TF is the total output of urban and rural food waste each year; UF is the the output of urban food waste; RF is the output of rural food waste; UG is the output of urban household garbage; RG is the output of rural household garbage; RP is the number of rural population; DG is the per capita daily garbage production; a1 is the proportion of urban food waste in urban household garbage (61.20%) (Gu et al. 2017); b1 is the proportion of rural food waste in rural household garbage (33.7%) (Wu et al. 2018); D is the total days of the year; w1 is water content (82%) (Gallipoli et al. 2020); TN1 is the total urban and rural food waste nitrogen reservoir; $\alpha1$ is average nitrogen content of food waste (2.07%) (Adhikari et al. 2009; Yang et al. 2013; Zhang et al. 2016).

2.3 Waste paper resources and the nitrogen reservoir

The total amount of waste paper resources was composed of urban waste paper and rural

waste one. The former was calculated by multiplying urban household garbage by the proportion of urban waste paper in urban household garbage. The latter was calculated by multiplying the rural population by the average daily waste generation per capita, the number of days per year and the ratio of rural waste paper to rural household garbage. The nitrogen reservoir of waste paper was calculated by multiplying the total amount of waste paper by its average nitrogen content.

$$TP = (UP + RP) \times (1 - w_2) = (UG \times a_2 + RG \times b_2) \times (1 - w_2) = (UG \times a_2 + RP \times DG \times D \times b_2) \times (1 - w_2) \quad (3)$$

$$TN_2 = TP \times \alpha_2 \quad (4)$$

Where TP is the total output of urban and rural waste paper each year; UP is the the output of urban waste paper; RP is the output of rural waste paper; UG is the output of urban household garbage; RG is the output of rural household garbage; RP is the number of rural population; DG is the per capita daily garbage production; a_2 is the proportion of urban waste paper in urban household garbage (9.6%) (Gu et al. 2017); b_2 is the proportion of rural waste paper in rural household garbage (10.75%) (Wu et al. 2018); D is the total days of the year; w_2 is water content (7.35%) (Ding et al. 2013); TN_2 is the total urban and rural waste paper nitrogen reservoir; α_2 is average nitrogen content of waste paper (0.23%) (Ding et al. 2013).

2.4 Wood chip resources and the nitrogen reservoir

The total amount of wood chips resources was composed of urban wood chips and rural ones. The former was calculated by multiplying urban household garbage by the proportion of urban wood chips in urban household garbage. The latter was calculated by multiplying the rural

population by the average daily waste generation per capita, the number of days per year and the ratio of rural wood chips to rural household garbage. The nitrogen reservoir of wood chips was calculated by multiplying the total amount of wood chips by their average nitrogen content.

$$TW = (UW + RW) \times (1 - w_3) = (UG \times a_3 + RG \times b_3) \times (1 - w_3) = (UG \times a_3 + RP \times DG \times D \times b_3) \times (1 - w_3) \quad (5)$$

$$TN_3 = TW \times \alpha_3 \quad (6)$$

Where TW is the total output of urban and rural wood chips each year; UW is the the output of urban wood chips; RW is the output of rural wood chips; UG is the output of urban household garbage; RG is the output of rural household garbage; RP is the number of rural population; DG is the per capita daily garbage production; a_3 is the proportion of urban wood chips in urban household garbage (1.8%) (Gu et al. 2017); b_3 is the proportion of rural wood chips in rural household garbage (3.23%) (Wu et al. 2018); D is the total days of the year; w_3 is water content (7.24%) (Zhou et al. 2018); TN_3 is the total urban and rural wood chips nitrogen reservoir; α_3 is average nitrogen content of wood chips (0.59%) (Zhou et al. 2018).

2.5 Total biodegradable household garbage nitrogen reservoir

The total nitrogen reservoir of urban and rural biodegradable household garbage included urban and rural food waste nitrogen reservoir, waste paper nitrogen reservoir, and wood chips nitrogen reservoir, which was calculated as followings:

$$TN = TN_1 + TN_2 + TN_3 \quad (7)$$

TN is the total urban and rural biodegradable household garbage nitrogen reservoir; TN_1 is the total urban and rural food waste nitrogen reservoir; TN_2 is the total urban and rural waste

paper nitrogen reservoir; TN3 is the total urban and rural wood chips nitrogen reservoir.

2.6 Statistical analysis

Microsoft Excel 2007 was applied to process the data. SPSS 20.0 (SPSS Inc, Chicago, IL, USA) was applied to analyse the data. Figures were generated using SigmaPlot 12.5 (Systat Software Inc., San Jose, CA, USA).

3 Results

3.1 Food waste resource and its nitrogen reservoir

The total amount of urban and rural household garbage resources in China dated from 2010 to 2019 was shown in Table 1. According to formula (1), the amount of urban and rural food waste (dry weight) in China had been increasing year by year, reaching to $3.16 \times 10^7 \text{ t} \cdot \text{a}^{-1}$ in 2019. As well, the amount of urban food waste continued to grow, reaching to $2.67 \times 10^7 \text{ t} \cdot \text{a}^{-1}$ in 2019. Compared with 2010, the urban food waste increased by 53%, accounting 88% of the total food waste. However, as more and more rural people flooded into cities, the amount of rural food waste decreased, from $5.94 \times 10^6 \text{ t} \cdot \text{a}^{-1}$ in 2010 to $4.89 \times 10^6 \text{ t} \cdot \text{a}^{-1}$ in 2019 (Fig 1). According to formula (2), the nitrogen reservoir of food waste had been also increasing yearly, reaching to $6.51 \times 10^5 \text{ t} \cdot \text{a}^{-1}$ in 2019 (Fig 1).

3.2 Waste paper resource and its nitrogen reservoir

It was noted that the amount of urban waste paper resources elevated from $1.41 \times 10^7 \text{ t} \cdot \text{a}^{-1}$ in 2010 to $2.15 \times 10^7 \text{ t} \cdot \text{a}^{-1}$ in 2019 after calculated by formula (3). The amount of rural waste paper resource decreased from $9.76 \times 10^6 \text{ t} \cdot \text{a}^{-1}$ to $8.02 \times 10^6 \text{ t} \cdot \text{a}^{-1}$. The total amount of urban and rural waste paper resources increased year by year, reaching to $2.96 \times 10^7 \text{ t} \cdot \text{a}^{-1}$ in 2019. According to

formula (4), the nitrogen reservoir of waste paper increased with the increase of waste paper resource, reaching to $6.80 \times 10^4 \text{ t} \cdot \text{a}^{-1}$ in 2019 (Fig 2).

3.3 Wood chips resource and its nitrogen reservoir

From 2010 to 2019, the amount of urban wood chips increased from $2.64 \times 10^6 \text{ t} \cdot \text{a}^{-1}$ to $4.04 \times 10^6 \text{ t} \cdot \text{a}^{-1}$ after calculated by formula (5). The amount of rural wood chips decreased from $2.94 \times 10^6 \text{ t} \cdot \text{a}^{-1}$ to $2.41 \times 10^6 \text{ t} \cdot \text{a}^{-1}$. The total amount of wood chips in urban and rural increased yearly, reaching to $6.45 \times 10^6 \text{ t} \cdot \text{a}^{-1}$ in 2019. According to formula (6), the wood chip nitrogen reservoir increased to $3.81 \times 10^4 \text{ t} \cdot \text{a}^{-1}$ in 2019 (Fig 3).

3.4 The total nitrogen reservoir of biodegradable household garbage and its components

It was found that the total nitrogen reservoir of urban and rural biodegradable household garbage in China had been increasing since 2010. In 2019, the nitrogen reservoir was $75.92 \times 10^4 \text{ t} \cdot \text{a}^{-1}$, increased by 33% compared with that of 2010 (Fig 4). For the components, urban and rural biodegradable household garbage were composed of food waste, waste paper and wood chips, among which food waste accounted for 84%-86%, waste paper 8%-9% and wood chips 5%-6%. The proportion of food waste had been increasing, while the proportion of waste paper and wood chips had been decreasing from 2010 to 2019 (Fig 4).

3.5 Yearly rates of changes in biodegradable household garbage resources

From 2010 to 2019, the yearly changes of urban biodegradable household garbage resources had increased while that of rural had decreased in China (Table 2). From 2014 to 2015, the yearly rate of increase in urban biodegradable household garbage resources was the maximum. From 2016 to 2017, however, the yearly rate of decrease in rural household garbage was the

maximum.

3.6 Per capita biodegradable household garbage resources in seven regions of China

In 2019, the amount of urban and rural food waste, waste paper, wood chips and the per capita biodegradable household garbage resources in seven regions of China were shown in Table 3. The largest amount of biodegradable household garbage and the per capita biodegradable household garbage were from east and southern China, respectively, with the least being from northwest China.

3.7 The relationship between per capital GDP and per capital discharge of food waste

Among the biodegradable household garbage, food waste accounted for the largest proportion, which was the most promising renewable resource as organic fertilizer. In 2019, the amount of urban and rural food waste resources in different provinces of China were shown in Table 4 (Provincial scale). The top five provinces with the largest amount of food waste were from Guangdong, Jiangsu, Shandong, Zhejiang and Henan. As one of most developed provinces in China, Guangdong had the maximum food waste discharge of $3.98 \times 10^6 \text{ t} \cdot \text{a}^{-1}$, accounting for 13% of the total food waste of the country.

There was a pretty significant linear positive correlation between the per capital GDP and per capital discharge of food waste ($P < 0.01$) (Fig 5). The per capital GDP values of different provinces were found to be mostly relevant with per capital food waste. The more GDP increased, the more foods were wasted, indicating that the fast development of economy was based on the huge waste of food resources which further led to environmental pollutions.

4. Discussion

4.1 Components and treatments of urban and rural degradable household garbage

Economic development, urbanization, and the living standards improvement of human being have led to a sharp increase in the amount of household garbage discharge, especially in developed countries (Breunig et al. 2017). To a certain extent, the discharge of household garbage is restricted by socio-economic conditions (Huang et al. 2013) and geographical locations (Han et al. 2015). Unfortunately, the developing countries have been following the footsteps of developed countries, with their garbage waste being increased rapidly (Rai et al. 2019). Most of cities in the world have struggled to “Garbage siege”. Unlike urban residents, rural residents have maintained a high utilization rate of those materials (Ma et al. 2018). In 2019, the total amount of urban and rural household garbage reached to $3.23 \times 10^8 \text{ t} \cdot \text{a}^{-1}$ in China (Table 1).

At present, the methods for treating urban household garbage mainly include incineration, landfill, composting, *etc.* (Zhang, 2019). In the rural China, however, there is still a shortage of appropriate infrastructure and solid waste management (Hiramatsu et al. 2009). Most rural household garbages are randomly discarded without any treatments, resulting in increase of environmental pollution and damage of human health (Cao et al. 2018).

Implementing household garbage classification which follows the principles of reduction, recycling, and harmlessness, is believed to be an effective approach to improve the urban and rural environments and promote resource recycling (Shi et al. 2020). However, it achieved limited effect, despite of enough mobilization has been done by the government, as urban and rural residents always believe that garbage disposal is the government's business not theirs.

According to the compositions and characteristics of household garbage, urban and rural degradable household garbages are divided into food waste, waste paper and wood chips (Li et al. 2019). We here found these resources in China were respectively 31.56 , 29.55 and 6.45×10^6 $t \cdot a^{-1}$. In 2019, the total amount of the biodegradable household garbage was 67.4×10^6 $t \cdot a^{-1}$ in China (Table 3). The urban biodegradable household garbage was on the rise over time (Table 2).

Urban and rural wood chips, as a kind of biomass, can be used as adsorbents for treating waste water, which reduces waste water treatment costs while increases environmental benefits (Li et al. 2010). Waste wood chips could be also converted into gaseous or liquid fuel, chemical raw materials and other products through thermochemical, chemical, and biological methods (Xu et al. 2018). Waste paper, however, is an ideal recyclable renewable resource that can be used for paper-making, wood production, or terated as various functional materials (Liu 2016). In order to increase the utilization rate of paper, those waste papers are normally reused, even recycled, which could reduce the amount of felling of trees and obtain more ecological benefits (Liu 2018). Therefore, the utilization of waste wood chips and waste paper follows a mode of circular economy. The application of such waste in the environmental protection industry should be continuously more strengthened rather than be used as fertilizers.

Food waste is the main component of urban and rural household garbage, which is also one of the most promising renewable resources (Cecchi and Cavinato 2019). In the U.S. wastes about 63×10^6 $t \cdot a^{-1}$ of food wasted (Dusoruth and Peterson 2020), South Korea watstes about 5×10^6 $t \cdot a^{-1}$ food (Lee et al. 2019), in Europe approximately 10.0×10^7 $t \cdot a^{-1}$ of food are wasted (Zhang et

al. 2018). Food waste reached to $3.16 \times 10^7 \text{ t} \cdot \text{a}^{-1}$ in 2019 in China (dry weight). Much of the food waste is a result of urbanization, for instance, urban food waste in China was as large as $2.67 \times 10^7 \text{ t} \cdot \text{a}^{-1}$, while rural food waste was only $4.89 \times 10^6 \text{ t} \cdot \text{a}^{-1}$ in the year of 2019 (Fig 1). Food waste can be treated by landfill, incineration and composting (Han et al. 2015). However, landfill is likely to bring about greenhouse gases emissions, generation of large amount of leachate, and finally environmental pollution (Ma and Liu 2019). Although incineration could reduce the volume of food waste, it needs a high demand for energy and might produce harmful substances and greenhouse gasses (Liu et al. 2019).

Aerobic composting is a relatively environmentally friend technology for food waste treatment, as foods contain high concentrations of easily degradable organic substances and nutrients which are easily to be decomposed (Hou et al. 2017). Studies have found that food waste composted with Chinese medicinal herbal residues (Zhou et al. 2018), green waste (Williams et al. 2019), sugarcane leaves (Shan et al. 2019), cattle manure (Xing et al. 2019), pig manure (Dennehy et al. 2017; Jiang et al. 2018), chicken manure (Saad et al. 2019) could be processed to form biological fertilizers, which would increase organic matters in the soil and improve the soil structure (Wang and Zeng 2018). Therefore, the degradable food waste, especially rural food waste could be used on-site as fertilizers for organic crop production, which can largely reduce transportation and treatment costs and increase farmers' income as well.

4.2 Nitrogen reservoir of biodegradable household garbage in different components

In China, food waste accounted for the largest proportion (84%-86%) of the nitrogen reservoir of biodegradable household garbage, followed by waste paper and wood chips. Along

with time, the proportion of food waste increased, while waste paper and wood chips declined (Fig 4). In 2019, we found that the nitrogen reservoir of food waste, waste paper and wood chips in China were 65.31×10^4 , 6.80×10^4 , and $3.81 \times 10^4 \text{ t} \cdot \text{a}^{-1}$, respectively.

The nitrogen in food waste was mainly organic one, which was found in various molecular forms, such as protein, amino acid, and nucleic acid, *etc.* (Wang and Zeng 2018). It was reported that the protein content of food waste was 20% (Waqas et al. 2019) and NH_4^+-N was 2800 mg kg^{-1} (Rigby and Smith 2013). Nitrate is the main form of nitrogen absorption and utilization by most crops in cultivated soils (Andrews et al. 2013). So, as a source of biomass nitrogen, food waste can be used in organic agriculture to improve soil quality and promote agricultural development.

4.3 Potential of biodegradable household garbage of replacing synthetic nitrogen fertilizers

Among the biodegradable household garbage, wood chips and waste paper could be recycled, which play more important roles in the environmental protection industry rather than used as fertilizers. Therefore, this article did not consider them as organic fertilizers. During fermentation process, various components in food waste are converted into stable humus-like substances and rapidly available nutrients, which can be quickly hydrolyzed (Bi et al. 2019), or directly absorbed by plants (Hou et al. 2017). Those materials could also improve nutrition levels of the soil (Du et al. 2018) and could be generally beneficial to soil microbial communities (Liu et al. 2020) which are closely related to soil fertility. Soil microorganisms provide a variety of services for agriculture ecosystem, such as cycling of nutrient elements, degradation of pesticides, suppression of plant diseases, and promotion of plant growth (Ding et al. 2019). In

food waste, the dissolved organic matter is very active, which directly provides energy for microbes (Shan et al. 2019). Therefore, food waste is regarded as an ideal and cheap raw material for the production of biological fertilizers (Ma and Liu 2019). It was documented that the rapid humification of food waste prepared as a soil conditioner could significantly improve the total organic carbon content in orchard soils (Jia et al. 2019). Some also found that food waste culture medium could replace inorganic culture medium as a nutrient supplement to cultivate chlorella and improve nutrient utilization efficiency (Chew et al. 2018). The organic fertilizer processed by food waste and substrate in different ratios could promote the growth of potted vegetables pepper (*Capsicum annuum*) and cabbage (*Brassica pekinensis*) (Li et al. 2020). In addition, some investigators who directly applied food waste to potting soil found that food waste promoted leaf growth of *Chlorophytum comosum*, and increased soil available nitrogen, phosphorus and potassium (Song et al. 2014). Therefore, the ability of food waste in producing organic foods suggested here might be an alternative and effective way of treating biodegradable household garbage in the future.

Just recently, our team had proposed that biomass resources originally produced by photosynthesis and their biodegradable derivatives such as biodegradable household garbage, can be all used as organic fertilizers which have huge potentials in replacing chemical ones (Cui et al. 2021). Under the action of microorganism, the organic matter in the food waste will be rapidly decomposed into substances that can be easily absorbed by agriculture plants, thus partly replacing the usage of chemical fertilizers. In China, the actual amount of chemically synthesized nitrogen fertilizer absorbed by agricultural plants nationwide was $6.20 \times 10^6 \text{ t} \cdot \text{a}^{-1}$. Food waste, if

applied as organic fertilizers, could replace 11% of chemical nitrogen application. Especially, the rural food waste when simply stacked, processed and returned directly to the farmland, could save processing and transportation costs. This will not only fully make use of household garbage, but also make up for the lack of organic fertilizers in the development of organic farming, so as to ensure food security.

4.4 The relationship between GDP and food waste

Food waste occurs at all stages of the supply chain which is affected by many factors, such as geography and economy, production systems, infrastructure, markets, and consumption (Bonadonna et al. 2019). Larger consumer market and more consumption input undoubtedly exacerbate the food waste (Di Talia et al. 2019).

Our study displayed that there was a very significant linear positive correlation between the per capital GDP and per capital food waste production (Fig 5). GDP is mainly based on the classification of cities based on the concentration of commercial resources, diversity of lifestyles, future plasticity, urban hubs, and urban occupant activity index in China (Wang 2018). Our findings demonstrated that the GDP and food waste generation of economically developed provinces were much higher than those of underdeveloped provinces (Fig 5). And the largest amount of per capita biodegradable household garbage was in the southern China of the seven regions of the country (Table 3). This might be owing to its own economic conditions and geographical location. Southern China has been mostly economically developed, with the amount of waste generated remaining the highest over the years (Tian et al., 2018). This situation is consistent with that of the United States (Breunig et al. 2017). Food waste contains biomass

nitrogen which could be used again for food production after simple treatment. If well paid, farmers might be actively mobilized and engaged in organic farming, so as to implement waste sorting and utilization at the source, reduce the load of rural waste entering to cities and cut down waste disposal costs.

4.5 Questions and suggestions

So far, most investigations on food waste treatments usually focus on energy recovery. However, they do not consider the economic feasibility of such an approach (Ma and Liu 2019). Food waste can be easily collected from various sources such as food processing industries, households, and hospitality sectors (Paritosh et al. 2017; Sindhu et al. 2019). Nevertheless, food waste may contain some inert materials, such as glass or plastic, and the distribution is somewhat difficult.

Although the prices of organic food are much higher and farmers are willing to invest their labors to use food waste in organic farming, correct collection, storage, and distribution are major obstacles of food waste management. To overcome those shortages, we here suggest: 1) The government should improve the household garbage collection and classification system in rural areas, and encourage farmers to use biodegradable household garbage in organic farming. By doing so, almost 80% of the degradable garbage in China could used as organic fertilizers to produce organic foods, thus uniting the ecological and economic chains together; 2) In cities, the government should encourage companies to build special food waste treatment plants to solve the problem of destination of urban household garbage by market-based means; 3) The government should supplement waste disposal subsidies to enterprises or farmers who produce

organic food using the degraded food waste, if they have actually reduced the amount of biodegradable waste.

5. Conclusions

The total amount of urban and rural biodegradable household garbage in China was $6.76 \times 10^7 \text{ t} \cdot \text{a}^{-1}$, with the nitrogen reservoir being $7.59 \times 10^5 \text{ t} \cdot \text{a}^{-1}$. The nitrogen reservoir of food waste potentially used as organic fertilizers reached to $6.53 \times 10^5 \text{ t} \cdot \text{a}^{-1}$, being equivalent to 11% of the amount of actual absorption for synthetic nitrogen fertilizers ($6.20 \times 10^6 \text{ t} \cdot \text{a}^{-1}$) by agriculture plants of the country. There was a significant correlation between per capital GDP and the per capital food waste. Food waste from household garbage should be classified and processed at the source, economically used as organic fertilizers to replace chemical ones. Our innovative solution might realize the recycling of biodegradable waste and the sustainable development of agriculture, ensure both food security and environmental protection.

Acknowledgement

We thank all the staffs of Hongyi Organic Farm for providing accommodation and the field facilities, especially Mr. Jiang Gaoliang and Mr. Jiang Shenglin. We also thank all the researchers who given us help during the study.

References

- Adhikari BK, Barrington S, Martinez J, King S. 2009. Effectiveness of three bulking agents for food waste composting. *Waste Management* 29:197-203.
DOI: 10.1016/j.wasman.2008.04.001
- Andrews M, Raven JA, Lea PJ. 2013. Do plants need nitrate? The mechanisms by which nitrogen form affects plants. *Annals of Applied Biology* 163:174-199.

DOI: 10.1111/aab.12045

Aulakh MS, Garg AK, Manchanda JS, Dercon G, Nguyen ML. 2017. Biological nitrogen fixation by soybean and fate of applied N¹⁵-fertilizer in succeeding wheat under conventional tillage and conservation agriculture practices. *Nutrient Cycling in Agroecosystems* 107:79-89. DOI: 10.1007/s10705-016-9816-8

Bi SJ, Hong XJ, Bai Y, Liu JL, Yu XH, Fang SM, Gao YM, Yan L, Wang ZH, Wang YJ, Wang WD. 2019. Methane production dynamics of co-digestion of cow manure and food waste under mesophilic condition. *Journal of Biobased Materials and Bioenergy* 13:257-263. DOI: 10.1166/jbmb.2019.1846

Bonadonna A, Matozzo A, Giachino C, Peira G. 2019. Farmer behavior and perception regarding food waste and unsold food. *British Food Journal* 121:89-103. DOI:10.1108/bfj-12-2017-0727

Breunig HM, Jin L, Robinson A, Scown CD. 2017. Bioenergy potential from food waste in California. *Environmental Science & Technology* 51:1120-1128. DOI: 10.1021/acs.est.6b04591

Cao S, Xu DD, Liu SQ. 2018. A study of the relationships between the characteristics of the village population structure and rural residential solid waste collection services: Evidence from China. *International Journal of Environmental Research and Public Health* 15:17. DOI: 10.3390/ijerph15112352

Cecchi F, Cavinato C. 2019. Smart approaches to food waste final disposal. *International Journal of Environmental Research and Public Health* 16:13. DOI: 10.3390/ijerph16162860

Chen HB, Jiang W, Yang Y, Yang Y, Man X. 2017. State of the art on food waste research: a bibliometrics study from 1997 to 2014. *Journal of Cleaner Production* 140:840-846. DOI:10.1016/j.jclepro.2015.11.085

- 426 Chew KW, Chia SR, Show PL, Ling TC, Arya SS, Chang JS. 2018. Food waste compost as an
427 organic nutrient source for the cultivation of *Chlorella vulgaris*. *Bioresource Technology*
428 267:356-362. DOI: 10.1016/j.biortech.2018.07.069
- 429 Cooper KA, Quested TE, Lanctuit H, Zimmermann D, Espinoza-Orias N, Roulin A. 2018.
430 Nutrition in the bin: A nutritional and environmental assessment of food wasted in the
431 UK. *Frontiers in Nutrition* 5:19 doi:10.3389/fnut.2018.00019
- 432 Cui XH, Guo LY, Li CH, Liu MZ, Wu GL, Jiang GM. 2021. The total biomass nitrogen
433 reservoir and its potential of replacing chemical fertilizers in China. *Renewable &*
434 *Sustainable Energy Reviews* 135:110215. DOI: 10.1016/j.rser.2020.110215
- 435 Dennehy C, Lawlor PG, Gardiner GE, Jiang Y, Shalloo L, Zhan X. 2017. Stochastic modelling
436 of the economic viability of on-farm co-digestion of pig manure and food waste in Ireland.
437 *Applied Energy* 205:1528-1537. DOI: 10.1016/j.apenergy.2017.08.101
- 438 Di Talia E, Simeone M, Scarpato D. 2019. Consumer behaviour types in household food waste.
439 *Journal of Cleaner Production* 214:166-172. DOI:10.1016/j.jclepro.2018.12.216
- 440 Ding GC, Bai MH, Han H, Li HX, Ding XY, Yang HF, Xu T, Li J. 2019. Microbial taxonomic,
441 nitrogen cycling and phosphorus recycling community composition during long-term
442 organic greenhouse farming. *FEMS Microbiology Ecology* 95:12.
443 DOI: 10.1093/femsec/fiz042
- 444 Ding K, Zhong ZP, Yu LL, Liu ZC. 2013. Pyrolysis characteristics and kinetic study of mixed
445 pyrolysis of municipal solid waste. *Journal of Southeast University* 43:130-135. DOI:
446 10.3969/j.issn.1001-0505.2013.01.025 (in Chinese)
- 447 Du CY, Abdullah JJ, Greetham D, Fu, DN, Yu MY, Ren LW, Li S, Lu DN. 2018. Valorization

of food waste into biofertiliser and its field application. *Journal of Cleaner Production* 187:273-284. DOI: 10.1016/j.jclepro.2018.03.211

Dusoruth V, Peterson HH. 2020. Food waste tendencies: Behavioral response to cosmetic deterioration of food. *PLoS One* 15:22. DOI:10.1371/journal.pone.02332870

Gallipoli A, Braguglia CM, Gianico A, Montecchio D, Pagliaccia P. 2020. Kitchen waste valorization through a mildtemperature pretreatment to enhance biogas production and fermentability: Kinetics study in mesophilic and thermophilic regimen. *Journal of Environmental Sciences* 89:167-179. DOI:10.1016/j.jes.2019.10.016

Ghinea C, Apostol LC, Prisacaru AE, Leahu A. 2019. Development of a model for food waste composting. *Environmental Science and Pollution Research* 26:4056-4069 doi:10.1007/s11356-018-3939-1

Giordano C, Alboni F, Falasconi L. 2019. Quantities, determinants, and awareness of households' food waste in Italy: A comparison between diary and questionnaires. *Quantities Sustainability* 11:15 doi:10.3390/su11123381

Gu BX, Jiang SQ, Wang HK, Wang ZB, Jia RF, Yang J, He S, Cheng R. 2017. Characterization, quantification and management of China's municipal solid waste in spatiotemporal distributions: A review. *Waste Management* 61:67-77. DOI:10.1016/j.wasman.2016.11.039

Guo JH, Liu XJ, Zhang Y, Shen JL, Han WX, Zhang WF, Christie P, Goulding KWT, Vitousek PM, Zhang FS. 2010. Significant acidification in major Chinese croplands. *Science* 327:1008-1010. DOI:10.1126/science.1182570

- 469 Guven H, Wang Z, Eriksson O. 2019. Evaluation of future food waste management alternatives
470 in Istanbul from the life cycle assessment perspective. *Journal of Cleaner Production*
471 239:12 doi:10.1016/j.jclepro.2019.117999
- 472 Han ZY, Dan Z, Shi GZ, Shen LK, Xu WL, Xie YH. 2015. Characteristics and management of
473 domestic waste in a rural area of the Tibetan Plateau. *Journal of the Air & Waste*
474 *Management Association* 65:1365-1375. DOI:10.1080/10962247.2015.1078859
- 475 Han ZY, Ye CW, Zhang Y, Dan Z, Zou ZY, Liu D, Shi GZ. 2019. Characteristics and
476 management modes of domestic waste in rural areas of developing countries: a case study
477 of China. *Environmental Science and Pollution Research* 26:8485-8501.
478 DOI:10.1007/s11356-019-04289-w
- 479 Hansen LB, Termansen M, Hasler B. 2019. The potential for nitrogen abatement trading in
480 agriculture: A hypothetical market experiment. *Journal of Agricultural Economics* 70:812-
481 839. DOI:10.1111/1477-9552.12319
- 482 Heidari A, Mirzaii F, Rahn timer M, Alidoost F. 2020. A theoretical framework for explaining the
483 determinants of food waste reduction in residential households: a case study of Mashhad,
484 Iran. *Environmental Science and Pollution Research* 27:6774-6784 doi:10.1007/s11356-
485 019-06518-8
- 486 Hiramatsu A, Hara Y, Sekiyama M, Honda R, Chiemchaisri C. 2009. Municipal solid waste flow
487 and waste generation characteristics in an urban-rural fringe area in Thailand. *Waste*
488 *Management & Research* 27:951-960. DOI:10.1177/0734242x09103819
- 489 Hou JQ, Li MX, Xi BD, Tan WB, Ding J, Hao Y, Liu DM, Liu HL. 2017. Short-duration

hydrothermal fermentation of food waste: preparation of soil conditioner for amending
organic-matter-impoverished arable soils. *Environmental Science and Pollution Research*
24:21283-21297. DOI:10.1007/s11356-017-9514-3

Huang KX, Wang JX, Bai JF, Qiu HG. 2013. Domestic solid waste discharge and its
determinants in rural China. *China Agricultural Economic Review* 5:512-525.
DOI:10.1108/caer-02-2012-0008

Jia X, Zhao B, Ren LH, Li MX, Zhang XH, Hou JQ, He L. 2019. Effects of food waste soil
conditioner on organic carbon distribution in orchard soil. *Research of Environmental*
Sciences 32:485-492. DOI: 10. 13198/j.issn.1001-6929.2018.06.23 (in Chinese)

Jiang Y, Dennehy C, Lawlor PG, Hu ZH, Yang QF, McCarthy G, Tan SP, Zhan XM, Gardiner
GE. 2018. Inactivation of Salmonella during dry co-digestion of food waste and pig for
manure. *Waste Management* 82:231-240. DOI:10.1016/j.wasman.2018.10.037

Lee, CH, Park SJ, Hwang H.Y, Kim MS, Jung, HI, Luyima D, Hong SY, Oh TK, Kim SH.
2019. Effects of food waste compost on the shift of microbial community in water saturated
and unsaturated soil condition. *Applied Biological Chemistry* 62:7 doi:10.1186/s13765-019-
0445-1

Li C. 2010. Application of native botanical materials in electroplating wastewater treatment.
Plating & Finishing 32:25-30 (in Chinese)

Li SF, Xu Z, Zhou HP, Wei GQ. 2020. A comparative study on the utilization of two kinds of
organic wastes in vegetable planting. *Vegetables* 10:8-13 (in Chinese)

Li XR, Bi F, Han ZD, Qin Y, Wang HS, Wu WX. 2019. Garbage source classification

performance, impact factor, and management strategy in rural areas of China: A case study
in Hangzhou. *Waste Management* 89:313-321. DOI:10.1016/j.wasman.2019.04.020

Liu HY. 2016. Recycling and sorting of waste paper. *Heilongjiang Pulp & Paper* 44:21-25 (in
Chinese)

Liu J, Zhang J, Li DM, Xu CX, Xiang XJ. 2020. Differential responses of arbuscular
mycorrhizal fungal communities to mineral and organic fertilization. *Microbiology Open*
9:10. DOI: 10.1002/mbo3.920

Liu JH, Li Q, Gu W, Wang C. 2019. The impact of consumption patterns on the generation of
municipal solid waste in China: Evidences from provincial data. *International Journal of*
Environmental Research and Public Health 19:1-19. DOI:10.3390/ijerph16101717

Liu SH. 2020. Chinese food chinese rice bowl. *Grain Science and Technology and Economy*
45:9-11 (in Chinese)

Liu XX. 2018. Turning waste into treasure-A discussion on the reuse technology of waste paper.
China Resources Comprehensive Utilization 36:67-69. DOI:1008-9500(2018)11-0067-03
(in Chinese)

Lu CQ, Tian HQ. 2017. Global nitrogen and phosphorus fertilizer use for agriculture production
in the past half century: shifted hot spots and nutrient imbalance. *Earth System Science*
Data 9:181-192. DOI:10.5194/essd-9-181-2017

Ma J, Hipel KW, Hanson ML. 2018. An evaluation of the social dimensions in public
participation in rural domestic waste source-separated collection in Guilin, China.
Environmental Monitoring and Assessment 190:14. DOI:10.1007/s10661-017-6405-5

Ma Y, Liu Y. 2019. Turning food waste to energy and resources towards a great environmental and economic sustainability: An innovative integrated biological approach. *Biotechnology Advances* 37:107414. DOI:10.1016/j.biotechadv.2019.06.013

Ministry of Housing and Urban-Rural Development of the People's Republic of China. China Statistical Yearbook on urban and rural construction 2011-2020 (in Chinese). Available online. 2011-2020. <http://www.mohurd.gov.cn/xytj/tjzljsxytjgb/jstjnj/index.html>.

National Bureau of Statistics of the People's Republic of China (NBS). China statistical Yearbook in 2020 (in Chinese). 2020. Available online, <http://www.stats.gov.cn/tjsj/ndsj/2019/indexch.htm>.

Parihar CM, Parihar MD, Sapkota TB, Nanwal RK, Singh AK, Jat SL, Nayak HS, Mahala DM, Singh LK, Kakraliya SK, Stirling CM, Jat ML. 2018. Long-term impact of conservation agriculture and diversified maize rotations on carbon pools and stocks, mineral nitrogen fractions and nitrous oxide fluxes in inceptisol of India. *Science of the Total Environment* 640:1382-1392. DOI:10.1016/j.scitotenv.2018.05.405

Paritosh K, Kushwaha SK, Yadav M, Pareek N, Chawade A, Vivekanand V. 2017. Food waste to energy: An overview of sustainable approaches for food waste management and nutrient recycling. *Biomed Research International*:19. DOI:10.1155/2017/2370927

Rao P, Rathod V. 2019. Valorization of food and agricultural waste: a step towards greener future. *Chemical Record* 19:1858-1871. DOI:10.1002/tcr.201800094

Rigby H, Smith SR. 2013. Nitrogen availability and indirect measurements of greenhouse gas

emissions from aerobic and anaerobic biowaste digestates applied to agricultural soils. Waste Management 33:2641-2652. DOI: 10.1016/j.wasman.2013.08.005

Saad MFM, Rahman NA, Yusoff MZM. 2019. Hydrogen and methane production from co-digestion of food waste and chicken manure. Polish Journal Of Environmental Studies 28:2805-2814. DOI:10.15244/pjoes/83670

Seo H, Cho K, Shin J, Lee M, Park J, Lee BC, Song KG. 2019. Linking process performances and core microbial community structures in anaerobic membrane bioreactor with rotatory disk (ARMBR) system fed with high-strength food waste recycling wastewater. Bioresource Technology 291:11 doi:10.1016/j.biortech.2019.121918

Shan GC, Xu JQ, Jiang ZW, Li MQ, Li QL. 2019. The transformation of different dissolved organic matter subfractions and distribution of heavy metals during food waste and sugarcane leaves co-composting. Waste Management 87:636-644. DOI:10.1016/j.wasman.2019.03.005

Sharma LK, Bali SK. 2018. A review of methods to improve nitrogen use efficiency in agriculture. Sustainability 10:23. DOI:10.3390/su10010051

Shi XX, Zheng GD, Shao ZZ, Gao D. 2020. Effect of source-classified and mixed collection from residential household waste bins on the emission characteristics of volatile organic compounds. Science of the Total Environment 707:9. DOI:10.1016/j.scitotenv.2019.135478

Sindhu R, Gnansounou E, Rebello S, Binod P, Varjani S, Thakur IS, Nair RB, Pandey A. 2019. Conversion of food and kitchen waste to value-added products. Journal of Environmental Management 241:619-630. DOI:10.1016/j.jenvman.2019.02.053

574 Song B, Nie LH, Yuan QN, Zhang X, Zhang GQ, Gao PX, Zhu JL. 2014. Effects of several
575 common kitchen wastes on the growth of *Chlorophytum comosum*. *Journal of Agriculture*
576 4:56-59 (in Chinese)

577 Stathopoulos CE. 2017. Utilisation of plant food. *Waste Foods* (Basel, Switzerland) 6
578 doi:10.3390/foods6060045

579 Stuart D, Houser M. 2018. Producing Compliant Polluters: Seed companies and nitrogen
580 fertilizer application in US corn agriculture. *Rural Sociology* 83:857-881.
581 DOI:10.1111/ruso.12212.

582 Sutton MA, Oenema O, Erisman JW, Leip A, van Grinsven H, Winiwarter W. 2011. Too much
583 of a good thing. *Nature* 472:159-161. DOI:10.1038/472159a

584 The World Bank. *Solid Waste Management* 2018.

585 Tian GJ, Kong LQ, Liu XJ, Yuan WP. 2018. The spatio-temporal dynamic pattern of rural
586 domestic solid waste discharge of China and its challenges. *Environmental Science and*
587 *Pollution Research* 25:10115-10125. DOI:10.1007/s11356-017-1154-0

588 Urso JH, Gilbertson LM. 2018. Atom Conversion Efficiency: A new sustainability metric
589 applied to nitrogen and phosphorus use in agriculture *ACS sustain. Chemical Engineering*
590 *Journal* 6:4453-4463. DOI:10.1021/acssuschemeng.7b03600

591 von Massow M, Parizeau K, Gallant M, Wickson M, Haines J, Ma DWL, Wallace A, Carroll N,
592 Duncan AM. 2019. Valuing the multiple impacts of household food. *Waste Frontiers in*
593 *Nutrition* 6:17 doi:10.3389/fnut.2019.00143

594 Wang JH. 2018. Research on the efficiency of urban green innovation and its influencing factors.

Xianan:Shaanxi Normal University (in Chinese)

Wang LE, Hou P, Liu X J, Cheng SK. 2018. The connotation and realization way of sustainable food consumption in China. *Resources Science* 40(8):1550-1559. DOI:10.18402/resci.2018.08.06

Wang SG, Zeng Y. 2018. Ammonia emission mitigation in food waste composting: A review. *Bioresource Technology* 248:13-19. DOI:10.1016/j.biortech.2017.07.050

Wang TX. The food wasted can feed 300 million people every year in the country. *China News Network*. <http://www.chinanews.com/gn/news/2010/03-10/2161052.shtml>. 2010-03-10.

Waqas M, Nizami AS, Aburiazaiza AS, Barakat MA, Asam ZZ, Khattak B, Rashid MI. 2019. Untapped potential of zeolites in optimization of food waste composting. *Journal of Environmental Management* 241:99-112. DOI:10.1016/j.jenvman.2019.04.014

Williams SR, Zhu-Barker X, Lew S, Croze BJ, Fallan KR, Horwath WR. 2019. Impact of composting food waste with green waste on greenhouse gas emissions from compost windrows. *Compost Science & Utilization* 27:35-45. DOI:10.1080/1065657x.2018.1550023

Wu XH, Yue B, Huang QF, Wang Q, Li ZL, Wang YT, Yu JY. 2018. Investigation of the physical and chemical characteristics of rural solid waste in China and its spatiotemporal distributions. *Environmental Science and Pollution Research* 25:17330-17342. DOI:10.1007/s11356-018-1432-5

Xing BS, Han YL, Wang XC, Ma Jing, Cao SF, Li Q, Wen JW, Yuan HL. 2019. Cow manure as additive to a DMBR for stable and high-rate digestion of food waste: Performance and

microbial community. *Water research* 168:115099. DOI:10.1016/j.watres.2019.115099

Xiong C. 2015. Experimental study on organic fertilizer production from kitchen waste by hydrothermal treatment. Hangzhou: Zhejiang University (in Chinese)

Xu HB, Hu JH, Liu HL, Li A, Jin LF, Liu JL. 2018. Experiment and thermodynamic analysis of the sawdust catalytic gasification with copper slag. *Chemical Industry and Engineering Progress* 35:3142-3148. DOI:10.16085/j.issn.1000-6613.2016.10.018 (in Chinese)

Yang F, Li GX, Yang QY, Luo WH. 2013. Effect of bulking agents on maturity and gaseous emissions during kitchen waste composting. *Chemosphere* 93:1393-1399. DOI:10.1016/j.chemosphere.2013.07.002

Yu XF, Guo LY, Jiang GM, Song YJ, Muminov MA. 2018. Advances of organic products over conventional productions with respect to nutritional quality and food security. *Acta Ecologica Sinica* 38:53–60. DOI:10.1016/j.chnaes.2018.01.009

Zeng J, Yue Fj, Li SL, Wang ZJ, Qin CQ, Wu QX, Xu S. 2020. Agriculture driven nitrogen wet deposition in a karst catchment in southwest China. *Agriculture, Ecosystems and Environment* 294:10. DOI: 10.1016/j.agee.2020.106883

Zhang HY, Li GX, Gu J, Wang GQ, Li YY, Zhang DF. 2016. Influence of aeration on volatile sulfur compounds (VSCs) and NH₃ emissions during aerobic composting of kitchen waste. *Waste Management* 58:369-375. DOI:10.1016/j.wasman.2016.08.022

Zhang LF. 2019. Research on the management of domestic waste disposal in Xiong'an. Shijiazhuang: Hebei University (in Chinese)

Zhang M, Gao M, Yue SY, Zheng TL, Gao Z, Ma XY, Wang QH. 2018. Global trends and

637 future prospects of food waste research: a bibliometric analysis. Environmental Science
 638 and Pollution Research 25:24600-24610 doi:10.1007/s11356-018-2598-6
 639 Zhang YF. 2019. Research on the management of domestic waste disposal in Xiong'an new
 640 distric (in Chinese). Baoding: Hebei University (in Chinese)
 641 Zhou Y, Selvam A, Wong JWC. 2018. Chinese medicinal herbal residues as a bulking agent for
 642 food waste composting. Bioresource Technology 249:182-188.
 643 DOI:10.1016/j.biortech.2017.09.212

Figure 1

The biomass and nitrogen reservoir of urban and rural food waste in China from 2010 to 2019




Figure 2

The biomass and nitrogen reservoir of urban and rural waste paper in China from 2010 to 2019




Figure 3

The biomass and nitrogen reservoir of urban and rural wood chips in China from 2010 to 2019




Figure 4

Changes of nitrogen reservoir and the proportion of biodegradable household garbage from 2010 to 2019 in China




Figure 5

The relationship between per capita GDP and per capita discharge of food waste




Table 1 (on next page)

Urban and rural household garbage resources in China from 2010 to 2019 (fresh weight)

Table 1 Urban and rural household garbage resources in China from 2010 to 2019 (fresh weight).

Household garbage		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Urban	Production (10 ⁴ t)	15438	15734	15805	16395	17081	17239	17860	19142	20362	21521
Rural	Population (10 ⁴)	67113	65656	64222	62961	61866	60346	58973	57661	56401	55162
	Per capita daily (kg/d)	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
	Days	365	365	366	365	365	365	366	365	365	365
	Production (10 ⁴ t)	9798	9586	9402	9192	9032	8811	8634	8419	8235	8054
Total	Production (10 ⁴ t)	25603	25981	26483	26431	26892	27953	28996	29940	31037	32260

Table 2(on next page)

Urban increase and rural decrease yearly rates of biodegradable household garbage resources in China from 2010 to 2019

Table 2 Urban increase and rural decrease yearly rates of biodegradable household garbage resources in China from 2010 to 2019

Year □	Biodegradable household garbage	
	Urban yearly increase rates (%)	Rural yearly decrease rates (%)
2010-2011	3.73	2.17
2011-2012	4.18	1.92
2012-2013	0.93	2.23
2013-2014	3.60	1.74
2014-2015	7.18	2.46
2015-2016	6.37	2.01
2016-2017	5.69	2.49
2017-2018	5.95	2.19
2018-2019	6.16	2.20

Table 3(on next page)

Urban and rural per capita biodegradable household garbage resources in seven regions of China in 2019

Table 3 Urban and rural per capita biodegradable household garbage resources in seven regions of China in 2019

		North	Northeast	East	Central	Southern	Southwest	Northwest
Urban	Food waste (10 ⁴ t)	331.42	219.45	884.92	318.36	451.82	292.54	165.08
	Waste paper (10 ⁴ t)	267.59	177.18	714.49	257.05	364.80	236.20	133.29
	Wood chips (10 ⁴ t)	50.23	33.26	134.13	48.25	68.48	44.34	25.02
Rural	Food waste (10 ⁴ t)	54.94	35.23	127.69	86.63	54.08	83.92	41.62
	Waste paper (10 ⁴ t)	90.21	57.85	209.66	142.24	88.79	137.79	68.33
	Wood chips (10 ⁴ t)	27.14	17.40	63.07	42.79	26.71	41.45	20.56
Biodegradable household								
Total	garbage (10 ⁴ t)	821.53	540.37	2133.95	895.34	1054.68	836.24	453.89
	Population (10 ⁴)	17577	10794	41423	22485	17426	20331	10349
	Per capita (kg a ⁻¹)	46.74	50.06	51.52	39.82	60.52	41.13	43.86

Table 4(on next page)

Gross domestic product (GDP) and food waste resources in 2019 in China

Table 4

Gross domestic product (GDP) and food waste resources in 2019 in China

Province	Population (10 ⁴)	GDP (billion, \$)	Urban and rural food waste production (10 ⁴ t)	Per capita GDP(\$)	Per capita food waste(kg)
Guangdong	11521	1648.74	397.92	14310.74	34.54
Jiangsu	8070	1525.63	220.35	18904.96	27.30
Shandong	10070	1088.24	231.16	10806.75	22.96
Zhejiang	5850	954.78	184.11	16321.03	31.47
Henan	9640	830.86	164.94	8618.88	17.11
Sichuan	8375	713.82	163.01	8523.22	19.46
Hubei	5927	701.76	128.43	11840.05	21.67
Fujian	3973	649.18	118.32	16339.79	29.78
Hunan	6918	608.71	111.63	8798.93	16.14
Shanghai	2428	584.26	85.21	24063.43	35.09
Anhui	6366	568.32	96.09	8927.43	15.09
Beijing	2154	541.63	113.95	25145.31	52.90
Hebei	7592	537.55	116.87	7080.48	15.39
Shaanxi	3876	394.96	83.76	10189.89	21.61
Liaoning	4352	381.43	120.84	8764.48	27.77
Jiangxi	4666	379.11	77.37	8124.95	16.58
Chongqing	3124	361.47	75.47	11570.74	24.16
Yunnan	4858	355.62	72.20	7320.30	14.86
Guangxi	4960	325.20	76.31	6556.45	15.39
Inner Mongolia	2540	263.57	51.70	10376.77	20.35
Shanxi	3729	260.73	68.48	6991.95	18.36
Guizhou	3623	256.78	56.52	7087.50	15.60
Tianjin	1562	215.98	35.36	13827.14	22.64
Heilongjiang	3751	208.45	70.67	5557.18	18.84
Xinjiang	2523	208.21	48.75	8252.48	19.32
Jilin	2691	179.57	63.16	6672.98	23.47
Gansu	2647	133.50	42.88	5043.45	16.20
Hainan	945	81.29	31.66	8602.12	33.50
Ningxia	695	57.40	16.85	8258.99	24.24
Qinghai	608	45.42	14.45	7470.39	23.77
Tibet	351	26.00	9.25	7407.41	26.35

Note: \$1.0=6.5305 Chinese Yuan.