

Unraveling the hidden diversity of the Middle Devonian (Emsian) crinoids (Crinoidea, Echinodermata) from Poland (#68648)

1

First submission

Guidance from your Editor

Please submit by **28 Dec 2021** for the benefit of the authors (and your \$200 publishing discount) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

! Custom checks

8 Figure file(s)

New species checks



Have you checked our [new species policies](#)?



Do you agree that it is a new species?



Is it correctly described e.g. meets ICZN standard?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Unraveling the hidden diversity of the Middle Devonian (Emsian) crinoids (Crinoidea, Echinodermata) from Poland

William Ausich¹, Mariusz A Salamon^{Corresp., 2}, Bartosz Płachno³, Tomasz Brachaniec², Wojciech Krawczyński², Andrzej Boczarowski², Karolina Paszcza², Magdalena Łukowiak⁴, Przemysław Gorzelak⁵

¹ University of Columbus, Columbus, USA

² University of Silesia in Katowice, Sosnowiec, Poland

³ Jagiellonian University Cracow, Kraków, Poland

⁴ Insitute of Paleobiology, Warszawa, Poland

⁵ Insitute of Paleobiology Polish Academy of Science, Warszawa, Poland

Corresponding Author: Mariusz A Salamon

Email address: paleo.crinoids@poczta.fm

Most previous publications on Devonian crinoids from the Holy Cross Mountains in Poland have concentrated on crinoid columns, and until now, little has been published about crinoid cups and calyxes. Herein, five crinoid taxa are described from an abundant occurrence of aboral cups and partial crowns from the Bukowa Góra Member (Emsian) in the Holy Cross Mountains of southern Poland. The following taxa are described: *Bactrocrinites* sp., *Codiocrinus sevastopuloi* n. sp., *Halocrinites geminatus* (Bohatý, 2005), *Halocrinites schlotheimii* Steininger, 1831, and a single brachial plate from a flexible crinoid placed in *Flexibilia incertae sedis*. Simple discoid holdfasts are also present encrusted to cylindrical stromatoporoids. These taxa are the first crinoids described from the remains of partial crowns from Emsian strata of Poland.

Unraveling the hidden diversity of the Middle Devonian (Emsian) crinoids (Crinoidea, Echinodermata) from Poland

William I. Ausich¹, Mariusz A. Salamon^{2*}, Bartosz J. Płachno³, Tomasz Brachaniec², Wojciech Krawczyński², Andrzej Boczarowski², Karolina Paszcza², Magdalena Łukowiak⁴ and Przemysław Gorzelak⁴

¹School of Earth Sciences, Ohio State University, 125 South Oval Mall, Columbus, Ohio, U.S.A.

^{2*}University of Silesia in Katowice, Faculty of Natural Sciences, Institute of Earth Sciences, Będzińska Street 60, 41-200 Sosnowiec, Poland, e-mail: paleo.crinoids@poczta.fm

³Jagiellonian University in Kraków, Faculty of Biology, Institute of Botany, Gronostajowa Street 9, 30-387 Cracow, Poland

⁴Institute of Paleobiology, Polish Academy of Sciences, Twarda 51/55, 00-818 Warsaw, Poland

Corresponding Author:

Mariusz A. Salamon²

²University of Silesia in Katowice, Faculty of Natural Sciences, Institute of Earth Sciences, Będzińska Street 60, 41-200 Sosnowiec, Poland

e-mail: paleo.crinoids@poczta.fm

ABSTRACT

Most previous publications on Devonian crinoids from the Holy Cross Mountains in Poland have concentrated on crinoid columns, and until now, little has been published about crinoid cups and calyxes. Herein, five crinoid taxa are described from an abundant occurrence of aboral cups and partial crowns from the Bukowa Góra Member (Emsian) in the Holy Cross Mountains of southern Poland. The following taxa are described: *Bactrocrinites* sp., *Codiocrinus sevastopuloi* sp. nov., *Halocrinites geminatus* (Bohatý, 2005), *Halocrinites schlotheimii* Steininger, 1831, and a single brachial plate from a flexible crinoid placed in *Flexibilia incertae sedis*. Simple discoid holdfasts are also present encrusted to cylindrical stromatoporoids. These taxa are the first crinoids described from the remains of partial crowns from Emsian strata of Poland.

Subjects Paleontology

Keywords Devonian, Emsian, Poland, crinoids, Echinodermata, diversity.

INTRODUCTION

Crinoid remains are abundant in Devonian (Emsian-Famennian) strata of Poland (Holy Cross Mountains, southern Poland; Cracow-Silesian area, southern Poland; Sudetes, southwestern Poland; Pomerania, northern Poland). Polish Devonian crinoids were mentioned initially by Dames (1868), Zeuschner (1867, 1869), Gürich (1896), and Sobolev (1909). Much later, Kongiel

(1958) and Piotrowski (1977) described the occurrence of the genus *Ammonocrinus* in the Holy Cross Mountains (see [Gorzelak et al., 2014](#)) a series of subsequent papers, Głuchowski (1980, 1981a-c, 1993a, 1993b, 2002, 2003; see also Hauser, 2002) listed ~50 crinoid taxa from Devonian (Lochkovian, Emsian-Famennian) strata of Poland (for summary see Fig. 1). Unfortunately, most of them were based on isolated skeletal remains, mainly columnals; and they were described using the principles of artificial classification of crinoid remains proposed by Moore and Jeffords (1968). Głuchowski (2003) added that the applicability of crinoid stems may be useful for stratigraphic and correlation purposes.

To date, only eight crinoid taxa have been identified on the basis of complete (or nearly complete) crowns and aboral cups with column from the Devonian of Poland. Among these is *Haplocrinites* sp. from Givetian-Frasnian of Holy Cross Mountains recorded by Głuchowski (1993a, 2003). Specimens of this taxon from late Frasnian Detrital Beds of the Holy Cross Mountains were later designated by Hauser (2002) as *Haplocrinites gluchowskii* Hauser, 2002. According to Głuchowski et al. (2006) Givetian and early Frasnian *Haplocrinites* sp. specimens differ from *H. gluchowskii* in having distinctly less prominent radial facets. Another haplocrinitid (Haplocrinitidae) species is *Haplocrinites aremoricensis* Le Menn, 1985 from the uppermost Givetian of the Holy Cross Mountains (Głuchowski, 1993a). Also the cupressocrinitid (Cupressocrinitidae) taxa have been described from the Holy Cross Mountains by Głuchowski (1993a). These are *Cupressocrinites* cf. *abbreviatus* Goldfuss (late Eifelian-late Givetian) (now *Halocrinites schlotheimii* Steininger, 1831), *C. inflatus* Schultze, 1866 (late Givetian) (now *Halocrinites inflatus*), and *C. sampelayoi* (Almela and Revilla, 1950) (now *Halocrinites minor* (Schultze, 1866) known from late Givetian (Głuchowski 1993a, 2002). The remaining two crinoid species belong to the Melocrinitidae. These are *Melocrinites* cf. *gibbosus* Goldfuss, 1831

and *M. cf. hieroglyphicus* Goldfuss, 1831, which were found in the sediments of the uppermost Frasnian.

Only four crinoid taxa are known from the Emsian of the Holy Cross Mountains and all were documented on the basis of isolated columnals or their impressions (casts). In particular, Głuchowski (1981b, 2002) listed the following taxa from the Bukowa Góra shales: *Acanthocrinus* sp. (col.), *Formosocrinus* cf. *formosus* (col.) (Yeltyscheva and Sisova, 1973), *Laudonomphalus humilicarinatus* (col.) (Yeltyscheva in Yeltyscheva and Dubatolova, 1961) [now *Hexacrinus*? *humilicarinatus* (col.)], and *Marettocrinus subbiconcavus* (col.) (Stukalina, 1965).

Here we report complete or almost complete Emsian cups associated with numerous isolated calyx and column remains from the Bukowa Góra Member in the Holy Cross Mountains of southern Poland. These include *Bactrocrinites* sp., *Codiocrinus sevastopuloi* sp. nov., *Halocrinites geminatus* (Bohatý, 2005), *Halocrinites schlotheimii* Steininger, 1831, and *Flexibilia incertae sedis*. Simple discoid holdfasts are also present. Remains of unidentifiable specimens indicate that several other crinoids also existed in the Bukowa Góra Member fauna.

Figure 1 around here

GEOLOGIC FRAMEWORK

The Holy Cross Mountains are located in the southern part of Poland. Their main element is the Paleozoic core, divided into two parts: the Łysogóry region (northern, connected with the Łysogóry Block) and the Kielce region (southern, connected with the Małopolska Block; see Fig. 2A). These regions differ from each other by facies development of contemporaneous deposits. Devonian sediments of the Łysogóry region were formed in the deeper basin in contrast to the shallower facies exposed in the Kielce region (Szulczewski, 1995).

Outcrops of Lower Devonian rocks in the Łysogóry Region are connected with the southern limb of the Bodzentyn Syncline. The sedimentary rocks of the upper Emsian are best exposed in the active quarry "Bukowa Góra", located about 16 km northeast of Kielce (see Fig. 2A). The section includes sediments belonging to *patulus* Conodont Zone and *douglastownense-euryptero* Miospore Zone (Malec, 2005; Filipiak, 2011; Fijałkowska-Madej & Malec, 2011).

In the lower part of the section, the 110 m thick Zagórze "formation" comprised of siliciclastic deposits is present (see Fig. 2B). They are mostly represented by quartzitic sandstones with abundant trace fossils and by claystones. Within sandstones, there are storm originated brachiopod coquinas with gastropods, bivalves, tentaculitids, crinoids, rarely trilobites, rugose corals, nautiloids, and ostracodes. At the top of the Zagórze "formation", conglomerates and sandstones of estuary facies crop out. Deposits of the Zagórze "formation" were formed in the shallow sea environment from the lagoon to the shoreface (Szulczewski & Porębski, 2008; see also Łobanowski, 1971, 1981).

As a result of the progressive deepening of the marine basin, deposits of the Grzegorzowice Formation were formed, which is also present in the Kielce region (Malec, 2005; Wójcik, 2015). In the Bukowa Góra quarry section the two members are present: the Bukowa Góra Member and the Kapkazy Member. The Bukowa Góra Member occurs only within the

patulus Conodont Zone in the western part of the Bodzentyn Syncline (see Fig. 2B). In the eastern part of the Bodzentyn Syncline, the Bukowa Góra Member appears earlier, i.e., in the *serotinus* Conodont Zone (Malec, 2005). Malec (2005) marked the lower boundary of the Bukowa Góra Member in the bottom of the complex of dark claystones, whereas Szulczewski & Porębski (2008) put this boundary on the pebble conglomerate that begins of the lower shoreface to offshore transitional series.

In the lower part of interbedded sandstones, the Bukowa Góra Member is comprised of sandstones, siltstones, and claystones, which are about 7 m thick (see Szulczewski & Porębski, 2008). Above them appears the offshore facies represented by black to dark-gray claystones and silty claystones with a thickness of ~13 m and containing discontinuous beds of dolomitic limestones up to 10 cm thick (see Fig. 2B). Both claystones and limestones contain a rich faunal assemblage related to the colonization of the soft sea bottom. There are massive colonies of both stromatoporoids and tabulate corals accompanied by solitary rugoses, brachiopods, crinoids, ostracods, gastropods and trilobites (Malec, 2005; see also Głuchowski, 1993b; Fijałkowska-Mader et al., 1997).

Claystones of the Bukowa Góra Member are overlain by sandstones of the Kapkazy Member, which is ~34 m thick. The lower part of the Kapkazy Member is comprised of coarse-grained and conglomeratic sandstones, containing rare crinoids, brachiopods and gastropods. Above this is fine-grained sandstone, which is indicative of a clear shallowing of the sea basin (Malec 2005).

Figure 2 around here

130

131 MATERIALS AND METHODS

132

133 The studied material from Bukowa Góra Quarry was collected in 2019 and 2021. The first step
134 consisted of examination of slab surfaces in the field. At this stage, numerous crinoid remains
135 (isolated columnals and complete or nearly complete crowns) were collected. The next step
136 consisted of soaking the respective samples (11 shales samples weighing each ca. 10 kg) only
137 with hot water. Limy samples (4 samples weighing each ca. 5 kg) were soaked with Glauber's
138 salt. These samples were then boiled and frozen (2-3 times). The residues were finally washed
139 with running tap water and sieved on a sieve column ($\varnothing$ 1.0, 0.315 and 0.1 mm mesh). The final
140 step consisted of drying the shaly and limy residues at 160°C. Residue was hand-picked from
141 each macerated sample for microscopic study.

142 All crinoids were photographed by a SONY DSC-RX10M3 digital camera. Specimens
143 discussed here are deposited in the University of Silesia in Katowice, Faculty of Natural
144 Sciences, Institute of Earth Sciences, Poland GIUS 4-3696) and in ~~a private collection~~
145 ~~(Nr. CREF; see Boháč 2005).~~

146 “The electronic version of this article in Portable Document Format (PDF) will represent
147 a published work according to the International Commission on Zoological Nomenclature
148 (ICZN), and hence the new names contained in the electronic version are effectively published
149 under that Code from the electronic edition alone. This published work and the nomenclatural
150 acts it contains have been registered in ZooBank, the online registration system for the ICZN.
151 The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information
152 viewed through any standard web browser by appending the LSID to the prefix

http://zoobank.org/. The LSID for this publication is: zoobank.org:pub:B89FD16E-2084-431A-ACE9-4E4362C6C3CD, and for taxonomic registration it is: urn:lsid:zoobank.org:act:66DBF909-CF1C-479C-BDA0-F324F4FFC15F. The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central SCIE and CLOCKSS”.

RESULTS

More than 1,000 of columnals and pluricolumnals, dozens of disarticulated ossicles from cups and arms, and 26 complete (or nearly complete) cups/calices were collected. As a result of our investigations, the following taxa were identified: *Bactrocrinites* sp., *Codiocrinus sevastopuloi* sp. nov., *Halocrinites geminatus* (Bohatý, 2005), *Halocrinites schlotheimii* Steininger, 1831, and *Flexibilia incertae sedis*. Simple discoid holdfasts are also described.

SYSTEMATIC PALEONTOLOGY

Abbreviations used for specimen measurements include ACH, aboral cup height; ACdistW, distal width of aboral cup; ACmaxW, maximum width of aboral cup; ACproxW, proximal width of aboral cup; BConW, basal concavity width; BH, basal plate height; BW, basal plate maximum width; CrW, crown width; 1stPBH, first primibrachial height; 1stBrW, first primibrachial width; 2ndPBH, second primibrachial height; 2ndPBdistW, second primibrachial distal width; 2ndPBproxW, second primibrachial proximal width; 5thPBH, fourth primibrachial height; 3rdPBW, third primibrachial width; 3rdPBH, third primibrachial height; 5thSBW, fifth primibrachial width.

All measurements are in mm. Terminology for encrusting organisms follows the recommendations of Taylor and Wilson (2002).

Class Crinoidea Miller, 1821

Subclass Pentacrinoidea Jaekel, 1918

Infraclass Inadunata Wachsmuth and Springer, 1885

Parvclass Cladida Moore and Laudon, 1943

Magnorder Euclidida Wright, 2017

Superorder Cyathiformes Wright et al., 2017

Superfamily Codiacrinoidea Bather, 1890

Family Codiocrinidae Bather, 1890

Subfamily Codiocrininae Bather, 1890

Genus *Codiocrinus* Schultze, 1866

Type species

Codiocrinus granulatus Schultze, 1866

Included species

193 *C. granulatus* Schultze, 1866; *C. nicolli* Jell and Jell, 1999; *C. ornatus* (Prokop, 1973); *C.*
 194 *piriformis* LeMenn, 1997; *C. procerus* (Prokop, 1973); *P. rarus* Jell and Holloway, 1983; *C.*
 195 *robustus* LeMenn, 1997; *C. schultzei* Follman, 1887; *C. secundus* Jell, 1999.

196

197 **Figure 3 around here**

198

199 *Codiocrinus sevastopuloi* sp. nov.

200 Fig. 3A1-A4, 3B1-B3, 3C

201 **Diagnosis**

202 Aboral cup medium bowl shape; three or more radiating ridges from center of basal plates that
 203 project onto radial and infrabasal plates, also very fine nodose sculpturing across calyx plates;
 204 basal plates largest plates of aboral cup; radial facets ~50% of distal width of radial plates
 205 (arms and column characters not known).

206 **Types**

207 Holotype: GIUS4-3693/Codiocrinus1; paratypes: GIUS4-3693/Codiocrinus2, GIUS4-
 208 3693/Codiocrinus3.

209 **Occurrence**

210 Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland.

211 **Description**

Aboral cup medium globe shaped (Fig. 3A4, 3B1), height to width ratio ~1.0, maximum width at middle aboral cup height; three or more radiating ridges from center of basal plates that project onto radial and infrabasal plates (Fig. 3A1), also very fine nodose sculpturing across calyx plates. Infrabasal circlet ~9% of aboral cup height, extends proximally in a short neck that is truncate proximally with a shallow, circular basal concavity that occupies ~75% of proximal aboral cup width (Fig. 4A2, 4B2). Five pentagonal infrabasal plates, ~3.8 times wider than high, outer surface concave, sculpturing irregular nodose. Basal circlet ~55% of aboral cup height; basal plates largest plates in aboral cup, hexagonal, ~1.2 times higher than wide; sculpturing with radiating ridges and nodes, ridges from near the center of the plates to ridges on adjoining proximal and distal plates. Radial circlet ~36% of aboral cup height; radial plates ~1.2 times wider than high, pentagonal; plate sculpturing with ridges and nodes, ridges diagonal from base of radial facet to like ridges on adjoining basal plates. Radial facets angustary (~52% of radial plate distal width), horseshoe shaped (Fig. 4A3, 4B3). Radial facets, arms, and column unknown.

Etymology

The species name is in recognition of the substantial contributions that George D. Sevastopulo made to crinoid paleobiology, as well as paleontology and stratigraphy in general.

Measurements

GIUS4-3693/Codiocrinus1 (holotype): ACH, 9.8; ACmaxW, 10.4; IH, 1.5; IW, 2.5; BH, 5.4; BW, 7.6; RH, 5.2; RmaxW, 6.0, RdistW, 5.3, RFW, 3.1. GIUS4-3693/Codiocrinus2 (paratype): ACH, 11.25; ACmaxW, 11.25*; IH, 1.4; IW, 5.3; BH, 8.4; BW, 6.8; RH, 5.4; RmaxW, 6.6, RdistW, 4.8, RFW, 2.5.

Remarks

One well-preserved and two poorly preserved aboral cups are assigned to *Codiocrinus* *sevastopuloi* sp. nov. Both poorly preserved specimens have their shapes distorted through compaction.

Ten species of *Codiocrinus*, including *C. sevastopuloi*, are recognized herein. *C.?* *weyeri* is excluded, and it is regarded either as an aberrant individual or a member of another genus. Of these ten species, only three have arms and proximal columnals preserved (*C. robustus*, *C. schultzei*, and *C. secundus*). Thus, species diagnoses are largely based on characters of the aboral cup, which vary widely.

The most noticeable character used to differentiate species of *Codiocrinus* is the aboral cup shape, which may be low bowl, high cone, medium globe, high globe, or medium vase in shape. *Codiocrinus granulatus*, *C. nicolli*, *C. robustus*, and *Codiocrinus sevastopuloi* sp. nov. all have a medium globe-shaped aboral cup. *Codiocrinus granulatus* has a medium globe-shaped aboral cup, two poorly defined radiating ridges from the base of the radial facet onto each subjacent basal plate and perhaps some poorly developed concentric ridges, radial plates are the largest plates in the aboral cup, and the radial facets occupy ~50% of the distal radial plate width. *Codiocrinus nicolli* has a medium globe-shaped aboral cup, very fine nodose sculpturing, radial plates are the largest plates in the aboral cup, and the radial facets occupy ~60% of the distal radial plate width. *Codiocrinus robustus* has a medium globe-shaped aboral cup, three radiating ridges from the basal plate center and otherwise smooth sculpturing, basal plates are the largest plates in the aboral cup, and the radial facets occupy ~50% of the distal radial plate width.

Alternatively, *Codiocrinus sevastopuloi* sp. nov. has a medium globe-shaped aboral cup, three or more radiating ridges from center of basal plates that project onto radial and infrabasal

plates, also very fine nodose sculpturing across calyx plates, basal plates are the largest plates in the aboral cup, and the radial facets occupy ~50% of the distal radial plate width.

Figure 4 around here

Superfamily Dendrocrinacea Wachsmuth and Springer, 1886

Family Dendrocrinidae Wachsmuth and Springer, 1886

Genus *Bactrocrinites* Schnur in Steininger, 1849

Type species

Poteriocrinus fusiformis (Roemer, 1844).

Included species

Bactrocrinites birmanicus Reed, 1908; *B. cyathus* Schmidt, 1942; *B. depressus* (Schultze, 1866); *B. fieldi* (Springer and Solcom, 1906); *B. fusiformis* (Roemer, 1844); *B. holstenfelsensis* Hauser, 2007; *B. jaekeli* (Schmidt, 1934); *B. muelleri* (Jaekel, 1895); *B. oklahomaensis* Strimple, 1952; *B. onondagensis* Goldring, 1954; *B. penaneachensis* LeMenn, 1985; *B. porectus* Bohatý, 2005; *B. tenuis* (Jaekel, 1895); *B.? trabicus* (Schmidt, 1934); *B. zeileri* (Mueller in Zeiler and Wirtgen, 1855)

Bactrocrinites sp.

Fig. 3E, 4

Description

Relatively large aboral cup, aboral cup plates with pustulose plate sculpturing (Fig. 3E). Infrabasal plates not known. Basal plates partially preserved, inferred to be the dominant plate circlet in aboral cup (Fig. 3E). C radial plate supported beneath by radianal plate and BC basal plate; D radial plate larger than C radial plate, supported beneath by CD and DE basal plates. Radial facets large, semicircular, angustate, declivate. Two anal plates in aboral cup. Radianal presumably tetragonal, below and to the left of the C radial plate and supports the anal X plate on the upper left (Fig. 4). Anal X plate hexagonal, supported beneath by the CD basal plate and the radianal, separates and articulates with lateral sides of the C and D radial plates.

Other aspects of the aboral cup, anal sac, arms, and column are not known.

Occurrence

Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland.

Remarks

Species diagnostic characters for Devonian *Bactrocrinites* include shape of the aboral cup, plate sculpturing, relative heights of aboral cup plates, and the dimensions of the infrabasal and basal plates. Unfortunately, the single specimen of *Bactrocrinites* from the Emsian of Poland is not complete (GIUS4-3693/*Bactrocrinites*), so aboral cup shape, relative proportions of aboral cup plates and the dimensions of the basal plates cannot be determined. The pustulose aboral cup plate sculpturing and what are inferred to be prominent (high) basal plates most closely ally this

specimen with *B. fusiformis*. However, a more complete accounting of the morphology of this Emsian specimen is required before a confident species assignment can be made.

Figure 5 and 6 around here

Superfamily Gasterocomoidea Roemer, 1854

Family Cupressocrinitidae Roemer, 1854

Subfamily Cupressocrinitinae Bohatý 2006

Genus *Halocrinites* Steininger, 1831

Type species

Halocrinites schlotheimi schlotheimi Steininger, 1831.

Included species

Halocrinites altus (Schultze, 1866); *H. assimilis* (Dubatolova, 1964); *H. geminatus* (Bohatý, 2005); *H. gibber* (Bather, 1919); *H. heinorum* Bohatý and Ausich (2021); *H. inflatus inflatus* (Schultze, 1866); *H. inflatus convexus* (Hauser, 2001); *H. inflatus cuneatus* Bohatý, 2006; *H. inflatus depressus* (Hauser, 2001); *H. minor* (Schultze, 1866); *H. nodosus* (Sandberger and Sandberger, 1856); *H. rectangularis* (Schmidt, 1941); *H. schlotheimii schlotheimii* Steininger, 1831; *H. schlotheimii granulosus* Schultze, 1866; *H. schreueri* Bohatý, 2006; *H. tesserula* (Goldfuss, 1831); *H. townsendi* (König, 1825); and *H. urogali* Roemer, 1850.

317 Remarks

318 As discussed in Bohatý and Ausich (2021), generic and specific assignments of the
 319 Cupressocrinitidae have been varied, commonly changed, and confused until recently (e.g.,
 320 Bohatý, 2005; Bohatý, 2006; Bohatý, 2009; Bohatý and Herbig, 2010; and Bohatý and Ausich,
 321 2021). Two species of *Halocrinites* are recognized from the Bukowa Góra Member in Poland,
 322 including *H. geminatus* (Bohatý, 2005) and *H. schlotheimii* (Steininger, 1831). Most
 323 *Halocrinites* specimens from Poland are lacking the exoplacoid layer or are sufficiently worn
 324 that the character of the exoplacoid layer cannot be determined. One exception is specimen
 325 GIUS 4-3696Hscholth6 (Fig. 5E), although even this specimen is worn. They are differentiated
 326 on the basis of aboral cup shape, basal plate morphology, size of the infrabasal circlet relative to
 327 the size of the proximalmost columnal, and the size of the basal concavity, as described below.
 328 *Halocrinites schlotheimii* has a bowl-shaped aboral cup with a ratio of aboral cup diameter
 329 versus crown height $\sim 1:1.15\text{--}2.0$; aboral cup typically ~ 2.0 times wider than high; infrabasal
 330 plates fused into a single pentagonal plate that is confined to the basal concavity; brachials wider
 331 than high (height to width ratio $\sim 1:2.0\text{--}2.5$); proximal columnal circular not filling entire basal
 332 concavity. In contrast, *H. geminatus* aboral cup bowl to moderately conical in shape; ratio of
 333 aboral cup diameter versus crown height $\sim 1:1.15\text{--}2.0$; typically 2.0 times wider than high;
 334 infrabasal plates fused into a single pentalobate plate that is confined to the basal concavity;
 335 brachials wider than high (height to width ratio $\sim 1:2.0\text{--}2.5$); proximal columnal circular not
 336 filling entire basal concavity.

337 Similar to the Cupressocrinitidae described  Bohatý and Ausich (2021), *Halocrinites*
 338 from Poland have a variety of epizoans encrusting the outer surface of crown plates. These
 339 include trepostome bryozoans encrusted on aboral cup and brachial plates. A presumable

microconchid that is attached to a radial plate, and a juvenile pelmatozoan holdfast is attached to a different radial plate. These encrustations did not induce a recognizable response from the crinoid host, so it is probable that these encrustations occurred after the death of the crinoid and, thus, are episkeletozoans (see Taylor and Wilson, 2000 

Halocrinites schlotheimii Steininger, 1831

Figs. 3F1–F3, 5A1–A3, 5B1–B2, 5C1–C2, 53, 6A1, 6A2

Type

The type specimens for this taxon are not known.

Diagnosis

Halocrinites with bowl-shaped aboral cup with a ratio of aboral cup diameter versus aboral cup height $\sim 1:1.15\text{--}2.0$; aboral cup typically ~ 2.0 times wider than high; infrabasal plates fused into a single pentagonal plate that is confined to the basal concavity; brachials wider than high (height to width ratio $\sim 1:2.0\text{--}2.5$); proximal columnal circular not filling entire basal concavity.

Occurrence

In Poland, *H. schlotheimii* is from the Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, Poland. Previously, this species has been described from the Eifelian and Givetian of Australia, China, Germany, Poland, and Spain (Webster and Webster, 2019).

Description

Calyx medium sized. Aboral cup low to very low bowl shape in adults with height to maximum width ratio 0.44–0.66 (Fig. 3F); perfect pentameral symmetry; deep, subpentalobate basal concavity occupies 62–66% of proximal aboral cup width (Fig. 5E). Plates gently convex; coarse multilaminar exoplacoid sculpturing preserved on only a few specimens.

Infrabasal circlet completely in basal concavity pentagonal. Infrabasal plates presumably five. Basal circlet ~58–60% of aboral cup height, present on base and on vertical sides of aboral cup (Figs. 3D, 3F, 5A–5C). Five basal plates, equal in size, wider than high, smaller than radial plates. Radial circlet 40–42% of calyx height (Fig. 6A). Radial plates five, pentagonal, largest plates in aboral cup, height to width ratio 0.48–0.58. Radial facets plenary, planate; radial facet topography not known. Posterior interray plates absent from aboral cup; anal sac, if present, unknown.

Arms five, atomous, brachials uniserial; V-shaped in cross section across width of brachial plate; incompletely known (preserved only through sixth primibrachial) (Fig. 6A). First primibrachial (articular plate, see Bohatý and Ausich, 2021), very low, full width of radial facet; subsequent brachials, flat sided, equal in width to distal edge of second primibrachials; height to width ratio ~0.65.

Proximal column narrow, attachment to base of aboral cup circular, occupies slightly more than one half of infrabasal circlet; remainder of column unknown.

Measurements

GIUS4-3693/Hschloth3: CrH, 33.0*; ACH, 7.5; ACmaxW, 11.3; 4thPBH, 6.0; 4thPBW, 4.0.

GIUS4-3693/Hschloth4: ACH, 6.3; ACdistW, 14.0; BH, 6.3; BW, 6.5; RH, 4.3; RW, 8.9.

Remarks

Six specimens of *H. schlotheimii* are known from the Emsian of Poland (GIUS4-3693/Hschloth1 to GIUS4-3693/Hschloth6). In the collection of Polish specimens, small individuals tend to have more pronounced convex basal plates that that nearly produce a central node.

Halocrinites geminatus (Bohatý, 2005)

Fig. 5D, 5H

Type

~~Holotype is  CREF34e-1.~~

Diagnosis

Halocrinites with aboral cup bowl to moderately conical in shape; ratio of aboral cup diameter versus aboral cup height ~1:1.15–2.0; typically 2.0 times wider than high; infrabasal plates fused into a single pentalobate plate that is confined to the basal concavity; brachials wider than high (height to width ratio ~1:2.0–2.5; proximal columnal circular not filling entire basal concavity.

Occurrence

In Poland, *H. geminatus* is from the Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains. Previously, it was known from the Eifelian to early Givetian of Germany (Webster and Webster, 2019).

Description

Calyx medium sized. Aboral cup very low bowl shape (Fig. 5D1), height to maximum width ratio 0.43; perfect pentameral symmetry; shallow, subpentalobate basal concavity occupies ~73% of proximal aboral cup width (Fig. 5D2). Plates gently convex.

Infrabasal circlet completely concealed in basal concavity; outer margin of basal concavity subtetragonal, entirely covered by proximal columnal. Infrabasal plates presumably five. Basal circlet ~56% of aboral cup height, present on base and on vertical sides of aboral cup. Five basal plates, equal in size, wider than high, much smaller than radial plates. Radial circlet ~44% of calyx height. Radial plates five, pentagonal, largest plates in aboral cup, height to width ratio 0.60. Radial facets plenary, planate; radial facet topography not known. Posterior interarray plates absent from aboral cup; anal sac, if present, unknown.

Arms five, atomous, brachials uniserial (Fig. 5H); V-shaped in cross section across width of brachial plate; incompletely known (preserved only through sixth primibrachial). First primibrachial (articular plate, see Bohatý and Ausich, 2021), very low (height to width ratio 0.16), full width of radial facet; subsequent brachials, flat sided, equal in width to distal edge of second primibrachials; height to width ratio ~1.2.

Proximal column attachment to base of aboral cup wide, tetralobate, fills entire basal concavity covering infrabasal plates; remainder of column unknown.

Measurements

GIUS4-3969/Hgem1: CrH, 41.0*; ACH, 9.0; ACmaxW, 21.0; ACproxW, 9.0; BConW, 6.5; BH, 7.6; BW, 7.25; RH, 5.9; RW, 12.4; 1stPbH, 1.5; 1stPbW, 13.0; 2ndPbH, 6.0; 2ndPbproxW, 13.0; 2ndPbdistW, 10.0; 3rdPBH, 5.9.0; 3rdPBW, 5.0.

Remarks

Bohatý (2005) illustrated individuals of *H. geminatus* with a wide variety of shapes. The description above is for the Poland specimens that all have a very low bowl shape. ~~No specimens from Poland preserve remnants of the exotheca sculpturing.~~

Figure 7 and 8 around here

Crinoidea Incertae Sedis

Figs. 5F, 5G, 6B-6D, 7A-7I, 8C, 8D

Occurrence

Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland.

Remarks

A single brachial plate is identified as a flexible crinoid (GIUS4-3693/flexible; Fig. 7I). The brachial plate is ~ 4.0 times deeper than high with, only the distal facet is visible and part of the sides of the plate are visible that includes the aboral indentation where a patelloid process from the distal adjoining brachial would reside. The brachial plate is as wide as deep. A crenulated articular ridge is present along the abaxial portion of the facet, and the lateral sides of the facet are crenulated. A narrow, shallow aboral groove is present along the adaxial margin of the facet.

Poorly preserved and unidentifiable remains of several additional taxa also occur in the Bukowa Góra Member. In addition to the flexible crinoid and holdfasts mentioned below, others include camerate crinoids (Fig. 5F, 5G) and various distinctive cladid crinoids (Fig. 6B-6D). Distinctive columnals and pluricolumnals are also present (Fig. 7A, 8C, 8D). The pluricolumnals

illustrated in Fig. 8C undoubtedly belong to the Platycrinidae and may be *Platycrinites*
minimalis (col.) Gluchowski, 1993a.

447

Simple discoid holdfasts

Fig. 8A, 8B

450

Occurrence

Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, Poland.

Remarks

Solitary rugose coral and a presumable stromatoporoid specimens associated with the described
 crinoids have small, discoid holdfasts cemented to their outer surface (Fig. 8A, 8B). These
 holdfasts are subcircular in outline and some have a slightly digitate outer margin. In one
 example, the holdfast articulation to the column was canted toward the long axis of a rugose
 coral, suggesting the crinoid was encrusted to the coral when both were alive. Therefore, these
 holdfasts should be considered epizoozoans (Taylor and Wilson, 2002). It is not possible to
 speculate on the identity of the crown that was attached to these holdfasts, and the smaller
 specimens may have been juveniles.

462

CONCLUDING REMARKS

The first Emsian crinoids described on the basis of aboral cups and crowns are reported here from the Bukowa Góra Member in the Holy Cross Mountains of southern Poland. Named taxa include *Bactrocrinites* sp., *Codiocrinus sevastopuloi* sp. nov., *Halocrinites geminatus* (Bohatý, 2005), *Halocrinites schlotheimii* Steininger, 1831. Taxa that can only be recognized as incertae sedis include one flexible crinoid, as many as three camerate crinoids, as many as four additional cladid crinoids, and a number of distinctive holdfasts, columnals, and pluricolumnals that cannot be matched with the crown to which they were attached. Additional collecting in the Bukowa Góra Member should yield remains of many crinoids.

Previously, *Halocrinites* (including *H. schlotheimii*) have been described from younger Devonian strata in Poland (see Fig. 1). Further, species of *Bactrocrinites*, *Codiocrinus*, and *Halocrinites* occur in other Devonian crinoid faunas from Germany and Spain (Webster and Webster, 2019).

The new crinoids reported here are from the Bukowa Góra Member of Poland (Emsian) and are an extension of the Lower to Middle Devonian crinoid faunas from across Europe, which are best represented by Emsian to Givetian crinoids from Germany and Spain (e.g., Bohatý, 2005, 2006, 2009, Bohatý and Herbig, 2010; Hauser, 2001, 2002, 2007). *Halocrinites* has been reported from Germany, Spain, Belgium, and Russia (Eifelian–Frasnian); whereas both *Codiocrinus* and *Bactrocrinites* have longer ranges and are cosmopolitan in distribution. In addition, to Western Europe, *Codiocrinus* is known from Gondwana terrane (northern Africa and Australia). The oldest recognized species of *Bactrocrinites* is from the middle Silurian of North America, and this genus is only known from North America and Europe. As known, *Bactrocrinites* became extinct at the Givetian–Frasnian extinction and *Bactrocrinites* and *Halocrinites* became extinct at the Frasnian–Famennian extinction.

487

488 ACKNOWLEDGMENTS

489 We thank Michał Stachacz, Kamil Moskowczenko, Damian Kuźma, Janusz Kucharski and
 490 Mateusz Syncerz for help during the field work. Jan Bohatý generously helped with
 491 identifications of taxa. Paweł Bącal is acknowledged for SEM imaging. We also thank XXX and
 492 XXX, who provided careful reviews that helped us improve this paper. This project was
 493 supported by NCN Grant no. 2018/31/B/ST10/00387.

494

495 REFERENCES

- 496 **Almela A, and Revilla J. 1950.** Especies fósiles nuevas del Devoniano de León. *Notas y*
 497 *Comunicaciones del Instituto Geológico y Minero de España* 20:45–60.
- 498 **Bather, FA. 1890.** British fossil crinoids. II. The classification of the Inadunata Fistulata
 499 (cont'd). *Annals and Magazine of Natural History*, ser. 6, 5:373–388.
- 500 **Bather FA. 1919.** *Cupressocrinus gibber* n. sp. du Dévonien supérieur de Belgique. *Bulletin de*
 501 *la Société Belge de Géologie, de Paléontologie et l'Hydrologie* 28:129–136.
- 502 **Bohatý, JA. 2005.** Doppellagige Kronenplatten: Ein neues anatomisches Merkmal paläozoischer
 503 Crinoiden und Revision der Familie Cupressocrinitidae (Devon). *Paläontologische*
 504 *Zeitschrift* 79(2):201–225.
- 505 **Bohatý, JA. 2009.** Pre- and postmortem skeletal modifications of the Cupressocrinitidae
 506 (Crinoidea, Cladida). *Journal of Paleontology* 83:45–62.
- 507 **Bohatý, JA, Herbig H-G. 2010.** Middle Givetian echinoderms from the Schlade Valley
 508 (Rhenish Massif, Germany): habitats, taxonomy and ecostratigraphy. *Paläontologische*
 509 *Zeitschrift* 84:365–386.

Bohatý JA. 2006. Neue Cupressocrinitidae (Crinoidea) aus den mitteldevischen Kalkmulden der Eifel (linksrheinisches Schiefergebirge, Deutschland). *Senckenbergiana Lethaea* 86(2):151–189.

Bohatý JA, Ausich WI. 2021. Revision of two Devonian cupressocrinitids from the Schultz Collection (Museum of Comparative Zoology, Harvard University) and description of a new *Halocrinites* (Crinoidea, Eucladida). *Journal of Paleontology*, 17 pp. [doi: 10.1017/jpa.2021.65]

Dames W. 1868. Über die in der Umgebung Freiburgs in Nieder-Schlesien auftretenden devonischen Ablagerungen. *Zeitschrift der Deutschen Geologischen Gesellschaft* 21:469–508.

Dubatolova YuA. 1964. Morskije lilii devona Kuzbassa. *Akademiya Nauk SSSR, Sibirskoe Otdeleniye Trudy Instituta Geologii i Geofiziki*, 153 pp.

Fijałkowska-Mader A, Malec J. 2011. Biostratigraphy of the Emsian to Eifelian in the Holy Cross Mountains (Poland). *Geological Quarterly* 55:109–138.

Fijałkowska-Mader A, Malec J, Tarnowska M, Turnau E. 1997. Stratygrafia dolnego dewonu w rejonie Bodzentyna – region łysogórski Gór Świętokrzyskich. *Posiedzenia Naukowe Państwowego Instytutu Geologicznego* 53:122–125.

Filipiak P. 2011. Palynology of the Lower and Middle Devonian deposits in southern and central Poland. *Review of Palaeobotany and Palynology* 166:213–252.

Follman O. 188  Interdevonische Crinoidea. *Verhandlungen des Naturhistorischen Vereins der Preussischen Rheinlande, Westfalens, and des Regierungs-bezirks Osnabruck*, 26 pp.

Gluchowski E. 1980. New taxa of Devonian and Carboniferous crinoid stem parts from Poland. *Bulletin of the Polish Academy of Science* 28:43–49.

- Gluchowski E. 1981a.** Stratigraphic significance of Paleozoic crinoid columnals from Poland. *Zeszyty Naukowe AGH* 7:89–110.
- Gluchowski E. 1981b.** Paleozoic crinoid columnals and pluricolumnals from Poland. *Zeszyty Naukowe AGH* 7:29–57.
- Gluchowski E. 1981c.** Człony łodyg liliowców z serii skalskiej (żywet) Gór Świętokrzyskich. *Sprawozdanie z posiedzeń Komisji Nauk Geologicznych PAN* 22:428–429.
- Gluchowski E. 1993a.** Crinoid assemblages in the Polish Givetian and Frasnian. *Acta Palaeontologica Polonica* 39:35–92.
- Gluchowski E. 1993b.** Upper Emsian crinoids from the Bukowa Góra quarry in the Klonów Range, Holy Cross Mts. *Prace Naukowe Uniwersytetu Śląskiego, 1331, Geologia* 12/13:159–174.
- Gluchowski E. 2002.** Crinoids from the Famennian of the Holy Cross Mountains, Poland. *Acta Palaeontologica Polonica* 47:319–328.
- Gluchowski E. 2003.** Gromada Crinoidea Miller, 1821. In: Malinowska L. (ed.), Budowa Geologiczna Polski. Atlas skamieniałości przewodnich i charakterystycznych. T. III. Dewon. Państwowy Instytut Wydawniczy, Warszawa, 458–475.
- Gluchowski E, Casier J-G, Olempska E. 2006.** Crinoid and ostracod succession within the Early–Middle Frasnian interval in the Wietrzna quarry, Holy Cross Mountains, Poland. *Acta Palaeontologica Polonica* 51:695–706.
- Goldfuss GA. 1826-1844.** Petrefacta Germaniae, tam ea, Quae in Museo Universitatis Regiae Borussicae Fridericiae Wilhelmae Rhenanea, servantur, quam alia quaecunque in Museis Hoeninghusiano Muensteriano aliisque, extant, iconibus et descriptions illustrata. -- Abbildungen und Beschreibungen der Petrefacten Deutschlands und der Angränzende

- Länder, unter Mitwirkung des Herrn Grafen Georg zu Münster, herausgegeben von August Goldfuss. v. 1 (1826–1833), Divisio prima. Zoophytorum reliquiae, p. 1–114; Divisio secunda. Radiariorum reliquiae, p. 115–221 [Echinodermata]; Divisio tertia. Annulatorium reliquiae, p. 222–242; v. 2 (1834–40), Divisio quarta. Molluscorum acephalicorum reliquiae. I. Bivalvia, p. 65–286; II. Brachiopoda, p. 287–303; III. (1841–44), Divisio quinta. Molluscorum gasteropodum reliquiae, p. 1–121; Düsseldorf, Arnz & Co.
- Goldring, W. 1954.** Devonian crinoids. new and old, II. *New York State Museum Circular* 37, 51 pp.
- Gorzelak P, Gluchowski E, Salamon MA. 2014.** Reassessing the improbability of a muscular crinoid stem. *Scientific Reports* 4:4069.
- Gürich G. 1896.** Das Paläozoicum im polnische Mittelgebirge. *Verhandlungen der Russischkaiserlichen mineralogischen gesellschaft zu St. Petersburg* 2, 168 pp.
- Hauser J. 2001.** Neubeschreibung mitteldevonischer Eifelcrinoiden aus der Sammlung Schultze (Museum of Comparative Zoology, The Agassiz Museum, Harvard University, Massachusetts, USA) nebst einer Zusammenstellung der Eifelcrinoiden (Holotypen) der Goldfuss-Sammlung. Published by author, Bonn, 86 pp.
- Hauser J. 2002.** Die Crinoiden der Frasnies–Stufe (Oberdevon) von Wallerheim/Loch (Prümer Mulde, Eifel) nebst einer Zusammenstellung sämtlicher Melocriniten weltweit. Published by author, Bonn, 69 pp.
- Hauser J. 2007.** Die Crinoidenwelt der Eifel vor 350.000.000 Jahren–Neue Crinoiden aus dem Mitteldevon der Eifel, Teil 2. Published by author, Bonn, 83 pp.

- 578 **Jaekel O. 1895.** Beiträge zur Kenntniss der palaeozoischen Crinoiden Deutschlands.
579 *Paläontologisches Abhandlungen* 7, (n.f. 3), p. 1–116.
- 580 **Jaekel O. 1918.** Phylogenie und System der Pelmatozoen. *Paläontologische Zeitschrift* 3:1–
581 128.
- 582 **Jell PA. 1999.** Silurian and Devonian crinoids from central Victoria. *Memoirs of the Queensland*
583 *Museum* 43:114 pp.
- 584 **Jell PA, Holloway DJ. 1983.** Devonian and ?Late Silurian palaeontology of the Winneke
585 Reservoir site, Christman Hills, Victoria. *Proceedings of the Royal Society of Victoria*
586 95:1–21.
- 587 **Jell PA, Jell JS. 1999.** Crinoids, a blastoid and a cyclocystoid from the Upper Devonian reef
588 complex of the Canning Basin, Western Australia. *Memoirs of the Queensland Museum*
589 43: 201–236.
- 590 **König CDE. 1825.** *Icones Fossilium Sectiles*. London, 4 pp.
- 591 **Kongiel R. 1958.** Nowy gatunek *Ammonocrinus* i jego występowanie w Polsce. *Prace Muzeum*
592 *Ziemi* 2:31–40.
- 593 **LeMenn J. 1985.** Les crinoides du Dévonien inférieur et moyen du massif Armoricaïn.
594 *Mémoires de la Société Géologique et Minéralogique de Bretagne* 30, 268 pp.
- 595 **LeMenn J. 1997.** Crinoïdes devoniens d'Afrique du nord. revision systematique et affinités
596 paleobiogeographiques. *Annales de la Société Géologique du Nord*, 2nd ser 5:129–139.
- 597 **Łobanowski H. 1971.** The Lower Devonian in the western part of the Klonów belt (Holy Cross
598 Mts). Part 1. Upper Emsian. *Acta Geologica Polonica* 21:629–686.

- 599 **Łobanowski H. 1981.** Bukowa Góra quarry: Lower Devonian sandstones. In: Żakowa H. (ed.),
600 Przewodnik 53 Zjazdu Polskiego Towarzystwa Geologicznego w Kielcach,
601 Wydawnictwa Geologiczne; Warszawa, 249–255.
- 602 **Malec J. 2005.** Lithostratigraphy of the Lower and Middle Devonian boundary interval in the
603 Łysogóry Region of the Holy Cross Mts (Poland). *Biuletyn Państwowego Instytutu*
604 *Geologicznego* 415:5–28.
- 605 **Marynowski L, Salamon MA, Narkiewicz M. 2002.** Thermal maturity and depositional
606 environments of organic matter in the post-Variscan succession on the Holy Cross
607 Mountains. *Geological Quarterly* 46:25–36.
- 608 **Miller JS. 1821.** *A natural history of the Crinoidea, or lily-shaped animals; with observations*
609 *on the genera, Asteria, Euryale, Comatula and Marsupites.* Bristol, C. Frost, 150 pp.
- 610 **Moore RC, Jeffords RM. 1968.** Classification and nomenclature of fossil crinoids based on
611 studies of dissociated parts of their columns. *The University of Kansas Paleontological*
612 *Contributions* 46:1–86.
- 613 **Moore RC, Laudon LR. 1943.** Evolution and classification of Paleozoic crinoids. *Geological*
614 *Society of America Special Paper* 46, 151 pp.
- 615 **Piotrowski A. 1977.** Genus *Ammonicrinus* (Crinoidea) from the Middle Devonian of the Holy
616 Cross Mountains, Poland. *Acta Palaeontologica Polonica* 22:205–218.
- 617 **Prokop RJ. 1973.** *Elicrinus* n. gen. from the Lower Devonian of Bohemia (Crinoidea). *Věstník*
618 *Ustředního Ústavu Geologického* 48(4):221–224.

- 619 **Reed FRC. 1908.** The Devonian faunas of the Northern Shan states. *India Geological Survey,*
620 *Memoirs* 2(5). 157 pp.
- 621 **Roemer, CF. 1844.** *Das Rheinische Übergangsgebirge. Eine palaeontologisch-geognostische*
622 *Darstellung.* Hannover, Hahn, 96 pp.
- 623 **Roemer, CF. 1850.** Über *Stephanocrinus*, eine fossile Crinoiden-Gattung aus der Familie der
624 Cystideen. *Wiegmanns Archiv für Naturgeschichte, Jahrgang* 16:365–375.
- 625 **Roemer CF. 1852–54.** Erste Periode, Kohlen-Gebirge. In *Lethaea Geognostica*, H. G. Bronn,
626 1851–1856, 3rd edit. Stuttgart, E. Schweizerbart, 2, 788 pp.
- 627 **Salamon MA, Gerrienne P, Steemans P, Gorzelak P, Filipiak P, Le Hérissé A, Paris F,**
628 **Cascales-Miñana B, Brachaniec T, Misz-Kennan M, Niedźwiedzki R, Trela W. 2018.**
629 Putative Late Ordovician land plants. *New Phytologist* DOI: doi.org/10.1111/nph.15091.
- 630 **Sandberger G, Sandberger F. 1850-1856.** *Die Versteinerungen des rheinischen*
631 *Schichtensystems in Nassau.* Wiesbaden, Kreidel and Niedner, 564 pp.
- 632 **Schmidt WE. 1934.** Die Crinoideen des Rheinischen Devons, I. Teil; Die Crinoideen des
633 Hunsrückschiefers. **Abhandlung der Preussischen Geologischen Landesanstalt** 163,
634 149 pp.
- 635 **Schmidt WE. 1941.** Die Crinoiden des Rheinischen Devons II. Teil. A. Nachtrag zu: Die
636 Crinoiden des Hunsrückschiefers; B. Die Crinoideen des Unterdevons bis zur
637 *cultrijugatus*-Zone (mit Ausschluß des Hunsrückschiefers). *Abhandlungen der*
638 *Preußischen Geologischen Landesanstalt* 182, 253 pp.

- Schmidt WE. 1942.** Die Crinoideen des Rheinischen Devons, II. Teil; A. Nachtrag zu Die Crinoideen des Hunsrückschiefers; B. Die Crinoideen des Unterdevon bis zur *Cultrijugatus* -Zone (mit Ausschluss des Hunsrückschiefers). *Abhandlungen der Reichsstelle für Bodenforschung, n. s.*, 182, 253 pp.
- Sobolev D. 1909.** Sredniy devon kelecko-sandomirskoho krjaza. *Materialy dla geologii Rossiji* 24:41–536.
- Schultze LJT. 1866.** *Monographie der Echinodermen des Eifler Kalkes*. Vienna, Carl Gerold's Sohn, 118 pp.
- Springer F and Slocom AW. 1906.** *Hypsocrinus*, a new genus of crinoids from the Devonian. *Field Columbia Museum Publication*, 114, Geology Series 2(9): 267–271,
- Steininger J. 1831.** Bemerkungen über die Versteinerungen, welche in dem Uebergangs-Kalkgebirge der Eifel gefunden werden. Eine Beilage zum Gymnasial-Program zu Trier, Trier, 44 pp.
- Steininger J. 1849.** Die Versteinerungen des Uebergangs-Gebirges der Eifel. *Jahresbericht über den Schul-Cursus 1848/49 an dem Gymnasium zu Trier*, 21–22.
- Strimple, HL. 1952.** Some new species of crinoids from the Henryhouse Formation of Oklahoma. *Journal of the Washington Academy of Science* 42:75–79.
- Stukalina, GA. 1965.** Novye vidji *Hexacrinites* tsentralnogo Kazakhstana [New species of *Hexacrinites* from central Kazakhstan]. *Ezhegodnik Vsesoyuznogo Paleontologičeskogo Obščestva*, v. 17, p. 188–195 (In Russian).
- Szulczewski M. 1995.** Depositional evolution of the Holy Cross Mts. (Poland) in the Devonian and Carboniferous – a review. *Geological Quarterly* 39:159–174.

Szulczewski M, Porębski S. 2008. Stop 1 – Bukowa Góra, Lower Devonian. In: Pieńkowski G. and Uchman A. (eds), Ichnological Sites of Poland – the Holy Cross Mountains and the Carpathian Flysch: The Second International Congress on Ichnology, The Pre-Congress and Post-Congress Field Trip Guide Book, Cracow, 18–37.

Taylor PD, Wilson MA. 2002. A new terminology for marine organisms inhabiting hard substrates. *PALAIOS* 17:522–525.

Wachsmuth C, Springer F. 1880–1886. Revision of the Palaeocrinoidea. Proceedings of the Academy of Natural Sciences of Philadelphia Pt. I. The families Ichthyocrinidae and Cyathocrinidae (1880), p. 226–378, (separate repaged p. 1–153). Pt. II. Family Sphaeroidocrinidae, with the sub-families Platycrinidae, Rhodocrinidae, and Actinocrinidae (1881), p. 177–411 (separate repaged, p. 1–237). Pt. III, Sec. 1. Discussion of the classification and relations of the brachiate crinoids, and conclusion of the generic descriptions (1885), p. 225–364 (separate repaged, 1–138). Pt. III, Sec. 2. Discussion of the classification and relations of the brachiate crinoids, and conclusion of the generic descriptions (1886), p. 64–226 (separate repaged to continue with section 1, 139–302).

Webster GD, Webster DW. 2019. Bibliography and index of Paleozoic crinoids, coronoids, and hemistreptocrinids, 1758–2012. <http://crinoids.azurewebsites.net/> (accessed September 2021).

Webster GD, Becker RT, Maples CG. 2005. Biostratigraphy, paleoecology, and taxonomy of Devonian (Emsian and Famennian) crinoids from southeastern Morocco. *Journal of Paleontology* 79: 1052–1071.

- Wójcik K. 2015.** The uppermost Emsian and lower Eifelian in the Kielce Region of the Holy Cross Mts. Part I: Lithostratigraphy. *Acta Geologica Polonica* 65:141–179.
- Wright DF. 2017.** Bayesian estimation of fossil phylogenies and the evolution of early to middle Paleozoic crinoids (Echinodermata). *Journal of Paleontology* 91:799–814, [doi: 10.1017/jpa.2016.141].
- Wright DF, Ausich WI, Cole SR, Peter ME, Rhenberg EC. 2017.** Phylogenetic taxonomy and classification of the Crinoidea (Echinodermata). *Journal of Paleontology* 91:829–846.
- Yeltyschewa RS, Dubalotova YA. 1961.** Morskie lilii V kn. Biostratigrafiia paleozoia Sayano-Altaiskoi gornoi oblasti, Srednii paleozoi, t. II [Crinoids. In Biostratigraphy of the Paleozoic of the Sayano Altai mountainous country, middle Paleozoic, v. II]. Moscow, *Izdatel'stvo Nauka, Trudy, pt. 20*:294–296, 552–560. (In Russian)
- Yeltyschewa RS. and Sisova EN. 1973.** Anthinocrinidae-Novoe Semeistvo Srednepaleozoiskikh Morskikh Lili V kn. Voprosy Stratifrafi i Tektoniki Vostochnogo Kazakhstana [Anthinocrinidae New Family of Middle Paleozoic Marine Lily. In Aleksandrova, M. I., ed, Questions on the Stratigraphy and Tectonics of East Kazakhstan]. *Trudy Vsesoyuznogo Ordena Lenina Nauchno-Issledovatel'skogo Geologicheskogo Instituta, n. s.* 160:86–99 (In Russian)
- Zeiler F, Wirtgen PW. 1855.** Bemerkungen über die Petreefacten der älteren devonischen Gebirge am Rheine, insbesondere über die Echinodermen in der Umgegend von Coblenz und in dem Eifeler Kalke vorkommenden Arten. *Verhandlungen des Naturhistorisch Vereins der Preussischen Rheinlands und Westfalen* 12:79–85.

704 **Zeuschner L. 1867.** Devonische Formation im Sandomirer-Chenciner Gebirge. *Neues Jahrbuch*
705 *für Mineralogie, Geologie und Paläontologie* 67:478–508.

706 **Zeuschner L. 1869.** Geognostische Beschreibung der mittleren devonischen Schichten zwischen
707 Grzegorzowice und Skąły-Zagaje, bei Nowa Słupia. *Zeitschrift der Deutschen*
708 *Geologischen Gesellschaft* 21:263–274.

709 **Zittel KA, von. 1895.** *Grundzüge der Palaeontologie (Palaeozoologie)*, 1st edit. R. Oldenbourg,
710 München, 971 pp.

711

712 Captions:

713 Figure 1 Stratigraphic ranges of crinoid taxa recorded in Devonian of Poland. Data compiled
714 after: Dames (1868), Zeuschner (1867, 1869), Gürich (1896), Sobolev (1909), Kongiel (1958),
715 Piotrowski (1977), Głuchowski (1980, 1981a-c, 1993, 2002, 2003), and Hauser (2002).

716

717 Figure 2 (A) The map of Poland with the Holy Cross Mountains area marked as grey rectangle.
718 B. The lithostratigraphical scheme of Middle and Upper Devonian in the Bukowa Góra Quarry.
719 Compiled after Marynowski et al. (2002), Malec (2005), Szulczewski & Porębski (2008), Wójcik
720 (2015), and Salamon et al. (2018).

721

722 Figure 3 (A1-A4) *Codiocrinus sevastopuloi* sp. nov. GIUS 4-3696/Codiocrinus1, holotype; (A1)
723 lateral view of aboral cup; (A2) basal view of aboral cup, note basal concavity bordered by ridge;
724 (A3), oral view of aboral cup; (A4), lateral view of aboral cup. (B1-B3) *Codiocrinus*

sevastopuloi sp. nov. GIUS 4-3696/Codiocrinus2. paratype; (B1) lateral view of aboral cup; (B2) basal view of aboral cup; (B3) oral view of aboral cup. (C) lateral view of an incomplete and compressed specimen of *Codiocrinus sevastopuloi* sp. nov. GIUS 4-3696/Codiocrinus3. (D1-D2) compressed specimen of *Halocrinites* sp. GIUS 4-3696/Hsp; (D1) lateral view of aboral cup with plate boundaries visible; (D2) basal view of cup. (E) *Bactrocrinites* sp. GIUS 4-3696/Bactrocrinites1; lateral view of incomplete aboral cup, note small radial plates and large basal plates. (F1-F3) compressed aboral cup of *Halocrinites schlotheimii*. GIUS 4-3696/Hschloth1 (F1) oblique basal view; (F2, F3) lateral views. All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. Scale bar equals 10 mm.

Figure 4. Plate drawing of the posterior of *Bactrocrinites* sp.

Figure 5 (A1-A3, B1-B2, C1-C2) *Halocrinites schlotheimii*. GIUS 4-3696/Hschloth2, 5 and 4 respectively; (A1, B1, C1) lateral views of aboral cup; (A2, B2, C2) basal views of aboral cup; (A3) oral view of aboral cup. (D, H) *Halocrinites geminatus*. GIUS 4-3696/Hgem2 and 1 respectively; (D1) lateral view of aboral cup; (D2) basal view of aboral cup. (E) *Halocrinites* with good exoplacoid sculpturing preserved; GIUS 4-3696/Hschloth6. (F, G) Crinoidea indeterminate (presumably remains of a camerate crinoid). GIUS 4-3696/indet1 and 2 respectively. All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. Scale bar equals 10 mm.

Figure 6 (A1-A2) *Halocrinites schlotheimii*; (A1) lateral view of partial crown, aboral cup plate boundaries distinct; (A2) basal view of a moderately compressed aboral cup. GIUS 4-3696/Hschloth3. (B-D) Crinoidea indeterminate; GIUS 4-3696/indet3, 4 and 5 respectively; (B, C) partial arms of an unknown cladid crinoids; (D) in upper left of specimen a pentalobate column presumably infrabasal plates attached; probably an unknown cladid). All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. Scale bar equals 10 mm.

Figure 7 (A-H) crinoid pluricolumnals. GIUS 4-3696/indet6-13. (A, B) pluricolumnal with numerous nodes around the periphery of each columnal; (C), set of pluricolumnals; (D) pluricolumnal with nodes and perhaps some spines around the periphery of each columnal; (E) numerous pluricolumnals of a column that lacks nodes; (F) lateral view of three-columnal pluricolumnal with a few nodes around the periphery of columnals that are offset in position from one columnal to the next; (G) columnal facet with a narrow peripheral lumen and a narrow, raised perilumen; (H) columnal with a wide lumen and a narrow, raised perilumen: (I) Flexible crinoid brachial. GIUS 4-3696/flexible; note crenulate sides and on the upper margin a notch to hold a patelloid process from the next highest columnal. All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. All scale bar equals 10 mm but in case of flexible crinoid brachial it is 1 mm.

Figure 8 (A, B) crinoid holdfast on stromatoporoids. GIUS 4-3696/holdfast1 and 2 respectively. (C) crinoid pluricolumnal presumably from the dististele with broken radices; (D) pluricolumnal

769 with elliptical columnals, presumably from a member of the Platycrinitidae; GIUS 4-
 770 3696/indet14 and 15 respectively. All specimens are from Bukowa Góra Member (Emsian),
 771 Bukowa Góra quarry, Holy Cross Mountains, southern Poland. Scale bar equals 10 mm.

Figure 1

Figure1. Stratigraphic ranges of crinoid taxa recorded in Devonian of Poland.

Data compiled after: Dames (1868), Zeuschner (1867, 1869), Gürich (1896), Sobolev (1909), Kongiel (1958), Piotrowski (1977), Głuchowski (1980, 1981a-c, 1993, 2002, 2003), and Hauser (2002).

Lochkovian	Pragian	Emsian	Eifelian	Givetian	Frasnian	Famenn.	Crinoid taxa
							<i>Kotancrinus</i> cf. <i>balaensis</i> (col.)
							<i>Poteroicrinites?</i> <i>morlieresensis</i>
							<i>Formosocrinus</i> cf. <i>formosus</i> (col.)
							<i>Acanthocrinus</i> sp. (col.)
							<i>Marettocrinus subbiconcavus</i> (col.)
							<i>Laudonomphalus humilicarinatus</i> (col.)
							<i>Procupressocrinites gracilis</i>
							<i>Mediocrinus microgrumosus</i> (col.)
							<i>Kazachstanocrinus acutilobus</i> (col.)
							<i>Asperocrinus brevispinosus</i> (col.)
							<i>Ricebocrinus kulagaiensis</i> (col.)
							<i>Ammonocrinus sulcatus</i> (col.)
							<i>Eocampocrinus fragilis</i> (col.)
							<i>Gilbertocrinus vetulus</i> (col.)
							<i>Myelodactylus canaliculatus</i> (col.)
							<i>Pentagonostipes petaloides</i> (col.)
							<i>Platycrinites minimalis</i> (col.)
							<i>Noctoicrinus?</i> <i>varius</i> (col.)
							<i>Marettocrinus angustannulus</i> (col.)
							<i>Halocrinites schlotheimi</i>
							<i>Praerocrinus polonicus</i> (col.)
							<i>Stenocrinus</i> cf. <i>bifurcatus</i> (col.)
							<i>Stenocrinus degratus</i> (col.)
							<i>Tantalocrinus scutellus</i> (col.)
							<i>Tjeecrinus crassijugatus</i> (col.)
							<i>Schyschcatocrinus creber</i> (col.)
							<i>Kazachstanocrinus tenuis</i> (col.)
							<i>Stenocrinus raricostatus</i> (col.)
							<i>Marettocrinus kartzevae</i> (col.)
							<i>Ammonocrinus kongieli</i> (col.)
							<i>Amurocrinus imatschensis</i> (col.)
							<i>Haplocrinites minor</i>
							<i>Laudonomphalus pinguicostatus</i> (col.)
							<i>Haplocrinites aremoricanus</i>
							<i>Anthinocrinus brevicostatus</i> (col.)
							<i>Anthinocrinus primaevus</i> (col.)
							<i>Halocrinites inflatus</i>
							<i>Ononicrinus gracilis</i> (col.)
							<i>Urushicrinus perbellus</i> (col.)
							<i>Ricebocrinus parvus</i> (col.)
							<i>Tjeecrinus simplex</i> (col.)
							<i>Schyschcatocrinus multififormis</i> (col.)
							<i>Haplocrinites</i> sp.
							<i>Schyschcatocrinus delicatus</i> (col.)
							<i>Wenjukowiocrinus wenjukowi</i> (col.)
							<i>Caleocrinus bicostatus</i> (col.)
							<i>Glyphidocrinus infimus</i> (col.)
							<i>Caleocrinus kielcensis</i> (col.)
							<i>Melocrinites</i> cf. <i>hieroglyphicus</i>
							<i>Melocrinites</i> cf. <i>gibbosus</i>
							<i>Exaesidiscus compositus</i> (col.)
							<i>Haplocrinites gluchowskii</i>
							<i>Tjeecrinus insectus</i> (col.)
							<i>Cosmocrinus polonicus</i> (col.)

Figure 2

Figure 2. (A) The map of Poland with the Holy Cross Mountains area marked as grey rectangle. (B) The lithostratigraphical scheme of Middle and Upper Devonian in the Bukowa Góra Quarry.

Compiled after Marynowski et al. (2002), Malec (2005), Szulczewski & Porębski (2008), Wójcik (2015), and Salamon et al. (2018).





Figure 3

Figure 3. (A1-A4) *Codiocrinus sevastopuloi* sp. nov. GIUS 4-3696/Codiocrinus1, holotype; (A1) lateral view of aboral cup; (A2) basal view of aboral cup, note basal concavity bordered by ridge; (A3), oral view of aboral cup; (A4), lateral

All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. Scale bar equals 10 mm.




Figure 4

Figure 4. Plate drawing of the posterior of *Bactrocrinites* sp.




Figure 5

Figure 5. (A1-A3, B1-B2, C1-C2) *Halocrinites schlotheimii*. GIUS 4-3696/Hschloth2, 5 and 4 respectively; (A1, B1, C1) lateral views of aboral cup; (A2, B2, C2) basal views of aboral cup; (A3) oral view of aboral cup. (D, H) [i]Halocrinites gem

All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. Scale bar equals 10 mm.




Figure 6

Figure 6. (A1-A2) *Halocrinites schlotheimii*; (A1) lateral view of partial crown, aboral cup plate boundaries distinct; (A2) basal view of a moderately compressed aboral cup. GIUS 4-3696/Hschloth3. (B-D) Crinoidea indeterminate; GIUS 4-3

All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. Scale bar equals 10 mm.




Figure 7

Figure 7. (A-H) crinoid pluricolumnals. GIUS 4-3696/indet6-13. (A, B) pluricolumnal with numerous nodes around the periphery of each columnal; (C), set of pluricolumnals; (D) pluricolumnal with nodes and perhaps some spines around the periphery of

All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. All scale bar equals 10 mm but in case of flexible crinoid brachial it is 1 mm.





Figure 8

Figure 8. (A, B) crinoid holdfast on stromatoporoids. GIUS 4-3696/holdfast1 and 2 respectively. (C) crinoid pluricolumnal presumably from the dististele with broken radices; (D) pluricolumnal with elliptical columnals, presumably from a member of t

All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. Scale bar equals 10 mm.



A



B



C



D