

# Full vaccination coverage and associated factors among children aged 12 to 23 months in remote rural area of Demba Gofa District, Southern Ethiopia

Tadele Dana Darebo<sup>Corresp., 1</sup>, Bahru Belachew Oshe<sup>2</sup>, Chala Wegi Diro<sup>1</sup>

<sup>1</sup> School of Public Health, Wolaita Sodo University, Wolaita Sodo, South, Ethiopia

<sup>2</sup> Maternal and child health department, Gofa zone health office, Gofa, South, Ethiopia

Corresponding Author: Tadele Dana Darebo

Email address: danatadele@gmail.com

**Background:** Full vaccination refers to the administration of vaccines/antigens recommended for children in the first year of life. However, little is known about full vaccination in remote, rural Ethiopia. This study aimed to measure full vaccination coverage among children aged 12 to 23 months and associated factors in Demba Gofa District, Southern Ethiopia. **Methods:** A cross-sectional community-based study was conducted in April and May 2019 using a multistage sampling technique to select 677 mothers with children 12-23 months of age. Data was collected using a pre-tested structured questionnaire, and data were edited, coded, entered, and cleaned using Epi Info v3.1 and analyzed using SPSS v20. Bivariate and multivariable logistic regression was used to understand associations between dependent and independent variables. **Results:** Three-hundred and nine children (47.0%) were fully vaccinated, 274 (41.7%) were partially vaccinated, and 74 (11.3%) were not vaccinated at all. Children were more likely to be vaccinated if decisions were made jointly with husbands (AOR=1.88, 95% CI: 1.06, 3.34), were made by mothers (AOR=4.03, 95% CI: 1.66, 9.78), followed postnatal care (AOR=5.02, 95% CI: 2.28, 11.05), if the child's age for completing vaccination was known (AOR=2.54, 95% CI: 1.04, 6.23), and if vaccinations did not make the child sick (AOR=0.32, 95% CI: 0.16, 0.64). **Conclusion:** Full vaccination coverage was less than average in the study district and far below the governmental target (90%) necessary for sustained control of vaccine-preventable diseases. Interventions targeted towards maternal healthcare decision-making, postnatal care, knowledge on vaccination timing, and importance should be prioritized to improve full vaccination coverage.

**Keywords:** Full vaccination, 12-23 months, children, Demba Gofa District, Southern Ethiopia

Full Vaccination Coverage and Associated Factors Among Children Aged 12 to 23 Months in  
remote rural area of Demba Gofa District, Southern Ethiopia

Tadele Dana Darebo<sup>1</sup>, Bahru Belachew Oshe<sup>2</sup>, Chala Wegi Diro<sup>1</sup>

<sup>1</sup> School of Public Health, Wolaita Sodo University, W/Sodo, South, Ethiopia

<sup>2</sup> Maternal and child health, Gofa zone health office, Gofa, South, Ethiopia

Corresponding Author:

Tadele Darebo

WSU street, Wolaita Sodo, South Ethiopia, Ethiopia

Email address: [danatadele@gmail.com](mailto:danatadele@gmail.com)

# Abstract

**Background:** Full vaccination refers to the administration of vaccines/antigens recommended for children in the first year of life. However, little is known about full vaccination in remote, rural Ethiopia. This study aimed to measure full vaccination coverage among children aged 12 to 23 months and associated factors in Demba Gofa District, Southern Ethiopia.

**Methods:** A cross-sectional community-based study was conducted in April and May 2019 using a multistage sampling technique to select 677 mothers with children 12-23 months of age. Data was collected using a pre-tested structured questionnaire, and data were edited, coded, entered, and cleaned using Epi Info v3.1 and analyzed using SPSS v20. Bivariate and multivariable logistic regression was used to understand associations between dependent and independent variables.

**Results:** Three-hundred and nine children (47.0%) were fully vaccinated, 274 (41.7%) were partially vaccinated, and 74 (11.3%) were not vaccinated at all. Children were more likely to be vaccinated if decisions were made jointly with husbands (AOR=1.88, 95% CI: 1.06, 3.34), were made by mothers (AOR=4.03, 95% CI: 1.66, 9.78), followed postnatal care (AOR=5.02, 95% CI: 2.28, 11.05), if the child's age for completing vaccination was known (AOR=2.54, 95% CI: 1.04, 6.23), and if vaccinations did not make the child sick (AOR=0.32, 95% CI: 0.16, 0.64).

**Conclusion:** Full vaccination coverage was less than average in the study district and far below the governmental target (90%) necessary for sustained control of vaccine-preventable diseases. Interventions targeted towards maternal healthcare decision-making, postnatal care, knowledge on vaccination timing, and importance should be prioritized to improve full vaccination coverage.

**Keywords:** Full vaccination, 12-23 months, children, Demba Gofa District, Southern Ethiopia

# Introduction

vaccination is the efficient way to protect people from infectious diseases by stimulating the host immune system to produce antibodies or other specific immune defenses. (1, 2). Vaccination has reduced and eliminated various childhood diseases globally, including diphtheria, tetanus, pertussis, polio, measles, and tuberculosis. Several new vaccines have now been added to public health programs (hepatitis B, *Haemophilus influenzae* B (HiB), pneumococcus, and rotavirus) (3-10). Global vaccination coverage has increased, contributing to decreases in child mortality from 9.6 million in 2000 to 5.9 million in 2015, especially low and middle-income countries (LMICs) (3,11-18).

The World Health Organization (WHO) recommends, which is also strictly endorsed by Ethiopia, states that a child is fully vaccinated if he/she has the BCG (tuberculosis) vaccine at birth; oral polio vaccine (OPV) at birth, 6, 10, and 14 weeks; pentavalent vaccine for diphtheria, tetanus, pertussis (whooping cough), hepatitis B, and HiB at 6, 10, and 14 weeks; pneumococcal conjugate vaccine (PCV) at 6, 10, and 14 weeks; rotavirus at 6 and 10 weeks; inactivated polio vaccine (IPV) at 14 weeks; and measles at 9 months (4-6). The Expanded Program on Immunization (EPI) agenda plans by applying WHO standards to give the primary vaccination series to at least 90% of children. Despite these efforts, the target has still to be attained in many LMICs (5,19-25). For instance, each day over 20,000 children die from preventable infectious diseases, and a larger part of these deaths occur in sub-Saharan Africa (SSA) and South East Asia (SEA) compared to the rest of the world (7,9,26-29).

Globally, 22.6 million infant children are partially protected through vaccination and over 25% of these are reported to be in LMICs (14-16). An estimated 2.5 million children aged less than five years die annually due to diseases that can be prevented by vaccination (11, 15,30). About 14.8 million children who are not vaccinated with pentavalent 3 before celebrating their first year are found only in ten countries, including Ethiopia (14,15,29,30). In spite of the fact that most under-five deaths can be easily tackled by vaccination, about half occur in SSA (4,6,31-38). The most common vaccine preventable diseases that result in morbidity and mortality under-five children are pneumonia, diarrheal diseases and measles. More than three million cases of pneumonia is reported eah year claiming the lives of about 20% of an estimated 40,000 annual underfie death. Diarrehal diseases follows accounting 1.7 million cases per year and measles

being the third with the 50 per one million population per year with the case fatality rate of 3-6%(39,40). Ethiopia aimed to target at least 90% of the population with all vaccines by 2020. However, studies conducted in Ethiopia in children aged 12-23 months have reported vaccination rates from 39% to 43%. Full vaccination coverage highly discrepant at the regional level lowest in Afar with 15% highest Addis Ababa 89%, , with different factors including sociodemographic and economic factors, maternal healthcare service utilization, knowledge and attitudes of mothers/caregivers towards vaccination, accessibility, and perceived availability identified to affect full vaccination coverage (21,22,38,39). However, there is still a lack of information on the prevalence and associated factors of full vaccination coverage across different regions of Ethiopia and areas of the country where epidemics intermittently occur. This study therefore aimed to measure the coverage of full vaccination and associated factors among children aged 12 to 23 months in Demba Gofa District, Southern Ethiopia to assist program implementers and frontline health workers understand the factors influencing full vaccination coverage and to plan for default tracing mechanisms.

## Materials & Methods

### Study setting and design

A community-based cross-sectional study was conducted in April and May 2019 in the Demba Gofa District, Ethiopia, which is located in Southern Ethiopia 514 km from Addis Ababa. The district contains 34 kebeles, all rural, with a total population of 96,427; 1,659 (1.72%) were children aged 12-23 months. Demba Gofa District has four governmental health centers, 34 health posts, and 15 private primary clinics. The EPI service was provided only by governmental health facilities (45).

### Patient and public involvement

Not applicable for this study.

### Source population

The source population was all children aged 12-23 months paired with their mothers/caregivers.

# 99 Inclusion/exclusion criteria

100 The inclusion criteria were all mothers/caregivers with children aged 12-23 months who were  
101 residents of selected kebeles (for at least six months) in the district, while exclusion criteria were  
102 mentally disabled/critically ill mothers/caretakers.

# 103 **Sample size determination**

104 The sample size was determined using the single population proportion formula and P=27.7% as  
105 the prevalence of full immunization coverage among children aged 12-23 months in Ambo  
106 district, Central Ethiopia (41). By using a 95% level of significance, a margin of error of 5%, a  
107 design effect of 2, and a non-response rate of 10%, the final sample size was 677:

$$108 \quad n = \frac{(Z\alpha/2)^2 p(1-P)}{d^2}$$

109

# 110 Sampling procedure

111 Multi-stage sampling was used to select the study population at the community level. In the first  
112 stage, ten kebeles were selected using a simple random sampling method from a total of 34  
113 kebeles in the district. The total sample size was divided by the size of all households that had  
114 children aged 12-23 months in each selected kebele. Then, the total sample size was  
115 proportionally allocated according to the number of children in each kebele. Finally, a simple  
116 random sampling method was employed to select children using the lottery method from each  
117 kebele. When there were two or more children in the same household, the lottery method was  
118 used to select only one. During absence of eligible child in the household, the next one was  
119 included. If eligible participants absent during data collection, households are revisited twice by  
120 the interviewers.

# 121 Data collection and quality control

122 Data were collected using a pre-tested and structured questionnaire developed from the related  
123 literature used to conduct similar study in elsewhere(8-11). The questionnaire was initially  
124 prepared in English and then translated to Gofatho (local Gofa language) and then translated  
125 back into English by the qualified translators to check for consistency. Data were collected  
126 through face-to-face interviews from mothers/caregivers. Training was given for two days before  
127 data collection by the principal investigator to the data collectors and supervisors on the purpose of

the study, data collection tools, and on minimizing the recall bias. Mothers/caregivers were asked to show vaccination cards (if available, the interviewer copied the vaccination dates), and for those who had no or lost vaccination cards, mothers/caregivers were asked about the vaccination status of their children using different recall mechanisms such as asking the route of administration, injection sites, examining the BCG scar, etc. Pretesting was performed in 5% (8,23) of the sample size in kebeles outside of the study area after which necessary modifications were undertaken. Daily check-ups were performed for data completeness, accuracy, and consistency.

# Operational definitions

**Fully vaccinated:** A child aged 12-23 months who received one dose of BCG, at least three doses of OPV, three doses of pentavalent, three doses of PCV, two doses of Rota, and one dose of measles vaccines before celebrating the first birth year(10).

**Partially vaccinated:** If a child missed at least one dose of the ten vaccines mentioned above(10).

**Unvaccinated:** A child who had not received any of the ten vaccines(10).

**Knowledge of vaccination:** A mother/caregiver answering a mean or above of knowledge questions was considered to have good knowledge, while a mother/caregiver answering below a mean of knowledge questions was considered to have poor knowledge(8,14).

**Attitude towards vaccination:** When the respondent answered the mean and above of the ten attitude questions were deemed to have a positive attitude and those answering below the mean were deemed to have a negative attitude(8,14).

# Data management and analysis

After data collection, data were edited and coded for completeness and consistency and then entered into Epi Data v3.1 and exported to SPSS v20 for analysis. After cleaning data for inconsistencies, errors, and missing values, descriptive statistics (mean, median, SD, percent, frequency) were calculated to visualize the overall distribution of the study subjects for the variables under study. Bivariate analysis was performed to determine associations between independent and full vaccination status. Multicollinearity was checked using a cutoff <10 based on the variance inflation factor (VIF) or tolerance test >0.1. The necessary assumptions of logistic regression were checked using Hosmer and Lemeshow tests to assess the fitness of the

model at a P-value < 0.05. All explanatory variables associated with the full vaccination status at p-value  $\leq 0.25$  were selected for multivariable analysis. Finally, multivariate analysis was used to measure the degree of association between independent and outcome variable. Adjusted odds ratio(AOR) with 95% CI was used to determine statistically significant.

## Ethical considerations

Ethical approval was obtained from the Research Ethical Review Committee of the College of Health Sciences and Medicine of Wolaita Sodo University under the number CHSM/ERC/42. An official letter of cooperation was taken to the Gofa Zone Health Department and Demba Gofa District Health Office and a permission letter was then received from each. Official letters also were written to the kebele administration and health posts. Informed written consent was obtained from participants after explaining the aims of the study, and the confidentiality and anonymity of information was preserved.

# Results

## Socio-demographic and economic characteristics of study participants

From a total of 677 mothers with children aged 12-23 months old, 657 were interviewed, a response rate of 97%. Most respondents (632, 96.2%) were mothers of the children, 77.8% were Protestant, 634 (96.5%) were married, and all were of Gofa ethnicity. The majority of mothers/caregivers (260, 39.6%) were illiterate, and most husbands (497, 75.6%) were farmers. 131 (19.9%) families were grouped into the richest wealth quantile. The average family size was 5.79 with a range from 2 to 12. Over half (376, 57.2%) of the children were male, and most belonged to families with multiple children (**Table 1**).

## Maternal healthcare service utilization

Four hundred and sixty-eight (72.0%) mothers sought healthcare services through joint decisions with their husbands. Most respondents (545, 83.8%) had received antenatal care (ANC) during their last pregnancy, most (517, 79.5%) had attended ANC at least two or more times, and 485 (74.6%) had taken tetanus injections. Four hundred and twenty-five (65.4%) mothers gave birth to their last baby in healthcare institutions, but only 62 (9.5%) mothers attended postnatal care at least once (**Table 2**).



# **Knowledge and attitudes of mothers/caregivers towards vaccination and vaccine-preventable disease**

Of 657 total respondents, 603 (92.8%) had ever heard about vaccination and vaccine-preventable diseases. The majority of respondents (549, 91.0%) heard about vaccination and vaccine-preventable diseases from health care workers. Nearly half of the respondents (308, 47.4%) mentioned measles as a specific vaccine-preventable disease, and 210 (32.3%) mentioned tetanus. Respondents were also asked the objective of vaccinating children, the age at which a child should begin vaccination, the sessions needed to complete vaccination, and the correct age at which a child completes its vaccination program. Regarding knowledge, almost three-quarters (487, 74.0%) of study participants have a good knowledge, whereas 469 (71.4%) of study subjects have a positive attitude towards vaccination.

## **Accessibility and vaccination coverage**

All respondents reported that they had access to a health facility that provided vaccination services near them. Over 90% of health facilities that provided vaccination services in the Demba Gofa District were reported to be health posts. With respect to vaccination delivery strategy, 351 (53.4%) were static, 158 (24.0%) were outreach, and 148 (22.5%) were home to home.

From a total of 657 children aged 12-23 months old, under half (309, 47.0%) were fully vaccinated or completed all the recommended vaccines before celebrating their first birthday. No child had a vaccination card, and all data were collected by recall alone. The reasons mentioned for partial and unvaccinated children were absenteeism of vaccinators, time inconvenience, not knowing the exact vaccination date, beliefs that vaccination has no use, fear of side effects, and religious and cultural issues.

## **Factors associated with full vaccination coverage**

In bivariate analysis, there were significant associations between full vaccination status and the type of respondent, level of education of mother/caregiver, maternal healthcare decision-making, number of ANC attendances, number of tetanus vaccinations, post-natal care attendance, having information on vaccination and vaccine-preventable diseases, aim for vaccination, number of vaccine-preventable diseases known by the respondent, age at which a child began vaccination, sessions needed to complete vaccination, age at which a child completed vaccination, and whether a vaccination made a child sick or not.

In multivariable analysis, maternal healthcare decisions made jointly with husbands [AOR = 1.88, 95% CI (1.06, 3.34)], maternal healthcare decision made by mothers [AOR = 4.03, 95% CI (1.66, 9.78)], number of post-natal care attendances [AOR = 5.02, 95% CI (2.28, 11.05)], age at which the child completed vaccination [AOR = 2.54, 95% CI (1.04, 6.23)], and vaccination not making the child sick [AOR = 0.32, 95% CI (0.16, 0.64)] were all significantly associated with full vaccination status (**Table 3**).

## Discussion

This study measured full vaccination coverage and associated factors among children aged 12-23 months in Demba Gofa District, Southern Ethiopia. In this study, only 47.0% of children were fully vaccinated. BCG, OPV3, pentavalent 3, PCV3, and Rota had a similar coverage of 88.7%, but measles coverage was only 59.4%, far below the others. Compared to the mini EDHS 2019 report, our results show slightly higher coverage (22) but equivalent coverage to results from Mizan Aman town (16) and Arba Minch Zuriya District (38) in Southern Ethiopia. However, our coverage was lower than those reported for North Gondar Lay Armachiho District, (33) and Sinana District, Southeast Ethiopia (36). Furthermore, coverage was very low compared to coverage reported for Tehulederie District, northeast Ethiopia (97%) (37), Debre Markos town, Amhara regional state (92%) (9), and studies conducted in Cameroon (47) and South Nigeria (23). The discrepancy may be due to differences in socioeconomic factors like residence and relatively poor healthcare systems as well as the settings in which the studies were conducted.

Different factors were examined with respect to associations with the full vaccination status. No socio-demographic characteristic was significantly associated with the full vaccination status of a child. This is in contrast to other studies conducted in different parts of Ethiopia (Arbegona district, South Ethiopia; Mizan Aman town, Bench Maji Zone, Southwest Ethiopia; Mecha District, Northwest Ethiopia; Jigjiga district, Somali national regional state, and Addis Ababa city (5, 16, 34, 35, 44)) which have shown that maternal age, maternal level of education, maternal occupation, and husbands' level of education were significantly associated with full vaccination coverage. A study from Bangladesh also showed that maternal age and maternal level of education were significantly associated with full vaccination coverage. The possible reasons for these differences may be due to differences in the study area, other sociodemographic characteristics of the participants, and the study settings.

We also found no associations between the children's characteristics and full vaccination coverage. However, studies from Arbegona District, Southern Ethiopia; Debre Markos town, Amhara regional state; Indonesia; Bangladesh; Sinana District, Southeast Ethiopia; and Addis Ababa town (5, 9, 13, 26, 36, 44) all showed that age, sex, and birth order of the children were significantly associated with full vaccination coverage.

Mothers who made healthcare decisions jointly with their husbands were 1.88 [95% CI (1.06, 3.34)] times more likely to vaccinate their children fully than when decisions were made by the husbands alone. Furthermore, mothers who made healthcare decisions themselves were 4.03 [AOR 95% CI(1.66, 9.78)] times more likely to fully vaccinate their children than when decisions were made by husbands alone. This is similar to a study from Indonesia (3) and indicates that maternal healthcare decisions made jointly with husbands and by mothers alone give mothers more freedom of health choices that benefit their children.

Mothers who attended post-natal care follow-up were 5.02 [AOR 95% CI (2.28, 11.05)] times more likely to fully vaccinate their children than mothers who did not. This is similar to a study from East-central Ethiopia (14). This may be because during postnatal care follow-up, mothers have a greater chance to obtain advice on vaccinations and exposure to the service that increases the probability of being fully vaccinated.

Respondents who replied that a child should complete vaccination by one year of age were 2.54 [95% CI (1.04, 6.23)] times more likely to fully vaccinate their children than those who did not know the exact age of vaccination completion. This is similar to studies from the North Gondar Lay Armachiho District, Northwest Ethiopia; Ambo District, Central Ethiopia; and Addis Ababa city (15, 41, 44). A possible explanation may be that having better knowledge on the timing of vaccination affected utilization of the service, as improved knowledge is expected to positively influence the uptake and utilization of a healthcare service.

Mothers who responded that vaccination will not make the child sick were less likely to vaccinate their children fully than those who thought that it would. Reasons given by respondents who did not complete vaccination were vaccinator absenteeism, a lack of awareness, time inconvenience, and not knowing the vaccination date. This was similar to a study from Laelay Adiabo District, Northern Tigray, Ethiopia (15). Furthermore, feeling that the vaccination had no use, a fear of side effects, religious and cultural refusals, a belief that vaccination hurts

the child, and a lack of awareness were reasons raised by mothers/caregivers who did not vaccinate their children at all. Similar reasons have been given in other studies. (9,14,15). The strength of this study are applying community-based study in a remote and rural part of the country and using a simple random sampling technique. However, recall bias of mothers could affect the study findings despite the efforts made to minimize them by asking the injection site, repetitions of the schedules, looking at the scars. Vaccine supply management, which could affect vaccination uptake, was not assessed.

## Conclusions

Less than half of children studied here were fully vaccinated and the prevalence of unvaccinated children was 11.3%, far below the government target of 90% necessary for sustained control of vaccine-preventable diseases. Maternal healthcare decision-making, postnatal care attendance, the objective of vaccination, the age at which a child begins and completes vaccination, and sessions needed to complete vaccination were significant factors associated with full vaccination. Absenteeism of vaccinators, a lack of awareness, time inconvenience, not knowing the exact date of vaccination, and local religious and cultural contexts were reasons given for not fully vaccinating their children. We advise that district health offices continuously conduct systematic supervision and periodic evaluation of vaccination performance of health facilities using a standardized checklist. Also, there should be an adequate supply of vaccination cards to health facilities (health centers and health posts). The health sector and other legal bodies are advised to improve maternal healthcare decision-making by empowering women on decision-making processes. Promoting institutional delivery and postnatal care services could improve full vaccination coverage. Health extension workers should emphasize explaining the specific names of vaccine-preventable diseases, the age at which a child begins and completes vaccination, and the number of sessions needed to complete vaccinations. Starting with senior management and extending to health extension workers, there is a need for improved planning of convenient times for the parent to attend vaccination, evaluating performance among target children, estimating the frequency of missed children, and providing the opportunity for re-attendance for defaulting children. The vaccine supply chain should also be studied further.

## Acknowledgements

We are grateful to Wolaita Sodo University, College of Health and Medical Science, School of Public Health for supporting this study. Our thanks extend to district health managers and staff for their support in facilitating the research process by timely approving and writing support letters. Finally, special thanks go to data collectors and study participants who contributed to this study

## References

1. Nancy PNaM. "Role(s) of vaccines and immunization programs in global disease control," Center for Strategic and International Studies. 2011.
2. UNICEF e. Integrating immunization and other services for women and children PMNCH Knowledge Summary 25. . 2013.
3. Putri Herliana AD. Determinants of immunization coverage of children aged 12–59 months in Indonesia. 25 August 2018.
4. WHO. Recommendations for routine immunization. 2015.
5. Abel Negussie WK, Sahilu Assegid and Ada K. Hagan. Factors associated with incomplete childhood immunization in Arbegona district , southern Ethiopia. BMC public health. 2016.
6. Ministry of Health SL. National Expanded Program on Immunization, Epidemiology Unit. 2012.
7. Gentile A. Pediatric disease burden and vaccination recommendations: understanding local differences. International journal of infectious diseases. [Review]. 2010.
8. Mouhamed Abdou Salam Mbengue, Aminata Mboup, Indou Deme Ly, Adama Faye, Fatou Bintou Niang Camara, Moussa, Thiam BPN, Tandakha Ndiaye Dieye, Souleymane Mboup. Vaccination coverage and immunization timeliness among children aged 12-23 months in Senegal. PanAfrican Medical Journal. 2017.
9. Dilie. TGaA. Vaccination Coverage and Associated Factors among Children Aged 12–23 Months in Debre Markos Town, Amhara Regional State, Ethiopia. Hindawi Advances in Public Health. 2017.
10. Federal Ministry of Health AA. Ethiopia National Expanded Program on Immunization. Comprehensive Multi-Year Plan 2016 – 2020. April, 2015.
11. Yemesrach A. Tefera ALW, Eyoel B. Mekonen, Bradley F. Carlson and Matthew L. Boulton. Predictors and Barriers to Full Vaccination among Children in Ethiopia. MDPI. 2018.
12. Md Illias Kanchan Sk MFS, Prashika Rajendra Kurlikar, Mithlesh Chourase, Rajaram Yadav, Abhijit Basu Biswas. Status and determinants of child immunisation coverage in three South Asian countries, India, Bangladesh and Nepal: Evidence from the Demographic and Health Survey. Sri Lanka Journal of Child Health. 2018.
13. Holipah AMaYK. Determinants of immunization status among 12- to 23-month-old children in Indonesia (2008–2013). BMC public health. 2018.
14. Melaku Kindie Yenit YAGaAMS. Mothers' health service utilization and attitude were the main predictors of incomplete childhood vaccination in east-central Ethiopia. Archives of Public Health. 2018.
15. Hailay Gebretnsae Aregawi TGG, Yamane Gebremariam Abebe, Kidanu Gebremariam Meles, Alem Desta Wuneh. Determinants of defaulting from completion of child immunization in Laelay Adiabo district , Tigray Region, Northern Ethiopia. PLOS ONE. 2017.

16. Asrat Meleko MG, and Frehiwot Birhanu. Assessment of Child Immunization Coverage and Associated Factors with Full Vaccination among Children Aged 12–23 Months at Mizan Aman Town, Bench Maji Zone , Southwest Ethiopia. *Hindawi International Journal of Pediatrics*. 2017.
17. Nepal Development Society B, Chitwan, Nepal, Centre for International Health, University of Bergen, Bergen, Norway, Department of Nutrition, School of Public Health, University of Kinshasa, Kinshasa, Democratic Republic of Congo, Fafo, Institute for Labour and Social Research, Oslo, Norway. Individual- and community-level determinants of child immunization in the Democratic Republic of Congo. *PLOS ONE*. 2018.
18. Didier K. Ekouevi FAG-K, Issifou Yaya, Wendpouiré I. Zida-Compaor,, Amevegbé Boko ES, Anani Lacle, Nicolas Ndibu, Yaovi Toke and Dadja E. Landoh. Incomplete immunization among children aged 12–23 months in Togo: a multilevel analysis of individual and contextual factors. *BMC public health*. 2018.
19. Aschalew Teka Bekele, Braka Fiona, Karengera Thomas, Aron Kassahun, Gallagher Kathleen, Peter Nsubuga, Yohannes, Ababu AL. Factors contributing to routine immunization performance in Ethiopia. *PanAfrican Medical Journal*. 2017.
20. Kristin VanderEnde PMG-D, MSc; Mamadou S. Diallo, PhD; Laura M. Conklin, MD; Aaron S. Wallace, PhD. Global routine vaccination coverage, *MMWR /November 21/2013*. ResearchGate. November 16, 2018.;67.
21. Central Statistical Agency:Addis Ababa E. Central Statistical Agency. Ethiopia Demographic and Health Survey2016. Addis Ababa, Ethiopia. July 2017.
22. Central Statistical Agency:Addis Ababa E. Mini Ethiopian Demographic and Health Survey. July 2019.
23. Adeleye O.A MN. Determinants of Full Vaccination Status in a Rural Community with Accessible Vaccination Services in South-South Nigeria. *JOURNAL OF COMMUNITY HEALTH and PRIMARY HEALTH CARE*. September 2016;27.
24. Abiyemi Benita Awoh EP. Immunisation coverage in rural–urban migrant children in low and middle-income countries (LMICs). 2015.
25. MMWR UDoHaHSCfDCaP. Global Routine Vaccination Coverage, Morbidity and Mortality Weekly Report 2017.
26. Nurnabi Sheikh MS, Nausad Ali, Raisul Akram, Rashidul Alam Mahumud, Muhammad Asaduzzaman and Abdur Razzaque Sarker. Coverage, Timelines, and Determinants of Incomplete Immunization in Bangladesh. *MDPI Tropical Medicine and Infectious Diseas*. 2018.
27. Fraidoon Farzad JAR, Eiko Yamamoto and Nobuyuki Hamajima. Socio-economic and demographic determinants of full immunization among children of 12–23 months in Afghanistan. 2017.
28. Enakshi Ganguly M, Rahul Gupta, MStat, Alik Widge, MD, PhD , R. Purushotham Reddy, MBA, K. Balasubramanian, PhD, and P. S. Reddy, MD. Increasing Full Child Immunization Rates by Government Using an Innovative Computerized Immunization Due List in Rural India. *The Journal of Health Care Organization, Provision, and Financing*. 2018;55.
29. Richard Mihigo JO, and Blanche Anya, Pascal Mkanda, Felicitas Zawaira. Challenges of immunization in the African region. *PanAfrican Medical Journal*. 2017.
30. Martin Nyaaba Adokiya BBaJAN. Evaluation of immunization coverage and its associated factors among children 12–23 months of age in Techiman Municipality, Ghana. *Archives of Public Health*. 2017.

31. Elizabeth B. Adedire IA, Olufunmilayo I. Fawole, Olufemi Ajumobi, Simon Kasasa, Peter Wasswa and Patrick Nguku. Immunisation coverage and its determinants among children aged 12-23 months in Atakumosa-west district , Osun State Nigeria. BMC public health. 2017.
32. Shellese A. Shemwella MBP, b, Lázaro González-Calvoa,c, Montse Renom-Llonchd, Alexandre Boond, Samuel Martinhoc, Charlotte B. Cherrya, Ann F. Greena,c and Troy D. Moona. Determinants of full vaccination status in children aged 12–23 months in Gurùé and Milange district s, Mozambique. 2017.
33. Melkamu Beyene Kassahun GABaAST. Level of immunization coverage and associated factors among children aged 12–23 months in Lay Armachiho district, North Gondar Zone , Northwest Ethiopia. BMC public health. 2015.
34. Ayal Debie BT. Assessment of full vaccination coverage and associated factors among children aged 12-23 months in Mecha District , North West Ethiopia. Science Publishing Group. 2014.
35. Abdi Nur Mohamud AF, Walelegn Worku, Manay Kifle and Hardeep Rai Sharma. Immunization coverage of 12–23 months old children and associated factors in Jigjiga district , Somali National Regional State, Ethiopia. BMC public health. 2014.
36. Dechasa ELaW. An assessment of child immunization coverage and its determinants in Sinana district , Southeast Ethiopia. BMC Pediatr. 2015.
37. Toyeb Yasine Ebrahim and WBS. childhood immunization coverage in tehulederie district, northeast of Ethiopia. International journal of current research. september, 2015;7.
38. Facha W. Full vaccination Coverage and Associated Factors among Children Aged 12 to 23 Months in Arba Minch Zuriya Woreda, Southern Ethiopia. Journal of Pharmacy and Alternative Medicine. 2015;vol.7.
39. Tahir Yousuf Nour , Alinoor Mohamed Farah, et al. Immunization coverage in Ethiopia among 12–23 month old children: systematic review and meta-analysis. BMC Public Health. (2020) 20:1134.
40. Patience Esohe Konwea FADaSEO. Determinants of compliance with child immunization among mothers of children under five years of age in Ekiti State, Nigeria. Emerald Insight. 2018.
41. Etana B, Deressa W. Factors associated with complete immunization coverage in children aged 12-23 months in Ambo Woreda, Central Ethiopia. BMC public health. 2012;12:566.
42. Abdirisak Mohamed Abdi PSKaDRUY. Barriers to Full Immunization Coverage of Under Five Years Children in Benadir Region, Somalia. Journal of Chemical, Biological and Physical Sciences. 2016;6.
43. Asamnew Zewdie MLaTM. Reasons for defaulting from childhood immunization program: from Hadiya Zone , Southern Ethiopia. BMC public health. 2016.
44. Shiferaw Birhanu AA, Yezabnesh Kibie, Ayalew Jejaw. Knowledge, Attitude and Practice of Mothers Towards Immunization of Infants in Health Centres at Addis Ababa, Ethiopia. 2016:1.
45. Office DGDH. Demba Gofa district health office health profile. 2018.
46. Elizabeth Kiptoo\* ME, George Kobia and Raphael Ngure. Factors Influencing Low Immunization Coverage Among Children Between 12 - 23 Months in East Pokot, Baringo Country, Kenya. International Journal of Vaccines and Vaccination. 2015;1(2).

442 47. Didier K. Fifonsi A. Issifou Y et al. Vaccine coverage and determinants of incomplete  
 443 vaccination in children aged 12–23 months in Dschang, West Region, Cameroon: a cross-  
 444 sectional survey during a polio outbreak. BMC Public Health. 2015;15:630.

445

446



**Table 1**(on next page)

Socio-demographic & economic characteristics of mothers/caregivers in Demba Gofa district, Southern Ethiopia, 2019 (N=657).

1 Table 1. Socio-demographic & economic characteristics of mothers/caregivers in Demba Gofa  
2 district, Southern Ethiopia, 2019 (N=657).

Variables	Category	Frequency (#)	Percent (%)
Respondent	Mother	632	96.2
	Caregiver	25	3.8
Age of mother/caregiver (yrs)	15-24	178	27.1
	25-34	331	50.4
	≥35	148	22.5
Religion	Protestant	511	77.8
	Orthodox	138	21.0
	Muslim	8	1.2
Marital status	Married (in union)	634	96.5
	Separated	12	1.8
	Divorced	4	0.6
	Others (not married)	7	1.1
Level of education of mothers /caregivers	Illiterate	260	39.6
	Can read & write	43	6.5
	Primary school (1-8)	173	26.3
	Secondary & pre. (9-12)	120	18.3
	Diploma & above	61	9.3
Occupation of mothers/caregivers	Housewife	548	83.4
	Merchant	67	10.2
	Student	22	3.3
	Gov't employee	12	1.8
	Daily laborer	3	0.5
	Others	5	0.8
Husbands' level of education	Illiterate	211	32.1
	Can read & write	51	7.8
	Primary school (1-8)	159	24.2
	Secondary & pre. (9-12)	145	22.1
	Diploma & above	81	12.3
Occupation of father	Farmer	497	75.6
	Merchant	68	10.4
	Student	24	3.7
	Gov't employee	40	6.1
	Daily laborer	10	1.5
	Others	8	1.2
Wealth index	Poorest	132	20.1
	Poorer	131	19.9
	Middle	132	20.1
	Richer	131	19.9
	Richest	131	19.9
	≤3	109	16.6

Number of people living in HHs	4-5	238	36.2
	$\geq 6$	310	47.2
Sex of child	Male	376	57.2
	Female	281	42.8
Birth order of last child	One	120	18.3
	Two-three	118	18.0
	Four-five	195	29.7
	Above five	224	34.1
Age of the child (in months)	12-14	222	33.8
	15-17	156	23.7
	18-20	147	22.3
	21-23	132	20.1

3

# **Table 2**(on next page)

Table 2. Maternal healthcare service utilization in Demba Gofa District, Southern Ethiopia, 2019 (N=657).

1 Table 2. Maternal healthcare service utilization in Demba Gofa District, Southern Ethiopia, 2019  
2 (N=657).

Variables	Category	#	%
Maternal healthcare decision making	By herself	84	12.8
	Jointly with their husband	473	72.0
	Husband alone	100	15.2
ANC attendance	Yes	551	83.9
	No	106	16.1
Number of ANC attendance	Once	28	5.1
	Twice	118	21.4
	Three times	193	35.0
	Four times	212	38.5
TT vaccination	Yes	490	88.9
	No	61	11.1
TT vaccination status	TT 1	79	16.1
	TT 2	257	52.4
	TT 3	113	23.1
	TT 4	41	8.4
Delivery place of last baby	Home	227	34.6
	Health institution	430	65.4
PNC attendance	Yes	63	9.6
	No	594	90.4
PNC	Once	36	57.1
	Twice	26	41.3
	Three times	1	1.6

3

4

# **Table 3**(on next page)

Table3. Factors associated with full vaccination coverage of children aged 12-23 months in Demba Gofa district, southern Ethiopia, 2019, (N=657).

1 Table3. Factors associated with full vaccination coverage of children aged 12-23 months in  
2 Demba Gofa district, southern Ethiopia, 2019, (N=657).

Variables	Category	Fully vaccination		Odds Ratio (95% CI)	
		Yes	No	COR	AOR
Respondent	Mother	289 (93.5)	343 (98.6)	0.21 (0.08, 0.57)	0.54 (0.13, 2.12)
	Caregiver	20 (6.5)	5 (1.4)	1	1
Level of education of mother/caregiver	Illiterate	114 (36.9)	146 (42.0)	1	1
	Read & write	29 (9.4)	14 (4.0)	2.65 (1.34, 5.25)	2.00 (0.72, 5.54)
	Primary school (1-8)	75 (24.3)	98 (28.2)	0.98 (0.66, 1.44)	1.03 (0.60, 1.75)
	Secondary & prep. (9-12)	57 (18.4)	63 (18.1)	1.16 (0.75, 1.78)	0.99 (0.57, 1.73)
	Diploma & above	34 (11.0)	27 (7.8)	1.61 (0.92, 2.82)	0.87 (0.41, 1.81)
Maternal healthcare decision making	Jointly with husband	227 (73.5)	246 (70.7)	1.71 (1.09, 2.68)	<b>1.88 (1.06, 3.34) *</b>
	By herself	47 (15.2)	37 (10.6)	2.36 (1.30, 4.27)	<b>4.03 (1.66, 9.78) *</b>
	Husband alone	35 (11.3)	65 (18.7)	1	1
Number of ANC attendance	Once	9 (3.2)	19 (7.0)	1	1
	Twice	40 (14.4)	78 (28.6)	1.08 (0.45, 2.61)	1.26 (0.31, 5.07)
	Three times	111 (39.9)	82 (30.0)	2.85 (1.23, 6.64)	2.64 (0.66, 10.43)
	Four times	118 (42.2)	94 (34.4)	2.65 (1.14, 6.12)	2.22 (0.54, 9.00)
Number of TT vaccination	1	30 (11.6)	49 (21.1)	1	1
	2	133 (51.6)	124 (53.4)	1.75 (1.04, 2.93)	1.31 (0.70, 2.47)
	3	71 (27.5)	42 (18.1)	2.76 (1.52, 4.99)	2.00 (0.94, 4.23)
	4	24 (9.3)	17 (7.3)	2.30 (1.06, 4.98)	0.99 (0.33, 2.98)
PNC attendance	Yes	46 (14.9)	17 (4.9)	3.40 (1.90, 6.08)	<b>5.02 (2.28,11.05)*</b>
	No	263 (85.1)	331 (95.1)	1	1
Information on vaccination & VPDs	Yes	302 (97.7)	307 (88.2)	0.17 (0.07, 0.39)	0.71 (0.16, 3.11)
	No	7 (2.3)	41 (11.8)	1	1
Objective of vaccination	To prevent disease	235 (76.1)	209 (60.1)	4.09 (2.04, 8.17)	3.44 (0.85, 13.81)
	For child health	63 (20.4)	99 (28.4)	2.31 (1.10, 4.84)	1.88 (0.47, 7.52)
	Do not know	11 (3.6)	40 (11.5)	1	1
Number of VPDs known by the respondent	Single disease	90 (29.1)	115 (33.0)	1.48 (0.99, 2.19)	0.90 (0.51, 1.60)
	More than one disease	145 (46.9)	93 (26.7)	2.95 (2.01, 4.33)	<b>1.82 (0.99, 3.34) *</b>
	Do not know	74 (23.9)	140 (40.2)	1	1
Age at which a child begins vaccination	Just after birth	155 (50.2)	132 (37.9)	4.69 (2.26, 9.75)	0.82 (0.19, 3.60)
	One month after birth	144 (46.6)	176 (50.6)	3.27 (1.58, 6.77)	0.65 (0.14, 2.87)
	Do not know	10 (3.2)	40 (11.5)	1	1
Sessions needed to complete vaccination	Three & less	48 (15.5)	64 (18.4)	1.72 (0.97, 3.04)	0.42 (0.16, 1.11)
	Four or five	212 (68.6)	183 (52.6)	2.66 (1.66, 4.27)	0.74 (0.33, 1.67)
	Six & above	19 (6.1)	32 (9.2)	1.36 (0.67, 2.78)	0.34 (0.12, 1.00)
	Do not know	30 (9.7)	69 (19.8)	1	1
Age to complete vaccination	Before one year	192 (62.1)	202 (58.0)	2.51 (1.55, 4.05)	0.92 (0.40, 2.09)
	One year & above	89 (28.8)	72 (20.7)	3.26 (1.91, 5.57)	<b>2.54 (1.04, 6.21) *</b>
	Do not know	28 (9.1)	74 (21.3)	1	1
Vaccination will not make child to sick	Yes	275 (89.0)	280 (80.5)	1.96 (1.26, 3.06)	<b>0.32 (0.16, 0.64) *</b>
	No	34 (11.0)	68 (19.5)	1	1

3 **Key** \* p-value<0.05, COR=Crude odds ratio, AOR=Adjusted odds ratio