

# Genome-wide identification of the *SWEET* gene family mediating the cold stress response in *Prunus mume*

Zhenying Wen<sup>1,2</sup>, Mingyu Li<sup>1,2</sup>, Juan Meng<sup>1,2</sup>, Ping Li<sup>1,2</sup>, Tangren Cheng<sup>1,2</sup>, Qixiang Zhang<sup>1,2</sup>, Lidan Sun<sup>Corresp. 1,2</sup>

<sup>1</sup> Beijing Key Laboratory of Ornamental Plants Germplasm Innovation & Molecular Breeding, School of Landscape Architecture, Beijing Forestry University, Beijing, China

<sup>2</sup> National Engineering Research Center for Floriculture, Beijing Laboratory of Urban and Rural Ecological Environment, Beijing Forestry University, Beijing, China

Corresponding Author: Lidan Sun  
Email address: sld656@126.com

The *SWEET* (Sugars Will Eventually be Exported Transporter) gene family encodes a family of sugar transporters that plays essential roles in plant growth, reproduction, and biotic and abiotic stresses. *Prunus mume* is a considerable ornamental wood plant with high edible and medicinal values; however, low temperature severely limits its geographical distribution. To investigate whether this gene family mediates *P. mume*'s response to cold stress, we identified its 17 *SWEET* genes from *P. mume* and divided them members into four groups. Sixteen of these genes were anchored on six chromosomes, and one gene was anchored on the scaffold with four pairs of segmental gene duplications and two pairs of tandem gene duplications. *Cis*-acting regulatory element analysis indicated that the *PmSWEET* genes are potentially involved in the *P. mume* developmental procedure, such as circadian control, abscisic acid-response and light-response, and responses to numerous stresses, such as low-temperature and drought. We performed low-temperature treatment in the cold-tolerant cultivar 'Songchun' and cold-sensitive cultivar 'Zaolve' and found that the expression of four of 17 *PmSWEETs* was either upregulated or downregulated with prolonged treatment times, which indicates that these family members may potentially play a role in cold stress responses in *P. mume*. Our study provides a basis for further investigation of the role of *SWEET* proteins in the development of *P. mume* and its responses to cold stress.

# Genome-wide identification of the *SWEET* gene family mediating the cold stress response in *Prunus mume*

Zhenying Wen<sup>1,2</sup>, Mingyu Li<sup>1,2</sup>, Juan Meng<sup>1,2</sup>, Ping Li<sup>1,2</sup>, Tangren Cheng<sup>1,2</sup>, Qixiang Zhang<sup>1,2</sup>, Lidan Sun<sup>1,2\*</sup>

<sup>1</sup>Beijing Key Laboratory of Ornamental Plants Germplasm Innovation & Molecular Breeding, School of Landscape Architecture, Beijing Forestry University, Beijing 100083, China

<sup>2</sup>National Engineering Research Center for Floriculture, Beijing Laboratory of Urban and Rural Ecological Environment, Beijing Forestry University, Beijing 100083, China

## Corresponding Author:

Lidan Sun<sup>1,2</sup>

No. 35 Qinghua East Road Haidian District, Beijing, 100083, China

Email address: [sunlidan@bjfu.edu.cn](mailto:sunlidan@bjfu.edu.cn)

## Abstract

The SWEET (Sugars Will Eventually be Exported Transporter) gene family encodes a family of sugar transporters that plays essential roles in plant growth, reproduction, and biotic and abiotic stresses. *Prunus mume* is a considerable ornamental wood plant with high edible and medicinal values; however, low temperature severely limits its geographical distribution. To investigate whether this gene family mediates *P. mume*'s response to cold stress, we identified its 17 *SWEET* genes from *P. mume* and divided them members into four groups. Sixteen of these genes were anchored on six chromosomes, and one gene was anchored on the scaffold with four pairs of segmental gene duplications and two pairs of tandem gene duplications. *Cis*-acting regulatory element analysis indicated that the *PmSWEET* genes are potentially involved in the *P. mume* developmental procedure, such as circadian control, abscisic acid-response and light-response, and responses to numerous stresses, such as low-temperature and drought. We performed low-temperature treatment in the cold-tolerant cultivar 'Songchun' and cold-sensitive cultivar 'Zaolve' and found that the expression of four of 17 *PmSWEETs* was either upregulated or downregulated with prolonged treatment times, which indicates that these family members may potentially play a role in cold stress responses in *P. mume*. Our study provides a basis for further investigation of the role of *SWEET* proteins in the development of *P. mume* and its responses to cold stress.

**Keywords:** *Prunus mume*, SWEET gene family, expression pattern, cold response.

## 1. Introduction

Sucrose is the main carbohydrate in most plants; it is synthesized in the leaves during photosynthesis and then transported by phloem sap to storage organs, such as roots, stems, flowers, seeds and fruits (Rennie and Turgeon, 2009; Lemoine et al., 2013). Sucrose provides energy and carbon sources for plants and acts as an important signal and resistance molecule that participates in the normal growth of higher plants (Chen et al., 2015). However, these sugars must be assisted by appropriate sugar transporters as it cannot be transported independently to the storage organs (Ainsworth and Bush, 2011). At present, three transporter families have been identified as essential sugar transporters: monosaccharide transporters (MSTs), sucrose transporters (SUTs), and Sugar Will Eventually be Exported transporters (SWEETs) (Chen et al., 2010; Chen et al., 2015; Eom et al., 2015). Of these three families, *SWEETs* were the final gene family to be uncovered and were first identified by Chen et al. in *Arabidopsis* (Chen et al., 2010). SWEET proteins act as sugar transporters that mediate the inflow or outflow of phloem parenchyma sugar into the phloem apoplast (Slewiniski, 2011; Braun, 2012; Chen, 2014). Unlike the SUT and MST families, which require energy to transport sugar across the plasma membrane

(Maynard and Lucas, 1982; Lemoine, 2000), SWEET proteins promote the diffusion of sugar across concentration gradients at the cellular membrane or vacuolar membrane, regardless of the proton gradient or pH of the cellular environment (Chen et al., 2012; Chen et al., 2015).

SWEET proteins are characterized by conserved MtN3\_saliva (MtN3\_slv) transmembrane (TM) domains (Chen et al., 2012), also known as PQ-loop repeats (Eom et al., 2015; Feng and Frommer, 2015). SWEETs in eukaryotes commonly consist of seven transmembrane helices (TMHs), which contain a pair of 3-TMH repeats detached by an added helix (Xuan et al., 2013), and this structure has been described as the “3-1-3” TM SWEET structure (Chen et al., 2010). In contrast to the structure of eukaryote SWEET proteins, prokaryote SWEET proteins, known as SemiSWEETs, are composed of only three TMHs (Xuan et al., 2013). In eukaryotes, proteins that contain 6 or 7 TMHs are prevalent, but SemiSWEETs with 3 or 4 TMHs have also been detected in plant genomes. In a study of SWEET genes from 25 plant genomes, 140 of the 411 SWEET sugar transporters identified were semiSWEET; with all of the identified semiSWEETs either lacking the first or second 3-TM domain or exist only in partial form (Patil et al., 2015). This data therefore demonstrates that the presence of semiSWEETs in higher plant genomes is not unusual, and further, that SWEETs may in actual fact have formed by direct fusion from SemiSWEETs (Jia et al., 2017). In addition, a novel extraSWEET protein consisting of 14 and 15 TMHs has been reported from *Vitis vinifera* (Patil et al., 2015) and *Oryza punctata* (Jia et al., 2017); it is speculated that this extraSWEET may have formed from the duplication of a 7 TMH SWEET gene in these two species. Recent research on 3,249 SWEET proteins also identified a superSWEET with > 18 TMHs in oomycetes, which carry 5–8 repeats of a semiSWEET (Jia et al., 2017). According to phylogenetic analysis, the SWEET genes in *Arabidopsis* can be divided into four clades: Clade I (SWEET1–3) and Clade II (SWEET4–8) mainly transport glucose, while Clade I also transports hexose (Chen et al., 2010; Lin et al., 2014). Clade III members (SWEET9–15) mainly transports sucrose (Chen et al., 2012; Eom et al., 2015), and Clade IV members (SWEET16–17), which are located on the tonoplast membrane, mainly transports fructose (Eom et al., 2015). The phylogenetic relationships of the SWEET genes described hereafter are all based on results from *Arabidopsis*.

Advances in whole-genome sequencing have enabled genome-wide identification of SWEET genes in numerous species. These include important crops, fruits and vegetables, such as rice (*Oryza sativa*) (Yuan and Wang, 2013), sorghum (*Sorghum bicolor*) (Mizuno et al., 2016), soybean (*Glycine max*) (Patil et al., 2015), apple (*Malus domestica*) (Wei et al., 2014), grape (*Vitis vinifera*) (Chong et al., 2014), banana (*Musa acuminata*) (Miao et al., 2017), tomato (*Solanum lycopersicum*) (Feng et al., 2015), rapeseed (*Brassica napus*) (Jian et al., 2016), potato (*Solanum tuberosum*) (Li et al., 2020) and valencia sweet orange (*Citrus sinensis*) (Yao et al., 2021). Additionally, many SWEET genes have been confirmed to play diverse and complex roles in physiological processes, such as nectar secretion (Ge et al., 2000; Lin et al., 2014),

pollen development (Sun et al., 2013), senescence (Quirino et al., 1999), and seed filling (Sosso et al., 2015). Moreover, SWEET genes are also involved in biotic and abiotic stress responses (Yuan and Wang, 2013), including the reaction of plants to stress at low temperatures. For example, overexpression of *AtSWEET16* and *AtSWEET17* increases cold tolerance (Chardon et al., 2013; Klemens et al., 2013; Guo et al., 2014); overexpression of *AtSWEET4* increases plant size and frost resistance (Chong et al., 2014; Liu et al., 2016); and *AtSWEET11* and *AtSWEET12* are involved in responses to stress caused by cold or dehydration (Le Hir et al., 2015; Durand et al., 2016). *AtSWEET15* is also known as SAG29 (where SAG stands for senescence-associated gene); however, its transcription level gradually increases at low temperature, high salinity, and drought during natural leaf senescence (Quirino et al., 1999). Cold stress significantly inhibits the expression of *CsSWEET2*, *CsSWEET3*, and *CsSWEET16* in *Camellia sinensis*, while the expression of *CsSWEET1* and *CsSWEET17* increases sharply (Yue et al., 2015). A functional study of *CsSWEET16* in *C. sinensis* revealed that it is located in the vacuolar membrane and regulates cold resistance in transgenic *Arabidopsis* plants (Wang et al., 2018). The transcriptional activity of many *SISWEET* genes increases under low-temperature stress in tomato (Feng et al., 2015). Studies have shown that expression of the *MaSWEET* gene in banana is upregulated in response to low temperature, salt, and osmotic stress (Miao et al., 2017). Using genome-wide analysis of the *BoSWEET* gene in *Brassica oleracea* var. *capitata*, five possible candidate genes were found to promote sugar transport and thereby enhance chilling tolerance in cabbage (Zhang et al., 2019).

*Prunus mume* is a traditional flower native to southwest China and the middle and lower reaches of the Yangtze River. In the northern China, low temperatures severely limit the growth and distribution of this species. Although SWEET sugar transporters have been associated with responses to cold in other species, little is known about the role of *PmSWEETs* in cold responses in *P. mume*. This study aims to conduct a genome-wide analysis of the SWEET gene family in *P. mume*, with a specific focus on SWEET gene transcriptional responses to cold stress, providing a starting point to study the detailed role of *PmSWEETs*.

## 2. Materials and Methods

### 2.1 Plant Genomic Resources

To explore the phylogeny of the SWEET genes in *P. mume* and other species, we downloaded SWEET proteins from two model plants (*Arabidopsis thaliana* and *Oryza sativa*, representing dicotyledons and monocotyledons, respectively) and eight other Rosaceae species. The protein sequences of 17 *AtSWEETs* and 21 *OsSWEETs* were downloaded from the TAIR 10 database (<http://www.arabidopsis.org/>) and TIGR (<http://rice.plantbiology.msu.edu/>), respectively. The *P. mume* genome sequence and annotation files were obtained from the *P. mume* genome project (<http://prunusmumegenome.bjfu.edu.cn/>); the genomes of eight other Rosaceae species, *Malus*

*domestica* (Daccord et al., 2017), *Prunus avium* (Shirasawa et al., 2017), *Prunus persica* (Verde et al., 2013), *Prunus yedoensis* (Baek et al., 2018), *Pyrus communis* (Linsmith et al., 2019), *Rosa chinensis* (Raymond et al., 2018), *Prunus salicina* (Liu et al., 2020), and *Prunus armeniaca* (Jiang et al., 2019), were downloaded from the Genome Database for Rosaceae (<https://www.rosaceae.org/>).

## 2.2 Identification of *SWEET* Genes in *P. mume* and Other Species

The hidden Markov model (HMM) profiles of the MtN3\_slv domain for the *SWEET* gene family (PF03083) were downloaded from the Pfam database (<http://pfam.xfam.org/>) and used as queries to search for *SWEET* proteins in the proteomes of *P. mume* and other species with HMMER software (version 3.1b2, <http://hmmer.org/>) (Finn et al., 2015). To ensure confidence, the E-value cutoff was set at  $10^{-5}$ . Then, all putative *SWEET* proteins were screened to confirm the presence of the MtN3\_slv domain by SMART (<http://smart.embl-heidelberg.de/>), the Pfam database (<http://pfam.xfam.org/>) and NCBI-CDD (<https://www.ncbi.nlm.nih.gov/cdd>), and sequences with MtN3\_slv domain were retained.

The *SWEET* genes were named based on their location information in the genome. In addition, the number of amino acids, molecular weight (MW) and isoelectric point (pi) were calculated using the online ExPASy program (<https://web.expasy.org/cgi-bin/protparam/protparam>). The distributions of TM helices were predicted by TMHMM Server v. 2.0 (<http://www.cbs.dtu.dk/services/TMHMM/>).

## 2.3 Phylogenetic and Conserved Domain Analysis

To examine the phylogeny between *SWEET* genes in *P. mume* and other species, alignment of full-length *SWEET* protein sequences from three species (*P. mume*, *A. thaliana*, and *O. sativa*) and eight Rosaceae species was performed by using MAFFT software with the FFT-NS-1 strategy (Katoh and Standley, 2013). Subsequently, maximum likelihood (ML) phylogenetic trees were constructed using FastTree (version 2.1.11) (Price et al., 2010) with default parameters. Then, iTOL v4.0 (<https://itol.embl.de/itol.cgi>) (Letunic and Bork, 2019) and AI CS6 software were used to annotate and embellish the phylogenetic tree.

## 2.4 Conserved Motif and Gene Structure Analysis

The conserved motifs of *PmSWEETs* were predicted by MEME Suite Version 5.3.3 (<https://meme-suite.org/meme/tools/meme>) (Bailey et al., 2009), where the maximum number of motifs for the conserved domains was set to 10, motif width was set to 6-50, and the residuals were designated as the default parameters. Gene structure data was extracted from the *P. mume* genome gff file, visualized using TBtools software (Chen et al., 2020), and then edited in AI CS6 software.

## 2.5 Chromosome Location, Duplication and Synteny Analysis

The location and chromosome length information of *PmSWEETs* was obtained from the gff file downloaded from the *P. mume* genome project (<http://prunusmumegenome.bjfu.edu.cn/>). A chromosomal location figure was drawn using the online tool MG2C ([http://mg2c.iask.in/mg2c\\_v2.0/](http://mg2c.iask.in/mg2c_v2.0/)). Gene tandem and segment replication events were analyzed using the Multiple Collinearity Scan Toolkit (MCScanX) and Circos in TBtools, respectively, with the default parameters. The synteny of the *PmSWEETs* across *A. thaliana*, *P. armeniaca*, and *P. salicina* was mapped using MCScanX in TBtools. The Ks and Ka values for duplicated gene pairs were calculated based on the coding sequence alignments using the Ka/Ks calculator in TBtools. According to two ordinary rates ( $\lambda$ ) of  $1.5 \times 10^{-8}$  and  $6.1 \times 10^{-9}$  substitutions per site per year (Lynch and Conery, 2000; Blanc and Wolfe, 2004), the formula  $t = Ks/2\lambda \times 10^{-6}$  Mya was used to calculate the divergence time.

## 2.6 Cis-Acting Element Analysis of *PmSWEET* Gene Promoter Regions

The upstream sequences (2.0 kb) of the *PmSWEETs* were retrieved from the genomic sequence data in TBtools and then submitted to the PlantCARE database (<http://bioinformatics.psb.ugent.be/webtools/plantcare/html/>) (Lescot et al., 2002) for *cis*-acting element analysis. We finally selected 12 elements, including those induced by hormones, such as methyl jasmonate (MeJA)-responsive, abscisic acid (ABA)-responsive, and stress-responsive elements; the stress-responsive factors included those involved in defense and stress, low temperature, and light. By combining these data with phylogenetic tree information (nwk file), the map was constructed by TBtools and edited by AI CS6 software.

## 2.7 *PmSWEET* Genes Expression Analysis

To investigate the function of *PmSWEETs* involved in tissue development and cold tolerance, we used root, stem, leaf, bud and fruit data from RNA sequencing (Zhang et al., 2012) to analyze the *PmSWEET* expression patterns in different tissues and then used flower bud dormancy data from RNA sequencing of *P. mume* ('Zaolve') (Zhang et al., 2018) to analyze *PmSWEET* responses to low temperature from November to February. Furthermore, we explored the expression of SWEET gene family members in the stem of *P. mume* ('Songchun') in geographically distinct locations, including Beijing (BJ, N39°54', E116°28'), Chifeng (CF, N42°17', E118°58') and Gongzhuling (ZGL, N43°42', E124°47') and for three different periods of the year, including cold acclimation (October, autumn), the final period of endo-dormancy (January, winter), and deacclimation (March, spring) (Jiang, 2020). TBtools (Chen et al., 2020) was used to create the heatmap.

## 2.8 qRT-PCR Analysis of *PmSWEET* Genes

To examine the response of *PmSWEET* to low temperature, the annual branches of the cold-sensitive cultivar ‘Zaolve’ and the cold-tolerant cultivar ‘Songchun’ were collected. Before chilling treatment, the shoots were incubated overnight at 22 °C and then transferred to 4 °C for 0, 1, 4, 6, 12, 24, 48, and 72 h under long-day conditions (16-h light/8-h dark). The stems were collected immediately stored in liquid nitrogen until their longterm storage at -80 °C in readiness for RNA extraction. Each treatment had three biological replicates.

Total RNA of each sample was extracted using the RNAPrep Pure Plant Plus Kit (Tiangen, Beijing, China). Complementary cDNA was synthesized using ReverTra Ace® qPCR RT Master Mix with gDNA Remover (Toyobo, Osaka, Japan). The specific primers were designed by Primer 3 (<https://bioinfo.ut.ee/primer3-0.4.0/>) based on the cDNA sequences (Table S1). The expression levels of *PmSWEETs* at low temperature were analyzed using quantitative real-time polymerase chain reaction (qRT-PCR) with a PikoReal real-time PCR system (Thermo Fisher Scientific, CA, USA) with SYBR® Green Premix *Pro Taq* HS qPCR kit (Accurate biology, China). The reactions were performed in a 10 µL volume, including 5.0 µL SYBR®Green Premix *Pro Taq* HS qPCR master mix, 0.5 µL each of forward and reverse primers, 1.0 µL of cDNA and 3.0 µL of ddH<sub>2</sub>O. The reactions were performed according to the following procedure: 95 °C for 30 s, followed by 40 cycles of 95 °C for 5 s and 60 °C for 30 s. Via the use of the phosphatase 2A gene of *P. mume* as the reference gene, the relative expression was calculated by using the formula  $2^{-\Delta\Delta C_t}$  method (Livak and Schmittgen, 2001). Each real-time qRT-PCR was conducted in three biological replicates. The statistical analyses of ‘Zaolve’ and ‘Songchun’ were independent carried out using SPSS22.0, the one-way ANOVA analysis of variance was calculated by least significant difference (LSD) and Student-Newman-Keuls test with significant difference at level  $p = 0.05$ . GraphPad Prism6 software was used to draw the diagram.

### 3. Results

#### 3.1 Identification of Members of the *Prunus mume* SWEET Gene Family

A total of 17 nonredundant *PmSWEETs* were detected in the *P. mume* genome (sequence information is shown in Supplement File S1), and 175 SWEETs were detected in the eight other species of Rosaceae, including 16 *SWEET* genes in *P. armeniaca*, 19 in *P. avium*, 19 in *P. persica*, 19 in *P. salicina*, 16 in *P. yedoensis*, 21 in *P. communis*, 29 in *M. domestica*, and 36 in *R. chinensis* with rigorous filtering. All the newly identified SWEET genes were named according to their chromosome location (Table 1 and Table S2). We determined that candidates with at least one MtN3\_slv domain were “genuine” *SWEETs*, all *SWEETs* contained MtN3\_slv domains (domain architecture of *PmSWEETs* is shown in Supplement File S2). The number of amino acids, molecular weight (MW), and isoelectric point (pI) were calculated on the basis of the protein sequences. As exhibited in Table 1, the predicted *PmSWEET* proteins ranged from



105 (*PmSWEET14*) to 580 (*PmSWEET8*) amino acids in length, with relative molecular weights ranging from 15.96 kDa (*PmSWEET11*) to 63.43 kDa (*PmSWEET8*), and theoretical pIs ranging from 8.30 (*PmSWEET4*) to 9.76 (*PmSWEET3*). The MW and pI of family member *PmSWEET14* could not be determined using this approach however due to the presence of four consecutive undefined amino acids (Table 1). Through prediction and analysis of TMHs of the 17 identified *PmSWEETs*, we found that these *PmSWEET* proteins were predicted to have 2–7 TMHs, and seven members of the *P. mume* SWEET gene family possess 7 TMHs, rarely, there may be only three or two TMHs. Detailed location information of the TMHs is shown in Table S3 and Figure S1.

### 3.2 Phylogenetic Analysis and Classification of *SWEET* Genes

To better understand the evolution of homologous *SWEET* genes, we used the ML method to create a phylogenetic tree of all *SWEET* sequences from *A. thaliana* (model dicots), *O. sativa* (model monocots), and *P. mume*. According to previously reported *AtSWEETs* and *OsSWEETs* (Chen et al., 2010; Yuan and Wang, 2013), the 17 identified *PmSWEETs* were divided into four clades (i.e., Clade I, Clade II, Clade III, and Clade IV) (Figure S2). To investigate the evolutionary relationships between *PmSWEETs* and the *SWEETs* of other species, an ML phylogenetic tree of *SWEETs* from 11 species, including 8 other Rosaceae species, was constructed. All members of the *SWEET* gene family in the 11 species were divided into four clades (Figure 1). The largest clade was Clade III, which comprised five *OsSWEET* genes, seven *AtSWEET* genes, and 68 Rosaceae *SWEET* genes; the specific number of genes is shown in Table S4. The smallest clade was Clade IV, which consisted of only two *A. thaliana* *SWEET* genes, one *O. sativa* gene, and 18 Rosaceae *SWEET* genes (Table S4), indicating that *SWEETs* were distributed unevenly among the clades. The numbers of genes in Clade I, II and III varied greatly, suggesting that the *SWEET* gene family expanded, especially in Clades I, II and III, during Rosaceae evolution. The *SWEETs* of Rosaceae were distributed uniformly across each small clade, whereas *SWEETs* from *O. sativa* tended to cluster together. The *PmSWEETs*, *PpSWEETs*, and *PavSWEETs* were clustered together and had similar distributions in the phylogenetic tree.

### 3.3 Conserved Motif and Gene Structure Analysis

To explore the sequence features of *PmSWEET* proteins, MEME software and TBtools were used to predict and draw conserved domains. As a consequence, ten distinct motifs were detected in *SWEET* proteins (Figure 2B), and a schematic diagram of *PmSWEET* protein motifs is shown in Figure S3. The number of *PmSWEETs* motifs was distinctive, ranging from 1 to 7. Of them, 12 *PmSWEETs* contained more than four motifs, 4 *PmSWEETs* harbored four motifs, and *PmSWEET14* contained only one motifs. Motifs 1, 2, 3, 4 and 6 were highly conserved and present in 15 *PmSWEET*, 13 *PmSWEET*, 16 *PmSWEET*, 11 *PmSWEET* and 12 *PmSWEET*

proteins, respectively; while motifs 7, 8 and 10 were relatively unique and existed in only 4 *PmSWEET*, 2 *PmSWEET* and 2 *PmSWEET* proteins, respectively. Intriguingly, aside from some unusual proteins, *SWEET* members of the same clade had similar conserved motifs, suggesting that they might have similar functions.

To elucidate the structural characteristics of the *PmSWEETs*, the exon-intron structure was further analyzed. As shown in Figure 2C, *PmSWEETs* in Clade II (except *PmSWEET10*) contained four introns. *PmSWEET1*, *PmSWEET9*, and *PmSWEET15* in Clade III had five introns, *PmSWEET8* contained the largest number of introns (12 introns), while *PmSWEET14* contained only one intron. All *PmSWEETs* in Clade IV had five introns. The number of introns in Clade I varied from just two to ten, *PmSWEET17* had two introns, *PmSWEET4* contained five introns, *PmSWEET11* and *PmSWEET12* contained three introns, *PmSWEET3* had ten introns. These results indicated that aside from some unusual proteins, genes clustered together generally exhibited similar gene structures.

### 3.4 Chromosomal Distribution and Tandem Duplication of *PmSWEET* gene family members

According to gene location information, all 17 *PmSWEETs* were mapped, showing that 16 *PmSWEETs* were located on chromosomes, and one *PmSWEET* gene was located on scaffold54 (Figure 3). The *PmSWEETs* on chromosomes 6 and 7 were clustered in the center of each chromosomes, and all contained four *PmSWEETs*. Two genes each were distributed on chromosomes 2, 3, 4 and 5. *PmSWEET11* and *PmSWEET12* and then *PmSWEET14* and *PmSWEET15* were clustered into two tandem duplication events on chromosomes 6 and 7, respectively. Based on the above results, some *PmSWEETs* gene family members were putatively generated by gene tandem duplication.

### 3.5 Segmental Duplication and Synteny of the *PmSWEET* Gene Family

Synteny analysis of *PmSWEETs* was performed using the Circos program of TBtools, four segmental duplication events, including *PmSWEET1/PmSWEET14*, *PmSWEET5/PmSWEET8*, *PmSWEET6/PmSWEET9* and *PmSWEET6/PmSWEET16* were detected, and further, each gene pair was located on a different chromosome, as shown with red lines in Figure 4. This finding strongly suggests that some *PmSWEETs* were probability generated by gene segmental duplication. In addition, the selection pressure and divergence time of the duplication events were estimated by the Ka (nonsynonymous) and Ks (synonymous) substitution ratio. In the evolutionary process, the Ka/Ks ratio > 1 indicates positive selection (adaptive evolution), a ratio = 1 indicates neutral evolution (drift), and a ratio < 1 indicates negative selection (conservation). Only one pair of segmentally duplicated *PmSWEETs* (*PmSWEET6/9*) had a Ka/Ks ratio of 0.45, which was significant, and indicated a synonymous change that has been selected during plant

genome evolution. The differentiation period of the *PmSWEET6/9* gene pair was 55.34~136.07 Mya.

To further examine the specific retention of *PmSWEETs*, their collinearity relationship with *AtSWEETs*, *PaSWEETs*, and *PsSWEETs* were detected using the MCScanX procedure of TBtools. A total of 16 homologous gene pairs were detected in *P. mume* and *A. thaliana*. Similarly, 16 pairs of homologous genes between *P. mume* and *P. armeniaca* and 20 between *P. mume* and *P. salicina* were detected (Figure 5, Table S5). The collinear complexity of *P. mume* with *P. salicina* was much higher than that with *P. armeniaca* and *A. thaliana*. These results suggested that *P. mume* was relatively distantly related to *A. thaliana* and *P. armeniaca*, but is more closely related to *P. salicina*.

### 3.6 Prediction Analysis of *Cis*-Acting Elements within *PmSWEETs* gene promoters

To further investigate the possible regulatory mechanism of *PmSWEETs* in the process of growth or in plant defence mechanisms, in particular the response to abiotic stress, such as low temperature, we submitted the 2.0 kb upstream sequence from the translation start site of each *PmSWEET* gene to the PlantCARE database to search for the presence of specific *cis*-elements. The *PmSWEET* promoters comprised several conserved regulatory elements that respond to plant hormones and environmental stress, and twelve of these were analyzed further (Figure 6, Table S6). Elements related to light response, anaerobic induction, and ABA response were widespread in the promoter areas of 17, 17 and 16 members of the *P. mume* SWEET gene family, respectively. According to the regulatory elements in their promoters, 14, 12, 11, 10, and 9 *P. mume* SWEET gene family members were sensitive to drought inducibility, MeJA, gibberellin, low temperatures and auxin, respectively. By combining these findings with the results of phylogenetic analysis, it was found that gene members of the same clade had similar *cis*-elements. These results indicated that *PmSWEET* genes were involved in the regulatory mechanisms of various stress responses.

### 3.7 Expression Pattern Analysis of *PmSWEETs*

To investigate the role of *PmSWEETs* in development and response to low temperature, the expression patterns of family members in the roots, stems, leaves, buds, fruits and flower buds of different stages of dormancy, were examined based on the RNA-seq dataset (Jiang, 2020), and their RPKM values are shown in Tables S7 and S8. As illustrated in Figure 7A, 14 of the *PmSWEET* genes were expressed in at least one tissue, whereas RNA-seq failed to detect the expression of three family members (*PmSWEET5*, *PmSWEET 10* and *PmSWEET 11*). Among them, five *PmSWEETs* presented relatively higher expression levels in fruits (*PmSWEET1*, *PmSWEET6*, *PmSWEET9*, *PmSWEET12* and *PmSWEET17*) and buds (*PmSWEET3*, *PmSWEET13*, *PmSWEET14*, *PmSWEET15* and *PmSWEET16*). Two *PmSWEETs* showed higher

expression levels in roots (*PmSWEET4* and *PmSWEET7*) and stems (*PmSWEET2* and *PmSWEET8*). Additionally, several genes (*PmSWEET2*, *PmSWEET3*, *PmSWEET4*, *PmSWEET7*, *PmSWEET8*, *PmSWEET12* and *PmSWEET13*) were expressed in leaves, but their expression levels were low.

Most *PmSWEETs* were expressed during the bud dormancy period (except *PmSWEET5* and *PmSWEET16*) as well as being expressed at specific stages of development (Figure 7B). Ten *PmSWEET* genes exhibited specifically higher expressions in the Natural flush (NF) stage (February), *PmSWEET9* was preferentially expressed in the Endo-dormancy I (EDI) stage (November), *PmSWEET10* and *PmSWEET12* showed the highest level of expression in the Endo-dormancy II (EDII) stage (December); and *PmSWEET1*, *PmSWEET3*, *PmSWEET6*, *PmSWEET12* and *PmSWEET13* showed upregulated expression in the Endo-dormancy III (EDIII) stage (January). Among these upregulated genes, eight *PmSWEETs* (*PmSWEET6*, *PmSWEET7*, *PmSWEET10*, *PmSWEET11*, *PmSWEET13*, *PmSWEET14*, *PmSWEET15* and *PmSWEET17*) (Table S6) contained low temperature response elements within their analyzed promoter regions.

To further investigate the expression patterns of *PmSWEETs* under cold exposure, we analyzed the stems of the cold-tolerant cultivar *P. mume* ‘Songchun’ at three geographical locations, and their FPKM values are displayed in Table S9. The expression of six *PmSWEET* genes (*PmSWEET5*, *PmSWEET6*, *PmSWEET11*, *PmSWEET14*, *PmSWEET16* and *PmSWEET17*) was not detected. Among the other 11 *PmSWEET* genes, seven *PmSWEETs* (*PmSWEET1*, *PmSWEET2*, *PmSWEET3*, *PmSWEET4*, *PmSWEET7*, *PmSWEET8* and *PmSWEET9*) showed higher expression in spring (3.2~5.3 °C). *PmSWEET13* expression was upregulated in autumn (6.1~7.9 °C) and winter in Beijing (-5.4 °C) and Chifeng (-11.4 °C), but downregulated in spring; the expression levels of *PmSWEET10*, *PmSWEET12* and *PmSWEET15* increased significantly in winter in Beijing (-5.4 °C) (Figure 8A). Among these genes with upregulated expression, four *PmSWEETs* (*PmSWEET7*, *PmSWEET10*, *PmSWEET13* and *PmSWEET15*) (Table S6) contained low-temperature response elements within their analyzed promoter regions. To compare the expression patterns of *PmSWEETs* during different times of the year, another heatmap was generated (Figure 8B). As shown in Figure 8B, *PmSWEETs* expression in the material sourced from the locations, Chifeng and Gongzhuling showed similar expression patterns at the same time of the year, while *PmSWEETs* expressed for the material sourced from the Beijing location showed higher expression in winter (Figure 8B). This may be related to the latitude of the three places, Gongzhuling has the highest latitude, followed by Chifeng and Beijing. There is little difference between the temperature in autumn and spring in these three places, but there is a big difference in winter. In winter, the temperature in Beijing (-5.4 °C) is higher than that in the other two places (Gongzhuling is -22.8 °C, Chifeng is -11.4 °C), which may be the temperature that induces some *PmSWEET* gene expression.

### 3.8 Expression Patterns of *P. mume* SWEETs under Cold Treatment

To investigate the role of *PmSWEETs* in response to cold stress, the expression patterns under imposed hypothermia (4 °C) (0, 1, 4, 6, 12, 24, 48 and 72 h) were examined by qRT-PCR using the cold-sensitive cultivar ‘Zaolve’ and the cold-tolerant cultivar ‘Songchun’. We performed a qRT-PCR assay on the 17 identified *P. mume* SWEETs, but the expression of only 11 *PmSWEETs* was detectable by this approach, while the remaining 6 *PmSWEETs* (*PmSWEET5*, *PmSWEET6*, *PmSWEET9*, *PmSWEET11*, *PmSWEET15* and *PmSWEET16*) were not detected, consistent with the transcriptome data (Figures 7, 8). As displayed in Figure 9, the changes in expression levels of the 11 *SWEET* genes in the two cultivars differed during the imposed cold stress treatment period. In two varieties, three genes (*PmSWEET2*, *PmSWEET7* and *PmSWEET8*) could be induced to downregulated in both ‘Songchun’ and ‘Zaolve’. In addition, the expression of *PmSWEET13* could be induced to upregulated in both ‘Songchun’ and ‘Zaolve’, which rose approximately 11-fold after 6 h of cold treatment in ‘Songchun’, while rose approximately 9-fold after 1 h, and then increased nearly 80-fold after 72 h of cold treatment in ‘Zaolve’. One gene (*PmSWEET3*) changed only slightly in both ‘Songchun’ and ‘Zaolve’. Six genes (*PmSWEET1*, *PmSWEET4*, *PmSWEET10*, *PmSWEET12*, *PmSWEET14*, and *PmSWEET17*) exhibited different expression patterns in the two cultivars. Among those, *PmSWEET1* and *PmSWEET12* were upregulated initially, then downregulated with increasing treatment duration in ‘Songchun’, while in ‘Zaolve’, there was no obvious change in early stage, but rapidly upgraded at 48 h and 72 h, respectively. *PmSWEET4* and *PmSWEET10* were dramatically downregulated with increased cold stress duration in ‘Songchun’, while they were upregulated within 6 h and then decreased with extended treatment in ‘Zaolve’. *PmSWEET14* was no obvious change in early stage, but rapidly upregulated at 72 h in ‘Songchun’, while it was rapidly upregulated at 24 h in ‘Zaolve’, and then downregulated with increasing treatment duration. *PmSWEET17* was upregulated firstly, then downregulated with increasing treatment duration in ‘Songchun’, while it was highly expressed only at 4 h in ‘Zaolve’.

## 4. Discussion

SWEET genes form a family of sugar transporters that play a role in the transportation of sugars, mainly sucrose, glucose and fructose (Chen et al., 2010; Chen et al., 2012; Feng and Frommer, 2015; Guo et al., 2014; Klemens et al., 2013; Le Hir et al., 2015), and participate in diverse physiological and biological processes in the growth and development of many plants and their responses to biotic and abiotic factors (Lemoine et al., 2013; Li et al., 2017; Li et al., 2018; Zhao et al., 2018). Previous studies have shown that *SWEETs* participate in cold stress responses in several plants (Chardon et al., 2013; Klemens et al., 2013; Guo et al., 2014; Chong et al., 2014; Liu et al., 2016; Le Hir et al., 2015; Yue et al., 2015; Wang et al., 2018; Feng et al., 2015; Miao et al., 2017; Zhang et al., 2019). However, little is known about the potential roles of

*P. mume* SWEET genes involved in cold stress. *P. mume* has a high ornamental value, and it can blossom at lower temperatures; but different varieties have different cold resistance, making it a very good material for studying the mechanisms of how *P. mume* SWEET genes function in cold responses. Understanding the link between SWEET genes of *P. mume* and cold-resistance could provide insights into cold-resistance molecular breeding in the future. In this research, we detected a total of 17 *PmSWEETs* in *P. mume*, as many as in *Arabidopsis*, and similar to the numbers in other species of *Prunus*, showing that SWEET genes are still relatively conserved in *Prunus*. The length of *PmSWEET* proteins ranges from 105 aa to 580 aa, and this range provides diversity in the number of TMHs (2–7). *PmSWEETs*, except for *PmSWEET14*, have a theoretical pI larger than 8.0. As an important parameter of proteins, pI is determined by the relative contents of amino acid residues at different pH values, which affects the stability, activity and function of proteins (Gasteiger, 2005). The pI of *PmSWEET14* was not detected, which may be due to its short amino acid sequence.

By predicting TMH domains, we found that the number of TMHs in *PmSWEET* genes ranged from 2 to 7, certain *P. mume* SWEETs with only two, three, four, five or six TMHs (Table 1). Fewer than seven TMHs in the eukaryotic SWEET family were also found in other plants, such as wheat (Gao et al., 2018; Gautam et al., 2019), walnut (Jiang et al., 2020), *Kentucky bluegrass* (Zhang et al., 2020) and soybean (Patil et al., 2015). To further validate the accuracy of the SWEET protein, we submitted the protein sequence to the NCBI-CDD and SMART online tools to predict its conserved domains, and it was found that each assessed family member contained the MtN3\_slv domain, and therefore, belonged to the SWEET family. The results means that duplication and fusion or genetic loss might take place in the *P. mume* genome. Similar to the case in other plants (Chen et al., 2010; Yuan and Wang, 2013; Patil et al., 2015), *PmSWEETs* can be classified into four clades, and the number of 11 species SWEET genes members in Clade III was larger than that in other clades (Figure 1), suggesting that Clade III may have expanded during evolution. Conserved motif analysis indicates that some special motifs only reside in some certain *PmSWEET* gene members. For instance, motif 8 was uniquely present in *PmSWEET11* and *PmSWEET17*; and motif 10 was uniquely present in *PmSWEET3* and *PmSWEET15*. These results are consistent with those of other plants, such as *Arabidopsis* (Chen et al., 2010), rice (Yuan and Wang, 2013), banana (Miao et al., 2017) and wheat (Gautam et al., 2019). Studies have disclosed that gene structural diversity and conserved protein motif divergence performed key roles in the evolution of the SWEET gene family (Xu et al., 2012), some *PmSWEETs* harbored unique conserved motifs, implying it may be responsible for the functional diversity of SWEET in *P. mume*.

Gene duplication, including tandem and segmental duplication events, is the origin of gene family expansion and genomic evolution in plants (Cannon et al., 2004; Ganko et al., 2007). In this study, two pairs of *PmSWEETs* were detected as tandem duplications, and four pairs of

*PmSWEET*s were segmental duplications. This outcome was consistent with those of other studies on *SWEET* duplication, including segmental and tandem duplications (Feng et al., 2015; Miao et al., 2017; Gao et al., 2018; Jiang et al., 2020).

The *cis*-elements in the promoter play an essential role in gene regulation. All *PmSWEET*s contain at least one light-responsive and anaerobically induced *cis*-element, suggesting that the two elements have an essential role in *PmSWEET* regulation. Moreover, 10 *PmSWEET*s contained one or more low-temperature responsive *cis*-elements (Table S6), indicating that these *PmSWEET*s may play important roles in the response to cold stress. However, whether and how these *cis*-elements work in *P. mume* requires further research.

Studies have shown that under low-temperature stress, the soluble sugar content in plants increases, and sugar transporters maintain the balance of osmotic potential through the balance and distribution of sugar, thus improving the cold tolerance of plants (Yamada et al., 2010). Numerous studies have also verified that *SWEET*s are involved in maintaining sugar homeostasis in plant organs and promoting plant adaptation to low temperatures (Seo et al., 2011; Chardon et al., 2013; Klemens et al. 2013; Chandran, 2015; Le Hir et al. 2015; Miao et al., 2017; Wang et al., 2018; Zhang et al., 2019; Zhang et al., 2020). Transcriptome analysis showed that *PmSWEET*s were differentially expressed in different tissues and during dormancy release and cold acclimation. *PmSWEET5* expression was not detected in any tissue/organ that we used, indicating that its expression may be variety -specific or time-specific. Some *PmSWEET*s had specific expression patterns in different organs (Figure 7A). For example, expression of *PmSWEET10* was detected only in ‘Zaolve’ buds at dormancy (stage EDII) and ‘Songchun’ stems in winter in Beijing; *PmSWEET16* expression was detected only in *P. mume* buds, which indicates that the genes are expressed only in specific tissues or varieties, such organ-specific expression patterns was also observed in wheat (Gao et al., 2018; Gautam et al., 2019), walnut (Jiang et al., 2020), tea (Wang et al., 2018) and cabbage (Zhang et al., 2019). *AtSWEET5*, the homologue of *PmSWEET10* and *PmSWEET16*, plays a key role in seed germination, and expressed at different stages of pollen development (Engel et al., 2005). The results from expression studies of different organs indicate a role for *PmSWEET10* and *PmSWEET16* in pollen development, suggesting they might have a similar role as *AtSWEET5*. *PmSWEET1*, *PmSWEET6*, *PmSWEET9*, *PmSWEET12* and *PmSWEET17* were strongly expressed in fruit, indicating that these genes may regulate sugar allocation during fruit ripening. Such specific high expression of *SWEET*s in fruits has also been found in pineapple (Guo et al., 2018), sweet orange (Zheng et al., 2014) and apple (Zhen et al, 2018), it can be inferred that *SWEET* protein plays an important role in fruit development and ripening.. *PmSWEET4* (Clade I) and *PmSWEET7* (Clade IV) were strongly expressed in roots, this results had similar expression patterns to previous studies, that *SWEET*s in Clade IV were highly expressed in the root cortex and encoded proteins such as specific fructose uniporters in the root vacuole membrane (Guo et al., 2014).

The present results also show that most of the *PmSWEET* genes are expressed more strongly at different endo-dormancy stages in flower bud and fruit tissues than in other tissues and that these genes are differentially expressed during flower development (Figure 7A, 7B). Together, these results suggest that the *P. mume* SWEET family is intimately associated with reproductive development and that different genes are specifically involved during different developmental stages. In rice, *Arabidopsis* and soybean, the expression of SWEET genes is also higher in reproductive tissues than in other tissues (Yuan et al., 2014; Patil et al., 2015). *PmSWEETs* also have different expression levels during dormancy release in flower buds (from November to February). Thus, we speculate that these *PmSWEETs* may participate in the cold reaction at low temperatures to protect the flower bud. In addition, some *PmSWEETs* were expressed more at colder temperatures in the spring (3.2~5.3 °C) and at approximately -5 °C in the winter (Figure 8A), indicating that these two temperatures may trigger their cold stress response and increase *PmSWEET* expression to reduce stress injury.

The qRT-PCR analysis suggested that six of 17 *PmSWEET* genes (*PmSWEET5*, *PmSWEET6*, *PmSWEET9*, *PmSWEET11*, *PmSWEET15*, and 16) were not expressed in the stem, which was consistent with the transcriptome data. *PmSWEETs* were activated by low temperature (4 °C) and increased or decreased in expression with the extension of treatment time (Figure 9). The expression levels of five *PmSWEETs* (*PmSWEET2*, *PmSWEET4*, *PmSWEET7*, *PmSWEET8*, and *PmSWEET10*) in ‘Songchun’ and three *PmSWEETs* (*PmSWEET2*, *PmSWEET7* and *PmSWEET8*) in ‘Zaolve’ decreased with increasing treatment times (Figure 9), which suggested that these genes might be negatively regulated by low temperatures and result in increased cold sensitivity. The expression levels of two *PmSWEETs* (*PmSWEET13* and *PmSWEET14*) in ‘Songchun’ and three *PmSWEETs* (*PmSWEET1*, *PmSWEET12*, and *PmSWEET13*) in ‘Zaolve’ increased with prolonged treatment (Figure 9), which suggested that these genes might be positively regulated by cold stress responses and increase cold sensitivity. The discrepancy in expression patterns between *PmSWEET1*, *PmSWEET4*, *PmSWEET10*, *PmSWEET12*, *PmSWEET14* and *PmSWEET17* is potentially due to genetic differences between ‘Songchun’ and ‘Zaolve’.

## 5. Conclusions

In summary, our study is the first to perform genome-wide identification and characterization of SWEETs in *P. mume*, including chromosomal location, duplicated genes, gene structure, phylogenetic relationships and conserved motifs. In addition, the expression profiles of the *PmSWEET* genes in different tissues and geographic locations were also examined based on the RNA-seq data. Furthermore, the expression profiles of these *PmSWEET* genes under cold stress conditions were analyzed by qRT-PCR assay. Our results could provide important information for further research on the biological functions of *PmSWEETs*.



515

## 516 **Funding**

517 This work was supported by Forestry and Grassland Science and Technology Innovation Youth  
518 Top Talent Project of China (No. 2020132608), the National Key Research and Development  
519 Program of China (2018YFD1000401), and the National Natural Science Foundation of China  
520 (No. 31870689).

## 521 **Conflict of Interest**

522 The authors declare that the research was conducted in the absence of any commercial or  
523 financial relationships that could be construed as a potential conflict of interest.

## 524 **Author Contributions**

525 LS: conceptualization. PL and ML: data curation. ZW: formal analysis and software. LS, QZ and  
526 TC: funding acquisition and writing reviews and editing. ZW and JM: methodology. ZW:  
527 validation, visualization, and drafting the manuscript. All authors contributed to writing and  
528 approved the final manuscript.

## 529 **Data Availability Statement**

530 The original contributions presented in the study are included in the article/Supplementary  
531 Material, further inquiries can be directed to the corresponding author/s.

## 532 **Supplementary Material**

533 Supplemental information for this article can be found online at

534 Supplementary Figure 1 | Schematic representation of *PmSWEET* proteins.

535 Supplementary Figure 2 | Phylogenetic trees of *Arabidopsis thaliana*, *Prunus mume* and Rice

536 Supplementary Figure 3 | Schematic diagram of *PmSWEET* protein motifs

537 Supplementary Table 1 | Primer sequences used for qRT-PCR

538 Supplementary Table 2 | Information for the proteins used in the present study

539 Supplementary Table 3 | TM helix Locus of *PmSWEETs*

540 Supplementary Table 4 | The specific number of genes in the Clades used in the present study

541 Supplementary Table 5 | Duplication events between *P. mume* and *A. thaliana*, *P. armeniaca* and  
542 *P. salicina*

543 Supplementary Table 6 | The data of cis-acting element in *PmSWEETs* promoters

544 Supplementary Table 7 | Expression profiles of 17 *PmSWEET* genes in five different tissues  
545 (root, stem, leaf, bud and fruit) (RPKM)

546 Supplementary Table 8 | Expression profiles of *PmSWEET* genes during the process of flower

547 bud dormancy release (RPKM)  
 548 Supplementary Table 9 | Expression profiles of 17 *PmSWEET* genes in different regions and  
 549 seasons (FPKM)  
 550 Supplementary Flie 1 | Protein sequences of *P. mume*  
 551 Supplementary Flie 2 | Domain architecture of *PmSWEETs*  
 552

# References

- Ainsworth, E.A., and Bush, D.R. (2011). Carbohydrate export from the leaf: A highly regulated process and target to enhance photosynthesis and productivity. *Plant Physiol.* 155(1), 64-69. doi: 10.1104/pp.110.167684s
- Baek, S., Choi, K., Kim, G.B., Yu, H.J., Cho, A., Jang, H., et al. (2018). Draft genome sequence of wild *Prunus yedoensis* reveals massive inter-specific hybridization between sympatric flowering cherries. *Genome Biol.* 19(1), 127. doi: 10.1186/s13059-018-1497-y
- Bailey, T.L., Boden, M., Buske, F.A., Frith, M., Grant, C.E., Clementi, L., et al. (2009). MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res.* 37(Web Server issue), W202-208. doi: 10.1093/nar/gkp335
- Blanc, G., and Wolfe, K.H. (2004). Widespread paleopolyploidy in model plant species inferred from age distributions of duplicate genes. *Plant Cell* 16(7), 1667-1678. doi: 10.1105/tpc.021345.
- Braun, D.M. (2012). SWEET! The pathway is complete. *Science* 335(6065), 173-174. doi: 10.1126/science.1216828
- Cannon, S.B., Mitra, A., Baumgarten, A., Young, N.D., and May, G. (2004). The roles of segmental and tandem gene duplication in the evolution of large gene families in *Arabidopsis thaliana*. *BMC Plant Biol.* 4, 10. doi: 10.1186/1471-2229-4-10
- Chardon, F., Bedu, M., Calenge, F., Klemens, P.A.W., Spinner, L., Clement, G., et al. (2013). Leaf fructose content is controlled by the vacuolar transporter *SWEET17* in *Arabidopsis*. *Curr. Biol.* 23(8), 697-702. doi: 10.1016/j.cub.2013.03.021
- Chandran, D. (2015). Co-option of developmentally regulated plant SWEET transporters for pathogen nutrition and abiotic stress tolerance. *IUBMB Life.* 67(7): 461-71. doi: 10.1002/iub.1394
- Chen, C., Chen, H., Zhang, Y., Thomas, H.R., Frank, M.H., He, Y., et al. (2020). TBtools: An integrative toolkit developed for interactive analyses of big biological data. *Mol. Plant.* 13(8), 1194-1202. doi: 10.1016/j.molp.2020.06.009
- Chen, L.Q. (2014). SWEET sugar transporters for phloem transport and pathogen nutrition. *New Phytol.* 201(4), 1150-1155. doi: 10.1111/nph.12445
- Chen, L.Q., Hou, B.H., Lalonde, S., Takanaga, H., Hartung, M.L., Qu, X.Q., et al. (2010). Sugar transporters for intercellular exchange and nutrition of pathogens. *Nature* 468(7323), 527-532. doi: 10.1038/nature09606
- Chen, L.Q., Cheung, L.S., Feng, L., Tanner, W., and Frommer, W.B. (2015). Transport of sugars. *Annu. Rev. Biochem.* 84, 865-894. doi: 10.1146/annurev-biochem-060614-033904
- Chen, L.Q., Qu, X.Q., Hou, B.H., Sossio, D., Osorio, S., Fernie, A.R., et al. (2012). Sucrose efflux mediated by SWEET proteins as a key step for phloem transport. *Science* 335(6065), 207-211. doi: 10.1126/science.1213351

- Chong, J., Piron, M.C., Meyer, S., Merdinoglu, D., Bertsch, C., and Mestre, P. (2014). The SWEET family of sugar transporters in grapevine: *VvSWEET4* is involved in the interaction with *Botrytis cinerea*. *J. Exp. Bot.* 65(22), 6589-6601. doi: 10.1093/jxb/eru375
- Daccord, N., Celton, J.M., Linsmith, G., Becker, C., Choisne, N., Schijlen, E., et al. (2017). High-quality de novo assembly of the apple genome and methylome dynamics of early fruit development. *Nat. Genet.* 49(7), 1099-1106. doi: 10.1038/ng.3886
- Durand, M., Porcheron, B., Hennion, N., Maurousset, L., Lemoine, R., and Pourtau, N. (2016). Water deficit enhances C export to the roots in *Arabidopsis thaliana* plants with contribution of sucrose transporters in both shoot and roots. *Plant Physiol.* 170(3), 1460-1479. doi: 10.1104/pp.15.01926
- Engel, M.L., Holmes-Davis, R., McCormick, S. (2005). Green Sperm. Identification of male gamete promoters in *Arabidopsis*. *Plant Physiol.* 138, 2124–2133. doi: 10.1104/pp.104.054213
- Eom, J.S., Chen, L.Q., Sosso, D., Julius, B.T., Lin, I.W., Qu, X.Q., et al. (2015). SWEETs, transporters for intracellular and intercellular sugar translocation. *Curr. Opin. Plant Biol.* 25, 53-62. doi: 10.1016/j.pbi.2015.04.005
- Feng, L., and Frommer, W.B. (2015). Structure and function of SemiSWEET and SWEET sugar transporters. *Trends Biochem. Sci.* 40(8), 480-486. doi: 10.1016/j.tibs.2015.05.005.
- Feng, C.Y., Han, J.X., Han, X.X., and Jiang, J. (2015). Genome-wide identification, phylogeny, and expression analysis of the SWEET gene family in tomato. *Gene* 573(2), 261-272. doi: 10.1016/j.gene.2015.07.055
- Finn, R.D., Clements, J., Arndt, W., Miller, B.L., Wheeler, T.J., Schreiber, F., et al. (2015). HMMER web server: 2015 update. *Nucleic Acids Res.* 43(W1), W30-W38. doi: 10.1093/nar/gkv397
- Ganko, E.W., Meyers, B.C., and Vision, T.J. (2007). Divergence in expression between duplicated genes in *Arabidopsis*. *Mol. Biol. Evol.* 24(10), 2298-2309. doi: 10.1093/molbev/msm158
- Gao, Y., Wang, Z.Y., Kumar, V., Xu, X.F., Yuan, P., Zhu, X.F., et al. (2018). Genome-wide identification of the SWEET gene family in wheat. *Gene* 642, 284-292. doi: 10.1016/j.gene.2017.11.044
- Gasteiger, E. (2005). Protein identification and analysis tools on the ExPASy server. *The proteomics protocols handbook*. Humana Press, pp. 571-607
- Ge, Y.X., Angenent, G.C., Wittich, P.E., Peters, J., Franken, J., Busscher, M., et al. (2000). NEC1, a novel gene, highly expressed in nectary tissue of *Petunia hybrida*. *Plant J.* 24(6), 725-734. doi: 10.1046/j.1365-3113x.2000.00926.x
- Guo, C., Li, H., Xia, X., Liu, X., and Yang, L. (2018) Functional and evolution characterization of SWEET sugar transporters in *Ananas comosus*. *Biochem. Biophys. Res. Commun.* 496(2): 407-414. doi: 10.1016/j.bbrc.2018.01.024

- Guo, W.J., Nagy, R., Chen, H.Y., Pfrunder, S., Yu, Y.C., Santelia, D., et al. (2014). *SWEET17*, a facilitative transporter, mediates fructose transport across the tonoplast of *Arabidopsis* roots and leaves. *Plant Physiol.* 164(2), 777-789. doi: 10.1104/pp.113.232751
- Jia, B., Zhu, X.F., Pu, Z.J., Duan, Y.X., Hao, L.J., Zhang, J., et al. (2017). Integrative view of the diversity and evolution of SWEET and SemiSWEET sugar transporters. *Front. Plant Sci.* 8, 2178. doi: 10.3389/fpls.2017.02178
- Jian, H., Lu, K., Yang, B., Wang, T., Zhang, L., Zhang, A., et al. (2016). Genome-wide analysis and expression profiling of the SUC and SWEET gene families of sucrose transporters in *Oilseed Rape (Brassica napus L.)*. *Front. Plant Sci.* 7, 1464. doi: 10.3389/fpls.2016.01464
- Jiang, F., Zhang, J., Wang, S., Yang, L., Yingfeng, L., Gao, S., et al. (2019). The apricot (*Prunus armeniaca L.*) genome elucidates Rosaceae evolution and beta-carotenoid synthesis. *Hortic. Res.* 6, 128. doi: 10.1038/s41438-019-0215-6
- Jiang, L. (2020). Physiological Changes and Gene Expression Pattern in Response to Low Temperature Stress in *Prunus mume*. Beijing: Beijing Forestry University. 44-50.
- Jiang, S., Balan, B., Assis, R.d.A.B., Sagawa, C.H.D., Wan, X., Han, S., et al. (2020). Genome-wide profiling and phylogenetic analysis of the SWEET sugar transporter gene family in walnut and their lack of responsiveness to *Xanthomonas arboricola* pv. *juglandis* infection. *Int. J. Mol. Sci.* 21(4), 1251. doi: 10.3390/ijms21041251
- Katoh, K., and Standley, D.M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30(4), 772-780. doi: 10.1093/molbev/mst010
- Klemens, P.A., Patzke, K., Deitmer, J., Spinner, L., Le Hir, R., Bellini, C., et al. (2013). Overexpression of the vacuolar sugar carrier *AtSWEET16* modifies germination, growth, and stress tolerance in *Arabidopsis*. *Plant Physiol.* 163(3), 1338-1352. doi: 10.1104/pp.113.224972
- Le Hir, R., Spinner, L., Klemens, P.A.W., Chakraborti, D., de Marco, F., Vilaine, F., et al. (2015). Disruption of the sugar transporters *AtSWEET11* and *AtSWEET12* affects vascular development and freezing tolerance in *Arabidopsis*. *Mol. Plant* 8(11), 1687-1690. doi: 10.1016/j.molp.2015.08.007
- Lemoine, R. (2000). Sucrose transporters in plants: update on function and structure. *BBA. Bioenergetics* 1465(1-2), 246-262. doi: 10.1016/s0005-2736(00)00142-5
- Lemoine, R., La Camera, S., Atanassova, R., Dedaldechamp, F., Allario, T., Pourtau, N., et al. (2013). Source-to-sink transport of sugar and regulation by environmental factors. *Front. Plant Sci.* 4, 272. doi: 10.3389/fpls.2013.00272
- Lescot, M., Dehais, P., Thijs, G., Marchal, K., Moreau, Y., Van de Peer, Y., et al. (2002). PlantCARE, a database of plant cis-acting regulatory elements and a portal to tools for in silico analysis of promoter sequences. *Nucleic Acids Res.* 30(1), 325-327. doi: 10.1093/nar/30.1.325

- Letunic, I., and Bork, P. (2019). Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic Acids Res.* 47(W1), W256-W259. doi: 10.1093/nar/gkz239
- Li, H., Li, X., Xuan, Y., Jiang, J., Wei, Y., and Piao, Z. (2018). Genome wide identification and expression profiling of SWEET genes family reveals its role during plasmodiophora brassicae-induced formation of clubroot in *Brassica rapa*. *Front. Plant Sci.* 9, 207. doi: 10.3389/fpls.2018.00207
- Li, J., Qin, M., Qiao, X., Cheng, Y., Li, X., Zhang, H., et al. (2017). A new insight into the evolution and functional divergence of SWEET transporters in Chinese white pear (*Pyrus bretschneideri*). *Plant Cell Physiol.* 58(4), 839-850. doi: 10.1093/pcp/pcx025
- Li, L., Sheen, J. (2016). Dynamic and diverse sugar signaling. *Curr. Opin. Plant Biol.* 33, 116–125. doi: 10.1016/j.pbi.2016.06.018
- Li, M., Xie, H., He, M., Su, W., Yang, Y., Wang, J., et al. (2020). Genome-wide identification and expression analysis of the *StSWEET* family genes in potato (*Solanum tuberosum* L.). *Genes Genom.* 42(2), 135-153. doi: 10.1007/s13258-019-00890-y
- Lin, I.W., Sosso, D., Chen, L.Q., Gase, K., Kim, S.G., Kessler, D., et al. (2014). Nectar secretion requires sucrose phosphate synthases and the sugar transporter *SWEET9*. *Nature* 508(7497), 546-549. doi: 10.1038/nature13082
- Linsmith, G., Rombauts, S., Montanari, S., Deng, C.H., Celton, J.M., Guerif, P., et al. (2019). Pseudo-chromosome-length genome assembly of a double haploid "Bartlett" pear (*Pyrus communis* L.). *Gigascience* 8(12), giz138. doi: 10.1093/gigascience/giz138
- Liu, C., Feng, C., Peng, W., Hao, J., Wang, J., Pan, J., et al. (2020). Chromosome-level draft genome of a diploid plum (*Prunus salicina*). *Gigascience* 9(12), giaa130. doi: 10.1093/gigascience/giaa130
- Liu, X., Zhang, Y., Yang, C., Tian, Z., and Li, J. (2016). *AtSWEET4*, a hexose facilitator, mediates sugar transport to axial sinks and affects plant development. *Sci. Rep.* 6, 24563. doi: 10.1038/srep24563
- Livak, K.J., Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-delta delta C(T)) method. *Methods.* 25, 402–408. doi: 10.1006/meth.2001.1262
- Lynch, M., and Conery, J.S. (2000). The evolutionary fate and consequences of duplicate genes. *Science* 290(5494), 1151-1155. doi: 10.1126/science.290.5494.1151
- Maynard, J.W., and Lucas, W.J. (1982). Sucrose and glucose uptake into *Beta vulgaris* leaf tissues : A case for general (apoplastic) retrieval systems. *Plant physiol.* 70(5), 1436-1443. doi: 10.1104/pp.70.5.1436
- Miao, H., Sun, P., Liu, Q., Miao, Y., Liu, J., Zhang, K., et al. (2017). Genome-wide analyses of SWEET family proteins reveal involvement in fruit development and abiotic/biotic stress responses in banana. *Sci. Rep.* 7(1), 3536. doi: 10.1038/s41598-017-03872-w

- Mizuno, H., Kasuga, S., and Kawahigashi, H. (2016). The sorghum SWEET gene family: stem sucrose accumulation as revealed through transcriptome profiling. *Biotechnol. Biofuels* 9, 127. doi: 10.1186/s13068-016-0546-6
- Patil, G., Valliyodan, B., Deshmukh, R., Prince, S., Nicander, B., Zhao, M., et al. (2015). Soybean (*Glycine max*) SWEET gene family: insights through comparative genomics, transcriptome profiling and whole genome re-sequence analysis. *BMC Genomics* 16(1), 520. doi: 10.1186/s12864-015-1730-y
- Price, M.N., Dehal, P.S., and Arkin, A.P. (2010). FastTree 2-approximately maximum-likelihood trees for large alignments. *PloS One* 5(3), e9490. doi: 10.1371/journal.pone.0009490
- Quirino, B.F., Normanly, J., and Amasino, R.M. (1999). Diverse range of gene activity during *Arabidopsis thaliana* leaf senescence includes pathogen-independent induction of defense-related genes. *Plant Mol.Biol.* 40(2), 267-278. doi: 10.1023/a:1006199932265
- Raymond, O., Gouzy, J., Just, J., Badouin, H., Verdenaud, M., Lemainque, A., et al. (2018). The Rosa genome provides new insights into the domestication of modern roses. *Nat. Genet.* 50(6), 772-777. doi: 10.1038/s41588-018-0110-3
- Rennie, E.A., and Turgeon, R. (2009). A comprehensive picture of phloem loading strategies. *P. Natl. Acad. Sci. U.S.A.* 106(33), 14162-14167. doi: 10.1073/pnas.0902279106
- Seo, P.J., Park, J.M., Kang, S.K., Kim, S.G., and Park, C.M. (2011). An *Arabidopsis* senescence-associated protein SAG29 regulates cell viability under high salinity. *Planta* 233(1), 189-200. doi: 10.1007/s00425-010-1293-8
- Shirasawa, K., Isuzugawa, K., Ikenaga, M., Saito, Y., Yamamoto, T., Hirakawa, H., et al. (2017). The genome sequence of sweet cherry (*Prunus avium*) for use in genomics-assisted breeding. *DNA Res.* 24(5), 499-508. doi: 10.1093/dnares/dsx020
- Slewinski, T.L. (2011). Diverse functional roles of monosaccharide transporters and their homologs in vascular plants: aphysiological perspective. *Mol. Plant* 4(4), 641-662. doi: 10.1093/mp/ssr051
- Sosso, D., Luo, D., Li, Q.B., Sasse, J., Yang, J., Gendrot, G., et al. (2015). Seed filling in domesticated maize and rice depends on SWEET-mediated hexose transport. *Nat. Genet.* 47(12), 1489-1493. doi: 10.1038/ng.3422
- Sun, M.X., Huang, X.Y., Yang, J., Guan, Y.F., Yang, Z.N. (2013). *Arabidopsis* RPG1 is important for primexine deposition and functions redundantly with RPG2 for plant fertility at the late reproductive stage. *Plant Reprod.* 26, 83–91. doi: 10.1007/s00497-012-0208-1
- Verde, I., Abbott, A.G., Scalabrin, S., Jung, S., Shu, S., Marroni, F., et al. (2013). The high-quality draft genome of peach (*Prunus persica*) identifies unique patterns of genetic diversity, domestication and genome evolution. *Nat. Genet.* 45(5), 487-494. doi: 10.1038/ng.2586
- Wang, L., Yao, L., Hao, X., Li, N., Qian, W., Yue, C., et al. (2018). Tea plant SWEET transporters: expression profiling, sugar transport, and the involvement of CsSWEET16 in modifying cold tolerance in *Arabidopsis*. *Plant Mol. Biol.* 96(6), 577-592. doi: 10.1007/s11103-018-0716-y



- 741 Wei, X., Liu, F., Chen, C., Ma, F., and Li, M. (2014). The *Malus domestica* sugar transporter gene  
742 family: identifications based on genome and expression profiling related to the  
743 accumulation of fruit sugars. *Front. Plant Sci.* 5, 569. doi: 10.3389/fpls.2014.00569
- 744 Xu, G., Guo, C., Shan, H., and Kong, H. (2012). Divergence of duplicate genes in exon-intron  
745 structure. *Proc. Natl. Acad. Sci. U.S.A.* 109(4), 1187-1192. doi: 10.1073/pnas.1109047109
- 746 Xuan, Y.H., Hu, Y.B., Chen, L.Q., Sosso, D., Ducat, D.C., Hou, B.H., et al. (2013). Functional  
747 role of oligomerization for bacterial and plant SWEET sugar transporter family. *Proc. Natl.*  
748 *Acad. Sci. U.S.A.* 110(39), E3685-E3694. doi: 10.1073/pnas.1311244110
- 749 Yamada, K., Osakabe, Y., Mizoi, J., Nakashima, K., Fujita, Y., Shinozaki, K., et al. (2010).  
750 Functional analysis of an *Arabidopsis thaliana* abiotic stress-inducible facilitated diffusion  
751 transporter for monosaccharides. *J. Biol. Chem.* 285(2), 1138-1146. doi:  
752 10.1074/jbc.M109.054288
- 753 Yao, T., Xie, R., Zhou, Y., Hu, J., Gao, Y., and Zhou, C. (2021). Genome-wide identification of  
754 SWEET gene family and its response to abiotic stresses in valencia sweet orange. *Plant*  
755 *Mol. Biol. Rep.* doi: 10.1007/s11105-020-01268-1
- 756 Yuan, M., and Wang, S. (2013). Rice MtN3/Saliva/SWEET family genes and their homologs in  
757 cellular organisms. *Mol. Plant* 6(3), 665-674. doi: 10.1093/mp/sst035
- 758 Yuan, M., Zhao, J., Huang, R., Li, X., Xiao, J., and Wang, S. (2014). Rice MtN3/saliva/SWEET  
759 gene family: Evolution, expression profiling, and sugar transport. *J. Integr. Plant Biol.*  
760 56(6), 559-570. doi: 10.1111/jipb.12173
- 761 Yue, C., Cao, H.L., Wang, L., Zhou, Y.H., Huang, Y.T., Hao, X.Y., et al. (2015). Effects of cold  
762 acclimation on sugar metabolism and sugar-related gene expression in tea plant during the  
763 winter season. *Plant Mol. Biol.* 88(6), 591-608. doi: 10.1007/s11103-015-0345-7
- 764 Zhang, R., Niu, K., Ma, H. (2020). Identification and expression analysis of the SWEET gene  
765 family from *Poa pratensis* under abiotic stresses. *Dna Cell Biol.* 39(9), 1-15. doi:  
766 10.1089/dna.2020.5418
- 767 Zhang, Q., Chen, W., Sun, L., Zhao, F., Huang, B., Yang, W., et al. (2012). The genome of *Prunus*  
768 *mume*. *Nat. Commun.* 3, 1318. doi: 10.1038/ncomms2290
- 769 Zhang, W., Wang, S., Yu, F., Tang, J., Shan, X., Bao, K., et al. (2019). Genome-wide  
770 characterization and expression profiling of SWEET genes in cabbage (*Brassica oleracea*  
771 var. *capitata* L.) reveal their roles in chilling and clubroot disease responses. *BMC*  
772 *Genomics* 20(1), 93. doi: 10.1186/s12864-019-5454-2
- 773 Zhang, Z., Zhuo, X., Zhao, K., Zheng, T., Han, Y., Yuan, C., et al. (2018). Transcriptome profiles  
774 reveal the crucial roles of hormone and sugar in the bud dormancy of *Prunus mume*. *Sci.*  
775 *Rep.* 8(1), 5090. doi: 10.1038/s41598-018-23108-9
- 776 Zhao, L., Yao, J., Chen, W., Yan, L.I., Youjun, L., Guo, Y., et al. (2018). A genome-wide analysis  
777 of SWEET gene family in cotton and their expressions under different stresses. *J. Cotton*  
778 *Res.* 1(2), 1-15. doi: 10.1186/s42397-018-0007-9

779 Zhen Q.L., Fang T., Peng Q., Liao L., Zhao L., Owiti A., and Han Y.P. (2018). Developing gene-  
 780 tagged molecular markers for evaluation of genetic association of apple SWEET genes  
 781 with fruit sugar accumulation. *Hortic. Res.* 5(1): 14. doi: 10.1038/s41438-018-0024-3  
 782 Zheng Q.M., Tang Z., Xu Q., and Deng X.X. (2014). Isolation, phylogenetic relationship and  
 783 expression profiling of sugar transporter genes in sweet orange (*Citrus sinensis*). *Plant*  
 784 *Cell.* 119(3): 609-624. doi: 10.1007/s11240-014-0560-y

# **Table 1**(on next page)

Table 1. The *PmSWEET* gene family members in *P. mume*.

**Table 1. The *PmSWEET* gene family members in *P. mume*.**

Name	Gene ID	Clade	CDS (bp)	No. of amino acids	Molecular weight (kDa)	Theoretical pI	TMHs	No. of MtN3/saliv a domain	Locus
PmSWEET1	Pm007067	III	849	282	31.38	8.34	7	2	Pa2:21184396..21186332
PmSWEET2	Pm008206	IV	759	252	27.74	8.50	7	2	Pa2:31718730..31721555
PmSWEET3	Pm010330	I	1248	415	46.25	9.76	8	2	Pa3:3891190..3895205
PmSWEET4	Pm011260	I	708	235	26.45	8.30	7	2	Pa3:9921623..9924001
PmSWEET5	Pm013198	II	519	172	19.42	8.97	5	1	Pa4:2433448..2434735
PmSWEET6	Pm015728	II	708	235	25.67	9.21	5	2	Pa4:21122646..21124537
PmSWEET7	Pm017566	IV	735	244	26.99	9.14	7	2	Pa5:12327097..12328384
PmSWEET8	Pm018875	III	1743	580	63.43	8.34	6	2	Pa5:20984940..20990591
PmSWEET9	Pm019954	III	828	275	30.68	9.20	7	2	Pa6:436315..437664
PmSWEET10	Pm021931	II	708	235	26.60	8.59	6	2	Pa6:12459796..12461199
PmSWEET11	Pm022695	I	417	138	15.96	9.74	3	1	Pa6:19934418..19935334
PmSWEET12	Pm022696	I	651	216	23.21	8.78	5	2	Pa6:19944525..19945680
PmSWEET13	Pm024167	II	780	259	28.66	9.37	6	2	Pa7:10796671..10798904
PmSWEET14	Pm024554	III	318	105	-	-	2	1	Pa7:13005181..13005663
PmSWEET15	Pm024555	III	891	296	33.14	8.61	7	2	Pa7:13012731..13014646
PmSWEET16	Pm024712	II	639	212	23.95	8.37	5	2	Pa7:13852243..13854234
PmSWEET17	Pm030352	I	510	169	19.26	9.14	4	1	scaffold54:138478..139392

# Figure 1

Figure 1. Phylogenetic tree of SWEET sequences from *P. mume* and other plant species.

Clades I, II, III, and IV are indicated by blue, indigo, orange and pale yellow branch lines, respectively. At, *A. thaliana*; Os, *O. sativa*; Pa, *P. armeniaca*; Pav, *P. avium*; Pc, *P. communis*; Pm, *P. mume*; Pp, *P. persica*; Ps, *P. salicina*; Py, *P. yedoensis* var. *nudiflora*; Md, *M. domestica*; Rc, *R. chinensis*.



# Figure 2

Figure 2. Phylogenetic relationship, conserved motif and gene structure analysis of *PmSWEET* genes.


A: The ML phylogenetic tree of *PmSWEET* genes. The SWEET genes were grouped into four clades, and blue, purple, red, and green represents Clades I, II, III, and IV, respectively. B: The motif composition of *PmSWEET* proteins. Ten motifs were displayed in different colored rectangles. Motif1: GVVWFLYGLLKKDLFIAIPNGLGFJLGLVQLILYAIYR, Motif2: TKKRSLIVGIJCIVFNIIMYASPLTIMKLVIKTKSVEYMPFYLSLFLFLN, Motif3: LVITINGFGAVIELIYJAIFIIYAPKKKRKKI, Motif4: APVPTFYRIKKKSTEEFQSVPYVAALLNMotif5: WYGMPFVHPDN, Motif6: FGILGNIISFLLFL, Motif7: STNWDDDD, Motif8: PMTTLKRIMKKNEFTEQYLSGIPYLMT, Motif9: AMLWLYYGLLKPN, Motif10: NCZGCKDQYQHPQKCKE. Detailed information is shown with logos obtained from the MEME Suite website in Supplementary Figure 3. C: Exon-intron organization of *PmSWEET* genes. Green and black correspond to exons and introns, respectively.



# Figure 3

Figure 3. Schematic representations of the chromosomal location of the *PmSWEET* genes.

The chromosome number is indicated on the top of each chromosome and scaffold. Scf54 indicates scaffold54. Green and red gene names indicate tandem duplicated gene pairs.






# Figure 4

Figure 4. The Circos figure for *PmSWEET* segmental duplication links.


The red lines indicate segmented duplicated gene pairs.



# Figure 5

Figure 5. Synteny of SWEET genes in different genome of *P. mume*, *A. thaliana*, *P. armeniaca* and *P. salicina*.


A: Synteny of *PmSWEET* and *AtSWEET* gene pairs. B: Synteny of *PmSWEET* and *PaSWEET* gene pairs. C: Synteny of *PmSWEET* and *PsSWEET* gene pairs.



# Figure 6

Figure 6. Predicted cis-elements responding to plant growth regulation, hormone response, and stresses response present in the promoter of *PmSWEET* genes


Different colored boxes represent different elements and their positions in each *PmSWEET* promoter. The SWEET genes are classified into four clades, and blue, indigo, purple red, and green represent Clades I, II, III, and IV, respectively.



# Figure 7

Figure 7. Expression profiles of *PmSWEET* genes in different tissues and different flower buds stage


A: Expression profiles of *PmSWEETs* in different tissues. B: Expression profiles of *PmSWEETs* in the flower bud during dormancy. EDI: Endo-dormancy I, November; EDII: Endo-dormancy II, December; EDIII: Endo-dormancy III, January; NF: Natural flush, February. A 2-based log function conversion is performed on the expression amount, and then normalized by row using min-max method. The color scale on the right of the heat map refers to relative expression level, and the color gradient from blue to red shows an increasing expression level.



# Figure 8

Figure 8. Expression profiles of *PmSWEET*s in stems in different seasons and regions

A: Expression profiles of *PmSWEET*s in stems of ‘Songchun’ in different regions (Beijing, Chifengand Gongzhuling) and seasons (autumn, winter and spring). B: Comparison of differential expressionprofiles of stems in Beijing, Chifeng and Gongzhuling during different seasons.A 2-based log function conversion is performed on the expression amount, and then normalized by row using min-max method. The color scale on the right of the heat map refers to relative expression level, and the color gradient from blue to red shows an increasing expression level. Aut, Autumn; Win, Winter; Spr, Spring. BJ, Beijing; CF, Chifeng; GZL, Gongzhuling.



# Figure 9

Figure 9. Expression patterns of 11 *PmSWEET* genes under artificial low temperature treatments

The relative quantification method ( $2^{-\Delta\Delta C_t}$ ) was used to evaluate the transcript levels of 11 *PmSWEET* genes. Error bars are standard deviation of three replicates. The statistical analyses of 'Zaolve' and 'Songchun' were independent carried out using SPSS22.0, the one-way ANOVA analysis of variance was calculated by least significant difference (LSD) and Student-Newman-Keuls test, different letters above the bars indicate significant differences ( $p = 0.05$ ). Black letters indicate 'Zaolve', red letters indicate 'Songchun'. GraphPad Prism6 software was used to draw the diagram.

