

The fecal microbiota of Thai school-aged children associated with demographic factors and diet

Lucsame Gruneck¹, Eleni Gentekaki^{1,2}, Kongkiat Kespechara³, Justin Denny⁴, Thomas J. Sharpton^{5,6}, Lisa K. Marriott⁴, Jackilen Shannon⁴, Siam Popluechai^{1,2}

¹Gut Microbiome Research Group, Mae Fah Luang University, Muang, Chiang Rai, Thailand

²School of Science, Mae Fah Luang University, Muang, Chiang Rai, Thailand

³Sooksatharana (Social Enterprise) Co., Ltd., Muang, Phuket, Thailand.

⁴OHSU-PSU School of Public Health, Oregon Health & Science University, Portland, OR, United States of America

⁵Department of Microbiology, Oregon State University, Corvallis, OR, 97331, USA

⁶Department of Statistics, Oregon State University, Corvallis, OR, 97331, USA

Corresponding Author:

Siam Popluechai^{1,2}

¹Gut Microbiome Research Group, Mae Fah Luang University, Muang, Chiang Rai, Thailand

²School of Science, Mae Fah Luang University, Muang, Chiang Rai, Thailand

Email address: siam@mfu.ac.th

Abstract

Background. Birth delivery method and breastfeeding practices contribute to microbiota colonization. Other factors, including diet and demographic factors, structure gut microbiome assembly and diversity through childhood development. The exploration of these factors, especially those which influence in Southeast Asian children, remains limited.

Methods. We investigated fecal microbiota of 127 Thai school-aged children in Thailand using quantitative PCR (qPCR) and to assessed the influence of dietary behaviors and demographic factors on the gut microbiota, using a multivariate Multivariate analysis (multiple factor analysis (MFA) and Partial Least Squares Discriminant Analysis (PLS-DA) was used link particular gut microbes to diet and demographic factors.

Results. Dietary behaviors and demographic factors were associated with variation among gut microbiota. The abundance of *Gammaproteobacteria* increased in children with infrequent intake of high fat foods. Obese children possessed a lower level of Firmicutes and *Ruminococcus*. *Bifidobacterium* was enriched in pre-teen aged children and detected at lower levels among formula-fed children. *Prevotella* was more abundant in children who were delivered vaginally. While ethnicity explained a small amount of variation in the gut microbiota, it nonetheless was found to be significantly associated with microbiome composition.

Conclusions. Our results clarify the extent to which exogenous and demographic factors associate with, and possibly drive, the assembly of the gut microbiome of an understudied population of Thai school-aged children in Thailand.

Introduction

Microbial colonization of the gut starts upon birth, and the composition of the microbiota community diversifies throughout childhood. Proteobacteria and Actinobacteria dominate the gut microbiome early in life (Zhuang et al., 2019). As infants develop, their gut microbiota community becomes more complex and, at two to three years of age, its structure and composition begins to more closely resemble that of adults (Rodríguez et al., 2015). Shifts in the microbiota composition occur during this process and are influenced by several factors such as mode of delivery, feeding type, and diet (Martin et al., 2016; Rutayisire et al., 2016; Iddrisu et al., 2021).

Early in life, the assembly of the gut microbiome highly depends on delivery mode and feeding practices (Li et al., 2020; Mitchell et al., 2020; Coker et al., 2021). The guts microbiome of infants born vaginally are more enriched in *Bifidobacterium* and/or *Bacteroides* compared with those infants delivered by cesarean section (Yang et al., 2019; Reyman et al., 2019; Niu et al., 2020). Over time, the association between gut microbiota and mode of delivery weakens (Rutayisire et al., 2016). Nonetheless, differences in the abundance of microbiota between

children of different modes of delivery ~~was seen-persists~~ in school-aged children ~~(i.e. seven-year olds)~~ (Salminen et al., 2004). *Bifidobacterium* dominates the gut of infants receiving breast milk as compared to formula fed children (van den Elsen et al., 2019; Lawson et al., 2020). The impact of breastfeeding duration persists later ~~in life as shown in a cohort study of early school-aged children~~ (Zhong et al., 2019). ~~Overall, these findings imply This suggests~~ that a history of delivery mode and feeding type could have a long-term impact on ~~the~~ diversity of one's gut microbiome.

Additional host-related factors such as ethnicity, age, and body mass index (BMI) contribute to alterations of gut microbiota. Geographical factors and ethnicity significantly affect gut microbiota profiles of school-aged children at the genus level (Liu et al., 2020). Although many studies have already monitored compositional changes in the gut microbiota depending on one's age (e.g., comparing between childhood and adulthood) (Derrien, Alvarez & de Vos, 2019), data on microbiota profiles among school-aged children remains limited (Odamaki et al., 2016). Variation in the microbiome among children has also been linked to BMI (Bervoets et al., 2013; Da Silva, Monteil & Davis, 2020; Shin & Cho, 2020). These changes have been associated with diets which are precursors to weight gain and shape the gut microbiome (Voreades, Kozil & Weir, 2014; Cho, 2021).

Diet can shape the human gut microbiota (Singh et al., 2017; Zmora, Suez & Elinav, 2019). In children, diet explains most of the variation in gut microbiota profiles between countries or continents (De Filippo et al., 2010; Nakayama et al., 2015) as already observed in adults (Yasir et al., 2015; Escobar et al., 2015; Ghosh et al., 2020). Moving away from traditional diets with a high concentration of fiber, fruits and vegetables towards a Western diet rich in animal protein, fat and sugar is a cause of concern as high fat diets have been shown to disrupt the balance of gut microbiota in animal models (Kim et al., 2012). This effect has also been observed in humans where a decrease in the abundance of butyrate-producing bacteria has been noted in populations consuming higher-fat diets (Wan et al., 2019). ~~In recent year~~ Recently Southeast Asian populations have begun to adopt the Western diet (Ooraikul, Sirichote & Siripongvutikorn, 2008). However, only few studies have investigated the effects of this dietary pattern on the gut microbiota of young Southeast Asians (Nakayama et al., 2017; Gollos-Gubat et al., 2020).

This study is the first to examine the impact of diet and demographic factors (gender, age, BMI z-score, birth records, feeding type, and ethnicity) on the gut microbiota of ~~Thai~~ school-aged children in Thailand. Multivariate analyses were implemented to determine the potential contribution of multiple factors on variations of microbiota profiles as well as identifying most relevant features (microbiota taxa) for each host variable. Our results provide a preliminary overview of the associations observed between the abundance of gut microbiota and investigated factors in school-aged children from Thailand.

Materials & Methods

Ethics approval

All participants provided written informed consent (Supplementary File 1) and the study was approved by the Ethics committee of Mae Fah Luang University (Ethics Registry: REH-61204). The study was conducted in accordance with the Declaration of Helsinki.

Study population and group definition

We recruited 127 children from Ban Huai Rai Samakee elementary school in Chiang Rai, Thailand. Recruitment of subjects was conducted by voluntary participation through the school's administration. Parents provided informed consent prior to participation. Demographic data collection included gender, age, weight, height, ethnicity, history of birth delivery mode and feeding practice (representing the feeding mode in infancy) (Supplementary File 2). The children's weight and height were measured by class instructors. Information on birth delivery method and feeding type were collected through child self-report and/or parental-report surveys. Body mass index (BMI) derived from the weight (kg) and height (m²) ratio was converted into gender-specific z-scores for BMI-for-age according to BMI cut-offs for children (5-19 years) set by World Health Organization (de Onis et al., 2007). Z-scores for BMI-for-age were classified into 5 groups: severe thinness (SVThinness; < -3 SD; n = 1), thinness (≥ -3 SD to < -2 SD; n = 5), normal weight (≥ -2 SD to ≤ +1.0 SD; n = 83), overweight (OV; > +1SD to ≤ +2SD; n = 20), and obese (OB; > +2 SD; n = 18) (Fig. S1). Age groups were defined according to interquartile range (IQR: 25%, 50%, and 75%): age_A (≤ 8.05 years; n = 32), age_B (8.05 < years < 11.06; n = 61), and age_C (≥ 11.06 years; n = 34). Five ethnic groups were recorded in this study: Akha (n = 39), Chinese (n = 34), Lahu (n = 5), Thai (n = 19), and Thai Yai (n = 30). Birth delivery mode comprised vaginal delivery (n = 85) and cesarean section (n = 42). Feeding types were categorized into three groups: breastfeeding (n = 98), formula feeding (n = 20), and mixed feeding (n = 9).

Dietary information

Dietary habits of children were surveyed using a Thai short dietary behaviors screener developed by *Let's Get Healthy!* For use in Thai ("LGH20 Food Behaviors Screener, Thai"; OHSU Institutional Review Board protocol #3694). The screener included 20 questions that grouped participants across five dietary behavior categories: Healthy eating behavior (HEB), fruits and vegetables (FV), high sugar foods and beverages (HSFB), high salt foods (HSF), and high fat foods (HFF) (Supplementary File 3A). Answer options measuring frequency of consumption were divided into four levels: Frequently (daily), sometimes (weekly), infrequently (monthly), and never. The scores for HEB and FV were assigned as 3 (daily), 2 (weekly), 1 (monthly), or 0 (never). The responses for HSFB, HSF, and HFF were reverse scored. Total component scores (i.e., a sum score for each category) were divided into quartiles to assign levels of risk (low, low to moderate, moderate to high, and high (Supplementary File 3B and 3C). Highest frequencies of HEB and FV consumption would be associated with low risk, while high risk would characterize

children eating mostly HSFB, HSF, and HFF. The instrument screens general dietary behaviors, but does not provide a quantitative assessment of portion size and frequency to permit quantification of a specific food or nutrient intake. Instead, intake rankings permit categorization of individuals according to overall dietary behaviors, such as healthy eating or high consumption of fatty foods.

Sample collection, DNA extraction, and quantitative PCR

Fecal samples were collected from all children in sterilized containers and immediately frozen at -80°C . Microbiota DNA was extracted from fecal samples using the innuPREP Stool DNA Kit (Analytik Jena Biometra, Jena, Germany) according to the manufacturer's instructions. DNA yield and purity were determined using the Take 3 Micro-Volume Plate (Biotek, Winooski, VT, USA). Absolute quantification of bacteria was then conducted by qPCR using Real-Time Thermal Cyclers CFX96 Touch™ (Bio-Rad, Singapore). Primers targeting microbiota 16S rRNA genes used in this study are summarized in Table S1. Reactions consisted of template DNA, forward and reverse primers, 1X SYBR green (2X SensiFAST™ SYBR No-ROX mix, BIOLINE, UK), and nuclease-free water. The assay conditions and calculations of microbiota copy numbers were performed according to previously described protocol (Chumponsuk et al., 2021). The average estimates of microbiota abundance by converting CT values were expressed as logarithmic copy number per gram of wet weight feces.

Statistical analysis

A sum score for dietary behaviors of children was visualized as a bar plot with ggplot2 (Wickham H, 2009). The association between dietary behavior was assessed using Spearman's rank correlation and visualized with corrrplot version 0.84 (Wei & Simko, 2017). Normality and homogeneity of variance were tested by Shapiro-Wilk test and Levene's test (stats package version 4.0.3) (R Core Team, 2020a). Differences in the abundance of gut microbiota (Supplementary File 4) between groups (dietary behaviors and demographic factors) were determined by one-way ANOVA, Welch's t-test, and Kruskal-Wallis rank sum test ($p < 0.05$) followed by multiple comparisons using Tukey's HSD test, pairwise t-tests, and Dunn's test with Benjamini-Hochberg (BH) p -value correction (hereafter referred to as q -value) (stats package version 4.0.3 (R Core Team, 2020a) and FSA package version 0.8.31 (Ogle, Wheeler & Dinno, 2020). Association between birth delivery mode and the abundance of gut microbiota was determined by permutational multivariate analysis of variance (PERMANOVA) with adjustment for covariates (age and feeding type). Group dispersions based on a maximum distance were measured by betadisper with 999 permutations in the R package [vegan \(version 2.5-6\)](#) (Oksanen et al., 2016). Multiple factor analysis (MFA) was performed to evaluate the influence of host variables (dietary behaviors and demographic factors) on variations of gut microbiota using FactorMine R version 2.3 (Lê, Josse & Husson, 2008). Contribution of variables to the data set was visualized with Factoextra version 1.0.7 (Kassambara & Mundt, 2020). To investigate the most relevant features (microbiota taxa) in characterizing each host factor, Partial Least Squares-

Discriminant Analysis (PLS-DA) was carried out by the mixOmics package version 6.12.2 (Rohart et al., 2017). Canonical mode with 100 iterations was used as a parameter for classifying classes (groups of samples). Receiver operating characteristic curve (ROC curve) and area under the curve (AUC) were also calculated to examine the validity of supervised classification results. Predicted scores of categorical outcomes were compared between one class versus the others by Wilcoxon test (Rohart et al., 2017). The classification accuracy of PLS-DA models is interpreted as follows: no discrimination (AUC 0.5), low discrimination (AUC 0.6 to 0.7), acceptable (AUC 0.7 to 0.8), excellent (AUC 0.8 to 0.9), and outstanding (AUC > 0.9) (Lobo, Jiménez-valverde & Real, 2008; Mandrekar, 2010). All analyses were performed in R software version 4.3 (R Core Team, 2020b). A more detailed explanation of multivariate analyses is described in Supplementary File 5.

Results

Dietary behaviors

The frequencies of dietary behaviors of children varied greatly in their score value (Fig. S2). To determine their relationship between diet behaviors, we performed a correlation analysis based on Spearman's rank correlation coefficient. After multiple testing corrections using the Benjamini-Hochberg method, we found that high sugar foods and beverages consumption were significantly correlated with high salty foods consumption ($\rho = 0.39$, $q < 0.0001$) and high fat foods ($\rho = 0.25$, $q = 0.01$, Fig. S3). A positive association between high salt and high fat behaviors was also detected ($\rho = 0.27$, $q = 0.01$). Moreover, the fruits and vegetables consumption were negatively correlated with every dietary behavior except for those associated with healthy eating behaviors ($\rho = 0.2$, $q = 0.04$). This healthy eating behavior was negatively correlated with consumption of fatty foods ($\rho = -0.23$, $q = 0.02$). Despite the strength of association being considerably weak, the results identified a trend in children reporting high unhealthy foods consumption (e.g., HSFB, HSF, HFF) also reporting low healthy foods behaviors (HEB and FV).

Gut microbiota associated with dietary behaviors

MFA constructed by integration of dietary behaviors and abundance of gut microbiota revealed variation in gut microbiota profiles of children (Supplementary File 6A). *Bacteroides* was highly correlated with dimension 1 (Dim 1; $r = 0.91$, $p < 0.0001$), followed by *Gammaproteobacteria* ($r = 0.90$, $p < 0.0001$) and total bacteria ($r = 0.89$, $p < 0.0001$) (Fig. 1A). Variation in the abundances of these taxa was best explained by HFF behaviors, with an increasing trend in microbial abundances indicated in HFF-low risk (coordinate = 1.43, $p = 0.02$; Fig. 1B). In Dim 2, the clusters were separated according to the number of individuals distributed in each diet category. *Ruminococcus* ($r = -0.21$, $p = 0.02$) and *Akkermansia* ($r = -0.26$, $p < 0.01$) described the distribution of HFF-low risk in Dim 3 (coordinate = 1.83, $p < 0.0001$) and Dim 4 (coordinate = 1.46, $p < 0.001$), respectively (Fig. 1C). Both genera were decreased in individuals with low HFF behaviors (Figure 1D). Other diet behaviors (HEB, FV, HSF, and HSFB), however, had a

lower coordinate on the first, third and fourth axes of the MFA factor map than HFF suggesting less contribution of these dietary behaviors to the variation in gut microbiota profiles of children in this study.

Further analysis of the association between gut microbiota and dietary behaviors using PLS-DA also identified relevant features (i.e., microbiota taxa) in classifying dietary behaviors based on the level of consumption. Total bacteria and *Gammaproteobacteria*, highly contributed to discrimination of samples along component 1 (Dim 1), and strongly characterized HFF-low risk (AUC = 0.81, $p = 0.04$, Figs. 2A, S4A). The abundances of total bacteria ($p = 0.02$, Fig. 2C), *Gammaproteobacteria* ($p < 0.0001$, Fig. 2E), and *Lactobacillus* ($p = 0.01$, Fig. 2D) were significantly different among HFF categories. After adjustment by multiple comparisons using the benjamini-Hochberg method, *Gammaproteobacteria* significantly increased in children with low HFF compared to those with high HFF ($q < 0.001$), moderate to high risk HFF ($q < 0.001$), and the highest HFF consumption ($q = 0.03$). In component 2 of PLS-DA for HFF, *Lactobacillus* and *Ruminococcus* were the most discriminative bacteria in children reporting low HFF consumption (AUC = 0.82, $p = 0.03$, Figs. 2B, S4B). However, a significant difference in the abundance of *Lactobacillus* was detected between low HFF to moderate and high HFF consumption after adjustment ($q = 0.05$, Fig. 2D). Moreover, PLS-DA for fruits and vegetables (FV) consumption showed that total bacteria, *Prevotella*, *Bacteroides*, and *Faecalibacterium* were the top three bacteria that separated children with high FV (FV-low risk) from those with lower FV consumption (low to moderate risk and moderate to high risk FV consumption) (Fig. S5A; AUC = 0.66, $p = 0.01$). The abundance of total bacteria was also significantly higher in those reporting high FV as compared to those reporting lower FV consumption ($q = 0.04$, Fig. S5C). Nevertheless, the classification was better in the second component where *Roseburia* and *Ruminococcus* contributed to high FV consumption (Figs. S5B, S5D; AUC = 0.70, $p < 0.001$). For high salty foods (HSF), *Faecalibacterium* characterized moderate to high HSF consumption followed by *Bifidobacterium* and *Roseburia* on component 2, whereas *Lactobacillus* was associated with low HSF consumption (Fig. S6; AUC = 0.70, $p < 0.001$). When considering healthy eating behavior (HEB) and consumption of high sugar foods and beverages (HSFB), the supervised analysis yielded no discrimination between classes (AUC < 0.6, $p > 0.05$). Regarding the observed variability of individuals with different levels of dietary consumption, both MFA and PLS-DA analyses suggested that the consumption of high fat foods had the highest influence on the gut microbiota abundances in children.

Associations between demographic factors and gut microbiota in children

Analysis of gut microbiota with integration of six demographic factors (gender, age, BMI Z-score, ethnicity, birth delivery records, and feeding type) illustrated differences of association patterns with the gut microbiota among the demographic categories (Fig. 3 and Supplementary File 7). The MFA explained 18.6% and 8.3% of the variance in Dim 1 and Dim 2, respectively (Fig. S7A). *Bacteroides*, *Gammaproteobacteria*, and total bacteria were the top three variables

that described individual variation in Dim 1 ($p < 0.0001$, Fig. S7B). Their abundances decreased in underweight (Thinness) and Thai ethnicity children, while an increasing trend contributed to normal weight (Table 1, Figs. 3A, 3B). In Dim 2, *Lactobacillus* mainly described the variation of individual profiles grouped by delivery mode ($R^2 = 0.37$, $p < 0.0001$), BMI z-score ($R^2 = 0.34$, $p < 0.0001$), and age tertile ($R^2 = 0.31$, $p < 0.0001$) (Figs. 3C, 3D). Abundance of *Lactobacillus* decreased in children delivered vaginally, and in those of normal weight, and oldest age (age_C) but increased in those delivered by cesarean section, OB (obese), and youngest age (age_A). Increased *Gammaproteobacteria* in middle age students (age_B), underweight (Thinness), and Thai ethnicity characterized Dim 3 (respectively, Figs. S8A–S8C), while this bacteria was decreased in Lahu ethnicity and oldest age (age_C). Variation of individuals in Dim 4 was mainly described by Firmicutes and ethnicity ($R^2 = 0.45$, $p < 0.0001$): the abundance of these bacteria was increased in children of Lahu and Thai ethnicity, but decreased in those of Chinese and Akha ethnicity. In Dim 5, OV (increased) had a contrasting profile of *Ruminococcus* to OB (decreased). A similar pattern of this bacterial genus was also described for mixed feeding (increased) and formula feeding (decreased) (Figs. S8D). Considering all demographic variables included in the MFA, gender had the least contribution to the variation in microbial abundances, while other factors were associated with subtle differences, which may be of relevance to profiling the gut microbiota in children.

Correlation between gut microbiota and BMI z-score

Comparisons of gut microbiota across BMI z-score groups showed a significant difference in the abundances of Firmicutes ($p < 0.01$) and *Ruminococcus* ($p = 0.01$) (Figs. 4A–4B). After adjustment by multiple comparisons, the abundance of Firmicutes and *Ruminococcus* were significantly higher in students of normal weight ($q < 0.01$) and OV ($q < 0.05$) compared to obese. The supervised analysis also indicated discriminations of these microbiota taxa between BMI z-score groups (Fig. 4C). Normal BMI was highly associated with increased abundance of *Ruminococcus* (component 1: AUC = 0.63, $p = 0.02$, Figs. 4D, S9), while low abundance of Firmicutes and *Ruminococcus* in OB discriminated them from those in other groups (component 1: AUC = 0.68, $p = 0.02$, Figs. 4D, S9). A decreasing trend in the abundance of *Gammaproteobacteria* and *Bacteroides* contributed to thinness (AUC = 0.76, $p = 0.04$, Fig. 4E), however, their association was less important.

Relation between gut microbiota abundance with age group

Differences in the abundance of Firmicutes ($p = 0.05$) and *Bifidobacterium* ($p = 0.02$) were detected at different age tertiles of school-aged children (Fig. S10). Significant increase in Firmicutes ($q = 0.04$) was found in oldest children over 11 years of age (age_C) compared to those in age_B (8.05 < age < 11.06 years) (Fig. 5A). Age_C also showed greater abundance of *Bifidobacterium* than age_A ($q = 0.02$) and age_B ($q = 0.04$) groups (Fig. 5B). Further evaluation of age-associated differences in the gut microbiota of children by PLS-DA revealed certain microbiota taxa contributing to the discrimination. The PLS-DA plot displayed variations

in microbiota profiles according to age tertile (Fig. 5C). Feature classification indicated Firmicutes, *Bacteroides*, *Roseburia*, *Prevotella*, and *Ruminococcus* as the top five more abundant microbiota taxa in the oldest school children (age_C) (Fig. 5D). Of these, Firmicutes had the highest contribution to age_C in component 1 (AUC = 0.62, $p = 0.03$, Fig. S11). The model supports that children over 11 years of age have a higher abundance of this microbiota phylum.

Comparison of microbiota abundance in different delivery mode~~*Prevotella* was enriched in children that were delivered vaginally~~

In this study, we included a record of childbirth to determine its association with the gut microbiota. A comparison of means between the two birth delivery modes showed no significant difference in their abundance of microbiota (Fig. S12). When we performed PERMANOVA with adjustment for covariates (age and feeding type; Supplementary File 8), the test indicated that birth delivery mode was significantly associated with the abundance of *Prevotella* ($p = 0.03$, Fig. S13A), while no influence of sample dispersions was detected ($p = 0.08$, and S13B). Further analyses using PLS-DA also revealed variations of gut microbiota abundance based on birth delivery mode (Fig. S13C). The enrichment of *Prevotella* in vaginal delivery was clearly distinguished from that observed in those delivered by cesarean section (component 1: AUC = 0.69, $p < 0.001$, Figs. S13D, S13E).

Differences in the abundance of gut microbiota of children associated with feeding type

The gut microbiota profile of children varied across feeding types (Fig. S14). A comparison of microbiota abundances among the three feeding types (breastfeeding, formula feeding, and mixed feeding) showed significant differences in the abundance of Firmicutes and *Bifidobacterium* ($p < 0.05$). Both bacterial taxa were significantly higher in mixed feeding children than in those receiving formula feeding ($q < 0.05$, Figs. 6C, 6D). Abundance of *Bifidobacterium* was significantly increased in children breastfed as infants compared to those formula fed as infants ($q = 0.01$, Fig. 6D). We then analyzed the association between gut microbiota and feeding type using PLS-DA to identify key-discriminatory microbiota taxa. Although the PLS-DA components displayed overlapping clusters (Fig. 6A), several differentially abundant bacteria that contributed to the variation in feeding type were indicated (Fig. 6B). The classification model suggested that *Faecalibacterium* (Fig. 6E), Firmicutes, *Roseburia* and *Bifidobacterium* increased following mixed feeding in component 1 (AUC = 0.60, $p = 0.31$, Fig. S15A). In component 2, a similar pattern was observed for Firmicutes and *Ruminococcus* (AUC = 0.71, $p = 0.03$), whereas *Gammaproteobacteria* increased in formula fed children (AUC = 0.79, $p < 0.0001$) (Figs. 15B, S16A).

The influence of gender towards gut microbiota profile in children

Comparisons of the abundances of gut microbiota found no significant difference between gender (Figs. S17). This factor, however, accounted for 47% of the variation in microbial abundances observed in component 1 of PLS-DA plots of gender (Figs. S18A). Classification

models further demonstrated that *Lactobacillus*, *Gammaproteobacteria*, and *Bacteroides* were the top three microbiota taxa associated with girls (Fig. S18B). Based on assessing the discriminative ability of these microbiota taxa for each class (categorical variables), the test indicated that the outcome had poor discrimination capacity to distinguish between classes (AUC < 0.6, $p > 0.05$, Figs. S18C). The model indicated that gender did not influence the gut microbiota profiles of children in this study.

Correlation between ethnicity and gut microbiota composition

No significant differences in the abundances of gut microbiota were found across ethnicity (Fig. S19). When we included ethnicity in the PLS-DA, the model demonstrated the association of this variable with the gut microbiota of children. While *Bacteroides* was the discriminative bacteria in Lahu ethnicity, *Gammaproteobacteria* was enriched in individuals of Akha ethnicity (component 1; AUC < 0.6, $p > 0.05$, Figs. S20A, S20B). However, a higher AUC value was obtained in component 2, where *Akkermansia* discriminated Thai Yai from others (AUC = 0.68, $p < 0.01$), while *Faecalibacterium* and *Roseburia* were the most discriminative bacteria in Akha ethnicity (AUC = 0.67, $p < 0.01$, Figs. S20C, S20D). These models implied that ethnicity had a slight influence on the gut microbiota of school-aged children.

Discussion

Our qPCR study of eleven microbiota taxa and total bacteria in the stool of 127 school-aged children revealed associations with dietary behaviors and demographic factors. Supervised analyses suggested that the gut microbiota profile was influenced by high fat foods consumption and the demographic factors of BMI z-score, age, mode of birth delivery, method of milk delivery, and ethnicity. Gender was not linked to variation in the gut microbiome in this study.

The human diet has a significant impact on the gut microbiota, as changes in the composition in response to food consumption have been extensively documented (David et al., 2014; Zmora, Suez & Elinav, 2019; Leeming et al., 2019). Here, we observed a significant increase in the abundance of *Gammaproteobacteria* in children who scored lower on high fat foods behaviors (characterized by the frequency of eating high fat foods, fried foods as well as food or dessert which was cooked with coconut milk, butter, or margarine). Previous studies in animals, and an in vitro model of the human gut using 16S rRNA gene sequencing, also showed that the abundance of bacteria belonging to the *Gammaproteobacteria* class increased following a high-fat diet consumption (Lecomte et al., 2015; Agans et al., 2018). To the best of our knowledge, a similar finding has not been previously reported based on qPCR. Whether decrease in abundance of *Gammaproteobacteria* is indeed associated with high fat diets in this population would require additional validation using larger sized cohorts, and ideally combination of both high throughput sequencing and qPCR for comparability across studies. An increased abundance of *Lactobacillus* and *Ruminococcus* were also associated with lower HFF consumption. The abundances of these

bacteria ~~was shown to be~~ disturbed in animal models fed ~~with~~ a high-fat diet (Daniel et al., 2014; Resch et al., 2021), which indicates that these taxa may not grow well in the gut in the presence of high-fat foods. Hence, in this study, lower reporting of HFF dietary behaviors among children perhaps favors the growth of these bacteria. Furthermore, ~~the frequent~~ reporting of high fruits and vegetables ~~consumption~~ seemed ~~to highly promote the enrichment of~~ ~~appeared~~ ~~associated with~~ total bacteria and *Prevotella* levels. These results are consistent with previous studies of ~~the gut microbiome of Asian populations, which address~~ the influence of shifting from traditional to Western diets (high-fat/low-fiber) on the gut microbiota ~~of Asian populations~~. *Prevotella*-type taxa were overrepresented in the gut of school-aged children in rural Thailand when compared with children in urban areas, who harbored more *Bacteroides*-type bacteria. Frequency of fruit and vegetable intake may therefore support different enterotypes, as was already reported for Filipino children from rural Baybay as well as for Thai vegetarians (Ruengsomwong et al., 2014; Nakayama et al., 2015, 2017). Outside of Asia, *Prevotella* dominates the microbiota communities of rural African children consuming diets high in fiber compared to those of European children (De Filippo et al., 2010). These converging findings emphasize the importance of a fiber-rich food diet ~~in order~~ to colonize the gut with *Prevotella* (Kisuse et al., 2018). High salty foods (HSF) intake affected the abundances of gut microbiota. Specifically, the butyrate producer *Faecalibacterium* and *Lactobacillus* were differentially associated with reported moderate to high salty foods and low salty foods consumption, respectively. A similar contrasting profile between *Roseburia* (another butyrate-producing bacterium) and *Lactobacillus* was previously shown in mice fed high- and low-salt diets. The former was enriched in mice fed high-salt diet (Wang et al., 2017), while the proportion of the latter was significantly reduced (Wang et al., 2017; Miranda et al., 2018). A similar finding has also been noted in humans (Wilck et al., 2017). These findings suggest that high salt food consumption impacts the abundance of specific gut microbiota members.

Changes in the gut microbiota profile of children have been associated with BMI status classified based on both centiles (Bervoets et al., 2013) and z-scores (Goloso-Gubat et al., 2020; Shin & Cho, 2020). In this study, a low abundance of Firmicutes and *Ruminococcus* was associated with obesity, while normal and overweight children had a high abundance of these bacteria. These findings are in contrast to previous studies based on 16S rRNA sequencing, whereby obese children had a high abundance of Firmicutes (Da Silva, Monteil & Davis, 2020), while *Ruminococcus* was nearly depleted in overweight/obese when compared to normal-weight children (Karvonen et al., 2019). A longitudinal study conducted in school-aged children with dietary records also highlighted a decrease of *Ruminococcaceae* in children who developed obesity and had a high calorie intake (high carbohydrate/high fat and high protein/high fat) associated with the obese status, (Rampelli et al., 2018). These findings suggest food intake and weight gain could contribute to variability in the gut microbiome (Rampelli et al., 2018). Despite unequal sample sizes and a different dietary assessment method herein, most obese children (72%) consumed high fat foods quite frequently (moderate to high risk) (Fig. S21), while only

33% ate fruits and vegetables (Fig. S22). Thus, the observed differences in microbiota abundance in our study were likely influenced by high-calorie diets, although further study with more participants, longer follow-up periods, and more extensive microbiome profiling is needed to verify this hypothesis.

The abundance of *Bifidobacterium* can vary across the stages of life (Arbolea et al., 2016; Saturio et al., 2021) and this genera is often enriched in the gut microbiota of children compared to those in (Derrien et al., 2019). A similar trend was also detected in our study with a high level of *Bifidobacterium* among school-aged children grouped by age tertile. The abundance of *Bifidobacterium* was significantly high in children aged over 11 years. Moreover, children in previous studies that fell into the same age categories as in this study also had a higher fecal concentration of *Bifidobacterium* compared to those that were older aged (Agans et al., 2011; Hollister et al., 2015; Zhong et al., 2019). Concerning age variables, a gap may exist with these findings as we stratified individuals by quantile ranges. Whether or not the presence of this particular bacterium is associated with age, changes in *Bifidobacterium* levels from childhood to adolescence using narrow-age ranges may be worth investigating in order to better comprehend this relationship.

Both birth delivery method and feeding type appears to have a strong influence on the early-life gut microbiota (Cukrowska et al., 2020; Mitchell et al., 2020). The impact of the former has been shown in a large longitudinal analysis of gut microbiota from 600 newborns and 175 mothers, which denoted significant differences in the composition of gut microbiota between cesarean section born and vaginally delivered infants (Shao et al., 2019). The latter type of birth was associated with a high abundance of *Prevotella*, as shown in newborns and during the first two years of life (Dominguez-Bello et al., 2010; Bokulich et al., 2016). Although our study was conducted in school-aged children, enrichment of this genus was still observed in those who were born vaginally. This result implies that the impact of method of delivery may continue beyond infancy. Furthermore, we found that the abundance of *Bifidobacterium* was lowest in children who were formula fed as infants when compared with children who were either breast fed or mixed fed during infancy. *Bifidobacterium* abundance is increased in the gut of breast-fed infants rather than in those that are formula-fed. It has been speculated that the bacterium utilizes human milk oligosaccharides (HMO) (Lee et al., 2015; Forbes et al., 2018; Lawson et al., 2020). Our data suggests that a lack of exposure to breast milk at an early age may reduce abundance of gut *Bifidobacterium*, while mixed-feeding may stabilize the abundance close to breastfeeding. As time progresses, however, many other factors including the influence of one's diet is expected to also influence the makeup of one's gut microbiome.

Ethnicity introduces variations in the gut microbiota profiles through diet (Khine et al., 2019; Dwiyanto et al., 2021). Considering the small sample size of our study, however, our findings did not have an adequate power to identify the associations between the consumption of ethnic-

based diets and the abundance of gut microbiota. We did, however, observe a trend when discriminating between ethnic groups. For instance, two genera within the phylum Firmicutes (*Faecalibacterium* and *Roseburia*) were associated with children of Akha ethnicity, whereas *Akkermansia* was mainly found to associate with the Thai-Yai ethnic group. These results are inconclusive due to a lack of dietary data relating to ethnic cultural practices.

Although our study demonstrated the independent effect of each host factor on the gut microbiota, our results should be interpreted with caution. Major limitations include the lack of sample size estimation and data on cultural practices (e.g., traditional diets, lifestyle, etc.). Since recruitment of subjects was based on voluntary participation, the number of subcategories was not homogeneous. In this regard, inter-individual variation was investigated using multivariate statistical analyses with all concerned factors. The same method has been implemented in our previous works (Gruneck et al., 2020; Chumponsuk et al., 2021). Moreover, we were unable to collect data on cultural practices due to the language barriers, which might link to dietary behaviors of these school-aged children. Both limitations described above serve to limit our ability to explore correlations between important risk factors and the gut microbiome of school-aged children. One such potential confounding factor, physical activity, should also be included with future studies to better understand the role this plays together with BMI and diet.

Conclusions

This study highlights how diet influences gut microbiota. A high abundance of *Gammaproteobacteria* was noted in children who reported consumption of few high fat foods. Demographic factors such as BMI z-score, age, and feeding type also demonstrated their potential associations with gut microbiota. Obese children were characterized by a low abundance of *Ruminococcus*. Those over 11 years of age were found to have a high level of *Bifidobacterium*, whereas this abundance decreased in children with a history of formula feeding. Moreover, birth mode and ethnicity displayed a trend towards the enrichment of gut microbiota. Considering all host variables, gender was not a determinant of microbiota profiles in this study.

Acknowledgements

The authors would like to thank all volunteers for providing fecal samples and dietary data. We would like to thank Mr. Channarong Wanthanjai for technical assistance.

Abbreviations

HEB – Healthy eating behavior
FV – fruits and vegetables
HSFB – high sugar foods and beverages
HSF – high salt foods
HFF – high fat foods
OV – overweight

OB – obese
age_A – age of children ≤ 8.05 years
age_B – age of children between 8.05 and 11.06 years ($8.05 < \text{years} < 11.06$)
age_C – age of children ≥ 11.06 years

References

- Agans R, Gordon A, Kramer DL, Perez-Burillo S, Rufián-Henares JA, Paliy O. 2018. Dietary fatty acids sustain the growth of the human gut microbiota. *Applied and Environmental Microbiology* 84(24):e01525-18. DOI: 10.1128/AEM.01525-18.
- Agans R, Rigsbee L, Kenche H, Michail S, Khamis HJ, Paliy O. 2011. Distal gut microbiota of adolescent children is different from that of adults. *FEMS Microbiology Ecology* 77(2):404–412. DOI: 10.1111/j.1574-6941.2011.01120.x.
- Arbolea S, Watkins C, Stanton C, Ross RP. 2016. Gut *Bifidobacteria* populations in human health and aging. *Frontiers in Microbiology* 7:1024. DOI: 10.3389/fmicb.2016.01204.
- Bervoets L, Van Hoorenbeeck K, Kortleven I, Van Noten C, Hens N, Vael C, Goossens H, Desager KN, Vankerckhoven V. 2013. Differences in gut microbiota composition between obese and lean children: A cross-sectional study. *Gut Pathogens* 5(4):10. DOI: 10.1186/1757-4749-5-10.
- Bokulich NA, Chung J, Battaglia T, Henderson N, Jay M, Li H, Lieber AD, Wu F, Perez-Perez GI, Chen Y, Schweizer W, Zheng X, Contreras M, Dominguez-Bello MG, Blaser MJ. 2016. Antibiotics, birth mode, and diet shape microbiome maturation during early life. *Science Translational Medicine* 8(343):343ra82. DOI: 10.1126/scitranslmed.aad7121.
- Cho KY. 2021. Lifestyle modifications result in alterations in the gut microbiota in obese children. *BMC Microbiology* 21(1). DOI: 10.1186/s12866-020-02002-3.
- Chumponsuk T, Gruneck L, Gentekaki E, Jitprasertwong P, Kullawong N, Nakayama J, Popluechai S. 2021. The salivary microbiota of Thai adults with metabolic disorders and association with diet. *Archives of Oral Biology* 122:105036. DOI: 10.1016/j.archoralbio.2020.105036.
- Coker MO, Laue HE, Hoen AG, Hilliard M, Dade E, Li Z, Palys T, Morrison HG, Baker E, Karagas MR, Madan JC. 2021. Infant Feeding-feeding Alters-alters the Longitudinal longitudinal impact-impact of Birth-birth Mode-mode on the Development-development of the Gut-gut Microbiota-microbiota in the First-first Year-year of Lifelife. *Frontiers in Microbiology* 12:642197. DOI: 10.3389/fmicb.2021.642197.
- Cukrowska B, Biera JB, Zakrzewska M, Klukowski M, Maciorkowska E. 2020. The Relationship between the Infant Gut Microbiota and Allergy. The Role of *Bifidobacterium* breve and Prebiotic Oligosaccharides in the Activation of Anti-Allergic Mechanisms in Early Life. *Nutrients* 12(4):946. DOI: 10.3390/nu12040946.
- Daniel H, Gholami AM, Berry D, Desmarchelier C, Hahne H, Loh G, Mondot S, Lepage P, Rothballer M, Walker A, Böhm C, Wenning M, Wagner M, Blaut M, Schmitt-Kopplin P, Kuster B, Haller D, Clavel T. 2014. High-fat diet alters gut microbiota physiology in mice. *The ISME Journal* 8(2):295–308. DOI: 10.1038/ismej.2013.155.
- David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling A V, Devlin AS, Varma Y, Fischbach MA, Biddinger SB, Dutton RJ, Turnbaugh PJ. 2014. Diet rapidly and reproducibly alters the human gut microbiome. *Nature* 505(7484):559–563. DOI: 10.1038/nature12820.

Derrien M, Alvarez AS, de Vos WM. 2019. The Gut Microbiota in the First Decade of Life. *Trends in Microbiology* 27(12):997–1010. DOI: 10.1016/j.tim.2019.08.001.

Dominguez-Bello MG, Costello EK, Contreras M, Magris M, Hidalgo G, Fierer N, Knight R. 2010. Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proceedings of the National Academy of Sciences of the United States of America* 107(26):11971–11975. DOI: 10.1073/pnas.1002601107.

Dwiyanto J, Hussain MH, Reidpath D, Ong KS, Qasim A, Lee SWH, Lee SM, Foo SC, Chong CW, Rahman S. 2021. Ethnicity influences the gut microbiota of individuals sharing a geographical location: a cross-sectional study from a middle-income country. *Scientific Reports* 11(1):2618. DOI: 10.1038/s41598-021-82311-3.

van den Elsen LWJ, Garssen J, Burcelin R, Verhasselt V. 2019. Shaping the gut microbiota by breastfeeding: The gateway to allergy prevention? *Frontiers in Pediatrics* 7:47. DOI: 10.3389/fped.2019.00047.

Escobar JS, Klotz B, Valdes BE, Agudelo GM. 2015. The gut microbiota of Colombians differs from that of Americans, Europeans and Asians. *BMC Microbiology* 14(1):311. DOI: 10.1186/s12866-014-0311-6.

De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S, Collini S, Pieraccini G, Lionetti P. 2010. Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proceedings of the National Academy of Sciences of the United States of America* 107(33):14691–14696. DOI: 10.1073/pnas.1005963107.

Forbes JD, Azad MB, Vehling L, Tun HM, Konya TB, Guttman DS, Field CJ, Lefebvre D, Sears MR, Becker AB, Mandhane PJ, Turvey SE, Moraes TJ, Subbarao P, Scott JA, Kozyrskyj AL, Anand SS, Befus AD, Brauer M, Brook JR, Chen E, Cyr MM, Daley D, Dell SD, Denburg JA, Duan QL, Eiwegger T, Grasmann H, HayGlass K, Hegele RG, Holness DL, Hystad P, Kobor M, Kollmann TR, Laprise C, Lou WYW, Macri J, Miller G, Pare P, Ramsey C, Ratjen F, Sandford A, Scott J, Silverman F, Simons E, Takaro T, Tebbutt SJ, To T. 2018. Association of exposure to formula in the hospital and subsequent infant feeding practices with gut microbiota and risk of overweight in the first year of life. *JAMA Pediatrics* 172(7):e181161. DOI: 10.1001/jamapediatrics.2018.1161.

Ghosh TS, Rampelli S, Jeffery IB, Santoro A, Neto M, Capri M, Giampieri E, Jennings A, Candela M, Turrone S, Zoetendal EG, Hermes GDA, Elodie C, Meunier N, Brugere CM, Pujos-Guillot E, Berendsen AM, De Groot LCPGM, Feskens EJM, Kaluza J, Pietruszka B, Bielak MJ, Comte B, Maijo-Ferre M, Nicoletti C, De Vos WM, Fairweather-Tait S, Cassidy A, Brigidi P, Franceschi C, O'Toole PW. 2020. Mediterranean diet intervention alters the gut microbiome in older people reducing frailty and improving health status: The NU-AGE 1-year dietary intervention across five European countries. *Gut* 69(7):1218–1228. DOI: 10.1136/gutjnl-2019-319654.

Goloso-Gubat MJ, Ducarmon QR, Tan RCA, Zwartink RD, Kuijper EJ, Nacis JS, Santos NLC. 2020. Gut Microbiota and Dietary Intake of Normal-Weight and Overweight Filipino Children. *Microorganisms* 8(7):1015. DOI: 10.3390/microorganisms8071015.

Gruneck L, Kullawong N, Kespechara K, Popluechai S. 2020. Gut microbiota of obese and diabetic Thai subjects and interplay with dietary habits and blood profiles. *PeerJ* 8. DOI: 10.7717/peerj.9622.

Hollister EB, Riehle K, Luna RA, Weidler EM, Rubio-Gonzales M, Mistretta TA, Raza S, Doddapaneni H V., Metcalf GA, Muzny DM, Gibbs RA, Petrosino JF, Shulman RJ,

- Versalovic J. 2015. Structure and function of the healthy pre-adolescent pediatric gut microbiome. *Microbiome* 3(1):36. DOI: 10.1186/s40168-015-0101-x.
- Karvonen AM, Sordillo JE, Gold DR, Bacharier LB, O'Connor GT, Zeiger RS, Beigelman A, Weiss ST, Litonjua AA. 2019. Gut microbiota and overweight in 3-year old children. *International Journal of Obesity* 43(4):713–723. DOI: 10.1038/s41366-018-0290-z.
- Kassambara A, Mundt F. 2020. factoextra: Extract and Visualize the Results of Multivariate Data Analyses. *R package version 1.0.7*.
- Khine WWT, Zhang Y, Goie GJY, Wong MS, Liong M, Lee YY, Cao H, Lee YK. 2019. Gut microbiome of pre-adolescent children of two ethnicities residing in three distant cities. *Scientific Reports* 9(1):7831. DOI: 10.1038/s41598-019-44369-y.
- Kim KA, Gu W, Lee IA, Joh EH, Kim DH. 2012. High Fat Diet-Induced Gut Microbiota Exacerbates Inflammation and Obesity in Mice via the TLR4 Signaling Pathway. *PLoS ONE* 7(10):e47713. DOI: 10.1371/journal.pone.0047713.
- Kisuse J, La-ongkham O, Nakphaichit M, Therdtatha P, Momoda R, Tanaka M, Fukuda S, Popluechai S, Kespechara K, Sonomoto K, Lee YK, Nitisinprasert S, Nakayama J. 2018. Urban diets linked to gut microbiome and metabolome alterations in children: A comparative cross-sectional study in Thailand. *Frontiers in Microbiology* 9:1345. DOI: 10.3389/fmicb.2018.01345.
- Lawson MAE, O'Neill IJ, Kujawska M, Gowrinadh Javvadi S, Wijeyesekera A, Flegg Z, Chalklen L, Hall LJ. 2020. Breast milk-derived human milk oligosaccharides promote Bifidobacterium interactions within a single ecosystem. *ISME Journal* 14(2):635–648. DOI: 10.1038/s41396-019-0553-2.
- Lê S, Josse J, Huisson F. 2008. FactoMineR: An R package for multivariate analysis. *Journal of Statistical Software* 25:1–18.
- Lecomte V, Kaakoush NO, Maloney CA, Raipuria M, Huinao KD, Mitchell HM, Morris MJ. 2015. Changes in gut microbiota in rats fed a high fat diet correlate with obesity-associated metabolic parameters. *PLoS ONE* 10(5):e0126931. DOI: 10.1371/journal.pone.0126931.
- Lee SA, Lim JY, Kim BS, Cho SJ, Kim NY, Kim O Bin, Kim Y. 2015. Comparison of the gut microbiota profile in breast-fed and formula-fed Korean infants using pyrosequencing. *Nutrition Research and Practice* 9(3):242–248. DOI: 10.4162/nrp.2015.9.3.242.
- Leeming ER, Johnson AJ, Spector TD, Roy CIL. 2019. Effect of diet on the gut microbiota: Rethinking intervention duration. *Nutrients* 11(12):2862. DOI: 10.3390/nu11122862.
- Li N, Yan F, Wang N, Song Y, Yue Y, Guan J, Li B, Huo G. 2020. Distinct Gut Microbiota and Metabolite Profiles Induced by Different Feeding Methods in Healthy Chinese Infants. *Frontiers in Microbiology* 11:714. DOI: 10.3389/fmicb.2020.00714.
- Liu K, Zhang Y, Li Q, Li H, Long D, Yan S, Huang W, Long R, Huang X. 2020. Ethnic Differences Shape the Alpha but Not Beta Diversity of Gut Microbiota from School Children in the Absence of Environmental Differences. *Microorganisms* 8(2):254. DOI: 10.3390/microorganisms8020254.
- Lobo JM, Jiménez-valverde A, Real R. 2008. AUC: A misleading measure of the performance of predictive distribution models. *Global Ecology and Biogeography* 17(2):145–151. DOI: 10.1111/j.1466-8238.2007.00358.x.
- Mandrekar JN. 2010. Receiver Operating Characteristic Curve in Diagnostic Test Assessment. *Journal of Thoracic Oncology* 5(9):1315–1316. DOI: <https://doi.org/10.1097/JTO.0b013e3181ec173d>.
- Miranda PM, De Palma G, Serkis V, Lu J, Louis-Auguste MP, McCarville JL, Verdu EF, Collins

SM, Bercik P. 2018. High salt diet exacerbates colitis in mice by decreasing *Lactobacillus* levels and butyrate production. *Microbiome* 6(1):57. DOI: 10.1186/s40168-018-0433-4.

Mitchell CM, Mazzoni C, Hogstrom L, Bryant A, Bergerat A, Cher A, Pochan S, Herman P, Carrigan M, Sharp K, Huttenhower C, Lander ES, Vlamakis H, Xavier RJ, Yassour M. 2020. Delivery Mode Affects Stability of Early Infant Gut Microbiota. *Cell Reports Medicine* 1(9):100156. DOI: 10.1016/j.xcrm.2020.100156.

Nakayama J, Watanabe K, Jiang J, Matsuda K, Chao SH, Haryono P, La-Ongkham O, Sarwoko MA, Sujaya IN, Zhao L, Chen KT, Chen YP, Chiu HH, Hidaka T, Huang N-X, Kiyohara C, Kurakawa T, Sakamoto N, Sonomoto K, Tashiro K, Tsuji H, Chen MJ, Leelavatcharamas V, Liao CC, Nitisinprasert S, Rahayu ES, Ren FZ, Tsai YC, Lee YK. 2015. Diversity in gut bacterial community of school-age children in Asia. *Scientific Reports* 5(1):8397. DOI: 10.1038/srep08397.

Nakayama J, Yamamoto A, Palermo-Conde LA, Higashi K, Sonomoto K, Tan J, Lee YK. 2017. Impact of westernized diet on gut microbiota in children on Leyte island. *Frontiers in Microbiology* 8:197. DOI: 10.3389/fmicb.2017.00197.

Niu J, Xu L, Qian Y, Sun Z, Yu D, Huang J, Zhou X, Wang Y, Zhang T, Ren R, Li Z, Yu J, Gao X. 2020. Evolution of the Gut Microbiome in Early Childhood: A Cross-Sectional Study of Chinese Children. *Frontiers in Microbiology* 11:439. DOI: 10.3389/fmicb.2020.00439.

Odamaki T, Kato K, Sugahara H, Hashikura N, Takahashi S, Xiao JZ, Abe F, Osawa R. 2016. Age-related changes in gut microbiota composition from newborn to centenarian: A cross-sectional study. *BMC Microbiology* 16(1):90. DOI: 10.1186/s12866-016-0708-5.

Ogle D, Wheeler P, Dinno A. 2020. FSA: Fisheries Stock Analysis. *R package version 0.8.30*.

Oksanen J, Kindt R, Pierre L, O'Hara B, Simpson GL, Solymos P, Stevens MH. HH, Wagner H, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MH. HH, Wagner H. 2016. *vegan: Community Ecology Package*, R package version 2.4-0. *R package version 2.2-1*.

de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. 2007. Development of a WHO growth reference for school-aged children and adolescents. *Bulletin of the World Health Organization* 85(9):660–667. DOI: 10.2471/blt.07.043497.

Ooraikul B, Sirichote A, Siripongvutikorn S. 2008. Southeast Asian Diets and Health Promotion. In: *Wild-Type Food in Health Promotion and Disease Prevention*. Totowa, NJ: Humana Press, 515–533. DOI: 10.1007/978-1-59745-330-1_36.

R Core Team. 2020a. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

R Core Team. 2020b. R: A Language and Environment for Statistical Computing.

Rampelli S, Guenther K, Turrone S, Wolters M, Veidebaum T, Kourides Y, Molnár D, Lissner L, Benitez-Paez A, Sanz Y, Fraterman A, Michels N, Brigidi P, Candela M, Ahrens W. 2018. Pre-obese children's dysbiotic gut microbiome and unhealthy diets may predict the development of obesity. *Communications Biology* 1(1):222. DOI: 10.1038/s42003-018-0221-5.

Resch C, Parikh M, Austria JA, Proctor SD, Netticadan T, Blewett H, Pierce GN. 2021. The influence of diet and sex on the gut microbiota of lean and obese jcr:La-cp rats. *Microorganisms* 9(5):1037. DOI: 10.3390/microorganisms9051037.

Reyman M, van Houten MA, van Baarle D, Bosch AATM, Man WH, Chu MLJN, Arp K, Watson RL, Sanders EAM, Fuentes S, Bogaert D. 2019. Impact of delivery mode-associated gut microbiota dynamics on health in the first year of life. *Nature*

Communications 10(1):4997. DOI: 10.1038/s41467-019-13014-7.
 Rohart F, Gautier B, Singh A, Lê Cao K-A. 2017. mixOmics: An R package for 'omics feature selection and multiple data integration. *PLOS Computational Biology* 13(11):e1005752-.
 Ruengsomwong S, Korenori Y, Sakamoto N, Wannissorn B, Nakayama J, Nitisinprasert S. 2014. Senior thai fecal microbiota comparison between vegetarians and non-vegetarians using PCR-DGGE and real-time PCR. *Journal of Microbiology and Biotechnology* 24(8):1026–1033. DOI: 10.4014/jmb.1310.10043.
 Rutayisire E, Huang K, Liu Y, Tao F. 2016. The mode of delivery affects the diversity and colonization pattern of the gut microbiota during the first year of infants' life: A systematic review. *BMC Gastroenterology* 16(1):86. DOI: 10.1186/s12876-016-0498-0.
 Salminen S, Gibson GR, McCartney AL, Isolauri E. 2004. Influence of mode of delivery on gut microbiota composition in seven year old children. *Gut* 53(9):1388–1389. DOI: 10.1136/gut.2004.041640.
 Saturio S, Nogacka AM, Suárez M, Fernández N, Mantecón L, Mancabelli L, Milani C, Ventura M, de los Reyes-Gavilán CG, Solís G, Arbolea S, Gueimonde M. 2021. Early-life development of the bifidobacterial community in the infant gut. *International Journal of Molecular Sciences* 22(7):3382. DOI: 10.3390/ijms22073382.
 Shao Y, Forster SC, Tsaliki E, Vervier K, Strang A, Simpson N, Kumar N, Stares MD, Rodger A, Brocklehurst P, Field N, Lawley TD. 2019. Stunted microbiota and opportunistic pathogen colonization in caesarean-section birth. *Nature* 574(7776):117–121. DOI: 10.1038/s41586-019-1560-1.
 Shin S, Cho KY. 2020. Altered Gut Microbiota and Shift in Bacteroidetes between Young Obese and Normal-Weight Korean Children: A Cross-Sectional Observational Study. *BioMed Research International* 2020:1–19. DOI: 10.1155/2020/6587136.
 Da Silva CC, Monteil MA, Davis EM. 2020. Overweight and Obesity in Children Are Associated with an Abundance of Firmicutes and Reduction of Bifidobacterium in Their Gastrointestinal Microbiota. *Childhood Obesity* 16(3):204–210. DOI: 10.1089/chi.2019.0280.
 Singh RK, Chang HW, Yan D, Lee KM, Ucmak D, Wong K, Abrouk M, Farahnik B, Nakamura M, Zhu TH, Bhutani T, Liao W. 2017. Influence of diet on the gut microbiome and implications for human health. *Journal of Translational Medicine* 15(1):73. DOI: 10.1186/s12967-017-1175-y.
 Voreades N, Kozil A, Weir TL. 2014. Diet and the development of the human intestinal microbiome. *Frontiers in Microbiology* 5:494. DOI: 10.3389/fmicb.2014.00494.
 Wan Y, Wang F, Yuan J, Li J, Jiang D, Zhang J, Li H, Wang R, Tang J, Huang T, Zheng J, Sinclair AJ, Mann J, Li D. 2019. Effects of dietary fat on gut microbiota and faecal metabolites, and their relationship with cardiometabolic risk factors: a 6-month randomised controlled-feeding trial. *Gut* 68(8):1417–1429. DOI: 10.1136/gutjnl-2018-317609.
 Wang C, Huang Z, Yu K, Ding R, Ye K, Dai C, Xu X, Zhou G, Li C. 2017. High-Salt Diet Has a Certain Impact on Protein Digestion and Gut Microbiota: A Sequencing and Proteome Combined Study. *Frontiers in Microbiology* 8. DOI: 10.3389/fmicb.2017.01838.
 Wei T, Simko V. 2017. R package "corrplot": Visualization of a Correlation Matrix. DOI: <https://github.com/taiyun/corrplot>.
 Wickham H. 2009. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag, New York.
 Wilck N, Matus MG, Kearney SM, Olesen SW, Forslund K, Bartolomeus H, Haase S, Mähler A, Balogh A, Markó L, Vvedenskaya O, Kleiner FH, Tsvetkov D, Klug L, Costea PI,

Sunagawa S, Maier L, Rakova N, Schatz V, Neubert P, Frätzer C, Krannich A, Gollasch M,
 Grohme DA, Côte-Real BF, Gerlach RG, Basic M, Typas A, Wu C, Titze JM, Jantsch J,
 Boschmann M, Dechend R, Kleinewietfeld M, Kempa S, Bork P, Linker RA, Alm EJ,
 Müller DN. 2017. Salt-responsive gut commensal modulates TH17 axis and disease. *Nature*
 551(7682):585–589. DOI: 10.1038/nature24628.
 Yang R, Gao R, Cui S, Zhong H, Zhang X, Chen Y, Wang J, Qin H. 2019. Dynamic signatures
 of gut microbiota and influences of delivery and feeding modes during the first 6 months of
 life. *Physiological Genomics* 51(8):368–378. DOI: 10.1152/physiolgenomics.00026.2019.
 Yasir M, Angelakis E, Bibi F, Azhar EI, Bachar D, Lagier JC, Gaborit B, Hassan AM, Jiman-
 Fatani AA, Alshali KZ, Robert C, Dutour A, Raoult D. 2015. Comparison of the gut
 microbiota of people in France and Saudi Arabia. *Nutrition and Diabetes* 5(4):e153. DOI:
 10.1038/nutd.2015.3.
 Zhong H, Penders J, Shi Z, Ren H, Cai K, Fang C, Ding Q, Thijs C, Blaak EE, Stehouwer CDA,
 Xu X, Yang H, Wang J, Wang J, Jonkers DMAE, Masclee AAM, Brix S, Li J, Arts ICW,
 Kristiansen K. 2019. Impact of early events and lifestyle on the gut microbiota and
 metabolic phenotypes in young school-age children. *Microbiome* 7(1):2. DOI:
 10.1186/s40168-018-0608-z.
 Zmora N, Suez J, Elinav E. 2019. You are what you eat: diet, health and the gut microbiota.
Nature Reviews Gastroenterology and Hepatology 16(1):35–56. DOI: 10.1038/s41575-018-
 0061-2.

Commented [LMG1]:

