

Two continents and two names for a Neotropical colletid bee species (Hymenoptera: Colletidae: Neopasiphaeinae): *Hoplocolletes ventralis* (Friese, 1924)

Eduardo A B Almeida, Fábio B Quinteiro

Neopasiphaeinae bees (Apoidea: Colletidae) are well known for their Amphinotic distribution in the Australian and Neotropical regions. Affinities between colletid taxa in Australia and South America have been speculated for decades, and have been confirmed by recent phylogenetic hypotheses that indicate a biogeographic scenario compatible with a trans-Antarctic biotic connection during the Paleogene. Despite this proximity, no species occurs on both sides of the Pacific Ocean, but the Neotropical species *Hoplocolletes ventralis* (Friese, 1924), which was described as an Australian taxon due to an error in the specimen labels. This mistake was recognized by C.D. Michener 50 years ago. We herein report that the same labeling problem also happened with *Dasycolletes chalceus* Friese, 1924, which remained as a tentatively placed species in the Australian genus *Leioproctus* until now. Moreover, *Dasycolletes chalceus* is interpreted as a synonym of *Dasycolletes ventralis*. We also provide a revised diagnosis for *Hoplocolletes*, describe the male of *H. ventralis* in detail for the first time, including a comparative study of its genitalia and associated sterna.

Two continents and two names for a Neotropical colletid bee species (Hymenoptera: Colletidae: Neopasiphaeinae): *Hoplocolletes ventralis* (Friese, 1924)

EDUARDO A. B. ALMEIDA* & FÁBIO B. QUINTEIRO

Laboratório de Biologia Comparada e Abelhas (LBCA) - Departamento de Biologia, Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto (FFCLRP), Universidade de São Paulo. 14040-600. Ribeirão Preto, SP. Brazil

* Corresponding author. E-mail: eduardo@ffclrp.usp.br

Abstract

Neopasiphaeinae bees (Apoidea: Colletidae) are well known for their Amphinotic distribution in the Australian and Neotropical regions. Affinities between colletid taxa in Australia and South America have been speculated for decades, and have been confirmed by recent phylogenetic hypotheses that indicate a biogeographic scenario compatible with a trans-Antarctic biotic connection during the Paleogene. Despite this proximity, no species occurs on both sides of the Pacific Ocean, but the Neotropical species *Hoplocolletes ventralis* (Friese, 1924), which was described as an Australian taxon due to an error in the specimen labels. This mistake was recognized by C.D. Michener 50 years ago. We herein report that the same labeling problem also happened with *Dasycolletes chalceus* Friese, 1924, which remained as a tentatively placed species in the Australian genus *Leioproctus* until now. Moreover, *Dasycolletes chalceus* is interpreted as a synonym of *Dasycolletes ventralis*. We also provide a revised diagnosis for *Hoplocolletes*, describe the male of *H. ventralis* in detail for the first time, including a comparative study of its genitalia and associated sterna.

Keywords: Apoidea, Australia, biogeography, Brazil, systematics, taxonomy

Introduction

Affinities between taxa of Colletidae distributed in Australia and South America have been speculated for decades (Michener, 1965, 1989), and have been confirmed by recent phylogenetic hypotheses that indicate a biogeographic scenario compatible with a trans-Antarctic biotic connection during the Paleogene (Almeida et al., 2012). *Dasycolletes ventralis* Friese, 1924 was described as an Australian colletid species based on a single female specimen labeled as having been collected in Sydney (Australia). The species actually is endemic to Brazil, and there is no species occurring in Australia that could be confused with it. The confusion certainly results from an error in the label, as concluded by Michener (1965: p.41), an interpretation followed by subsequent authors (e.g., Moure, Graf & Urban, 2007; Rasmussen & Ascher, 2008). After the species description, it was moved to the genus *Paracolletes* by Cockerell (1929), and later placed in *Leioproctus* (*Hoplocolletes*), created by Michener (1965) to accommodate it considering its clear affinities to other taxa classified as *Leioproctus*, but also recognizing its uniqueness (see also Michener 1989, 2007). *Hoplocolletes* remains as monotypic taxon in Neopasiphaeinae (Colletidae), but has often been classified as genus instead of subgenus (e.g., Silveira, Melo & Almeida, 2002; Moure, Graf & Urban, 2007; Almeida & Danforth, 2009; Almeida et al., 2012), a position followed in this paper.

Hoplocolletes ventralis has been recorded in three states in southeastern Brazil: Espírito Santo, Minas Gerais, Rio de Janeiro (Silveira, Melo & Almeida, 2002; Moure, Graf & Urban,



2007). *Hoplocolletes* remains a poorly known genus, with relatively little distributional information, male undescribed for decades since *H. ventralis* was recognized as a new species, host-plant preferences unknown, and the only piece of bionomical information for this species is that it is a soil nesting bee (EAB Almeida, pers.observation). The phylogenetic affinities of *Hoplocolletes* and other neopasiphaeinae taxa were uncertain until molecular phylogenetic hypotheses placed this taxon in a clade ~~also~~ comprising *Eulonchopria* and *Nomiocolletes* (Almeida & Danforth, 2009; Almeida et al., 2012). Michener (1989: p.630) suggested that *Hoplocolletes* could be part of a "Basal Group", characterized by the fully developed sternal scopa. Based on the phylogenetic hypotheses currently available, it seems that this scopa arose multiple times in the Neopasiphaeinae clade, since *Hoplocolletes*, *Cephalocolletes*, *Reedapis*, and *Tetraglossula* are not close relatives (Almeida & Danforth, 2009; Almeida et al., 2012).

The aim of this work is twofold: resolve a taxonomic problem related to a new synonymy involving *Hoplocolletes ventralis* and increase the knowledge about the morphology and distinctiveness of *Hoplocolletes*, particularly by providing a novel description of the male genital complex for this species. We report that the abovementioned labeling problem that made the taxonomic history of *Hoplocolletes ventralis* problematic also happened with *Dasycolletes chalceus* Friese, 1924, which remained as a tentatively placed species in the Australian genus *Leioproctus* until now (Michener, 1965; Cardale, 1993; Almeida, 2008; Rasmussen & Ascher, 2008). *Dasycolletes chalceus* is interpreted as a synonym of *Dasycolletes ventralis*.

Material & Methods

Part of the material studied is deposited in the Entomological Collection "Prof. J.M.F.Camargo" [RPSP] in Departamento de Biologia (FFLRP/USP, Ribeirão Preto, Brazil).

A male specimen of *Hoplocolletes ventralis* was obtained as a loan from Entomological Collection "Pe.J.S.Moure" [DZUP], Departamento de Zoologia (UFPR, Curitiba, Brazil), and the female type specimen of *Dasycolletes chalceus* Friese, 1924 was studied and photographed at the entomological collection of Museum für Naturkunde [ZMB] (Berlin, Germany). Photographs of the female specimen of *Dasycolletes ventralis* Friese, 1924, deposited at the American Museum of Natural History (AMNH) collection, were kindly taken and made available for this study.

The general morphological terminology follows Michener (2007). Antennal flagellomeres are indicated as F1, F2, etc.; metasomal terga and sterna, respectively, as T1 to T7, and S1 to S8. The density of punctation and intervals between the punctures are based on relative puncture diameter, pd (e.g., <1pd: less than 1x the puncture diameter between the punctures). Color images were obtained on a Zeiss Axiocam 206 color camera associated to a Zeiss Discovery.V12 stereomicroscope, or with an AmScope MU1000A Digital Camera adapted onto a Leica MZ6 stereomicroscope; pictures were assembled with the software Helicon Focus 6.2.

Results

~~The *Dasycolletes chalceus* Friese, 1924 case.~~ The species *Dasycolletes chalceus* was not studied after its original description. It was described in the same publication and same page as *Dasycolletes ventralis* (Friese, 1924: p. 218). After 1924, it was only mentioned in catalogues and revisionary works (e.g., Michener, 1965; Cardale, 1993; Almeida, 2008; Rasmussen & Ascher, 2008), but the type specimen was never studied again. The only exemplar of *Dasycolletes chalceus* located and bearing Friese's original labels is deposited in ZMB (Fig.1). It

clearly has all diagnostic characters for *Hoplocolletes* as currently conceived, and no differences were found in relation to *Hoplocolletes ventralis* either. The only known specimen of *Dasycolletes ventralis* bearing Friese's original labels is in the American Museum of Natural History collection (New York, USA) (Fig. 2) and it is the same female studied by Michener (1965) that lead him to conclude it was not an Australian taxon, as indicated by the collecting labels, but a specimen probably collected in Brazil. The interpretation of Friese's types is a controversial subject and it is likely that the AMNH specimen is a duplicate, not the primary type (Rasmussen & Ascher, 2008; J.S.Ascher, pers.comm.). But, so far, it is the only specimen labeled by Friese himself as *Dasycolletes ventralis* available for study. It is worth noting that both specimens were probably collected together, have locality labels that are identical, "Australia \\\ Sydney \\\ 14.9/06". The collector's name is lacking in the *D. chalceus* specimen label but not in this species' description (Friese, 1924: p.218): "von Sydney im September, Frank leg."

***Hoplocolletes ventralis* (Friese, 1924)**

Dasycolletes ventralis Friese, H. (1924) [218].

Type data: holotype AMNH <F>.

Type locality: 'Australia, Sydney'.

Dasycolletes chalceus Friese, H. (1924) [218].

Type data: holotype ZMB <F>.

Type locality: 'Australia, Sydney'.

Description of male: Approximate body length: 10 mm; length of forewing: 7.7 mm; maximum width of metasoma (T2): 2.5 mm. *Color:* predominantly black; apical half of mandible, ventral

116 surface of F2-F11, tibiae, femora, trochanters, S2-S3, apical margins of terga dark reddish brown.
 117 Tarsi light brown. Tegula, pterostigma and wing veins dark brown; wing membrane brown
 118 infumated. *Pubescence*: predominantly pale yellowish or cream on entire body. Face and
 119 pronotal lobe with abundant pubescence; clypeus with decumbent to semidecumbent pilosity (0.5
 120 mm in length), more erect and shorter on paraocular area and frons (0.3-0.45 mm in length).
 121 Mesoscutum with scarce pilosity. Lateral pilosity of mesepisternum semidecumbent and sparse
 122 (0.25-0.35 mm in length). *Integumental surface*: coarse and dense on clypeus (≤ 1 pd), finer and
 123 denser frons (<1 pd), on vertex variable (denser medially, sparser [≤ 1 pd] laterally as well as on
 124 gena) integument smooth and shiny between punctures; coarse and dense on mesosoma, sparser
 125 on disc of mesoscutum and scutellum, and inferiorly on mesepisternum; metapostnotum smooth
 126 and shiny, delimited from pronotum by a pit-row; T1 smooth and shiny, with very sparse (2-7
 127 pd) moderately coarse punctation, transversal line of barely aligned punctures delimiting
 128 marginal region of T1; on T2 slightly denser than on T1, but punctation leaving broad shiny
 129 areas as well; T3 and T4 with basal portion finely and densely punctated, sparser and coarser
 130 distad. *Structure* (measurements in mm): head about 1.1x wider than long (2.66:2.43); inner
 131 orbits converging below (upper to lower interorbital distance, 1.76:1.47), inner margin almost
 132 straight; eye about 3.6x longer than its maximum width in frontal view (1.76:0.48), in lateral
 133 view about 1.2x wider than gena (0.74:0.64). Vertex well developed above ocelli (distance
 134 between upper margin of lateral ocellus and vertex = 0.53), comparable to ocelloocular distance
 135 (0.51); interocellar distance = 0.14; diameter of median ocellus = 0.25. Approximate length of
 136 antenna = 4.0, length and maximum width of scape = 0.73, 0.2; of pedicel = 0.16; of F1 = 0.19;
 137 F2 about 1.5x wider than long (0.18:0.27); F3 about 1.5x longer than wide (0.30:0.21).
 138 Mesoscutum length = 1.83, intertegular distance = 1.75. Genital capsule and male S7 and S8 as

illustrated in Figs 5-6 (see discussion about the male terminalia below, in the 'Revised Diagnosis' for *Hoplocolletes*).

***Hoplocolletes* Michener, 1965**

Revised diagnosis for the genus (characters apply to both sexes unless otherwise stated).

Length 10-12 mm. Body black to dark brown; head and mesosoma with conspicuous coarse punctation; T1 and T2 largely impunctate, smooth and shining (remaining terga rather finely and closely punctate); pubescence short, sparse, blackish to dark brown on female (light yellow to fulvous on male), except on hind legs and metasomal sterna where there are long, pale hairs; metasomal hair bands absent, male clypeus with plumose and semidecumbent pubescence. Mandible with an ordinary preapical tooth. Inner orbits subparallel (female, Figs 1D, 2C) or converging below (male, Fig.3C). Facial fovea absent; clypeus weakly convex; labrum with apical margin concave medially, elevated zone highest medially, occupying about basal half of labrum (Michener, 1989: Fig.7q). Preoccipital carina absent; malar area linear; clypeus little protuberant. Male flagellum elongate (approximately 3.0 mm long), F2 longer than wide. Vertex produced behind ocelli and eyes (Figs 1A,D, 2A,C, 3A,B,C). Apex of scape reaching upper margin of median ocellus (Figs1D, 2C); antennae arising about middle of face. Dorsolateral angle of pronotum low, rounded, scarcely evident; metapostnotum smooth, marginal line pitted, its basal part slightly longer than metanotum. Femoral scopa sparse, formed by long delicate branched hairs, those behind corbicula and on trochanter long but simple. Tibial scopal hairs dividing to form few major branches. Female basitibial plate ~~of female~~ distinct, hairs short, appressed, different from those of adjacent areas, marginal carinae clearly exposed. Inner hind tibial spur of female coarsely pectinate with 5-6 teeth (Michener, 1989: Fig.7q). Forewing with

three submarginal cells, second much shorter than third and receiving recurrent vein beyond middle (Figs 1F, 3A); basal vein of forewing meeting cu-v (Fig. 1); stigma large, long, not quite parallel sided, two-thirds as long as costal side of marginal cell, marginal cell longer than distance between its apex and wing apex. T1 dorsally approximately twice wider than long; S3-S5 of female with dense, long (shorter than exposed part of sternum), pale yellow, simple hairs (some hooked at tips) forming band occupying apical half of each sternum, female S2 with similar but sparser hair band (Figs 1C, 2D); S3-S5 of male with a longer hairs near apical margin, S5 with distinct apical fringe.

Male genital capsule and associated sterna of *Hoplocolletes ventralis* are illustrated in Figs 5-6 along with exemplar species of two other neopasiphaeinae genera: *Nomiocolletes joergenseni* (Fries, 1908) and *Reedapis semicyanea* (Spinola, 1851). According to the phylogenetic hypotheses of Almeida & Danforth (2009) and Almeida et al. (2012), *Hoplocolletes* and *Nomiocolletes* are closely related lineages, whereas *Reedapis* is part of a more distantly related clade. The interpretation of homologies for the male terminalia in this comparative context makes the understanding of relevant characters of *Hoplocolletes* more defensible. Apical process of male S7 ~~of~~ comprising two lobes on each side: one apicolateral more developed and hairier (Fig. 5: lateral lobe - LLb) ~~and one~~ closer to the base of this process (Fig. 5: basal lobe - BLb), *Nomiocolletes* is distinctive ~~for~~ having a bilobed lateral lobe; *Hoplocolletes* does not have apical protuberances as found in other Neopasiphaeinae (Fig. 5: apical lobe - ALb); apodeme of S7 relatively long in relation to the apical process. Median process of male S8 (Fig. 5: MPr) similar in length to the remainder of S8; spiculum ordinary (not as produced as in *Nomiocolletes* or *Reedapis*). Gonobase of male genitalia (Fig. 6: Gbs) less than 1/4 of total length of genital capsule; gonostylus and gonocoxite (Fig. 6: Gns, Gcx) fused on dorsal surface but separable on

ventral surface, apex of gonostylus rounded and not bent ventrad (directed mesad and ventrad in the other two species); apex of penis valve (Fig.6: PV) bent ventrad, ventral spine well-developed (Fig.6: SPV).

Acknowledgements

We are grateful to Gabriel A. R. Melo (Universidade Federal do Paraná - [DZUP]) who kindly provided information about a locality where *Hoplocolletes ventralis* is known to occur in Minas Gerais state, and loaned a male specimen used in this study. We are also indebted to Frank Koch for hosting EABAlmeida during a visit to the Museum für Naturkunde [ZMB], and to Hadel H. Go and Jerome G. Rozen, Jr. (American Museum of Natural History - AMNH) for taking photographs and making them available for this work. Our thanks to John S. Ascher (National University of Singapore) and Gabriel A. R. Melo for valuable discussion on the systematics of neopasiphaeine bees and the work of H.Friese, and to Diego S. Porto for critical comments on this manuscript.

REFERENCES

- Almeida EAB. 2008. Revised species checklist of the Paracolletinae (Hymenoptera, Colletidae) of the Australian Region, with the description of new taxa. *Zootaxa* 1891:1-24.
- Almeida EAB, Danforth BN. 2009. Phylogeny of colletid bees (Hymenoptera: Colletidae) inferred from four nuclear genes. *Molecular Phylogenetics and Evolution* 50:290-309. DOI:10.1016/j.ympev.2008.09.028

- 206 Almeida EAB, Pie MR, Brady, SG, Danforth BN. 2012. Biogeography and diversification of
207 colletid bees (Hymenoptera: Colletidae): Emerging patterns from the Southern End of the
208 World. *Journal of Biogeography* 39:526-544. DOI:10.1111/j.1365-2699.2011.02624.x
- 209 Cardale JC. 1993. Hymenoptera: Apoidea. In: Houston WWK, Maynard GV, eds. *Zoological*
210 *Catalogue of Australia, Vol. 10*. Canberra: Government Publishing Service, ix + 405 pp.
- 211 Cockerell TDA. 1929. Bees, chiefly Australian species, described or determined by Dr. H.
212 Friese. *American Museum Novitates* 343:1-20.
- 213 Danforth BN, Cardinal S, Praz C, Almeida EAB, Michez D. 2013. The impact of molecular data
214 on our understanding of bee phylogeny and evolution. *Annual Review of Entomology* 58:57-
215 78. DOI:10.1146/annurev-ento-120811-153633
- 216 Danforth BN, Sipes S, Fang J, Brady SG. 2006. The history of early bee diversification based on
217 five genes plus morphology. *Proceedings of the National Academy of Sciences USA*
218 103:15118-15123. doi:10.1073/pnas.0604033103
- 219 Friese H. 1924. Ueber die Bienen Australiens. *Konowia: Zeitschrift für systematische*
220 *Insektenkunde* 3:216-249.
- 221 Michener CD. 1965. A classification of the bees of the Australian and South Pacific regions.
222 *Bulletin of the American Museum of Natural History* 130:1-362 + 15 plates.
- 223 Michener CD. 1989. Classification of the American Colletinae. *University of Kansas Science*
224 *Bulletin* 53:622-703.
- 225 Michener CD. 2007. *The Bees of the World* (2nd Ed.). Baltimore: John Hopkins University Press.
- 226 Moure JS, Graf V, Urban D. 2007. Paracolletini Cockerell, 1934. In: Moure JS, Urban D, Melo
227 GAR, eds. *Catalogue of Bees (Hymenoptera, Apoidea) in the Neotropical Region*. Curitiba:
228 Sociedade Brasileira de Entomologia, 723-743.

- 229 Rasmussen C, Ascher JS. 2008. Heinrich Friese (1860–1948): Names proposed and notes on a
- 230 pioneer melittologist (Hymenoptera, Anthophila). *Zootaxa* 1833: 1-118.
- 231 Silveira FA, Melo GAR, Almeida EAB. 2002. *Abelhas Brasileiras: Sistemica e Identificação*.
- 232 Belo Horizonte: F.A.Silveira.
- 233 Spinola M. 1851. Hymenópteros, in C. Gay, *Historia Fisica y Politica de Chile, Zoologia, Vol. 6*.
- 234 Paris: Casa del autor, 153-569.

FIGURE CAPTIONS

Figure 1. Female specimen of *Dasycolletes chalceus* Friese, 1924 deposited at the Museum für Naturkunde collection [ZMB] (photo credit: Eduardo A.B. Almeida): **A** - Dorsal habitus (scale bar = 1 mm), **B** - lateral habitus, **C** - face, **D** - ventral metasomal scopa, **E** - specimen labels.

Figure 2. Female specimen of *Dasycolletes ventralis* of Friese, 1924 deposited at the American Museum of Natural History collection [AMNH] (photo credit: Hadel Go): **A** - Dorsal habitus, **B** - lateral habitus, **C** - face, **D** - magnified view of ventral metasomal scopa, **E** - specimen labels.

Figure 3. Male specimen of *Hoplocolletes chalceus* (Friese, 1924) from Itapina, ES, Brazil [DZUP] (photo credit: Eduardo A.B. Almeida): **A** - Lateral habitus, **B** - dorsal habitus, **C** - face, **D** - mesosoma and anterior metasoma; scale bars = 1 mm.

Figure 4. Comparative morphology of male metasomal sterna S7 and S8 (dorsal views shown on left) of *Hoplocolletes ventralis* (Friese, 1924), *Nomiocolletes joergenseni* (Friese, 1908), and *Reedapis semicyanea* (Spinola, 1851). ALb = apical lobe of S7, BLb = basal lobe of S7, LLb = lateral lobe of S7, LPr = lateral process of S8, MPr = median process of S8; scale bars = 0.5 mm. Cladogram represents a hypothesis for the phylogenetic relationships among these three taxa (Almeida & Danforth, 2009).

Figure 5. Comparative morphology of male genitalia (dorsal views shown on left) of *Hoplocolletes ventralis* (Friese, 1924), *Nomiocolletes joergenseni* (Friese, 1908), and *Reedapis*

259 *semicyanea* (Spinola, 1851). ApP = apodeme of penis valve, Cs = cuspis of volsella, Dg =
 260 digitus of volsella, Gbs = gonobase, Gcx = gonocoxa, Gns = gonostyle, PV = penis valve, SPV =
 261 ventral spine of penis valve; scale bars = 0.5 mm. Cladogram represents a hypothesis for the
 262 phylogenetic relationships among these three taxa (Almeida & Danforth, 2009).

1

Female specimen of *Dasycolletes chalceus* Friese, 1924 deposited at the Museum für Naturkunde collection [ZMB].

Female specimen of *Dasycolletes chalceus* Friese, 1924 deposited at the Museum für Naturkunde collection [ZMB] (photo credit: Eduardo A.B. Almeida): A - Dorsal habitus (scale bar = 1 mm), B - lateral habitus, C - face, D - ventral metasomal scopa, E - labels.



2

Female specimen of *Dasycolletes ventralis* of Friese, 1924 deposited at the American Museum of Natural History collection [AMNH].


Female specimen of *Dasycolletes ventralis* of Friese, 1924 deposited at the American Museum of Natural History collection [AMNH] (photo credit: Hadel Go): A - Dorsal habitus, B - lateral habitus, C - face, D - magnified view of ventral metasomal scopa, E - labels.



3

Male specimen of *Hoplocolletes chalceus* (Fries, 1924) from Itapina, ES, Brazil [DZUP].




Male specimen of *Hoplocolletes chalceus* (Fries, 1924) from Itapina, ES, Brazil [DZUP] (photo credit: Eduardo A.B. Almeida): A - Lateral habitus, B - dorsal habitus, C - face, D - mesosoma and anterior metasoma; scale bars = 1 mm.



4

Comparative morphology of male metasomal sterna S7 and S8 (dorsal views shown on left) of *Hoplocolletes ventralis* (Friese, 1924) and related neopasiphaeine taxa.


Comparative morphology of male metasomal sterna S7 and S8 (dorsal views shown on left) of *Hoplocolletes ventralis* (Friese, 1924), *Nomiocolletes joergenseni* (Friese, 1908), and *Reedapis semicyanea* (Spinola, 1851). ALb = apical lobe of S7, BLb = basal lobe of S7, LLb = lateral lobe of S7, LPr = lateral process of S8, MPr = median process of S8; scale bars = 0.5 mm. Cladogram represents a hypothesis for the phylogenetic relationships among these three taxa (Almeida & Danforth, 2009).




5

Comparative morphology of male genitalia (dorsal views shown on left) of *Hoplocolletes ventralis* (Friese, 1924) and related neopasiphaeinae taxa.


Comparative morphology of male genitalia (dorsal views shown on left) of *Hoplocolletes ventralis* (Friese, 1924), *Nomiocolletes joergenseni* (Friese, 1908), and *Reedapis semicyanea* (Spinola, 1851). ApP = apodeme of penis valve, Cs = cuspis of volsella, Dg = digitus of volsella, Gbs = gonobase, Gcx = gonocoxa, Gns = gonostyle, PV = penis valve, SPV = ventral spine of penis valve; scale bars = 0.5 mm. Cladogram represents a hypothesis for the phylogenetic relationships among these three taxa (Almeida & Danforth, 2009).



Hoplocolletes ventralis



Nomocolletes joergenseni



Reedapis semicyanea