

Gene *ssa-miR-301a-3p* improves rainbow trout (*Oncorhynchus mykiss*) resistance to heat stress by targeting *hsp90b2*

Zhe Liu¹, Fang Ma², Yujun Kang¹ and Xiaoxia Liu¹

¹Gansu Agricultural University, Lanzhou, China

²Tianshui Normal University, Tianshui, China

ABSTRACT

Rainbow trout (*Oncorhynchus mykiss*) is a cold-water fish that is commonly harmed by high temperatures. MicroRNAs (miRNAs) are being investigated intensively because they act as essential metabolic regulators and have a role in the heat stress response. Although there have been numerous studies on rainbow trout heat stress, research on miRNA implicated in rainbow trout heat stress is quite restricted. Rainbow trout were sampled at 18 and 24 °C, respectively, to examine the mechanism of miRNA under heat stress, and we identified a heat stress-induced miRNA, *ssa-miR-301a-3p*, for further investigation based on our bioinformatics analysis of rainbow trout small RNA sequencing data. Bioinformatics research suggested that *hsp90b2* is a probable target gene for *ssa-miR-301a-3p*. QRT-PCR was used to confirm the expression levels of *ssa-miR-301a-3p* and *hsp90b2*. Meanwhile, the dual-luciferase reporter assay was employed to validate the *ssa-miR-301a-3p-hsp90b2* targeted connection. The results indicated that at 24 °C, the relative expression of *ssa-miR-301a-3p* was considerably lower than at 18 °C. On the other hand, *hsp90b2* expression, followed the opposite pattern. The binding of *ssa-miR-301a-3p* to the 3'-UTR of *hsp90b2* resulted in a substantial decrease in luciferase activity. The findings showed that *ssa-miR-301a-3p* was implicated in heat stress, and our findings provide fresh insights into the processes of miRNA in response to heat stress in rainbow trout.

Submitted 26 November 2021

Accepted 1 May 2022

Published 5 July 2022

Corresponding author

Zhe Liu, liuz@gsau.edu.cn

Academic editor

Barbara Nowak

Additional Information and
Declarations can be found on
page 10

DOI 10.7717/peerj.13476

© Copyright

2022 Liu et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Aquaculture, Fisheries and Fish Science, Biochemistry, Genetics, Molecular Biology, Freshwater Biology

Keywords Expression regulation, *ssa-miR-301a-3p*, *hsp90b2*, HEK 293T cells, Rainbow trout

INTRODUCTION

Clusters consisting of 21–23 nucleotides and single-stranded non-coding small RNAs are called miRNAs. They can specifically bind to the 3'-untranslation regions (3'-UTRs) of their target genes by a base pair-mediated fashion, and negatively regulate downstream target gene expression at the post-transcriptional level (Bartel, 2004; Baulcombe, 2004; Carrington & Ambros, 2003; Chen & Meister, 2005; Lee & Ambros, 2001; Matzke et al., 2004). In recent years, a number of studies have focused on the relationship between miRNA and temperature stresses, and many studies have demonstrated that the expression levels of miRNA are regulated by temperature stress (Yu et al., 2010; Zhang et al., 2014; Zhang et al., 2017). The expressions of miR-199a-3p and miR-34a-5p are most

significantly different between heat-tolerant (TOL) mice and heat-intolerant (INT) mice. Other miRNAs including miR-19a, miR-19b, miR-27b, miR-30a-5p, miR-181a, miR-181b, miR-345-3 and miR-1246 were found in the serum of Holstein cows under heat stress (Zheng *et al.*, 2014). Moreover, miR-23a is involved in heat stress, miR-23a can regulate the expression of CDK5, and the decrease of miR-23a expression increases the expression of CDK5 protein to resist heat stress (Morey *et al.*, 2015).

As a member of the miR-130 family, ssa-miR-301a-3p is highly conserved in rats, chickens and cattle and can promote the proliferation, migration and angiogenesis of vascular endothelial cells (Zhou *et al.*, 2012). After water temperature increased, miR-301 expression in gonads of zebrafish was significantly down-regulated (Ikert & Craig, 2020). In the carp study, miR-301a in the liver was found to regulate the perception of cellular nutrients, oxygen, energy levels, and the activation of insulin receptors and insulin-like growth factor 1 receptors when affected by temperature (Sun *et al.*, 2019). In fish, miR-301a-3p is also implicated in the regulation of oxidative stress, inflammation, and apoptosis, the expression level of miR-301a-3p is significantly up-regulated under hypoxia pressure (Jiang & Lv, 2021; Xia *et al.*, 2020). Intriguingly, the expression of miR-301a-3p, as well as the expressions of other miR-130a family members including miR-130b-3p, miR-130a-3p, and miR-301b-3p, were significantly upregulated by heat stress (Zhang *et al.*, 2020). Taken together, data above may suggest a potential relationship among miR-301a-3p and heat stress. Although hundreds of miRNAs have been identified since the discovery of the first member 20 years ago (Bushati & Cohen, 2007), studies on the function of miRNAs in animals under heat stress are limited, especially fish. Based on our bioinformatics analysis of small RNA sequencing data of rainbow trout (Huang *et al.*, 2018), we identified one miRNA, ssa-miR-301a-3p, which was significantly down-regulated after heat stress. In this study, our aim was to verify that ssa-miR-301a-3p is involved in heat stress in rainbow trout. We predicted the target gene of ssa-miR-301a-3p by using bioinformatics analysis to screen RNA-seq data (Li *et al.*, 2017).

Because global warming threatens animal health and animal husbandry economy, heat stress, as one of the important abiotic stresses, interferes with cellular homeostasis and induces physiological and biochemical changes in animals (St-Pierre, Cobanov & Schnitkey, 2003; Quinteiro-Filho *et al.*, 2010). Exposure to high temperatures may alter gene expression, leading to the synthesis of stress-related proteins, such as heat shock proteins (HSPs), which are known as key regulators in the heat stress response (Iba, 2002) and will be up-regulated when the organism suffers high temperatures (Lindquist, 1986). HSPs are highly conservative proteins that are induced when the organism is affected by virus infection, temperature variation, heavy metals, starvation and osmotic shock, hypoxia, and oxidative stress in the environment (Haase & Fitze, 2016; Basu *et al.*, 2002; Cara *et al.*, 2005; Kregel, 2002). Heat shock protein 90 (hsp90) is involved in protein sorting, maturation, stabilization and activation (Jackson, 2013; Hong *et al.*, 2013). In vertebrates, *hsp90* has six major isoforms (*hsp90a1a*, GenBank accession no. KC150880. *hsp90a1b*, GenBank accession no. KC150881. *hsp90a2a*, GenBank accession no. KC150878. *hsp90a2b*, GenBank accession no. KC150879. *hsp90b1*, GenBank accession no. KC150882 and *hsp90b2*, GenBank accession no. KC150883), and a previous study showed

that mRNA level of *hsp90b2* in rainbow trout hepatocytes was significantly increased after heat shock (Jia, Cavileer & Nagler, 2016).

Rainbow trout is a major commercial and cold-water fish species and has a preferred water temperature. The appropriate growth temperature for rainbow trout is 12–18 °C, and the water temperature above 25 °C is perilous (Matthews & Berg, 1997). Our previous studies have found that the growth and development of rainbow trout will be severely affected when water temperature increases (Xia et al., 2017). High temperatures can lead to economic losses due to lower productivity and reproductivity, as well as increased health burden. Heat stress is one of the most characteristic environmental stressors. Individuals respond to heat stress by increasing body temperature, reducing feed intake and changing physiological state, which is called physiological adaptation response. However, other protective measures have been developed in animals to cope with the challenges of heat stress, mainly manifested as altered transcription and gene expression levels. Our previous analysis showed that 24 °C is a key high-temperature point that the organism changed from adaptive regulation to injury (Xia et al., 2018). Thus, we chose 18 and 24 °C as the control and heat treatment temperature, respectively. In this study, we are investigating miRNAs that are induced at optimal (18 °C) and high (24 °C) temperatures in the liver of rainbow trout (Huang et al., 2018). We used bioinformatics software to predict their target genes and found that miR-301a-3p regulate *hsp90b2*. Herein our core objective was to validate if *hsp90b2* is indeed the target genes of miR-301a-3p, and for this purpose, we used the Dual-Luciferase® Reporter Assay System. Subsequently, we combined the results of Gene Ontology (go) and Kyoto Encyclopedia of genes and genomes (KEGG) pathway enrichment (Huang et al., 2018) to analyze the main biological functions of the miR-301a-3p and *hsp90b2*, and to evaluate the related regulatory mechanisms. The objective of this study was to validate targets of ssa-miR-301a-3p and evaluate the possible role of ssa-miR-301a-3p in the liver of rainbow trout under heat stress by repressing *hsp90b2*. The results of this study will provide useful information of miRNA in response to heat stress in rainbow trout liver, and provide a reference for the studies of heat stress of this species in the future.

MATERIALS AND METHODS

Ethical approval

This study was conducted in strict accordance with the guidelines for the Care and Use of Laboratory Animals in China and was approved by the Institutional Ethic Committee of Gansu Agricultural University.

Oncorhynchus mykiss preparation

Healthy rainbow trout (average weight: 400 ± 10.5 g) belonging to full-sibling inbred families were obtained from the trout cage aquafarm in Liujiaxia reservoir (Gansu, China) and temporarily sustained in a 3,000 L aerated water tank at 18 °C for 1 week. A total of 120 healthy fish were chosen for subsequent experiments. Those fish were randomly divided into six groups and followed by an adaptation for another 7 days at 18 °C. Three groups were maintained at 18 °C and the other three groups were treated with a thermal

Table 1 Primers used for the quantitative real-time PCR (qRT-PCR).

Gene name	Accession no.	Primer sequence (5' to 3')
<i>Hsp90b2</i>	KC150883	F: 5'-GTTTGGCGTGGGTTTCTACTCT-3' R: 5'-CCACCTCCTTGACCCTCTTCT-3'
<i>ACTB</i>	AB196465	F: 5'-TGGGGCAGTATGGCTTGTATG-3' R: 5'-CTCTGGCACCCCTAATCACCTCT-3'
Ssa-miR-301a-3p	MIMAT0032593	F: 5'-AGTGCCATAGTATTGTCATAGC-3' R: mRQ 3'primer
<i>U6</i>	XM_021614042	F: 5'-GGAACGATACAGAGAAGATTAGC-3' R: 5'-TGGAACGCTTCACGAATTTGCG-3'

Note:

F, forward; R, reverse.

stress. The temperature system continued to increase with a heating rate of 1 °C/24 h from 18 to 24 °C. After anesthetizing the fish with a lethal dose of MS-222, the liver samples were harvested from female fish at 18 (control group, CL) and 24 °C (heat stress group, HL) respectively. The tissues were immediately flash frozen in liquid nitrogen, and stored at -80 °C until RNA extraction.

Target genes prediction

Based on our miRNA-seq data, we found that the expression level of one miRNA, ssa-miR-301a-3p, was significantly downregulated in rainbow trout livers under heat stress. In order to understand the mechanism of ssa-miR-301a-3p, the potential target genes of ssa-miR-301a-3p were predicted by using TargetScan (www.targetscan.org), PicTar (<https://pictar.mdc-berlin.de/>), and miRanda (<https://cbio.mskcc.org/miRNA2003/miranda.html>) analysis of RNA-seq data, which was the result of our previous research group (Li *et al.*, 2017).

Quantitative real-time RT-PCR analysis

For the ssa-miR-301a-3p, the first-strand cDNA was synthesized from the total RNA using Mir-X miRNA First-Strand Synthesis Kit (TaKaRa, Dalian, China). Specific primer for the ssa-miR-301a-3p was designed and primer sequences were listed in Table 1. The qRT-PCR was performed using Mir-X miRNA qRT-PCR SYBR Kit (TaKaRa, Dalian, China) and LightCycler® 480 Instrument II (Roche, Basel, Switzerland). *U6* was used as an endogenous control. The cycling parameters were as follows: 95 °C for 10 s, 40 cycles at 95 °C for 5 s and 60 °C for 20 s; melting curve analyses at 95 °C for 60 s, 55 °C for 30 s, and 95 °C for 30 s.

For *hsp90b2*, the PrimerScript™RT Reagent Kit with gDNA Eraser (TaKaRa, Dalian, China) was used to synthesize first-strand cDNA, and SYBR® Premix Ex Taq kit (TaKaRa, Dalian, China) and LightCycler® 480 Instrument II were used for qRT-PCR. The *ACTB* was taken as a reference gene. Specific primer for the *hsp90b2* was listed in Table 1. The cycling parameters were as follows: 95 °C for 10 s, 40 cycles at 95 °C for 5 s and 61 °C for 20 s; melting curve analyses at 95 °C for 60 s, 55 °C for 30 s, and 95 °C for 30 s.

```

Hsp90b2: 3' UTR WT  5'-AAAACAAGTACCCTTTTGCCTGA-3'
                    |||||
ssa-miR-301a-3p:  3'-CGAUACUGUUAUGAUAACGUGAC-5'
Hsp90b2: 3' UTR MUT 5'-AAAACAAGTACCCTTAACGTGACA-3'

```

Figure 1 3'-UTR binding site of *hsp90b2* mRNA and ssa-miR-301a-3p.

Full-size  DOI: 10.7717/peerj.13476/fig-1

Construction of recombinant plasmid

3'-UTR segment of the *hsp90b2* with the ssa-miR-301a-3p binding site was cloned into pmirGLO dual-luciferase vector (Promega, Madison, WI, USA), which was digested with restriction enzymes *SacI* and *XhoI*, and produced a wild-type structure pmirGLO-*hsp90b2*-wt-3'-UTR (*hsp90b2* WT) by using ClonExpress® Entry One Step Cloning Kit (GenePharma, Shanghai, China). 3'-UTR mutation sequence of *hsp90b2* pmirGLO-*hsp90b2*-mut-3'-UTR (*hsp90b2* MUT) constructed by binding site (TTGCACTG) within the 3'-UTR of the *hsp90b2* was changed using Site-Directed Mutagenesis Kit (Beyotime, Shanghai, China) and took a place of (AACGTGAC) (Fig. 1). At the same time, reverse complementary sequences of ssa-miR-301a-3p were cloned to pmirGLO dual-luciferase vector to construct positive control (ssa-miR-301a-3p PC), which was transferred to HEK 293T cells and was processed to siRNA (small interference RNA) segments of 21–23 bp in length by Dicer, and then siRNA recognized ssa-miR-301a-3p mimic with homologous sequences through base complementary ligation and mediated ssa-miR-301a-3p mimic degradation. All recombinant plasmids were tested for enzyme digestion and sequencing.

Cell culture and transfection

HEK 293T cells were purchased from cell Bank of Chinese Academy of Sciences. HEK 293T cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) containing 10% fetal bovine serum (FBS) at 37 °C with 5% CO₂ saturated humidity incubator. Data were collected as previously described in Liu *et al.* (2021), ssa-miR-301a-3p mimic and mimic negative-control (NC) were synthesized from GenePharma (Shanghai, China). The sequences are as follows: ssa-miR-301a-3p mimic sense: 5'-CAGUGCAAUAGUAUU GUCAUAGC-3', ssa-miR-301a-3p mimic antisense: 5'-UAUGACAAUACUAUUGCA CUGUU-3'; mimic NC sense: 5'-UUCUCCGAACGUGUCACGUTT-3', mimic NC antisense: 5'-ACGUGACACGUUCGGAGAATT-3'.

The transfection group consisted of mimic NC + *hsp90b2* WT, mimic NC + *hsp90b2* MUT, ssa-miR-301a-3p mimic + *hsp90b2* WT, ssa-miR-301a-3p mimic + *hsp90b2* MUT, mimic NC + ssa-miR-301a-3p PC, and ssa-miR-301a-3p mimic + ssa-miR-301a-3p PC.

Dual-luciferase reporter assay

The HEK 293T cells were seeded at a density of 5.0×10^5 cells per well in a 12-well culture plate, incubated for 24 h before transfection, and then transfected with 10 μ l of ssa-miR-301a-3p mimic, 10 μ l of mimic NC (GenePharma, Shanghai, China), 1.6 μ g of *hsp90b2* WT, 1.6 μ g of *hsp90b2* MUT and 1.6 μ g ssa-miR-301a-3p PC plasmid simultaneously.

Table 2 The target gene sequencing results of ssa-miR-301a-3p.

Gene name	Log ₂ FoldChange	Padj
<i>Hsp90b2</i>	1.5767	1.47E-07
<i>TBCC1</i>	1.8655	0.034978
<i>P4HAI</i>	3.0113	4.41E-09
<i>Hsp90b1</i>	1.2616	0.000139

Note:

Log₂FoldChange: $\log_2(\text{Sample1}/\text{Sample2})$, sample1 represents the readcount value of 24 °C in sequencing results, sample2 represents the readcount value of 18 °C in sequencing results; Padj: corrected pvalue. The smaller the corrected pvalue, the more significant the change in gene expression at 24 °C compared with 18 °C.

Meanwhile, a total of 4 µl of Lipofectamine 2000 (Invitrogen, Waltham, MA, USA) was added to another Eppendorf tube containing 100 µl of DMEM without serum. After 5 min at room temperature, the two tubes were mixed and placed at room temperature for another 20 min. The mixture was added to the 12-well plate and transfected for 5 h, and then cultured for 24 and 48 h with 500 µl fresh DMEM culture medium containing 10% FBS, and received, respectively.

Data were collected as previously described in [Liu et al. \(2021\)](#), Dual-luciferase assays were carried out in a 96-well plate (Promega, Madison, WI, USA). The activity of renilla luciferase was normalized to that of the firefly luciferase and displayed as a percentage of the control. For each treatment, three biological replications were used.

Statistical analysis

Expression of ssa-miR-301a-3p and *hsp90b2* was calculated in relative quantitation ($RQ = 2^{-\Delta\Delta CT}$). All data were analyzed using SPSS 19.0 software (IBM, Armonk, NY, USA). The relative luciferase activity value was calculated according to the formula of Renilla luciferase activity value/firefly luciferase activity value. Data of the treatment group and control group were compared using t-test, and values are expressed as mean \pm standard deviation. *P*-values less than 0.05 were considered to be statistically significant and all data were shown as mean \pm standard error (SE).

RESULTS

Potential target gene prediction of ssa-miR-301a-3p

A total of four putative ssa-miR-301a-3p target genes were obtained by biological analysis of RNA-seq data. These four genes were *hsp90b2*, *TBCC1*, *P4HAI*, and *hsp90b1*, respectively. Among them, *hsp90b2* and *hsp90b1* were involved in heat stress-related signaling pathway, while *TBCC1* and *P4HAI* were down-regulated after heat stress. The change in expression of *hsp90b2* was much more significant than that of *hsp90b1* under heat stress (Table 2). Moreover, 3'-UTR of *hsp90b2* had a complete match with the ssa-miR-301a-3p sequence and the expression level of *hsp90b2* was significantly increased in heat stress group (HL) compared with the control group (CL). Therefore, in this study, based on the negative regulation relationship between miRNA and target genes, we chose *hsp90b2* as the target gene of ssa-miR-301a-3p. The binding site of ssa-miR-301a-3p in the 3'-UTR of *hsp90b2* was in the 247–254 position (Fig. 1).




Figure 2 *ssa-miR-301a-3p* and *hsp90b2* mRNA in the liver of rainbow trout at 18 and 24 °C. Expression of *ssa-miR-301a-3p* and *hsp90b2* mRNA in the liver of rainbow trout at 18 and 24 °C. (A) The high-throughput sequencing result of *ssa-miR-301a-3p*; (B) The qRT-PCR validation result of *ssa-miR-301a-3p*; (C) The high-throughput sequencing result of *hsp90b2*; (D) The qRT-PCR validation result of *hsp90b2*. *Statistically significant difference between CL and HL $P < 0.05$. **Statistically significant difference between CL and HL $P < 0.01$. [Full-size !\[\]\(fd7fe780e8fd8eece60268c87d0c3e04_img.jpg\) DOI: 10.7717/peerj.13476/fig-2](https://doi.org/10.7717/peerj.13476/fig-2)

Expression of *ssa-miR-301a-3p* and *hsp90b2* in the liver of rainbow trout

We examined the relative expression level of *ssa-miR-301a-3p* and *hsp90b2* in the CL and HL by qRT-PCR. Results indicated that *ssa-miR-301a-3p* expression level had a significant downgrade in HL ($P < 0.05$) compared with the CL (Fig. 2B), and the relative expression of *hsp90b2* mRNA in HL was significantly up-regulated compared to the CL ($P < 0.01$) (Fig. 2D). In addition, we listed the high-throughput sequencing results of *ssa-miR-301a-3p* and *hsp90b2* and found that the sequencing results of qRT-PCR were basically consistent with those of high-throughput sequencing (Figs. 2A, 2C). The inverse relationship between the expressions of *ssa-miR-301a-3p* and *hsp90b2* mRNA in CL and HL, suggests that there might be a negative correlation between *ssa-miR-301a-3p* and *hsp90b2*.




Figure 3 Dual-luciferase reporter assay analysis of the target relationship of ssa-miR-301a-3p with *hsp90b2* in HEK 293T cells. Relative luciferase reporter expression was normalized to NC. Each treatment was repeated in triplicates. Data are depicted as mean \pm standard error. An asterisk (*) indicates $P < 0.05$ vs the mimic NC. Full-size [DOI: 10.7717/peerj.13476/fig-3](https://doi.org/10.7717/peerj.13476/fig-3)

Analysis of the ssa-miR-301a-3p binding site in the 3'-UTR of *hsp90b2*

We performed luciferase activity assays using HEK 293T cells to verify whether ssa-miR-301a-3p binds to the 3'-UTR of *hsp90b2*. There was no significant change in the expression of luciferase in the mimic NC + ssa-miR-301a-3p PC ($P > 0.05$), while the co-transfection of ssa-miR-301a-3p mimic with ssa-miR-301a-3p PC revealed that luciferase activity was inhibited ($P < 0.05$) (Fig. 3), which excluded the influence of transfection agents, RNA extracts, and detection methods in the experiment. The recombinant plasmids of *hsp90b2* WT and *hsp90b2* MUT were also co-transfected into HEK 293T cells along with either the ssa-miR-301a-3p mimic or mimic NC. There was no significant change in the expression of luciferase in the mimic NC + *hsp90b2* WT and mimic NC + *hsp90b2* MUT groups ($P > 0.05$) (Fig. 3). The inhibition of luciferase activity could be found through co-transfection of ssa-miR-301a-3p mimic with *hsp90b2* WT ($P < 0.05$), but luciferase activity was not significantly affected by co-transfection of ssa-miR-301a-3p mimic and *hsp90b2* MUT ($P > 0.05$). These results demonstrated that ssa-miR-301a-3p was involved in regulation of *hsp90b2* expression by binding with the putative binding site in the 3'-UTR of the *hsp90b2*.

DISCUSSION

In this study, we used three methods including TargetScan, PicTa, and miRanda to predict target genes for ssa-miR-301a-3p, and found that the 3'-UTR of *hsp90b2* and complementary region of ssa-miR-301a-3p was completely complementary. We observed a significant down-regulation in luciferase activity of ssa-miR-301a-3p mimics group transfection of pmirGLO-wt-*hsp90b2* 3'-UTR; consequently, the expression of *hsp90b2* mRNA was not effectively inhibited. These data indicate ssa-miR-301a-3p can complement with the 3'-UTR site of *hsp90b2*, which makes the luciferase activity decrease significantly and ssa-miR-301a-3p participates in the heat stress response of rainbow trout by targeting *hsp90b2*. In addition, as a positive control group, ssa-miR-301a-3p mimics can significantly reduce the luciferase activity of pmirGLO-ssa-miR-301a-3p in the same

experimental system, which proves that the experimental system is reasonable and the experimental results are stable and accurate. Therefore, the results showed that *hsp90b2* gene was a target gene for ssa-miR-301a-3p.

As a potential biomarker of fish's response to thermal attack, *hsp90* provides protection for cellular internal tissues and regulates important cellular functions (Sharma, Singh & Chakrabarti, 2017). As the main cell defense against acute heat stress, HSPs can be synthesized quickly (Kregel, 2002). In this study, significantly increased expression level of *hsp90b2* was found in the liver of rainbow trout after heat stress. Like our results many other investigators also reported upregulation of *hsp90* in liver of other fish species subjected to heat shock e.g., carp (Hermesz, Abraham & Nemcsok, 2001), *C. idela* (Wu et al., 2012; Sharma, Singh & Chakrabarti, 2017), *Channa striatus* (Purohit et al., 2014), *Puntius sophore* (Mahanty et al., 2017) and *Labeo Rohita*, *Catla Catla* (Ahmad et al., 2020). In fish, *hsp90* is involved in embryonic muscle development, metamorphosis, oxidative stress and anti-bacterial infection (Du et al., 2008; Manchado et al., 2008; Padmini & Rani, 2011; Cha et al., 2013). By participating in the aggregation of heat shock transcription factors to activate target genes, *hsp90* protects cells in heat shock conditions and also protects cells by inhibiting heat shock-induced apoptosis (Ali, Bharadwaj & Ovsenek, 1998; Yavelsky et al., 2004). Research of Rendell et al. (2006) showed that the *hsp90* synthesis in rainbow trout at different growth stages was significantly increased in the liver under heat stress. In addition, some studies have found that *hsp90b2* was involved in the endoplasmic reticulum-associated degradation (ERAD) of protein processing in endoplasmic reticulum pathway (Hirsch et al., 2009; Xie & Ng, 2010). In our previous study, GO and KEGG pathway enrichment analyses indicated that the target gene *hsp90b2* is regulated by miR-301a-3p, a process that involves the heat stress-related ER protein processing pathway (Li et al., 2017). The ubiquitin ligase Hrd1 recognizes *hsp90b2* (Bordallo et al., 1998; Bays et al., 2001), and translocation is mediated through the Cdc48-Ufd1-Npl4-p97AT pathway. Ubc7p and Hrd1p mediate ubiquitination. Finally, Rad23p and Dsk2p regulate the proteasome.

At optimum temperature (18 °C), miR-301a-3p inhibited the expression of *hsp90b2* by binding to the 3'-UTR of *hsp90b2*; while under heat stress (24 °C), miR-301a-3p induced the expression of the target gene *hsp90b2*. Furthermore, under heat stress, the number of aberrant and misfolded proteins in rainbow trout liver increases significantly. The buildup of these misfolded proteins accumulates significantly, affecting the normal function of the ER, inducing ER stress, and finally leading to cell death (Huang et al., 2018). The elimination of misfolded peptides/proteins is so critical. Under heat stress conditions, we detected a considerable downregulation of miR-301a-3p expression; as a result, *hsp90b2* expression was not successfully suppressed. These findings suggest that miR-301a-3p plays a role in rainbow trout heat stress response by targeting *hsp90b2*, which is enriched in the ERAD pathway, to increase the degradation of misfolded proteins, hence regulating the adaptive protection mechanism.

Fish growth is influenced by various aspects such as temperature variation, hypoxia, salinity and heavy metal (Myers, 1998; Topal et al., 2017). There is abundant evidence that miRNA can regulate gene regulation through direct or indirect means in the face of various

kinds of stress (Bartel, 2009). Studies have shown that miRNAs have the ability to regulate the expression levels of *Hsp60* and *Hsp70* in skeletal muscle and cardiac muscle cells (Shan et al., 2010; Kukreti et al., 2013). In the present study, we found a targeted relationship between ssa-miR-301a-3p and *hsp90b2*, and when the expression of ssa-miR-301a-3p was down-regulated, the expression of *hsp90b2* was significantly up-regulated, indicating there is a negative regulatory relationship between ssa-miR-301a-3p and *hsp90b2*. When the expression of *hsp90b2* is up-regulated, the misfolding of protein may be corrected, and the immune and endocrine systems may mediate under stress conditions and resistance of rainbow trout to heat stress is enhanced.

CONCLUSION

The expression level of ssa-miR-301a-3p was down-regulated and *hsp90b2* was exactly the opposite of ssa-miR-301a-3p under heat stress. There might be negative regulatory relations between ssa-miR-301a-3p and *hsp90b2* through binding with the 3'-UTR region of *hsp90b2*. Our findings provide evidence for the role of ssa-miR-301a-3p in the liver and will help understand the mechanism of miRNA-mediated gene regulation in response to heat stress in rainbow trout.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by grants from the National Natural Science Foundation of China (Grant No. 31660735). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
National Natural Science Foundation of China: 31660735.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Zhe Liu conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Fang Ma performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Yujun Kang analyzed the data, prepared figures and/or tables, and approved the final draft.
- Xiaoxia Liu performed the experiments, prepared figures and/or tables, and approved the final draft.

Animal Ethics

The following information was supplied relating to ethical approvals (*i.e.*, approving body and any reference numbers):

Care and Use of Laboratory Animals in China and was approved by the Institutional Ethic Committee of Gansu Agricultural University.

Data Availability

The following information was supplied regarding data availability:

The data is available at NCBI SRA: [SRP109730](https://www.ncbi.nlm.nih.gov/sra/SRP109730).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.13476#supplemental-information>.

REFERENCES

- Ahmad M, Zuberi A, Ali M, Syed A, Murtaza MUH, Khan A, Kamran M. 2020. Effect of acclimated temperature on thermal tolerance, immune response and expression of HSP genes in *Labeo rohita*, *Catla catla* and their intergeneric hybrids. *Journal of Thermal Biology* **89**:102570 DOI [10.1016/j.jtherbio.2020.102570](https://doi.org/10.1016/j.jtherbio.2020.102570).
- Ali A, Bharadwaj SR, Ovsenek N. 1998. HSP90 interacts with and regulates the activity of heat shock factor 1 in *Xenopus* oocytes. *Molecular and Cellular Biology* **18**(9):4949–4960 DOI [10.1128/mcb.18.9.4949](https://doi.org/10.1128/mcb.18.9.4949).
- Bartel DP. 2004. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* **116**(2):281–297 DOI [10.1016/S0092-8674\(04\)00045-5](https://doi.org/10.1016/S0092-8674(04)00045-5).
- Bartel DP. 2009. MicroRNAs: target recognition and regulatory functions. *Cell* **136**(2):215–233 DOI [10.1016/j.cell.2009.01.002](https://doi.org/10.1016/j.cell.2009.01.002).
- Basu N, Todgham AE, Ackerman PA, Bibeau MR, Nakano K, Schulte PM, Iwama GK. 2002. Heat shock protein genes and their functional significance in fish. *Gene* **295**:173–183 DOI [10.1016/s0378-1119\(02\)00687-x](https://doi.org/10.1016/s0378-1119(02)00687-x).
- Baulcombe D. 2004. RNA silencing in plants. *Nature* **431**(7006):356–363 DOI [10.1038/nature02874](https://doi.org/10.1038/nature02874).
- Bays NW, Gardner RG, Seelig LP, Joazeiro CA, Hampton RY. 2001. Jrd1p/Der3p is a membrane-anchored ubiquitin ligase required for ER-associated degradation. *Nature Cell Biology* **3**(1):24–29 DOI [10.1038/35050524](https://doi.org/10.1038/35050524).
- Bordallo J, Plemper RK, Finger A, Wolf DH. 1998. Der3p/Hrd1p is required for endoplasmic reticulum-associated degradation of misfolded luminal and integral membrane proteins. *Molecular Biology of the Cell* **9**(1):209–222 DOI [10.1091/mbc.9.1.209](https://doi.org/10.1091/mbc.9.1.209).
- Bushati N, Cohen SM. 2007. microRNA functions. *Annual Review of Cell and Developmental Biology* **23**(1):175–205 DOI [10.1146/annurev.cellbio.23.090506.123406](https://doi.org/10.1146/annurev.cellbio.23.090506.123406).
- Cara JB, Aluru N, Moyano FJ, Vijayan MM. 2005. Food-deprivation induces HSP70 and HSP90 protein expression in larval gilthead sea bream and rainbow trout. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* **142**(4):426–431 DOI [10.1016/j.cbpb.2005.09.005](https://doi.org/10.1016/j.cbpb.2005.09.005).
- Carrington JC, Ambros V. 2003. Role of microRNAs in plant and animal development. *Science* **301**(5631):336–338 DOI [10.1126/science.1085242](https://doi.org/10.1126/science.1085242).

- Cha IS, Kwon J, Park SB, Jang HB, Nho SW, Kim YK, Hikima J, Aoki T, Jung TS. 2013.** Heat shock protein profiles on the protein and gene expression levels in olive flounder kidney infected with *Streptococcus parauberis*. *Fish & Shellfish Immunology* **34**(6):1455–1462 DOI [10.1016/j.fsi.2013.03.355](https://doi.org/10.1016/j.fsi.2013.03.355).
- Chen PY, Meister G. 2005.** microRNA-guided posttranscriptional gene regulation. *Biological Chemistry* **386**(12):1205–1218 DOI [10.1515/BC.2005.139](https://doi.org/10.1515/BC.2005.139).
- Du SJ, Li HQ, Bian YH, Zhong YW. 2008.** Heat-shock protein 90 α 1 is required for organized myofibril assembly in skeletal muscles of zebrafish embryos. *Proceedings of the National Academy of Sciences of the United States of America* **105**(2):554–559 DOI [10.1073/pnas.0707330105](https://doi.org/10.1073/pnas.0707330105).
- Haase M, Fitze G. 2016.** HSP90AB1: helping the good and the bad. *Gene* **575**(2):171–186 DOI [10.1016/j.gene.2015.08.063](https://doi.org/10.1016/j.gene.2015.08.063).
- Hermesz E, Abraham M, Nemcsok J. 2001.** Identification of two *hsp90* genes in carp. *Comparative Biochemistry Physiology Part C: Toxicology and Pharmacology* **129**:397–407 DOI [10.1016/S1532-0456\(01\)00216-2](https://doi.org/10.1016/S1532-0456(01)00216-2).
- Hirsch C, Gauss R, Horn SC, Neuber O, Sommer T. 2009.** The ubiquitylation machinery of the endoplasmic reticulum. *Nature* **458**(7237):453–460 DOI [10.1038/nature07962](https://doi.org/10.1038/nature07962).
- Hong DS, Banerji U, Tavana B, George GC, Aaron J, Kurzrock R. 2013.** Targeting the molecular chaperone heat shock protein 90 (HSP90): lessons learned and future directions. *Cancer Treatment Reviews* **39**(4):375–387 DOI [10.1016/j.ctrv.2012.10.001](https://doi.org/10.1016/j.ctrv.2012.10.001).
- Huang JQ, Li YJ, Ma F, Kang YJ, Liu XX, Liu Z, Wang JF. 2018.** Identification and characterization of microRNAs in the liver of rainbow trout in response to heat stress by high-throughput sequencing. *Gene* **679**(2):274–281 DOI [10.1016/j.gene.2018.09.012](https://doi.org/10.1016/j.gene.2018.09.012).
- Iba K. 2002.** Acclimative response to temperature stress in higher plants: approaches of gene engineering for temperature tolerance. *Annual Review of Plant Biology* **53**(1):225–245 DOI [10.1146/annurev.arplant.53.100201.160729](https://doi.org/10.1146/annurev.arplant.53.100201.160729).
- Ikert H, Craig PM. 2020.** Chronic exposure to venlafaxine and increased water temperature reversibly alters microRNA in zebrafish gonads (*Danio rerio*). *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics* **33**:100634 DOI [10.1016/j.cbd.2019.100634](https://doi.org/10.1016/j.cbd.2019.100634).
- Jackson SE. 2013.** Hsp90: structure and function. *Topics in Current Chemistry* **328**(12):155–240 DOI [10.1007/128_2012_356](https://doi.org/10.1007/128_2012_356).
- Jia Y, Cavileer TD, Nagler JJ. 2016.** Acute hyperthermic responses of heat shock protein and estrogen receptor mRNAs in rainbow trout hepatocytes. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **201**(2):156–161 DOI [10.1016/j.cbpa.2016.04.023](https://doi.org/10.1016/j.cbpa.2016.04.023).
- Jiang H, Lv J. 2021.** MicroRNA-301a-3p increases oxidative stress, inflammation and apoptosis in ox-LDL-induced HUVECs by targeting KLF7. *Experimental and Therapeutic Medicine* **21**(6):569 DOI [10.3892/etm.2021.10001](https://doi.org/10.3892/etm.2021.10001).
- Kregel KC. 2002.** Heat shock proteins: modifying factors in physiological stress responses and acquired thermotolerance. *Journal of Applied Physiology* **92**(5):2177–2186 DOI [10.1152/jappphysiol.01267.2001](https://doi.org/10.1152/jappphysiol.01267.2001).
- Kukreti H, Amuthavalli K, Harikumar A, Sathiyamoorthy S, Feng PZ, Anantharaj R, Tan SL, Lokireddy S, Bonala S, Sriram S, McFarlane C, Kambadur R, Sharma M. 2013.** Muscle-specific microRNA1 (miR1) targets heat shock protein 70 (HSP70) during dexamethasone-mediated atrophy. *Journal of Biological Chemistry* **288**(9):6663–6678 DOI [10.1074/jbc.M112.390369](https://doi.org/10.1074/jbc.M112.390369).

- Lee RC, Ambros V. 2001. An extensive class of small RNAs in *Caenorhabditis elegans*. *Science* 294(5543):862–864 DOI 10.1126/science.1065329.
- Li YJ, Huang JQ, Liu Z, Zhou YJ, Xia BP, Wang YJ, Kang YJ, Wang JF. 2017. Transcriptome analysis provides insights into hepatic responses to moderate heat stress in the rainbow trout (*Oncorhynchus mykiss*). *Gene* 619:1–9 DOI 10.1016/j.gene.2017.03.041.
- Lindquist S. 1986. The heat-shock response. *Annual Review of Biochemistry* 55(1):1151–1191 DOI 10.1146/annurev.bi.55.070186.005443.
- Liu X, Wang Y, Liu Z, Kang Y, Ma F, Luo Z, Wang J, Huang J. 2021. miR-434 and miR-242 have a potential role in heat stress response in rainbow trout (*Oncorhynchus mykiss*). *Journal of Fish Biology* 99(6):1798–1803 DOI 10.1111/jfb.14881.
- Mahanty A, Purohit GK, Yadav RP, Mohanty S, Mohanty BP. 2017. hsp90 and hsp47 appear to play an important role in minnow *Puntius sophore* for surviving in the hot spring run-off aquatic ecosystem. *Fish Physiology and Biochemistry* 43(1):89–102 DOI 10.1007/s10695-016-0270-y.
- Manchado M, Salas-Leiton E, Infante C, Ponce M, Asensi E, Crespo A, Zuasti E, Canavate JP. 2008. Molecular characterization, gene expression and transcriptional regulation of cytosolic HSP90 genes in the flatfish Senegalese sole (*Solea senegalensis* Kaup). *Gene* 416(1–2):77–84 DOI 10.1016/j.gene.2008.03.007.
- Matthews KR, Berg NH. 1997. Rainbow trout responses to water temperature and dissolved oxygen stress in two southern California stream pools. *Journal of Fish Biology* 50(1):50–67 DOI 10.1111/j.1095-8649.1997.tb01339.x.
- Matzke M, Aufsatz W, Kanno T, Daxinger L, Papp I, Mette MF, Matzke AJ. 2004. Genetic analysis of RNA-mediated transcriptional gene silencing. *Biochimica et Biophysica Acta* 1677(1–3):129–141 DOI 10.1016/j.bbaexp.2003.10.015.
- Morey TM, Roufayel R, Johnston DS, Fletcher AS, Mosser DD. 2015. Heat shock inhibition of CDK5 increases NOXA levels through miR-23a repression. *Journal of Biological Chemistry* 290(18):11443–11454 DOI 10.1074/jbc.M114.625988.
- Myers RA. 1998. When do environment-recruitment correlations work? *Reviews in Fish Biology and Fisheries* 8(3):285–305 DOI 10.1023/A:1008828730759.
- Padmini E, Rani MU. 2011. Heat-shock protein 90 alpha (HSP90 α) modulates signaling pathways towards tolerance of oxidative stress and enhanced survival of hepatocytes of *Mugil cephalus*. *Cell Stress and Chaperones* 16(4):411–425 DOI 10.1007/s12192-011-0255-9.
- Purohit GK, Mahanty A, Suar M, Sharma AP, Mohanty BP, Mohanty S. 2014. Investigating hsp gene expression in liver of *Channa striatus* under heat stress for understanding the upper thermal acclimation. *BioMed Research International* 2014:1–10 DOI 10.1155/2014/381719.
- Quinteiro-Filho WM, Ribeiro A, FerrazdePaula V, Pinheiro ML, Sakai M, Sá LRM, Ferreira AJP, Palermo-Neto J. 2010. Heat stress impairs performance parameters, induces intestinal injury, and decreases macrophage activity in broiler chickens. *Poultry Science* 89(9):1905–1914 DOI 10.3382/ps.2010-00812.
- Rendell JL, Fowler S, Cockshutt A, Currie S. 2006. Development-dependent differences in intracellular localization of stress proteins (hsps) in rainbow trout, *Oncorhynchus mykiss*, following heat shock. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics* 1(2):238–252 DOI 10.1016/j.cbd.2005.12.004.
- Shan ZX, Lin QX, Deng CY, Zhu JN, Mai LP, Liu JL, Fu YH, Liu XY, Li YX, Zhang YY, Lin SG, Yu XY. 2010. miR-1/miR-206 regulate Hsp60 expression contributing to glucose-mediated apoptosis in cardiomyocytes. *FEBS Letters* 584(16):3592–3600 DOI 10.1016/j.febslet.2010.07.027.

- Sharma J, Singh SP, Chakrabarti R. 2017. Effect of temperature on digestive physiology, immune-modulatory parameters, and expression level of Hsp and LDH genes in *Catla catla* (Hamilton, 1822). *Aquaculture* 479:134–141 DOI 10.1016/j.aquaculture.2017.05.031.
- St-Pierre NR, Cobanov B, Schnitkey G. 2003. Economic losses from heat stress by US livestock industries. *Journal of Dairy Science* 86:E52–E77 DOI 10.3168/jds.S0022-0302(03)74040-5.
- Sun J, Zhao L, Wu H, Lian W, Cui C, Du Z, Luo W, Li M, Yang S. 2019. Analysis of miRNA-seq in the liver of common carp (*Cyprinus carpio* L.) in response to different environmental temperatures. *Functional & Integrative Genomics* 19(2):265–280 DOI 10.1007/s10142-018-0643-7.
- Topal A, Atamanalp M, Qruc E, Erol HS. 2017. Physiological and biochemical effects of nickel on rainbow trout (*Oncorhynchus mykiss*) tissues: assessment of nuclear factor kappa B activation, oxidative stress and histopathological changes. *Chemosphere* 166(5):445–452 DOI 10.1016/j.chemosphere.2016.09.106.
- Wu CX, Zhao FY, Zhang Y, Zhu YJ, Ma MS, Mao HL, Hu CY. 2012. Overexpression of Hsp90 from grass carp (*Ctenopharyngodon idella*) increases thermal protection against heat stress. *Fish & Shellfish Immunology* 33(1):42–47 DOI 10.1016/j.fsi.2012.03.033.
- Xia BP, Liu Z, Zhou YJ, Wang YJ, Huang JQ, Li YJ, Kang YJ, Wang JF, Liu XX. 2018. Effects of heat stress on biochemical parameters and heat shock protein family A (*Hsp70*) member 5 (*HSPA5*) mRNA expression in rainbow trout (*Oncorhynchus mykiss*). *Marine & Freshwater Research* 69(1):1674–1680 DOI 10.1016/j.fsi.2012.03.033.
- Xia X, Wang S, Ni B, Xing S, Cao H, Zhang Z, Yu F, Zhao E, Zhao G. 2020. Hypoxic gastric cancer-derived exosomes promote progression and metastasis via MiR-301a-3p/PHD3/HIF-1 α positive feedback loop. *Oncogene* 39(39):6231–6244 DOI 10.1038/s41388-020-01425-6.
- Xia X, Zhang K, Luo G, Cen G, Cao J, Huang K, Qiu Z. 2017. Downregulation of miR-301a-3p sensitizes pancreatic cancer cells to gemcitabine treatment via PTEN. *American Journal of Translational Research* 9:1886–1895.
- Xie W, Ng DT. 2010. ERAD substrate recognition in budding yeast. *Seminars in Cell & Developmental Biology* 21(5):533–539 DOI 10.1016/j.semcdb.2010.02.007.
- Yavelsky V, Vais O, Piura B, Wolfson M, Rabinovich A, Fraifeld V. 2004. The role of Hsp90 in cell response to hyperthermia. *Journal of Thermal Biology* 29(7–8):509–514 DOI 10.1016/j.jtherbio.2004.08.078.
- Yu J, Liu F, Yin P, Zhu X, Cheng G, Wang N, Lu A, Luan W, Zhang N, Li J, Guo K, Yin Y, Wang H, Xu J. 2010. Integrating miRNA and mRNA expression profiles in response to heat stress-induced injury in rat small intestine. *Functional & Integrative Genomics* 11(2):203–213 DOI 10.1007/s10142-010-0198-8.
- Zhang H, Chen T, Xiong J, Hu B, Luo J, Xi Q, Jiang Q, Sun J, Zhang Y. 2020. MiR-130a-3p inhibits PRL expression and is associated with heat stress-induced PRL reduction. *Frontiers in Endocrinology* 11:92 DOI 10.3389/fendo.2020.00092.
- Zhang C, Tong C, Tian F, Zhao K. 2017. Integrated mRNA and microRNA transcriptome analyses reveal regulation of thermal acclimation in *Gymnocypris przewalskii*: a case study in Tibetan Schizothoracine fish. *PLOS ONE* 12(10):e0186433 DOI 10.1371/journal.pone.0186433.
- Zhang R, Zhang Q, Niu JY, Lu K, Xie B, Cui DS, Xu SJ. 2014. Screening of microRNAs associated with Alzheimer's disease using oxidative stress cell model and different strains of senescence accelerated mice. *Journal of the Neurological Sciences* 338(1–2):57–64 DOI 10.1016/j.jns.2013.12.017.

Zheng Y, Chen KL, Zheng XM, Li HX, Wang GL. 2014. Identification and bioinformatics analysis of microRNAs associated with stress and immune response in serum of heat-stressed and normal Holstein cows. *Cell Stress and Chaperones* **19**(6):973–981
DOI [10.1007/s12192-014-0521-8](https://doi.org/10.1007/s12192-014-0521-8).

Zhou P, Jiang W, Wu L, Chang R, Wu K, Wang Z. 2012. miR-301a is a candidate oncogene that targets the homeobox gene Gax in human hepatocellular carcinoma. *Digestive Diseases and Sciences* **57**(5):1171–1180 DOI [10.1007/s10620-012-2099-2](https://doi.org/10.1007/s10620-012-2099-2).