

Evaluating a linkage between obesity and the occurrence of dental caries among school going children in Sakaka, Al Jouf, Kingdom of Saudi Arabia (#71641)

1

First submission

Guidance from your Editor

Please submit by **21 Apr 2022** for the benefit of the authors (and your \$200 publishing discount) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

4 Table file(s)

1 Raw data file(s)



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

- 1. Your most important issue*
- 2. The next most important item*
- 3. ...*
- 4. The least important points*

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Evaluating a linkage between obesity and the occurrence of dental caries among school going children in Sakaka, Al Jouf, Kingdom of Saudi Arabia

Osama Khattak^{Corresp., 1}, Azhar Iqbal¹, Farooq Ahmad Chaudhary^{Corresp., 2}, Jamaluddin Syed^{3, 4}, Thani AlSharari⁵, Sudhakar Vundavalli⁶, Bayan Abdullah Sadiq Aljahdali⁷, Ahmed Eidan Abdullah AlZahrani⁸, Rakhi Issrani⁶, Sherif Elsayad Sultan⁹

¹ Department of Restorative Dentistry, Jouf University, Sakaka, Saudi Arabia

² School of Dentistry (SOD), Federal Medical Teaching Institution (FMTI)/PIMS, Shaheed Zulfiqar Ali Bhutto Medical University (SZABMU), Islamabad, Pakistan

³ Oral Basic Clinical Sciences Department, faculty of Dentistry, King AbdulAziz university, Jeddah, Saudi Arabia

⁴ Department of Regenerative Dentistry, University of Bari, Bari, Italy

⁵ Restorative and Dental Materials Department, Faculty of Dentistry, Taif University, Taif, Saudi Arabia

⁶ Department of Preventive Dentistry, College of Dentistry, Jouf University, Sakaka, Saudi Arabia

⁷ General dentist, Arar, Saudi Arabia

⁸ General dentist, Taif, Saudi Arabia

⁹ Department of fixed Prosthodontics, Tanta University, Tanta, Egypt

Corresponding Authors: Osama Khattak, Farooq Ahmad Chaudhary
Email address: dr.osama.khattak@jodent.org, chaudhary4@hotmail.com

Background: Obesity and dental caries are global public health problems. There are conflicting reports about the relationship between caries and obesity. The aim of this study was to analyse the type of relationship between the dental caries and obesity among school children in Al-Jouf region of Saudi Arabia. **Methods:** This cross-sectional study was conducted among 400 participants aged 6 to 14 years. The study involved measuring caries (dmft/DMFT), assessing body mass index (BMI), and administering a self-completion questionnaire. The Independent *t*-test, one-way ANOVA, and multivariate logistic regression analyses were performed. **Results:** A total of 380 children agreed to participate in the study. Comparison between the dental caries (dmft/DMFT) and oral hygiene habits like frequency of brushing, usage of dental floss and mouthwash/ rinsing habits was found to be insignificant ($P=0.23$). Mean DMFT values were directly proportional to frequency of intake of sucrose diet in permanent dentition. Statistically significant association ($P<0.001$) was found between overweight (mean DMFT=2.8) and obesity (mean DMFT=3.9) with caries scores in permanent dentition only. Odds ratio for BMI to caries score in permanent dentition was found to be 3.2 which was statistically significant whereas other variables were found to be not significant. **Conclusion:** There was a positive association between obesity and dental caries in school going Saudi Arabian children. Comprehensive

multidisciplinary approach by health professionals is recommended for preventive public health issues related to caries and obesity in teenagers.

Evaluating a linkage between obesity and the occurrence of dental caries among school going children in Sakaka , AlJouf, Kingdom of Saudi Arabia.

Osama Khattak^{1*}, Azhar Iqbal¹, Farooq Ahmad Chaudhary², Jamaluddin Syed^{3,4}, Thani AlSharari⁵, Sudhakar Vundavalli⁶, Bayan Abdullah Sadiq Aljahdali⁷, Ahmed Eidan Abdullah AlZahrani⁸, Rakhi Issrani⁶, Sherif Elsayad Sultan⁹

¹ Department of Restorative Dentistry, Jouf University, Sakaka, Saudi Arabia.
² School of Dentistry (SOD), Federal Medical Teaching Institution (FMTI)/PIMS, Shaheed Zulfiqar Ali Bhutto Medical University (SZABMU), Islamabad, Pakistan.
³ Oral Basic Clinical Sciences Department, faculty of Dentistry, King AbdulAziz university, Jeddah, Saudi Arabia.
⁴ Department of Regenerative Dentistry, University of Bari, Italy.
⁵ Restorative and Dental Materials Department , Faculty of Dentistry , Taif University, Saudi Arabia
⁶ Department of Preventive Dentistry, College of Dentistry, Jouf University; Saudi Arabia.
⁷ General dentist, Arar, Saudi Arabia; Bayan.alj90@gmail.com
⁸ General dentist, Taif, Saudi Arabia; Dr.ahmedana@hotmail.com

Corresponding Author:
 Osama Khattak^{1*}
 Department of Restorative Dentistry, Jouf University, Sakaka, Saudi Arabia.
 Email address: dr.osama.khattak@jodent.org

Abstract

Background: Obesity and dental caries are global public health problems. There are conflicting reports about the relationship between caries and obesity. The aim of this study was to analyse the type of relationship between the dental caries and obesity among school children in Al-Jouf region of Saudi Arabia.

Methods: This cross-sectional study was conducted among 400 participants aged 6 to 14 years. The study involved measuring caries (dmft/DMFT), assessing body mass index (BMI), and administering a self-completion questionnaire. The Independent *t*-test, one-way ANOVA, and multivariate logistic regression analyses were performed.

Results: A total of 380 children agreed to participate in the study. Comparison between the dental caries (dmft/DMFT) and oral hygiene habits like frequency of brushing, usage of dental floss and mouthwash/ rinsing habits was found to be insignificant ($P=0.23$). Mean DMFT values were directly proportional to frequency of intake of sucrose diet in permanent dentition. Statistically significant association ($P< 0.001$) was found between overweight (mean DMFT=2.8) and obesity (mean DMFT=3.9) with caries scores in permanent dentition only. Odds ratio for BMI to caries score in permanent dentition was found to be 3.2 which was statistically significant whereas other variables were found to be not significant.

Conclusion: There was a positive association between obesity and dental caries in school going Saudi Arabian children. Comprehensive multidisciplinary approach by health professionals is recommended for preventive public health issues related to caries and obesity in teenagers.

48 **Introduction**

49 Numerous changes have been observed over the last few decades in the diet and lifestyle of the
 50 general population due to rapid urbanization, industrialization, and economic
 51 development(Swinburn et al. 2011). These changes include consumption of high amounts of
 52 carbohydrates and lower physical activity, which negatively impact their general and dental health,
 53 especially in young age groups, causing dental caries and obesity(Fernandez et al. 2017). These
 54 two conditions are considered chronic, widespread, and multifactorial, having a devastating impact
 55 on the lives of children and youth (Fernandez et al. 2017).

56 Dental caries have become a global public health issue for pre-school children and lead to pain,
 57 chewing difficulties, and disorders of general health affecting the growth and development of
 58 children(Fernandez et al. 2017; Hayden et al. 2013). Studies from various regions of the world
 59 reported the prevalence of dental caries in children ranges from 40-90%, and in Saudi Arabia its
 60 prevalence is reported to be around 80% in primary and 70% in permanent dentition (Al Agili
 61 2013; Fernandez et al. 2017; Tsai et al. 2000).

62 Similar to dental caries prevalence of obesity is also very high and it is becoming one of the most
 63 common general health problems among school children(Akpata et al. 1992). A cross-sectional
 64 study in America reported the prevalence of obesity among children of age 4 at 18%, in Taiwan
 65 among 3-10 years children it was in the range of 4% to 17%, and in Saudia Arabia obesity has
 66 become an emerging public health problem among the children due to its alarming high rate (Al
 67 Othaimen et al. 2007; Anderson & Whitaker 2009; Tsai et al. 2000). If obesity remains
 68 uncontrolled, it can lead to devastating health conditions such as high blood pressure, diabetes,
 69 stroke, and cardiovascular disease. All these conditions ultimately can lead to disabilities and
 70 death.(Sturm 2002) Studies have shown that obesity and dental caries have become common

childhood diseases that are affecting the overall health and development of children(Wake et al. 2007). Many contributing factors are common for both these conditions, these include lifestyle, biological, dietary, genetic, cultural, socioeconomic, and environmental factors (Chi et al. 2017; Spiegel & Palmer 2012). Therefore an association between dental caries and obesity seems logical and knowledge about this relationship is very important so that more effective and targeted public measures could be initiated to reduce its prevalence (Chi et al. 2017). It is also believed that understanding more about this relationship can help to control both of these conditions at once (Scorzetti et al. 2013). Previous studies that examined this relationship provided conflicting findings and the relationship between obesity and dental caries is still unclear (Fernandez et al. 2017; Scorzetti et al. 2013).

No such study has been conducted in the Al-Jouf region of Saudi Arabia; therefore, no data is available about the association between dental caries and obesity in the school children in this region. Hence, the present study was designed to examine and report the type of relationship between dental caries and obesity among school children in the Al-Jouf region of Saudi Arabia.

Materials & Methods

This cross-sectional study was carried out between March to May 2021 among the children of Sakaka, Al-Jouf province, Saudi Arabia. The children aged between 6-14 years old and permanent residents of the Sakaka region were selected for this study using stratified cluster random sampling method. Initially schools were selected through simple random (lottery) method from list of the schools in Sakaka. From the selected schools children were selected using systematic random sampling method. Ethical approval was obtained from the ethical local committee of bioethics,

Jouf University (Ref: 07-06-43). The children who were medically compromised and had any psychological disorders were excluded from this study.

The sample size calculation was based on the prevalence of dental caries reported by Gudipani RK et al. (2021)(Gudipani et al. 2021). For this investigation, separate sample sizes were determined for distinct caries prevalence rates for different BMI groups, with the largest calculated sample size among all being used. Considering 95% confidence levels, and 80% power, 350 children were sufficient to detect the clinically significant difference of 10% in the prevalence of caries. The total sample size was adjusted to 400, expecting the fact that there were approximately 10-15% chances of non-participation/rejection.

Participation was voluntary, both parents and children were informed about the purpose of the study before taking oral and written consent from children and either of the parents. It was assured that the results of the study would be only presented or published as aggregate data maintaining the confidentiality of personal information.

All the selected children underwent intraoral examination by two trained and calibrated dentists. Dental caries was recorded using **dmft/DMFT** (decayed, missing, filled teeth for deciduous and permanent dentition) based on WHO methods for oral health surveys(Organization 2013). For the calibration process the inter-and intra-examiner reliability was analysed using Cohen's kappa coefficient that was ranging from 0.78 to 0.82 indicates fair to a good agreement.

The body mass index (BMI) was used to evaluate obesity among children by calculating their weight in kg divided by the square of height in meters (kg/m²). The weight of the children was recorded on a physician's scale and a stadiometer was used to measure the height. The BMI scores of children were categorized according to the cut-off values described by International Obesity Task Force, as follows: BMI group- Moderate malnutrition (BMI 16-16.99), Mild malnutrition

(BMI 17-18.49), Normal (BMI 18.5–24.99), Overweight (BMI 25–29), Obese (BMI 30≥)..(James 2004)

The self-administered questionnaire collected information regarding demographic characteristics (age), dietary (sugar consumption), and oral hygiene habits (frequency of tooth brushing, use of dental floss, and mouthwash/ rinsing habits). Quantification was done on tooth brushing frequency and the amount of sugar consumed/day. Sugar consumption was determined from 24-hour dietary recall chart filled by parents on two different days with one week gap and average of both charts was taken as sugar consumption value for the participant. Face-to-face interviews of 20 parents were conducted to assess the reliability of the questionnaire and Cronbach's alpha of 0.75 was obtained in this study.

Statistical analysis

The analysis of collected data was carried out in IBM SPSS software version 24.0 with the significance level set at 5%. The comparison of mean dental caries scores with gender was examined using an independent sample t-test and the relationship of mean caries scores with BMI scores was analysed using Pearson's correlation coefficient. Multiple regression was performed for caries scores as a dependent variable and various independent variables which were significant in bivariate analysis.

Results

Out of 400 children contacted, 380 children were consented to take part in this study with a response rate of 95%. The mean age of the sample was 9.8 years, and it was consisted of 211 (55.5%) males and 169 (44.5%) female participants. [Table 1]

Mean dmft/DMFT values [dmft=4.2 and DMFT=3.2] were highest among the children who didn't brush their teeth regularly compared to the children who brushed regularly which was non-significant statistically. A minor difference was reported in mean dmft/DMFT values between the children who floss regular and who didn't floss i.e., dmft of 3.8 vs 3.6 and DMFT of 2.4 vs 2.2. Children who used mouthwash/who have mouth rinsing habits had fewer caries in permanent dentition [2 vs 2.8] and deciduous dentition [3.9 vs 3.2]. Mean DMFT values were directly proportional to the frequency of intake of sucrose diet in permanent dentition, mean DMFT of 2 in children who consumed sucrose diet once a day to 2.6 in people who consumed sucrose diet more than 6 times a day. Table 2 presents the distribution of mean caries scores (dmft/DMFT) according to oral hygiene habits, and sugar consumption.

BMI values of the children were compared to mean caries scores as shown in Table 3. A statistically significant association was found between overweight (mean DMFT=2.8) and obesity (mean DMFT=3.9) with caries scores in permanent dentition and also for deciduous dentition but was statistically insignificant.

Multiple regression was performed for the statistically significant independent variables observed in bivariate analysis. Gender, BMI values, and sugar intake were included in the model. The odds ratio for BMI to caries score in permanent dentition was found to be 3.2 which was statistically significant whereas other variables were not significant. [Table 4]

Discussion

A plausible indirect biological association of dental caries and obesity has been argued in literature

considering the fact that poor dietary habits and inappropriate diet not only promotes obesity but also create favourable condition for dental caries to flourish, especially with increased intake of Sugary food (Gerdin et al. 2008). However further research and investigation are needed to establish this relationship due to its controversial status. Hence, this study was designed to analyses this type of relationship between caries and obesity in Saudi Arabian 6 to 14 years old school-going children.

The mean dmft/DMFT in this study was higher in those children who brushed less than once a day as compared to those who brushed once and more per day. This finding was statistically insignificant. This result was in line with the results of the previous studies where a greater risk of dental caries was related to irregular tooth brushing habits (Ashour et al. 2018; Chaudhary & Ahmad 2021; Chaudhary et al. 2019; Farooqi et al. 2015). However, other oral hygiene habits like usage of dental floss and mouthwash/ rinsing habits were found to be not significant among the study participants.

On the other hand, a comparison between caries (dmft/DMFT) and dietary habits among the study participants showed that mean DMFT values were directly proportional to the frequency of intake of sucrose diet in permanent dentition. A similar finding was noted in the study done by Ashour NA et al. (2018) in which the subjects with sugar consumption were more likely to have caries than subjects with no sugar consumption(Ashour et al. 2018). In the last 30 years, the Saudi Arabian population has witnessed marked changes in their lifestyle and dietary habits resulting in deteriorating health behaviours with a more sedentary lifestyle, reduce physical activity, and fat-rich diet (Alshihri et al. 2019).

Results of this study showed that the obese children were 3.2 times more likely to have caries than the non-obese children in their permanent dentition. The association of weight and caries in the

literature still remains controversial, however, the results of this study indicated a strong relationship between overweight status and dental caries. A systematic review by Hayden C et al. (2013) had shown a similar association of obesity and increased level of dental caries in permanent dentition (Hayden et al. 2013). A previous study in the Al-Khobar region of Saudi Arabia also showed a similar association between obesity and caries experience (Al-Ansari & Nazir 2020).

In this study, dental caries in the primary dentition showed no correlation with the BMI scores, and similar results were found in another study conducted by Hayden C et al. (2013), suggesting that older children will be more likely to be obese due to increasingly sedentary lifestyle with age (Hayden et al. 2013).

Few studies have reported an inverse association between caries and BMI (Cheng et al. 2019; Fernandez et al. 2017; Kopycka-Kedzierawski et al. 2008). However, the mechanism of this inverse relationship is difficult to explain as it is evident from the literature that consumption of sugary food and refined carbohydrates contributes to the development of dental caries and obesity (Fernandez et al. 2017). A plausible argument that can support this inverse association can be attributable to the dietary patterns of obese. People with obesity do not necessarily consume more sugary and refined carbs diet rather they consume more fatty, fried, and unrefined carbohydrates foods. This dietary pattern could certainly increase obesity, but not necessarily effects dental caries (Alshihri et al. 2019).

The other perception related to the diet is that people who frequently consume snacks have a greater risk of obesity and caries, but many studies that examine this relationship of snacking habits with the weight status found either no relationship or reported opposite relationship where young people who consumed snacks in between the meals were less likely to be obese (Fernandez et al. 2017). Apart from the diet itself the process of mastication is greatly affected by caries which can

lead to feeding difficulties and nutritional intake deficiency mostly in children and older people (Alshihri et al. 2019). Study by Gilchrist F et al. (2015) showed that children with untreated caries had difficulties in chewing hard food along with food getting stuck in teeth which restricted their practice of eating a healthy diet (Gilchrist et al. 2015). The consequences of these dietary limitations are not just limited to weight only but would reduce consumption of important dietary nutrients such as calcium, phosphate, and vitamins A and D, which are essential for developing and maintaining healthy teeth and gums (Alshihri et al. 2019). Another aspect of an inverse association of snacking with dental caries is the increased secretion of saliva due to more food consumption by obese people that could eventually reduce the incidence of caries owing to its protective effect. Additionally, studies have mentioned other factors such as poverty and low socioeconomic status that explain the relationship of being underweight and having more dental caries (Chaudhary & Ahmad 2021; Chaudhary et al. 2019; Farsi et al. 2016; Kumar et al. 2017). Few studies have reported that BMI did not correlate with caries, such as studies in Saudi Arabia and Brazil on school-going children showed no association between BMI and dental caries in both primary and permanent dentition (Alves et al. 2013; da Silva et al. 2016). In a systematic review of caries and adiposity reported a similar finding of no relationship between overweight and dental caries that was assessed by BMI (Kantovitz et al. 2006). One possible explanation for this no correlation might be due to the multifactorial etiological nature of both dental caries and obesity that are impacted by various genetic and environmental factors (Alshihri et al. 2019). Some studies showed a positive association between BMI and caries; however, this relationship is stronger and more reliable in permanent than in primary dentition (Al-Ansari & Nazir 2020). The burden of caries and obesity usually increases with age as these diseases gradually accumulate during the life course, therefore a stronger association is observed with an increase in age as a

burden of these diseases also increases with age. In this study similar pattern was observed as obesity was statistically significantly associated with dental caries in permanent teeth. There are some common risk factors mentioned in the literature that can increase the likelihood of both obesity and caries (Hayden et al. 2013). These risk factors can either be modifiable such as behavioural and psychological factors or can be non-modifiable such as biological, genetic, socio-demographic, and cultural (Ashour et al. 2018; Gerdin et al. 2008; Sakeenabi et al. 2012) . Higher sugar consumption is one of the common examples of behavioural modifiable factors that negatively affect both conditions (Hayden et al. 2013). Therefore, the research focusing on dental caries and obesity should include the risk factors and use the integrating and collaborating strategies in prevention of it both locally and globally. The cross-sectional studies design which we used in this study is not suitable for establishing the causal relationship. However, this type of study can provide very important information regarding the prevalence, risk indicators and provide useful data that can encourage further longitudinal studies and help in monitoring the oral health conditions of Saudi Arabian children.

Conclusions

In this study, a significant association was found between obesity and caries experience among school children of Al-Jouf, KSA. This finding provides an important information for the prevention and management of dental caries in children by focusing on specific risk factors associated with these two diseases having common risk determinants. There is a need for a comprehensive and integrated effort by both medical and dental healthcare professionals for the prevention and management of obesity and caries in children by implementing practical solutions. Future research should investigate the specific factors of overweight that can play the protective role against dental caries in permanent dentition.

References

- Akpata E, Al-Shammery AR, and Saeed HI. 1992. Dental caries, sugar consumption and restorative dental care in 12–13-year-old children in Riyadh, Saudi Arabia. *Community dentistry and oral epidemiology* 20:343-346.
- Al-Ansari A, and Nazir M. 2020. Relationship between obesity and dental caries in Saudi male adolescents. *International journal of dentistry* 2020.
- Al Agili DE. 2013. A systematic review of population-based dental caries studies among children in Saudi Arabia. *The Saudi dental journal* 25:3-11.
- Al Othaimen A, Al Nozha M, and Osman A. 2007. Obesity: an emerging problem in Saudi Arabia. Analysis of data from the National Nutrition Survey. *EMHJ-Eastern Mediterranean Health Journal*, 13 (2), 441-448, 2007.
- Alshihri AA, Rogers HJ, Alqahtani MA, and Aldossary MS. 2019. Association between dental caries and obesity in children and young people: a narrative review. *International journal of dentistry* 2019.
- Alves LS, Susin C, Damé-Teixeira N, and Maltz M. 2013. Overweight and obesity are not associated with dental caries among 12-year-old South Brazilian schoolchildren. *Community dentistry and oral epidemiology* 41:224-231.
- Anderson SE, and Whitaker RC. 2009. Prevalence of obesity among US preschool children in different racial and ethnic groups. *Archives of pediatrics & adolescent medicine* 163:344-348.
- Ashour NA, Ashour AA, and Basha S. 2018. Association between body mass index and dental caries among special care female children in Makkah City. *Annals of Saudi medicine* 38:28-35.
- Chaudhary FA, and Ahmad B. 2021. The relationship between psychosocial distress and oral health status in patients with facial burns and mediation by oral health behaviour. *BMC oral health* 21:1-9.
- Chaudhary FA, Ahmad B, and Bashir U. 2019. Dental health status and oral health behaviours of patients with facial burn in Pakistan. *BMC Oral Health* 19:1-10.
- Cheng Y-h, Liao Y, Chen D-y, Wang Y, and Wu Y. 2019. Prevalence of dental caries and its association with body mass index among school-age children in Shenzhen, China. *BMC oral health* 19:1-9.
- Chi DL, Luu M, and Chu F. 2017. A scoping review of epidemiologic risk factors for pediatric obesity: implications for future childhood obesity and dental caries prevention research. *Journal of Public Health Dentistry* 77:S8-S31.
- da Silva RAB, Barreiros D, Oliveira S, da Silva LAB, Nelson-Filho P, and Küchler EC. 2016. Association between body mass index and caries experience in Brazilian children and adolescents. *Journal of Dentistry for Children* 83:146-151.
- Farooqi FA, Khabeer A, Moheet IA, Khan SQ, and Farooq I. 2015. Prevalence of dental caries in primary and permanent teeth and its relation with tooth brushing habits among schoolchildren in Eastern Saudi Arabia. *Saudi medical journal* 36:737.
- Farsi DJ, Elkhodary HM, Merdad LA, Farsi NM, Alaki SM, Alamoudi NM, Bakhaidar HA, and Alolayyan MA. 2016. Prevalence of obesity in elementary school children and its association with dental caries. *Saudi medical journal* 37:1387.
- Fernandez MR, Goettems ML, Demarco FF, and Correa MB. 2017. Is obesity associated to dental caries in Brazilian schoolchildren? *Brazilian oral research* 31.
- Gerdin EW, Angbratt M, Aronsson K, Eriksson E, and Johansson I. 2008. Dental caries and body mass index by socio-economic status in Swedish children. *Community dentistry and oral epidemiology* 36:459-465.

- Gilchrist F, Marshman Z, Deery C, and Rodd HD. 2015. The impact of dental caries on children and young people: what they have to say? *International journal of paediatric dentistry* 25:327-338.
- Gudipani RK, Albilasi RM, HadiAlrewili O, Alam MK, Patil SR, and Saeed F. 2021. Association of Body Mass Index and Waist Circumference With Dental Caries and Consequences of Untreated Dental Caries Among 12-to 14-Year-old Boys: A Cross-Sectional Study. *International Dental Journal*.
- Hayden C, Bowler JO, Chambers S, Freeman R, Humphris G, Richards D, and Cecil JE. 2013. Obesity and dental caries in children: a systematic review and meta-analysis. *Community dentistry and oral epidemiology* 41:289-308.
- James PT. 2004. Obesity: the worldwide epidemic. *Clinics in dermatology* 22:276-280.
- Kantovitz KR, Pascon FM, Rontani RMP, Gavião MBD, and Pascon FM. 2006. Obesity and dental caries--A systematic review. *Oral health & preventive dentistry* 4.
- Kopycka-Kedzierawski D, Auinger P, Billings R, and Weitzman M. 2008. Caries status and overweight in 2-to 18-year-old US children: findings from national surveys. *Community dentistry and oral epidemiology* 36:157-167.
- Kumar S, Kroon J, Lalloo R, Kulkarni S, and Johnson NW. 2017. Relationship between body mass index and dental caries in children, and the influence of socio-economic status. *International Dental Journal* 67:91-97.
- Organization WH. 2013. *Oral health surveys: basic methods*: World Health Organization.
- Sakeenabi B, Swamy HS, and Mohammed RN. 2012. Association between obesity, dental caries and socioeconomic status in 6-and 13-year-old school children. *Oral health & preventive dentistry* 10.
- Scorzetti L, Marcattili D, Pasini M, Mattei A, Marchetti E, and Marzo G. 2013. Association between obesity and periodontal disease in children. *Eur J Paediatr Dent* 14:181-184.
- Spiegel KA, and Palmer CA. 2012. Childhood dental caries and childhood obesity. Different problems with overlapping causes. *American journal of dentistry* 25:59-64.
- Sturm R. 2002. The effects of obesity, smoking, and drinking on medical problems and costs. *Health affairs* 21:245-253.
- Swinburn BA, Sacks G, Hall KD, McPherson K, Finegood DT, Moodie ML, and Gortmaker SL. 2011. The global obesity pandemic: shaped by global drivers and local environments. *The Lancet* 378:804-814.
- Tsai A, Hsiang C, and Johnsen D. 2000. Caries levels and patterns in the primary dentition of preschool children in Taiwan. *Chang Gung medical journal* 23:22-27.
- Wake M, Nicholson JM, Hardy P, and Smith K. 2007. Preschooler obesity and parenting styles of mothers and fathers: Australian national population study. *Pediatrics* 120:e1520-e1527.

Table 1 (on next page)

Participant characteristics (N=380).

Table 1. Participant characteristics (N=380).

Gender	N (%)
Boys	211 (55.5%)
Girls	169 (44.5%)
Total	380
Age (years)	
6	24 (6.3%)
7	31 (8.9%)
8	42 (11%)
9	62 (16.3%)
10	32 (8.4%)
11	41 (10.6%)
12	62 (16.1%)
13	44 (11.4%)
14	42 (11%)
Mean age	9.8 Years
Range	(6-14 years)
BMI Scores	Mean (Range)
Male	28.2 (17-34)
Female	24.3 (16-30)
Overall	26.4 (16-34)

Table 2(on next page)

Comparison between dental caries, oral hygiene practices and dietary habits of participants.

Table 2. Comparison between dental caries, oral hygiene practices and dietary habits of participants.

	Mean dmft/DMFT scores		Caries prevalence (dmft/DMFT≥1)
	Deciduous	Permanent	
Gender			
Male	3.8±1.2	3±1.3	79.4%
Female	3.6±1.4	2.7±0.8	72.6%
Overall	3.7±1.6	2.8±1.0	76.1%
Unpaired 't' test (t value)	1.54	3.22	
P- value	0.69	0.04*	
Brush teeth per day			
Less than once	4.2±1.8	3.2±1.6	81.3%
Once a day	3.4± 0.8	2.2± 1	73.2%
Twice a day	2.1± 1.2	2±1.2	76.5%
More than Twice a day	1.9±0.9	1.8±1.4	75.1%
ANOVA (F value)	1.29	0.51	
P-value	0.27	0.72	
Dental floss			
Yes	3.8±1.4	2.4±1.8	74.8%
No	3.6±1.6	2.2±1.4	78.5%
Unpaired 't' test (t value)	-0.92	-0.62	
P- value	0.43	0.53	
Mouthwash/rinsing habit			
Yes	3.2±1.2	2.0±0.8	75%
No	3.9±1.4	2.8±1.2	77.3%
Unpaired 't' test	0.51	1.24	
P- value	0.82	0.07	
Frequency of consumption of Sucrose containing foods			
Once a day	3.4±2.1	2±0.8	66.4%
2-4 times a day	3.2±1.8	2.2±1.2	70%
4-6 times a day	2.8±1.2	2.3±1.6	74.2%
More than 6 times a day	3.8± 1.0	2.6±1.8	81.3%
ANOVA (F value)	0.99	3.82	
P-value	0.23	0.01*	

SD- Standard Deviation, * Statistically Significant.

Table 3(on next page)

Relationship between BMI and dental caries experience among study participants.

Table 3. Relationship between BMI and dental caries experience among study participants.

	BMI group					ANOVA	
						F value	P value
	Moderate malnutrition	Mild malnutrition	Normal	Over weight	Obese		
Mean dmft	3.4±2.1	3±1.6	3.6±1.8	3.6±2.2	4.8±2.8	1.23	0.08
Mean DMFT	2±1.1	2.4±1.4	2.4±0.8	2.8±1.6	3.9±2.2*	4.36	0.001*
Caries Prevalence	64.2%	65.6%	67.3%	78%	83.6%		

* Statistically Significant

Table 4(on next page)

Multiple regression analysis to identify factors influencing DMFT score.

Table 4. Multiple regression analysis to identify factors influencing DMFT score.

Model	Unstandardized Coefficients		Standardized coefficients	P-value	95% CI
	b	S.E	b		
1	(Constant)	2.111	.383	--	0.001*
	Sugar intake	0.43	.080	0.42	0.29
	BMI	3.20	1.69	0.82	0.001*
	Gender	0.04	.076	0.04	0.52

* Statistically Significant