

# Curcumin effect on *Acanthamoeba triangularis* encystation under nutrient starvation

Rachasak Boonhok<sup>1</sup>, Suthinee Sangkanu<sup>2</sup>, Suganya Phumjan<sup>1</sup>, Ramita Jongboonjua<sup>1</sup>, Nawarat Sangnopparat<sup>1</sup>, Pattamaporn Kwankaew<sup>1</sup>, Aman Tedasen<sup>1</sup>, Chooi Ling Lim<sup>3</sup>, Maria de Lourdes Pereira<sup>4</sup>, Mohammed Rahmatullah<sup>5</sup>, Polrat Wilairatana<sup>6</sup>, Christophe Wiart<sup>7</sup>, Karma G. Dolma<sup>8</sup>, Alok K. Paul<sup>9</sup>, Madhu Gupta<sup>10</sup> and Veeranoot Nissapatorn<sup>2</sup>

<sup>1</sup> Department of Medical Technology, School of Allied Health Sciences, and Research Excellence Center for Innovation and Health Products (RECIHP), Walailak University, Thai Buri, Nakhon Si Thammarat, Thailand

<sup>2</sup> School of Allied Health Sciences, Southeast Asia Water Team (SEA Water Team) and World Union for Herbal Drug Discovery (WUHeDD), and Research Excellence Center for Innovation and Health Products (RECIHP), Walailak University, Thai Buri, Nakhon Si Thammarat, Thailand

<sup>3</sup> Division of Applied Biomedical Science and Biotechnology, School of Health Sciences, International Medical University, Kuala Lumpur, Malaysia

<sup>4</sup> CICECO-Aveiro Institute of Materials and Department of Medical Sciences, University of Aveiro, Aveiro, Portugal

<sup>5</sup> Department of Biotechnology and Genetic Engineering, University of Development Alternative, Dhaka, Bangladesh

<sup>6</sup> Department of Clinical Tropical Medicine, Faculty of Tropical Medicine, Mahidol University, Rachathewe, Bangkok, Thailand

<sup>7</sup> The Institute for Tropical Biology and Conservation, Universiti Malaysia Sabah, Kota Kinabalu, Sabah, Malaysia

<sup>8</sup> Department of Microbiology, Sikkim Manipal Institute of Medical Sciences, Sikkim, India

<sup>9</sup> School of Pharmacy and Pharmacology, University of Tasmania, Hobart, Tasmania, Australia

<sup>10</sup> Department of Pharmaceutics, Delhi Pharmaceutical Sciences and Research University, New Delhi, India

Submitted 25 March 2022

Accepted 9 June 2022

Published 4 July 2022

Corresponding authors

Polrat Wilairatana,  
polrat.wil@mahidol.ac.th  
Veeranoot Nissapatorn,  
veeranoot.ni@wu.ac.th

Academic editor

Erika Braga

Additional Information and  
Declarations can be found on  
page 18

DOI 10.7717/peerj.13657

© Copyright

2022 Boonhok et al.

Distributed under

Creative Commons CC-BY 4.0

## ABSTRACT

**Background:** Curcumin is an active compound derived from turmeric, *Curcuma longa*, and is known for its benefits to human health. The amoebicidal activity of curcumin against *Acanthamoeba triangularis* was recently discovered. However, a physiological change of intracellular pathways related to *A. triangularis* encystation mechanism, including autophagy in the surviving amoeba after curcumin treatment, has never been reported. This study aims to investigate the effect of curcumin on the survival of *A. triangularis* under nutrient starvation and nutrient-rich condition, as well as to evaluate the *A. triangularis* encystation and a physiological change of *Acanthamoeba* autophagy at the mRNA level.

**Methods:** In this study, *A. triangularis* amoebas were treated with a sublethal dose of curcumin under nutrient starvation and nutrient-rich condition and the surviving amoebas was investigated. Cysts formation and vacuolization were examined by microscopy and transcriptional expression of autophagy-related genes and other encystation-related genes were evaluated by real-time PCR.

## OPEN ACCESS

**Results:** *A. triangularis* cysts were formed under nutrient starvation. However, in the presence of the autophagy inhibitor, 3-methyladenine (3-MA), the percentage of cysts was significantly reduced. Interestingly, in the presence of curcumin, most of the parasites remained in the trophozoite stage in both the starvation and nutrient-rich condition. In vacuolization analysis, the percentage of amoebas with enlarged vacuole was increased upon starvation. However, the percentage was significantly declined in the presence of curcumin and 3-MA. Molecular analysis of *A. triangularis* autophagy-related (ATG) genes showed that the mRNA expression of the ATG genes, ATG3, ATG8b, ATG12, ATG16, under the starvation with curcumin was at a basal level along the treatment. The results were similar to those of the curcumin-treated amoebas under a nutrient-rich condition, except AcATG16 which increased later. On the other hand, mRNA expression of encystation-related genes, cellulose synthase and serine proteinase, remained unchanged during the first 18 h, but significantly increased at 24 h post treatment.

**Conclusion:** Curcumin inhibits cyst formation in surviving trophozoites, which may result from its effect on mRNA expression of key *Acanthamoeba* ATG-related genes. However, further investigation into the mechanism of curcumin in *A. triangularis* trophozoites arrest and its association with autophagy or other encystation-related pathways is needed to support the future use of curcumin.

**Subjects** Cell Biology, Microbiology, Molecular Biology, Parasitology

**Keywords** Autophagy, *Acanthamoeba triangularis*, Curcumin, Encystation, Nutrient Starvation, Real-time PCR

## INTRODUCTION

*Acanthamoeba* spp. are free-living amoebas present in the environment, particularly in soil and water (Siddiqui & Khan, 2012). Several species of *Acanthamoeba* have been characterized (Chelkha et al., 2020), and most of the human pathogenic species are classified into T4 genotype, for example, *Acanthamoeba castellanii*, *A. polyphaga*, and *A. triangularis* (Guimaraes et al., 2016; Juarez et al., 2018; Hussain et al., 2020). *Acanthamoeba* spp. are transmitted to humans by different routes (Neelam & Niederkorn, 2017; de Lacerda & Lira, 2021) and lead to various clinical presentations, especially in immunocompromised individuals who may have granulomatous amebic encephalitis (Matson et al., 1988), chronic sinusitis (Kim et al., 2000), or cutaneous lesions (Morrison et al., 2016). In addition, for healthy individuals who wear contact lenses, this group is at risk of *Acanthamoeba* infection if they have poor personal hygiene habits and *Acanthamoeba* keratitis (AK), a well-known ocular disease caused by this protozoan parasite, usually present in the group of people (Lorenzo-Morales, Khan & Walochnik, 2015; Neelam & Niederkorn, 2017; Khan, Anwar & Siddiqui, 2019). Regarding *Acanthamoeba* life cycle, the amoeba usually presents a trophozoite stage, a metabolically active form and can multiply within the human host. However, after being exposed to a stressful condition, it can transform into a cyst form with a double wall that is more resistant to a harsh environment (Anwar, Khan & Siddiqui, 2018). This form is a major barrier to *Acanthamoeba* treatment. So far, several drugs have been approved by the

United States Food and Drug Administration, but standard therapeutic management of *Acanthamoeba*-infected patients is not yet available (Elsheikha, Siddiqui & Khan, 2020). The two most commonly used first-line drugs for *Acanthamoeba* treatment, especially in AK patients, still use chlorhexidine and polyhexamethylene biguanide. However, identifying new compounds and screening natural extracts for amoebicidal activity are still attractive approaches for further studies. It could provide an alternative drug for *Acanthamoeba* treatment or be used as a complementary treatment for *Acanthamoeba* infection in the future.

Autophagy is a lysosomal degradation pathway for intracellular cytosolic materials (Yorimitsu & Klionsky, 2005; Feng et al., 2014). This mechanism is essential for all eukaryotic cells to supply energy and support cell survival. In humans, a defect of the autophagy mechanism is associated with several diseases, for example, neurodegenerative diseases (Menzies, Fleming & Rubinsztein, 2015), non-alcoholic fatty liver disease (Khambu et al., 2018), or infectious diseases (Castillo et al., 2012; Brinck Andersen et al., 2020). Starvation or nutrient depletion is a classical stress condition for autophagy induction both *in vitro* and *in vivo* (Mizushima et al., 2004; Suzuki, 2013). Several autophagy-related (Atg) proteins participate in the formation of a double-membrane vacuole called autophagosome (Eskelinen, 2005; Feng et al., 2014). In mammals, more than 30 Atg proteins have been identified (Feng et al., 2014). However, a partial list of ATG genes is conserved in free-living amoeba, including *Acanthamoeba* spp., and some Atg proteins have been identified (Picazarri, Nakada-Tsukui & Nozaki, 2008; Moon et al., 2009; Song et al., 2012; Kim et al., 2015). *Acanthamoeba* autophagy is of interest as a number of Atg proteins have been partially characterized and reported to be involved with *Acanthamoeba* encystation, a mechanism in which trophozoites transform to cysts (Moon et al., 2011; Song et al., 2012; Moon et al., 2013; Kim et al., 2015). Hence, a study of autophagy at both transcriptional and protein levels is needed to understand its biological functions and interaction with other intracellular pathways, which further extends to its association with their pathogenesis in humans.

Curcumin, an active compound obtained from turmeric, *Curcuma longa* (Kocaadam & Şanlıer, 2017), contains several pharmacological activities, for example, anti-inflammatory (Wal et al., 2019), antioxidant (Jakubczyk et al., 2020), anti-cancer (Vallianou et al., 2015; Tomeh, Hadianamrei & Zhao, 2019), and antimicrobial activities (Cui, Miao & Cui, 2007; Martins et al., 2009; Teow et al., 2016; Yang et al., 2016; Mitsuwan et al., 2020). The amoebicidal activity of curcumin against *A. triangularis* trophozoites and cysts was recently identified (Mitsuwan et al., 2020). It reveals another property of curcumin against this water-borne parasitic pathogen and could be a promising compound for further drug development against *Acanthamoeba* infection. In this study, we investigated the effect of curcumin on surviving *A. triangularis* trophozoites after being exposed to a sublethal dose of curcumin under nutrient starvation and nutrient-rich condition. Cyst formation and vacuolization were examined by microscopic observation and molecular analysis of autophagy-related as well as other encystation-related genes at the transcriptional level, was investigated by real-time PCR. This raises another point of concern, in addition to the killing activity by plant extract or compound where surviving amoebas after the treatment

are likely to transform into a cyst, with an emphasis on *Acanthamoeba* autophagy, which is one of the pathways involved with *Acanthamoeba* encystation.

## MATERIALS AND METHODS

### *A. triangularis* cultivation

PYG medium, a nutrient-rich condition or full medium, (2% (w/v) proteose peptone, 0.1% (w/v) yeast extract, 400  $\mu$ M CaCl<sub>2</sub>, 4 mM MgSO<sub>4</sub>, 2.5 mM Na<sub>2</sub>HPO<sub>4</sub>, 2.5 mM KH<sub>2</sub>PO<sub>4</sub>, 50  $\mu$ M (NH<sub>4</sub>)<sub>2</sub>Fe(SO<sub>4</sub>)<sub>2</sub>, 100 mM glucose) was used to grow *A. triangularis* trophozoites, strain WU19001 (*Mitsuwan et al., 2020*). The parasite was maintained at room temperature (RT) without shaking (*Taravaud, Loiseau & Pomel, 2017*). The culture medium was replaced with fresh PYG every 2 days until trophozoites harvesting. To induce *A. triangularis* cysts, trophozoites were washed and grown in PAS supplemented with 5% glucose, a nutrient-depleted condition, called starvation, which was modified from Aqeel and colleagues based on our initial in-house laboratory trials (*Aqeel et al., 2013*). The PAS powder, obtained from HiMedia, Mumbai, India, consisted of NaCl, MgSO<sub>4</sub>·7H<sub>2</sub>O, CaCl<sub>2</sub>·2H<sub>2</sub>O, Na<sub>2</sub>HPO<sub>4</sub> and KH<sub>2</sub>PO<sub>4</sub>.

### Curcumin preparation and determination of the half-maximal inhibitory concentration (IC<sub>50</sub>)

Curcumin powder was commercially purchased (Sigma Aldrich, St. Louis, MI, USA). The curcumin was dissolved in 100% DMSO and prepared at stock 750 mg/mL. This was further diluted to the working concentration with medium. The identification of the IC<sub>50</sub> against *A. triangularis* trophozoites was performed in 96-well black plate (SPL Life Sciences, Seoul, Korea). Curcumin concentration was prepared with 2-fold serial dilution with starting final concentration of 8,000  $\mu$ g/mL. Thus, the maximum of the final %DMSO was 2.13. Trophozoites were harvested and washed with fresh AnaeroGRO™ Peptone Yeast Extract Glucose Broth: Proteose peptone and yeast extract were purchased from HiMedia Laboratories, Mumbai, India. Sodium citrate dihydrate (C<sub>6</sub>H<sub>5</sub>Na<sub>3</sub>O<sub>7</sub>·2H<sub>2</sub>O), disodium phosphate (NaHPO<sub>4</sub>), sodium chloride (NaCl), calcium chloride (CaCl<sub>2</sub>), and glucose were purchased from Sigma Chemical Co. (St. Louis, MO, USA). Potassium dihydrogen phosphate (KH<sub>2</sub>PO<sub>4</sub>) and magnesium sulfate heptahydrate (MgSO<sub>4</sub>·7H<sub>2</sub>O) were procured from Labscan (Bangkok, Thailand). Trypan blue (0.4%) was obtained from Gibco BRL (Grand Island, NY, USA). All chemicals and medium components used were of analytical grade and added at  $2 \times 10^4$  cells/well. A control group of untreated cells and PYG medium alone was included. All edge wells were filled with Page's saline (PAS) buffer. After 24 h post-treatment, the parasite viability was analyzed by PrestoBlue® reagent (Invitrogen, Waltham, MA, USA) staining according to the manufacturer's protocol. The plate was incubated for 30 min at 37 °C incubators, and fluorescence intensity was measured at excitation/emission wavelength of 535/615 nm by a microplate reader (BioTek SynergyTMMX microplate reader, Winooski, VT, USA). The IC<sub>50</sub> of curcumin was then calculated by prism5 software (GraphPad Software, San Diego, CA, USA). The experiments were conducted in triplicate with three independent experiments.

### Analysis of cysts formation and vacuolization

*A. triangularis* trophozoites were cultured in PYG medium or PAS supplemented with 5% glucose, and the curcumin was added at a final concentration of 50 µg/mL for 24 h. After Trypan Blue staining, the parasites were assessed for cysts formation and vacuolization every 6 h after the curcumin treatment. At least 200 viable cells per condition were investigated, and different forms of the parasites *i.e.* irregular trophozoites, rounded trophozoites, and cysts, were identified under a light microscope. The percentage of cyst and proportion of parasite forms at each time point was calculated. The irregular trophozoites were further evaluated for their vacuole formation, and the surviving trophozoites of at least 100 cells per condition were analyzed. Trophozoites with vacuoles, regardless of their size, as well as the trophozoites containing enlarged vacuoles, were examined. An enlarged vacuole (EV) was defined as a vacuole with a diameter of at least 5 µm, and the trophozoite containing at least one EV was counted as 1 (Boonhok *et al.*, 2021b). The experiment was performed with three independent experiments.

### Determination of minimal inhibitory concentration (MIC) and drug combination assay

A drug combination study of chlorhexidine, a standard anti-*Acanthamoeba* drug, and curcumin was performed for their amoebicidal activity. The minimum inhibitory concentration (MIC) of curcumin and chlorhexidine was identified along with the microtiter broth dilution method (Mitsuwan *et al.*, 2020). The drug/compound was prepared in 96-well clear plate (SPL Life Sciences, Seoul, Korea), and trophozoites of  $2 \times 10^5$  cells/100 µL were then added into each well. The plates were incubated at RT in the dark for 24 h. The parasite viability was quantified by Trypan Blue staining under a light microscope, Eclipse TE2000-S (Nikon, Konan, Minato-ku, Tokyo, Japan). The MIC value in our study referred to the lowest concentration with *A. triangularis* growth inhibition greater than 90%. Thus, the MIC of curcumin and chlorhexidine was 250 and 16 µg/mL, respectively. Their MICs were used in drug combination assay as a starting concentration. The drug and curcumin were prepared in 96-well plate with 2-fold serial dilution. Then, the trophozoites were added and incubated at RT for 24 h before quantification of parasite viability. To examine the effect of curcumin in combination with autophagy inhibitors, 3-methyladenine (3MA) and wortmannin, which were purchased from Sigma Aldrich (St. Louis, MI, USA), the drug combination assay was then performed as mentioned above. However, the starting concentration of the inhibitors was used at 20 µg/mL to cover the concentration tested in this study.

### Preparation of total RNA and cDNA synthesis

*A. triangularis* trophozoites were cultured with PYG medium or PAS supplemented with 5% glucose medium in a 24-well transparent plate with a final number of  $2 \times 10^5$  cells per well. Parasites were treated with curcumin at a final concentration of 50 µg/mL and incubated at room temperature for 24 h. Then, the parasites were harvested every 6 h after treatment (6, 12, 18, 24 h). Parasite preparation for RNA extraction and cDNA synthesis

were performed as described by Boonhok and his colleagues (Boonhok et al., 2021a). The cDNA was kept at  $-20^{\circ}\text{C}$  until use.

### Validation of PCR primers


All specific primers against *Acanthamoeba* genes used in this study are listed in Table S4. The target genes were ATG3 (GenBank accession no. GU270859), ATG8b (GenBank accession no. KC524507.1), ATG12 (GenBank accession no. HQ830265.1), ATG16 (GenBank accession no. FJ906697), cellulose synthase (CS) (GenBank accession no. EDCBI66TR), serine proteinase (SP) (GenBank accession no. EU365404), metacaspase (MCA) (GenBank accession no. AF480890), interleukin-1 converting enzyme-like protease (IL) (GenBank accession no. XM\_004338552), and 18S rRNA was used as a reference gene. These primers were tested against *A. triangularis* strain WU19001 DNA. To confirm primer specificity, the PCR product was sent for sequencing (Apical Scientific Sdn. Bhd., Seri Kembangan, Selangor, Malaysia), and the DNA sequence was then analyzed and blasted against *A. castellanii* NCBI databases before performing a quantitative PCR (Boonhok et al., 2021a).

### Analysis of gene expression by quantitative PCR

iTaq Universal SYBR Green Supermix Kit obtained from Bio-Rad (Bio-Rad, Hercules, CA, USA) was used to prepare the quantitative PCR reaction (qPCR) along with the manufacturer's instruction. The 18S rDNA was used as a reference gene. The reaction was conducted in a PCR tube, which consisted of 10  $\mu\text{L}$  of 2 $\times$  iTaq Universal SYBR Green Supermix, 100 ng cDNA, and 1  $\mu\text{L}$  of 200 mM F+R primers. The total volume was adjusted with DEPC water to 20  $\mu\text{L}$ . Thermal cycler, StepOnePlus Real-time PCR systems (Applied Biosystems, Waltham, MA, USA), the software was set as follows; holding stage  $95^{\circ}\text{C}$  for 30 s, cycling stage for 40 cycles at  $95^{\circ}\text{C}$  for 15 s,  $60^{\circ}\text{C}$  for 60 s, and melting curve stage at  $95^{\circ}\text{C}$  for 15 s,  $60^{\circ}\text{C}$  for 60 s,  $95^{\circ}\text{C}$  for 15 s with a temperature increase of  $0.3^{\circ}\text{C}$ . The average deltaCt ( $\Delta\text{Ct}$ ) was obtained by the thermal cycler. The delta-delta Ct ( $\Delta\Delta\text{Ct}$ ) and a relative expression of the mRNA were calculated as follows, the  $\Delta\Delta\text{Ct} = ((\text{Ct of treated sample GOI} - \text{Ct of treated sample housekeeper}) - (\text{Ct of untreated control GOI} - \text{Ct of untreated control housekeeper}))$  where GOI refers to the gene of interest. The relative expression of mRNA = 2 to the power of (minus X) or  $2^{-X}$  where X is  $\Delta\Delta\text{Ct}$ . The interpretation of the result is if the value  $>1$ ,  $<1$ , and 1 means that the expression is increased, decreased, and constant, respectively.

### Statistical data analysis

Experiments were conducted with 2–3 technical replicates in three independent experiments. All data were recorded in Microsoft Excel 2016 (Microsoft Corporation, Redmond, Washington, USA). The statistical analysis was done using Prism five software (Graphpad Software, San Diego, CA, USA), and the mean  $\pm$  SD or  $\pm$  SEM was used, including a two-tailed unpaired Student's t-test. *P* values less than 0.05 were considered statistically significant.



**Figure 1** *A. triangularis* cysts formation under starvation. (A) The trophozoites were cultured in the starvation medium, PAS supplemented with 5% glucose, for 24 h. The parasites grew in PYG medium, or full medium was used as a control. The parasites were stained with Trypan Blue, and the viable parasites were analyzed under a microscope at the indicated time points. *A. triangularis* cysts were counted and represented as mean percentage  $\pm$  SD. Data were obtained from three independent experiments. \*\*\* $P < 0.001$ . (B) Representative images of parasites cultured in full and starved medium. Bar 20  $\mu$ M. White and black arrowheads indicate the irregular trophozoites and rounded trophozoites, respectively, while the unfilled arrowhead indicates cysts. [Full-size !\[\]\(fd7fe780e8fd8eece60268c87d0c3e04\_img.jpg\) DOI: 10.7717/peerj.13657/fig-1](https://doi.org/10.7717/peerj.13657/fig-1)

## RESULTS

### Starvation induces *A. triangularis* encystation

Starvation or nutrient-depleted condition was used to induce *A. triangularis* encystation. The mean percentage of *A. triangularis* cysts was approximately 53.30%, and it was significantly different from that of the nutrient-rich condition or full medium (Fig. 1A).


The representative images of the parasites under starved and full conditions are shown in [Fig. 1B](#).

### Effect of curcumin on *A. triangularis* autophagy under starvation

The  $IC_{50}$  of curcumin against *A. triangularis* trophozoites under the full condition at 24 h treatment was first identified, and the  $IC_{50}$  was  $48.64 \pm 30.86$   $\mu\text{g/mL}$ . The representative data is shown in [Fig. S1](#). The curcumin concentration of 50  $\mu\text{g/mL}$  was then used as representative curcumin concentration throughout this study. Starvation alone was included as a positive control for *A. triangularis* encystation. Starvation is known as a classical autophagy inducer in several eukaryotic cells ([Kamada, Sekito & Ohsumi, 2004](#); [Díaz-Troya et al., 2008](#)); thus, in this assay, we included autophagy inhibitors, 3MA, and wortmannin, to examine a physiological and morphological change of *A. triangularis* upon the treatment. Our result showed that in the presence of 1 mM 3MA, the encystation was significantly decreased, whereas the 1  $\mu\text{M}$  wortmannin was slightly impacted to the encystment. The percentage of cysts under starvation + 3MA or wortmannin was approximately 22.55% and 41.30%, respectively ([Fig. 2A](#)). We next investigated whether curcumin stress supports cysts formation under starved conditions. Interestingly, in the presence of 50  $\mu\text{g/mL}$  curcumin, the percentage of cysts was approximately 1.39%, the surviving amoeba remained in the trophozoites stage ([Fig. 2A](#)). The representative images of the curcumin-treated parasite under starved conditions are shown in [Fig. 2B](#). In addition, the parasites treated with a combination of curcumin and autophagy inhibitors were included in this experiment to test whether this could completely inhibit the encystation or not. The result showed that the percentage of cysts in curcumin + 3MA and curcumin + wortmannin remained at the basal level similar to that of curcumin-treated alone ([Fig. 2A](#)). Moreover, different viable forms of *A. triangularis*, namely, irregular trophozoites, rounded trophozoites, and cysts under different conditions, starvation alone, starvation + 3MA, and starvation + curcumin, were quantified under the microscope every 6 h. In starvation alone, the irregular trophozoites started to round off at 6 h. Cysts were seen 12 h after culture, and at 24 h, the percentage of cysts was approximately 50%. In the presence of 3MA, cysts were clearly seen at 18 h post-treatment, and at 24 h, the percentage of cysts was approximately 20%. Under curcumin-treated conditions, the parasites were mainly in irregular trophozoites, approximately 90%, and the percentage of cysts was approximately 2% ([Fig. S2](#)).

The evaluation of vacuolization in surviving trophozoites was further performed. The number of trophozoites containing vacuole and trophozoites with enlarged vacuole were analyzed under the light microscope. The percentage of trophozoites with vacuoles in the curcumin-treated condition was almost 100% along with the 24 h treatment, and the percentage was similar to that of starvation alone and starvation + 3MA ([Fig. S3A](#)). The percentage of trophozoites with enlarged vacuoles was further investigated. In starvation alone, the percentage increased significantly with the treatment, and the mean percentage at 24 h was approximately 22.41%. Interestingly, in the curcumin-treated condition, the mean percentage along the treatment was significantly reduced and





**Figure 2** *A. triangularis* response to curcumin under starved condition. (A) Cysts formation, the trophozoites were cultured in a starvation medium, PAS+5% glucose, with autophagy inhibitors or 50  $\mu\text{g}/\text{mL}$  curcumin  $\pm$  autophagy inhibitors for 24 h. Starvation alone was included as a positive control. Cyst was quantified every 6 h post-treatment. The percentage of cyst was calculated and represented as mean  $\pm$  SD. Data were obtained from three independent experiments. NS, not significant;  $**P < 0.01$ ;  $***P < 0.001$ . (B) Representative image of curcumin-treated parasites under starved condition. Bar 20  $\mu\text{m}$ . (C) Vacuolization in surviving trophozoites, at least 100 cells, the trophozoites per condition were examined for enlarged vacuole, a diameter of at least 5  $\mu\text{m}$ . Data were obtained from three independent experiments and represented as a mean percentage  $\pm$  SD.  $**P < 0.01$ ;  $***P < 0.001$ . (D) Transcriptional expression of autophagy-related genes after curcumin treatment, *A. triangularis* trophozoites were cultured in starvation medium with or without 50  $\mu\text{g}/\text{mL}$  curcumin for 24 h. The parasites were harvested every 6 h, and the mRNA level of ATG3, ATG8b, ATG12, ATG16 genes were analyzed by qPCR. Their expression at each time point was expressed as a relative mRNA expression. The 18S rRNA was included as a reference gene. The expression at time 0 h was set to 1. The data were obtained from three independent experiments. Bar graphs represent mean  $\pm$  SEM.

Full-size DOI: [10.7717/peerj.13657/fig-2](https://doi.org/10.7717/peerj.13657/fig-2)

maintained in the range of 7.89–12.16%. The result was similar to the 3MA-treated condition where the percentage was in the range of 6.79–10.92% (Fig. 2C).

The molecular analysis of *A. triangularis* autophagy-related genes, ATG3, ATG8b, ATG12, and ATG16, was conducted at transcriptional level upon curcumin treatment. The validation of PCR primers (Table S4) by conventional PCR against *A. triangularis* DNA was performed, and the target genes were successfully amplified. The gel result representing PCR products is shown in Fig. S4. In addition, analysis of DNA sequencing of the amplicons was performed, and the results are shown in Table S5. Then, the quantitative PCR was performed, and the results showed that the mRNA expression of all tested ATG genes was unchanged along with the treatment and maintained at the basal level (Fig. 2D). In addition, the expression of these ATG genes under 3MA-treated conditions was investigated. As expected, the expression of the ATG genes was at the basal level along with the treatment (Fig. S5). The overall results demonstrated the inhibitory effect of curcumin on the surviving amoebas against *A. triangularis* encystation even under starvation.

### Effect of curcumin on *A. triangularis* autophagy under a nutrient-rich condition


To measure the effect of curcumin alone without the stress of starvation, *A. triangularis* trophozoites were cultured in PYG, a nutrient-rich medium. As expected, curcumin did not activate cyst formation. The percentage of cysts was at a basal level and was not significantly different from that of the full medium alone (Fig. 3A). The representative image of the parasites under curcumin treatment is shown in Fig. 3B. The surviving parasites remained in the trophozoite stage.

Vacuolization in the surviving trophozoites was then analyzed. The percentage of trophozoites with vacuoles under curcumin treatment was nearly 100%, and the percentage was at a level comparable to full medium alone (Fig. 3C). To investigate the maturation of vacuole, parasites with enlarged vacuoles were examined. The percentage of trophozoites with enlarged vacuoles was consistent with the treatment, in the range of 6.23–6.81%.

Molecular analysis by qPCR revealed that mRNA expression of the autophagy-related genes, that is ATG3, ATG8b, ATG12 genes, were at the basal level throughout the treatment while ATG16 mRNA expression was increased at 18 and 24 h post-treatment (Fig. 3D).

### Effect of curcumin on *A. triangularis* encystation-related genes under a nutrient-rich condition


Apart from autophagy, we further assessed other *A. triangularis* encystation-related genes, namely, cellulose synthase (CS) and serine proteinase (SP) upon curcumin treatment. The mRNA expression pattern of both genes was similar in that expression was slightly changed during the first 18 h, and significantly increased at 24 h post treatment (Fig. 4).



**Figure 3** *A. triangularis* response to curcumin under a nutrient-rich condition. (A) Cyst formation, the *Acanthamoeba* trophozoites were cultured in a PYG medium with or without 50  $\mu\text{g}/\text{mL}$  curcumin for 24 h. Starvation was included as a positive control for cysts formation. Cysts were quantified every 6 h post treatment. A percentage of cysts was calculated and represented as mean  $\pm$  SD. Data obtained from three independent experiments. NS, not significant;  $***P < 0.001$ . (B) Representative image of curcumin-treated parasites under a full condition. Bar 20  $\mu\text{m}$ . (C) Vacuolization in surviving trophozoites, at least 100 cells the trophozoites per condition were examined for enlarged vacuoles, a diameter of at least 5  $\mu\text{m}$ . Data obtained from three independent experiments and represented as a mean percentage  $\pm$  SD. NS, not significant. (D) Transcriptional expression of autophagy-related genes after curcumin treatment, *A. triangularis* trophozoites were cultured in PYG medium with or without 50  $\mu\text{g}/\text{mL}$  curcumin for 24 h. The parasites were harvested every 6 h and the mRNA level of ATG3, ATG8b, ATG12, ATG16 genes were analyzed by qPCR. Their expression at each time point was expressed as a relative mRNA expression. 18S rRNA was included as a reference gene. The expression at time 0 h was set to 1. Data were obtained from three independent experiments. Bar graphs represent mean  $\pm$  SEM.


Full-size DOI: [10.7717/peerj.13657/fig-3](https://doi.org/10.7717/peerj.13657/fig-3)

Moreover, due to a crosstalk between autophagy and apoptosis in other eukaryotic cells, we also observed mRNA expression of genes involved in the apoptosis pathway, namely, metacaspase (MCA) and interleukin-1 converting enzyme-like protease (IL) in response to curcumin. MCA mRNA expression was slightly changed, but still at the basal level along with the treatment. The expression of IL mRNA was rapidly increased at 6 h post-treatment and gradually declined at later time points. However, the increase expression was again observed at 24 h post-treatment (Fig. S6). The response of surviving *A. triangularis* to curcumin either under nutrient starvation or nutrient-rich condition was illustrated in Fig. 5.



**Figure 4** Transcriptional expression of other encystation-related genes of *A. triangularis* in response to curcumin under a nutrient-rich condition. Investigation of cellulose synthase (CS) and serine proteinase (SP) mRNA expression was carried out. cDNA samples were shared with autophagy analysis. The qPCR was performed, and 18S rRNA was included as a reference gene. Data were obtained from three independent experiments. Bar graphs displayed mean  $\pm$  SEM. \*\* $P < 0.01$ ; \*\*\* $P < 0.001$ .

Full-size DOI: 10.7717/peerj.13657/fig-4



**Figure 5** *A. triangularis* response to curcumin. Curcumin at the sublethal dose was used for *A. triangularis* trophozoites treatment. Approximately 50% of the parasites died, while the surviving parasites remained in the trophozoites stage. Transcriptional expression of tested autophagy-related genes was at the basal level in both starvation and a nutrient-rich condition except *AcATG16*, which increased at later time points under the nutrient-rich condition. Other *A. triangularis* encystation-related genes tested in this study, cellulose synthase and serine proteinase, were also increased at a later time point. Inducing an arrest in the trophozoites by curcumin is possibly resulting in the deactivation of the ATG genes and subsequent inhibition of vacuoles maturation.

Full-size DOI: 10.7717/peerj.13657/fig-5

### Curcumin-based drug combination study

A drug combination study between curcumin and chlorhexidine was performed, and our previous results on the co-treatment of curcumin and autophagy inhibitors, 3MA and wortmannin, did not completely inhibit *A. triangularis* encystation, thus these combinations were included in this assay to explore their interaction. The concentration of compound/drug was varied based on their MICs, except for the autophagy inhibitors which were designed to cover the concentration used in the previous experiment. The MICs of curcumin and chlorhexidine were started at 250 and 16  $\mu\text{g}/\text{mL}$ , respectively. These were used as a starting concentration in the drug combination assay while the starting concentration of 3MA and wortmannin was used at 20 mM and 20  $\mu\text{M}$ , respectively. The results of parasite viability were represented as mean  $\pm$  SD. In the curcumin-chlorhexidine combination assay, at maximum concentrations of curcumin (MIC 250  $\mu\text{g}/\text{mL}$ ) and chlorhexidine (MIC 16  $\mu\text{g}/\text{mL}$ ), the percentage of trophozoites viability was in the range of 5–8%. Reduction concentration of chlorhexidine to 8  $\mu\text{g}/\text{mL}$  in combination with different concentrations of curcumin, the percentage was increased to the range of 42–53%, but their percentages were similar to those of chlorhexidine alone, at 52%. At lower concentrations of chlorhexidine (4, 2, 1  $\mu\text{g}/\text{mL}$ ), a pattern of the percentage viability at certain concentration of chlorhexidine was similar, and its percentage viability was gradually increased when the curcumin concentration was reduced (Table S1). In the curcumin-3MA (Table S2) and curcumin-wortmannin (Table S3) combinations, the pattern of results was similar. At certain 3MA or wortmannin concentrations below the MIC of curcumin, the percentage viability was gradually increased by reducing the concentration of curcumin. For the combinations that were close to our interest, 62.5  $\mu\text{g}/\text{mL}$  curcumin-1.25 mM 3MA or 1.25  $\mu\text{M}$  wortmannin, the percentage viability was comparable to curcumin alone.

### DISCUSSION

The cystic stage of *Acanthamoeba* is one of the main obstacles for therapeutic use as the penetration of anti-*Acanthamoeba* drugs across a double-layered cyst wall is fairly difficult (Turner et al., 2000; Abjani et al., 2016). The identification of new active compounds and drug repurposing with amoebicidal activity are urgently needed. In addition, the compound/drug that is able to prolong the trophozoite stage may be useful for drug combination purposes in the therapy of *Acanthamoeba* infection. In this study, an  $\text{IC}_{50}$  of curcumin against *A. triangularis* was identified. The killing activity of curcumin was confirmed, and interestingly, the surviving amoebas were arrested in the trophozoite stage after curcumin treatment at the sublethal dose. The dual benefits of the curcumin, amoebicidal activity, and arresting cyst transformation against *Acanthamoeba* sp. gain more attention. Regarding a long history of medicinal use of curcumin, it contains several pharmacological activities, for example, anti-inflammatory (Wal et al., 2019), antioxidant (Jakubczyk et al., 2020), anti-cancer (Vallianou et al., 2015; Tomeh, Hadianamrei & Zhao, 2019), and antimicrobial activities (Cui, Miao & Cui, 2007; Martins et al., 2009; Teow et al., 2016; Yang et al., 2016; Mitsuwan et al., 2020). For its anti-parasitic effect, the curcumin has been very-well studied in many parasites for example

*Schistosomiasis mansoni* (de Paula Aguiar et al., 2016; Hussein et al., 2017), *Besnoitia besnoiti* (Cervantes-Valencia et al., 2019), *Giardia lamblia* (Gutiérrez-Gutiérrez et al., 2017), *Leishmania major* (Koide et al., 2002), *Plasmodium falciparum* (Cui, Miao & Cui, 2007; Mishra et al., 2008), and *Trypanosoma cruzi* (Novaes et al., 2016). However, the killing mechanism by curcumin has been partially characterized in some parasites, but in *Acanthamoeba*, the documentation is largely unidentified.

Encystation refers to a mechanism in which amoeba trophozoites are transformed into cysts under stress conditions (Schaap & Schilde, 2018). In *Acanthamoeba*, several pathways, for example, actin dynamics, glycolysis, proteolysis (Bouyer et al., 2009), proteins such as cyst specific protein 21 (Chen et al., 2004), serine protease (Dudley, Alsam & Khan, 2008; Moon et al., 2008), cysteine protease (Leitsch et al., 2010; Moon et al., 2012), glycogen phosphorylase (Lorenzo-Morales et al., 2008), sirtuin proteins (Joo et al., 2020), and Shwachman-Bodian-Diamond syndrome protein (Wang, Lin & He, 2021) have been reported to be involved with this mechanism. However, the coordination and crosstalk between these pathways to support the encystation are still unknown. Single or multiple pathways may be required for cyst formation and the cyst induction probably depends on the strength and specificity of the cyst formation signal. Autophagy is an intracellular stress-sensing mechanism that occurs rapidly in response to stimuli such as rapamycin, starvation, or cytokines (Kamada, Sekito & Ohsumi, 2004; Kroemer, Mariño & Levine, 2010). So far, more than 30 autophagy-related (Atg) proteins have been identified in yeast and humans, and their roles in this pathway have been extensively studied (Kamada, Sekito & Ohsumi, 2004; Feng et al., 2014; Galluzzi et al., 2017). However, a partial list of Atg proteins has been characterized in *Acanthamoeba* i.e. Atg3 (Moon et al., 2011), Atg8 (Moon et al., 2009; Moon et al., 2013), Atg12 (Kim et al., 2015), Atg16 (Fujita et al., 2008) and they were reported to be associated with *Acanthamoeba* encystation.

Starvation or a nutrient-depleted condition is a classical autophagy inducer in several eukaryotic cells (Kamada, Sekito & Ohsumi, 2004; Mizushima et al., 2004; Mejlvang et al., 2018). In *Acanthamoeba*, starvation conditions are able to induce *Acanthamoeba* encystation at different degrees, depending on the medium formulation, time, and *Acanthamoeba* spp. (Aqeel et al., 2013; Sohn et al., 2017; Boonhok et al., 2021a). In our study, starvation by Page's Saline buffer (PAS) supplemented with 5% glucose was utilized. Approximately 40–50% of cysts were observed at 24 h after initiation of the culture, and the percentage of trophozoites containing enlarged vacuoles was increased significantly. Autophagy inhibitors, 3MA, and wortmannin, which are known to inhibit phosphatidylinositol 3-kinase (PI3K) activity in the autophagy pathway (Wu et al., 2010), have also been applied to see an autophagic response in *A. triangularis*. The 3MA significantly inhibited the formation of *A. triangularis* cysts while wortmannin was slightly affected. The different degrees of inhibition may result from the specificity of binding to its PI3K substrate, and the concentration used in the assay. In the presence of curcumin at the sublethal dose under starved conditions, most of the parasites remained in the trophozoite, not transforming into cyst stages. However, the mechanism of action of curcumin remains to be elucidated and needs further investigation. The curcumin may

bind to *Acanthamoeba* surface or intracellular proteins including cell cycle proteins and autophagy-related proteins which leads to the cell cycle arrest and inhibition of encystation, respectively. Basically, induction of autophagy, a double membrane autophagosome or vacuole is formed (Huang & Klionsky, 2002; Nakatogawa et al., 2009), and in *Acanthamoeba*, the formation of vacuoles including autophagosome and autolysosome is associated with the cyst wall formation (Bowers & Korn, 1969). The percentage of trophozoites containing vacuoles or enlarged vacuoles was thus analyzed by microscopy in our study. Due to a highly active trophozoite stage (Alves et al., 2017), analysis of the trophozoites containing vacuoles, almost 100% of the trophozoites contained vacuoles, and the percentage of trophozoites containing vacuoles regardless of the vacuole size made a difference between tested conditions. However, analyzing trophozoites containing enlarged vacuoles, the percentage was significantly reduced upon autophagy inhibitor or curcumin treatment. However, the combination of curcumin with autophagy inhibitor did not completely inhibit cysts formation. This may result from the dose of drug/compound tested in this study. Moreover, our drug combination data also demonstrated no synergistic, additive, or antagonistic effects in any drug combinations against *A. triangularis* trophozoites. This indicates that the outcome observed in our study is derived from a single drug. Regarding the effect of curcumin under microscopic examination which markedly inhibited cysts formation and reduced vacuolization in surviving trophozoites, molecular analysis of *A. triangularis* autophagy mRNA expression was performed to assess a physiological change, in response to the curcumin. Considering *Acanthamoeba* autophagy, Moon and his colleague first characterized Atg8 in *Acanthamoeba castellanii* (Moon et al., 2009). AcAtg8 was distributed in the amoeba cytosol, and its expression was peaked during encystation. In addition, intracellular colocalization of AcAtg8 and lysosome on the membrane has been demonstrated (Moon et al., 2009). An AcAtg8 isoform, AcAtg8b was later identified. This isoform was highly expressed during encystation and was required for *Acanthamoeba* encystation (Moon et al., 2013). Atg3, an E2 ubiquitin-like conjugating enzyme, is known to play a role in the Atg8 conjugation system (Feng et al., 2014). In *A. castellanii*, AcAtg3 was investigated by Moon and his colleagues and found that its mRNA expression was not increased during the encystation, but the depletion of AcAtg3 affected the maturation of cysts (Moon et al., 2011). Atg12 plays a role in autophagosome formation by forming an Atg12-Atg5-Atg16L1 complex and acting as an E3-like enzyme to promote Atg8 lipidation in the autophagosomal membrane (Yin, Pascual & Klionsky, 2016). At the early phase of encystation, *Acanthamoeba* Atg12 was consistently distributed in trophozoites. Later, it was formed as a puncta and co-located with an autophagic membrane. Even its mRNA expression was not increased during encystation as expected, but it was crucial for the encystation as the down-regulation of AcAtg12 in trophozoites inhibited cyst formation (Kim et al., 2015). *Acanthamoeba* Atg16 was partially colocalized with autophagolysosome and highly expressed during *A. castellanii* encystation (Song et al., 2012). Depletion of AcAtg16 inhibited the formation of autophagosomes and further disrupted the encystation mechanism (Song et al., 2012). As expected, all tested genes were at the basal level in 3MA-or curcumin-treated conditions. The inhibition of key ATG mRNA expression thus

supports the attenuation of *A. triangularis* encystation as well as cyst production. Autophagy is a tightly regulated pathway and its response depends on the strength and specificity of signals (Kroemer, Mariño & Levine, 2010; Simon et al., 2017), thus our data indicate that the signal strength of curcumin is higher than starvation signal and its underlying mechanism, curcumin may specifically interact with proteins associated with the inhibition of *Acanthamoeba* encystment or cell cycle arrest (Bínová, Bína & Nohtyfňnková, 2021).

Next, we investigated the effect of a single curcumin signal under a nutrient-rich or full condition using PYG medium. The cyst formation in response to curcumin was at the basal level similar to full medium alone. The percentage of surviving trophozoites with enlarged vacuoles was also at the basal level and was not different between curcumin-treated and untreated conditions. In addition, the real-time PCR analysis revealed that the tested ATG genes were similar to those of curcumin treatment under starved condition except AcATG16 that up-regulated at later time points. The increased expression of AcATG16 was also observed in *Peganum harmala* seed extract-treated *A. triangularis* (Boonhok et al., 2021a); however, in *Cassia angustifolia* extract treatment, the increase in AcATG16 mRNA was not observed (Boonhok et al., 2021b). This may indicate a role of Atg16 in *A. triangularis* responses to the specific stress signal in autophagy or other cellular pathways, which requires further investigations. Under the nutrient-rich condition, the mRNA expression of other *A. triangularis* encystation-related genes was investigated. Both cellulose synthase (EDCBI66TR) and serine proteinase (EU365404) were slightly changed in the first 18 h, to our surprise, at 24 h, their expression was significantly increased even the microscopic examination showed that there was no cyst induction at this time. Cellulose is the main component of cyst wall, and three enzymes namely, glycogen phosphorylase, UDP-glucose pyrophosphorylase, and cellulose synthase, are required for cellulose synthesis during *Acanthamoeba* encystation (Moon & Kong, 2012; Garajová et al., 2019). In addition to cellulose synthase, investigation on the mRNA expression of another two genes is required to predict cyst formation after 24 h conclusively; otherwise, this may indicate an additional function of the cellulose synthase. On the other hand, serine proteinase that increased at 24 h post curcumin treatment may indicate its role in other cellular activities in addition to cell differentiation (Blaschitz et al., 2006; Rascon & McKerrow, 2013). Moreover, we observed metacaspase, which is known to be involved in apoptosis-like cell death in several microorganisms and associated with *A. castellanii* encystation (Trzyna, Legras & Cordingley, 2008; Saheb, Trzyna & Bush, 2014), as well as interleukin-1 converting enzyme-like protease, known as caspase-1, has a role in programmed cell death of parasites (Kosec et al., 2006; Wu et al., 2018). The mRNA expression of metacaspase (AF480890) was consistent over the time period of curcumin treatment which may support no cyst formation. However, interleukin-1 converting enzyme-like protease (XM004338552) was a quick response to curcumin as its mRNA expression was immediately increased at 6 h post-treatment. However, at later time points, its expression was declined to the basal level. The increase of this gene at an early time point may indicate an apoptotic cell death by curcumin. However, to confirm this type of cell death, an apoptosis assay is required. Once the amoeba is able to cope with



the curcumin stress, the interleukin-1 converting enzyme-like protease expression is gradually declined, which reveals an ability of *A. triangularis* trophozoites to overcome the curcumin stress or a death signal.

In the line of curcumin effect on autophagy, curcumin is known to modulate autophagy (Shakeri et al., 2019), and the outcome is varied depending on cell type and curcumin concentration as described herein. In human endothelial cells, EA.hy926 and HUVECs, 5 or 20  $\mu\text{M}$  curcumin induced autophagy to reduce oxidative stress-induced cell damage (Han et al., 2012; Guo et al., 2016). An amount of 40  $\mu\text{M}$  of curcumin was able to induce autophagy which is partially involved with anticancer activity in human lung adenocarcinoma cell line, A549 (Liu et al., 2017). In human colon cancer cells, HCT116, 40  $\mu\text{M}$  curcumin-induced reactive oxygen species (ROS) production, which further activated autophagy followed by cell death (Liu et al., 2017). On the other hand, in mouse hippocampal neuronal cell line, HT-22, 10 or 15  $\mu\text{M}$  curcumin promoted cell recovery in  $\text{A}\beta\text{1-42}$ -treated condition by inhibiting autophagy (Zhang et al., 2018). At 5  $\mu\text{M}$  curcumin, it reduced apoptosis and inhibited autophagy and hypoxia-inducible factor 1-alpha in rat adrenal pheochromocytoma cell, PC12, model of oxygen-glucose deprivation/reperfusion (OGD/R) condition (Hou, Wang & Feng, 2019). Along with the OGD/R model, 10  $\mu\text{M}$  curcumin was able to increase the resistance of cortical neurons by reducing autophagy and cell apoptosis in an mTOR-dependent manner (Shi et al., 2019). Even autophagy is a quick response to various stimuli, but its mechanism is tightly regulated and be more selective in which Atg proteins work together in a specific manner and coordinate with other pathways or proteins to create a wide variety of physiological processes in cells (Wang & Qin, 2013; Galluzzi et al., 2017) and the autophagic response might be varied depending on the cell type and the dose of curcumin. Investigation of function and physiological change of *A. triangularis* Atg proteins in response to stresses, including the curcumin stress, is needed. Regarding the ability of curcumin in cell arrest, several studies have mentioned this pharmacological activity. Curcumin treatment caused cell cycle arrest at G1/S and G2/M phases and activated a caspase-3 pathway, resulting in human osteosarcoma (HOS) cell death (Lee, Lee & Kim, 2009). In human cervical carcinoma cells, SiHa cells, curcumin activated ROS production, apoptosis, autophagy, cell cycle arrest, and cellular senescence. These activities co-occurred with the upregulation of p53 and p21 proteins (Wang & Wu, 2020). In colon cancer cells, HT-29, curcumin-induced ROS production led to apoptotic cell death and cell cycle inhibition (Agarwal et al., 2018). The similar results were observed in another colon cancer cell line, MC38. The mechanism of action of curcumin was also partially characterized and shown to down-regulate several cell cycle proteins *i.e.* cyclin A2, cyclin E1, cell cycle dependent kinase 2 (CDK2), and transcription factor E2F1 (Li et al., 2022). Altogether, *Acanthamoeba* autophagy and/or cell cycle pathway may be involved in our finding.

Curcumin and curcumin derivatives have so far been extensively studied for therapeutic purposes, especially in parasitic infections (Din et al., 2016). A successful development of a new class of curcumin has been reported against *Trypanosoma cruzi* (Matiadis et al., 2021). In *Plasmodium* infection, several strategies have been developed to increase the effectiveness of curcumin for example nanotized curcumin (Ghosh et al., 2014), curcumin

containing liposomes (Martí Coma-Cros et al., 2018), among others. The strategies open another direction in drug development that could be applied in *Acanthamoeba* research. Moreover, drug combination strategy by targeting the autophagy pathway in other models has been reported (Zanotto-Filho et al., 2015), and this strategy may be applied in *Acanthamoeba* infection in the future. Taken together, evaluation of *Acanthamoeba* cyst formation and analysis of expression of autophagy-related genes or proteins in the surviving amoebas during drug or natural compound screening may help to assess the risk of *Acanthamoeba* encystation and can be a useful information for drug combination study to improve therapeutic efficacy and help reducing the drug resistance cycle in the area of infectious diseases (Hill & Cowen, 2015).

## CONCLUSIONS

Curcumin has a wide range of pharmacological activities and medicinal properties against numerous diseases. In *A. triangularis*, an amoebicidal activity of curcumin was recently demonstrated. Our study revealed that curcumin at sublethal dose is able to inhibit a transformation of *A. triangularis* trophozoites into cysts even under nutrient starvation. This may result from an attenuation of *Acanthamoeba* autophagy. However, an underlying mechanism of curcumin in *A. triangularis* trophozoites arrest is still unknown and needs further investigation. Overall, a dual benefit of curcumin, amoebicidal activity and arresting cyst transformation, may be another evidence to support drug development and future use of curcumin in *Acanthamoeba* infection therapy.

## ACKNOWLEDGEMENTS

We thank the Research Institute of Health Science (RIHS) staff at Walailak University.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This research was funded by Walailak University grant No. WU-IRG-63-073, The Royal Patronage of Her Royal Highness Princess Maha Chakri Sirindhorn-Botanical Garden of Walailak University, Nakhon Si Thammarat, under the project entitled: Medicinal Thai Native Plants against *Acanthamoeba triangularis* as a serious eye infection (WUBG 031-2565), Thailand. M.d.L.P. thanks to Project CICECO-Aveiro Institute of Materials, UIDB/50011/2020, UIDP/50011/2020, and LA/P0006/2020, financed by national funds through the FCT/MEC (PIDDAC). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Walailak University: WU-IRG-63-073 and WUBG 031-2565.

Project CICECO-Aveiro Institute of Materials: UIDB/50011/2020, UIDP/50011/2020, and LA/P0006/2020, FCT/MEC (PIDDAC).

## Competing Interests

The authors declare that they have no competing interests.

## Author Contributions

- Rachasak Boonhok conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, financial support, and approved the final draft.
- Suthinee Sangkanu conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Suganya Phumjan performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Ramita Jongboonjua performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Nawarat Sangnopparat performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Pattamaporn Kwankaew performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Aman Tedasen performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Chooi Ling Lim conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Maria de Lourdes Pereira conceived and designed the experiments, authored or reviewed drafts of the article, financial support, and approved the final draft.
- Mohammed Rahmatullah conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Polrat Wilairatana conceived and designed the experiments, authored or reviewed drafts of the article, financial support, and approved the final draft.
- Christophe Wiart conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Karma G. Dolma conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Alok K. Paul conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Madhu Gupta conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Veeranoot Nissapatorn conceived and designed the experiments, authored or reviewed drafts of the article, financial support, and approved the final draft.

## Data Availability

The following information was supplied regarding data availability:

The raw data are available in the [Supplemental File](#).

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.13657#supplemental-information>.

## REFERENCES

- Abjani F, Khan NA, Yousuf FA, Siddiqui R. 2016. Targeting cyst wall is an effective strategy in improving the efficacy of marketed contact lens disinfecting solutions against *Acanthamoeba castellanii* cysts. *Contact Lens and Anterior Eye* 39:239–243 DOI 10.1016/j.clae.2015.11.004.
- Agarwal A, Kasinathan A, Ganesan R, Balasubramanian A, Bhaskaran J, Suresh S, Srinivasan R, Aravind K, Sivalingam N. 2018. Curcumin induces apoptosis and cell cycle arrest via the activation of reactive oxygen species-independent mitochondrial apoptotic pathway in Smad4 and p53 mutated colon adenocarcinoma HT29 cells. *Nutrition Research* 51:67–81 DOI 10.1016/j.nutres.2017.12.011.
- Alves DDSMM, Alves LM, da Costa TL, de Castro AM, Vinaud MC. 2017. Anaerobic metabolism in T4 *Acanthamoeba* genotype. *Current Microbiology* 74:685–690 DOI 10.1016/j.nutres.2017.12.011.
- Anwar A, Khan NA, Siddiqui R. 2018. Combating *Acanthamoeba* spp. cysts: what are the options? *Parasites & Vectors* 11:26 DOI 10.1186/s13071-017-2572-z.
- Aqeel Y, Siddiqui R, Iftikhar H, Khan NA. 2013. The effect of different environmental conditions on the encystation of *Acanthamoeba castellanii* belonging to the T4 genotype. *Experimental Parasitology* 135:30–35 DOI 10.1016/j.exppara.2013.05.017.
- Bínová E, Bína D, Nohtyfťnková E. 2021. DNA content in *Acanthamoeba* during two stress defense reactions: encystation, pseudocyst formation and cell cycle. *European Journal of Protistology* 77:125745 DOI 10.1016/j.ejop.2020.125745.
- Blaschitz M, Köhler M, Aspöck H, Walochnik J. 2006. Detection of a serine proteinase gene in *Acanthamoeba* genotype T6 (Amoebozoa: Lobosea). *Experimental Parasitology* 114:26–33 DOI 10.1016/j.exppara.2006.02.004.
- Boonhok R, Sangkanu S, Chuprom J, Srisuphanunt M, Norouzi R, Siyadatpanah A, Mirzaei F, Mitsuan W, Wisessombat S, de Lourdes Pereira M. 2021a. *Peganum harmala* extract has antiamebic activity to *Acanthamoeba triangularis* trophozoites and changes expression of autophagy-related genes. *Pathogens* 10:842 DOI 10.3390/pathogens10070842.
- Boonhok R, Sangkanu S, Norouzi R, Siyadatpanah A, Mirzaei F, Mitsuan W, Charong N, Wisessombat S, de Lourdes Pereira M, Rahmatullah M. 2021b. Amoebicidal activity of *Cassia angustifolia* extract and its effect on *Acanthamoeba triangularis* autophagy-related gene expression at transcriptional level. *Parasitology* 148(9):1–36 DOI 10.1017/S0031182021000718.
- Bouyer S, Rodier M-H, Guillot A, Héchard Y. 2009. *Acanthamoeba castellanii*: proteins involved in actin dynamics, glycolysis, and proteolysis are regulated during encystation. *Experimental Parasitology* 123:90–94 DOI 10.1016/j.exppara.2009.06.006.
- Bowers B, Korn ED. 1969. The fine structure of *Acanthamoeba castellanii* (Neff strain) II encystment. *The Journal of Cell Biology* 41:786–805 DOI 10.1083/jcb.41.3.786.
- Brinck Andersen N-S, Jorgensen SE, Skipper KA, Larsen SM, Heinz J, Thomsen MM, Farahani E, Cai Y, Hait AS, Kay L. 2020. Essential role of autophagy in restricting poliovirus infection revealed by identification of an ATG7 defect in a poliomyelitis patient. *Autophagy* 17(9):2449–2464 DOI 10.1080/15548627.2020.1831800.
- Castillo EF, Dekonenko A, Arko-Mensah J, Mandell MA, Dupont N, Jiang S, Delgado-Vargas M, Timmins GS, Bhattacharya D, Yang H. 2012. Autophagy protects against

- active tuberculosis by suppressing bacterial burden and inflammation. *Proceedings of the National Academy of Sciences of the United States of America* **109**:E3168–E3176 DOI [10.1073/pnas.1210500109](https://doi.org/10.1073/pnas.1210500109).
- Cervantes-Valencia ME, Hermosilla C, Alcalá-Canto Y, Tapia G, Taubert A, Silva LM. 2019.** Antiparasitic efficacy of curcumin against *Besnoitia besnoiti* tachyzoites *in vitro*. *Frontiers in Veterinary Science* **5**:333 DOI [10.3389/fvets.2018.00333](https://doi.org/10.3389/fvets.2018.00333).
- Chelkha N, Jardot P, Moussaoui I, Levasseur A, La Scola B, Colson P. 2020.** Core gene-based molecular detection and identification of *Acanthamoeba* species. *Scientific Reports* **10**:1–9 DOI [10.1038/s41598-020-57998-5](https://doi.org/10.1038/s41598-020-57998-5).
- Chen L, Orfeo T, Gilmartin G, Bateman E. 2004.** Mechanism of cyst specific protein 21 mRNA induction during *Acanthamoeba* differentiation. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research* **1691**:23–31 DOI [10.1016/j.bbamcr.2003.11.005](https://doi.org/10.1016/j.bbamcr.2003.11.005).
- Cui L, Miao J, Cui L. 2007.** Cytotoxic effect of curcumin on malaria parasite *Plasmodium falciparum*: inhibition of histone acetylation and generation of reactive oxygen species. *Antimicrobial Agents and Chemotherapy* **51**:488–494 DOI [10.1128/AAC.01238-06](https://doi.org/10.1128/AAC.01238-06).
- de Lacerda AG, Lira M. 2021.** *Acanthamoeba* keratitis: a review of biology, pathophysiology and epidemiology. *Ophthalmic & Physiological Optics* **41**:116–135 DOI [10.1111/opo.12752](https://doi.org/10.1111/opo.12752).
- de Paula Aguiar D, Brunetto Moreira Moscardini M, Rezende Morais E, Graciano de Paula R, Ferreira PM, Afonso A, Belo S, Tomie Ouchida A, Curti C, Cunha WR. 2016.** Curcumin generates oxidative stress and induces apoptosis in adult *Schistosoma mansoni* worms. *PLOS ONE* **11**:e0167135 DOI [10.1371/journal.pone.0167135](https://doi.org/10.1371/journal.pone.0167135).
- Díaz-Troya S, Pérez-Pérez ME, Florencio FJ, Crespo JL. 2008.** The role of TOR in autophagy regulation from yeast to plants and mammals. *Autophagy* **4**:851–865 DOI [10.4161/auto.6555](https://doi.org/10.4161/auto.6555).
- Din ZU, dos Santos A, Trapp MA, Lazarin-Bidóia D, Garcia FP, Peron F, Nakamura CV, Rodrigues-Filho E. 2016.** Curcumin inspired synthesis of unsymmetrical diarylpentanoids with highly potent anti-parasitic activities: *in silico* studies and DFT-based stereochemical calculation. *Medicinal Chemistry Communication* **7**:820–831 DOI [10.1039/C5MD00599J](https://doi.org/10.1039/C5MD00599J).
- Dudley R, Alsam S, Khan NA. 2008.** The role of proteases in the differentiation of *Acanthamoeba castellanii*. *FEMS Microbiology Letters* **286**:9–15 DOI [10.1111/j.1574-6968.2008.01249.x](https://doi.org/10.1111/j.1574-6968.2008.01249.x).
- Elsheikha HM, Siddiqui R, Khan NA. 2020.** Drug discovery against *Acanthamoeba* infections: present knowledge and unmet needs. *Pathogens* **9**:405 DOI [10.3390/pathogens9050405](https://doi.org/10.3390/pathogens9050405).
- Eskelinen E-L. 2005.** Maturation of autophagic vacuoles in mammalian cells. *Autophagy* **1**(1):1–10 DOI [10.4161/auto.1.1.1270](https://doi.org/10.4161/auto.1.1.1270).
- Feng Y, He D, Yao Z, Klionsky DJ. 2014.** The machinery of macroautophagy. *Cell Research* **24**:24–41 DOI [10.1038/cr.2013.168](https://doi.org/10.1038/cr.2013.168).
- Fujita N, Itoh T, Omori H, Fukuda M, Noda T, Yoshimori T. 2008.** The Atg16L complex specifies the site of LC3 lipidation for membrane biogenesis in autophagy. *Molecular Biology of the Cell* **19**:2092–2100 DOI [10.1091/mbc.e07-12-1257](https://doi.org/10.1091/mbc.e07-12-1257).
- Galluzzi L, Baehrecke EH, Ballabio A, Boya P, Bravo-San Pedro JM, Cecconi F, Choi AM, Chu CT, Codogno P, Colombo MI. 2017.** Molecular definitions of autophagy and related processes. *The EMBO Journal* **36**(13):1811–1836 DOI [10.15252/embj.201796697](https://doi.org/10.15252/embj.201796697).
- Garajová M, Mrva M, Vaškovicová N, Martinka M, Melicherová J, Valigurová A. 2019.** Cellulose fibrils formation and organisation of cytoskeleton during encystment are essential for *Acanthamoeba* cyst wall architecture. *Scientific Reports* **9**(1):1–21 DOI [10.1038/s41598-019-41084-6](https://doi.org/10.1038/s41598-019-41084-6).

- Ghosh A, Banerjee T, Bhandary S, Surolia A. 2014.** Formulation of nanotized curcumin and demonstration of its antimalarial efficacy. *International Journal of Nanomedicine* **9**(1):5373 DOI [10.2147/IJN.S62756](https://doi.org/10.2147/IJN.S62756).
- Guimaraes AJ, Gomes KX, Cortines JR, Peralta JM, Peralta RHS. 2016.** *Acanthamoeba* spp. as a universal host for pathogenic microorganisms: one bridge from environment to host virulence. *Microbiological Research* **193**:30–38 DOI [10.1016/j.micres.2016.08.001](https://doi.org/10.1016/j.micres.2016.08.001).
- Guo S, Long M, Li X, Zhu S, Zhang M, Yang Z. 2016.** Curcumin activates autophagy and attenuates oxidative damage in EA. hy926 cells via the Akt/mTOR pathway. *Molecular Medicine Reports* **13**:2187–2193 DOI [10.3892/mmr.2016.4796](https://doi.org/10.3892/mmr.2016.4796).
- Gutiérrez-Gutiérrez F, Palomo-Ligas L, Hernández-Hernández JM, Pérez-Rangel A, Aguayo-Ortiz R, Hernández-Campos A, Castillo R, González-Pozos S, Cortés-Zárate R, Ramírez-Herrera MA. 2017.** Curcumin alters the cytoskeleton and microtubule organization on trophozoites of *Giardia lamblia*. *Acta Tropica* **172**:113–121 DOI [10.1016/j.actatropica.2017.04.027](https://doi.org/10.1016/j.actatropica.2017.04.027).
- Han J, Pan X-Y, Xu Y, Xiao Y, An Y, Tie L, Pan Y, Li X-J. 2012.** Curcumin induces autophagy to protect vascular endothelial cell survival from oxidative stress damage. *Autophagy* **8**(5):812–825 DOI [10.4161/auto.19471](https://doi.org/10.4161/auto.19471).
- Hill JA, Cowen LE. 2015.** Using combination therapy to thwart drug resistance. *Future Microbiology* **10**(11):1719–1726 DOI [10.2217/fmb.15.68](https://doi.org/10.2217/fmb.15.68).
- Hou Y, Wang J, Feng J. 2019.** The neuroprotective effects of curcumin are associated with the regulation of the reciprocal function between autophagy and HIF-1 $\alpha$  in cerebral ischemia-reperfusion injury. *Drug Design, Development and Therapy* **13**:1135 DOI [10.2147/DDDT.S194182](https://doi.org/10.2147/DDDT.S194182).
- Huang W-P, Klionsky DJ. 2002.** Autophagy in yeast: a review of the molecular machinery. *Cell Structure and Function* **27**:409–420 DOI [10.1247/csf.27.409](https://doi.org/10.1247/csf.27.409).
- Hussain RHM, Isa NSM, Kamaruddin KA, Ghani MKA, Khan NA, Siddiqui R, Anuar TS. 2020.** Morphological and molecular characterization of *Acanthamoeba* isolated from contact lens paraphernalia in Malaysia: highlighting the pathogenic potential of T4 genotype. *Asian Pacific Journal of Tropical Medicine* **13**(12):542 DOI [10.4103/1995-7645.296722](https://doi.org/10.4103/1995-7645.296722).
- Hussein A, Rashed S, El Hayawan I, El-Sayed R, Ali H. 2017.** Evaluation of the anti-schistosomal effects of turmeric (*Curcuma longa*) versus praziquantel in *Schistosoma mansoni* infected mice. *Iranian Journal of Parasitology* **12**(4):587–596.
- Jakubczyk K, Druźga A, Katarzyna J, Skonieczna-Żydecka K. 2020.** Antioxidant potential of curcumin—a meta-analysis of randomized clinical trials. *Antioxidants* **9**(11):1092 DOI [10.3390/antiox9111092](https://doi.org/10.3390/antiox9111092).
- Joo S-Y, Aung JM, Shin M, Moon E-K, Kong H-H, Goo Y-K, Chung D-I, Hong Y. 2020.** The role of the *Acanthamoeba castellanii* Sir2-like protein in the growth and encystation of *Acanthamoeba*. *Parasites & Vectors* **13**(1):1–14 DOI [10.1186/s13071-020-04237-5](https://doi.org/10.1186/s13071-020-04237-5).
- Juarez MM, Tártara LI, Cid AG, Real JP, Bermúdez JM, Rajal VB, Palma SD. 2018.** *Acanthamoeba* in the eye, can the parasite hide even more? Latest developments on the disease. *Contact Lens and Anterior Eye* **41**(3):245–251 DOI [10.1016/j.clae.2017.12.017](https://doi.org/10.1016/j.clae.2017.12.017).
- Kamada Y, Sekito T, Ohsumi Y. 2004.** Autophagy in yeast: ATOR-mediated response to nutrient starvation. In: *TOR. Current Topics in Microbiology and Immunology*. Berlin, Heidelberg: Springer, 73–84.
- Khambu B, Yan S, Huda N, Liu G, Yin X-M. 2018.** Autophagy in non-alcoholic fatty liver disease and alcoholic liver disease. *Liver Research* **2**:112–119 DOI [10.1016/j.livres.2018.09.004](https://doi.org/10.1016/j.livres.2018.09.004).

- Khan NA, Anwar A, Siddiqui R. 2019.** *Acanthamoeba* keratitis: current status and urgent research priorities. *Current Medicinal Chemistry* **26**:5711–5726  
DOI [10.2174/0929867325666180510125633](https://doi.org/10.2174/0929867325666180510125633).
- Kim S-H, Moon E-K, Hong Y, Chung D-I, Kong H-H. 2015.** Autophagy protein 12 plays an essential role in *Acanthamoeba* encystation. *Experimental Parasitology* **159**:46–52  
DOI [10.1016/j.exppara.2015.08.013](https://doi.org/10.1016/j.exppara.2015.08.013).
- Kim SY, Syms MJ, Holtel MR, Nauschuetz KK. 2000.** *Acanthamoeba* sinusitis with subsequent dissemination in an AIDS patient. *Ear, Nose & Throat Journal* **79**:168–174  
DOI [10.1177/014556130007900310](https://doi.org/10.1177/014556130007900310).
- Kocaadam B, Şanlıer N. 2017.** Curcumin, an active component of turmeric (*Curcuma longa*), and its effects on health. *Critical Reviews in Food Science and Nutrition* **57**(13):2889–2895  
DOI [10.1080/10408398.2015.1077195](https://doi.org/10.1080/10408398.2015.1077195).
- Koide T, Nose M, Ogihara Y, Yabu Y, Ohta N. 2002.** Leishmanicidal effect of curcumin *in vitro*. *Biological and Pharmaceutical Bulletin* **25**(1):131–133 DOI [10.1248/bpb.25.131](https://doi.org/10.1248/bpb.25.131).
- Kosec G, Alvarez VE, Agüero F, Sánchez D, Dolinar M, Turk B, Turk V, Cazzulo JJ. 2006.** Metacaspases of *Trypanosoma cruzi*: possible candidates for programmed cell death mediators. *Molecular and Biochemical Parasitology* **145**(1):18–28 DOI [10.1016/j.molbiopara.2005.09.001](https://doi.org/10.1016/j.molbiopara.2005.09.001).
- Kroemer G, Mariño G, Levine B. 2010.** Autophagy and the integrated stress response. *Molecular cell* **40**:280–293 DOI [10.1016/j.molcel.2010.09.023](https://doi.org/10.1016/j.molcel.2010.09.023).
- Lee DS, Lee MK, Kim JH. 2009.** Curcumin induces cell cycle arrest and apoptosis in human osteosarcoma (HOS) cells. *Anticancer Research* **29**:5039–5044.
- Leitsch D, Köhler M, Marchetti-Deschmann M, Deutsch A, Allmaier G, Duchêne M, Walochnik J. 2010.** Major role for cysteine proteases during the early phase of *Acanthamoeba castellanii* encystment. *Eukaryotic Cell* **9**:611–618 DOI [10.1128/EC.00300-09](https://doi.org/10.1128/EC.00300-09).
- Li P, Pu S, Lin C, He L, Zhao H, Yang C, Guo Z, Xu S, Zhou Z. 2022.** Curcumin selectively induces colon cancer cell apoptosis and S cell cycle arrest by regulates Rb/E2F/p53 pathway. *Journal of Molecular Structure* **1263**:133180 DOI [10.1016/j.molstruc.2022.133180](https://doi.org/10.1016/j.molstruc.2022.133180).
- Liu F, Gao S, Yang Y, Zhao X, Fan Y, Ma W, Yang D, Yang A, Yu Y. 2017.** Curcumin induced autophagy anticancer effects on human lung adenocarcinoma cell line A549. *Oncology Letters* **14**:2775–2782 DOI [10.3892/ol.2017.6565](https://doi.org/10.3892/ol.2017.6565).
- Lorenzo-Morales J, Khan NA, Walochnik J. 2015.** An update on *Acanthamoeba* keratitis: diagnosis, pathogenesis and treatment. *Parasite* **22**:10 DOI [10.1051/parasite/2015010](https://doi.org/10.1051/parasite/2015010).
- Lorenzo-Morales J, Kliessikova J, Martinez-Carretero E, De Pablos LM, Profotova B, Nohynkova E, Osuna A, Valladares B. 2008.** Glycogen phosphorylase in *Acanthamoeba* spp.: determining the role of the enzyme during the encystment process using RNA interference. *Eukaryotic Cell* **7**:509–517 DOI [10.1128/EC.00316-07](https://doi.org/10.1128/EC.00316-07).
- Martins C, Da Silva D, Neres A, Magalhaes T, Watanabe G, Modolo L, Sabino A, De Fátima A, De Resende M. 2009.** Curcumin as a promising antifungal of clinical interest. *Journal of Antimicrobial Chemotherapy* **63**:337–339 DOI [10.1093/jac/dkn488](https://doi.org/10.1093/jac/dkn488).
- Martí Coma-Cros E, Biosca A, Lantero E, Manca ML, Caddeo C, Gutiérrez L, Ramírez M, Borgheti-Cardoso LN, Manconi M, Fernández-Busquets X. 2018.** Antimalarial activity of orally administered curcumin incorporated in Eudragit®-containing liposomes. *International Journal of Molecular Sciences* **19**:1361 DOI [10.3390/ijms19051361](https://doi.org/10.3390/ijms19051361).
- Matiadis D, Saporiti T, Aguilera E, Robert X, Guillon C, Cabrera N, Pérez-Montfort R, Sagnou M, Alvarez G. 2021.** Pyrazol (in) e derivatives of curcumin analogs as a new class of anti-*Trypanosoma cruzi* agents. *Future Medicinal Chemistry* **13**:701–714  
DOI [10.4155/fmc-2020-0349](https://doi.org/10.4155/fmc-2020-0349).

- Matson DO, Rouah E, Lee RT, Armstrong D, Parke JT, Baker CJ. 1988.** *Acanthamoeba* meningoencephalitis masquerading as neurocysticercosis. *The Pediatric Infectious Disease Journal* 7:121–124 DOI 10.1097/00006454-198802000-00011.
- Mejlvang J, Olsvik H, Svenning S, Bruun J-A, Abudu YP, Larsen KB, Brech A, Hansen TE, Brenne H, Hansen T. 2018.** Starvation induces rapid degradation of selective autophagy receptors by endosomal microautophagy. *Journal of Cell Biology* 217:3640–3655 DOI 10.1083/jcb.201711002.
- Menzies FM, Fleming A, Rubinsztein DC. 2015.** Compromised autophagy and neurodegenerative diseases. *Nature Reviews Neuroscience* 16:345–357 DOI 10.1038/nrn3961.
- Mishra S, Karmodiya K, Surolia N, Surolia A. 2008.** Synthesis and exploration of novel curcumin analogues as anti-malarial agents. *Bioorganic & Medicinal Chemistry* 16:2894–2902 DOI 10.1016/j.bmc.2007.12.054.
- Mitsuwan W, Bunsuwansakul C, Leonard TE, Laohaprapanon S, Hounkong K, Bunluepuech K, Chalernpol K, Mahboob T, Sumudi Raju C, Dhobi M. 2020.** *Curcuma longa* ethanol extract and Curcumin inhibit the growth of *Acanthamoeba triangularis* trophozoites and cysts isolated from water reservoirs at Walailak University, Thailand. *Pathogens and Global Health* 114:1–11 DOI 10.1080/20477724.2020.1755551.
- Mizushima N, Yamamoto A, Matsui M, Yoshimori T, Ohsumi Y. 2004.** *In vivo* analysis of autophagy in response to nutrient starvation using transgenic mice expressing a fluorescent autophagosome marker. *Molecular Biology of the Cell* 15:1101–1111 DOI 10.1091/mbc.e03-09-0704.
- Moon E-K, Chung D-I, Hong Y-C, Kong H-H. 2008.** Characterization of a serine proteinase mediating encystation of *Acanthamoeba*. *Eukaryotic Cell* 7:1513–1517 DOI 10.1128/EC.00068-08.
- Moon E-K, Chung D-I, Hong Y-C, Kong H-H. 2009.** Autophagy protein 8 mediating autophagosome in encysting *Acanthamoeba*. *Molecular and Biochemical Parasitology* 168:43–48 DOI 10.1016/j.molbiopara.2009.06.005.
- Moon E-K, Chung D-I, Hong Y, Kong H-H. 2011.** Atg3-mediated lipidation of Atg8 is involved in encystation of *Acanthamoeba*. *The Korean Journal of Parasitology* 49:103 DOI 10.3347/kjp.2011.49.2.103.
- Moon E-K, Chung D-I, Hong Y, Kong H-H. 2012.** Protein kinase C signaling molecules regulate encystation of *Acanthamoeba*. *Experimental Parasitology* 132:524–529 DOI 10.1016/j.exppara.2012.07.008.
- Moon E-K, Hong Y, Chung D-I, Kong H-H. 2013.** Identification of atg8 isoform in encysting *Acanthamoeba*. *The Korean Journal of Parasitology* 51:497 DOI 10.3347/kjp.2013.51.5.497.
- Moon E-K, Kong H-H. 2012.** Short-cut pathway to synthesize cellulose of encysting *Acanthamoeba*. *The Korean Journal of Parasitology* 50:361 DOI 10.3347/kjp.2012.50.4.361.
- Morrison AO, Morris R, Shannon A, Lauer SR, Guarner J, Kraft CS. 2016.** Disseminated *Acanthamoeba* infection presenting with cutaneous lesions in an immunocompromised patient: a case report, review of histomorphologic findings, and potential diagnostic pitfalls. *American Journal of Clinical Pathology* 145:266–270 DOI 10.1093/ajcp/aqv081.
- Nakatogawa H, Suzuki K, Kamada Y, Ohsumi Y. 2009.** Dynamics and diversity in autophagy mechanisms: lessons from yeast. *Nature Reviews Molecular Cell Biology* 10:458–467 DOI 10.1038/nrm2708.
- Neelam S, Niederkorn JY. 2017.** Pathobiology and immunobiology of *Acanthamoeba keratitis*: Insights from animal models. *Yale Journal of Biology and Medicine* 90(2):261–268.



- Novaes RD, Sartini MVP, Rodrigues JPF, Gonçalves RV, Santos EC, Souza RLM, Caldas IS. 2016. Curcumin enhances the anti-*Trypanosoma cruzi* activity of benzimidazole-based chemotherapy in acute experimental Chagas disease. *Antimicrobial Agents and Chemotherapy* 60:3355–3364 DOI 10.1128/AAC.00343-16.
- Picazarri K, Nakada-Tsukui K, Nozaki T. 2008. Autophagy during proliferation and encystation in the protozoan parasite *Entamoeba invadens*. *Infection and Immunity* 76:278–288 DOI 10.1128/IAI.00636-07.
- Rascon AA, McKerrow JH. 2013. Synthetic and natural protease inhibitors provide insights into parasite development, virulence and pathogenesis. *Current Medicinal Chemistry* 20:3078–3102 DOI 10.2174/0929867311320250005.
- Saheb E, Trzyna W, Bush J. 2014. Caspase-like proteins: *Acanthamoeba castellanii* metacaspase and *Dictyostelium discoideum* paracaspase, What are their functions? *Journal of Biosciences* 39:909–916 DOI 10.1007/s12038-014-9486-0.
- Schaap P, Schilde C. 2018. Encystation: the most prevalent and underinvestigated differentiation pathway of eukaryotes. *Microbiology* 164:727–739 DOI 10.1099/mic.0.000653.
- Shakeri A, Cicero AF, Panahi Y, Mohajeri M, Sahebkar A. 2019. Curcumin: a naturally occurring autophagy modulator. *Journal of Cellular Physiology* 234:5643–5654 DOI 10.1002/jcp.27404.
- Shi Q, Zhang Q, Peng Y, Zhang X, Wang Y, Shi L. 2019. A natural diarylheptanoid protects cortical neurons against oxygen-glucose deprivation-induced autophagy and apoptosis. *Journal of Pharmacy and Pharmacology* 71:1110–1118 DOI 10.1111/jphp.13096.
- Siddiqui R, Khan NA. 2012. Biology and pathogenesis of *Acanthamoeba*. *Parasites & Vectors* 5:6 DOI 10.1186/1756-3305-5-6.
- Simon H-U, Friis R, Tait SW, Ryan KM. 2017. Retrograde signaling from autophagy modulates stress responses. *Science Signaling* 10:eaag2791 DOI 10.1126/scisignal.aag2791.
- Sohn H-J, Kang H, Seo G-E, Kim J-H, Jung S-Y, Shin H-J. 2017. Efficient liquid media for encystation of pathogenic free-living amoebae. *The Korean Journal of Parasitology* 55:233 DOI 10.3347/kjp.2017.55.3.233.
- Song S-M, Han B-I, Moon E-K, Lee Y-R, Yu HS, Jha BK, Danne D-BS, Kong H-H, Chung D-I, Hong Y. 2012. Autophagy protein 16-mediated autophagy is required for the encystation of *Acanthamoeba castellanii*. *Molecular and Biochemical Parasitology* 183:158–165 DOI 10.1016/j.molbiopara.2012.02.013.
- Suzuki K. 2013. Selective autophagy in budding yeast. *Cell Death & Differentiation* 20:43–48 DOI 10.1038/cdd.2012.73.
- Taravaud A, Loiseau PM, Pomel S. 2017. *In vitro* evaluation of antimicrobial agents on *Acanthamoeba* sp. and evidence of a natural resilience to amphotericin B. *The International Journal for Parasitology: Drugs and Drug Resistance* 7:328–336 DOI 10.1016/j.ijpddr.2017.09.002.
- Teow S-Y, Liew K, Ali SA, Khoo AS-B, Peh S-C. 2016. Antibacterial action of curcumin against *Staphylococcus aureus*: a brief review. *Journal of Tropical Medicine* 2016:2853045 DOI 10.1155/2016/2853045.
- Tomeh MA, Hadianamrei R, Zhao X. 2019. A review of curcumin and its derivatives as anticancer agents. *International Journal of Molecular Sciences* 20:1033 DOI 10.3390/ijms20051033.
- Trzyna WC, Legras XD, Cordingley JS. 2008. A type-1 metacaspase from *Acanthamoeba castellanii*. *Microbiological Research* 163:414–423 DOI 10.1016/j.micres.2006.06.017.
- Turner N, Russell A, Furr J, Lloyd D. 2000. Emergence of resistance to biocides during differentiation of *Acanthamoeba castellanii*. *Journal of Antimicrobial Chemotherapy* 46:27–34 DOI 10.1093/jac/46.1.27.

- Vallianou NG, Evangelopoulos A, Schizas N, Kazazis C. 2015. Potential anticancer properties and mechanisms of action of curcumin. *Anticancer Research* 35(2):645–651.
- Wal P, Saraswat N, Pal RS, Wal A, Chaubey M. 2019. A detailed insight of the anti-inflammatory effects of curcumin with the assessment of parameters, sources of ROS and associated mechanisms. *Open Medicine Journal* 6:64–76 DOI 10.2174/1874220301906010064.
- Wang Y-J, Lin W-C, He M-S. 2021. The *Acanthamoeba* SBDS, a cytoskeleton-associated gene, is highly expressed during phagocytosis and encystation. *Journal of Microbiology, Immunology and Infection* 54:482–489 DOI 10.1016/j.jmii.2019.11.003.
- Wang Y, Qin Z-H. 2013. Coordination of autophagy with other cellular activities. *Acta Pharmacologica Sinica* 34:585–594 DOI 10.1007/978-981-15-0602-4\_30.
- Wang T, Wu X. 2020. Curcumin induces G2/M arrest and triggers autophagy, ROS generation and cell senescence in cervical cancer cells. *Journal of Cancer* 11:6704 DOI 10.7150/jca.45176.
- Wu D, Qiao K, Feng M, Fu Y, Cai J, Deng Y, Tachibana H, Cheng X. 2018. Apoptosis of *Acanthamoeba castellanii* trophozoites induced by oleic acid. *Journal of Eukaryotic Microbiology* 65:191–199 DOI 10.1111/jeu.12454.
- Wu Y-T, Tan H-L, Shui G, Bauvy C, Huang Q, Wenk MR, Ong C-N, Codogno P, Shen H-M. 2010. Dual role of 3-methyladenine in modulation of autophagy via different temporal patterns of inhibition on class I and III phosphoinositide 3-kinase. *Journal of Biological Chemistry* 285(14):10850–10861 DOI 10.1074/jbc.M109.080796.
- Yang M, Lee G, Si J, Lee S-J, You HJ, Ko G. 2016. Curcumin shows antiviral properties against norovirus. *Molecules* 21(10):1401 DOI 10.3390/molecules21101401.
- Yin Z, Pascual C, Klionsky DJ. 2016. Autophagy: machinery and regulation. *Microbial Cell* 3(12):588–596 DOI 10.15698/mic2016.12.546.
- Yorimitsu T, Klionsky DJ. 2005. Autophagy: molecular machinery for self-eating. *Cell Death and Differentiation* 12(S2):1542–1552 DOI 10.1038/sj.cdd.4401765.
- Zanotto-Filho A, Braganhol E, Klafke K, Figueiró F, Terra SR, Paludo FJ, Morrone M, Bristot IJ, Battastini AM, Forcelini CM. 2015. Autophagy inhibition improves the efficacy of curcumin/temozolomide combination therapy in glioblastomas. *Cancer Letters* 358(2):220–231 DOI 10.1016/j.canlet.2014.12.044.
- Zhang L, Fang Y, Cheng X, Lian Y, Zeng Z, Wu C, Zhu H, Xu H. 2018. The potential protective effect of curcumin on amyloid- $\beta$ -42 induced cytotoxicity in HT-22 cells. *BioMed Research International* 2018(21):1–8 DOI 10.1155/2018/8134902.