

Biological age for chronic kidney disease patients using index model

Shaiful Anuar Abu Bakar ^{Corresp., Equal first author, 1}, **Sharifah Nazatul Shima Syed Mohamed Shahrudin** ^{Equal first author, 1}, **Noriszura Ismail** ², **Wan Ahmad Hafiz Wan Md Adnan** ³

¹ Institute of Mathematical Sciences, Universiti Malaya, Kuala Lumpur, Kuala Lumpur, Malaysia

² Department of Mathematical Sciences, Universiti Kebangsaan Malaysia, Bangi, SELANGOR, Malaysia

³ Department of Medicine, Universiti Malaya, Kuala Lumpur, Kuala Lumpur, Malaysia

Corresponding Author: Shaiful Anuar Abu Bakar

Email address: saab@um.edu.my

Estimation of biological age (BA) facilitates an advantage of embedding information on the health being in addition to an individual lifetime. Factors that contribute towards profiling the human biological age can be diverse. Therefore, this study focuses on developing a BA model for patients with Chronic Kidney Disease (CKD). The procedure commences with the selection of significant biomarkers using a correlation test. Appropriate weightage is then assigned to each selected biomarker using the indexing method to produce a BA index. The BA index is matched to age variation within the sample to acquire additional terms for the chronological age leading ultimately to the estimated BA. From a sample of 190 patients (133 trained data and 57 testing data) obtained from the University of Malaya Medical Centre (UMMC), Malaysia, the intensity of the BA is found to be between 3 to 9 years from the chronological age. Visual observations further validate the high similarities between the training and testing data sets.

Biological age for chronic kidney disease patients using index model

Shaiful Anuar Abu Bakar¹, Sharifah Nazatul Shima Syed Mohamed Shahrudin¹, Noriszura Ismail², Wan Ahmad Hafiz Wan Md Adnan³

¹ Institute of Mathematical Sciences, Faculty of Science, Universiti Malaya, Malaysia.

² Department of Mathematical Sciences, Universiti Kebangsaan Malaysia, Malaysia.

³ Department of Medicine, Faculty of Medicine, Universiti Malaya, Malaysia.

Corresponding Author:

Shaiful Anuar Abu Bakar¹

Universiti Malaya, 50603 Kuala Lumpur, Malaysia

Email address: saab@um.edu.my

Abstract

Estimation of biological age (BA) facilitates an advantage of embedding information on the health being in addition to an individual lifetime. Factors that contribute towards profiling the human biological age can be diverse. Therefore, this study focuses on developing a BA model for patients with Chronic Kidney Disease (CKD). The procedure commences with the selection of significant biomarkers using a correlation test. Appropriate weightage is then assigned to each selected biomarker using the indexing method to produce a BA index. The BA index is matched to age variation within the sample to acquire additional terms for the chronological age leading ultimately to the estimated BA. From a sample of 190 patients (133 trained data and 57 testing data) obtained from the University of Malaya Medical Centre (UMMC), Malaysia, the magnitude of the difference for the BA is found to be between 3 to 9 years from the chronological age. Visual observations further validate the high similarities between the training and testing data sets.

Introduction

Estimation of biological age (BA) is becoming increasingly popular as an asymptotic measure to understand changes in physical functionality as well as the ageing process of a living being. Over the chronological age (CA), which exhibits an exact figure signifying the period between birth and present time of an individual, the BA is widely used to indicate the healthy and unhealthy ageing through variables that contribute to healthspan.¹ Furthermore, BA shows the health state of each individual and serves as a comparative measure between individuals of the same age and gender.² Thus, it describes one lifetime behaviour informatively provided the common premise that physical functionality decline is parallel to deterioration of health condition and age increment.

Unhealthy individuals demonstrate only decreased function in CA, but functional BA is intended to represent different stages of ageing.³ Group of individuals that perceived ill-health exhibited higher biological ages compared to the healthier groups. In this respect, those with more significant functional biological age have a higher chance of death because they reached a more advanced ageing stage earlier than others. Thus, the traditional approach in measuring individual health according to CA is less appealing in today's highly dynamic and rapidly changing global lifestyle.

Researchers have recently employed several statistical techniques to develop the BA model, primarily by incorporating multiple relevant biomarkers into the model. The BA model does not only predict ageing-related diseases but also considers the functional status during ageing.⁴ Several articles have been published on the measurement of BA using statistical methods. Multiple Linear Regression (MLR) remains one of the most widely used methods for calculating BA.⁵⁻¹³ Nevertheless, MLR has been criticized for the multicollinearity risk besides the potential for estimates to regress toward the mean.⁶ These suggest the MLR equation underestimates the individual BA in the older age while overestimating the younger age.¹² Principal Component Analysis (PCA) was proposed to overcome the disadvantage of MLR in the development of the BA formula.^{2,6,7,9-12} However, the PCA cannot avoid some of the statistical deficiencies of MLR.¹⁴ An alternative to this, Klemmer & Doubal Method (KDM) provides better precision in estimating BA than MLR and PCA methods.^{6,7,9,10} Although KDM gives the most reliable estimates in BA prediction, it involves complex calculations.⁶

This study focuses on developing the BA model for patients with Chronic Kidney Disease (CKD). The public-health effect from mortality due to this disease has not been fully assessed.¹⁵ Furthermore, CA does not give a good reflection on the time-dependent changes in kidney function.¹⁶ To better understand an individual degree of ageing or life span and how CKD influences an individual degree of ageing, a new approach needs to be developed.

In this study, we develop the BA using the indexing method. An index number is the most common statistical method to measure changes in a set of data points besides summarizing and ranking a particular data set. Moreover, measuring BA by examining the index number keeps track of the original representation of the data and thus ensures the output resembles the empirical structure closely. During this indexing process, each selected biomarker is given a

unique treatment corresponding to its severity level. Visual observations are also presented to justify the appropriateness of the method used.

Materials & Methods

IRB/Ethics approval

The data used in this study was approved by The Medical Research Ethics Committee, University of Malaya Medical Centre (MREC ID NO 2018428-6258). The committee granted permission to carry out the study within its facilities with common terms including; to adhere the instruction, guidelines and requirement by the committee. The Patient Information Sheet and Consent Form were waived by the committee. This retrospective study uses the data based on the earlier initiated and completed studies in a new outlook.

Measure of correlation

The strength between two variables can be measured using Pearson's correlation coefficient.¹⁷ A representative measure for this, the r -value, signifies both the magnitude and direction of the strength, that is, a closer value to ± 1 indicate high strength in the positive or negative direction. The r -value can be computed as:

$$r = \frac{n \sum_{i=1}^n x_i y_i - \left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n y_i \right)}{\sqrt{\left(n \sum_{i=1}^n x_i^2 - \left(\sum_{i=1}^n x_i \right)^2 \right) \left(n \sum_{i=1}^n y_i^2 - \left(\sum_{i=1}^n y_i \right)^2 \right)}} \quad (1)$$

where x and y are the value for the two variables and n is the total number of samples.

This study considers ten biomarker relationships with the CA; height, weight, gender, BMI, creatinine, e-GFR, PB Systolic, BP diastolic, CTCA calcium score and CKD stage from CKD patients. BA biomarkers that have an absolute r -value greater than 0.15 were selected for inclusion in BA calculation.^{7,12}

Weighted average method

This study proposes a weighted average method to estimate the weight for each significant biomarker. Note that the weight ranges from 0 to 1. Higher weight signifies a higher association between the health biomarkers to the BA index. The weight for biomarker i is computed as follows:

$$w_i = \frac{|r_i|}{\sum_{j=1}^n |r_j|} \quad (2)$$

where r_i is the correlation coefficient of the i^{th} biomarker computed using equation (1) and n is the total number of significant biomarkers. Note that the sum for all w_i 's equals to one, $\sum_{i=1}^n w_i = 1$.

Indexing method

This study uses the indexing method for BA calculation. The index produced from this method represents the amount of change with respect to the base value. For each biomarker, the base value is set to be the normal value or the favourable health condition. Thus, each biomarker has a unique indexing assignment based on the medical measurement they carry. In brief,

$$Index_{(i)} = \left| \frac{\text{Measured value } i - \text{Normal value } i}{\text{Normal value } i} \right| \quad (3)$$

where measured value is the current health reading of the patient while the normal value is the normal reading level for health biomarker i . Table 1 summarizes several common reading levels based on the standard clinical practice as well as work carried out in literature studies. The health biomarkers are categorized into several reading levels based on the severity of the postulated measurements.

Index equations are then developed based on Table 1. Note that the normal reference value is taken as the mid-point of the normal reading level. The index equation for the BMI is given by:

$$Index_{BMI} = \begin{cases} 1 - \left[\frac{4}{13}(x - 18.5) \right], & \text{if } 18.5 \leq x \leq 21.75, \\ 1 + \left[\frac{4}{33}(x - 30) \right], & \text{if } 21.75 < x \leq 30, \\ 1, & \text{otherwise.} \end{cases} \quad (4)$$

The index equation for the systolic blood pressure is given by:

$$Index_{SBP} = \begin{cases} 1 - \left[\frac{1}{5}(x - 120) \right], & \text{if } 120 \leq x \leq 125, \\ 1 + \left[\frac{1}{15}(x - 140) \right], & \text{if } 125 < x \leq 140, \\ 1, & \text{otherwise.} \end{cases} \quad (5)$$

The index equation for the diastolic blood pressure is given by:

$$Index_{DBP} = \begin{cases} 1 - \left[\frac{1}{2}(x - 80) \right], & \text{if } 80 \leq x \leq 82, \\ 1 + \left[\frac{1}{8}(x - 90) \right], & \text{if } 82 < x \leq 90, \\ 1, & \text{otherwise.} \end{cases} \quad (6)$$

129 The index equation for the eGFR is given by:

$$Index_{eGFR} = \begin{cases} 1, & \text{if } x \leq 15, \\ \frac{90 - x}{75}, & \text{if } 15 < x < 90, \\ 0, & \text{if } x \geq 90. \end{cases} \quad (7)$$

130 The index equation for the CTCA is given by:

$$Index_{CTCA} = \begin{cases} 0, & \text{if } x \leq 100, \\ \frac{x - 100}{300}, & \text{if } 100 < x < 400, \\ 1, & \text{if } x \geq 400. \end{cases} \quad (8)$$

131 Equations (4) to (8) formulate the framework for BA index calculation similar to medical
132 practices for six biomarkers to detect the severity level.

133 Figure 1 (a) to (e) show the aforementioned state graphically. Note that the health
134 biomarkers index ranges from 0 to 1 where 0 indicates a favourable level of the health
135 biomarker, index value from 0 to 1 indicates deteriorating health condition, while index value of
136 1 indicates a critically ill stage. The medical measures for each of the biomarkers are unique.
137 Therefore, the index value can be a useful comparative tool across these measurements.

138 In order to produce the biological index for each individual, each health biomarker index
139 is multiplied by its corresponding weight. The weight proportionately signifies the contribution
140 of each biomarker index to the BA index. Mathematically, the overall index for individual x is
141 computed as follows:

$$I_{BioAge \text{ for } x} = w_1 Index_{1,x} + w_2 Index_{2,x} + \dots + w_n Index_{n,x} \quad (9)$$

142 where w_i and $Index_{i,x}$ are the weight and index for the i^{th} health biomarker of individual x ,
143 respectively.

144

145 BA Estimation

146 Several methods have been proposed to estimate the BA. Among these are the multiple
147 linear regression (MLR) ⁹ and principal component analysis (PCA) ¹¹. PCA method is derived
148 from MLR and it reduces the effects of underestimated or overestimated BA.⁹ Both methods
149 show a linear relationship between BA and health parameters.

BA index models developed for predicting BA in this study also follow a linear relationship for individual general health status. It is developed based on the mathematical settings of the index method. The method suggests combining all individual subcomponents indices in one principal component (in this case, all health biomarkers into a single BA index). The subcomponent index measures the changes for each representative group of the biomarkers from the CA. Note however that the value of the index is not in year term. A common approach translating it into meaningful year-unit is by equation $BA_x = (I_{BioAge\ for\ x} \times standard\ deviation) + mean\ of\ CA$. Because the sample data focuses on the individual kidney patients, the following adjustment was made to the BA_x .

$$BA_x = (I_{BioAge\ for\ x} \times SD) + CA_x \quad (10)$$

where BA_x and CA_x are the biological age and chronological age for individual x , respectively. The standard deviation SD is computed based on the chronological age of the sample.

Bland-Altman analysis

It is vital to observe the mean values to understand the nature of a prediction model. It follows that the degree of dispersion from the mean indicates the fitness of individual BA values.⁸ Degrees of dispersion between BA and CA are commonly presented using the Bland-Altman plots. Bland-Altman plots the difference of CA and BA against the mean of the two measurements where it is easier to measure the magnitude, spot outlier, and see the data trend.¹⁸ If the differences are normally distributed, the mean differences should lie between $\bar{d} - 1.96s$ and $\bar{d} + 1.96s$ (95% confidence interval) where \bar{d} is the estimated mean difference and s is the standard deviation for the differences¹⁹.

Results and Discussion

Data

The study population consists of 190 patients subject to stage 1 to stage 4 CKD. Patients were recruited from the inpatient clinic, University of Malaya Medical Center (UMMC), Kuala Lumpur, Malaysia. Chronological age varied from 35 to 82 years, comprising males (115 patients) and females (75 patients). The age range was chosen to ensure that the population is old enough to be experiencing age-related changes in biomarkers.

Physical measurements include gender, height, weight, body mass index (BMI), systolic blood pressure (SBP), diastolic blood pressure (DBP), CTCA Calcium Score (CTCA), Creatinine, CKD stage and eGFR. It is observed that the CKD stage for the data involved is between 2 and 5. More specifically, 4% are in stage 2, 50% are in stage 3, 38% are in stage 4 and 8% are in stage 5. Table 2 shows the mean result, standard deviation (SD) and data range of CKD patients. For validation purposes of the BA model, 70 percent of the data are used as the training set (i.e., 133 data) while the remaining 30 percent for the testing set (i.e., 57 data).

Findings

Development of BA using the indexing method involves several sequences; correlation analysis, computation of weighted average for selected health biomarkers, construction of BA indices from the index equations and estimation of BA based on sample variations.

Biomarkers that show an absolute correlation coefficient value exceeding 0.15 are selected for BA estimation. An associated significance test that examine the correlation between the biomarkers and the chronological age for the training data set is summarized in Table 3. Seven biomarkers are found significant for inclusion in the BA estimation. It is observed that creatinine, weight, BMI, eGFR, BP systolic and BP diastolic decrease as age increases, while only the CTCA Calcium increases with age. Height, CKD stage and gender have a low correlation with age and are thus excluded from the BA estimation.

Note that some biomarker features have almost similar clinical implications and express high inter-correlation. It indicates the existence of redundancy. To cater for this redundancy, one biomarker with more substantial significance is selected. It is observed from Table 4 that weight and BMI have an absolute correlation coefficient of more than 0.15 (-0.234 and -0.173, respectively) and both show high correlation with each other (0.787). Due to the marginal difference in their correlation with age and clinical significance of the BMI, it is selected for BA estimation. Furthermore, BMI is more reliable to indicate the individual weight, either normal, overweight or obese ²⁰. Similar procedure is used for selection between the creatinine and eGFR. Both biomarkers represent the renal function. Creatinine and eGFR show mild inter-correlation (0.641) in addition to the absolute correlation coefficient of more than 0.15 (-0.202 and -0.233, respectively). Therefore, eGFR is selected for the BA estimation. With respect to its significance, measurement of eGFR is the most reliable assessment of renal function in CKD ²¹ where it is used as an index of renal function in clinical practice ²².

The weight for each parameter is derived based on the correlation analysis. The higher the correlation, the higher the weighted for the BA parameters. As shown in Table 5, BP Diastolic, CTCA and eGFR are the three highest contributors to the index value for predicting the BA.

Accordingly, five biomarkers including BMI, CTCA calcium, eGFR, BP systolic and BP diastolic are selected to estimate the BA index. The weightage for each of the selected biomarkers is then computed using equation (2) to arrive at the following BA index:

$$I_{BioAge} = 0.1422I_{BMI} + 0.2046I_{CTCA} + 0.1915I_{eGFR} + 0.1241I_{BPSystolic} + 0.3377I_{BPDiastolic} \quad (11)$$

Table 6 summarizes the estimated BA based on the BA index of equation (11). It is evident that all estimated BAs are higher than their corresponding CAs for CKD patients. Note however that the increase in BA varies for patients with identical CKD stages, acknowledging other competing factors that compensate for the difference. It is observed that the gain in BA for the 133 training data set ranges from 3 to 9 years with a mean of 7 years.

Overall, the BA for patients suffering kidney disease increased by 5% to 16% from its CA. Figure 2 shows that about 65% of kidney patients in stage 3 and stage 4 increase 9% to 12% from their CA. It indicates that on average the CKD patients at these stages gain between 5 to 9 years from their CA biologically.

Bland-Altman plot in Figure 3 exhibits the differences in CA and BA against the mean with a 95% confidence interval. All plots are shown below the zero value because this study utilizes data for patients diagnosed with CKD. Note however, the plot does not indicate whether the limits are acceptable or not. The judgment is based on clinical necessity. It is observed that most of the plots lie between the 95% confidence interval (CI), that is, they are inside the limit of agreement (LOA).

To explain further, the differences between CA and BA are plotted against the z-score of a standard normal distribution in Figure 4. It is observed that the majority of the plots fall along the straight line, which suggests the difference between CA and BA to follow the normal distribution. Furthermore, the correlation is found to be 0.9715 and the bell test using kurtosis test (0.170) indicates the degree of tailedness in the frequency is close to perfect normal distribution. In addition, the skewness test gives a value of 0.395 where the test value is between -0.5 and 0.5 or nearly zero, which justifies the assumption of normal distribution.

To further validate the BA index model, the testing dataset with 57 CKD patients is examined. Figure 5 shows the proximity between BA for the testing dataset and BA for the training dataset. BA of CKD patients increases as the CA increase due to the severity of the biomarkers measurements. On average, the age for patients diagnosed with CKD (stage 1 to stage 4) increases by seven years from their CA (64 years old) to BA (71 years old). Nevertheless, we conceived the small size of the sample and its representation of the population as the limitation of the study.

Conclusions

Recent studies have shown the importance of assessing the functional biomarkers for predicting ageing-related diseases. Ageing-related disease influences the BA of a person. Several statistical approaches have been observed in constructing the BA model, namely, the MLR, the PCA and the KDM.¹⁰ Each method owns its deficiency in the measurement of BA. Therefore, the indexing method is proposed to address issues, especially with redundant biomarkers (through manual examination of redundant biomarkers and selection of the relevant biomarker by experience), over or underestimated BA for a particular age group (by concentration to a particular disease group), and complicated calculations (through a tractable computation for each biomarker).

Estimation of BA using the indexing method proposed in this study facilitates a tractable form for each health biomarker. The initially calculated BA index represents the illness severity level for the CKD patients which contributes to proportionate gain in the BA. The level of

severity is categorized into high risk, normal risk and at risk so that each patient may be assessed individually. Ten biomarkers have been examined to see their appropriateness for inclusion in the BA estimation.

The results of this study show that patients with CKD between stage 2 to stage 5 experience gain in BA between 3 to 9 years. This finding may serve the medical practitioners a precaution for treatment in addition to current measurement facilities as it provides a comprehensive reading of the manifold biomarkers. The result is further validated with trained data and visual observations. Notwithstanding, increasing the sample size for the study and inclusion of diverse population may enhance the reliability of the end result.

Besides its practicality in the medical field, BA can be potentially useful in many areas, including the insurance industry where age and health play a central role in the premium calculation. Mortality projection based on the BA can be an exciting exploration of human lifetime behaviour that incorporates health biomarkers. In addition, an interesting future work is to develop methods which enables a fair comparison between the various biological age approaches.

Acknowledgements

The authors acknowledge the University of Malaya Medical Center (UMMC) for providing data for this research. The authors also thank the Editor and the Reviewers for their thoughtful comments and constructive suggestions.

References

1. Kim S, Jazwinski SM. Quantitative measures of healthy aging and biological age. *Healthy Aging Res.* 2015;4doi:10.12715/har.2015.4.26
2. Kang YG, Suh E, Chun H, et al. Models for estimating the metabolic syndrome biological age as the new index for evaluation and management of metabolic syndrome. *Clin Interv Aging.* 2017;12:253-261. doi:10.2147/CIA.S123316
3. Hayflick L. Biological aging is no longer an unsolved problem. *Annals of the New York academy of Sciences.* 2007;1100(1):1-13.
4. Jia L, Zhang W, Jia R, et al. Construction Formula of Biological Age Using the Principal Component Analysis. *Biomed Res Int.* 2016;2016:4697017. doi:10.1155/2016/4697017
5. Bae CY, Kang YG, Kim S, et al. Development of models for predicting biological age (BA) with physical, biochemical, and hormonal parameters. *Arch Gerontol Geriatr.* Sep-Oct 2008;47(2):253-65. doi:10.1016/j.archger.2007.08.009
6. Cho IH, Park KS, Lim CJ. An empirical comparative study on biological age estimation algorithms with an application of Work Ability Index (WAI). *Mech Ageing Dev.* Feb 2010;131(2):69-78. doi:10.1016/j.mad.2009.12.001


7. Jee H. Selection of a set of biomarkers and comparisons of biological age estimation models for Korean men. *J Exerc Rehabil.* Feb 2019;15(1):31-36. doi:10.12965/jer.1836644.322
8. Jee H, Park J. Selection of an optimal set of biomarkers and comparative analyses of biological age estimation models in Korean females. *Arch Gerontol Geriatr.* May - Jun 2017;70:84-91. doi:10.1016/j.archger.2017.01.005
9. Jia L, Zhang W, Chen X. Common methods of biological age estimation. *Clin Interv Aging.* 2017;12:759-772. doi:10.2147/CIA.S134921
10. Levine ME. Modeling the rate of senescence: can estimated biological age predict mortality more accurately than chronological age? *J Gerontol A Biol Sci Med Sci.* Jun 2013;68(6):667-74. doi:10.1093/gerona/gls233
11. Nakamura E, Miyao K. A method for identifying biomarkers of aging and constructing an index of biological age in humans. *J Gerontol A Biol Sci Med Sci.* Oct 2007;62(10):1096-105. doi:10.1093/gerona/62.10.1096
12. Park J, Cho B, Kwon H, et al. Developing a biological age assessment equation using principal component analysis and clinical biomarkers of aging in Korean men. *Arch Gerontol Geriatr.* Jul-Aug 2009;49(1):7-12. doi:10.1016/j.archger.2008.04.003
13. Krøll J, Saxtrup O. On the use of regression analysis for the estimation of human biological age. *Biogerontology.* 2000;1(4):363-368.
14. Klemra P, Doubal S. A new approach to the concept and computation of biological age. *Mech Ageing Dev.* Mar 2006;127(3):240-8. doi:10.1016/j.mad.2005.10.004
15. Wen CP, Cheng TY, Tsai MK, et al. All-cause mortality attributable to chronic kidney disease: a prospective cohort study based on 462 293 adults in Taiwan. *Lancet.* Jun 28 2008;371(9631):2173-82. doi:10.1016/S0140-6736(08)60952-6
16. Rowland J, Akbarov A, Maan A, et al. Tick-Tock Chimes the Kidney Clock - from Biology of Renal Ageing to Clinical Applications. *Kidney Blood Press Res.* 2018;43(1):55-67. doi:10.1159/000486907
17. Wackerly D, Mendenhall W, Scheaffer RL. *Mathematical statistics with applications.* Cengage Learning; 2014.
18. Altman DG, Bland JM. Measurement in medicine: the analysis of method comparison studies. *Journal of the Royal Statistical Society: Series D (The Statistician).* 1983;32(3):307-317.
19. Giavarina D. Understanding Bland Altman analysis. *Biochem Med (Zagreb).* 2015;25(2):141-51. doi:10.11613/BM.2015.015
20. Obesity NAAftSo, Heart N, Lung, et al. *The practical guide: identification, evaluation, and treatment of overweight and obesity in adults.* National Institutes of Health, National Heart, Lung, and Blood Institute, NHLBI Obesity Education Initiative, North American Association for the Study of Obesity.; 2000.
21. Bostom AG, Kronenberg F, Ritz E. Predictive performance of renal function equations for patients with chronic kidney disease and normal serum creatinine levels. *Journal of the American Society of Nephrology.* 2002;13(8):2140-2144.
22. Perrone RD, Madias NE, Levey AS. Serum creatinine as an index of renal function: new insights into old concepts. *Clinical chemistry.* 1992;38(10):1933-1953.
23. Fontana L, Hu FB. Optimal body weight for health and longevity: bridging basic, clinical, and population research. *Aging Cell.* Jun 2014;13(3):391-400. doi:10.1111/acer.12207

24. Walpole SC, Prieto-Merino D, Edwards P, et al. The weight of nations: an estimation of adult human biomass. *BMC Public Health*. Jun 18 2012;12:439. doi:10.1186/1471-2458-12-439
25. Ar AR, Y K, Chia YC. Management of hypertension. *Malaysian Family Physician*. 06/01 2011;6(1)
26. Rastogi A, Linden A, Nissenson AR. Disease management in chronic kidney disease. *Adv Chronic Kidney Dis*. Jan 2008;15(1):19-28. doi:10.1053/j.ackd.2007.10.011
27. Bhulani N, Khawaja A, Jafferani A, et al. Coronary calcium scoring: are the results comparable to computed tomography coronary angiography for screening coronary artery disease in a South Asian population? *BMC Res Notes*. Jul 19 2013;6:279. doi:10.1186/1756-0500-6-279
28. Neves PO, Andrade J, Monção H. Coronary artery calcium score: current status. *Radiologia brasileira*. 2017;50:182-189.


Figure 1

Biomarkers severity index


(a) BMI severity




(b) BP Systolic severity



(c) BP Diastolic severity



(d) eGFR severity



(e) CTCA calcium score severity




Figure 2

BA increment from CA by CKD stage




Figure 3

Bland-Altman plot




Figure 4

Scatter plot for differences between CA and BA versus the Z-score




Figure 5

CA versus BA

$$\bar{x}_{BA} = 71$$

$$\bar{x}_{CA} = 64$$




Table 1 (on next page)

Reading level for body mass index, blood pressure, eGFR, and calcium score

1 **Table 1:** Reading level for body mass index, blood pressure, eGFR, and calcium score

Health biomarkers	Reading level (Severity level)					Researchers
Body Mass Index (BMI) (in kg/m ²)	less than 18.5 (at risk)	18.5 to 24.9 (normal)	25 to 29.9 (moderate risk)	greater than 30 (high risk)		23,24
Blood Pressure (Systolic) (in mmHg)	less than 120 (at risk)	120 to 130 (normal)	130 to 139 (moderate risk)	greater than 140 (high risk)		25
Blood Pressure (Diastolic) (in mmHg)	less than 80 (at risk)	80 to 84 (normal)	85 to 89 (moderate risk)	greater than 90 (high risk)		25
eGFR ml/min/1.73m ²	≥ 90 Stage 1 (normal)	60 to 89 Stage 2 (low risk)	30 to 59 Stage 3 (moderate risk)	15 to 29 Stage 4 (moderately high risk)	<15 Stage 5 (high risk)	26
CTCA Calcium Score	0 (no risk)	less than 100 (low risk)	101 to 400 (moderately high risk)	> 400 (high risk)		27,28

2

3

Table 2(on next page)

General characteristics of the CKD patient biomarkers

1

Table 2: General characteristics of the CKD patient biomarkers.

Parameters	Male (n=115)	Female (n=75)	Combined (n=190)
Age (years)	62.88 ± 9.463	61.97 ± 9.988	62.52 ± 9.657
Weight (kg)	76.77 ± 12.305	68.60 ± 14.347	73.54 ± 13.709
Height (m)	1.66 ± 0.0647	1.523 ± 0.060	1.604 ± 0.091
BMI (kg/m ²)	27.93 ± 0.392	29.469 ± 5.109	28.535 ± 4.6328
CTCA Calcium Score	558.997 ± 733.855	291.853 ± 470.753	453.546 ± 654.786
Creatinine (μmol/L)	242.11 ± 130.025	229.59 ± 105.253	237.17 ± 120.718
eGFR (mL/min/1.73m ²)	35.513 ± 14.857	28.481 ± 12.432	32.737 ± 14.336
BPSystolic (mmHg)	146.88 ± 23.808	148.20 ± 21.075	147.4 ± 22.720
BPDiastolic (mmHg)	76.47 ± 14.906	76.23 ± 13.736	76.37 ± 14.419
CKD Stage	3.41 ± 0.687	3.67 ± 0.081	3.51 ± 0.703

2 **Note:** Mean±SD

3

Table 3(on next page)

p-values for the significance test

1

Table 3: p-values for the significance test

Parameters	p-values
Weight (kg)	0.003
Height (m)	0.056
BMI (kg/m ²)	0.023
CTCA Calcium Score	0.002
Creatinine (μmol/L)	0.010
eGFR (mL/min)	0.004
BPSystolic (mmHg)	0.041
BPDiastolic (mmHg)	0.000
CKD Stage	0.081
GenderT	0.421

Note: Trained data, n=133

2

Table 4(on next page)

Correlation coefficients (Pearson) between CA and biomarkers

1

Table 4: Correlation coefficients (Pearson) between CA and biomarkers

Parameter	Age	CTCA Calcium Score	Creatinine	Weight	Height	BMI	eGFR	BPSystolic	BPDiastolic	CKD Stage	Gender
Age	1.000	0.249	-0.202	-0.234	-0.138	-0.173	-0.233	-0.151	-0.411	0.122	0.017
		1.000	0.021	0.045	0.106	0.007	-0.056	0.009	-0.134	0.062	0.226
			1.000	0.042	0.113	-0.018	-0.641	0.179	0.021	0.716	0.066
				1.000	0.515	0.787	0.544	0.012	0.095	-0.452	0.369
					1.000	-0.108	0.292	0.087	0.115	-0.287	0.713
						1.000	0.399	-0.045	0.023	-0.306	-0.081
							1.000	-0.083	0.203	-0.862	0.276
								1.000	0.542	0.121	-0.016
									1.000	-0.138	0.053
										1.000	-0.259
											1.000

2

Table 5(on next page)

Average weighted for significant BA biomarkers

1

Table 5: Average weighted for significant BA biomarkers

Parameters	Weightage
Weight	0.142
BMI	0.105
CTCA	0.151
Creatinine	0.122
eGFR (mL/min)	0.141
BP Systolic (mmHg)	0.091
BPDiastolic (mmHg)	0.249

2

Table 6(on next page)

BA estimated using the BA index

1

Table 6: BA estimated using the BA index

No.	CA	CKD Stage	BA Index	BA	No.	CA	CKD Stage	BA Index	BA
1	64	4	0.5240	69	68	70	3	0.8971	78
2	60	4	0.6699	66	69	56	3	0.9104	65
3	48	3	0.5611	53	70	64	5	0.9575	73
4	65	4	0.5914	71	71	65	3	0.9392	74
5	67	4	0.7351	74	72	37	3	0.6988	44
6	63	5	0.9507	72	73	53	3	0.6813	59
7	60	5	0.7954	67	74	81	3	0.8380	89
8	59	3	0.7737	66	75	36	2	0.6562	42
9	70	4	0.9129	79	76	73	4	0.8726	81
10	66	3	0.9264	75	77	68	3	0.6789	74
11	61	3	0.7143	68	78	70	4	0.8573	78
12	82	4	0.7948	89	79	74	3	0.6681	80
13	57	5	0.8876	65	80	48	4	0.9191	57
14	48	2	0.6292	54	81	64	3	0.8854	72
15	67	4	0.6962	74	82	61	4	0.7756	68
16	43	3	0.8947	51	83	37	4	0.6919	44
17	81	5	1.0000	90	84	65	4	0.7499	72
18	55	3	0.3852	59	85	78	3	0.7795	85
19	63	3	0.7392	70	86	77	4	0.9145	86
20	63	3	0.7570	70	87	69	4	0.8449	77
21	68	4	0.9649	77	88	58	3	0.7076	65
22	60	4	0.7578	67	89	63	3	0.7245	70
23	67	4	0.7214	74	90	63	3	0.9308	72
24	58	5	0.9680	67	91	70	3	0.7933	77
25	58	4	0.7490	65	92	74	4	0.8170	82
26	56	4	0.8093	64	93	53	4	0.6986	60
27	43	4	0.9002	51	94	67	4	0.7660	74
28	62	4	0.8951	70	95	71	3	0.7973	79
29	75	3	0.8297	83	96	53	5	0.6726	59
30	55	4	0.9661	64	97	52	3	0.6456	58
31	62	4	0.6336	68	98	60	3	0.8970	68
32	63	4	0.7361	70	99	73	3	0.7774	80
33	72	4	0.7405	79	100	61	5	0.5983	67
34	76	4	0.8005	84	101	66	3	0.5996	72

35	43	4	0.6770	49	102	71	4	0.9006	79
36	59	3	0.9323	68	103	78	3	0.6414	84
37	72	3	0.6619	78	104	64	4	0.6862	70
38	64	4	0.6324	70	105	50	3	0.6637	56
39	72	4	0.8376	80	106	68	3	0.7547	75
40	46	3	0.7040	53	107	53	3	0.8595	61
41	75	4	0.7902	82	108	75	3	0.7372	82
42	69	3	0.6955	76	109	63	3	0.9176	72
43	66	4	0.6984	73	110	58	4	0.6760	64
44	68	2	0.4181	72	111	82	4	0.9827	91
45	64	3	0.6099	70	112	68	3	0.7254	75
46	55	4	0.9653	64	113	67	3	0.7050	74
47	63	4	0.7009	70	114	61	5	0.8571	69
48	63	3	0.3643	66	115	51	4	0.7512	58
49	69	4	0.8109	77	116	54	5	0.7234	61
50	69	3	0.8070	77	117	72	3	0.5938	78
51	76	4	0.6379	82	118	67	3	0.7171	74
52	68	4	0.7109	75	119	77	4	0.8781	85
53	65	3	0.5285	70	120	60	3	0.6174	66
54	65	3	0.8299	73	121	64	3	0.7517	71
55	59	3	0.7763	66	122	62	3	0.5667	67
56	69	4	0.8914	77	123	82	4	0.9089	91
57	76	3	0.9309	85	124	74	4	0.8258	82
58	69	4	0.7874	76	125	74	4	0.8620	82
59	63	5	0.6161	69	126	62	3	0.7566	69
60	48	3	0.6912	55	127	63	4	0.7086	70
61	59	5	0.9720	68	128	67	3	0.7509	74
62	53	2	0.4258	57	129	59	3	0.6114	65
63	56	2	0.6054	62	130	71	5	0.9078	80
64	65	2	0.7176	72	131	71	4	0.6842	77
65	69	4	0.9307	78	132	65	3	0.6418	71
66	58	3	0.6751	64	133	71	3	0.9161	80
67	57	3	0.6600	63					