

# Characterization of salivary microbiome in patients with pancreatic cancer

Pedro J. Torres, Erin M. Fletcher, Sean M. Gibbons, Michael Bouvet, Kelly S. Doran, Scott T. Kelley

Clinical manifestations of pancreatic cancer often do not occur until the cancer has undergone metastasis resulting in a very low survival rate. In this study, we investigated whether salivary bacterial profiles might provide useful biomarkers for early detection of pancreatic cancer. Using high-throughput sequencing of bacterial small subunit ribosomal RNA (16S rRNA) gene, we characterized the salivary microbiota of patients with pancreatic cancer and compared them to healthy patients and patients with digestive or non-digestive diseases. A total of 146 patients were enrolled at the UCSD Moores Cancer Center where saliva and demographic data were collected from each patient. Of these, we analyzed the salivary microbiome of 108 patients: 8 had been diagnosed with pancreatic cancer, 13 with pancreatic disease, 53 with digestive diseases (including cancer), 12 with non-digestive diseases (including non-digestive cancer) and 22 were classified as non-diseased (healthy) controls. Bacterial 16S rRNA sequences were amplified directly from salivary DNA extractions and subjected to high-throughput sequencing (HTS). We found significant differences in the abundances of several bacterial genera between patients with pancreatic cancer and patients in other categories, including *Porphyromonas*, *Haemophilus*, and *Campylobacter*. In particular, we found the bacterial ratio of *Leptotrichia* to *Porphyromonas* was consistently higher in pancreatic cancer patients compared to those with pancreatic disease, other digestive diseases and healthy controls. Interestingly, we did not find lower abundances of *Neisseria elongata* and *Streptococcus mitis* or higher numbers of *Granulicatella adiacens* reported previously using microarray methods. While salivary bacterial biomarkers appear promising, larger sample studies and repeat sampling of individuals are needed to confirm the validity of particular markers.

Characterization of the Salivary Microbiome in Patients with Pancreatic Cancer

Pedro J. Torres<sup>1</sup>, Erin M. Fletcher<sup>1</sup>, Sean M. Gibbons<sup>2,3</sup>, Michael Bouvet<sup>4</sup>, Kelly S. Doran<sup>1,5</sup>,  
Scott T. Kelley<sup>1</sup>

<sup>1</sup> Department of Biology, San Diego State University, 5500 Campanile Drive, San Diego, CA  
92182-4164, USA

<sup>2</sup> Graduate Program in Biophysical Sciences, University of Chicago, Chicago, IL 60637, USA

<sup>3</sup> Institute for Genomics and Systems Biology, Argonne National Laboratory, 9700, S. Cass Ave.  
Lemont, IL 60439, USA

<sup>4</sup> Department of Surgery, University of California, 3855 Health Sciences Drive, La Jolla, CA  
92093-0987, USA

<sup>5</sup> Department of Pediatrics, University of California San Diego School of Medicine, La Jolla,  
CA, 92093, USA.

**Running Title:** Salivary Microbial Biomarkers for Pancreatic Cancer

**#Correspondence to:** Dr. Scott T. Kelley, Department of Biology, San Diego State University,  
5500 Campanile Drive, San Diego, CA 92182-4164, USA; Tel: 619-594-5371, Fax: 619-594-  
5676, [skelley@mail.sdsu.edu](mailto:skelley@mail.sdsu.edu)

**Word count:** 3,599

**Total number of figures and tables:** 7

# Abstract

Clinical manifestations of pancreatic cancer often do not occur until the cancer has undergone metastasis, resulting in a very low survival rate. In this study, we investigated whether salivary bacterial profiles might provide useful biomarkers for early detection of pancreatic cancer. Using high-throughput sequencing of bacterial small subunit ribosomal RNA (16S rRNA) gene, we characterized the salivary microbiota of patients with pancreatic cancer and compared them to healthy patients and patients with other diseases, including pancreatic disease, non-pancreatic digestive disease/cancer and non-digestive disease/cancer. A total of 146 patients were enrolled at the UCSD Moores Cancer Center where saliva and demographic data were collected from each patient. Of these, we analyzed the salivary microbiome of 108 patients: 8 had been diagnosed with pancreatic cancer, 78 with other diseases and 22 were classified as non-diseased (healthy) controls. Bacterial 16S rRNA sequences were amplified directly from salivary DNA extractions and subjected to high-throughput sequencing (HTS). Several bacterial genera differed in abundance in patients with pancreatic cancer. We found a significantly higher ratio of *Leptotrichia* to *Porphyromonas* in the saliva of patients with pancreatic cancer than in the saliva of healthy patients or those with other disease (Kruskal-Wallis Test;  $P < 0.001$ ). *Leptotrichia* abundances were confirmed using real-time qPCR with *Leptotrichia* specific primers. Similar to

previous studies, we found lower relative abundances of *Neisseria* and *Aggregatibacter* in the saliva of pancreatic cancer patients, though these results were not significant at the  $P < 0.05$  level (K-W Test;  $P = 0.07$  and  $P = 0.09$  respectively). However, the relative abundances of other previously identified bacterial biomarkers, e.g. *Streptococcus mitis* and *Granulicatella adiacens*, were not significantly different in the saliva of pancreatic cancer patients. Overall, this study supports the hypothesis that bacteria abundance profiles in saliva are useful biomarkers for pancreatic cancer though much larger patient studies are needed to verify their predictive utility.

# Introduction

In the United States, approximately 40,000 people die every year from pancreatic adenocarcinoma, making it the fourth leading cause of cancer related death. Patients diagnosed in the early stage of pancreatic cancer have a 5-year survival rate of 24%, compared to 1.8% when diagnosed in the advanced stage (1). Clinical manifestations of pancreatic cancer do not appear until after the cancer has undergone metastasis (2), emphasizing the need for early detection biomarkers. The etiology of pancreatic cancer remains elusive, with cigarette smoking being the most established risk factor (3–6), although links have also been made to diabetes (7; 8), obesity (9), and chronic pancreatitis (10). Recent research has also shown that men with periodontal disease have a two-fold greater risk of developing pancreatic cancer after adjusting for smoking, diabetes, and body mass index (11).

The human oral cavity harbors a complex microbial community (microbiome) known to contain over 700 species of bacteria, more than half of which have not been cultivated (12). Researchers have identified a core microbial community in healthy individuals (13) and shifts from this core microbiome have been associated with dental caries and periodontitis (14). The composition of bacterial communities in saliva seems to reflect health status under certain circumstances (15), making the analysis of salivary microbiomes a promising approach for disease diagnostics. A study by Mittal *et al.* (2011) found that increases in the numbers of *Streptococcus mutans* and lactobacilli in saliva have been associated with oral disease prevalence (16), while another study showed that high salivary counts of *Capnocytophaga gingivalis*, *Prevotella melaninogenica* and *Streptococcus mitis* may be indicative of oral cancer (17).

A recent study by Farrell *et al.* (2012), suggested that the abundances of specific salivary bacteria could be used as biomarkers for early-stage pancreatic cancer. Using the Human Oral

Microbe Identification Microarray (HOMIM), researchers observed decreased levels of *Neisseria elongata* and *Streptococcus mitis* in patients with pancreatic cancer compared with healthy individuals, while levels of *Granulicatella adiacens* were significantly higher in individuals with pancreatic cancer (18). The HOMIM's ability to detect 300 of the most prevalent oral bacterial species has made it a suitable method for assessing community profiles at the phylum level as well as many common taxa at the genus level. However, the HOMIM microarray method fails to detect approximately half of the bacterial species commonly present in saliva (19).

In this study, we applied high-throughput sequencing (HTS) of the bacterial small-subunit ribosomal RNA (16S rRNA) genes to determine the salivary profiles of patients with and without pancreatic cancer. The use of HTS to sequence 16S rRNA bacterial genes from entire salivary microbial communities allows for a more comprehensive profile of the microbiome in health and disease (20). During this study, we collected 146 saliva samples from patients at the UCSD Moores Cancer Center. HTS was used to characterize the salivary microbiome of patients with pancreatic cancer and compare them to patients with other diseases (including pancreatic disease, non-pancreatic digestive disease/cancer and non-digestive disease/cancer) as well as non-diseased (healthy) controls. This allowed us to test the hypothesis that patients with pancreatic cancer may have a distinct microbial community profiles compared to non-diseased controls and to other forms of digestive and non-digestive diseases. Our results demonstrated that patients with pancreatic cancer had a significantly higher abundance ratio of particular bacterial genera.

## Materials and Methods

### Sample collection and patient information

This study was approved by the University of California San Diego (UCSD) and San Diego State University (SDSU) joint Institutional Review Board (IRB Approval #120101). Patients recruited for the study were being clinically evaluated at the UCSD Moores Cancer Center or were undergoing endoscopy procedures by UCSD Gastroenterologists in the Thornton Hospital Pre-Procedure Clinic between May 2012 and August 2013. All patients were required to fast for 12-hours prior to cancer evaluation and endoscopy procedures. To avoid bias during enrollment, the research coordinator responsible for recruiting participants' was unaware of patient diagnosis at time of sample collection. Consenting participants were provided with IRB-approved consent forms, and HIPAA forms, as well as an optional, voluntary written survey in which they could share relevant information about antibiotic, dental and smoking history. All participants gave informed consent and their identities were withheld from the research team. Each subject was free to withdraw from the study at any time. Participants were asked to give a saliva sample into a 50 mL conical tube. If the amount of saliva exceeded 55 uL, 10 uL was transferred into tube containing Brain-Heart Infusion media (BHI) and glycerol for future culturing. The remaining saliva was broken up into 55 uL aliquots and stored in sterile cryovials. Both BHI and saliva samples were then immediately stored at -80 °C until further processing.

Of the 146 participants, three subjects voluntarily withdrew and seven were not included in the study due insufficient production of saliva (<55 uL) leaving 136 saliva samples. After sample collection, the research coordinator accessed the participants' medical records electronically for patient diagnosis information that was included under a novel subject ID number. Diagnosis was used to determine health status and assess the stage of disease when each sample was taken. The various diagnoses were grouped into the following categories: pancreatic cancer, other disease (including pancreatic disease, non-pancreatic digestive disease/cancer and

non-digestive disease/cancer), and healthy (non-diseased) controls. Healthy individuals were defined as participants with no documented chronic digestive or non-digestive disease, and a 5-year resolution of any previously documented digestive or non-digestive disease. Exclusion criteria included participants undergoing active chemotherapy or radiation therapy or use of antibiotics two weeks prior to saliva collection as well as invasive surgery in the past year.

### **DNA Isolation, PCR and 16S rRNA Sequencing**

Bacterial DNA was extracted directly from 50 uL of patient saliva using the MoBio PowerSoil DNA Extraction Kit (Catalogue 12888-05, Mo Bio Laboratories, Carlsbad, CA, USA) following the manufacturer's protocol. Genomic DNA was quantified using the NanoDrop™ Spectrophotometer and stored at -20 °C.

The 16S ribosomal RNA (rRNA) amplicon region was amplified using barcoded 'universal' bacterial primer 515F (5'- AATGATACGGCGACCACCGAGATCTACAC TATGGTAATT GT GTGCCAGCMGCCGCGGTAA-3') and 806R (5'- CAAGCAGAAGACGGCATACGAGAT XXXXXXXXXXXXX AGTCAGTCAG CC GGACTACHVGGGTWTCTAAT-3') (X's indicate the location of the 12-bp barcode) with Illumina adaptors used by the Earth Microbiome Project (<http://www.earthmicrobiome.org/emp-standard-protocols/16s>). The barcoded primers allow pooling of multiple PCR amplicons in a single sequencing run. PCR was carried out using the reaction conditions outlined by the Earth Microbiome Project. Thermocycling parameters were as follows: 94°C for 3 minutes (denaturing) followed by amplification for 35 cycles at 94°C for 45 s, 50°C for 60 s and 72°C for 90 s, and a final extension of 72°C for 10 minutes (21). PCR amplicons were then sequenced on



the Illumina MiSeq platform at the Argonne National Laboratory Core sequencing facility (Lemont, IL).

# **Sequence Analysis**

16S rRNA sequences were de-multiplexed using the Quantitative Insights Into Microbial Ecology (QIIME v. 1.8.0, <http://www.qiime.org>) pipeline. Sequences were grouped into operational taxonomic units (OTUs) at 97% sequence similarity using the Greengenes reference database. OTUs that did not cluster with known taxa at 97% identity or higher in the database were clustered *de novo* (UCLUST (22)). Representative sequences for each OTU were then aligned using PyNast (23), and taxonomy was assigned using the RDP classifier (Version 2.2) (24). A phylogenetic tree was built using FastTree (25). Before performing downstream analysis, patient samples were rarefied to 10,000 sequences per sample, singletons and OTUs present in <25% of samples were removed prior to rarefaction. Chimeric sequences were identified using ChimeraSlayer in QIIME, as well as with DECIPHER (26), and subsequently removed. Alpha diversity metrics were computed using QIIME. Beta diversity distance between samples (weighted and unweighted UniFrac) were computed and used to account for both differences in relative abundance of taxa and phylogeny (27). Beta diversity comparisons were done using analysis of similarities (ANOSIM). We also tested whether there were significant differences in abundance ratios of particular genera between our different categories with GraphPad Prism version 6.0 using the Kruskal-Wallis test followed by Dunn's multiple comparison correction. Statistical significance was accepted at a  $p < 0.05$ . Analysis and identification of potential contaminants was done using SourceTracker (28).

# **Quantitative PCR (qPCR)**

*Leptotrichia* abundance was determined using qPCR. Briefly, for each sample we estimated *Leptotrichia* abundance using *Leptotrichia* specific 16S primers and normalized their values to overall bacterial abundance estimated using qPCR with universal bacterial 16S primers (5'-TCCTACGGGAGGCAGCAGT-3' forward primer, and 5'-GGACTACCAGGGTATCTAATCCTGTT-3' reverse primer) developed by Nadkarni et al. (2002) (29). qPCR was performed on a Bio-Rad CFX96 Touch™ Real-Time PCR Detection Instrument. The maximum C<sub>t</sub> (threshold cycle) for the universal 16S primers was set to 35 cycles and C<sub>t</sub> levels above this threshold were considered background noise. Genus-specific primers for amplification of *Leptotrichia* were designed using 16S rRNA sequences obtained from the RDP classifier (Version 2.2) (24). Primer3 online software was used for primer selection, and conditions were settled following the recommendations of Thorton and Basu, 2011 (30). The *Leptotrichia* forward primer sequence (5'- GGAGCAAACAGGATTAGATACCC - 3') and the *Leptotrichia* reverse primer sequence (5'- TTCGGCACAGACACTCTTCAT- 3') generated an amplicon of 87 bp. The PCR reaction contained 1 uM of both forward and reverse *Leptotrichia* primers with thermocycling parameters of 50°C for 2 min, 95°C for 10 min and 40 cycles of 95°C for 15 sec and 62.5°C for 1 min. The amplification reactions for the universal primers and *Leptotrichia* primers were carried out in at least duplicate using 25 uL of SYBR Green Master Mix (Bio-Rad) and 0.85 ng/uL of extracted DNA as template. Various online tools, including *In silico* PCR Amplification (31) and Ribosomal Database Project (24) were used to check the specificities of the oligonucleotide primer sequences for the target organism. A saliva sample was sequenced (Eton Bioscience, San Diego, CA) using our novel primers and primer specificity was further confirmed with a 16S rRNA database BlastN search.

# Results

Salivary microbial diversity profiles were generated for a total of 108 patients. 8 patients were diagnosed with pancreatic cancer (P), 78 were diagnosed with other diseases (including cancer) (O), and 22 were considered healthy (non-diseased) controls (H). Table 1 details the individual clinical characteristics, including, gender and ethnicity. Of the 108 patients, 23 patients in pancreatic, digestive, and non-digestive disease categories were diagnosed with having cancer. Table 2 details the types of cancer, as well as category groupings and the mean age of the cancer patients in each category.

Illumina sequencing yielded approximately 6.8 million sequences across all samples. The sequences are available on FigShare (<http://dx.doi.org/10.6084/m9.figshare.1422174>) along with the mapping file (<http://dx.doi.org/10.6084/m9.figshare.1422175>). An analysis of potential sample contamination using SourceTracker (28) identified some evidence of human skin and/or environmental contamination. The sequences associated with OTUs identified as contaminants, mostly *Staphylococcus* (skin) and *Cyanobacteria* (chloroplasts), were removed from all subsequent analyses. From these data, we identified a total of 12 bacterial phyla and 139 genera. Proteobacteria, Actinobacteria, Bacteroidetes, Firmicutes, and Fusobacteria were the 5 major phyla, accounting for 99.3% of oral bacteria (Figure 1). The mean relative abundance of Proteobacteria was lower in pancreatic cancer patients relative to other sample categories, while Firmicutes tended to be higher, though these were not significant after adjusting for multiple comparisons (FDR). The pancreatic cancer group also had higher levels of *Leptotrichia*, as well as lower levels of *Porphyromonas*, and *Neisseria* (Figure 2). In general, multi-level taxonomic profiles of the healthy group resembled the ‘other’ disease group, while the pancreatic cancer group was readily distinguishable (Fig. S1). However, there were no significant differences

217 among the three main groupings (H, O, and P) in either beta diversity (ANOSIM;  $P=0.1$ ) or  
218 alpha diversity (Chao1, K-W test;  $P=0.6$ ; Faith's PD, K-W test;  $P=0.56$ ).

219 As in previous studies by Farrell *et al.* (2012) and Lin *et al.* (2013) (32), we saw lower  
220 relative abundances of *Neisseria* and *Aggregatibacter*, although these differences were not  
221 significant (K-W test;  $P=0.07$  and  $P=0.09$  respectively). *Bacteriodes* was more abundant in  
222 pancreatic cancer patients compared to healthy individuals, similar to what Lin *et al.* observed,  
223 although this too was not significant (K-W test;  $P=0.27$ ). We did not see any difference in the  
224 relative abundance of *Streptococcus* or *Granulicatella*, which were shown to differ in a prior  
225 pancreatic cancer study (18). Additional analytical targets were based on a preliminary study  
226 consisting of our first 61 saliva samples (including 3 from pancreatic cancer patients), which  
227 showed significantly higher *Leptotrichia* and lower *Porphyrromonas* in pancreatic cancer patient  
228 saliva.

229 The abundance ratio of *Leptotrichia*, specifically two OTUs (arbitrarily named OTU  
230 31235 and OTU 4443207), to *Porphyrromonas* was significantly higher in pancreatic cancer  
231 patients (Figure 3). A BLAST comparison of these OTUs to the 16S sequence in the Human Oral  
232 Microbiome database (33) (HOMD RefSeq Version 13.2) found OTU 31235 to be 100% similar  
233 to *Leptotrichia sp. Oral taxon 221*, while OTU 4443207 was 99.3% similar to *Leptotrichia*  
234 *hongkongensis*. We found a strong positive correlation (Pearson's correlation  $r=0.903$ ,  
235  $P=0.0000001$ ) between *Leptotrichia* abundances obtained from 16S rRNA sequencing (OTU  
236 relative abundances) and from real-time qPCR (Figure 4).

# Discussion

Our analysis of salivary microbial profiles supports prior work suggesting that salivary microbial communities of patients diagnosed with pancreatic cancer are distinguishable from salivary microbial communities of healthy patients or patients with other diseases, including non-pancreatic cancers. At the phylum level, pancreatic cancer patients tended to have higher proportions of Firmicutes and lower proportions of Proteobacteria (Figure 1). At finer taxonomic levels, we observed differences in the mean relative abundances of particular genera in pancreatic cancer patients compared to other patient groups (Figure 2). For instance, there was a higher proportion of *Leptotrichia* in pancreatic cancer patients, while the proportion of *Porphyromonas* and *Neisseria* were lower in these patients.

The most striking difference we found between the microbial profiles of pancreatic cancer patients and other patient groups was in the ratio of the bacterial genera *Leptotrichia* and *Porphyromonas* (LP ratio) (Figure 3). The LP ratio had been identified as a potential biomarker from a preliminary analysis and an analysis of the full dataset found significantly higher LP ratio in pancreatic cancer patient saliva than in other patient groups. To verify these differences using another method, we cross-validated the relative abundances of *Leptotrichia* (Figure 4). Interestingly, during the analysis of the 16S rRNA data, we successfully used the LP ratio to reclassify one of the patients in the non-pancreatic cancer disease group. This particular individual had been initially diagnosed as having an unknown digestive disease, but the patient's high LP ratio suggested pancreatic cancer (Figure 3). Subsequently, the patient was re-evaluated and diagnosed with pancreatic cancer, supporting the notion that the LP ratio may serve as a pancreatic cancer biomarker.

Despite the small cohort of patients in this study, we believe our results are especially noteworthy because we were able to distinguish between patients with pancreatic cancer and patients with a variety of other diseases (including non-pancreatic cancer), in addition to healthy controls. Other researchers have proposed the use of ratios of bacterial taxa previously. Galimanas *et al.* (2014) suggested using salivary bacteria abundance ratios as a means for differentiating between healthy and diseased patients (34). Taxonomic ratios have been used to differentiate between subjects in studies of obesity (35), diabetes (36), and periodontal disease (37). Ratio comparisons also help to control for high levels of taxonomic variability among individuals (38–41).

A review of the literature revealed that *Leptotrichia*'s role in oral health remains elusive. However, these bacteria have been found in the bloodstream of immune-compromised patients (42) and co-occur significantly with colorectal tumors (43). *Leptotrichia* have been isolated from cardiovascular and gastrointestinal abscesses, from systemic infections, and are thought to be pathogenic (44). In regards to *Porphyromonas*, antibodies to *Porphyromonas gingivalis* have been directly associated with pancreatic cancer (45). A European cohort study measured plasma antibodies to 25 oral bacteria in pre-diagnostic blood samples from 405 pancreatic cancer patients and 416 matched controls and found a >2-fold increase in risk of pancreatic cancer among those with higher antibody titers to a pathogenic strain of *P. gingivalis* (45). At first glance, it appears contradictory that individuals with higher *Porphyromonas* antibody titers would have lower oral abundances. However, studies of systemic immunization of animals to particular periopathogens including *Porphyromonas* have shown reduced colonization of these bacteria in the mouth and a reduction of periodontitis (46–48). Similarly, higher *Porphyromonas*

antibody titers in individuals with pancreatic cancer may decrease their oral abundance, though this connection needs to be formally tested.

Shifts in salivary microbial diversity could also be a systematic response to pancreatic cancer. Pancreatic cancer is known to weaken the immune system (49), which could lead to overgrowth of oral bacteria and a shift towards systemically invasive periodontal pathogens. The proliferation of bacterial pathogens could assist cancer progression through systemic inflammation (50) or immune distraction (51). Thus, an initial increase in *Porphyromonas* might be followed by a decrease due to systemic invasion and antibody production. Indeed, inflammation is thought to play a significant role in the development of pancreatic cancer (52).

We also compared the relative abundances of several other bacterial genera that were indicated as potential biomarkers in previous work by Farrell *et al.* (2012). Like Farrell *et al.*, we found a lower proportion of *Neisseria* in pancreatic cancer patient saliva compared with the healthy and other disease category, though this trend was not significant. However, we did not find the same results as Farrell *et al.* for the other bacterial genera they identified. Our data also showed an increase in *Bacteroides* and decrease in the abundance of the bacterial genus *Aggregatibacter* in patients with pancreatic cancer, supporting the results of a pilot study by Lin *et al.* (2013), though neither trends were significant.

Methodological differences between our study and the Farrell *et al.* study in particular, may partially explain our divergent results. For instance, the inability of the V4 region of the 16S rRNA gene to discriminate *Streptococcus mitis* from other *Streptococcus* species may have prevented us from detecting difference in this species' abundance (18). Additionally, our study had a broader array of patient categories and cancers were not always confined to the pancreas at the time of sampling.

Interestingly, since the completion of our study, Mitsuhashi *et al.* (2015) reported the detection of oral *Fusobacterium* in pancreatic cancer tissue (53). A retrospective review of our abundance data also found a lower relative abundance of *Fusobacterium* in pancreatic cancer patients compared to other patient categories (Figure 2; K-W test,  $P=0.03$  prior to FDR correction) suggesting the processes driving differences in *Fusobacterium* may be similar to our proposed mechanism for *Porphyromonas*. Although the result was not significant after adjusting for multiple-comparisons (FDR), we suggest *Fusobacterium* abundance should be considered as a potential biomarker target for future studies with larger patient cohorts.

Overall, our study suggests that members of the salivary microbiome have promise as potential pancreatic cancer biomarkers and we may have uncovered an important new prospect in this regard (i.e., the LP ratio). However, our relatively small number of samples from pancreatic cancer patients and the discrepancies between our findings and previous work indicate that much larger patient cohorts will be needed to determine whether salivary biomarkers are diagnostically useful. Future studies should focus on improved metadata collection, including diet and oral health information (i.e. periodontal disease), which would make it possible to run statistical analyses that control for multiple factors involved in shaping oral microbial diversity. It will also be important to sample the same individual's saliva over time to assess whether we can distinguish between disease stages and also to control for intra-individual variation. Further, it is possible that single biomarkers may never be able to consistently identify pancreatic patients from other conditions. Thus, we may need more complex metrics that combine the abundances of multiple salivary bacteria, metabolite profiles, and detailed patient metadata. Effective diagnostic biomarkers for pancreatic cancers have been difficult to find, but are sorely needed and have the potential to save thousands of lives each year.



328

# 329 Acknowledgements

330 We thank S. Owens and J. Marcel for their help in our sequencing runs. We also thank M.  
331 Wachter for assistance in enrolling patients into the study. Thanks to all the patients who  
332 volunteered to be a part of this study.

333

334

# 335 References:

- 336 1. Li D, Xie K, Wolff R, Abbruzzese JL. Pancreatic cancer. *Lancet* 2004;**363**:1049-57.
- 337 2. Holly EA, Chaliha I, Bracci PM, Gautam M. Signs and symptoms of pancreatic  
338 cancer: a population-based case-control study in the San Francisco Bay area. *Clin Gastroenterol*  
339 *H* 2004;**2**:510-17.
- 340 3. Vrieling A, Bueno-de-Mesquita HB, Boshuizen HC, Michaud DS, Severinsen  
341 MT, Overvad K, Olsen A, Tjønneland A, Clavel-Chapelon F, Boutron-Ruault MC, Kaaks  
342 R, Rohrmann S, Boeing H, Nöthlings U, Trichopoulou A, Moutsiou E, Dilis V, Palli D, Krogh  
343 V, Panico S, Tumino R, Vineis P, van Gils CH, Peeters PH, Lund E, Gram IT, Rodríguez  
344 L, Agudo A, Larrañaga N, Sánchez MJ, Navarro C, Barricarte A, Manjer J, Lindkvist B, Sund  
345 M, Ye W, Bingham S, Khaw KT, Roddam A, Key T, Boffetta P, Duell EJ, Jenab M, Gallo  
346 V, Riboli E. Cigarette smoking, environmental tobacco smoke exposure and pancreatic cancer  
347 risk in the European Prospective Investigation into Cancer and Nutrition. *Int J Cancer*  
348 2010;**126**:2394–2403.
- 349 4. Nakamura K, Nagata C, Wada K, Tamai Y, Tsuji M, Takatsuka N, Shimizu H. Cigarette  
350 Smoking and Other Lifestyle Factors in Relation to the Risk of Pancreatic Cancer Death: A

- 351 Prospective Cohort Study in Japan. *Jpn J Clin Oncol* 2011;**41**:225–231.
- 352 5. Fuchs CS, Colditz GA, Stampfer MJ. A prospective study of cigarette smoking and the  
353 risk of pancreatic cancer. *Arch Intern Med* 1996;**156**:2255-60.
- 354 6. Zheng W, McLaughlin JK, Gridley G, Bjelke E. A cohort study of smoking, alcohol  
355 consumption, and dietary factors for pancreatic cancer (United States). *Cancer Cause Control*  
356 1993;**4**:477-82.
- 357 7. Haugvik S-PP, Hedenström P, Korsæth E, Valente R, Hayes A, Siuka D, Maisonneuve  
358 P, Gladhaug IP, Lindkvist B, Capurso G. Diabetes, Smoking, Alcohol and Family History of  
359 Cancer as Risk Factors for Pancreatic Neuroendocrine Tumors: A Systematic Review and Meta-  
360 Analysis. *Neuroendocrinology* 2015 Jan 22. [Epub ahead of print].
- 361 8. Liu X, Hemminki K, Försti A, Sundquist K, Sundquist J, Ji J. Cancer risk in patients with  
362 type 2 diabetes mellitus and their relatives. *Int J Cancer* 2015 Jan 21. [Epub ahead of print].
- 363 9. Bracci PM. Obesity and pancreatic cancer: overview of epidemiologic evidence and  
364 biologic mechanisms. *Mol Carcinogen* 2012;**51**:53-63.
- 365 10. Malka D, Hammel P, Maire F, Rufat P, Madeira I. Risk of pancreatic adenocarcinoma  
366 in chronic pancreatitis. *Gut* 2002;**51**:849-52.
- 367 11. Michaud DS, Joshupura K, Giovannucci E, Fuchs CS. A prospective study of periodontal  
368 disease and pancreatic cancer in US male health professionals. *J Natl Cancer I* 2007;**99**:171–5.
- 369 12. Aas JA, Paster BJ, Stokes LN, Olsen I. Defining the normal bacterial flora of the oral  
370 cavity. *J Clin Microbiol* 2005;**43**:5721-32.
- 371 13. Zaura E, Keijser B, Huse SM. Defining the healthy "core microbiome" of oral microbial  
372 communitites. *BMC Microbiol* 2009;**9**:259.
- 373 14. Berezow AB, Darveau RP. Microbial shift and periodontitis. *Periodontol* 2000

- 2011;**55**:36-47.
15. Yamanaka W, Takeshita T, Shibata Y, Matsuo K. Compositional stability of a salivary bacterial population against supragingival microbiota shift following periodontal therapy. *PLoS One* 2012;**7**:e42806.
16. Mittal S, Bansal V, Garg S, Atreja G, Bansal S. The diagnostic role of Saliva-A Review. *J Clin Exp Dent* 2011;**3**:e314–20.
17. Mager D, Haffajee A, Devlin P, Norris C, Posner M, Goodson J. The salivary microbiota as a diagnostic indicator of oral cancer: A descriptive, non-randomized study of cancer-free and oral squamous cell carcinoma subjects. *J Transl Med* 2005;**3**:27.
18. Farrell J, Zhang L, Zhou H, Chia D, Elashoff D, Akin D, Paster BJ, Joshipura K, Wong DT. Variations of oral microbiota are associated with pancreatic diseases including pancreatic cancer. *Gut* 2012;**61**:582–8.
19. Ahn J, Yang L, Paster BJ, Ganly I, Morris L, Pei Z. Oral microbiome profiles: 16S rRNA pyrosequencing and microarray assay comparison. *PLoS One* 2011;**6**:e22788.
20. Kuczynski J, Lauber CL, Walters WA. Experimental and analytical tools for studying the human microbiome. *Nat Rev Genet* 2011;**13**:47-58.
21. Caporaso J, Lauber C, Walters W, Lyons-Berg D, Lozupone CA, Turnbaugh PJ, Fierer N, Knight R. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *P Natl Acad Sci USA* 2011;**108**:4516–22.
22. Edgar RC. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 2010;**26**:2460-1.
23. Caporaso JG, Bittinger K, Bushman FD. PyNAST: a flexible tool for aligning sequences to a template alignment. *Bioinformatics* 2010;**26**:266-7

- 397 24. Cole JR, Chai B, Marsh TL, Farris RJ. The Ribosomal Database Project (RDP-II):  
398 previewing a new autoaligner that allows regular updates and the new prokaryotic taxonomy.  
399 *Nucleic Acids Res* 2003;**31**:442-3
- 400 25. Price MN, Dehal PS, Arkin AP. FastTree: computing large minimum evolution trees with  
401 profiles instead of a distance matrix. *Mol Biol Evol* 2009;**26**:1641–50.
- 402 26. Wright ES, Yilmaz LS, Noguera DR. DECIPHER, A Search-Based Approach to Chimera  
403 Identification for 16S rRNA Sequences. *Appl Environ Microbiol* 2012;  
404 doi:10.1128/AEM.06516-11.
- 405 27. Vázquez-Baeza Y, Pirrung M, Gonzalez A, Knight R. EMPeror: a tool for visualizing  
406 high-throughput microbial community data. *Gigascience* 2013;**2**:16.
- 407 28. Knights D, Kuczynski J, Charlson ES, Zaneveld J, Mozer MC, Collman RG, Bushman  
408 FD, Knight R, Kelley ST. Bayesian community-wide culture-independent microbial source  
409 tracking. *Nat Methods* 2011;**8**:761–3.
- 410 29. Nadkarni MA, Martin FE, Jacques NA, Hunter N. Determination of bacterial load by  
411 real-time PCR using a broad-range (universal) probe and primers set. *Microbiol-UK*  
412 2002;**148**:257–66.
- 413 30. Thornton B, Basu C. Real-time PCR (qPCR) primer design using free online software.  
414 *Biochem Mol Biol Edu* 2011;**39**:145–54.
- 415 31. Bikandi J, San Millán R, Rementeria A, Garaizar J. In silico analysis of complete  
416 bacterial genomes: PCR, AFLP-PCR and endonuclease restriction. *Bioinformatics* 2004;**20**:798–  
417 9.
- 418 32. Lin IH, Wu J, Cohen, SM, Chen C, Bryk D, Marr M, Melis M, Newman E, Patcher HL,  
419 Alekseyenko AV, Hayes RB, Ahn J. Abstracts 101: Pilot study of oral microbiome and risk of

pancreatic cancer. *Cancer Res* 2013; doi: 10.1158/1538-7445.AM2013-101.

33. Chen T, Yu W-HH, Izard J, Baranova OV, Lakshmanan A, Dewhirst FE. The Human Oral Microbiome Database: a web accessible resource for investigating oral microbe taxonomic and genomic information. *Database* 2010;**2010**:baq013.

34. Galimanas V, Hall MW, Singh N, Lynch MD, Goldberg M, Tenenbaum H, Cvitkovitch DG, Neufeld JD, Senadheera DB. Bacterial community composition of chronic periodontitis and novel oral sampling sites for detecting disease indicators. *Microbiome* 2014;**2**:32.

35. Lazarevic V, Whiteson K, Gaia N, Hernandez D, Farinelli L, Osterås M, François P, Schrenzel J. Analysis of the salivary microbiome using culture-independent techniques. *J Clin Bioinforma* 2012;**2**:4.

36. Zhang Y, Zhang H. Microbiota associated with type 2 diabetes and its related complications. *Food Sci Hum Wellness* 2013;**2**:167-72.

37. Moolya NN, Shetty A, Gupta N, Gupta A, Jalan V, Sharma R. Orthodontic bracket designs and their impact on microbial profile and periodontal disease: A clinical trial. *J Orthod Sci* 2014;**3**:125–31.

38. Ding T, Schloss PD. Dynamics and associations of microbial community types across the human body. *Nature* 2014;**509**:357-60.

39. Grice EA, Segre JA. The human microbiome: our second genome. *Annu Rev Genom Hum G* 2012;**13**:151-70.

40. Schwarzborg K, Le R, Bharti B, Lindsay S, Casaburi G, Salvatore F, Saber MH, Alonaizan F, Slots J, Gottlieb RA, Caporaso JG, Kelley ST. The Personal Human Oral Microbiome Obscures the Effects of Treatment on Periodontal Disease. *PLoS One* 2014;**9**:e86708.

- 443 41. Wang J, Qi J, Zhao H, He S, Zhang Y, Wei S, Zhao F. Metagenomic sequencing  
444 reveals microbiota and its functional potential associated with periodontal disease. *Sci Rep*  
445 2013;**3**:1843.
- 446 42. Eribe E, Olsen I. Leptotrichia species in human infections. *Anaerobe* 2008;**14**:131–7.
- 447 43. Warren RL, Freeman DJ, Pleasance S, Watson P, Moore RA, Cochrane K, Allen-Vercoe  
448 E, Holt RA. Co-occurrence of anaerobic bacteria in colorectal carcinomas. *Microbiome*  
449 2013;**1**:16.
- 450 44. Han YW, Wang X. Mobile Microbiome Oral Bacteria in Extra-oral Infections and  
451 Inflammation. *J Dent Res* 2013;**92**:485-91.
- 452 45. Michaud DS, Izard J, Wilhelm-Benartzi CS, You DH, Grote VA, Tjønneland A, Dahm  
453 CC, Overvad K, Jenab M, Fedirko V, Boutron-Ruault MC, Clavel-Chapelon F, Racine A, Kaaks  
454 R, Boeing H, Foerster J, Trichopoulou A, Lagiou P, Trichopoulos D, Sacerdote C, Sieri S, Palli  
455 D, Tumino R, Panico S, Siersema PD, Peeters PH, Lund E, Barricarte A, Huerta JM, Molina-  
456 Montes E, Dorronsoro M, Quirós JR, Duell EJ, Ye W, Sund M, Lindkvist B, Johansen D, Khaw  
457 KT, Wareham N, Travis RC, Vineis P, Bueno-de-Mesquita HB, Riboli E. Plasma antibodies to  
458 oral bacteria and risk of pancreatic cancer in a large European prospective cohort study. *Gut*  
459 2012;**62**:1764-70.
- 460 46. Evans RT, Klausen B, Sojar HT, et al. Immunization with Porphyromonas (Bacteroides)  
461 gingivalis fimbriae protects against periodontal destruction. *Infect Immun* 1992;**60**:2926–35.
- 462 47. Persson GR, Engel D, Whitney C, Darveau R, Weinberg A, Brunsvold M, Page RC.  
463 Immunization against Porphyromonas gingivalis inhibits progression of experimental  
464 periodontitis in nonhuman primates. *Infect Immun* 1994;**62**:1026–31.
- 465 48. Clark WB, Magnusson I, Beem JE, Jung JM, Marks RG, McArthur WP. Immune

- 466 modulation of *Prevotella intermedia* colonization in squirrel monkeys. *Infect Immun*  
467 1991;**59**:1927–31.
- 468 49. von Bernstorff W, Voss M, Freichel S, Schmid A. Systemic and local  
469 immunosuppression in pancreatic cancer patients. *Clin Cancer Res* 2001;**7**:925-32s.
- 470 50. El-Shinnawi U, Soory M. Associations between periodontitis and systemic  
471 inflammatory diseases: response to treatment. *Recent Pat Endocr Metab Immune Drug Discov*  
472 2013;**7**:169-88.
- 473 51. Feurino LW, Zhang Y, Bharadwaj U. IL-6 stimulates Th2 type cytokine secretion and  
474 upregulates VEGF and NRP-1 expression in pancreatic cancer cells. *Cancer Biol Ther*  
475 2007;**6**:1096-100.
- 476 52. Farrow B, Evers MB. Inflammation and the development of pancreatic cancer. *Surg*  
477 *Oncol* 2002;**10**:153–69.
- 478 53. Mitsuhashi K, Nosho K, Sukawa Y, Matsunaga Y, Ito M, Kurihara H, Kanno S, Igarashi  
479 H, Naito T, Adachi Y, Tachibana M, Tanuma T, Maguchi H, Shinohara T, Hasegawa T, Imamura  
480 M, Kimura Y, Hirata K, Maruyama R, Suzuki H, Imai K, Yamamoto H, Shinomura Y.  
481 Association of *Fusobacterium* species in pancreatic cancer tissues with molecular features and  
482 prognosis. *Oncotarget* 2015;**30**:7209-20.


485

**Figure 1**(on next page)

Relative abundance of phyla identified in patient saliva summarized by diagnosis group.

Relative abundance of phyla in oral communities from 108 study patients summarized by diagnosis group (H= healthy control, O= other disease, and P= pancreatic cancer).






## Figure 2 (on next page)

Mean relative abundances of particular genera in pancreatic cancer patients (P) compared to healthy (H) and other disease (O) patient groups.


Relative abundances of genera in oral communities from 108 patients. Arrows point to specific genera that showed interesting trends across diagnosis groups.[b]



# Figure 3 (on next page)

Abundance ratio of *Leptotrichia* to *Porphyromonas* between different patient categories.


Each symbol represents the ratio of *Leptotrichia* Oral Taxon 221 and *Leptotrichia hongkongensis* to *Porphyromonas* for an individual patient (n=108). Patients are grouped into 3 different categories depending on their diagnosis: healthy control (H), other diseases (including cancer) (O), and pancreatic cancer (P). Horizontal bar and error bars represent the mean and SEM, respectively. \*\*\*p<0.001 (Kruskal-Wallis test followed by Dunn's multiple-comparison test).



# **Figure 4**(on next page)

Correlation between *Leptotrichia* abundance from 16S rRNA sequences and from real-time qPCR.

Cross validation of total *Leptotrichia* OTU abundance using real-time qPCR. After using 16S rRNA as a reference gene for normalization of the levels of *Leptotrichia* genus, data was normalized by fold change to three healthy controls with relatively low *Leptotrichia* OTU abundance. Each symbol represents a patient: P=6, and O=12. *Leptotrichia* OTU abundance was correlated with qPCR fold change according to Pearson's correlation ( $r=0.903$ ).



**Table 1**(on next page)

Clinical characteristics of study sample (n=108)



Demographics	Pancreatic Cancer (P) n=8	Other Disease (O) n=78	Healthy Control (H) n=22	Total n=108
Sex				
Male	6	38	12	56
Female	2	40	10	52
Ethnicity				
Caucasian	6	56	15	77
Hispanic	2	6	5	13
Asian	0	4	1	5
Unknown	0	12	1	13

1

**Table 2**(on next page)

Types of identified cancers (n=23)

Cancer by Category	Age Mean	N
Pancreas (n=8)	71.1	
Pancreatic Cancer		8
Digestive (n=9)	64.7	
Ampullary		3
Esophageal		3
Stomach		1
Rectal		2
Non-Digestive (n=6)	54.8	
Breast		1
Skin		1
Testicular		1
Thyroid		3