

Identification and characterization of ARF family members involved in Fig (*Ficus carica* L.) fruit development

Haomiao Wang¹, Hantang Huang¹, Yongkai Shang¹, Miaoyu Song^{Corresp., 1}, Huiqin Ma^{Corresp. 1, 2}

¹ College of Horticulture, China Agricultural University, Beijing, Beijing, China

² State Key Laboratory of Agrobiotechnology, China Agricultural University, Beijing, Beijing, China

Corresponding Authors: Miaoyu Song, Huiqin Ma
Email address: songmiaoyu@cau.edu.cn, hqma@cau.edu.cn

Auxin response factor (ARF) combine with AuxREs cis-acting elements in response to auxin to regulate plant development. To date, no comprehensive analysis of ARF genes expressed during fruit development has been conducted for common fig (*Ficus carica* L.). In this study, members of the FcARF gene family were screened, identified in the fig genome database and their features characterized using bioinformatics. Twenty FcARF genes were clustered into three classes, with almost similar highly conserved DBD (B3-like DNA binding domain), AUX/IAA (auxin/indole-3-acetic acid gene family) and MR domain structure among class members. Analysis of amino acid species in MR domain revealed 10 potential transcription activators and 10 transcription inhibitors, and 17 FcARF members were predicted to be located in the nucleus . DNA sequence analysis showed that the ARF gene family consisted of 4 - 25 exons, and the promoter region contained 16 cis-acting elements involved in stress response, hormone response and flavonoid biosynthesis. ARF genes were expressed in most tissues of fig, especially flower and peel. Transcriptomics analysis results showed that *FcARF2*, *FcARF11* and *FcARF12*, belonging to class-Ia, were stably and highly expressed in the early development stage of flower and peel of 'Purple peel' fig . However, their expression levels decreased after maturity. Expression of class-Ic member *FcARF3* conformed to the regularity of fig fruit development . These 4 potential transcription inhibitors may regulate fruit growth and development of 'Purple peel' fig. This study provides comprehensive information on the fig ARF gene family, including gene structure, chromosome position, phylogenetic relationship and expression pattern. Our work provides a foundation for further research on auxin-mediated fig fruit development.

Identification and characterization of ARF family members involved in Fig (*Ficus carica* L.) fruit development

Haomiao Wang¹, Hantang Huang¹, Yongkai Shang¹, Miaoyu Song¹, Huiqin Ma^{1,2}

¹ College of Horticulture, China Agricultural University, Beijing, China,

² State Key Laboratory of Agrobiotechnology, China Agricultural University, Beijing, China

Corresponding Author:

Miaoyu Song¹, Huiqin Ma¹

West Yuanmingyuan Road, Beijing, Country Email address:

Email address: songmiaoyu@cau.edu.cn; hqma@cau.edu.cn

Abstract: Auxin response factor (ARF) combine with AuxREs cis-acting elements in response to auxin to regulate plant development. To date, no comprehensive analysis of ARF genes expressed during fruit development has been conducted for common fig (*Ficus carica* L.). In this study, members of the FcARF gene family were screened, identified in the fig genome database and their features characterized using bioinformatics. Twenty FcARF genes were clustered into three classes, with almost similar highly conserved DBD (B3-like DNA binding domain), AUX/IAA (auxin/indole-3-acetic acid gene family) and MR domain structure among class members. Analysis of amino acid species in MR domain revealed 10 potential transcription activators and 10 transcription inhibitors, and 17 FcARF members were predicted to be located in the nucleus. DNA sequence analysis showed that the ARF gene family consisted of 4 - 25 exons, and the promoter region contained 16 cis-acting elements involved in stress response, hormone response and flavonoid biosynthesis. ARF genes were expressed in most tissues of fig, especially flower and peel. Transcriptomics analysis results showed that *FcARF2*, *FcARF11* and *FcARF12*, belonging to class-Ia, were stably and highly expressed in the early development stage of flower and peel of 'Purple peel' fig. However, their expression levels decreased after maturity. Expression of class-Ic member *FcARF3* conformed to the regularity of fig fruit development. These 4 potential transcription inhibitors may regulate fruit growth and development of 'Purple peel' fig. This study provides comprehensive information on the fig ARF gene family, including gene structure, chromosome position, phylogenetic relationship and expression pattern. Our work provides a foundation for further research on auxin-mediated fig fruit development.

Key words: *Ficus carica* L.; Auxin response factor; Gene family; Expression analysis; Fruit development

Introduction

Auxin is an important plant hormone, which directly regulates cell growth, division and specific differentiation, participating in embryogenesis, flower and fruit development, root formation and vascular bundle development (Du et al. 2012). At the molecular level, auxin regulates plant growth and development via a signal transduction pathway mediated by Aux/IAA proteins and auxin response factor (ARF) (Liu et al. 2015; Woodward et al. 2005). ARFs typically consist of three domains: an N-terminal B3-like DNA binding domain (DBD), a C-terminal dimerization domain (CTD), which forms a heterodimer with Aux/IAA family genes, and the middle domain (MR) with an activation (AD) or inhibition (RD) role (Guilfoyle & Hagen 2007; Kim et al. 1997; Li et al. 2015). The DBD domain can identify Auxin Response Element (AuxRE, ‘TGTCTC’) in auxin response gene promoter. The CTD of ARF protein contains two conserved motifs, motif III and motif IV, which facilitate formation of homo- and hetero-dimerization among ARFs and Aux/IAAs (Zhou et al. 2017). The MR domain either activates or inhibits transcription depending on its amino acid composition (Ye et al. 2020). Therefore, these domains are necessary for efficient regulatory function of ARF (Wang et al. 2007).

The ARF gene family has been identified in many plants. There are 23 ARFs in *Arabidopsis thaliana* (Okushima et al. 2005), 12 in *Fragaria vesca* (Wang et al. 2019), 19 in *Vitis vinifera* (Wan et al. 2014), 19 in *Citrus sinensis* (Li et al. 2015), 19 in *Capsicum annuum* L. (Zhang et al. 2017), 22 in *Lycopersicon esculentum* (Kumar et al. 2011), 25 in *Oryza sativa* L. (Wang et al. 2007) and 37 in *Populus alba* × *P. glandulosa* (Yang et al. 2014). Based on evolutionary relationship analysis, ARF genes in *Arabidopsis thaliana* cluster into three classes. Class I is the largest with 15 members belonging to three classes. Members in class-Ia are mainly involved in regulating flower and root development. For example, *AtARF1* and *AtARF2* promote flower senescence and abscission (Ellis et al. 2005), whereas *AtARF11* promotes lateral root formation (Feng et al. 2012). These genes share a high degree of similarity among their amino acid and nucleotide sequences. In addition, *AtARF23* is a pseudogene (Okushima et al. 2005). Class-Ic has two members, *AtARF3* and *AtARF4*, which regulate floral organogenesis and the timing and pattern of development (Ellis et al. 2005; Harper et al. 2000; Hunter et al. 2006; Pekker et al. 2005; Sessions et al. 1997). Members of class-II are ARF^{ClassA} transcription activators, which regulate auxin-mediated flower development, morphogenesis, lateral root formation and other processes (Truskina et al. 2021). Class-III consists of three most structurally distinct members that regulate pollen wall pattern formation, root cap formation or seed germination (Ellis et al. 2005; Guilfoyle & Hagen 2007).

As important plant growth regulators, ARF genes are expressed and play regulatory roles in different tissues (**Table S1 and Table S2**), especially floral organs and fruit. *AtARF2* was found to promote the development of stamens and participate in the senescence and abscission of floral organs (Okushima et al. 2005). *AtARF4* regulates flower patterns and plays a significant role in organ polarity development (Hunter et al. 2006; Liu et al. 2014; Pekker et al. 2005). *AtARF6/8*

and *SLARF6/8* can promote maturation of floral organs and regulate fruit and seed carpel development (Goetz et al. 2006; Wu et al. 2006). In fruiting plants, *SLARF3* plays multiple roles in tomato fruit development and participates in the formation of epidermal cells and trichomes (Zhang et al. 2015). *SLARF9* regulates cell differentiation during early tomato fruit development and *SLARF4* controls sugar metabolism during maturation (de Jong et al. 2015; Sagar et al. 2013). Six CiARF genes (*CiARF1/2/6/7/12/18*) are highly expressed in fruit receptacle, where they regulate the citrus pulp and peel development (Li et al. 2016). Seven PpARF genes (*PpARF1/2/3/4/5/6/9*) perform similar functions in *Prunus persica* (Hou et al. 2021). Studies on apples show that *MdARF5* induces ethylene synthesis by directly promoting the expression of ethylene synthesis genes *MdACS3a*, *MdACSl* and *MdACO1*, which then promote apple fruit ripening (Yue et al. 2020).

ARF proteins are negatively regulated by auxin/indole-3-acetic acid (Aux/IAA) proteins. The potential for interaction between ARF and Aux/IAA proteins is high as their C-terminal domains are homologous (Shen et al. 2010). Aux/IAA genes play a major role in regulating many auxin processes in plants. Aux/IAA genes exist as a family in many plants and are associated with leaf curl patterns, plant phototropism and tropism, plant height, and especially lateral root formation (Reed 2001). In *Arabidopsis thaliana*, *axr5 / iaa1* mutant showed auxin resistance and loss of root and stem geotropism and hypocotyl phototropism (Yang et al. 2004). Mutations in *AUXIN RESISTANT1 (AUX1)/LIKE-AUX1 (LAX)* genes result in auxin-related developmental defects and have been implicated in regulating root development, female gametophyte development and embryo development in *Arabidopsis thaliana* (Swarup & Bhosale 2019). *AtIAA8* regulates lateral root formation by interacting with auxin receptor *AtTIR1* and ARF transcription factors in the nucleus (Arase et al. 2012). *MdARF1* regulates anthocyanin accumulation in fruit by interacting with *MdIAA121* in apples (Wang et al., 2018).

As an important plant growth regulator, the role of ARF gene family in fruit ripening and development has been investigated in many species but not in fig fruits. With its origins in the Mediterranean coastal region, the common fig (*Ficus carica* L.) is one of the widely cultivated ancient fruit trees in the world. The fig fruit (syconia), an aggregated fruit formed by the expansion of female flower tissue and receptacle, has high antioxidant activity, great taste and high nutritional value (Song et al. 2021). The fig fruit exhibits a typical double sigmoid growth curve, including two rapid growth phase (I and III) separated by slower growth phase (II) (Song et al. 2021). The fig has the advantages of short juvenile period, easy vegetative growth and reproduction, and is gradually becoming a new model plant for studying fruit development mechanisms (Wang et al. 2018). Presently, the genome-wide characterization of fig auxin signal transduction-related components and their expression patterns has not been carried out. In this study, genome-wide data of ‘Horaishi’ and ‘Dottato’ (Mori et al. 2017; Usai et al. 2020) and ‘Purple Peel’ fig transcriptome data at different developmental stages (NCBI acceptance No. PRJNA723733) were used to predict ARF genes. In total, 20 ARF genes were identified in common fig, and their features, including chromosome position, phylogenetic relationship, gene structure and motif characteristics, characterized. In addition, we described the transcriptional accumulation patterns of ARF genes in different tissues, including at different developmental

and maturation stages of female flowers and peel. Our study provides a good basis for further exploring the role of *FcARFs* and characterizing their functions.

Materials & methods

2.1 Plant materials

The fig cultivar ‘Purple Peel’ plants were established at Shangzhuang Experimental Station of China Agricultural University, Beijing, China. The roots, stems, leaves of ‘Purple Peel’, and fruits at different developmental stages were collected for qRT-PCR analysis. Fruits were sampled at six stages based on the characteristics of fruit development. Stage 1 represented phase I (the first rapid growth period), stages 2, 3, and 4 were the early, middle, and late stages of phase II (slow growth period), and stages 5 and 6 represented phase III (the second rapid growth period). At each stage, 50 fruits were sampled. The peel and flower were separated onsite during sampling. Fresh samples were quick-frozen with liquid nitrogen and stored at -80°C until RNA extraction.

2.2 Identification of ARF transcription factor family in fig

Genome sequences of *cv. Horaishi* and *cv. Dottato* fig were downloaded from the National Center for Biotechnology Information (NCBI) (<https://www.ncbi.nlm.nih.gov/genome/?term=Figus+carica>) (Mori et al. 2017; Usai et al. 2020). The sequences were blasted (E-value = -5) using the hidden Markov model (HMM) files of B3, Auxin_resp and AUX_IAA domains in Pfam (<http://pfam.xfam.org/search/sequence>) with entry numbers PF02362, PF06507 and PF02309, respectively (El-Gebali et al. 2019). Candidate genes containing known conserved domains were retained, and further screened for ARF domains against three databases: Pfam, NCBI conserved domains (<http://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi>) and SMART (<http://smart.embl.de>) (Letunic et al. 2021). ARF family genes in *Arabidopsis thaliana* were downloaded from the Arabidopsis database (TAIR; <https://www.Arabidopsis.org/>). Fig ARF homologs were compared with Arabidopsis ARFs using BLASTP with default parameters to obtain annotation and grouping information. *FcARFs* sequences were analyzed using bioinformatics methods, and the physicochemical parameters, MW, pI and GRAVY of translated proteins were calculated using ExPASy (http://www.expasy.ch/tools/pi_tool.html) (Guo et al. 2014).

2.3 Multiple sequence alignment and phylogenetic analysis

Using Clustal 2.0, multiple-sequence alignments of amino acid sequences of predicted ARF genes in fig was carried out, with default parameters (Song et al. 2021). ClustalX 2.11 was used to analyze phylogenetic relationships among ARF gene families in Arabidopsis, fig, apple, grape and strawberry. (Liu et al. 2015; Luo et al. 2014; Sun et al. 2015a). The phylogenetic tree of *FcARFs* was constructed using MEGA 7.0 via the Neighbor-Joining method with calibration test parameter “Bootstrap = 1000 (1000 replicates)” (Tamura et al. 2011; Kumar et al. 2016). The analysis involved 87 amino acid sequences and ARF gene families in five species including fig were divided following the division of ARF gene families in *Arabidopsis thaliana*. Percentage confidence values were shown on branches and the evolutionary distances used to infer the phylogenetic tree via Poisson correction method (Zuckerkandl & Pauling 1965). Gaps were treated as a complete deletion. There were a total of 1619 positions in the final dataset.

2.4 Gene structure and conserved motif analysis

FcARFs gene sequences were downloaded from fig genome database(<https://www.ncbi.nlm.nih.gov/gen-ome/?term=Ficus+carica>), and the intron/exon structure map of the fig ARFs was generated online using the Gene Structure Display Server (GSDS: <http://gsds.cbi.pku.edu.ch>) (Guo et al. 2007). *FcARFs* amino acid sequences were referenced to fig genome database protein RefSeq. Because ARF family genes contain highly conserved domains, DNAMAN was used for multiple sequence alignment of *FcARF* protein sequences. The length and sequence information of other conserved motifs except the ARF conserved domain were obtained using the online MEME4.11.2 tool (<http://meme.sdsc.edu/meme/itro.html>) with the following parameters: number of repetitions “any,” highest motif number “15,” motif length “6–100,” and default values for the other parameters.

2.5 Chromosome mapping and collinearity analysis

Tandem replication of ARF genes usually refers to adjacent homologous genes on the same chromosome, whereas gene duplication events are regarded as repeated homologous genes on different chromosomes (Liu et al. 2011). Whether *FcARF* gene is considered an important event of tandem repeat between genes depends on its location on the chromosome. To study the distribution of ARF genes on chromosomes, their chromosomal location and annotation information were utilized. An openly available database and Map Draw software were used to generate physical location information for homolinear analysis of *ARF* genes in fig. The genome data of *Ficus hispida* and *Ficus microcarpa* were downloaded from the National Genomics Data Center database (<https://bigd.big.ac.cn/search/?dbId=gwh&q=PRJCA002187&page=1>) (Zhang et al. 2020). An interspecies collinearity analysis of ARFs in fig, *F. hispida*, and *F. macrocarpa* was performed using MCscanX and TBtools (Song et al., 2021). The final map was generated with Circos version 0.63 (<http://circos.ca/>). The non-synonymous replacement rate (Ka) and synonymous replacement rate (Ks) of replicated gene pairs were calculated using KaKs_Calculator 2.0 (Wang et al., 2010), with Ka/Ks ratio taken as environmental selection pressure.

2.6 Protein functional connection network

Since fig species was not included in STRING database, the search species was set to *Arabidopsis thaliana*. The interaction networks of 20 ARF proteins were analyzed using the STRING protein interaction database (<http://string-db.org/>), which is an online protein-protein interaction networks functional enrichment analysis prediction site (Szklarczyk et al. 2021). Protein-protein interaction networks were built with an interaction score of -1 to 1. In total, 20 fig ARF proteins were aligned with AtARF proteins using Blastp alignment with E-value of 1e-0 (Rosas-Camacho 2009, Chen et al. 2015). The interaction network of *FcARF* protein was constructed using homologous and *Arabidopsis thaliana* ARF protein, and genes of interest were visualized using Cytoscape software (Doncheva et al. 2019).

2.7 Expression analysis of *FcARFs* using RNA-seq and qRT-PCR

TBtools was used to analyze the expression patterns of *FcARFs* in ‘Purple-Peel’ fig fruit based on RNA-seq library (NCBI Accession No. PRJNA723733). Their peel and female flower tissue

were collected at six stages of fruit development and significant differential expression was determined with $p < 0.05$ and $|\log_2(\text{fold change})| \geq 1$. Different tissues and fruits were sampled from ‘Purple Peel’ plant at six stages for qRT-PCR. The peel and flower were separated during sampling. Fresh samples were quick-frozen with liquid nitrogen and stored at -80°C for RNA extraction. Total RNA was extracted using the CTAB method and reversed transcribed using reverse transcription kit (TaKaRa, Dalian, China). The specific primers for qRT-PCR are shown in **Table S5**. The design parameters for the primers include amplicon length, 100-150 bp; primer length, 15-25 bp; melting temperature (T_m), 57-61 $^\circ\text{C}$; GC base content, 40-60%; and excluding conserved domain (Li et al. 2019). Following a previously published method (Song et al. 2021), all qRT-PCR reaction volumes were set at 10 μL . The cycle parameters were as follows: pre-denaturation at 94 $^\circ\text{C}$ for 15 s, annealing at 60 $^\circ\text{C}$ for 30 s and extension at 72 $^\circ\text{C}$ for 1 min, for 40 cycles. qRT-PCR reaction for each gene was replicated three times. *Fc18S* and *FcActin* were used as an internal standard to calculate relative fold differences based on comparative cycle threshold values. We calculated relative expression levels using the comparative C_t ($2^{-\Delta\Delta C_t}$) method. SPSS 26.0 software was used for significance analysis.

Results

3.1 ARF family members identified in fig

Twenty ARF genes were identified in published fig genome and named *FcARF1* to *FcARF20* following the equivalent classification for *Arabidopsis thaliana* (**Figure 1A and Table 1**). Detailed information for each candidate gene, including gene name, gene ID, number of exons, protein length, protein molecular weight (kDa), isoelectric point (pI), instability coefficient, Aliphatic index, GRAVY and location in cells, is shown in **Table 1**. The length of amino acids encoded by *FcARFs* ranged from 588 (*FcARF10*) to 1150 (*FcARF7*), averaging 782.95 amino acids. The number of exons and predicted molecular weight ranged from 3 (*FcARF10*) to 17 (*FcARF18*) and 52.73 (*FcARF4*) to 127.06 kDa (*FcARF7*), respectively. The isoelectric point (pI) ranged from 5.03 (*FcARF5*) to 9.52 (*FcARF11*), and acidic amino acids were enriched in the protein molecules. In total, 80.95% of fig ARF family proteins had isoelectric points less than 7, with most of them being weakly acidic, consistent with pI of ARF proteins in citrus and strawberry (Li et al. 2015; Wang et al. 2019). On average, the total hydrophilicity of ARF family proteins in fig was negative, ranging from -0.726 (*FcARF11*) to -0.322 (*FcARF17*), which indicated that these proteins were hydrophilic. The instability index ranged from 42.05 (*FcARF17*) to 72.35 (*FcARF15*), suggesting that all these proteins were stable ($II > 40$). The Aliphatic index ranged from 63.06 (*FcARF11*) to 84.46 (*FcARF18*). Twenty *FcARF* proteins were localized in subcellular structures by online software Target P and CELLO v2.5. The results showed that they were mostly distributed in the nucleus and cytoplasm.

3.2 Phylogenetic analysis of FcARFs

In total, 87 ARF proteins comprising 23 from *Arabidopsis thaliana*, 20 from fig, 12 from strawberry, 19 from grape and 13 from apple were obtained from NCBI database to construct a phylogenetic tree using Neighbor-Joining approach. *FcARF* proteins were divided into three

classes (**Figure 1**). Class-I was further divided into three classes, including class-Ia with the following five members: *FcARF1*, *FcARF2*, *FcARF9*, *FcARF11* and *FcARF18*. Class-Ib included three members (*FcARF12*, *FcARF14* and *FcARF20*) whereas class-Ic included two (*FcARF3* and *FcARF4*). Class-II had six members and class-III had four. Compared with the equivalent *Arabidopsis* class, the number of ARF members in fig class-Ib was less. These results indicated that gene loss or gain occurred during the evolution of ARF family genes in fig.

3.3 Analysis of ARF gene family structure

Typical ARF protein has three conserved domains: N-terminal B3-like DNA binding domain (DBD), C-terminal dimerization domain (CTD), and a middle region (MR) (Xing et al. 2011a). **Figure 2** shows that all FcARFs contained MR as an intermediate domain. All proteins contained a complete CTD structure except class-III members. Similarly, DBD structure existed in all FcARFs proteins except *FcARF18*. The branches of the same phylogenetic tree always had similar structures, but some genes in the same class exhibited different gene structures. Among them, four FcARFs (*FcARF17*, *FcARF10*, *FcARF13* and *FcARF16*) had truncated CTD domains distributed in class-III of the phylogenetic tree (**Figure 2A**). To exhaustively verify the accuracy of FcARF conserved domains, the amino acid sequence alignment results of FcARFs and identified the location of conserved domain were aligned using DNAMAN software (**Figure S1**). The active MR domain has been found to be rich in glutamine (Q) whereas the inhibitory domain is rich in proline (P), serine (S) and threonine (T) (Wu et al., 2011; Tiwari et al., 2003).

Considering these characteristics, we analyzed the amino acid sequence and composition of the MR domain (**Figure S2**). Subsequently, we predicted 10 potential transcription activators (*FcARF4*, *FcARF5*, *FcARF7*, *FcARF9*, *FcARF10*, *FcARF13*, *FcARF16*, *FcARF17*, *FcARF18* and *FcARF19*) and 10 inhibitors (*FcARF1*, *FcARF2*, *FcARF3*, *FcARF6*, *FcARF8*, *FcARF11*, *FcARF12*, *FcARF14*, *FcARF15* and *FcARF20*) (**Figure 2B**).

3.4 Conservative FcARFs motif and cis-acting element analysis

The exon-intron structures of genes in the same gene family are similar, reflecting the close evolutionary relationship among all its members. The number of introns in *FcARF* ranged from 3 to 17 (**Figure 3A**). *FcARF18* gene in class-Ia had the most introns (17), whereas class-III members contained the least, suggesting that members have experienced intron loss during evolution. Conserved motifs 1-15 of *FcARFs* were analyzed (**Table S3**, **Figure S3**). Motif 2 and motif 8 belonging to DBD and MR domains, respectively, were highly conserved in all FcARF proteins. Our analysis of conserved motifs in the database revealed that conserved motifs in FcARF are unstable, and their positions had changed except motif 2 and motif 8. In addition, motifs 15 only existed in specific genes, which agrees with previously reported results (Wei et al. 2016). This indicates that these motifs may be essential for specific regulatory functions of these genes.

We found 2 kb sequence upstream of the initiation codon of *FcARF* gene in PLACE database that predicted many cis-acting elements larger than 4 bp. TGA-element and TGA-box were associated with auxin response and transcriptional activation. There were at least 16 cis-acting regulatory elements in the promoter region of *FcARF* family members in fig (**Figure 3B**). They were involved in abiotic stress responses, including light response, anaerobic inducibility, anoxic

specific inducibility, defense and stress response, drought induction, low temperature responsive and wound response. In addition, these cis-acting regulatory elements were involved in hormone response, (abscisic acid response, auxin response, gibberellin response, MeJA response and salicylic acid response), circadian control, flavonoid biosynthesis regulation, meristem expression and other regulatory elements. Drought stress-related MYBHv1 binding sites were found in the promoter regions of three ARF genes (*FcbARF2*, *FcARF14* and *FcARF18*).

3.5 Chromosome localization and distribution of FcARFs

Distributions of ARF transcription factors in fig linkage groups are shown in **Figure S4**. *FcARF1* ~ *FcARF20* were unevenly distributed in 12 chromosomes, apart from chromosomes 2 and 4. Chromosomes 1, 6, 7, 10, 11 and 12 had one ARF gene each, which were *FcARF6*, *FcARF7*, *FcARF5*, *FcARF9*, *FcARF14* and *FcARF20*, respectively. Chromosomes 3, 5 and 9 had 3 ARF genes each, which were the most abundant. Tandem replication occurs when the distance between genes is less than 100 kb and exists on the same chromosomes, only *FcARF3* and *FcARF16* on Chr3 occurred. The *FcARF* gene family had 3 pairs of genes from fragment replication (**Figure 4A**). During intraspecific evolution, the Ka/Ks values can explain positive selection ($Ka/Ks > 1$), neutral selection ($Ka/Ks = 1$) and negative selection ($Ka/Ks < 1$). Nineteen pairs (43.18%) of ARF genes in fig species had Ka/Ks greater than 1, while 25 pairs (56.82%) had Ka/Ks less than 1 (**Table S4**), indicating the existence of 2 evolutionary modes of positive and negative selection in fig, with negative selection being the main evolutionary mode. Collinearity analysis identified 18 separate orthologous genes between *F. carica* and *F. macrocarpa* as well as between *F. carica* and *F. hispida*, indicating that evolutionary distance between fresh fig and two evergreen *Ficus* species was similar (**Figure 4B**). *FcARF20* did not exhibit a collinear relationship with either of the two *Ficus* species, implying that there might be a unique ARF in Fig evolution.

3.6 Protein interaction network of FcARFs

The ARF protein interacts with other proteins, such as IAA and AUX1 proteins, to regulate downstream gene expressions and exert biological functions. Correlation analysis of ARF proteins and interacting proteins in figs, as well as the gene expression correlation network (Song et al 2022), was analyzed based on gene expressions in the developmental stages of fig fruits (**Figure 5B**). A protein interaction network was constructed by STRING, which included 5 ARF proteins and 9 other proteins (5 IAA proteins). Expressions of the five *FcARFs* were positively correlated with each other, co-expression scores were all above 0.7, with *FcIAA1*, *FcBZR1* and *FcAUX1* being negatively correlated with expressions of the five *FcARFs*.

In **Figure 5A**, thickness of lines represents protein interaction strengths. IAA8 interacted with five *FcARFs*, and their expressions were positively correlated, including *FcARF15*-IAA8. In *Arabidopsis*, interactions of *AtARF15* and *AtIAA8* proteins respond to early auxin responses and cooperatively regulates lateral root growth, development as well as morphogenesis (Arase et al. 2012). There was a strong interaction between *ARF6*-*PIF4*, with an expression correlation coefficient of 0.808, implying a positive correlation. Direct inhibition of auxin-responsive gene expressions mediated by interactions of *AtPIF4* and *AtARF6* in *Arabidopsis* constitutes a new layer in regulatory mechanisms of photoinhibition of auxin-induced hypocotyl elongation (Oh et

al. 2014). IAA1 was predicted to interact with FcARF8 and FcARF19, and expression correlation indices were -0.582 and -0.571, respectively, implying negative correlations. This is consistent with the prediction that IAA interacts with ARF and negatively regulates its expressions. In the regulatory network composed of fig ARFs, there were strong positive correlations among FcARFs levels, indicating functional complementarity and redundancy among them. The expression levels and interaction coefficients predict that FcIAA8 is a key positive cooperative factor in the network, while IAA1 is a negative regulatory cooperative factor.

3.7 Expressions of *FcARFs* in different tissues

Expression patterns of 20 FcARFs in fig's root (RO), stem (ST), old leaf (OL), young leaf (YL), flower (FL) and peel (PE) were analyzed by qRT-PCR (**Figure 6**). Twenty *FcARFs* showed tissue and organ expression specificity. In class-I, *FcARF1/12/14/18/20* were highly expressed in female flowers and peel, the major fruit components. *FcARF 2/3/4/9/11* were highly expressed in the stem. In class-II, *FcARF 5/8/15* were highly expressed in female flowers and peel while *FcARF 6/7/19* were highly expressed in the stem. In class-III, all FcARFs (*FcARF 10/13/16/17*) were highly expressed in female flowers and peel. These findings imply that expressions of these *FcARF* genes are high in reproductive organs, but low in vegetative organs, such as leaves (**Figure 6**).

3.8 Expressions of *FcARFs* during fig fruit ripening

Expressions patterns of *FcARFs* in 'Purple peel' fig peel and flower tissues at different growth stages were constructed by TBtools (**Figure 7**). Based on FPKM values, they were divided into 4 groups, among which 4 members greater than 100 came from 3 classes, respectively. Class-I had 4 members (*FcARF2*, *FcARF3*, *FcARF11*, and *FcARF12*) expressed in early stages of flower and peel development. The 50-100 group consists of 3 members (*FcARF1*, *FcARF8* and *FcARF18*) from class-Ia and class-II. Expression patterns of *FcARF1* and *FcARF18* were consistent, being highest in early stages of female flower development. The rest of the members with FPKM values between 10 and 50.

During the 6 growth stages of female flowers and peel, *FcARF3* from class-Ic was stably expressed, with FPKM values > 50. In addition, expressions of *FcARF3* in fast-growing phase I and III were higher than those in slow-growing phase II. Expressions of class-Ia members *FcARF2/9/11* were highest in the early developmental stages of fruits and were decreased after maturity. *FcARF11* levels were highest at P1, P2 and P4, while *FcARF9* levels were highest at P3. In this study, *FcARF* gene expressions in peel and flower tissues of fig fruit at 6 different stages of ripening levels were evaluated by qRT-PCR (**Figure S5 and S6**). Expression profiles of *FcARF* in female flowers and peel tissues obtained from transcription sequencing data should be verified.

Discussion

To establish the significance of ARF genes in the auxin signal transduction pathway in fig, we evaluated the main structural characteristics of the ARF gene family in fig. Using genome sequencing results, we isolated and identified 20 fig ARF genes.

Phylogenetic tree and protein structure predictions of ARF protein function

Phylogenetic trees have been used to predict the functions of different genes (Pandey et al. 2018). **Figure 1** shows that fig is highly homologous to *Arabidopsis thaliana*, a model plant, with 6 sister pairs. Phylogenetic tree analysis of ARF proteins in *Arabidopsis* and fig can predict the functions of FcARFs by clarifying their phylogenetic relationships. For instance, *FcARF3* and *FcARF4* of fig belong to class-Ic with *AtARF3* and *AtARF4*, which may be involved in regulation of flower formation, development and patterns. Class-Ia, which consists of *FcARF1*, *FcARF2*, *FcARF9*, and *FcARF11* may have a role in regulating leaf senescence, floral organ shedding and auxin homeostasis. *FcARF10*, *FcARF13*, *FcARF16* and *FcARF17* belong to class-III with *AtARF10*, *AtARF16* and *AtARF17*, which may be involved in regulation of pollen tube wall formation, root cap formation and seed germination.

Apart from *FcARF18*, which does not have the DBD domain, other FcARFs (*FcARF1* ~ *17*, *19,20*) have DBD and MR domains that determine their functions based on amino acid compositions (Sun et al. 2015b; Wang et al. 2007; Okushima et al. 2005). Through domain comparisons (**Figure S3** and **Table S3**), it can be found that *FcARF18* lacks motif 2 on the DBD domain, which inhibits its ability to recognize and bind DNA by forming a helix-turn-helix domain (Tiwari et al. 2003; Roosjen et al. 2018). In addition, most plant ARF gene families have an ARF protein without the CTD domain. All class-III FcARFs have truncated CTD domains, which is similar in *Arabidopsis thaliana* and other species (**Figure 2A**). Responses and signal transduction of plant cells to auxin are mainly mediated by the TIR1/AFB/-Aux/IAA-ARF signaling pathway in the nucleus. ARF participates in the auxin signaling pathway by forming a heterodimer with CTD regions of the Aux/IAA protein. The C-terminal domains of proteins are homologous. Without a complete CTD domain, *FcARF10/13/16/17* may not participate in the auxin signaling pathway or perform its independent functions (Liu et al. 2011).

The ARF genes are transcriptional activators or inhibitors. The activation domain (AD) in glutamine (Q) and leucine (L) in the middle region of the ARF protein can act as a transcription activator, while the repression domain (RD) rich in proline (P), serine (S) and threonine (T) can act as a transcription inhibitor (Tiwari et al. 2003; Zouine et al. 2014). In this study, the intermediate region of transcription factors in class-Ia and class-II was found to be rich in glutamine (Q), and it is postulated that class-II members have transcriptional activation functions. *AtARF5*, *AtARF6*, *AtARF7*, *AtARF8* and *AtARF19* in *Arabidopsis thaliana*, which are homologous to fig class-II, are ARF^{ClassA} transcription activators. Middle regions of classes-I and -III are rich in proline (P), serine (S) and threonine (T). Therefore, members of class-Ia and class-II inhibit transcription. Our findings should be verified by studies of regulatory mechanisms (Roosjen et al. 2018).

Evolution and Gene Structures of the ARF Family

Gene replication, including tandem repeat and large fragment replication, is the most important mechanism of gene recombination and amplification (Vision et al. 2000). Tandem repeat refers to two adjacent genes on the same chromosome, while large fragment replication events occur on different chromosomes, which is an important way of gene expansion and species evolution (Liu et al. 2011; Xing et al. 2011b). Compared to other plants, there are 20 ARF genes in fig, which

are less than those in other plants, such as 23 in *Arabidopsis thaliana*, 25 in rice and 39 in poplar. Combined with collinearity analysis, there are probably no large-scale gene repetition events in fig evolution. Collateral gene pairs (*FcARF13* and *FcARF10*) were located on chromosome 1, while *FcARF19*, *FcARF20*, *FcARF13* and *FcARF17* were located on chromosomes 9, 11, 13 and 3, respectively. The former belongs to tandem replication, while the latter belongs to large fragment replication. In this study, 3 (15%) of 20 FcARFs were established to be involved in tandem repeat or large fragment replication events, while 33% of MdARFs in apples were associated with gene replication (Luo et al. 2014). Even though there are few gene replication events in fig, we postulated that gene replication is the main mode of ARF gene family amplification, implying that different types of ARFs gene family members have different evolutionary paths and rates.

FcARFs are mainly involved in plant hormone responses, growth regulation, light response and environmental stress responses. As trans-acting factors, hormones can bind hormone response elements in promoter sequences, thereby regulating the transcriptional activities of target genes, and can participate in plant growth as well as environmental stress processes, such as salt and temperature as signal transduction factors (Gunes et al. 2007). We found that the ARF gene family has various hormone-responsive elements, hat hormones induce ARF gene family expressions in fig, participating in plant growth and development.

FcARFs play important roles in fig fruit development

Expression levels of ARF family members in different parts of fig were analyzed to elucidate the function of *FcARFs*. There were spatial expressions of *FcARFs* in roots, stems, old leaves, young leaves, female flowers and peel at the transcription level. Most ARF genes were highly expressed in female flowers and peels, followed by stems and roots, and least expressed in old leaves, indicating that ARF family members have important roles in fig fruit growth and development. Besides, *AtARF1* is involved in floral organ abscission (Ellis et al. 2005), and its homologous gene (*FcARF1*) is highly expressed in female flower tissues, implying a role in flower growth. Moreover, *FcARFs* (*FcARF 1/2/3/4/5/6/7/8/9/11/12/14/15/18/19/20*) exhibited tissue-specific transcript accumulation patterns. It has been reported that *AtARF11* is highly expressed in the roots and promotes lateral root formation (**Table S1**). *FcARF11* has a high homology with *AtARF11* and has high transcript accumulations in stem and roots (**Figure 6**), indicating that the roles of ARF during root development in fig are similar to those in *Arabidopsis thaliana*.

ARF7/ARF19 mutants in *Arabidopsis* exhibit strong auxin-related phenotypes, including during severe damage to root formation (Okushima et al. 2007; Okushima et al. 2005). *AtARF19* are homologous and have high transcript expressions in roots. It has been postulated that *FcARF19* has important roles in root formation and fig development.

Fruit development and ripening is a coordinated process of reproductive organ cell division, differentiation and expansion that is regulated by transcriptional regulatory networks (Uchiumi & Okamoto 2010). Auxin promotes the growth of female flowers as well as receptacles and causes fruit enlargement that plays a key role in fruit ripening (Flaishman et al. 2008; Kumar et al. 2011). In this study, all *FcARFs* were expressed from the first stage (phase I) to sixth stage (end of phase III) of fruit development. The fruits of figs went through a fast-slow-fast

developmental period. Sensing and transmitting signals have been shown to reveal changes in auxin expression patterns (Estrada-Johnson et al. 2017). *AtARF2* has an important role in regulating flower formation as well as senescence of *Arabidopsis thaliana* (Ellis et al. 2005). Its homologous gene (*FcARF2*) was highly expressed in the early flower and peel development stages, which may regulate fig flower morphologies. *FcARF11* and *FcARF12* from class-Ia were abundantly expressed during the early stages of fig fruit female flower as well as peel development and were predicted to function as transcriptional repressors (**Figure 2B**). It has not been established whether FcARF inhibition acts synergistically with other TFs.

AtARF3 has a major role in nutrition and reproductive development, regulation of gene and embryonic development, and determining the morphological patterns of pistil (Harper et al. 2000; Sessions et al. 1997; Xiao et al. 2004). Its homologous gene (*FcARF3*) is highly expressed during whole growth periods of fig fruits, and suppressed in middle slow growth periods, in tandem with the fig fruit development curve (Song et al. 2021). Moreover, it may be involved in various reproductive and developmental processes, such as female flower development in figs. The fig fruit development candidates obtained from fig ARF gene family analysis is the basis for the next step in fruit development, such as expansion mechanisms caused by female flower development.

Conclusions

The ARF genes are mainly expressed in fruits and flowers and are involved in development and morphogenesis. The common fig fruit contains enlarged female flowers that are closely correlated with development and maturity. Twenty FcARFs containing similar conserved motif structures were identified from fig genome databases, and were divided into three classes (I, II and III) by phylogenetic analyses. They were differentially expressed in tissues, with *FcARF1/5/8/10/12/13/14/15/16/17/18/20* being highly expressed in female flowers and peels, indicating their possible importance in fruit development. Further analysis of the *FcARFs* expression levers in female flower and peel during common fig cv. Purple Peel fruit development. *FcARF2*, *FcARF11* and *FcARF12* are class-Ia members that are mainly involved in regulating flower development. They were highly expressed in early stages of female flower development. Expression patterns of *FcARF3* were in line with fig fruit development curve, implying that it may be involved in the rate-limiting process of fig fruit development as a repressor. This study forms the basis for further functional verification of the fig ARF gene family.

Acknowledgments

Not applicable.

Table S1 (Chen et al. 2021; de Jong et al. 2015; Diao et al. 2020; Ellis et al. 2005; Ge et al. 2016; Goetz et al. 2006; Kelley et al. 2012; Krogan et al. 2014; Kumar et al. 2011; Li et al. 2016; Lim et al. 2010; Liu et al. 2015; Luo et al. 2018; Okushima et al. 2005; Pekker et al. 2005; Sagar et al. 2013; Sessions et al. 1997; Shen et al. 2010; Song et al. 2015; Wan et al. 2014; Wang et al. 2021; Wang et al. 2005; Wang et al. 2010; Wang et al. 2018; Yang et al. 2013; Ye et al. 2016; Zhang et al. 2020; Zhang et al. 2021; Zhang et al. 2015)

Reference

- Arase F, Nishitani H, Egusa M, Nishimoto N, Sakurai S, Sakamoto N, and Kaminaka H. 2012. IAA8 involved in lateral root formation interacts with the TIR1 auxin receptor and ARF transcription factors in Arabidopsis. *PLoS One* 7:e43414. 10.1371/journal.pone.0043414
- Chen Y, Liu B, Zhao Y, Yu W, and Si W. 2021. Whole-Genome Duplication and Purifying Selection Contributes to the Functional Redundancy of Auxin Response Factor (ARF) Genes in Foxtail Millet (*Setaria italica* L.). *Int J Genomics* 2021:2590665. 10.1155/2021/2590665
- Chen Z, Chen X, Yan H, Li W, Li Y, Cai R, and Xiang Y. 2015. The Lipxygenase Gene Family in Poplar: Identification, Classification, and Expression in Response to MeJA Treatment. *PLoS One* 10:e0125526. 10.1371/journal.pone.0125526
- de Jong M, Wolters-Arts M, Schimmel BC, Stultiens CL, de Groot PF, Powers SJ, Tikunov YM, Bovy AG, Mariani C, Vriezen WH, and Rieu I. 2015. *Solanum lycopersicum* AUXIN RESPONSE FACTOR 9 regulates cell division activity during early tomato fruit development. *J Exp Bot* 66:3405-3416. 10.1093/jxb/erv152
- Diao D, Hu X, Guan D, Wang W, Yang H, and Liu Y. 2020. Genome-wide identification of the ARF (auxin response factor) gene family in peach and their expression analysis. *Mol Biol Rep* 47:4331-4344. 10.1007/s11033-020-05525-0
- Doncheva NT, Morris JH, Gorodkin J, and Jensen LJ. 2019. Cytoscape StringApp: Network Analysis and Visualization of Proteomics Data. *J Proteome Res* 18:623-632. 10.1021/acs.jproteome.8b00702
- Du H, Wu N, Fu J, Wang SP, Li XH, Xiao JH, and Xiong LZ. 2012. A GH3 family member, OsGH3-2, modulates auxin and abscisic acid levels and differentially affects drought and cold tolerance in rice. *Journal of Experimental Botany* 63:6467-6480. 10.1093/jxb/ers300
- El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M, Richardson LJ, Salazar GA, Smart A, Sonnhammer ELL, Hirsh L, Paladin L, Piovesan D, Tosatto SCE, and Finn RD. 2019. The Pfam protein families database in 2019. *Nucleic Acids Res* 47:D427-D432. 10.1093/nar/gky995
- Ellis CM, Nagpal P, Young JC, Hagen G, Guilfoyle TJ, and Reed JW. 2005. AUXIN RESPONSE FACTOR1 and AUXIN RESPONSE FACTOR2 regulate senescence and floral organ abscission in Arabidopsis thaliana. *Development* 132:4563-4574. 10.1242/dev.02012
- Estrada-Johnson E, Csukasi F, Pizarro CM, Vallarino JG, Kiryakova Y, Vioque A, Brumos J, Medina-Escobar N, Botella MA, Alonso JM, Fernie AR, Sanchez-Sevilla JF, Osorio S, and Valpuesta V. 2017. Transcriptomic Analysis in Strawberry Fruits Reveals Active Auxin Biosynthesis and Signaling in the Ripe Receptacle. *Frontiers in Plant Science* 8. ARTN 88910.3389/fpls.2017.00889
- Feng ZH, Zhu J, Du XL, and Cui XH. 2012. Effects of three auxin-inducible LBD members on lateral root formation in Arabidopsis thaliana. *Planta* 236:1227-1237. 10.1007/s00425-012-1673-3
- Flaishman MA, Rodov V, and Stover E. 2008. The Fig: Botany, Horticulture, and Breeding. *Horticultural Reviews*, Vol 34 34:113-196.
- Ge J, Li B, Shen D, Xie J, Long J, and Dong H. 2016. Tobacco TTG2 regulates vegetative growth and seed production via the predominant role of ARF8 in cooperation with ARF17 and ARF19. *BMC Plant Biol* 16:126. 10.1186/s12870-016-0815-3
- Goetz M, Vivian-Smith A, Johnson SD, and Koltunow AM. 2006. AUXIN RESPONSE FACTOR8 is a negative

regulator of fruit initiation in Arabidopsis. *Plant Cell* 18:1873-1886. 10.1105/tpc.105.037192

Guilfoyle TJ, and Hagen G. 2007. Auxin response factors. *Current Opinion in Plant Biology* 10:453-460. 10.1016/j.pbi.2007.08.014

Gunes A, Inal A, Alpaslan M, Eraslan F, Bageci EG, and Cicek N. 2007. Salicylic acid induced changes on some physiological parameters symptomatic for oxidative stress and mineral nutrition in maize (*Zea mays* L.) grown under salinity. *Journal of Plant Physiology* 164:728-736. 10.1016/j.jplph.2005.12.009

Guo AY, Zhu QH, Chen X, and Luo JC. 2007. [GSDS: a gene structure display server]. *Yi Chuan* 29:1023-1026.

Guo C, Guo R, Xu X, Gao M, Li X, Song J, Zheng Y, and Wang X. 2014. Evolution and expression analysis of the grape (*Vitis vinifera* L.) WRKY gene family. *J Exp Bot* 65:1513-1528. 10.1093/jxb/eru007

Harper RM, Stowe-Evans EL, Luesse DR, Muto H, Tatematsu K, Watahiki MK, Yamamoto K, and Liscum E. 2000. The NPH4 locus encodes the auxin response factor ARF7, a conditional regulator of differential growth in aerial Arabidopsis tissue. *Plant Cell* 12:757-770. 10.1105/tpc.12.5.757

Hou Q, Qiu Z, Wen Z, Zhang H, Li Z, Hong Y, Qiao G, and Wen X. 2021. Genome-Wide Identification of ARF Gene Family Suggests a Functional Expression Pattern during Fruitlet Abscission in *Prunus avium* L. *Int J Mol Sci* 22. 10.3390/ijms222111968

Hunter C, Willmann MR, Wu G, Yoshikawa M, de la Luz Gutierrez-Nava M, and Poethig SR. 2006. Trans-acting siRNA-mediated repression of ETTIN and ARF4 regulates heteroblasty in Arabidopsis. *Development* 133:2973-2981. 10.1242/dev.02491

Kelley DR, Arreola A, Gallagher TL, and Gasser CS. 2012. ETTIN (ARF3) physically interacts with KANADI proteins to form a functional complex essential for integument development and polarity determination in Arabidopsis. *Development* 139:1105-1109. 10.1242/dev.067918

Krogan NT, Yin XJ, Ckurshumova W, and Berleth T. 2014. Distinct subclades of Aux/IAA genes are direct targets of ARF5/MP transcriptional regulation. *New Phytologist* 204:474-483. 10.1111/nph.12994

Kim J, Harter K, and Theologis A. 1997. Protein-protein interactions among the Aux/IAA proteins. *Proceedings of the National Academy of Sciences of the United States of America* 94:11786-11791. DOI 10.1073/pnas.94.22.11786

Kumar R, Tyagi AK, and Sharma AK. 2011. Genome-wide analysis of auxin response factor (ARF) gene family from tomato and analysis of their role in flower and fruit development. *Mol Genet Genomics* 285:245-260. 10.1007/s00438-011-0602-7

Kumar S, Stecher G, and Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 33:1870-1874. 10.1093/molbev/msw054

Letunic I, Khedkar S, and Bork P. 2021. SMART: recent updates, new developments and status in 2020. *Nucleic Acids Res* 49:D458-D460. 10.1093/nar/gkaa937

Li C, Xu J, Deng Y, Sun H, and Li Y. 2019. Selection of reference genes for normalization of cranberry (*Vaccinium macrocarpon* Ait.) gene expression under different experimental conditions. *PLoS One* 14:e0224798. 10.1371/journal.pone.0224798

Li SB, OuYang WZ, Hou XJ, Xie LL, Hu CG, and Zhang JZ. 2015. Genome-wide identification, isolation and expression analysis of auxin response factor (ARF) gene family in sweet orange (*Citrus sinensis*). *Front Plant Sci* 6:119. 10.3389/fpls.2015.00119

Li SB, Xie ZZ, Hu CG, and Zhang JZ. 2016. A Review of Auxin Response Factors (ARFs) in Plants. *Front Plant Sci* 7:47. 10.3389/fpls.2016.00047

- Lim PO, Lee IC, Kim J, Kim HJ, Ryu JS, Woo HR, and Nam HG. 2010. Auxin response factor 2 (ARF2) plays a major role in regulating auxin-mediated leaf longevity. *J Exp Bot* 61:1419-1430. 10.1093/jxb/erq010
- Liu K, Yuan C, Li H, Lin W, Yang Y, Shen C, and Zheng X. 2015. Genome-wide identification and characterization of auxin response factor (ARF) family genes related to flower and fruit development in papaya (*Carica papaya* L.). *BMC Genomics* 16:901. 10.1186/s12864-015-2182-0
- Liu N, Wu S, Van Houten J, Wang Y, Ding B, Fei Z, Clarke TH, Reed JW, and van der Knaap E. 2014. Down-regulation of AUXIN RESPONSE FACTORS 6 and 8 by microRNA 167 leads to floral development defects and female sterility in tomato. *J Exp Bot* 65:2507-2520. 10.1093/jxb/eru141
- Liu Y, Jiang HY, Chen WJ, Qian YX, Ma Q, Cheng BJ, and Zhu SW. 2011. Genome-wide analysis of the auxin response factor (ARF) gene family in maize (*Zea mays*). *Plant Growth Regulation* 63:225-234. 10.1007/s10725-010-9519-0
- Luo J, Zhou JJ, and Zhang JZ. 2018. Aux/IAA Gene Family in Plants: Molecular Structure, Regulation, and Function. *Int J Mol Sci* 19. 10.3390/ijms19010259
- Luo XC, Sun MH, Xu RR, Shu HR, Wang JW, and Zhang SZ. 2014. Genomewide identification and expression analysis of the ARF gene family in apple. *J Genet* 93:785-797. 10.1007/s12041-014-0462-0
- Mori K, Shirasawa K, Nogata H, Hirata C, Tashiro K, Habu T, Kim S, Himeno S, Kuhara S, and Ikegami H. 2017. Identification of RAN1 orthologue associated with sex determination through whole genome sequencing analysis in fig (*Ficus carica* L.). *Sci Rep* 7:41124. 10.1038/srep41124
- Oh E, Zhu JY, Bai MY, Arenhart RA, Sun Y, and Wang ZY. 2014. Cell elongation is regulated through a central circuit of interacting transcription factors in the *Arabidopsis* hypocotyl. *Elife* 3. ARTN e0303110.7554/eLife.03031
- Okushima Y, Fukaki H, Onoda M, Theologis A, and Tasaka M. 2007. ARF7 and ARF19 regulate lateral root formation via direct activation of LBD/ASL genes in *Arabidopsis*. *Plant Cell* 19:118-130. 10.1105/tpc.106.047761
- Okushima Y, Overvoorde PJ, Arima K, Alonso JM, Chan A, Chang C, Ecker JR, Hughes B, Lui A, Nguyen D, Onodera C, Quach H, Smith A, Yu G, and Theologis A. 2005. Functional genomic analysis of the AUXIN RESPONSE FACTOR gene family members in *Arabidopsis thaliana*: unique and overlapping functions of ARF7 and ARF19. *Plant Cell* 17:444-463. 10.1105/tpc.104.028316
- Pandey SK, Lee HW, Kim MJ, Cho C, Oh E, and Kim J. 2018. LBD18 uses a dual mode of a positive feedback loop to regulate ARF expression and transcriptional activity in *Arabidopsis*. *Plant J* 95:233-251. 10.1111/tpj.13945
- Pekker I, Alvarez JP, and Eshed Y. 2005. Auxin response factors mediate *Arabidopsis* organ asymmetry via modulation of KANADI activity. *Plant Cell* 17:2899-2910. 10.1105/tpc.105.034876
- Reed JW. 2001. Roles and activities of Aux/IAA proteins in *Arabidopsis*. *Trends Plant Sci* 6:420-425. 10.1016/s1360-1385(01)02042-8
- Roosjen M, Pague S, and Weijers D. 2018. Auxin Response Factors: output control in auxin biology. *Journal of Experimental Botany* 69:179-188. 10.1093/jxb/erx237
- Rosas-Camacho O. 2009. Deterministic Modeling of the Corrosion of Low-Carbon Steel by Dissolved Carbon Dioxide and the Effect of Acetic Acid. *ECS Transactions* 143-165.
- Sagar M, Chervin C, Mila I, Hao Y, Roustan JP, Benichou M, Gibon Y, Biais B, Maury P, Latche A, Pech JC, Bouzayen M, and Zouine M. 2013. SlARF4, an auxin response factor involved in the control of sugar metabolism during tomato fruit development. *Plant Physiol* 161:1362-1374. 10.1104/pp.113.213843
- Sessions A, Nemhauser JL, McColl A, Roe JL, Feldmann KA, and Zambryski PC. 1997. ETTIN patterns the

- Arabidopsis floral meristem and reproductive organs. *Development* 124:4481-4491. 10.1242/dev.124.22.4481
- Song M, Wang H, Wang Z, Huang H, Chen S, and Ma H. 2021. Genome-Wide Characterization and Analysis of bHLH Transcription Factors Related to Anthocyanin Biosynthesis in Fig (*Ficus carica* L.). *Front Plant Sci* 12:730692. 10.3389/fpls.2021.730692
- Shen C, Wang S, Bai Y, Wu Y, Zhang S, Chen M, Guilfoyle TJ, Wu P, and Qi Y. 2010. Functional analysis of the structural domain of ARF proteins in rice (*Oryza sativa* L.). *J Exp Bot* 61:3971-3981. 10.1093/jxb/erq208
- Song J, Gao ZH, Huo XM, Sun HL, Xu YS, Shi T, and Ni ZJ. 2015. Genome-wide identification of the auxin response factor (ARF) gene family and expression analysis of its role associated with pistil development in Japanese apricot (*Prunus mume* Sieb. et Zucc). *Acta Physiologiae Plantarum* 37. ARTN 14510.1007/s11738-015-1882-z
- Sun H, Fan HJ, and Ling HQ. 2015a. Genome-wide identification and characterization of the bHLH gene family in tomato. *BMC Genomics* 16:9. 10.1186/s12864-014-1209-2
- Sun R, Wang K, Guo T, Jones DC, Cobb J, Zhang B, and Wang Q. 2015b. Genome-wide identification of auxin response factor (ARF) genes and its tissue-specific prominent expression in *Gossypium raimondii*. *Funct Integr Genomics* 15:481-493. 10.1007/s10142-015-0437-0
- Swarup R, and Bhosale R. 2019. Developmental Roles of AUX1/LAX Auxin Influx Carriers in Plants. *Front Plant Sci* 10:1306. 10.3389/fpls.2019.01306
- Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, Doncheva NT, Legeay M, Fang T, Bork P, Jensen LJ, and von Mering C. 2021. The STRING database in 2021: customizable protein-protein networks, and functional characterization of user-uploaded gene/measurement sets. *Nucleic Acids Res* 49:D605-D612. 10.1093/nar/gkaa1074
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, and Kumar S. 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28:2731-2739. 10.1093/molbev/msr121
- Tang H, Wang X, Bowers JE, Ming R, Alam M, and Paterson AH. 2008. Unraveling ancient hexaploidy through multiply-aligned angiosperm gene maps. *Genome Research* 18:1944-1954. 10.1101/gr.080978.108
- Tiwari SB, Hagen G, and Guilfoyle T. 2003. The roles of auxin response factor domains in auxin-responsive transcription. *Plant Cell* 15:533-543. 10.1105/tpc.008417
- Truskina J, Han J, Chrysanthou E, Galvan-Ampudia CS, Laine S, Brunoud G, Mace J, Bellows S, Legrand J, Bagman AM, Smit ME, Smetana O, Stigliani A, Porco S, Bennett MJ, Mahonen AP, Parcy F, Farcot E, Roudier F, Brady SM, Bishopp A, and Vernoux T. 2021. Publisher Correction: A network of transcriptional repressors modulates auxin responses. *Nature* 589:E7. 10.1038/s41586-020-03066-x
- Uchiumi T, and Okamoto T. 2010. Rice fruit development is associated with an increased IAA content in pollinated ovaries. *Planta* 232:579-592. 10.1007/s00425-010-1197-7
- Usai G, Mascagni F, Giordani T, Vangelisti A, Bosi E, Zuccolo A, Ceccarelli M, King R, Hassani-Pak K, Zambrano LS, Cavallini A, and Natali L. 2020. Epigenetic patterns within the haplotype phased fig (*Ficus carica* L.) genome. *Plant J* 102:600-614. 10.1111/tpj.14635
- Vision TJ, Brown DG, and Tanksley SD. 2000. The origins of genomic duplications in *Arabidopsis*. *Science* 290:2114-2117. 10.1126/science.290.5499.2114
- Wan S, Li W, Zhu Y, Liu Z, Huang W, and Zhan J. 2014. Genome-wide identification, characterization and expression

- analysis of the auxin response factor gene family in *Vitis vinifera*. *Plant Cell Rep* 33:1365-1375. 10.1007/s00299-014-1622-7
- Wang D, Pei K, Fu Y, Sun Z, Li S, Liu H, Tang K, Han B, and Tao Y. 2007. Genome-wide analysis of the auxin response factors (ARF) gene family in rice (*Oryza sativa*). *Gene* 394:13-24. 10.1016/j.gene.2007.01.006
- Wan SB, Li WL, Zhu YY, Liu ZM, Huang WD, and Zhan JC. 2014. Genome-wide identification, characterization and expression analysis of the auxin response factor gene family in *Vitis vinifera*. *Plant Cell Reports* 33:1365-1375. 10.1007/s00299-014-1622-7
- Wang CK, Han PL, Zhao YW, Yu JQ, You CX, Hu DG, and Hao YJ. 2021. Genome-wide analysis of auxin response factor (ARF) genes and functional identification of MdARF2 reveals the involvement in the regulation of anthocyanin accumulation in apple. *New Zealand Journal of Crop and Horticultural Science* 49:78-91. 10.1080/01140671.2020.1779756
- Wang JW, Wang LJ, Mao YB, Cai WJ, Xue HW, and Chen XY. 2005. Control of root cap formation by MicroRNA-targeted auxin response factors in *Arabidopsis*. *Plant Cell* 17:2204-2216. 10.1105/tpc.105.033076
- Wang L, Feng Z, Wang X, Wang X, and Zhang X. 2010. DEGseq: an R package for identifying differentially expressed genes from RNA-seq data. *Bioinformatics* 26:136-138. 10.1093/bioinformatics/btp612
- Wang YC, Wang N, Xu HF, Jiang SH, Fang HC, Su MY, Zhang ZY, Zhang TL, and Chen XS. 2018. Auxin regulates anthocyanin biosynthesis through the Aux/IAA-ARF signaling pathway in apple. *Horticulture Research* 5. ARTN 5910.1038/s41438-018-0068-4
- Wang R, Zhao P, Kong N, Lu R, Pei Y, Huang C, Ma H, and Chen Q. 2018. Genome-Wide Identification and Characterization of the Potato bHLH Transcription Factor Family. *Genes (Basel)* 9. 10.3390/genes9010054
- Wang SX, Shi FY, Dong XX, Li YX, Zhang ZH, and Li H. 2019. Genome-wide identification and expression analysis of auxin response factor (ARF) gene family in strawberry (*Fragaria vesca*). *Journal of Integrative Agriculture* 18:1587-1603. 10.1016/S2095-3119(19)62556-6
- Wang YN, Li KX, Chen L, Zou YM, Liu HP, Tian YP, Li DX, Wang R, Zhao F, Ferguson BJ, Gresshoff PM, and Li X. 2015. MicroRNA167-Directed Regulation of the Auxin Response Factors GmARF8a and GmARF8b Is Required for Soybean Nodulation and Lateral Root Development. *Plant Physiology* 168:101-+. 10.1104/pp.15.00265
- Wei W, Hu Y, Cui MY, Han YT, Gao K, and Feng JY. 2016. Identification and Transcript Analysis of the TCP Transcription Factors in the Diploid Woodland Strawberry *Fragaria vesca*. *Front Plant Sci* 7:1937. 10.3389/fpls.2016.01937
- Woodward C, Bemis SM, Hill EJ, Sawa S, Koshida T, and Torii KU. 2005. Interaction of auxin and ERECTA in elaborating *Arabidopsis* inflorescence architecture revealed by the activation tagging of a new member of the YUCCA family putative flavin monooxygenases. *Plant Physiology* 139:192-203. 10.1104/pp.105.063495
- Wu MF, Tian Q, and Reed JW. 2006. *Arabidopsis* microRNA167 controls patterns of ARF6 and ARF8 expression, and regulates both female and male reproduction. *Development* 133:4211-4218. 10.1242/dev.02602
- Xiao JN, Huang XL, Huang X, and Li XJ. 2004. [Identification of an auxin response factor-like protein cDNA from mango cotyledon section]. *Sheng Wu Gong Cheng Xue Bao* 20:59-62.
- Xing H, Pudake RN, Guo G, Xing G, Hu Z, Zhang Y, Sun Q, and Ni Z. 2011a. Genome-wide identification and expression profiling of auxin response factor (ARF) gene family in maize. *BMC Genomics* 12:178. 10.1186/1471-2164-12-178
- Xing HY, Pudake RN, Guo GG, Xing GF, Hu ZR, Zhang YR, Sun QX, and Ni ZF. 2011b. Genome-wide identification

and expression profiling of auxin response factor (ARF) gene family in maize. *BMC Genomics* 12. Artn 178
10.1186/1471-2164-12-178

Yang C, Xu M, Xuan L, Jiang X, and Huang M. 2014. Identification and expression analysis of twenty ARF genes in
Populus. *Gene* 544:134-144. 10.1016/j.gene.2014.04.067

Yang X, Lee S, So JH, Dharmasiri S, Dharmasiri N, Ge L, Jensen C, Hangarter R, Hobbie L, and Estelle M. 2004.
The IAA1 protein is encoded by AXR5 and is a substrate of SCF(TIR1). *Plant J* 40:772-782. 10.1111/j.1365-
313X.2004.02254.x

Ye YJ, Gong ZY, Lu X, Miao DY, Shi JM, Lu J, and Zhao Y. 2016. Germostatin resistance locus 1 encodes a PHD
finger protein involved in auxin-mediated seed dormancy and germination. *Plant Journal* 85:3-15.
10.1111/tpj.13086

Ye Y, Wang J, Wang W, and Xu LA. 2020. ARF family identification in *Tamarix chinensis* reveals the salt responsive
expression of TcARF6 targeted by miR167. *PeerJ* 8:e8829. 10.7717/peerj.8829

Yue P, Lu Q, Liu Z, Lv T, Li X, Bu H, Liu W, Xu Y, Yuan H, and Wang A. 2020. Auxin-activated MdARF5 induces
the expression of ethylene biosynthetic genes to initiate apple fruit ripening. *New Phytol* 226:1781-1795.
10.1111/nph.16500

Zhang F, Tao WQ, Sun RQ, Wang JX, Li CL, Kong XP, Tian HY, and Ding ZJ. 2020. PRH1 mediates ARF7-LBD
dependent auxin signaling to regulate lateral root development in *Arabidopsis thaliana*. *Plos Genetics* 16.
ARTN e100804410.1371/journal.pgen.1008044

Zhang HX, Cao N, Dong CJ, and Shang QM. 2017. Genome-wide Identification and Expression of ARF Gene Family
during Adventitious Root Development in Hot Pepper (*Capsicum annuum*). *Horticultural Plant Journal*
3:151-164. 10.1016/j.hpj.2017.07.001

Zhang X, Cao J, Huang C, Zheng Z, Liu X, Shangguan X, Wang L, Zhang Y, and Chen Z. 2021. Characterization of
cotton ARF factors and the role of GhARF2b in fiber development. *BMC Genomics* 22:202. 10.1186/s12864-
021-07504-6

Zhang X, Wang G, Zhang S, Chen S, Wang Y, Wen P, Ma X, Shi Y, Qi R, Yang Y, Liao Z, Lin J, Lin J, Xu X, Chen
X, Xu X, Deng F, Zhao L, Lee YL, Wang R, Chen XY, Lin YR, Zhang J, Tang H, Chen J, and Ming R. 2020.
Genomes of the Banyan Tree and Pollinator Wasp Provide Insights into Fig-Wasp Coevolution. *Cell*
183:875-889 e817. 10.1016/j.cell.2020.09.043

Zhang XL, Yan F, Tang YW, Yuan YJ, Deng W, and Li ZG. 2015. Auxin Response Gene SlARF3 Plays Multiple
Roles in Tomato Development and is Involved in the Formation of Epidermal Cells and Trichomes. *Plant*
and Cell Physiology 56:2110-2124. 10.1093/pcp/pcv136

Zhou Z, Schenke D, Miao Y, and Cai D. 2017. Investigation of the crosstalk between the flg22 and the UV-B-induced
flavonol pathway in *Arabidopsis thaliana* seedlings. *Plant Cell Environ* 40:453-458. 10.1111/pce.12869

Zouine M, Fu Y, Chateigner-Boutin AL, Mila I, Frasse P, Wang H, Audran C, Roustan JP, and Bouzayen M. 2014.
Characterization of the tomato ARF gene family uncovers a multi-levels post-transcriptional regulation
including alternative splicing. *PLoS One* 9:e84203. 10.1371/journal.pone.0084203

Zuckerkandl E, and Pauling L. 1965. Evolutionary Divergence and Convergence in Proteins. In: Bryson V, and Vogel
HJ, eds. *Evolving Genes and Proteins*. Academic Press, 97-166.

Figure 1

Figure 1. Phylogenetic analysis of auxin response factors from *Ficus carica* L., *Arabidopsis thaliana*, *Fragaria vesca*, *Vitis vinifera* and *Malus domestica*.


The phylogenetic tree was constructed using the neighbor-joining method in MEGA 7.0. Bootstrap values from 1,000 replicates were indicated at each branch. Gene groups were indicated with different colors and were classified into four groups: class-I, class-II and class-III. class-I includes three subclasses: class-Ia class-Ib and class-Ic.



Figure 2

Figure 2. Analysis of the sequence and structure of the ARF gene family in Fig.

(A) Analysis of protein domains and gene structures. The B3 DNA binding domain (DBD), middle region (MR), and AUX/IAA (CTD) family domain were shown in green, purple and pink, respectively. The orange circle represents a transcription activator, and the blue circle represents a transcription inhibitor. (B) Distribution of the conserved motifs of FcARFs. Sequence alignment of the conserved B3 DNA binding domain (DBD), middle region (MR), and AUX/IAA (CTD) were performed by CLUSTALW.



B

Middle Region (MR)

CoARF1 VISSSHMHGLVATASHVLTGTTIVYHRS RTSPSSVVFIFRVMSVK.SNYSIOBKKRMKDEAEAEQRC.TGIIIGLAEHSGIIPASBACIKVGNASSNPVFLPBLRSHLSINFTYHNRNLF... FKKKFP
CoARF2 IVRNSHMHGLVATASHVLTGTTIVYHRS RTSPSSVVFIFRVMSVK.SNYSIOBKKRMKDEAEAEQRC.TGIIIGLAEHSGIIPASBACIKVGNASSNPVFLPBLRSHLSINFTYHNRNLF... FKKKFP
CoARF3 IIVNNSYENVSLVNAVSTKSMHVPSSRATHABVIFVFKYKVSIT.NPVSIGRETRRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF4 IIE.KNSCNSILVNAVSTKSMHVPSSRATHABVIFVFKYKVSIT.SPVCIGRETRRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF5 VISSSHMHGLVIAAAAHAAANSSTIETINE.RASPSVVFIFLAKYKRVHVTYVSGRERMLRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF6 VISSSHMHGLVIAAAAHAAANSSTIETINE.RASPSVVFIFLAKYKRVHVTYVSGRERMLRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF7 VISSSHMHGLVIAAAAHAAANSSTIETINE.RASPSVVFIFLAKYKRVHVTYVSGRERMLRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF8 VISSSHMHGLVIAAAAHAAANSSTIETINE.RASPSVVFIFLAKYKRVHVTYVSGRERMLRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF9 VISSSHMHGLVATASHVLTGTTIVYHRS.RTS..GTISNKKYLEALN.NKFCIGRETRRMBRDEAEFFRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF10 NYSRGGVRAESVGEAVLAAAGCEHVEYHRS.RASTPFGVCSAVRAAGM..IRWFAGRETRRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF11 VISSSHMHGLVATASHVLTGTTIVYHRS RTSPSSVVFIFRVMSVK.SNYSIOBKKRMKDEAEAEQRC.TGIIIGLAEHSGIIPASBACIKVGNASSNPVFLPBLRSHLSINFTYHNRNLF... FKKKFP
CoARF12 VISSSHMHGLVATASHVLTGTTIVYHRS RTSPSSVVFIFRVMSVK.SNYSIOBKKRMKDEAEAEQRC.TGIIIGLAEHSGIIPASBACIKVGNASSNPVFLPBLRSHLSINFTYHNRNLF... FKKKFP
CoARF13 IIVKGGKVPFVESVETIETIASNGCEHEVYHRS.RASTPFGVCSAVRAAGM..IRWFAGRETRRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF14 IMKSHKVPFVESVETIETIASNGCEHEVYHRS.RASTPFGVCSAVRAAGM..IRWFAGRETRRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF15 IIVKSHKVPFVESVETIETIASNGCEHEVYHRS.RASTPFGVCSAVRAAGM..IRWFAGRETRRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF16 MRGTGVRPEAVIAAAAHAAANSSTIETINE.RASPSVVFIFLAKYKRVHVTYVSGRERMLRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF17 ..SNGLSVAEVCLVAKVLAHKNKCTIVYHRS.RDGAEPFCVSAVRAALG.VQMGIGRETRRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF18 VISSSHMHGLVATASHVLTGTTIVYHRS RTSPSSVVFIFRVMSVK.LHLSVIGRETRRMBRDEEFFERRC...GEG...TTSGAGSWSRSLVQDSESLTCHP.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF19 VISSSHMHGLVIAAAAHAAANSSTIETINE.RASPSVVFIFLAKYKRVHVTYVSGRERMLRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP
CoARF20 VISSSHMHGLVATASHVLTGTTIVYHRS.RTSABVIFVCFPMKSTK.NPVSIGRETRRMBRDEEFFERRC.SGVITGICSLDEPVNSHRSRGNVQDGEASTAGER.PSRVSHKEELFPAALPAPNLFV... FKKKFP

DNA Binding Domain (DBD)

[illegible]

AUX/IAA Family (CTD)

[illegible]


Figure 3

Figure 3 □ Analysis of gene structures and cis-acting elements for FcARF genes.

(A) Analysis of exon-intron structures for FcARF genes. Blue rectangles, yellow rectangles and blank lines were indicated upstream/downstream, exons and introns, respectively.

The lengths of exons and introns are indicated by the scale bar. (B) The distribution of cis-elements in the 2000 bp upstream promoter regions of FcARF genes were analyzed. The *cis*-elements were predicted at PlantCARE ([https:// bioinformatics. psb. ugent. be/ webto ols/ plant care/ html/](https://bioinformatics.psb.ugent.be/webtools/plantcare/html/)). Different *cis*-elements were presented with different shapes and colors.

A



B




Figure 4

Figure 4. Chromosomal localization and collinearity analysis of the FcARF gene family.


(A) Collinearity analysis of FcARF genes, the circle plot was created with the MCScanX tool. Identified collared genes are linked by colored lines. (B) Collinearity relationship of ARF genes among *Ficus carica* (Fc), *Ficus hispida* (Fh), and *Ficus microcarpa* (Fm). Identified collinear genes are linked by blue lines.

Figure 5

Figure 5. Correlation analysis network among auxinsignaling-relatedpredicted proteins .

(A) The FcARFs and other predicted proteins interaction network were analyzed in the STRING database. Green color indicated FcARFs and the gray color indicated other auxinsignaling-related predicted proteins. Athickerlineindicates astrongerinteractionbetweenthe two proteins. (B) The auxinsignaling-relatedpredicted genes expression correlation analysis was performed by Pearson algorithm, based on these gene's expression of the female flower and peel during the fig fruit development, the data in the cells were indicated the correlations scores .

A



B




Figure 6

Figure 6. Tissue-specific transcript expression patterns of 20 ARF genes in the 'Purple Peel' fig.

Different plant tissues. RO: Root; ST: Stem; OL: Old Leaf; YF: Young Leaf; FL: Flower; PE: Peel. Relative transcript levels are calculated by qRT-PCR with *β-actin* as a standard. Data are means \pm SD of three separate measurements. The qRT-PCR data were analyzed by relative quantification using $2^{-\Delta\Delta C_t}$. Different letters above the bars indicate statistically significant differences ($P=0.05$).




Figure 7

Figure 7: Transcript expression pattern of 20 ARF genes in the 'Purple Peel' fig during different fruit development stages.

Fruit developmental stages. Stage 1: represented phase I (the first rapid growth period); stages 2, 3, and 4: were the early, middle, and late stages of phase II (slow growth period); stages 5 and 6: represented phase III (the second rapid growth period).




Table 1(on next page)

Table 1 FcARF genes encoding ARF proteins along with their molecular details.

1

Table 1 FcARF genes encoding ARF proteins along with their molecular details.

Gene name	Gene ID	Exon No.	Length (AA) ^a	Mw (k Da) ^b	Atomic Formula	pI ^c	Instability index	Aliphatic index	GRAVY	Localization
<i>FcARF1</i>	FCD_00029581	16	897	99.27	C ₃₂₇₉ H ₅₁₂₆ N ₉₂₈ O ₁₀₁₉ S ₃₄	6.04	71.85	73.03	-0.463	Nuclear
<i>FcARF2</i>	FCD_00016093	14	843	94.02	C ₄₁₁₆ H ₆₄₂₃ N ₁₁₉₁ O ₁₂₇₁ S ₃₄	6.72	55.25	64.06	-0.654	Nuclear
<i>FcARF3</i>	FCD_00001359	12	813	89.96	C ₃₉₄₆ H ₆₁₆₇ N ₁₁₁₁ O ₁₂₂₉ S ₃₅	6.55	56.15	71.5	-0.465	Nuclear
<i>FcARF4</i>	FCD_00032414	11	768	52.73	C ₂₃₃₅ H ₃₆₂₃ N ₆₄₇ O ₇₁₄ S ₁₇	7.21	57.64	69.49	-0.510	Nuclear
<i>FcARF5</i>	FCD_00003481	15	942	103.77	C ₄₅₄₄ H ₇₁₃₉ N ₁₂₄₁ O ₁₄₄₈ S ₄₅	5.03	58.47	78.33	-0.370	Nuclear
<i>FcARF6</i>	FCD_00013415	14	987	99.27	C ₄₃₆₀ H ₆₇₉₈ N ₁₂₃₄ O ₁₃₄₇ S ₃₈	6.04	71.85	73.03	-0.463	Nuclear
<i>FcARF7</i>	FCD_00017652	13	1150	127.056	C ₅₅₃₁ H ₈₇₂₃ N ₁₅₉₅ O ₁₇₃₇ S ₅₃	6.57	63.56	73.5	-0.511	Nuclear
<i>FcARF8</i>	FCD_00034325	14	899	99.39	C ₄₃₆₇ H ₆₇₈₁ N ₁₂₂₉ O ₁₃₆₂ S ₃₄	5.86	71.46	72.45	-0.491	Nuclear
<i>FcARF9</i>	FCD_00028996	14	687	76.74	C ₃₃₆₆ H ₅₃₁₁ N ₉₄₉ O ₁₀₅₄ S ₂₅	6.22	52.53	72.63	-0.523	Nuclear
<i>FcARF10</i>	FCD_00021037	3	588	65.32	C ₂₉₀₁ H ₄₄₇₈ N ₈₀₆ O ₈₇₁ S ₂₃	6.95	43.7	70.1	-0.475	Nuclear
<i>FcARF11</i>	FCD_00022972	11	595	67.15	C ₂₉₆₉ H ₄₆₃₈ N ₈₇₀ O ₈₇₈ S ₁	9.52	60.21	63.06	-0.726	Nuclear
<i>FcARF12</i>	FCD_00032300	12	689	78.08	C ₃₄₅₃ H ₅₄₃₂ N ₁₀₁₂ O ₁₀₂₀ S ₂₀	9.42	54.24	72.12	-0.655	Nuclear
<i>FcARF13</i>	FCD_00005247	5	699	76.48	C ₃₃₇₈ H ₅₂₃₃ N ₉₄₅ O ₁₀₂₅	6.36	50.71	71.43	-0.390	Cytoplasmic

					S ₃₁					
					C ₃₅₇₂ H ₅₅₀₄					
<i>FcARF14</i>	FCD_00 007016	14	721	81.01	N ₁₀₀₀ O ₁₀₈₁	6.35	60.08	66.81	-0.519	Nuclear
					S ₃₉					
					C ₄₃₈₁ H ₆₈₀₇					
<i>FcARF15</i>	FCD_00 016691	14	902	99.73	N ₁₂₃₃ O ₁₃₆₈	5.83	72.35	72.64	-0.493	Nuclear
					S ₃₄					
					C ₃₃₉₉ H ₅₃₁₀					
<i>FcARF16</i>	FCD_00 001287	4	708	77.28	N ₉₇₂ O ₁₀₃₁	7.62	44.78	72.57	-0.366	Cytoplasmic
					S ₃₁					
					C ₂₉₆₀ H ₄₆₃₉					
<i>FcARF17</i>	FCD_00 001289	4	620	67.06	N ₈₁₉ O ₉₃₀ S	5.90	42.05	79.69	-0.322	Nuclear
					S ₁₅					
					C ₃₀₅₄ H ₄₈₁₂					
<i>FcARF18</i>	FCD_00 035227	17	617	69.43	N ₈₆₂ O ₉₂₁ S	6.04	56.32	84.46	-0.326	Cytoplasmic
					S ₃₄					
					C ₅₄₂₃ H ₈₄₆₆					
<i>FcARF19</i>	FCD_00 010593	13	1115	124.55	N ₁₅₇₆ O ₁₇₁₂	6.03	70.65	68.66	-0.704	Nuclear
					S ₄₄					
					C ₃₅₇₆ H ₅₅₆₈					
<i>FcARF20</i>	FCD_00 013930	14	719	80.78	N ₁₀₁₄ O ₁₀₇₄	6.28	51.39	75.26	-0.554	Nuclear
					S ₂₆					

2

3 ^a Length of open reading frame in base pairs

4 ^b Molecular weight (Kilodaltons)

5 ^c Isoelectric point

6