

Moonlighting proteins in medically relevant fungi

Verania J. Arvizu-Rubio, Laura C. García-Carnero and Héctor Manuel Mora-Montes

Department of Biology, Universidad de Guanajuato, Guanajuato, Mexico

ABSTRACT

Moonlighting proteins represent an intriguing area of cell biology, due to their ability to perform two or more unrelated functions in one or many cellular compartments. These proteins have been described in all kingdoms of life and are usually constitutively expressed and conserved proteins with housekeeping functions. Although widely studied in pathogenic bacteria, the information about these proteins in pathogenic fungi is scarce, but there are some reports of their functions in the etiological agents of the main human mycoses, such as *Candida* spp., *Paracoccidioides brasiliensis*, *Histoplasma capsulatum*, *Aspergillus fumigatus*, *Cryptococcus neoformans*, and *Sporothrix schenckii*. In these fungi, most of the described moonlighting proteins are metabolic enzymes, such as enolase and glyceraldehyde-3-phosphate dehydrogenase; chaperones, transcription factors, and redox response proteins, such as peroxiredoxin and catalase, which moonlight at the cell surface and perform virulence-related processes, contributing to immune evasion, adhesions, invasion, and dissemination to host cells and tissues. All moonlighting proteins and their functions described in this review highlight the limited information about this biological aspect in pathogenic fungi, representing this a relevant opportunity area that will contribute to expanding our current knowledge of these organisms' pathogenesis.

Submitted 11 July 2022

Accepted 13 August 2022

Published 13 September 2022

Corresponding authors

Laura C. García-Carnero,
laura_cgc@hotmail.com

Héctor Manuel Mora-Montes,
hmora@ugto.mx

Academic editor

Nancy Keller

Additional Information and
Declarations can be found on
page 16

DOI 10.7717/peerj.14001

© Copyright
2022 Arvizu-Rubio et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Biochemistry, Cell Biology, Microbiology, Molecular Biology, Mycology

Keywords Virulence factor, Cell wall, Alternative function, Host-fungus interaction, Pathogenesis, Adhesin, Chaperone

INTRODUCTION

Moonlighting proteins are defined as multifunctional proteins with two or more biochemical or biophysical functions, that are not due to gene fusion, splicing, pleiotropic effects, proteolysis, or promiscuous enzymatic activities (*Huberts & van der Klei, 2010; Copley, 2012; Amblee & Jeffery, 2015; Jeffery, 2015, 1999*). According to the MoonProt database (<http://www.moonlightingproteins.org>), there are over 500 proteins that have been experimentally identified as moonlighting (*Chen et al., 2021*), and some of them have been found in pathogenic organisms, playing important roles during the host-pathogen interaction.

Many of the known moonlighting proteins are highly conserved and constitutively expressed housekeeping proteins, found in many organisms and cell types, functioning as chaperones or metabolic enzymes (*Gancedo & Flores, 2008; Huberts & van der Klei, 2010; Gancedo et al., 2010*).

Henderson & Martin, 2011), with alternative functions related to transcription regulation, intron splicing, DNA protection and repair, assembly of organelles, chaperones, adhesion, and cell surface receptors (*Gancedo, Flores & Gancedo, 2016*). These multitasking proteins may perform their alternative functions at the same time or may change from one to another, representing a switch point among functions that allows the cell to quickly respond to different environmental conditions at a low energetic cost (*Jeffery, 2004*; *Karkowska-Kuleta & Kozik, 2014*; *Jeffery, 2018*). Whether one function or the other is being performed is highly related to (i) the protein cellular localization and cell type, (ii) the available concentrations of the protein partners, (iii) the protein oligomeric state, (iv) post-translational modifications that the protein may undergo, or (v) a combination of these mechanisms (*Jeffery, 2018*).

A current working model trying to explain moonlighting functions indicates that these are dependent on the ability of the protein to perform them, developed after the canonical function and caused by a small number of mutations with little effect on this first function, which results in improved organism fitness, being therefore enhanced and selected during evolution (*Aharoni et al., 2005*; *Gancedo & Flores, 2008*; *Huberts & van der Klei, 2010*).

Even though many moonlighting proteins and their canonical functions are conserved among organisms, their alternative functions are not, for which it is not possible to predict these second functions using homology, since they do not depend on conserved motifs and domains (*Huberts & van der Klei, 2010*; *Copley, 2012*). However, it has been observed that many of these proteins, with cytoplasmic and extracellular functions, share certain structural and biophysical characteristics in their primary amino acid sequences and tertiary structures, such as the lack of signal peptide and anchors or motifs that attach them to the cell surface, high aliphatic indexes, high molecular weights, acidic isoelectric points, and negative hydropathic scores, features usually observed in cytosolic proteins (*Amblee & Jeffery, 2015*; *Jeffery, 2018*). An increasing number of moonlighting proteins have been identified as intrinsically disordered proteins or with intrinsically disordered domains or regions (IDDs) often related to the interaction of the protein with one or many partners, due to disorder-to-order transitions and alternative conformations that provide them with structural malleability (*Jeffery, 2015*; *Liu & Jeffery, 2020*; *Tompa, Szász & Buday, 2005*). Therefore, moonlighting functions can derive from modifications of the protein sequence, (i) which can be large conformational changes or transitions between IDDs and multiple distinct folded structures, so that different protein structure conformations can perform different functions, (ii) or can be subtle changes in small portions of amino acids in the protein sequence needed for the second function without altering the protein structure or affecting the region responsible for the canonical function (*Copley, 2012*; *Jeffery, 2018*).

Targeting of moonlighting proteins to the cell surface is still poorly understood, since they lack conventional secretion signals, such as the signal peptide, but it has been well proven that the finding of these proteins at the surface is due to secretion and not to cell leakage or adsorption from surrounding dead cells (*Jeffery, 2018*). Therefore, secretion systems in charge of targeting cytosolic moonlighting proteins to the surface are capable of distinguishing those that should stay inside the cell to perform their canonical functions from proteins that should be extracellularly located (*Jeffery, 2018*; *Copley, 2012*). Some

non-conventional secretory pathways have been suggested for the transport of these and other proteins that lack signal peptides, including (i) passive protein transfer through the plasma membrane, (ii) membrane flipping with the transfer of attached cytoplasmic proteins to the surface, (iii) translocation *via* soluble or membrane-bound transporters ([Karkowska-Kuleta & Kozik, 2014](#)), (iv) secretory vesicles or endosomal sub-compartments engaged in secretion ([Karkowska-Kuleta & Kozik, 2014](#)), (v) exosome-like vesicles ([Rodrigues & Djordjevic, 2012](#)), (vi) secretory lysosomes ([Nickel & Rabouille, 2009](#)), (vii) and post-translational translocation into the endoplasmic reticulum to enter the secretory pathway ([Hernández-Chávez et al., 2014](#); [Schmoll et al., 2016](#)).

Over the last few years, moonlighting proteins have been shown to play important roles not only in the maintenance of basic cellular functions but also during the host-pathogen interaction, representing an important and wide set of virulence factors that enhance the pathogen's ability to cause infection.

Moonlighting proteins as fungal virulence factors

Although the role of moonlighting proteins during the pathogen-host interaction is widely reported for bacteria ([Henderson & Martin, 2011](#)), this is a phenomenon less explored in fungi, probably due to the difficulty to study protein function in eukaryotic pathogens. However, some moonlighting proteins participating during fungal infection have been suggested and confirmed in pathogenic fungi, such as *Candida* spp., *Paracoccidioides brasiliensis*, *Histoplasma capsulatum*, *Aspergillus fumigatus*, *Cryptococcus neoformans*, and *Sporothrix schenckii*.

The fungal cell wall, and the capsule in capsulated fungal species, is the first structure that encounters the host cells and molecules during infection, and participates in the pathogen adherence to the host tissues and interaction with the host immune effectors. This structure is composed of polysaccharides, pigments, and proteins, which work as virulence determinants or factors during infection ([Díaz-Jiménez et al., 2012](#); [Mora-Montes et al., 2009](#); [Teixeira et al., 2014](#); [Tóth et al., 2019](#); [Gómez-Gaviria & Mora-Montes, 2020](#); [Navarro-Arias et al., 2019](#)). Some of the cell wall proteins have been reported to be conserved intracellular proteins moonlighting at the cell surface, with housekeeping canonical functions related to the following ([Satala et al., 2020a](#)):

- (a) Metabolic pathways, including (i) glycolysis and gluconeogenesis: fructose-bisphosphate aldolase (Fba), phosphoglycerate mutase (Gpm), glyceraldehyde-3-phosphate dehydrogenase (GAPDH or Tdh3), phosphoglycerate kinase (Pgk), glucose-6-phosphate isomerase 1 (Gpi1), triosephosphate isomerase (Tpi), glycerol-3-phosphate dehydrogenase 2 (Gpd2), enolase (Eno), trehalose-6-phosphate phosphatase (Tsl), and fructose-1,6-bisphosphatase (Fbp); (ii) fermentation: alcohol dehydrogenase (Adh); (iii) pentose phosphate pathway: 6-phosphogluconate dehydrogenase (Gnd), transketolase, and transaldolase; and (iv) Krebs and glyoxylate cycles: malate synthase.

(b) Protein synthesis, that include (i) ribosomal proteins, (ii) elongation factors: elongation factor 2 (Eft2), transcription elongation factor (Tef1); and (iii) chaperones: Ssa1 protein, Ssa2 protein, and heat shock proteins 60 (Hsp60) and 70 (Hsp70).

(c) Redox homeostasis: peroxiredoxin (Tsa), and peroxisomal catalase (Cta).

Most of the moonlighting proteins reported in pathogenic fungi working as virulence factors have been found on the cell surface, associated with the cell wall, or secreted. Thus, discrimination between proteins that have been indeed transported to the cell surface unconventionally and those that come from cell lysis or experimental artifacts needs to be done, to confirm their moonlighting status. Several methods have been used for this, which include protein immunolocalization at the cell wall in intact cells by flow cytometry ([Long et al., 2003](#); [Stie, Bruni & Fox, 2009](#); [Luo et al., 2013](#); [Dasari et al., 2019](#); [Gozalbo et al., 1998](#); [Gil-Navarro et al., 1997](#)), and electron and fluorescence microscopy ([Gozalbo et al., 1998](#); [Long et al., 2003](#); [Barbosa et al., 2006](#); [Nogueira et al., 2010](#); [Silveira et al., 2013](#); [da Silva Jde et al., 2013](#); [Crowe et al., 2003](#); [Angiolella et al., 1996](#)); surface protein biotinylation and precipitation ([López-Ribot et al., 1996](#); [Long et al., 2003](#); [Lopez et al., 2014](#); [Karkowska-Kuleta et al., 2011](#); [Kozik et al., 2015](#); [Karkowska-Kuleta et al., 2016](#)); isolation of the cell wall or cell wall components generating intact protoplasts ([Gil-Navarro et al., 1997](#); [Pitarch et al., 2002](#); [Li et al., 2003](#); [Castillo et al., 2008](#); [Longo et al., 2014](#); [García-Carnero et al., 2021](#)); and isolation of extracellular vesicles ([Vallejo et al., 2012](#)). As mentioned, the panorama of moonlighting proteins in fungi is limited when compared to bacteria, and a significant amount of information we currently have is on non-pathogenic fungal species, such as *Saccharomyces cerevisiae* ([Gancedo, Flores & Gancedo, 2016](#); [Flores & Gancedo, 2011](#); [Gancedo & Flores, 2008](#); [Rodríguez-Saavedra et al., 2021](#)). The information about moonlighting proteins in fungal pathogens of medical relevance is currently limited and searches in bibliography repositories indicate that the newest review manuscript on this subject was published in 2014 ([Karkowska-Kuleta & Kozik, 2014](#)), with a broad vision of eukaryotic pathogens, and when considering particular pathogenic species, the most recent review report is from 2020, gathering information about moonlighting proteins in the *C. albicans* cell wall ([Satala et al., 2020a](#)). Thus, a focused bibliographic search for moonlighting proteins in medically relevant fungal species has not been reported previously, and this review manuscript offers a thorough bibliographic revision on this subject ([Table 1](#)). Because of the reason above mentioned, this information will be of interest to the audience specialized in fungal biology, medical mycology, and medical microbiology. In a broader scope, it will be also of interest to the audience working in the cell biology and biochemistry fields, where moonlighting proteins are relevant too.

SURVEY METHODOLOGY

The PubMed (<https://pubmed.ncbi.nlm.nih.gov/>) and Google Scholar (<https://scholar.google.com/>) repositories were used to search for the following terms together with relevant Boolean Operators and MeSH terms identified for individual databases: moonlighting protein, moonlighting protein and fungi, moonlighting protein and fung*,

Table 1 Moonlighting proteins in medically relevant fungal species.

Protein <i>Candida</i> spp.	Canonical function	Moonlighting function	References
GAPDH	Glycolysis and gluconeogenesis	Binding to fibronectin, laminin, and plasminogen	Gozalbo <i>et al.</i> (1998), Crowe <i>et al.</i> (2003)
Eno1	Glycolysis and gluconeogenesis	Binding to plasmin, plasminogen, fibronectin, vitronectin, laminin, and kininogen Binding to medical devices	Jong <i>et al.</i> (2003), Kozik <i>et al.</i> (2015)
Gpm1	Glycolysis and gluconeogenesis	Binding to plasminogen, vitronectin, FH, FHL-1, and kininogen	Poltermann <i>et al.</i> (2007), Lopez <i>et al.</i> (2014)
Gpd2	Glycolysis and gluconeogenesis	Binding to plasminogen, FH, and FHL-1	Luo <i>et al.</i> (2013)
Adh1	Fermentation	Binding to plasminogen	Crowe <i>et al.</i> (2003)
Cta1	Redox homeostasis	Binding to plasminogen	Crowe <i>et al.</i> (2003)
Tef1	Elongation factor	Binding to plasminogen	Crowe <i>et al.</i> (2003), Poltermann <i>et al.</i> (2007)
Fba1 and Pgk1	Glycolysis and gluconeogenesis	Binding to plasminogen Binding to medical devices	Crowe <i>et al.</i> (2003); Poltermann <i>et al.</i> (2007)
Tsa1	Redox homeostasis	Binding to plasminogen, and kininogen	Crowe <i>et al.</i> (2003)
Eft2	Elongation factor	Binding to kininogen	Seweryn <i>et al.</i> (2015), Kozik <i>et al.</i> (2015)
Tpi1	Glycolysis and gluconeogenesis	Bind to kininogen, vitronectin, fibronectin, collagen, laminin, and elastin	Seweryn <i>et al.</i> (2015), Satala <i>et al.</i> (2021)
Gpi1	Glycolysis and gluconeogenesis	Binding to kininogen, fibronectin, vitronectin, and laminin	Seweryn <i>et al.</i> (2015), Kozik <i>et al.</i> (2015)
Gnd1	Pentose phosphate pathway	Binding to kininogen, fibronectin, vitronectin, and laminin	Seweryn <i>et al.</i> (2015), Kozik <i>et al.</i> (2015)
Fbp	Glycolysis and gluconeogenesis	Bind to fibronectin, vitronectin, and laminin	Kozik <i>et al.</i> (2015)
Malate synthase	Krebs and glyoxylate cycles	Bind to fibronectin, vitronectin, and laminin	Kozik <i>et al.</i> (2015)
Transketolase and transaldolase	Pentose phosphate pathway	Bind to fibronectin, vitronectin, and laminin	Kozik <i>et al.</i> (2015)
Ssa1	Chaperone	Binding to endothelial N-cadherin, epithelial E-cadherin, and histatin 5	Sun <i>et al.</i> (2010), Li <i>et al.</i> (2003)
Ssa2	Chaperone	Binding to idem	Sun <i>et al.</i> (2008), Li <i>et al.</i> (2003)
Als3	Adhesin	Ferritin receptor	Almeida <i>et al.</i> (2008)
<i>Paracoccidioides brasiliensis</i>			
Eno	Glycolysis and gluconeogenesis	Binding to laminin, fibronectin, plasminogen, and collagen type I and IV	Donofrio <i>et al.</i> , 2009, Nogueira <i>et al.</i> (2010)
Tpi	Glycolysis and gluconeogenesis	Binding to laminin	Pereira <i>et al.</i> (2007)
Malate synthase	Krebs and glyoxylate cycles	Binding to fibronectin, and collagen type I and IV Interaction with Eno and Tpi	da Silva Neto <i>et al.</i> (2009), de Oliveira <i>et al.</i> (2013)
GADPH	Glycolysis and gluconeogenesis	Binding to laminin, fibronectin, and collagen type I	Barbosa <i>et al.</i> (2006)
Fba	Glycolysis and gluconeogenesis	Binding to plasminogen Interaction with macrophages	Chaves <i>et al.</i> (2015)

(Continued)

Table 1 (continued)

Protein <i>Candida</i> spp.	Canonical function	Moonlighting function	References
14-3-3 protein	Regulation of many vital processes	Binding to laminin, and fibronectin	Andreotti et al. (2005) , Marcos et al. (2016)
<i>Histoplasma capsulatum</i>			
HIS-62	Chaperone	Binding to macrophage CR3 (CD11/CD18)	Long et al. (2003)
<i>Aspergillus fumigatus</i>			
Eno	Glycolysis and gluconeogenesis	Binding to plasminogen, FH, FHL-1, and C4BP Allergen	Dasari et al. (2019)
Tsl	Trehalose biosynthesis	Chitin synthase regulation	Thammahong et al. (2017)
<i>Cryptococcus neoformans</i>			
Hsp70	Chaperone	Macrophage and monocyte interaction and activation through the CD14 receptor Plasminogen binding	Asea et al. (2000) , Silveira et al. (2013)
Pgk, Fba, and pyruvate kinase	Glycolysis and gluconeogenesis	Binding to plasminogen	Stie, Bruni & Fox (2009)
Hsp60	Chaperone	Binding to plasminogen	Stie, Bruni & Fox (2009)
Transaldolase	Pentose phosphate pathway	Binding to plasminogen	Stie, Bruni & Fox (2009)
ATP-synthase alpha and beta subunits	Energy transduction	Binding to plasminogen	Stie, Bruni & Fox (2009)
Response to stress-related protein	Stress signaling pathways	Binding to plasminogen	Stie, Bruni & Fox (2009)
Glutamate dehydrogenase	Glutamate metabolism	Binding to plasminogen	Stie, Bruni & Fox (2009)
<i>Sporothrix schenckii</i>			
Hsp60	Chaperone	Binding to laminin, elastin, fibrinogen, and fibronectin	García-Carnero et al. (2021)

Note:

GAPDH, glyceraldehyde-3-phosphate dehydrogenase; Eno, enolase; Gpm, phosphoglycerate mutase; Gpd2, glycerol-3-phosphate dehydrogenase 2; Adh, alcohol dehydrogenase; Cta, peroxisomal catalase; Tef1, transcription elongation factor; Fba, fructose-bisphosphate aldolase; Pgk, phosphoglycerate kinase; Tsa, peroxiredoxin; Eft2, elongation factor 2; Tpi, triosephosphate isomerase; Gpi1, glucose-6-phosphate isomerase 1; Gnd, 6-phosphogluconate dehydrogenase; Fbp, fructose-1,6-bisphosphatase; Ssal and Ssa2: Hsp70, Als3: cell wall agglutinin-like sequence protein 3, HIS-60: Hsp60, Tsl, trehalose-6-phosphate phosphatase.

fungal cell wall and moonlighting protein, cell wall and moonlighting, *Candida* and moonlighting, *Paracoccidioides* and moonlighting, *Histoplasma* and moonlighting, *Aspergillus* and moonlighting, *Cryptococcus* and moonlighting, *Sporothrix* and moonlighting, yeast and moonlighting protein, hypha and moonlighting protein, filament cell and moonlighting protein, atypical cell wall protein, multifunctional protein and fung*, multitasking protein and fung*, and noncanonical cell wall protein.

Candida spp

Candida species are opportunistic fungal pathogens that most frequently establish the infective process in immunocompromised individuals, causing superficial and systemic infections, and representing an important health threat worldwide ([Spampinato & Leonardi, 2013](#); [Martínez-Duncker, Díaz-Jiménez & Mora-Montes, 2014](#)). The main reported species causing infection is *Candida albicans*, but recently, the incidence of other *Candida* species has increased, and these include *Candida glabrata*, *Candida tropicalis*,

Cell surface




Figure 1 Schematic representation of fungal moonlight proteins localization and function. Fungal moonlight proteins related to virulence are mostly adhesins with their moonlight function in the cell surface and their canonical function in the cytosol. An asterisk (*) indicates that Als3 and Tsl are moonlight proteins that perform both functions, canonical and moonlight, in the same cellular compartment.

Full-size DOI: 10.7717/peerj.14001/fig-1

Candida parapsilosis, *Candida krusei*, *Candida guilliermondii*, *Candida auris*, and *Candida lusitaniae* (Gómez-Gaviria & Mora-Montes, 2020; Tóth et al., 2019; Turner & Butler, 2014; Pappas et al., 2010). For the establishment of candidiasis, both pathogen and host-related factors are required and important. Many *Candida* virulence factors and determinants have been already described, including the expression of adhesins and secreted hydrolytic enzymes, dimorphism, and biofilm formation (Ciurea et al., 2020). The *Candida* cell wall contains canonical wall proteins found in other yeast-like organisms, but proteomic analyses have revealed the presence of cytosolic proteins in the cell wall (Pitarch et al., 2002; Castillo et al., 2008; Hernández et al., 2010; Gil-Bona et al., 2018), which can be classified according to their canonical functions into (i) enzymes involved in metabolic pathways, (ii) factors associated with protein synthesis, (iii) chaperones, (iv) and redox homeostasis enzymes (Satala et al., 2020a). Many of these proteins are still considered putative moonlighting proteins, but for some of them, the alternative function has been experimentally confirmed (Table 1 and Fig. 1).

In *C. albicans*, GAPDH protein found at the cell surface, especially on blastoconidia, participates in the adhesion to fibronectin, laminin, and plasminogen (Gozalbo et al., 1998; Crowe et al., 2003), contributing to fungal adhesion, dissemination, and damage to host tissue. The cell wall-associated GAPDH protein represents about 20–35% of the total amount of this protein in the cell and has been reported to be an important immunogenic antigen that elicits a humoral immune response (Gil-Navarro et al., 1997; Seidler, 2013). However, this protein does not seem to be a suitable therapeutic target against candidiasis, since active immunization with the recombinant protein, opsonization of yeasts with

anti-GAPDH antibodies, and passive transfer of polyclonal anti-GAPDH antibodies failed to generate a protective immune response in the host ([Gil et al., 2006](#)).

Eno1 has been characterized as a cell surface plasmin(ogen)-binding protein that enhances fibrinolysis and the fungal ability to cross the blood-brain barrier *in vitro*, participating in tissue invasion and dissemination ([Jong et al., 2003](#)). Deletion of its gene impairs *C. albicans* growth on glucose-containing media, alters hyphal formation, increases drug susceptibility, and eliminates the fungal ability to kill the host ([Ko et al., 2013](#)). This protein binds to the cell wall glucans, being tightly attached to the wall's inner layer ([Angiolella et al., 1996](#)). Although Eno1 is an immunodominant antigen that stimulates cellular and humoral responses during infection ([Sundstrom, Jensen & Balish, 1994](#)), its recombinant version is only capable of providing modest protection against candidiasis ([Montagnoli et al., 2004](#)). Intracellularly, this protein has been reported to present two enzymatic activities given by two different catalytic sites, as enolase, which is essential for the glycolytic and gluconeogenesis pathways ([Satala et al., 2020b](#)), and as transglutaminase, participating in fungal growth and morphogenesis, cell division and osmotic regulation ([Reyna-Beltrán et al., 2018](#)). Blocking of Eno1 transglutaminase activity, alone or in combination with fluconazole, provided an antifungal activity against *C. albicans* *in vitro* and *in vivo*, proposing this protein as a putative therapeutic target ([Li et al., 2019](#)). In addition, human Eno has also been found anchored on the cell surface of human cells, where it binds to plasminogen and represents a local fibrinolysis regulator ([Miles et al., 1991](#)). This moonlight function seems to be conserved among organisms, and is has been proven, at least in humans and some pathogenic organisms, to be due to the presence of lysine residues on the C-terminal end of the protein, which appear to function as plasminogen and extracellular matrix (ECM) components binding sites ([Miles et al., 1991; Bergmann et al., 2003; Derbise et al., 2004; Vanegas et al., 2007; Rahi et al., 2018; Satala et al., 2020a](#)).

The *C. albicans* alcohol dehydrogenase 1 (Adh1) catalyzes the conversion of acetaldehyde to ethanol and is also involved in the NADH-dependent methylglyoxal dehydrogenase activity that generates pyruvate ([Kwak, Ku & Kang, 2014](#)). The *adh1Δ* null mutants showed an increment in the methylglyoxal levels, growth defects, and virulence attenuation ([Kwak, Ku & Kang, 2014](#)).

Gpm1 protein is found on the *C. albicans* yeast and hypha cell surface, where it binds to plasminogen, vitronectin, and the host complement regulators Factor H (FH) and Factor H-like binding protein (FHL-1), facilitating attachment to endothelial cells and keratinocytes, degradation of the host ECM, and promoting immune evasion ([Poltermann et al., 2007; Lopez et al., 2014](#)). *GMP1* gene deletion decreased fungal binding to endothelial cells, confirming the importance of the encoded protein for *C. albicans* virulence ([Lopez et al., 2014](#)). Also, *C. albicans* secreted Gpd2 protein constitutes a plasminogen-, FH-, and FHL-1-binding protein ([Luo et al., 2013](#)).

Eno1 protein, GAPDH protein, and Gpm1 protein, along with the moonlighting proteins Adh1, Tsa1, and Cta1 represent the major plasminogen-binding proteins, contributing to 85% of the plasminogen binding in *C. albicans* cell wall protein extracts

([Crowe et al., 2003](#)). Tef1 protein, Pgk1 protein, and Fba1 protein also participate in plasminogen binding, though to a lesser extent ([Crowe et al., 2003; Poltermann et al., 2007](#)).

C. albicans Eno1 protein, Gmp1 protein, Gpi1 protein, Eft2 protein, Tpi1 protein, Ts1 protein, and Gnd1 protein are atypical cell wall proteins and have all been identified as kininogen-binding proteins ([Karkowska-Kuleta et al., 2011; Seweryn et al., 2015](#)). Kininogen is an important protein in a proteolytic cascade of human plasma that participates in the host pro-inflammatory, antimicrobial, and anti-adhesive responses ([Lalmanach et al., 2010](#)). *C. albicans* and *C. glabrata* Tpi1 protein has also been proven to bind to vitronectin, fibronectin, collagen, laminin, and elastin, suggesting its contribution to fungal adhesion ([Satala et al., 2021](#)).

C. parapsilosis and *C. tropicalis* pseudohyphae also have cell surface moonlighting adhesins that include malate synthase, Gpi1 protein, Gnd1 protein, Eno1 protein, Fbp protein, transketolase, transaldolase, and Eft2 protein, which bind to fibronectin, vitronectin, and laminin ([Kozik et al., 2015](#)). In addition, *C. tropicalis* Eno1 protein and Gpm1 protein, and *C. parapsilosis* Gnd1 protein also bind to high-molecular-mass kininogen ([Karkowska-Kuleta et al., 2016; Karkowska-Kuleta et al., 2017](#)).

These moonlighting proteins not only mediate the binding of *Candida* spp. cells to host components but also to medical devices, which is an important trait during biofilm formation. Eno1 was found to participate in the adhesion of *C. albicans*, *C. glabrata*, *C. krusei*, and *C. parapsilosis* to plastic surfaces; Fba1 protein favors the adhesion of *C. albicans*, *C. glabrata*, and *C. krusei*; and Pgk1 protein is involved in the adhesion of *C. parapsilosis* to different components of medical devices ([Núñez-Beltrán, López-Romero & Cuéllar-Cruz, 2017](#)).

Ssa1, a member of the Hsp70 family, has been localized throughout the *C. albicans* cell wall ([López-Ribot et al., 1996](#)), where it binds to endothelial cells N-cadherin and epithelial cells E-cadherin, thus being essential for host cell endocytosis and invasion ([Sun et al., 2010](#)). SSA1 gene deletion decreased *C. albicans* adhesion, damage, and invasion of the epithelium and endothelium, resulting in lower virulence ([Sun et al., 2010](#)), confirming the important role of this protein during infection. However, Ssa1 and Ssa2 proteins, another Hsp70, are reported to also have a detrimental effect on the fungus, due to their ability to bind to histatin 5, a salivary histidine-rich antifungal peptide, that causes cell death when adsorbed to the cell surface ([Sun et al., 2008; Li et al., 2003, 2006](#)). Ssa1 and Ssa2 proteins are considered the main immunogenic antigens in the *C. albicans* cell wall and are capable of eliciting cell-mediated and humoral immune responses ([Martínez et al., 1998; López-Ribot et al., 2004](#)). However, vaccination with the recombinant proteins failed to protect against candidiasis, but instead enhanced the infection ([Bromuro et al., 1998](#)).

Als3 protein, a member of the *C. albicans* cell wall agglutinin-like sequence (Als) family ([Hoyer et al., 1998](#)), is an important multifunctional protein in the hyphae and pseudohyphae surface ([Hoyer et al., 1998; Argimón et al., 2007](#)). Als3 has a canonical function of adhesin and invasin that binds to gelatin, fibronectin, fibrinogen, type IV collagen, laminin, salivary pellicle, and E- and N-cadherin, thus mediating adherence and internalization to endothelial and epithelial cells ([Sheppard et al., 2004; Phan et al., 2007; Nobbs, Vickerman & Jenkinson, 2010](#)), and also plays an important role in biofilm

formation (Nobile *et al.*, 2006; Zhao *et al.*, 2006). However, Als3 protein has an additional function as a ferritin receptor in *C. albicans* hyphae, where it binds to purified ferritin and ferritin contained within epithelial cells, helping the fungus to obtain iron from the host (Almeida *et al.*, 2008). ALS3 gene deletion decreased fungal adherence to endothelial and oral epithelial cells (Zhao *et al.*, 2004), decreased endocytosis by oral epithelial cells and vascular endothelial cells *in vitro*, altered biofilm structures (Nobile *et al.*, 2006; Zhao *et al.*, 2006), and diminished iron acquisition *in vitro* (Almeida *et al.*, 2008), demonstrating its multiple functions and their importance during infection. Als3 protein is also an immunodominant antigen, and vaccination with its recombinant version protected against vaginal and disseminated candidiasis (Spellberg *et al.*, 2006).

Paracoccidioides brasiliensis

The thermodimorphic fungi *P. brasiliensis* and *Paracoccidioides lutzii* are the etiological agents of paracoccidioidomycosis, an important human systemic disease with multiple clinical forms, endemic to Latin America (San-Blas, Niño-Vega & Iturriaga, 2002; Mendes *et al.*, 2017). The interaction of these fungi with their host and their successful colonization depends on many regulatory mechanisms and virulence factors that allow them to cause infection, which include morphogenesis, changes in cell wall polysaccharide composition, adhesion, adaptation to environmental stresses, and production of extracellular vesicles (de Oliveira *et al.*, 2015; Camacho & Niño-Vega, 2017).

Proteomic analyses of *P. brasiliensis* yeasts and mycelium cell walls, and vesicle and vesicle-free extracellular content have revealed the presence of many cytoplasmic proteins, mostly related to carbohydrate and protein metabolism, stress response, oxidation/reduction, translation, nucleic acid binding, and cellular architecture (Table 1) (Vallejo *et al.*, 2012; Longo *et al.*, 2014). Also, proteomic analysis of the *Paracoccidioides* secretome identified 15 plasminogen binding proteins, many of which are cytoplasmic proteins (Chaves *et al.*, 2015). Out of these proteins, some have already been proven to be moonlighting proteins that participate as adhesins during infection (de Oliveira *et al.*, 2015; Marcos *et al.*, 2014). Eno protein, Fba protein, GADPH protein, Tpi protein, and malate synthase have been found in the *Paracoccidioides* cell wall and extracellular vesicles, mediating adhesion to the host tissues (Vallejo *et al.*, 2012). Eno protein was found to be highly expressed when *P. brasiliensis* grows in the presence of blood and has the ability to bind to laminin, fibronectin, plasminogen, and type I and IV collagens (Donofrio *et al.*, 2009; Nogueira *et al.*, 2010; Marcos *et al.*, 2012), representing 80% of fungal adhesion to epithelial cells (Donofrio *et al.*, 2009), which demonstrates its essential role in fungal attachment, internalization, and invasion to the host tissues. Tpi protein is a yeast cytosolic and cell wall antigen that reacts with sera of patients with paracoccidioidomycosis and that participates in fungal binding to epithelial cells, with affinity to laminin (da Fonseca *et al.*, 2001; Pereira *et al.*, 2007). Malate synthase is also a *P. brasiliensis* cell wall and secreted protein that functions as an adhesin that binds to fibronectin and type I and IV collagens (da Silva Neto *et al.*, 2009). In addition, this protein interacts with other cell wall adhesins, such as Eno protein and Tpi protein, enhancing the fungal adhesion (de Oliveira *et al.*, 2013). GAPDH protein was found in *P. brasiliensis* extracellular vesicles and cell wall, with

increased expression in the yeast morphology and during the mycelium-yeast transition ([Barbosa et al., 2006, 2004; Longo et al., 2014](#)). On the cell surface, this protein binds to laminin, fibronectin, and type-I collagen, being involved in the fungal adhesion to pneumocytes during the early stages of infection ([Barbosa et al., 2006](#)). Fab protein binds to plasminogen and participates in the fungal interaction with macrophages, increasing fibrin degradation and internalization by phagocytes ([Chaves et al., 2015](#)).

A cytosolic 30 kDa glycoprotein, found highly expressed in the *P. brasiliensis* cell wall during infection and in high virulent strains, was identified as a member of the 14-3-3 protein family ([da Silva Jde et al., 2013; Andreotti et al., 2005](#)) of conserved regulatory eukaryotic proteins involved in transcription, signal transduction, protein localization and degradation, cell cycle, apoptosis, and many other processes ([Sluchanko & Gusev, 2017](#)). The cell surface 14-3-3 protein is an important *Paracoccidioides* adhesin that binds to laminin and fibronectin ([Andreotti et al., 2005; Marcos et al., 2016](#)), and the silencing of its gene affected the yeast morphology, impaired morphological switching, and decreased laminin binding, resulting in a significant virulence reduction ([Marcos et al., 2016](#)), underlining its importance to the *P. brasiliensis* virulence.

Histoplasma capsulatum

H. capsulatum is an environmental thermodimorphic fungus with two pathogenic varieties, *H. capsulatum var capsulatum*, and *H. capsulatum var duboissi*, that cause human histoplasmosis ([Antinori, 2014](#)), a worldwide distributed pulmonary and systemic disease with a mortality rate up to 7% ([Kauffman, 2007; Armstrong et al., 2018](#)). Disease development and severity depend on the host immune status, virulence of the fungal strain, and the fungal load inoculated ([Knox & Hage, 2010](#)). Only a handful of *H. capsulatum* virulence factors have been identified, and these include the presence of α -1,3-glucan at the cell surface, thermotolerance, melanin, and surface ligands for immune receptors ([Mihu & Nosanchuk, 2012](#)). Thus far, only one of these virulence factors has been proven to be a moonlighting protein ([Table 1](#)).

At the first stage of histoplasmosis infection, *H. capsulatum* gets recognized and internalized by alveolar macrophages through the integrin receptor CR3 (CD11/CD18). Once inside, this fungus impairs macrophage activation and phagolysosomal fusion, and replicates, killing the alveolar macrophages and invading neighboring and recruited macrophages, thus disseminating to other organs from the lungs ([Bullock & Wright, 1987; Newman et al., 1990](#)). This defective macrophage response against *H. capsulatum* depends, in part, on the receptor that the fungus uses to enter phagocytes, and Hsp60, also known as HIS-62, has been reported to be the major *H. capsulatum* surface protein that interacts with CD11/CD18 ([Long et al., 2003](#)). This intracellular chaperone has been found moonlighting on the outer surface of the *H. capsulatum* cell wall, and although HIS-62 protein lacks the classical secretion signal, it shows a unique leader sequence upstream of the amino terminus of the mature protein, with high homology to leader sequences of other pathogenic fungi ([Long et al., 2003](#)), which might be related to its secretion. Also, the cell wall-associated Hsp60 was found to be *N*-linked glycosylated, while the intracellular

protein is not, suggesting that the surface HIS-62 protein enters the secretory pathway ([Long et al., 2003](#)).

Hsp60 has been reported as an immunogenic antigen during histoplasmosis, and antibodies against this protein provided a protective immune response in the host, by reducing the fungal burden and increasing phagolysosomal fusion *in vitro* ([Guimarães et al., 2009](#)). Also, vaccination with the native and recombinant HIS-62 protected against pulmonary and systemic histoplasmosis, which suggests that this protein is an important target for the cellular and humoral immune response ([Gomez, Gomez & Deepe, 1991](#); [Gomez, Allendoerfer & Deepe, 1995](#)).

Aspergillus fumigatus

Aspergillus species are environmental molds found worldwide ([Hope, Walsh & Denning, 2005](#)). *A. fumigatus* is the most common species associated with aspergillosis syndromes ([Park & Mehrad, 2009](#)), which range from allergic noninvasive clinical forms in immunocompetent patients to systemic disease under immunosuppression conditions, with mortality rates up to 100% if the diagnosis is missed or delayed ([Brown et al., 2012](#); [Latgé & Chamilos, 2019](#); [Barnes & Marr, 2006](#)). The *Aspergillus* spp. pathogenicity is dependent on the host immune response and pulmonary microbiome, and on the presence of fungal virulence factors that might be in part enhanced by some host effectors ([Raksha, Singh & Urhekar, 2017](#); [Kolwijk & van de Veerdonk, 2014](#)). Different *A. fumigatus* virulence factors have been identified, which include adhesins, pigments, hydrolytic enzymes, catalases and superoxide dismutases, mycotoxins and non-protein metabolites, thermotolerance, allergens, and zinc acquisition ([Raksha, Singh & Urhekar, 2017](#); [Szalewski et al., 2018](#)).

One of the main *A. fumigatus* adhesin and allergen is the moonlighting protein Eno ([Table 1](#)), expressed at the swollen conidia and hypha cell surface, where it binds to plasminogen and represents a ligand for FH, FHL-1, and C4BP (C4 binding protein) ([Dasari et al., 2019](#)). The affinity of this protein to the plasma complement regulators contributes to the fungal immune evasion, since its binding to FH and FHL-1, and C4PB inactivates C3b and C4b, respectively, while binding to plasminogen degrades C3 and C3b. Thus, Eno protein contributes to cell attachment and invasion, tissue damage, and immune evasion ([Dasari et al., 2019](#)). Eno protein is also an important *Aspergillus* allergen that causes a hypersensitive immune reaction in the host and is highly recognized by IgE ([Lai et al., 2002](#); [Gautam et al., 2007](#); [Shah & Panjabi, 2014](#)).

The *A. fumigatus* Tsl protein was identified as an important enzyme for trehalose biosynthesis and cell wall structure. This protein directly interacts with chitin synthase and regulates its activity and localization, as a moonlight function, affecting the cell wall integrity and structure ([Thammahong et al., 2017](#)). Deletion of its encoding gene reduced trehalose content in conidia and mycelium but also altered the cell wall integrity, by increasing chitin, β-glucan, and galactosaminogalactan content and exposure ([Thammahong et al., 2017](#)). Also, these mutants trigger a much stronger pro-inflammatory response in the host, probably due to the exposure of those cell wall components at the cell surface, but without affecting the fungal virulence ([Thammahong et al., 2017](#)).

Other pathogenic fungi

The encapsulated yeast *C. neoformans* is an opportunistic fungus that can cause cryptococcal meningoencephalitis in immunocompromised patients (Brown *et al.*, 2012). Macrophages play an important role in fungal killing, but also in the replication and dissemination of *C. neoformans* (García-Rodas & Zaragoza, 2012), for which fungal surface components that interact and attach to phagocytes are essential for *C. neoformans* pathogenesis. Hsp70 has been detected on the *C. neoformans* cell surface where it is involved in macrophage and monocyte interaction and activation, through the CD14 receptor (Table 1) (Asea *et al.*, 2000; Silveira *et al.*, 2013). *C. neoformans* Hsp70 inhibits macrophage activation, thus helping the fungus to evade killing by the phagolysosome oxidative response, and also promotes attachment to epithelial cells through a still unidentified receptor that might be shared with glucuronoxylosmannan, the main capsular antigen (Silveira *et al.*, 2013). In addition, many cytosolic proteins with plasminogen affinity have been identified on the *C. neoformans* cell surface, including Pfk protein, Fba protein, Hsp60, Hsp70, transaldolase, pyruvate kinase, ATP-synthase alpha and beta subunits, the response to stress-related protein, and glutamate dehydrogenase (Stie, Bruni & Fox, 2009).

The thermotolerant fungal species of the *Sporothrix* pathogenic clade are the causative agents of sporotrichosis, a human and animal subcutaneous mycosis that can disseminate and cause systemic infections (Lopes-Bezerra *et al.*, 2018; Lopez-Romero *et al.*, 2011; Mora-Montes *et al.*, 2015; Nava-Pérez *et al.*, 2022). *Sporothrix schenckii* and *Sporothrix brasiliensis* are the most commonly isolated species and also the most studied, but only a few of their virulence factors and determinants have been identified, which include adhesins, peptidoglycan, morphological switching, and melanin (Tamez-Castrellon *et al.*, 2020; García-Carnero & Martínez-Álvarez, 2022). Some intracellular proteins, such as Hsp60, Hsp70, Tef1 protein, mitochondrial peroxiredoxin, lipase 1, Eno1 protein, and pyruvate kinase, have been found in cell wall preparations from *S. schenckii* cells growing in the presence of different stressors (Ruiz-Baca *et al.*, 2019), however, because the preparations were performed using whole-cell homogenates, it remains to be confirmed whether these are indeed proteins found on the *S. schenckii* surface with moonlighting functions. Proteomic analysis of the *S. schenckii* yeast-like cells' peptidoglycan revealed that it is composed of 325 possible proteins, many of which are intracellular proteins with housekeeping functions and without conventional signal sequences, that may be moonlighting proteins related to virulence on the cell surface (García-Carnero *et al.*, 2021). One of the main peptidoglycan proteins is Hsp60, which was proven to function as a cell surface adhesin with binding affinity to laminin, elastin, fibrinogen, and fibronectin, being essential for fungal virulence (Table 1) (García-Carnero *et al.*, 2021). Also, this protein appears to be highly immunogenic, and immunization with its recombinant version or treatment with anti-Hsp60 antibodies can protect laboratory animals against a lethal infection with *S. schenckii* (García-Carnero *et al.*, 2021).




Figure 2 Schematic representation of the host ligands that bind to the different fungal moonlighting proteins. The host ligands are located in many cell types and compartments, and many of the pathogen moonlighting proteins are capable of binding to more than one ligand, facilitating the attachment and dissemination of the fungus to different tissues. All the proteins are present on the cell surface of the pathogenic fungus, which is arbitrarily represented by yeast cells. GAPDH, glyceraldehyde-3-phosphate dehydrogenase; Eno, enolase; Gpm, phosphoglycerate mutase; Gpd2, glycerol-3-phosphate dehydrogenase 2; Adh, alcohol dehydrogenase; Cta, peroxisomal catalase; Tef1, transcription elongation factor; Fba, fructose-bisphosphate aldolase; Pgk, phosphoglycerate kinase; Tsa, peroxiredoxin; Eft2, elongation factor 2; Tpi, triosephosphate isomerase; Gpi1, glucose-6-phosphate isomerase 1; Gnd, 6-phosphogluconate dehydrogenase; Fbp, fructose-1, 6-bisphosphatase; Ssa1 and Ssa2: Hsp70, Als3: cell wall agglutinin-like sequence protein 3; HIS-60: Hsp60.

[Full-size](#) DOI: 10.7717/peerj.14001/fig-2

The host perspective

It is widely known that the adherence of a pathogen to its host represents an essential step to invade, colonize, evade the immune response, and cause infection, as already mentioned. Most of the moonlighting proteins that participate in fungal virulence processes functions as adhesins that bind to important components of the host ECM and of homeostatic and immune cascades (Fig. 2).

The ECM is a dynamic network of macromolecules that provides structural support for cells and tissues, composed mainly of proteoglycans and glycosaminoglycans, elastin and elastic fibers, collagens, laminins, fibronectin, thrombospondin, and vitronectin (Karamanos *et al.*, 2021). Through signal transduction, the ECM regulates many cellular functions, that include growth, proliferation, cell migration, and differentiation, playing an essential role in cell homeostasis (Karamanos *et al.*, 2021). Therefore, disruption of any of the ECM components alters tissue function and regeneration, thus facilitating infection by pathogenic fungi (Tomlin & Piccinini, 2018). Disruption of the ECM during infection can be caused directly by the degradation of its components after attachment through the fungus proteases, elastases, or collagenases; or indirectly through the host catabolic machinery that gets altered upon infection or by the activity of activated macrophages that destroy the ECM (Tomlin & Piccinini, 2018). In addition, fungal binding to plasminogen causes its cleavage to plasmin, an important protease of the fibrinolytic system that degrades fibrin (Castellino & Ploplis, 2005), which participates in blood coagulation, cell

migration, and tissue repair; activates collagenases that degrade collagens and activates complement mediators; and degrades several ECM components, such as fibronectin, laminin, and thrombospondin, thus degrading the host cell barriers and ECM (Singh *et al.*, 2012). Finally, the ECM also participates in the host immune response, by providing important signaling to the immune cells for their proliferation, differentiation, activation, and migration, which gets altered with the ECM disruption (Tomlin & Piccinini, 2018).

High molecular weight kininogen is a member of the human contact system, that upon activation cleaves kininogen and releases antimicrobial peptides and bradykinin, involved in the regulation of inflammatory processes, vascular permeability, and blood pressure (Ponczek, 2021). The binding of pathogenic fungi to kininogens through different ligands triggers inflammatory responses through bradykinin, resulting in vascular leakage, vasodilatation, and increased blood flow, enabling the inflow of plasma nutrients, and enhancing pathogen dissemination and invasion (Karkowska-Kuleta & Kozik, 2014; Oehmcke-Hecht & Köhler, 2018).

The complement system is an important effector of the innate immune response, that participates in the elimination of pathogenic organisms through proinflammatory responses, which include attraction and activation of phagocytes, and enhanced opsonization (Friese *et al.*, 1999). The plasma proteins FH and FHL-1 are regulatory elements that control the alternative complement activation at the level of C3b, an active complement-derived product, and the binding of pathogenic fungi to these regulators mediates immune evasion and disruption of the complement activation (Behnse *et al.*, 2008), thus promoting invasion and dissemination.

CONCLUSIONS

Although long known, moonlighting proteins have received little attention when participating in virulence processes of pathogenic fungi. To date, many moonlighting proteins have been suggested to function as fungal virulence factors, but only a few have been experimentally confirmed, with very limited information about their alternative functions, the mechanisms they use to perform them, and their cellular localization and trafficking. The information that we know about these moonlighting functions during infection highlights the importance these proteins have in fungal pathogenesis, by enabling fungal attachment, invasion, and dissemination in the host, along with helping with the fungal immune evasion. Therefore, they are key factors for understanding fundamental and clinically relevant biological processes in these organisms. Even though their study should be a priority, it has been neglected. *C. albicans* is perhaps the most studied species in medical mycology, and therefore, most of the available information about moonlighting proteins has been generated on this organism. However, as we know, fungal infections are not reduced to only candidemia/candidiasis, other fungal pathogens are relevant in this field. As reported in this review article, the information about moonlighting proteins in other fungal species is limited when compared with that generated in *Candida* species, stressing that more efforts are required to understand the contribution of these proteins to fungal pathogenesis, in a broader sense.

These proteins represent an example of genetic economy and directed evolution, being a low-cost adaptation mechanism for the cell when facing stressful conditions, such as those generated in the host during infection, and at the same time, contributing to the fungal virulence. As reviewed in this text, a great percentage of the reported fungal moonlighting proteins are conserved metabolic enzymes and chaperones, with alternative functions related to adhesion, tissue damage, and immunoevasion. The moonlighting function may likely have an impact on other processes during the interaction with the host, such as cell wall remodeling, posttranscriptional regulatory events, apoptosis, and cancer development, as reported for bacterial moonlighting proteins ([Podobnik et al., 2009](#); [Banerjee et al., 2007](#); [Lin et al., 2010](#); [Basak et al., 2005](#)). Thus, this represents an area of opportunity and a challenge in the years to come.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by Consejo Nacional de Ciencia y Tecnología (FC 2015-02-834 and Ciencia de Frontera 2019-6380), and Red Temática Glicociencia en Salud (CONACYT-México). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Consejo Nacional de Ciencia y Tecnología: FC 2015-02-834.
Ciencia de Frontera: 2019-6380.
Red Temática Glicociencia en Salud (CONACYT-México).

Competing Interests

Héctor M. Mora-Montes is an Academic Editor for PeerJ.

Author Contributions

- Verania J. Arvizu-Rubio performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Laura C. García-Carnero conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Héctor Manuel Mora-Montes conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the article, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

This is a review article.

REFERENCES

- Aharoni A, Gaidukov L, Khersonsky O, Gould SMQ, Roodveldt C, Tawfik DS. 2005. The 'evolvability' of promiscuous protein functions. *Nature Genetics* 37(1):73–76 DOI [10.1038/ng1482](https://doi.org/10.1038/ng1482).
- Almeida RS, Brunke S, Albrecht A, Thewes S, Laue M, Edwards JE Jr., Filler SG, Hube B. 2008. The Hyphal-associated adhesin and invasin Als3 of *Candida albicans* mediates iron acquisition from host ferritin. *PLOS Pathogens* 4(11):e1000217 DOI [10.1371/journal.ppat.1000217](https://doi.org/10.1371/journal.ppat.1000217).
- Amblee V, Jeffery CJ. 2015. Physical features of intracellular proteins that moonlight on the cell surface. *PLOS ONE* 10(6):e0130575 DOI [10.1371/journal.pone.0130575](https://doi.org/10.1371/journal.pone.0130575).
- Andreotti PF, Monteiro da Silva JL, Bailão AM, Soares CM, Benard G, Soares CP, Mendes-Giannini MJ. 2005. Isolation and partial characterization of a 30 kDa adhesin from *Paracoccidioides brasiliensis*. *Microbes and Infection* 7(5–6):875–881 DOI [10.1016/j.micinf.2005.02.005](https://doi.org/10.1016/j.micinf.2005.02.005).
- Angioletta L, Facchin M, Stringaro A, Maras B, Simonetti N, Cassone A. 1996. Identification of a glucan-associated enolase as a main cell wall protein of *Candida albicans* and an indirect target of lipopeptide antimycotics. *Journal of Infectious Diseases* 173(3):684–690 DOI [10.1093/infdis/173.3.684](https://doi.org/10.1093/infdis/173.3.684).
- Antinori S. 2014. *Histoplasma capsulatum*: more widespread than previously thought. *American Journal of Tropical Medicine and Hygiene* 90(6):982–983 DOI [10.4269/ajtmh.14-0175](https://doi.org/10.4269/ajtmh.14-0175).
- Argimón S, Wishart JA, Leng R, Macaskill S, Mavor A, Alexandris T, Nicholls S, Knight AW, Enjalbert B, Walmsley R, Odds FC, Gow NA, Brown AJ. 2007. Developmental regulation of an adhesin gene during cellular morphogenesis in the fungal pathogen *Candida albicans*. *Eukaryotic Cell* 6(4):682–692 DOI [10.1128/ec.00340-06](https://doi.org/10.1128/ec.00340-06).
- Armstrong PA, Jackson BR, Haselow D, Fields V, Ireland M, Austin C, Signs K, Fialkowski V, Patel R, Ellis P, Iwen PC, Pedati C, Gibbons-Burgener S, Anderson J, Dobbs T, Davidson S, McIntyre M, Warren K, Midla J, Luong N, Benedict K. 2018. Multistate epidemiology of histoplasmosis, United States, 2011–2014. *Emerging Infectious Diseases* 24(3):425–431 DOI [10.3201/eid2403.171258](https://doi.org/10.3201/eid2403.171258).
- Asea A, Kraeft SK, Kurt-Jones EA, Stevenson MA, Chen LB, Finberg RW, Koo GC, Calderwood SK. 2000. HSP70 stimulates cytokine production through a CD14-dependant pathway, demonstrating its dual role as a chaperone and cytokine. *Nature Medicine* 6(4):435–442 DOI [10.1038/74697](https://doi.org/10.1038/74697).
- Banerjee S, Nandyala AK, Raviprasad P, Ahmed N, Hasnain SE. 2007. Iron-dependent RNA-binding activity of *Mycobacterium tuberculosis* aconitase. *Journal of Bacteriology* 189(11):4046–4052 DOI [10.1128/jb.00026-07](https://doi.org/10.1128/jb.00026-07).
- Barbosa MS, Bão SN, Andreotti PF, de Faria FP, Felipe MSS, dos Santos Feitosa L, Mendes-Giannini MJS, de Almeida Soares CM. 2006. Glyceraldehyde-3-phosphate dehydrogenase of *Paracoccidioides brasiliensis* is a cell surface protein involved in fungal adhesion to extracellular matrix proteins and interaction with cells. *Infection and Immunity* 74(1):382–389 DOI [10.1128/iai.74.1.382-389.2006](https://doi.org/10.1128/iai.74.1.382-389.2006).
- Barbosa MS, Passos DAC, Felipe MSS, Jesuino RSA, Pereira M, de Almeida Soares CM. 2004. The glyceraldehyde-3-phosphate dehydrogenase homologue is differentially regulated in phases of *Paracoccidioides brasiliensis*: molecular and phylogenetic analysis. *Fungal Genetics and Biology* 41(7):667–675 DOI [10.1016/j.fgb.2004.02.002](https://doi.org/10.1016/j.fgb.2004.02.002).
- Barnes PD, Marr KA. 2006. Aspergillosis: spectrum of disease, diagnosis, and treatment. *Infectious Disease Clinics of North America* 20(3):545–561 DOI [10.1016/j.idc.2006.06.001](https://doi.org/10.1016/j.idc.2006.06.001).

- Basak C, Pathak SK, Bhattacharyya A, Pathak S, Basu J, Kundu M.** 2005. The secreted peptidyl prolyl cis,trans-isomerase HP0175 of *Helicobacter pylori* induces apoptosis of gastric epithelial cells in a TLR4- and apoptosis signal-regulating kinase 1-dependent manner. *Journal of Immunology* 174(9):5672–5680 DOI 10.4049/jimmunol.174.9.5672.
- Behnson J, Hartmann A, Schmaler J, Gehrke A, Brakhage AA, Zipfel PF.** 2008. The opportunistic human pathogenic fungus *Aspergillus fumigatus* evades the host complement system. *Infection and Immunity* 76(2):820–827 DOI 10.1128/iai.01037-07.
- Bergmann S, Wild D, Diekmann O, Frank R, Bracht D, Chhatwal GS, Hammerschmidt S.** 2003. Identification of a novel plasmin(ogen)-binding motif in surface displayed alpha-enolase of *Streptococcus pneumoniae*. *Molecular Microbiology* 49(2):411–423 DOI 10.1046/j.1365-2958.2003.03557.x.
- Bromuro C, La Valle R, Sandini S, Urbani F, Ausiello CM, Morelli L, d'Ostiani Fé C, Romani L, Cassone A.** 1998. A 70-kilodalton recombinant heat shock protein of *Candida albicans* is highly immunogenic and enhances systemic murine candidiasis. *Infection and Immunity* 66(5):2154–2162 DOI 10.1128/iai.66.5.2154-2162.1998.
- Brown GD, Denning DW, Gow NA, Levitz SM, Netea MG, White TC.** 2012. Hidden killers: human fungal infections. *Science Translational Medicine* 4(165):165rv13 DOI 10.1126/scitranslmed.3004404.
- Bullock WE, Wright SD.** 1987. Role of the adherence-promoting receptors, CR3, LFA-1, and p150,95, in binding of *Histoplasma capsulatum* by human macrophages. *Journal of Experimental Medicine* 165(1):195–210 DOI 10.1084/jem.165.1.195.
- Camacho E, Niño-Vega GA.** 2017. Paracoccidioides spp.: virulence factors and immune-evasion strategies. *Mediators Inflamm* 2017(6):5313691 DOI 10.1155/2017/5313691.
- Castellino FJ, Ploplis VA.** 2005. Structure and function of the plasminogen/plasmin system. *Thromb Haemost* 93(4):647–654 DOI 10.1160/th04-12-0842.
- Castillo L, Calvo E, Martínez AI, Ruiz-Herrera J, Valentín E, Lopez J, Sentandreu R.** 2008. A study of the *Candida albicans* cell wall proteome. *Proteomics* 8(18):3871–3881 DOI 10.1002/pmic.200800110.
- Chaves EG, Weber SS, Bão SN, Pereira LA, Bailão AM, Borges CL, Soares CM.** 2015. Analysis of *Paracoccidioides* secreted proteins reveals fructose 1, 6-bisphosphate aldolase as a plasminogen-binding protein. *BMC Microbiology* 15(1):53 DOI 10.1186/s12866-015-0393-9.
- Chen C, Liu H, Zabad S, Rivera N, Rowin E, Hassan M, Gomez De Jesus SM, Llinás Santos PS, Kravchenko K, Mikhova M, Ketterer S, Shen A, Shen S, Navas E, Horan B, Raudsepp J, Jeffery C.** 2021. MoonProt 3.0: an update of the moonlighting proteins database. *Nucleic Acids Research* 49(D1):D368–D372 DOI 10.1093/nar/gkaa1101.
- Ciurea CN, Kosovski IB, Mare AD, Toma F, Pintea-Simon IA, Man A.** 2020. *Candida* and candidiasis-opportunism versus pathogenicity: a review of the virulence traits. *Microorganisms* 8(6):857 DOI 10.3390/microorganisms8060857.
- Copley SD.** 2012. Moonlighting is mainstream: paradigm adjustment required. *Bioessays* 34(7):578–588 DOI 10.1002/bies.201100191.
- Crowe JD, Sievwright IK, Auld GC, Moore NR, Gow NA, Booth NA.** 2003. *Candida albicans* binds human plasminogen: identification of eight plasminogen-binding proteins. *Molecular Microbiology* 47(6):1637–1651 DOI 10.1046/j.1365-2958.2003.03390.x.
- da Fonseca CA, Jesuino Rália SA, Felipe MSS, Cunha DA, Brito WA, Soares Célia MA.** 2001. Two-dimensional electrophoresis and characterization of antigens from *Paracoccidioides brasiliensis*. *Microb Infect* 3(7):535–542 DOI 10.1016/S1286-4579(01)01409-5.

- da Silva Jde F, de Oliveira HC, Marcos CM, da Silva RA, da Costa TA, Calich VL, Almeida AM, Mendes-Giannini MJ.** 2013. *Paracoccidioides brasiliensis* 30 kDa adhesin: identification as a 14-3-3 protein, cloning and subcellular localization in infection models. *PLOS ONE* **8**(4):e62533 DOI [10.1371/journal.pone.0062533](https://doi.org/10.1371/journal.pone.0062533).
- da Silva Neto BR, de Fátima da Silva J, Mendes-Giannini MJ, Lenzi HL, de Almeida Soares CM, Pereira M.** 2009. The malate synthase of *Paracoccidioides brasiliensis* is a linked surface protein that behaves as an anchorless adhesin. *BMC Microbiology* **9**(1):272 DOI [10.1186/1471-2180-9-272](https://doi.org/10.1186/1471-2180-9-272).
- Dasari P, Koleci N, Shopova IA, Wartenberg D, Beyersdorf N, Dietrich S, Sahagún-Ruiz A, Figge MT, Skerka C, Brakhage AA, Zipfel PF.** 2019. Enolase from *Aspergillus fumigatus* is a moonlighting protein that binds the human plasma complement proteins factor H, FHL-1, C4BP, and plasminogen. *Frontiers in Immunology* **10**:2573 DOI [10.3389/fimmu.2019.02573](https://doi.org/10.3389/fimmu.2019.02573).
- de Oliveira HC, Assato PA, Assato Patrícia A, Marcos CM, Scorzoni L, de Paula E Silva ACA, De Fátima Da Silva J, de Lacorte Singulani J, Alarcon KM, Fusco-Almeida AM, Mendes-Giannini MJS.** 2015. Paracoccidioides-host interaction: an overview on recent advances in the paracoccidioidomycosis. *Frontiers in Microbiology* **6**(29):1319 DOI [10.3389/fmicb.2015.01319](https://doi.org/10.3389/fmicb.2015.01319).
- de Oliveira KM, da Silva Neto BR, Parente JA, da Silva RA, Quintino GO, Voltan AR, Mendes-Giannini MJ, de Almeida Soares CM, Pereira M.** 2013. Intermolecular interactions of the malate synthase of *Paracoccidioides* spp. *BMC Microbiology* **13**(1):107 DOI [10.1186/1471-2180-13-107](https://doi.org/10.1186/1471-2180-13-107).
- Derbise A, Song YP, Parikh S, Fischetti VA, Pancholi V.** 2004. Role of the C-terminal lysine residues of streptococcal surface enolase in Glu- and Lys-plasminogen-binding activities of group A *streptococci*. *Infection and Immunity* **72**(1):94–105 DOI [10.1128/iai.72.1.94-105.2004](https://doi.org/10.1128/iai.72.1.94-105.2004).
- Díaz-Jiménez DF, Pérez-García LA, Martínez-Álvarez JA, Mora-Montes HM.** 2012. Role of the fungal cell wall in pathogenesis and antifungal resistance. *Current Fungal Infection Reports* **6**(4):275–282 DOI [10.1007/s12281-012-0109-7](https://doi.org/10.1007/s12281-012-0109-7).
- Donofrio FC, Calil ACA, Miranda ET, Almeida AMF, Benard G, Soares CP, Veloso SN, Soares CMA, Mendes Giannini MJS.** 2009. Enolase from *Paracoccidioides brasiliensis*: isolation and identification as a fibronectin-binding protein. *Journal of Medical Microbiology* **58**(Pt 6):706–713 DOI [10.1099/jmm.0.003830-0](https://doi.org/10.1099/jmm.0.003830-0).
- Flores CL, Gancedo C.** 2011. Unraveling moonlighting functions with yeasts. *IUBMB Life* **63**(7):457–462 DOI [10.1002/iub.454](https://doi.org/10.1002/iub.454).
- Friese MA, Hellwage J, Jokiranta TS, Meri S, Peter HH, Eibel H, Zipfel PF.** 1999. FHL-1/reconectin and factor H: two human complement regulators which are encoded by the same gene are differently expressed and regulated. *Molecular Immunology* **36**(13–14):809–818 DOI [10.1016/s0161-5890\(99\)00101-7](https://doi.org/10.1016/s0161-5890(99)00101-7).
- Gancedo C, Flores CL.** 2008. Moonlighting proteins in yeasts. *Microbiology and Molecular Biology Reviews* **72**(1):197–210 DOI [10.1128/mmbr.00036-07](https://doi.org/10.1128/mmbr.00036-07).
- Gancedo C, Flores C-L, Gancedo JM.** 2016. The expanding landscape of moonlighting proteins in yeasts. *Microbiology and Molecular Biology Reviews* **80**(3):765–777 DOI [10.1128/MMBR.00012-16](https://doi.org/10.1128/MMBR.00012-16).
- García-Carnero LC, Martínez-Álvarez JA.** 2022. Virulence factors of *Sporothrix schenckii*. *Journal of Fungi* **8**(3):318 DOI [10.3390/jof8030318](https://doi.org/10.3390/jof8030318).
- García-Carnero LC, Salinas-Marín R, Lozoya-Pérez NE, Wrobel K, Wrobel K, Martínez-Duncker I, Niño-Vega GA, Mora-Montes HM.** 2021. The heat shock protein 60 and Pap1 participate in the *Sporothrix schenckii*-host interaction. *Journal of Fungi* **7**(11):960 DOI [10.3390/jof7110960](https://doi.org/10.3390/jof7110960).

- García-Rodas R, Zaragoza O.** 2012. Catch me if you can: phagocytosis and killing avoidance by *Cryptococcus neoformans*. *FEMS Immunology & Medical Microbiology* **64**(2):147–161 DOI [10.1111/j.1574-695X.2011.00871.x](https://doi.org/10.1111/j.1574-695X.2011.00871.x).
- Gautam P, Sundaram CS, Madan T, Gade WN, Shah A, Sirdeshmukh R, Sarma PU.** 2007. Identification of novel allergens of *Aspergillus fumigatus* using immunoproteomics approach. *Clinical and Experimental Allergy* **37**(8):1239–1249 DOI [10.1111/j.1365-2222.2007.02765.x](https://doi.org/10.1111/j.1365-2222.2007.02765.x).
- Gil-Bona A, Amador-García A, Gil C, Monteoliva L.** 2018. The external face of *Candida albicans*: a proteomic view of the cell surface and the extracellular environment. *Journal of Proteomics* **180**(1):70–79 DOI [10.1016/j.jprot.2017.12.002](https://doi.org/10.1016/j.jprot.2017.12.002).
- Gil-Navarro I, Gil ML, Casanova M, O'Connor JE, Martínez JP, Gozalbo D.** 1997. The glycolytic enzyme glyceraldehyde-3-phosphate dehydrogenase of *Candida albicans* is a surface antigen. *Journal of Bacteriology* **179**(16):4992–4999 DOI [10.1128/jb.179.16.4992-4999.1997](https://doi.org/10.1128/jb.179.16.4992-4999.1997).
- Gil ML, Dagan S, Eren R, Gozalbo D.** 2006. Evaluation of the usefulness of anti-glyceraldehyde-3-phosphate dehydrogenase antibodies as a treatment for invasive candidiasis in a murine model. *Antonie van Leeuwenhoek* **89**(3):345–350 DOI [10.1007/s10482-005-9037-7](https://doi.org/10.1007/s10482-005-9037-7).
- Gómez-Gaviria M, Mora-Montes HM.** 2020. Current aspects in the biology, pathogeny, and treatment of *Candida krusei*, a neglected fungal pathogen. *Infect Drug Resist* **13**:1673–1689 DOI [10.2147/idr.S247944](https://doi.org/10.2147/idr.S247944).
- Gomez FJ, Allendoerfer R, Deepe GS Jr.** 1995. Vaccination with recombinant heat shock protein 60 from *Histoplasma capsulatum* protects mice against pulmonary histoplasmosis. *Infection and Immunity* **63**(7):2587–2595 DOI [10.1128/iai.63.7.2587-2595.1995](https://doi.org/10.1128/iai.63.7.2587-2595.1995).
- Gomez FJ, Gomez AM, Deepe GS Jr.** 1991. Protective efficacy of a 62-kilodalton antigen, HIS-62, from the cell wall and cell membrane of *Histoplasma capsulatum* yeast cells. *Infection and Immunity* **59**(12):4459–4464 DOI [10.1128/iai.59.12.4459-4464.1991](https://doi.org/10.1128/iai.59.12.4459-4464.1991).
- Gozalbo D, Gil-Navarro I, Azorín I, Renau-Piquer J, Martínez JP, Gil ML.** 1998. The cell wall-associated glyceraldehyde-3-phosphate dehydrogenase of *Candida albicans* is also a fibronectin and laminin binding protein. *Infection and Immunity* **66**(5):2052–2059 DOI [10.1128/iai.66.5.2052-2059.1998](https://doi.org/10.1128/iai.66.5.2052-2059.1998).
- Guimarães AJ, Frases S, Gomez FJ, Zancopé-Oliveira RM, Nosanchuk JD.** 2009. Monoclonal antibodies to heat shock protein 60 alter the pathogenesis of *Histoplasma capsulatum*. *Infection and Immunity* **77**(4):1357–1367 DOI [10.1128/iai.01443-08](https://doi.org/10.1128/iai.01443-08).
- Henderson B, Martin A.** 2011. Bacterial virulence in the moonlight: multitasking bacterial moonlighting proteins are virulence determinants in infectious disease. *Infection and Immunity* **79**(9):3476–3491 DOI [10.1128/iai.00179-11](https://doi.org/10.1128/iai.00179-11).
- Hernández ML, Ximénez-Eembún P, Martínez-Gomariz M, Gutiérrez-Blázquez MD, Nombela C, Gil C.** 2010. Identification of *Candida albicans* exposed surface proteins *in vivo* by a rapid proteomic approach. *Journal of Proteomics* **73**(7):1404–1409 DOI [10.1016/j.jprot.2010.02.008](https://doi.org/10.1016/j.jprot.2010.02.008).
- Hernández-Chávez MJ, González-Hernández RJ, Trujillo-Esquivel JE, Hernández-Cervantes A, Mora-Montes HM.** 2014. Chapter 9-The secretory pathway in the filamentous fungus Trichoderma. In: Gupta VK, Schmoll M, Alfredo Herrera-Estrella RSU, Druzhinina I, Tuohy MG, eds. *Biotechnology and Biology of Trichoderma*. Amsterdam: Elsevier, 115–121.
- Hope WW, Walsh TJ, Denning DW.** 2005. Laboratory diagnosis of invasive aspergillosis. *Lancet Infectious Diseases* **5**(10):609–622 DOI [10.1016/s1473-3099\(05\)70238-3](https://doi.org/10.1016/s1473-3099(05)70238-3).
- Hoyer LL, Payne TL, Bell M, Myers AM, Scherer S.** 1998. *Candida albicans* ALS3 and insights into the nature of the ALS gene family. *Current Genetics* **33**(6):451–459 DOI [10.1007/s002940050359](https://doi.org/10.1007/s002940050359).

- Huberts DH, van der Klei IJ.** 2010. Moonlighting proteins: an intriguing mode of multitasking. *Biochim Biophys Acta* **1803**(4):520–525 DOI [10.1016/j.bbamcr.2010.01.022](https://doi.org/10.1016/j.bbamcr.2010.01.022).
- Jeffery CJ.** 1999. Moonlighting proteins. *Trends in Biochemical Sciences* **24**(1):8–11 DOI [10.1016/s0968-0004\(98\)01335-8](https://doi.org/10.1016/s0968-0004(98)01335-8).
- Jeffery CJ.** 2004. Molecular mechanisms for multitasking: recent crystal structures of moonlighting proteins. *Current Opinion in Structural Biology* **14**(6):663–668 DOI [10.1016/j.sbi.2004.10.001](https://doi.org/10.1016/j.sbi.2004.10.001).
- Jeffery CJ.** 2015. Why study moonlighting proteins? *Frontiers in Genetics* **6**:211 DOI [10.3389/fgene.2015.00211](https://doi.org/10.3389/fgene.2015.00211).
- Jeffery CJ.** 2018. Protein moonlighting: what is it, and why is it important? *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **373**(1738):20160523 DOI [10.1098/rstb.2016.0523](https://doi.org/10.1098/rstb.2016.0523).
- Jong AY, Chen SHM, Stins MF, Kim KS, Tuan TL, Huang SH.** 2003. Binding of *Candida albicans* enolase to plasmin(ogen) results in enhanced invasion of human brain microvascular endothelial cells. *Journal of Medical Microbiology* **52**(Pt 8):615–622 DOI [10.1099/jmm.0.05060-0](https://doi.org/10.1099/jmm.0.05060-0).
- Karamanos NK, Theocharis AD, Piperigkou Z, Manou D, Passi A, Skandalis SS, Vynios DH, Orian-Rousseau V, Ricard-Blum S, Schmelzer CEH, Duca L, Durbeej M, Afratis NA, Troeberg L, Franchi M, Masola V, Onisto M.** 2021. A guide to the composition and functions of the extracellular matrix. *The FEBS Journal* **288**(24):6850–6912 DOI [10.1111/febs.15776](https://doi.org/10.1111/febs.15776).
- Karkowska-Kuleta J, Kedracka-Krok S, Rapala-Kozik M, Kamysz W, Bielinska S, Karafova A, Kozik A.** 2011. Molecular determinants of the interaction between human high molecular weight kininogen and *Candida albicans* cell wall: identification of kininogen-binding proteins on fungal cell wall and mapping the cell wall-binding regions on kininogen molecule. *Peptides* **32**(12):2488–2496 DOI [10.1016/j.peptides.2011.10.021](https://doi.org/10.1016/j.peptides.2011.10.021).
- Karkowska-Kuleta J, Kozik A.** 2014. Moonlighting proteins as virulence factors of pathogenic fungi, parasitic protozoa and multicellular parasites. *Molecular Oral Microbiology* **29**(6):270–283 DOI [10.1111/omi.12078](https://doi.org/10.1111/omi.12078).
- Karkowska-Kuleta J, Zajac D, Bras G, Bochenska O, Rapala-Kozik M, Kozik A.** 2017. Binding of human plasminogen and high-molecular-mass kininogen by cell surface-exposed proteins of *Candida parapsilosis*. *Acta Biochimica Polonica* **64**(3):391–400 DOI [10.18388/abp.2017_1609](https://doi.org/10.18388/abp.2017_1609).
- Karkowska-Kuleta J, Zajac D, Bras G, Bochenska O, Seweryn K, Kedracka-Krok S, Jankowska U, Rapala-Kozik M, Kozik A.** 2016. Characterization of the interactions between human high-molecular-mass kininogen and cell wall proteins of pathogenic yeasts *Candida tropicalis*. *Acta Biochimica Polonica* **63**(3):427–436 DOI [10.18388/abp.2016_1353](https://doi.org/10.18388/abp.2016_1353).
- Kauffman CA.** 2007. Histoplasmosis: a clinical and laboratory update. *Clinical Microbiology Reviews* **20**(1):115–132 DOI [10.1128/cmr.00027-06](https://doi.org/10.1128/cmr.00027-06).
- Knox KS, Hage CA.** 2010. Histoplasmosis. *Proceedings of the American Thoracic Society* **7**(3):169–172 DOI [10.1513/pats.200907-069AL](https://doi.org/10.1513/pats.200907-069AL).
- Ko H-C, Hsiao T-Y, Chen C-T, Yang Y-L.** 2013. *Candida albicans* ENO1 null mutants exhibit altered drug susceptibility, hyphal formation, and virulence. *Journal of Microbiology* **51**(3):345–351 DOI [10.1007/s12275-013-2577-z](https://doi.org/10.1007/s12275-013-2577-z).
- Kolwijk E, van de Veerdonk FL.** 2014. The potential impact of the pulmonary microbiome on immunopathogenesis of *Aspergillus*-related lung disease. *European Journal of Immunology* **44**(11):3156–3165 DOI [10.1002/eji.201344404](https://doi.org/10.1002/eji.201344404).
- Kozik A, Karkowska-Kuleta J, Zajac D, Bochenska O, Kedracka-Krok S, Jankowska U, Rapala-Kozik M.** 2015. Fibronectin-, vitronectin- and laminin-binding proteins at the cell walls

of *Candida parapsilosis* and *Candida tropicalis* pathogenic yeasts. *BMC Microbiology* **15**(1):197 DOI [10.1186/s12866-015-0531-4](https://doi.org/10.1186/s12866-015-0531-4).

Kwak MK, Ku M, Kang SO. 2014. NAD(+)-linked alcohol dehydrogenase 1 regulates methylglyoxal concentration in *Candida albicans*. *FEBS Letters* **588**(7):1144–1153 DOI [10.1016/j.febslet.2014.02.042](https://doi.org/10.1016/j.febslet.2014.02.042).

Lai HY, Tam MF, Tang RB, Chou H, Chang CY, Tsai JJ, Shen HD. 2002. cDNA cloning and immunological characterization of a newly identified endolase allergen from *Penicillium citrinum* and *Aspergillus fumigatus*. *International Archives of Allergy and Immunology* **127**(3):181–190 DOI [10.1159/000053862](https://doi.org/10.1159/000053862).

Lalmanach G, Naudin C, Lecaille F, Fritz H. 2010. Kininogens: more than cysteine protease inhibitors and kinin precursors. *Biochimie* **92**(11):1568–1579 DOI [10.1016/j.biochi.2010.03.011](https://doi.org/10.1016/j.biochi.2010.03.011).

Latgé JP, Chamilos G. 2019. *Aspergillus fumigatus* and aspergillosis in 2019. *Clinical Microbiology Reviews* **33**(1):e00140-18 DOI [10.1128/cmr.00140-18](https://doi.org/10.1128/cmr.00140-18).

Li L, Zhang T, Xu J, Wu J, Wang Y, Qiu X, Zhang Y, Hou W, Yan L, An M, Jiang Y. 2019. The synergism of the small molecule ENOblock and fluconazole against fluconazole-resistant *Candida albicans*. *Frontiers in Microbiology* **10**:2071 DOI [10.3389/fmicb.2019.02071.2071](https://doi.org/10.3389/fmicb.2019.02071.2071).

Li XS, Reddy MS, Baev D, Edgerton M. 2003. *Candida albicans* Ssa1/2p is the cell envelope binding protein for human salivary histatin 5. *Journal of Biological Chemistry* **278**(31):28553–28561 DOI [10.1074/jbc.M300680200](https://doi.org/10.1074/jbc.M300680200).

Li XS, Sun JN, Okamoto-Shibayama K, Edgerton M. 2006. *Candida albicans* cell wall ssa proteins bind and facilitate import of salivary histatin 5 required for toxicity. *Journal of Biological Chemistry* **281**(32):22453–22463 DOI [10.1074/jbc.M604064200](https://doi.org/10.1074/jbc.M604064200).

Lin CS, He PJ, Tsai NM, Li CH, Yang SC, Hsu WT, Wu MS, Wu CJ, Cheng TL, Liao KW. 2010. A potential role for *Helicobacter pylori* heat shock protein 60 in gastric tumorigenesis. *Biochemical and Biophysical Research Communications* **392**(2):183–189 DOI [10.1016/j.bbrc.2010.01.010](https://doi.org/10.1016/j.bbrc.2010.01.010).

Liu H, Jeffery CJ. 2020. Moonlighting proteins in the fuzzy logic of cellular metabolism. *Molecules* **25**(15):3440 DOI [10.3390/molecules25153440](https://doi.org/10.3390/molecules25153440).

Long KH, Gomez FJ, Morris RE, Newman SL. 2003. Identification of heat shock protein 60 as the ligand on *Histoplasma capsulatum* that mediates binding to CD18 receptors on human macrophages. *Journal of Immunology* **170**(1):487–494 DOI [10.4049/jimmunol.170.1.487](https://doi.org/10.4049/jimmunol.170.1.487).

Longo LVG, da Cunha JPC, Sobreira TJP, Puccia R. 2014. Proteome of cell wall-extracts from pathogenic *Paracoccidioides brasiliensis*: comparison among morphological phases, isolates, and reported fungal extracellular vesicle proteins. *EuPA Open Proteomics* **3**:216–228 DOI [10.1016/j.euprot.2014.03.003](https://doi.org/10.1016/j.euprot.2014.03.003).

Lopes-Bezerra LM, Mora-Montes HM, Zhang Y, Nino-Vega G, Rodrigues AM, de Camargo ZP, de Hoog S. 2018. Sporotrichosis between 1898 and 2017: the evolution of knowledge on a changeable disease and on emerging etiological agents. *Medical Mycology* **56**(suppl_1):S126–S143 DOI [10.1093/mmy/myx103](https://doi.org/10.1093/mmy/myx103).

López-Ribot JL, Alloush HM, Masten BJ, Chaffin WL. 1996. Evidence for presence in the cell wall of *Candida albicans* of a protein related to the hsp70 family. *Infection and Immunity* **64**(8):3333–3340 DOI [10.1128/iai.64.8.3333-3340.1996](https://doi.org/10.1128/iai.64.8.3333-3340.1996).

López-Ribot JL, Casanova M, Murgui A, Martínez JP. 2004. Antibody response to *Candida albicans* cell wall antigens. *FEMS Immunology & Medical Microbiology* **41**(3):187–196 DOI [10.1016/j.femsim.2004.03.012](https://doi.org/10.1016/j.femsim.2004.03.012).

Lopez-Romero E, Reyes-Montes Mdel R, Perez-Torres A, Ruiz-Baca E, Villagomez-Castro JC, Mora-Montes HM, Flores-Carreon A, Toriello C. 2011. *Sporothrix schenckii* complex and

sporotrichosis, an emerging health problem. *Future Microbiology* **6**(1):85–102
DOI [10.2217/fmb.10.157](https://doi.org/10.2217/fmb.10.157).

- Lopez CM, Wallich R, Riesbeck K, Skerka C, Zipfel PF.** 2014. *Candida albicans* uses the surface protein Gpm1 to attach to human endothelial cells and to keratinocytes via the adhesive protein vitronectin. *PLOS ONE* **9**(3):e90796 DOI [10.1371/journal.pone.0090796](https://doi.org/10.1371/journal.pone.0090796).
- Luo S, Hoffmann R, Skerka C, Zipfel PF.** 2013. Glycerol-3-phosphate dehydrogenase 2 is a novel factor H-, factor H-like protein 1-, and plasminogen-binding surface protein of *Candida albicans*. *Journal of Infectious Diseases* **207**(4):594–603 DOI [10.1093/infdis/jis718](https://doi.org/10.1093/infdis/jis718).
- Marcos CM, de Fátima da Silva J, de Oliveira HC, Moraes da Silva RA, Mendes-Giannini MJ, Fusco-Almeida AM.** 2012. Surface-expressed enolase contributes to the adhesion of *Paracoccidioides brasiliensis* to host cells. *FEMS Yeast Research* **12**(5):557–570 DOI [10.1111/j.1567-1364.2012.00806.x](https://doi.org/10.1111/j.1567-1364.2012.00806.x).
- Marcos CM, Silva Jde F, Oliveira HC, Assato PA, Singulani Jde L, Lopez AM, Tamayo DP, Hernandez-Ruiz O, McEwen JG, Mendes-Giannini MJ, Fusco-Almeida AM.** 2016. Decreased expression of 14-3-3 in *Paracoccidioides brasiliensis* confirms its involvement in fungal pathogenesis. *Virulence* **7**(2):72–84 DOI [10.1080/21505594.2015.1122166](https://doi.org/10.1080/21505594.2015.1122166).
- Marcos CM, de Oliveira HC, da Silva JdF, Assato PA, Fusco-Almeida AM, Mendes-Giannini MJS.** 2014. The multifaceted roles of metabolic enzymes in the *Paracoccidioides* species complex. *Frontiers in Microbiology* **5**(29):719 DOI [10.3389/fmicb.2014.00719](https://doi.org/10.3389/fmicb.2014.00719).
- Martínez-Duncker I, Díaz-Jiménez DF, Mora-Montes HM.** 2014. Comparative analysis of protein glycosylation pathways in humans and the fungal pathogen *Candida albicans*. *International Journal of Microbiology* **2014**:267497 DOI [10.1155/2014/267497.2014](https://doi.org/10.1155/2014/267497).
- Martínez JP, Gil ML, López-Ribot JL, Chaffin WL.** 1998. Serologic response to cell wall mannoproteins and proteins of *Candida albicans*. *Clinical Microbiology Reviews* **11**(1):121–141 DOI [10.1128/cmrr.11.1.121](https://doi.org/10.1128/cmrr.11.1.121).
- Mendes RP, Cavalcante RS, Marques SA, Marques MEA, Venturini J, Sylvestre TF, Paniago AMM, Pereira AC, da Silva JF, Fabro AT, Bosco SMG, Bagagli E, Hahn RC, Levorato AD.** 2017. Paracoccidioidomycosis: current perspectives from Brazil. *The Open Microbiology Journal* **11**(1):224–282 DOI [10.2174/1874285801711010224](https://doi.org/10.2174/1874285801711010224).
- Mihu MR, Nosanchuk JD.** 2012. Histoplasma virulence and host responses. *International Journal of Microbiology* **2012**:268123 DOI [10.1155/2012/268123.2012](https://doi.org/10.1155/2012/268123).
- Miles LA, Dahlberg CM, Plescia J, Felez J, Kato K, Plow EF.** 1991. Role of cell-surface lysines in plasminogen binding to cells: identification of alpha-enolase as a candidate plasminogen receptor. *Biochemistry* **30**(6):1682–1691 DOI [10.1021/bi00220a034](https://doi.org/10.1021/bi00220a034).
- Montagnoli C, Sandini S, Bacci A, Romani L, Valle RL.** 2004. Immunogenicity and protective effect of recombinant enolase of *Candida albicans* in a murine model of systemic candidiasis. *Medical Mycology* **42**(4):319–324 DOI [10.1080/13693780310001644653](https://doi.org/10.1080/13693780310001644653).
- Mora-Montes HM, Dantas Ada S, Trujillo-Esquivel E, de Souza Baptista AR, Lopes-Bezerra LM.** 2015. Current progress in the biology of members of the *Sporothrix schenckii* complex following the genomic era. *FEMS Yeast Research* **15**(6):fov065 DOI [10.1093/femsyr/fov065](https://doi.org/10.1093/femsyr/fov065).
- Mora-Montes HM, Ponce-Noyola P, Villagomez-Castro JC, Gow NA, Flores-Carreon A, Lopez-Romero E.** 2009. Protein glycosylation in *Candida*. *Future Microbiology* **4**(9):1167–1183 DOI [10.2217/fmb.09.88](https://doi.org/10.2217/fmb.09.88).

- Nava-Pérez N, Neri-García LG, Romero-González OE, Terrones-Cruz JA, García-Carnero LC, Mora-Montes HM.** 2022. Biological and clinical attributes of *Sporothrix globosa*, a causative agent of sporotrichosis. *Infect Drug Resist* 15:2067–2090 DOI [10.2147/idr.s362099](https://doi.org/10.2147/idr.s362099).
- Navarro-Arias MJ, Hernández-Chávez MJ, García-Carnero LC, Amezcu-Hernández DG, Lozoya-Pérez NE, Estrada-Mata E, Martínez-Duncker I, Franco B, Mora-Montes HM.** 2019. Differential recognition of *Candida tropicalis*, *Candida guilliermondii*, *Candida krusei*, and *Candida auris* by human innate immune cells. *Infect Drug Resist* 12:783–794 DOI [10.2147/idr.S197531](https://doi.org/10.2147/idr.S197531).
- Newman SL, Bucher C, Rhodes J, Bullock WE.** 1990. Phagocytosis of *Histoplasma capsulatum* yeasts and microconidia by human cultured macrophages and alveolar macrophages. Cellular cytoskeleton requirement for attachment and ingestion. *Journal of Clinical Investigation* 85(1):223–230 DOI [10.1172/jci114416](https://doi.org/10.1172/jci114416).
- Nickel W, Rabouille C.** 2009. Mechanisms of regulated unconventional protein secretion. *Nature Reviews Molecular Cell Biology* 10(2):148–155 DOI [10.1038/nrm2617](https://doi.org/10.1038/nrm2617).
- Nobbs AH, Vickerman MM, Jenkinson HF.** 2010. Heterologous expression of *Candida albicans* cell wall-associated adhesins in *Saccharomyces cerevisiae* reveals differential specificities in adherence and biofilm formation and in binding oral *Streptococcus gordonii*. *Eukaryotic Cell* 9(10):1622–1634 DOI [10.1128/ec.00103-10](https://doi.org/10.1128/ec.00103-10).
- Nobile CJ, Andes DR, Nett JE, Smith FJ, Yue F, Phan QT, Edwards JE, Filler SG, Mitchell AP.** 2006. Critical role of Bcr1-dependent adhesins in *C. albicans* biofilm formation *in vitro* and *in vivo*. *PLOS Pathogens* 2(7):e63 DOI [10.1371/journal.ppat.0020063](https://doi.org/10.1371/journal.ppat.0020063).
- Nogueira SV, Fonseca FL, Rodrigues ML, Mundodi V, Abi-Chacra EA, Winters MS, Alderete JF, de Almeida Soares CM.** 2010. *Paracoccidioides brasiliensis* enolase is a surface protein that binds plasminogen and mediates interaction of yeast forms with host cells. *Infection and Immunity* 78(9):4040–4050 DOI [10.1128/iai.00221-10](https://doi.org/10.1128/iai.00221-10).
- Núñez-Beltrán A, López-Romero E, Cuéllar-Cruz M.** 2017. Identification of proteins involved in the adhesion of *Candida* species to different medical devices. *Microb Pathog* 107(9):293–303 DOI [10.1016/j.micpath.2017.04.009](https://doi.org/10.1016/j.micpath.2017.04.009).
- Oehmcke-Hecht S, Köhler J.** 2018. Interaction of the human contact system with pathogens—an update. *Frontiers in Immunology* 9:3389 DOI [10.3389/fimmu.2018.00312](https://doi.org/10.3389/fimmu.2018.00312).
- Pappas PG, Alexander BD, Andes DR, Hadley S, Kauffman CA, Freifeld A, Anaissie EJ, Brumble LM, Herwaldt L, Ito J, Kontoyiannis DP, Lyon GM, Marr KA, Morrison VA, Park BJ, Patterson TF, Perl TM, Oster RA, Schuster MG, Walker R, Walsh TJ, Wannemuehler KA, Chiller TM.** 2010. Invasive fungal infections among organ transplant recipients: results of the transplant-associated infection surveillance network (TRANSNET). *Clinical Infectious Diseases* 50(8):1101–1111 DOI [10.1086/651262](https://doi.org/10.1086/651262).
- Park SJ, Mehrad B.** 2009. Innate immunity to *Aspergillus* species. *Clinical Microbiology Reviews* 22(4):535–551 DOI [10.1128/cmr.00014-09](https://doi.org/10.1128/cmr.00014-09).
- Pereira LA, Bão SN, Barbosa MS, da Silva JL, Felipe MS, de Santana JM, Mendes-Giannini MJ, de Almeida Soares CM.** 2007. Analysis of the *Paracoccidioides brasiliensis* triosephosphate isomerase suggests the potential for adhesin function. *Fems Yeast Research* 7(8):1381–1388 DOI [10.1111/j.1567-1364.2007.00292.x](https://doi.org/10.1111/j.1567-1364.2007.00292.x).
- Phan QT, Myers CL, Fu Y, Sheppard DC, Yeaman MR, Welch WH, Ibrahim AS, Edwards JE, Filler SG.** 2007. Als3 is a *Candida albicans* invasin that binds to cadherins and induces endocytosis by host cells. *PLOS Biology* 5(3):e64 DOI [10.1371/journal.pbio.0050064](https://doi.org/10.1371/journal.pbio.0050064).

- Pitarch A, Sánchez M, Nombela C, Gil C.** 2002. Sequential fractionation and two-dimensional gel analysis unravels the complexity of the dimorphic fungus *Candida albicans* cell wall proteome. *Molecular & Cellular Proteomics* 1(12):967–982 DOI [10.1074/mcp.m200062-mcp200](https://doi.org/10.1074/mcp.m200062-mcp200).
- Podobnik M, Tyagi R, Matange N, Dermol U, Gupta AK, Mattoo R, Seshadri K, Visweswariah SS.** 2009. A mycobacterial cyclic AMP phosphodiesterase that moonlights as a modifier of cell wall permeability. *Journal of Biological Chemistry* 284(47):32846–32857 DOI [10.1074/jbc.M109.049635](https://doi.org/10.1074/jbc.M109.049635).
- Poltermann S, Kunert A, von der Heide M, Eck R, Hartmann A, Zipfel PF.** 2007. Gpm1p is a factor H-, FHL-1-, and plasminogen-binding surface protein of *Candida albicans*. *Journal of Biological Chemistry* 282(52):37537–37544 DOI [10.1074/jbc.M707280200](https://doi.org/10.1074/jbc.M707280200).
- Ponczek MB.** 2021. High molecular weight kininogen: a review of the structural literature. *International Journal of Molecular Sciences* 22(24):13370 DOI [10.3390/ijms222413370](https://doi.org/10.3390/ijms222413370).
- Rahi A, Dhiman A, Singh D, Lynn AM, Rehan M, Bhatnagar R.** 2018. Exploring the interaction between *Mycobacterium tuberculosis* enolase and human plasminogen using computational methods and experimental techniques. *Journal of Cellular Biochemistry* 119(2):2408–2417 DOI [10.1002/jcb.26403](https://doi.org/10.1002/jcb.26403).
- Raksha, Singh G, Urhekar AD.** 2017. Virulence factors detection in Aspergillus isolates from clinical and environmental samples. *Journal of Clinical and Diagnostic Research* 11(7):Dc13–Dc18 DOI [10.7860/jcdr/2017/24055.10211](https://doi.org/10.7860/jcdr/2017/24055.10211).
- Reyna-Beltrán E, Iranzo M, Calderón-González KG, Mondragón-Flores R, Labra-Barrios ML, Mormeneo S, Luna-Arias JP.** 2018. The *Candida albicans* ENO1 gene encodes a transglutaminase involved in growth, cell division, morphogenesis, and osmotic protection. *Journal of Biological Chemistry* 293(12):4304–4323 DOI [10.1074/jbc.M117.810440](https://doi.org/10.1074/jbc.M117.810440).
- Rodrigues ML, Djordjevic JT.** 2012. Unravelling secretion in *Cryptococcus neoformans*: more than one way to skin a cat. *Mycopathologia* 173(5–6):407–418 DOI [10.1007/s11046-011-9468-9](https://doi.org/10.1007/s11046-011-9468-9).
- Rodríguez-Saavedra C, Morgado-Martínez LE, Burgos-Palacios A, King-Díaz B, López-Coria M, Sánchez-Nieto S.** 2021. Moonlighting proteins: the case of the hexokinases. *Frontiers in Molecular Biosciences* 8:701975 DOI [10.3389/fmolb.2021.701975](https://doi.org/10.3389/fmolb.2021.701975).
- Ruiz-Baca E, Leyva-Sánchez H, Calderón-Barraza B, Esquivel-Naranjo U, López-Romero E, López-Rodríguez A, Cuéllar-Cruz M.** 2019. Identification of proteins in *Sporothrix schenckii sensu stricto* in response to oxidative stress induced by hydrogen peroxide. *Revista Iberoamericana de Micología* 36(1):17–23 DOI [10.1016/j.riam.2018.10.004](https://doi.org/10.1016/j.riam.2018.10.004).
- San-Blas G, Niño-Vega G, Iturriaga T.** 2002. *Paracoccidioides brasiliensis* and paracoccidioidomycosis: molecular approaches to morphogenesis, diagnosis, epidemiology, taxonomy and genetics. *Medical Mycology* 40(3):225–242 DOI [10.1080/mmy.40.3.225.242](https://doi.org/10.1080/mmy.40.3.225.242).
- Satala D, Karkowska-Kuleta J, Zelazna A, Rapala-Kozik M, Kozik A.** 2020a. Moonlighting proteins at the candidal cell surface. *Microorganisms* 8(7):1046 DOI [10.3390/microorganisms8071046](https://doi.org/10.3390/microorganisms8071046).
- Satala D, Satala G, Karkowska-Kuleta J, Bukowski M, Kluza A, Rapala-Kozik M, Kozik A.** 2020b. Structural insights into the interactions of candidal enolase with human vitronectin, fibronectin and plasminogen. *International Journal of Molecular Sciences* 21(21):7843 DOI [10.3390/ijms21217843](https://doi.org/10.3390/ijms21217843).
- Satala D, Satala G, Zawrotniak M, Kozik A.** 2021. *Candida albicans* and *Candida glabrata* triosephosphate isomerase-a moonlighting protein that can be exposed on the candidal cell surface and bind to human extracellular matrix proteins. *BMC Microbiology* 21(1):199 DOI [10.1186/s12866-021-02235-w](https://doi.org/10.1186/s12866-021-02235-w).

- Schmoll M, Dattenböck C, Carreras-Villaseñor N, Mendoza-Mendoza A, Tisch D, Alemán MI, Baker SE, Brown C, Cervantes-Badillo MG, Gomez-Rodriguez EY, Greenwood DR, Hernández-Oñate M, Kruszewska JS, Lawry R, Mora-Montes HM, Muñoz-Centeno T, Nieto-Jacobo MT, Sánchez-Arreguin JA, Seidl-Seiboth V, Stewart A, Uresti-Rivera EE, Wang CL, Wang TF, Zeilinger S, Casas-Flores S, Herrera-Estrella A. 2016. The genomes of three uneven siblings: footprints of the lifestyles of three *Trichoderma* species. *Microbiology and Molecular Biology Reviews* **80**(1):205–327 DOI [10.1128/mmbr.00040-15](https://doi.org/10.1128/mmbr.00040-15).
- Seidler NW. 2013. GAPDH, as a virulence factor. *Advances in Experimental Medicine and Biology* **985**(Pt 4):149–178 DOI [10.1007/978-94-007-4716-6_5](https://doi.org/10.1007/978-94-007-4716-6_5).
- Seweryn K, Karkowska-Kuleta J, Wolak N, Bochenska O, Kedracka-Krok S, Kozik A, Rapala-Kozik M. 2015. Kinetic and thermodynamic characterization of the interactions between the components of human plasma kinin-forming system and isolated and purified cell wall proteins of *Candida albicans*. *Acta Biochimica Polonica* **62**(4):825–835 DOI [10.18388/abp.2015_1142](https://doi.org/10.18388/abp.2015_1142).
- Shah A, Panjabi C. 2014. Allergic aspergillosis of the respiratory tract. *European Respiratory Review* **23**(131):8–29 DOI [10.1183/09059180.00007413](https://doi.org/10.1183/09059180.00007413).
- Sheppard DC, Yeaman MR, Welch WH, Phan QT, Fu Y, Ibrahim AS, Filler SG, Zhang M, Waring AJ, Edwards JE Jr. 2004. Functional and structural diversity in the Als protein family of *Candida albicans*. *Journal of Biological Chemistry* **279**(29):30480–30489 DOI [10.1074/jbc.M401929200](https://doi.org/10.1074/jbc.M401929200).
- Silveira CP, Piffer AC, Kmetzsch L, Fonseca FL, Soares DA, Staats CC, Rodrigues ML, Schrank A, Vainstein MH. 2013. The heat shock protein (Hsp) 70 of *Cryptococcus neoformans* is associated with the fungal cell surface and influences the interaction between yeast and host cells. *Fungal Genetics and Biology* **60**:53–63 DOI [10.1016/j.fgb.2013.08.005](https://doi.org/10.1016/j.fgb.2013.08.005).
- Singh B, Fleury C, Jalalvand F, Riesbeck K. 2012. Human pathogens utilize host extracellular matrix proteins laminin and collagen for adhesion and invasion of the host. *FEMS Microbiology Reviews* **36**(6):1122–1180 DOI [10.1111/j.1574-6976.2012.00340.x](https://doi.org/10.1111/j.1574-6976.2012.00340.x).
- Sluchanko NN, Gusev NB. 2017. Moonlighting chaperone-like activity of the universal regulatory 14-3-3 proteins. *FEBS Journal* **284**(9):1279–1295 DOI [10.1111/febs.13986](https://doi.org/10.1111/febs.13986).
- Spampinato C, Leonardi D. 2013. *Candida* infections, causes, targets, and resistance mechanisms: traditional and alternative antifungal agents. *Biomed Research International* **2013**(12):204237 DOI [10.1155/2013/204237](https://doi.org/10.1155/2013/204237).
- Spellberg BJ, Ibrahim AS, Avanesian V, Fu Y, Myers C, Phan QT, Filler SG, Yeaman MR, Edwards JE Jr. 2006. Efficacy of the anti-*Candida* rAls3p-N or rAls1p-N vaccines against disseminated and mucosal candidiasis. *Journal of Infectious Diseases* **194**(2):256–260 DOI [10.1086/504691](https://doi.org/10.1086/504691).
- Stie J, Bruni G, Fox D. 2009. Surface-associated plasminogen binding of *Cryptococcus neoformans* promotes extracellular matrix invasion. *PLOS ONE* **4**(6):e5780 DOI [10.1371/journal.pone.0005780](https://doi.org/10.1371/journal.pone.0005780).
- Sun JN, Li W, Jang WS, Nayyar N, Sutton MD, Edgerton M. 2008. Uptake of the antifungal cationic peptide Histatin 5 by *Candida albicans* Ssa2p requires binding to non-conventional sites within the ATPase domain. *Molecular Microbiology* **70**(5):1246–1260 DOI [10.1111/j.1365-2958.2008.06480.x](https://doi.org/10.1111/j.1365-2958.2008.06480.x).
- Sun JN, Solis NV, Phan QT, Bajwa JS, Kashleva H, Thompson A, Liu Y, Dongari-Bagtzoglou A, Edgerton M, Filler SG. 2010. Host cell invasion and virulence mediated by *Candida albicans* Ssa1. *PLOS Pathogens* **6**(11):e1001181 DOI [10.1371/journal.ppat.1001181](https://doi.org/10.1371/journal.ppat.1001181).

- Sundstrom P, Jensen J, Balish E.** 1994. Humoral and cellular immune responses to enolase after alimentary tract colonization or intravenous immunization with *Candida albicans*. *Journal of Infectious Diseases* **170**(2):390–395 DOI [10.1093/infdis/170.2.390](https://doi.org/10.1093/infdis/170.2.390).
- Szalewski DA, Hinrichs VS, Zinniel DK, Barletta RG.** 2018. The pathogenicity of *Aspergillus fumigatus*, drug resistance, and nanoparticle delivery. *Canadian Journal of Microbiology* **64**(7):439–453 DOI [10.1139/cjm-2017-0749](https://doi.org/10.1139/cjm-2017-0749).
- Tamez-Castrellon AK, Romeo O, Garcia-Carnero LC, Lozoya-Perez NE, Mora-Montes HM.** 2020. Virulence factors in *Sporothrix schenckii*, one of the causative agents of sporotrichosis. *Current Protein & Peptide Science* **21**(3):295–312 DOI [10.2174/138920372066191007103004](https://doi.org/10.2174/138920372066191007103004).
- Teixeira MM, de Almeida LG, Kubitschek-Barreira P, Alves FL, Kioshima ES, Abadio AK, Fernandes L, Derengowski LS, Ferreira KS, Souza RC, Ruiz JC, de Andrade NC, Paes HC, Nicola AM, Albuquerque P, Gerber AL, Martins VP, Peconick LD, Neto AV, Chaucanez CB, Silva PA, Cunha OL, de Oliveira FF, dos Santos TC, Barros AL, Soares MA, de Oliveira LM, Marini MM, Villalobos-Duno H, Cunha MM, de Hoog S, da Silveira JF, Henrissat B, Nino-Vega GA, Cisalpino PS, Mora-Montes HM, Almeida SR, Stajich JE, Lopes-Bezerra LM, Vasconcelos AT, Felipe MS.** 2014. Comparative genomics of the major fungal agents of human and animal sporotrichosis: *Sporothrix schenckii* and *Sporothrix brasiliensis*. *BMC Genomics* **15**(1):943 DOI [10.1186/1471-2164-15-943](https://doi.org/10.1186/1471-2164-15-943).
- Thammahong A, Caffrey-Card AK, Dhingra S, Obar JJ, Cramer RA.** 2017. *Aspergillus fumigatus* trehalose-regulatory subunit homolog moonlights to mediate cell wall homeostasis through modulation of chitin synthase activity. *mBio* **8**(2):e00056-17 DOI [10.1128/mBio.00056-17](https://doi.org/10.1128/mBio.00056-17).
- Tomlin H, Piccinini AM.** 2018. A complex interplay between the extracellular matrix and the innate immune response to microbial pathogens. *Immunology* **155**(2):186–201 DOI [10.1111/imm.12972](https://doi.org/10.1111/imm.12972).
- Tompa P, Szász C, Buday L.** 2005. Structural disorder throws new light on moonlighting. *Trends in Biochemical Sciences* **30**(9):484–489 DOI [10.1016/j.tibs.2005.07.008](https://doi.org/10.1016/j.tibs.2005.07.008).
- Tóth R, Nosek J, Mora-Montes HM, Gabaldon T, Bliss JM, Nosanchuk JD, Turner SA, Butler G, Vágvölgyi C, Gácser A.** 2019. *Candida parapsilosis*: from Genes to the Bedside. *Clinical Microbiology Reviews* **32**(2):e00111-18 DOI [10.1128/cmrr.00111-18](https://doi.org/10.1128/cmrr.00111-18).
- Turner SA, Butler G.** 2014. The *Candida* pathogenic species complex. *Cold Spring Harbor Perspectives in Medicine* **4**(9):a019778 DOI [10.1101/cshperspect.a019778](https://doi.org/10.1101/cshperspect.a019778).
- Vallejo MC, Nakayasu ES, Matsuo AL, Sobreira TJ, Longo LV, Ganiko L, Almeida IC, Puccia R.** 2012. Vesicle and vesicle-free extracellular proteome of *Paracoccidioides brasiliensis*: comparative analysis with other pathogenic fungi. *Journal of Proteome Research* **11**(3):1676–1685 DOI [10.1021/pr200872s](https://doi.org/10.1021/pr200872s).
- Vanegas G, Quiñones W, Carrasco-López C, Concepción JL, Albericio F, Avilán L.** 2007. Enolase as a plasminogen binding protein in *Leishmania mexicana*. *Parasitology Research* **101**(6):1511–1516 DOI [10.1007/s00436-007-0668-7](https://doi.org/10.1007/s00436-007-0668-7).
- Zhao X, Daniels KJ, Oh SH, Green CB, Yeater KM, Soll DR, Hoyer LL.** 2006. *Candida albicans* Als3p is required for wild-type biofilm formation on silicone elastomer surfaces. *Microbiology (Reading)* **152**(Pt 8):2287–2299 DOI [10.1099/mic.0.28959-0](https://doi.org/10.1099/mic.0.28959-0).
- Zhao X, Oh SH, Cheng G, Green CB, Nuessen JA, Yeater K, Leng RP, Brown AJP, Hoyer LL.** 2004. ALS3 and ALS8 represent a single locus that encodes a *Candida albicans* adhesin; functional comparisons between Als3p and Als1p. *Microbiology (Reading)* **150**(Pt 7):2415–2428 DOI [10.1099/mic.0.26943-0](https://doi.org/10.1099/mic.0.26943-0).