

# Genome-wide identification, characterization and expression analysis of *HAK* genes and decoding their role in responding to potassium deficiency and abiotic stress in *Medicago truncatula*

Yanxue Zhao<sup>1</sup>, Lei Wang<sup>1</sup>, Pengcheng Zhao<sup>2</sup>, Zhongjie Liu<sup>3</sup>, Siyi Guo<sup>1</sup>, Yang Li<sup>Corresp., 1</sup>, Hao Liu<sup>Corresp., 1</sup>

<sup>1</sup> State Key Laboratory of Crop Stress Adaptation and Improvement, School of Life Sciences, Henan University, Kaifeng, China

<sup>2</sup> College of Grassland Science, Nanjing Agricultural University, Nanjing, China

<sup>3</sup> Key Laboratory of Genetics and Fruit Development, College of Horticulture, Nanjing Agricultural University, Nanjing, China

Corresponding Authors: Yang Li, Hao Liu

Email address: liyang0378@henu.edu.cn, 10140026@vip.henu.edu.cn

**Background.** The HAK family is the largest potassium ( $K^+$ ) transporter family, vital in  $K^+$  uptake, plant growth, and both plant biotic and abiotic stress responses. Although *HAK* family members have been characterized and functionally investigated in many species, these genes are still not studied in detail in *Medicago truncatula*, a good model system for studying legume genetics.

**Methods.** In this study, we screened the *M. truncatula* *HAK* family members (*MtHAKs*). Furthermore, we also conducted the identification, phylogenetic analysis, and prediction of conserved motifs of *MtHAKs*. Moreover, we studied the expression levels of *MtHAKs* under  $K^+$  deficiency, drought, and salt stresses using quantitative real-time PCR (qRT-PCR).

**Results.** We identified 20 *MtHAK* family members and classified them into three clusters based on phylogenetic relationships. Conserved motif analyses showed that all *MtHAK* proteins besides *MtHAK10* contained the highly conserved  $K^+$  transport domain (GVVYGD LGTSPLY). qRT-PCR analysis showed that several *MtHAK* genes in roots were induced by abiotic stress. In particular, *MtHAK15*, *MtHAK17*, and *MtHAK18* were strongly up-regulated in the *M. truncatula* roots under  $K^+$  deficiency, drought, and salt stress conditions, thereby implying that these genes are good candidates for high-affinity  $K^+$  uptake and therefore have essential roles in drought and salt tolerance.

**Discussions.** Our results not only provide the first genetic description and evolutionary relationships of the  $K^+$  transporter family in *M. truncatula*, but also provide potential information responding to  $K^+$  deficiency and abiotic stresses, thereby laying the foundation for molecular breeding of stress-resistant legume crops in the future.

# Genome-wide identification, characterization and expression analysis of *HAK* genes and decoding their role in responding to potassium deficiency and abiotic stress in *Medicago truncatula*

Yanxue Zhao<sup>1</sup>, Lei Wang<sup>1</sup>, Pengcheng Zhao<sup>2</sup>, Zhongjie Liu<sup>3</sup>, Siyi Guo<sup>1</sup>, Yang Li<sup>1\*</sup>, Hao Liu<sup>1\*</sup>

<sup>1</sup> State Key Laboratory of Crop Stress Adaptation and Improvement, School of Life Sciences, Henan University, Kaifeng, China

<sup>2</sup> College of Grassland Science, Nanjing Agricultural University, Nanjing, China

<sup>3</sup> Key Laboratory of Genetics and Fruit Development, College of Horticulture, Nanjing Agricultural University, Nanjing, China

Corresponding Author:

Hao Liu

Jinming Road, Kaifeng, Henan, 475004, China

Email address: [10140026@vip.henu.edu.cn](mailto:10140026@vip.henu.edu.cn)

Yang Li

Jinming Road, Kaifeng, Henan, 475004, China

Email address: [liyang0378@henu.edu.cn](mailto:liyang0378@henu.edu.cn)

## Abstract

**Background.** The HAK family is the largest potassium (K<sup>+</sup>) transporter family, vital in K<sup>+</sup> uptake, plant growth, and both plant biotic and abiotic stress responses. Although *HAK* family members have been characterized and functionally investigated in many species, these genes are still not studied in detail in *Medicago truncatula*, a good model system for studying legume genetics.

**Methods.** In this study, we screened the *M. truncatula* *HAK* family members (*MtHAKs*). Furthermore, we also conducted the identification, phylogenetic analysis, and prediction of conserved motifs of *MtHAKs*. Moreover, we studied the expression levels of *MtHAKs* under K<sup>+</sup> deficiency, drought, and salt stresses using quantitative real-time PCR (qRT-PCR).

**Results.** We identified 20 *MtHAK* family members and classified them into three clusters based on phylogenetic relationships. Conserved motif analyses showed that all MtHAK proteins besides MtHAK10 contained the highly conserved K<sup>+</sup> transport domain (GVVYGD LGTSPLY). qRT-PCR analysis showed that several *MtHAK* genes in roots were induced by abiotic stress. In particular, *MtHAK15*, *MtHAK17*, and *MtHAK18* were strongly up-regulated in the *M. truncatula* roots under K<sup>+</sup> deficiency, drought, and salt stress conditions, thereby implying that these genes are good candidates for high-affinity K<sup>+</sup> uptake and therefore have essential roles in drought and salt tolerance.

**Discussions.** Our results not only provide the first genetic description and evolutionary relationships of the K<sup>+</sup> transporter family in *M. truncatula*, but also provide potential information responding to K<sup>+</sup> deficiency and abiotic stresses, thereby laying the foundation for molecular breeding of stress-resistant legume crops in the future.

## Introduction

K<sup>+</sup> is an essential macronutrient for various plant physiological functions, like ion homeostasis and the transport of nitrate and sugars (Li et al. 2018). Due to the limited K<sup>+</sup> resource, plants have evolved a series of K<sup>+</sup> transport systems to mediate its uptake and transport (Amrutha et al. 2007; Ashley et al. 2005; Gierth et al. 2005; Very et al. 2014). In plants, K<sup>+</sup> transporters are classified into four major families: 1) HAK (high-affinity K<sup>+</sup>)/KUP (K<sup>+</sup> uptake)/KT (K<sup>+</sup> transporter), 2) Trk/HKT, 3) CHX (cation/hydrogen exchanger), and 4) efflux antiporters KEA (K<sup>+</sup> efflux antiporter) (Gupta et al. 2008). Among them, the HAK/KUP/KT (HAK) family constitutes the largest K<sup>+</sup> transporter family that is ubiquitously present in plant genomes with varied numbers, e.g., 13 genes present in *Arabidopsis*, 27 in maize, and 27 in rice (Ahn et al. 2004; Corratge-Faillie et al. 2010; Gupta et al. 2008; Rubio et al. 2000; Zhang et al. 2012).

*HAK* genes have been found to play key roles in plant development and stress-related responses. For instance, *AtKUP4/TRH1* (*Tiny Root Hairs 1*) maintained the polar localization of AtPIN1 along with the auxin homeostasis and maxima in the root apex, thereby promoting root gravitropism response and root hair elongation (Rigas et al. 2001; Rigas et al. 2012; Vicente-Agullo et al. 2004). *VvKUP2* (*Vitis vinifera*) promoted the expansion of berry epidermal cells (Davies et al. 2006). *AtKUP2/SHY3* (*Short Hypocotyl 3*) mediated K<sup>+</sup>-dependent cell expansion in growing tissues, with the *shy3-1* mutant plants having shorter hypocotyls, smaller leaves, and

shorter flowering stems than the wild-type plants (Elumalai et al. 2002b). *Arabidopsis* *Kup2/6/8* triple mutants displayed larger plant bodies, thus suggesting their roles as negative regulators during the turgor pressure–dependent growth (Osakabe et al. 2013). *Arabidopsis* root meristem activity was maintained by *KUP9* by regulating the  $K^+$  level and auxin homeostasis at a low  $K^+$  level (Zhang et al. 2020). *HAK5* promoted the expression of *INTEGRIN-LINKED KINASE1* (*ILK1*) to positively regulate plant innate immunity and abiotic stress response in *Arabidopsis* (Brauer et al. 2016).  $K^+$  deficiency stress induced the expression of rice *OsHAK1*, and when overexpressed in plants they display enhanced salt and drought tolerance (Chen et al. 2015; Chen et al. 2017; Chen et al. 2018). Additionally, constitutive overexpression or mutation analysis of *OsHAK5*, *OsHAK21*, and *OsHAK16* demonstrated their role in  $K^+$  homeostasis and salt tolerance (Feng et al. 2019; Horie et al. 2011; Shen et al. 2015). *HvHAK1* confers salt and drought tolerance in barley by enhancing the leaf mesophyll  $H^+$  homeostasis and improving  $K^+$  nutrition (Feng et al. 2020; Mangano et al. 2008).

In plants, the HAK family is the homolog of the bacterial  $K^+$  transporter KUP and fungal  $K^+$  transporter HAK (Bañuelos et al. 1995; Schleyer & Bakker 1993; Very et al. 2014). Based on their hydropathy profiles, the plant HAK proteins were predicted to have 10–14 transmembrane (TM) domains, including a conserved  $K^+$  transport domain (GVVYGD LGTSPLY) (Gierth M and Mäser P 2007; Alonso and Rodríguez-Navarro 2000). Mutation assay analysis revealed that the role of  $K^+$  transport capacity is determined by both the 8<sup>th</sup> TM domain and the C-terminus of HAKs (Alonso and Rodríguez-Navarro 2000; Gomez-Porras et al. 2012; Mangano et al. 2008). Based on phylogenetic analysis, HAK family genes were generally classified into four clusters (I–IV) (Bañuelos et al. 1995; Gupta et al. 2008). HAK family members exhibit significant diversity in their subcellular localizations, including the plasma membrane, tonoplast, endoplasmic reticulum, and other endomembranes (Osakabe et al. 2013; Rigas et al. 2012). Expression analysis revealed that many members of the HAK family were also expressed in the root hairs and root tip cells, thereby implying the HAK family members are involved in  $K^+$  uptake (Ahn et al. 2004; Elumalai et al. 2002b; Qin et al. 2019; Yang et al. 2014). Indeed, several HAK family members have been shown to participate in  $K^+$  uptake and translocation in a few model plants, including *Arabidopsis*, rice, barley, maize, and tomato (Very et al. 2014).

*M. truncatula* has been regarded as a model system for studying legume genetics and its relatively small genome size helps understand nodule symbiosis (Young et al. 2011). Despite the

functional importance of the HAK genes, surprisingly little is known about their family members in *M. truncatula*. In this study, we performed comprehensive genome-wide analyses of the *M. truncatula* HAK family genes including phylogenetic relationships, chromosomal distributions, gene duplications, gene structures, *cis*-acting regulatory elements, and expression patterns in response to both K<sup>+</sup> deficiency and abiotic stress. Finally, these results not only elucidated the structures and expression patterns of 20 *MtHAKs* genes but also laid the foundation for their future functional analysis in *M. truncatula*.

## Materials & Methods

### Identification and sequence analysis of *MtHAKs*

*MtHAKs* sequences were obtained from the *Medicago truncatula* genome databases (HAPMAP, <https://medicagohapmap2.org>). The amino acid (aa) sequences of *Arabidopsis* (TAIR, <http://www.arabidopsis.org/>) and rice (TIGR, <http://rice.plantbiology.msu.edu/>) HAKs were used as the reference sequences for searching predicted homolog sequences in *M. truncatula* using the HMMER3.0 software (<http://hmmer.org/>). Subsequently, the genes were screened using a threshold of < 1e-100 E-value (full sequence and best 1 domain). Candidate protein members were verified using the SMART database (<http://smart.embl-heidelberg.de/>) and NCBI-Conserved Domain Database (CDD, <https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi>) (Zhao et al. 2021), with proteins with shorter aa length (< 400 aa) and those containing incomplete K<sup>+</sup> transporter domains being discarded. The longest gene was chosen for further analysis only if it had alternative splicing variants. Subcellular localization of *MtHAK* proteins was predicted using the WOLF PSORT software (<https://www.genscript.com/wolf-psort.html>) and the TMHMM Server 2.0 online tool (<https://services.healthtech.dtu.dk/service.php?TMHMM-2.0>) was used for predicting the protein transmembrane helices.

### Construction of *MtHAKs* phylogenetic tree

HAK protein sequences of *Arabidopsis* and rice were retrieved from the NCBI database (<https://www.ncbi.nlm.nih.gov>) (Table S1), while multiple sequence alignment was conducted using the ClustalW program (Version 2.1; <http://www.clustal.org/>). MEGA7.0 was used to construct the phylogenetic tree using the Neighbor-Joining method along with the bootstrap replicates being up to 1000 (Liu et al. 2019; Liu et al. 2020).

## Gene structure and conserved motif analysis

Gene structure and conserved motifs were visualized using the TBtools (Toolkit for Biologists integrating various biological data-handling tools) software (Chen et al. 2020). The conserved and identified motifs of protein sequences were predicted via the MEME (Multiple Expectation Maximization for motif Elicitation) program (Version 5.1.1), with the maximum protein motif number being set as 10, and the other parameters set as default (<http://meme-suite.org/tools/meme>) (Bailey et al. 2009).

## Chromosomal location and synteny analysis

The *MtHAK* chromosomal location was illustrated by the circos diagram by annotating genes to their specific chromosomal location in their genome sequences by using the TBtools software. These syntenic analyses were carried out by using the MCScanX with gene duplication parameters (Wang et al. 2012).

## Analysis of *cis*-acting regulatory elements in *MtHAKs* promoter regions

Putative *cis*-acting regulatory elements were analyzed using the PlantCARE online software (<http://bioinformatics.psb.ugent.be/webtools/plantcare/html/>). The 2.0 kb promoter sequences located upstream of the transcription starting site in each *MtHAK* gene were extracted from the *M. truncatula* genome database.

## Analysis of microarray expression profile

The microarray data of the expression profiles of *MtHAKs* in the roots, vegetative bud, stem, petiole, leaf, flower, pods, and seeds and their responses to abiotic stress were obtained from the MtGEA (Benedito et al. 2008). When a gene corresponded to multiple probes, the maximum value of the probe was selected for the subsequent analysis. The normalized microarray data was used to create the heatmap through the TBtools software, based on the mean value of each gene expression in all the analyzed organs. The expression patterns of *MtHAKs* in response to salt, drought, and cold stresses were obtained from the NCBI under GEO accession number GSE136739 (Song et al. 2017). The expression abundance of each *MtHAK* gene was represented by fragments per kilobase million (FPKM). The relative stress-induced expression levels were calculated by comparing with the control samples. The clustered heatmap was generated using the TBtools software and based on their relative expression.

## Stress treatment and qRT-PCR

For K<sup>+</sup> deficiency stress treatment, two-week-old seedlings were grown in 1/2 Hoagland nutrient medium without K<sup>+</sup> for 0 (control), 1, 6, 12, 24, and 48 h, respectively. For the salt stress treatment, two-week-old seedlings were grown in 1/2 Hoagland nutrient medium containing 300 mM NaCl for 0, 1, 6, 12, 24, and 48 h, respectively. For drought stress treatment, two-week-old seedlings were grown in 1/2 Hoagland nutrient medium containing 18% PEG6000 for 0, 1, 6, 12, 24, and 48 h, respectively. The root samples were subsequently cut, then snap frozen in liquid nitrogen, and finally stored at −80°C until further use. The qRT-PCR analysis was performed in triplicates for each of the biological replicates. Their relative expression levels were calculated using the 2<sup>−ΔΔCt</sup> analysis method (Liu et al. 2019; Zhao et al. 2022). The expression levels of the control samples were normalized to 1, with the *MtActin* gene being used as the internal control. Standard deviations and the significant differences were indicated by error bars and “\*” (*p* < 0.05), respectively.

## Results

### Identification of HAK members in *M. truncatula*

To identify *M. truncatula* HAK genes, we conducted a genome-wide search using the HMMER3.0 Software (<http://hmmer.org/>) based on the *M. truncatula* genome sequences along with the *Arabidopsis* and rice HAK genes as subjected queries. Then, we identified 20 nucleotide sequences with a typical canonical K<sup>+</sup> transporter domain (Pfam accession no. PF02705) using the Pfam and SMART databases, and they were subsequently designated as *MtHAK1* to *MtHAK20* depending on their chromosomal positions (Table 1). Detailed information on the 20 HAK genes is listed in Table S1. The number of protein transmembrane segments (TMS) ranged between 10 and 13, with the most common being 12–13 (70%). All the examined HAK proteins were predicted to be mainly localized in the plasma membrane using a PSORT analysis (<http://www.psort.org>). The protein length of the 20 identified HAK proteins ranged from 619 aa (MtHAK3) to 856 aa (MtHAK2) with an average length of 778 aa. Their relative molecular weights (MW) varied from 69.03 kDa (MtHAK3) to 95.67 kDa (MtHAK2). The isoelectric points (pI) ranged from 5.44 (MtHAK8) to 9.39 (MtHAK19).

### HAKs phylogenetic relationship among *M. truncatula*, *Arabidopsis* and rice

To analyze the evolutionary relationships of the MtHAK proteins, we conducted phylogenetic analyses of 60 HAK amino acid sequences (20 - *M. truncatula*, 13 - *Arabidopsis*, and 27 - rice)

to construct a phylogenetic tree using the Neighbor-Joining method. According to the evolutionary tree, we classified all *HAK* members into four major groups: Groups I–IV. Furthermore, we classified the MtHAK proteins into three clusters (from I to III): Cluster I (MtHAK6, 14, 15, 17, and 18), Cluster II (MtHAK4, 7, 9, 10, 11, 12, 13, 19, and 20), and Cluster III (MtHAK1, 2, 3, 17, 5, and 8) (Figure 1). All members in group IV belong to rice. The most members existed in Cluster II in *M. truncatula*, thus comprising 45% of all MtHAKs. The phylogenetic tree showed that MtHAKs were most closely related to *Arabidopsis* KUPs than those of rice HAKs, thereby indicating that MtHAKs might share evolutionary functional similarities with *Arabidopsis* KUPs. All MtHAKs in cluster I were distributed together with the already-identified AtHAK5, which suggested that they may be crucial for K<sup>+</sup> uptake from a low-K<sup>+</sup> level soil (Lara et al. 2020). Among cluster II members, MtHAK4 and MtHAK19 shared high sequence identity with AtKUP2 (Elumalai et al. 2002), and AtKUP4 (Rigas et al. 2001; Vicente-Agullo et al. 2004), respectively, thus implying they are likely to be involved in plant development processes. Additionally, among cluster III, MtHAK1 and MtHAK8 clustered together with AtKUP7 (Han et al. 2016), thereby suggesting their role in K<sup>+</sup> acquisition and translocation under low K<sup>+</sup> concentration.

### Gene structure and motif composition of *MtHAK* genes

MtHAK proteins were listed in order based on the phylogenetic analysis (Figure 2A), which was consistent with the results in Figure 1. Closely related members shared similar exon/intron structures, which were related to their biological functions. Gene structures of the *MtHAKs* were obtained based on the arrangement of the untranslated region, exon, and intron sequences generated using the TBTools software. As shown in Figure 2B, the exon number of *MtHAK* genes varied from 8 to 10, and the longest exon existed in the end of a gene except for *MtHAK2*, which is consistent with previous reported data (He et al. 2012; Hyun et al. 2014). Additionally, most *MtHAKs* in the same cluster shared high exon-intron structure similarity (Figure 2B).

To study the structural features, we analyzed conserved protein motifs of MtHAKs using the MEME program. We identified the conserved protein motifs varying from 29 to 50 aa in lengths and designated them as motifs 1–10. Conserved protein motif information is shown in Table S2. The highly conserved K<sup>+</sup> transport domain (GVVYGD LGTSPLY), included in motif 9, existed in all MtHAK proteins besides MtHAK10 (Figure2, Table S2). Motifs 1, 2, 3, 4, 5, 6, 7, 8, and 10 were almost evenly distributed along with a feature domain of K<sup>+</sup> transporter (Figure



2C, Table S2) in all the MtHAK proteins. Therefore, the motifs of conserved K<sup>+</sup> transporter and similarities of gene structure in the same cluster together implied the closing function among these HAK members.

### **Chromosomal distribution and synteny analysis of *MtHAK* genes**

All identified *MtHAK* genes were mapped onto chromosomes from the *M. truncatula* genome database to identify and locate their chromosomal distribution. Results showed that *MtHAKs* were distributed on seven of the eight chromosomes, with chromosome 8 containing the highest number of six *MtHAK* genes (Figure 3). Five *MtHAK* genes were located on chromosome 5, three on chromosome 2, two on chromosome 4 and 6, one on chromosome 3 and 7, and no gene was allocated on chromosome 1 (Figure 3). These results indicated that *MtHAKs* were scattered randomly onto different chromosome locations.

We further performed synteny analysis between *M. truncatula* and *Arabidopsis* to verify the evolutionary relationships and history of the *MtHAKs*. Subsequently, we found seven collinear gene pairs between *M. truncatula* and *Arabidopsis* in the dataset (Figure 3 and Table S3). This indicated that these identified genes might already have existed before protein structure divergence, thereby further implying a strong phylogenetic relationship. Furthermore, only one gene pair (*MtHAK2/MtHAK5*) existed as paralogs in *M. truncatula*.

### **Analysis of *cis*-acting elements in the promoter region of *MtHAK* genes**

To further investigate the gene function and regulatory mechanism of *MtHAKs*, we analyzed the 2 kb regions upstream of the translation start site of the 20 *MtHAK* genes using the PlantCARE database. We identified 73 putative *cis*-elements in the *MtHAK* promoters based on functional annotation, and the major types of *cis*-elements are shown in Figure 4 and Table S4. Post analysis, we identified *cis*-elements corresponding to different plant hormones like auxin (TGA-element and AuxRE-core), gibberellin (GARE-motif and P-box), MeJA (TGACG-motif and CGTCA-motif), ethylene (ERE-box), ABA (ABRE), and salicylic acid (TCA-element), in the promoter regions of all *MtHAKs* genes except *MtHAK20*, thereby suggesting that *MtHAKs* expression may be regulated by different phytohormones. Furthermore, we also found abiotic stress-responsive elements, including STRE, ARE, WRE3, WUN-motif, MBS, LTR, DRE-core, DRE1, and TC-rich repeats, in all the *MtHAKs* promoter regions except *MtHAK20*. Additionally, zein metabolism regulation element (O2-site), endosperm expression element (GCN4-motif and AACA-motif), palisade mesophyll cells element (HD-Zip 1), meristem expression element

(CAT-box and CCGTCC-motif), and seed regulation element RY-element were also abundant in the promoter of *MtHAKs* except *MtHAK20*. However, the *MtHAK20* promoter region was abundant in light-responsive elements (Table S4).

### **Spatial expression profiles of *MtHAK* genes**

To gain further insights into the potential biological function of *MtHAK* genes, we used the publicly available microarray data of the *Medicago truncatula* Gene Expression Atlas (MtGEA, <https://mtgea.noble.org/v3/>) to investigate the temporal and spatial expression pattern of the *MtHAKs*. *MtHAK4* showed relatively high expression in all tissues, while that of *MtHAK18* was low in all tissue (Table S5). Notably, *MtHAK6* and *MtHAK16* were expressed preferentially in the roots, thereby implicating their role in K<sup>+</sup> uptake from the soil (Figure 5, Table S5).

Both cluster III genes, *MtHAK2* and *MtHAK3* exhibited similar expression patterns and relatively high expression in leaves. *MtHAK13* was exclusively and highly expressed in floral organs, whereas *MtHAK8* showed the same in immature seeds (Figure 5, Table S5). Interestingly, *MtHAK5* and *MtHAK12* exhibited high and gradually increased expression patterns during the reproductive stages and finally peaked at 24 days after pollination (DAP). Contrastingly, *MtHAK15* was specifically highly expressed in immature seeds (10 DAP) with the expression pattern gradually decreasing along with seed maturation. Therefore, the spatial and temporal expression profiles indicated the functional diversity of *MtHAK* genes in *Medicago truncatula* development.

### **Expression patterns of *MtHAK* genes under K<sup>+</sup> deficiency**

Due to the major function of the HAK family being K<sup>+</sup> transport, we investigated the expression profiles of *MtHAK* genes in the roots under K<sup>+</sup> deficient conditions using qRT-PCR. As shown in Figure 6, among the 20 *MtHAK* genes, we obtained eight genes that showed upregulated expression patterns post K<sup>+</sup> deficiency treatment. *MtHAK6*, *MtHAK7*, and *MtHAK17* expression slightly increased and finally peaked at 48 h post treatment. *MtHAK15* and *MtHAK18* showed nearly the same expression pattern at the five different time points. *MtHAK9*, *MtHAK10*, and *MtHAK11* transcripts were strongly upregulated at 6 h, then peaking at 12 h and 24 h, and finally went down at 48 h. Therefore, these results suggested that these *MtHAK* genes were K<sup>+</sup> deficiency-responsive. Furthermore, it was noteworthy that *MtHAK6* was highly and specifically expressed in *Medicago truncatula* roots and also significantly upregulated in response to K<sup>+</sup> deficiency.

## Expression patterns of *MtHAK* genes under salt and drought stresses

Several *HAK* genes have been reported to participate in abiotic stresses (Elumalai et al. 2002; Vicente-Agullo et al. 2004; Chen et al. 2015; Shen et al. 2015). To verify this hypothesis, we evaluated the expression profiles of eight K<sup>+</sup> deficiency responsive genes via qRT-PCR under salt and drought stress treatments. The results revealed that all eight genes were induced by salt and drought stresses to different extents (Figures 7 and 8).

We determined the expression profile of *MtHAK* genes in *Medicago truncatula* roots at different times (0 h, 1 h, 6 h, 12 h, 24 h, and 48 h) under salt treatment (300 mM NaCl in nutrient solution). The results showed that expression of *MtHAK7*, *MtHAK9*, *MtHAK15*, *MtHAK17*, and *MtHAK18* exhibited significant upregulation. Interestingly, *MtHAK7* and *MtHAK18* were quickly and continuously upregulated from 1 h and subsequently increased at 48 h (Figure 7).

Additionally, we analyzed the expression profiles of *MtHAK* genes in *Medicago truncatula* roots under drought treatment simulated by 18% PEG6000 at different times (0 h, 1 h, 6 h, 12 h, 24 h, and 48 h). Under drought treatment, all selected genes besides *MtHAK9* were upregulated, albeit to different levels at different times (Figure 8). In particular, *MtHAK10*, *MtHAK15*, and *MtHAK18* rapidly responded to dehydration at 1 h. Contrastingly, *MtHAK17* was moderately upregulated from 6 h to 48 h. Both *MtHAK6* and *MtHAK7* exhibited highly induced expression at 24 h.

Interestingly, we found that *MtHAK15*, *MtHAK17*, and *MtHAK18* were strongly upregulated by both salt and drought stresses. The expression level of *MtHAK18* increased rapidly at 1 h as compared to the control, under both salt and drought treatments (Figures 7 and 8).

## Discussion

*HAK* family genes play key roles not only in K<sup>+</sup> acquisition and uptake, but also in plant growth, development, and abiotic stress response (Osakabe et al. 2013; Zhao et al. 2016). Although comprehensive genome-wide analysis of the *HAK* gene family has been widely reported in various plants, studies of the *HAK* gene family in the model legume *M. truncatula* were still lacking (Ahn et al. 2004; Gupta et al. 2008; Zhang et al. 2012). The release of the *M. truncatula* genome information makes it possible to systematically characterize and identify the *HAK* genes.

In this study, we identified 20 *HAK* genes in *M. truncatula*. We characterized their genetic structures as well as their expression patterns in different tissues and also during stress responses.

We classified the 20 identified *HAK* members into three clusters (clusters I to III) based on the evolutionary relationships, which was consistent with the previous classification in *Arabidopsis* (Figure 1) (Rubio et al. 2000). Phylogenetic analysis of *HAK* proteins revealed that *MtHAKs* shared higher similarity with *AtHAKs* as compared to the *OsHAKs* (Figure 1), thereby suggesting that *MtHAKs* may share similar functionality with *Arabidopsis AtHAKs*. Gene structure analysis showed that *MtHAK* genes contained 8–10 exons, with the last exon being the longest. However, the exception was *MtHAK2*, which was consistent with the previously reported exon-intron structure of *HAKs* (He et al. 2012; Hyun et al. 2014). Conserved protein motif analysis indicated that all the identified *MtHAKs* had at least five typical K<sup>+</sup> transporter motifs.

The tissue-specific gene expression patterns reflect their function and potential biological roles in plants. Approximately 10 of the 13 *Arabidopsis AtHAK* genes were strongly expressed in the root (Ahn et al. 2004). *MtHAK6* was preferentially highly expressed in the roots, and it belonged to the same clades of *AtHAK5* in the phylogenetic tree, which was also expressed in roots and mediated high-affinity root K<sup>+</sup> uptake (Lara et al. 2020), thereby implicating their role in K<sup>+</sup> acquisition from the soil (Figure 5, Table S5). *MtHAK16* shared high similarity with *AtKUP12*, which showed root hair-specific expression (Ahn et al. 2004). Therefore, these results may help elucidate the biological function of *Arabidopsis* orthologous *MtHAK* genes in K<sup>+</sup> acquisition in *M. truncatula*.

Some plant *HAK* genes were shown to participate in plant growth and development. For instance, *AtKUP4/TRH1* mutation impaired the root gravitropism response and root hair elongation (Rigas et al. 2001; Rigas et al. 2012; Vicente-Agullo et al. 2004). Knockout of *AtKT2/KUP2* caused shorter hypocotyl length, small rosette leaves, and short flowering stem phenotype (Elumalai et al. 2002). *MtHAK13* was exclusively and highly expressed in the floral organs, while showing low expression levels in other tissues, thereby suggesting its critical role in floral development. *MtHAK5*, *MtHAK8*, *MtHAK12*, and *MtHAK15* were specifically and highly expressed during the reproductive stages, thus implying their roles in facilitating seed maturation and maintaining fertility. The varied tissue expression pattern of the *MtHAK* genes indicated their diverse functions in plants.

Under  $K^+$  deficiency conditions, plants maintain cytosolic  $K^+$  homeostasis by uptaking  $K^+$  through HAKs, and these  $K^+$  transporter genes represent a major transcriptional regulation mechanism during low-  $K^+$  stress. AtHAK5 and AtKT1 are two essential transporters mediating high-affinity  $K^+$  uptake in the *Arabidopsis* roots, with the roots of their double-mutant unable to sustain plant growth (Lara et al. 2020). ZmHAK5 was characterized as a high-affinity  $K^+$  transporter in maize (Qin et al. 2019). The expression of OsHAK1 and OsHAK5 were significantly upregulated in roots under low  $K^+$  conditions, thereby maintaining the  $K^+$  uptake and translocation from the root to the shoot (Chen et al. 2015; Chen et al. 2017; Chen et al. 2018; Yang et al. 2014). We found that  $K^+$  deficiency upregulated the root-specific expression of eight *MtHAKs*, especially *MtHAK10* and *MtHAK11* (Figure 6). *MtHAK6* was preferentially and highly expressed in the roots, and it increased under  $K^+$  deficiency stress (Figure 6). Therefore, we expect that several *HAK* genes could increase the  $K^+$  absorption capacity during  $K^+$  deficiency. Previous studies reported that *HAK* genes were crucial for regulating water potential and turgor pressure during osmotic adjustment. These genes also positively regulated plant stress responses by regulating the balance of  $K^+$  influx/efflux balance, e.g., *OsHAK1* expression increases in the rice roots post the  $K^+$ -deficient condition and it positively regulated the salt and drought stress tolerance response (Chen et al. 2015; Chen et al. 2017; Chen et al. 2018). Consistent with the above results, in our study, e.g., *MtHAK7*, *MtHAK9*, *MtHAK15*, *MtHAK17*, and *MtHAK18* exhibited significantly upregulated expression levels in *M. trunculata* roots under salt stress (Figure 7). Interestingly, many *cis*-acting elements related to phytohormones, plant growth and development, and abiotic stress response, were extensively distributed in the promoter regions of the *MtHAKs* (Figure 4). ABREs are drought-stress responsive elements (Sah et al. 2016). *MtHAK* genes contained ABRE elements in their promoters, which further implied that *MtHAKs* participate in drought responses (Figures 4 and 6). Moreover, the qRT-PCR analysis showed that the expressions of most of the selected *MtHAK* genes were noticeably upregulated after drought stress. Notably, *MtHAK6*, which was preferentially highly expressed in the roots, was also significantly upregulated post drought stresses (Figure 8). In particular, the expression of *MtHAK15*, *MtHAK17*, and *MtHAK18* were strongly and specifically upregulated in *M. truncatula* roots under  $K^+$  deficiency, salt, and drought stress conditions, thus implying that these genes are potential candidates for high-affinity  $K^+$  uptake while also being essential in salt and drought tolerance.

## Conclusions

Based on phylogenetic analysis, we identified and characterized 20 MtHAK protein sequences from *M. truncatula* which were grouped into three clusters. Furthermore, we analyzed the chromosome location, conserved protein motif, and gene structure of all the *M. truncatula* HAK genes. The *cis*-acting elements regulating plant growth and development, or those responsive to phytohormone and abiotic stress were abundant in the promoter regions of MtHAKs. Gene expression analysis assay revealed that MtHAKs exhibited diverse tissue-specific expression patterns in various tissues using the publicly available RNA-seq data. Additionally, eight upregulated genes showed varied expression patterns post the K<sup>+</sup> deficiency treatment. The expression pattern analysis under K<sup>+</sup> deficiency, drought, and salt stress suggested that these genes are candidates for high-affinity K<sup>+</sup> uptake that are also crucial in drought and salt tolerance. Therefore, these results provide the first genetic description of the K<sup>+</sup> transporter family in *M. truncatula*, while also laying the foundation for molecular breeding of stress-resistant legume crops in the future.

## Acknowledgements

We would like to thank the reviewers for their comments and suggestion.

## References

- Gierth M and Mäser P. 2007. Potassium transporters in plants – Involvement in K<sup>+</sup> acquisition, redistribution and homeostasis. *FEBS Letters* **581**:2348-2356.
- Ahn SJ, Shin R, and Schachtman DP. 2004. Expression of KT/KUP genes in *Arabidopsis* and the role of root hairs in K<sup>+</sup> uptake. *Plant Physiology* **134**:1135-1145.
- Alonso and Rodríguez-Navarro. 2000. Potassium transport in fungi and plant. *Biochimica Et Biophysica Acta Reviews on Biomembranes*. **1469** (1):1-30.
- Amrutha RN, Sekhar PN, Varshney RK, and Kishor PBK. 2007. Genome-wide analysis and identification of genes related to potassium transporter families in rice (*Oryza sativa* L.). *Plant Science* **172**:708-721.
- Ashley MK, Grant M, and Grabov A. 2005. Plant responses to potassium deficiencies: a role for potassium transport proteins. *Journal of Experimental Botany* **57**:425-436.
- Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW, and Noble

- WS. 2009. MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Research* **37**:W202-208.
- Bañuelos MA, Klein RD, Alexander-Bowman SJ, and Rodríguez-Navarro A. 1995. A potassium transporter of the yeast *Schwanniomyces occidentalis* homologous to the Kup system of *Escherichia coli* has a high concentrative capacity. *Embo Journal* **14**:3021-3027.
- Benedito VA, Torres-Jerez I, Murray JD, Andriankaja A, and Udvardi MK. 2008. A gene expression atlas of the model legume *Medicago truncatula*. *The Plant Journal* **55**:504-513.
- Brauer EK, Ahsan N, Dale R, Kato N, Coluccio A, Piñeros MA, Kochian LV, Thelen JJ, and Popescu SC. 2016. The raf-like kinase ILK1 and the high affinity K<sup>+</sup> transporter HAK5 are required for innate immunity and abiotic stress response. *Plant Physiology* **171**(2):1470-1484.
- Chen C, Chen H, Zhang Y, Thomas HR, Frank MH, He Y, and Xia R. 2020. TBtools: an integrative toolkit developed for interactive analyses of big biological data. *Molecular Plant* **13**:1194-1202.
- Chen G, Hu Q, Luo L, Yang T, Zhang S, Hu Y, Yu L, and Xu G. 2015. Rice potassium transporter OsHAK1 is essential for maintaining potassium-mediated growth and functions in salt tolerance over low and high potassium concentration ranges. *Plant Cell and Environment* **38**:2747-2765.
- Chen G, Liu C, Gao Z, Zhang Y, Jiang H, Zhu L, Ren D, Yu L, Xu G, and Qian Q. 2017. OsHAK1, a high-affinity potassium transporter, positively regulates responses to drought stress in rice. *Frontiers in Plant Science* **8**:1885.
- Chen G, Zhang Y, Ruan B, Guo L, Zeng D, Gao Z, Zhu L, Hu J, Ren D, Yu L, Xu G, and Qian Q. 2018. OsHAK1 controls the vegetative growth and panicle fertility of rice by its effect on potassium-mediated sugar metabolism. *Plant Science* **274**:261-270.
- Corratge-Faillie C, Jabnune M, Zimmermann S, Very AA, Fizames C, and Sentenac H. 2010. Potassium and sodium transport in non-animal cells: the Trk/Ktr/HKT transporter family. *Cellular and Molecular Life Science* **67**:2511-2532.
- Davies C, Shin R, Liu W, Thomas MR, and Schachtman DP. 2006. Transporters expressed during grape berry (*Vitis vinifera* L.) development are associated with an increase in berry

- size and berry potassium accumulation. *Journal of Experimental Botany* **57**:3209-3216.
- Elumalai RP, Nagpal P, and Reed JW. 2002a.** A mutation in the *Arabidopsis* *KT2/KUP2* potassium transporter gene affects shoot cell expansion. *Plant Cell* **14**:119-131.
- Feng H, Tang Q, Cai J, Xu B, Xu G, and Yu L. 2019.** Rice *OsHAK16* functions in potassium uptake and translocation in shoot, maintaining potassium homeostasis and salt tolerance. *Planta* **250**:549-561.
- Feng X, Liu W, Qiu CW, Zeng F, Wang Y, Zhang G, Chen ZH, and Wu F. 2020.** *HvAKT2* and *HvHAK1* confer drought tolerance in *barley* through enhanced leaf mesophyll  $H^{(+)}$  homeostasis. *Plant Biotechnology Journal* **18**:1683-1696.
- Gierth M, Mäser P, and Schroeder JI. 2005.** The potassium transporter *AtHAK5* functions in  $K^{(+)}$  deprivation-induced high-affinity  $K^{(+)}$  uptake and *AKT1*  $K^{(+)}$  channel contribution to  $K^{(+)}$  uptake kinetics in *Arabidopsis* roots. *Plant Physiology* **137**:1105-1114.
- Gomez-Porras JL, Riaño-Pachón D, Benito B, Haro R, Sklodowski K, Rodríguez-Navarro A, and Dreyer I. 2012.** Phylogenetic analysis of  $K^{+}$  transporters in bryophytes, lycophytes, and flowering plants indicates a specialization of vascular plants. *Frontiers in Plant Science* **3**:167.
- Gupta M, Qiu XH, Wang L, Xie WB, Zhang CJ, Xiong LZ, Lian XM, and Zhang QF. 2008.** KT/HAK/KUP potassium transporters gene family and their whole-life cycle expression profile in rice (*Oryza sativa*). *Molecular Genetics and Genomics* **280**:437-452.
- Han M, Wu W, Wu WH, and Wang Y. 2016.** Potassium transporter KUP7 Is involved in  $K^{(+)}$  acquisition and translocation in *Arabidopsis* root under  $K^{(+)}$ -Limited conditions. *Molecular Plant* **9**:437-446.
- He CY, Cui K, Duan AG, Zeng YF, and Zhang JG. 2012.** Genome-wide and molecular evolution analysis of the poplar KT/HAK/KUP potassium transporter gene family. *Ecology and Evolution* **2**:1996-2004.
- Horie T, Sugawara M, Okada T, Taira K, Kaothien-Nakayama P, Katsuhara M, Shinmyo A, and Nakayama H. 2011.** Rice sodium-insensitive potassium transporter, OsHAK5, confers increased salt tolerance in tobacco BY2 cells. *Journal of Bioscience and Bioengineering* **111**:346-356.
- Hyun TK, Rim Y, Kim E, and Kim JS. 2014.** Genome-wide and molecular evolution analyses of the KT/HAK/KUP family in tomato (*Solanum lycopersicum* L.). *Genes & Genomics*



- 36:365-374.
- Lara A, Rodenas R, Andres Z, Martinez V, Quintero FJ, Nieves-Cordones M, Botella MA, and Rubio F. 2020.** *Arabidopsis* K<sup>+</sup> transporter HAK5-mediated high-affinity root K<sup>+</sup> uptake is regulated by protein kinases CIPK1 and CIPK9. *Journal of Experimental Botany* **71**:5053-5060.
- Li W, Xu G, Alli A, and Yu L. 2018.** Plant HAK/KUP/KT K<sup>+</sup> transporters: function and regulation. *Seminars in Cell & Developmental Biology* **74**:133-141.
- Liu H, Guo S, Lu M, Zhang Y, Li J, Wang W, Wang P, Zhang J, Hu Z, Li L, Si L, Zhang J, Qi Q, Jiang X, Botella JR, Wang H, and Song CP. 2019.** Biosynthesis of DHGA12 and its roles in *Arabidopsis* seedling establishment. *Nature Communications* **10**:1768.
- Liu Z, Haider MS, Khan N, and Fang J. 2020.** Comprehensive sequence analysis of IQD gene family and their expression profiling in Grapevine (*Vitis vinifera*). *Genes* **11**(2):235.
- Mangano S, Silberstein S, and Santa-María G. 2008.** Point mutations in the barley HvHAK1 potassium transporter lead to improved K<sup>+</sup>-nutrition and enhanced resistance to salt stress. *FEBS Letters* **582**(28):3922-3929.
- Osakabe Y, Arinaga N, Umezawa T, Katsura S, Nagamachi K, Tanaka H, Ohiraki H, Yamada K, Seo SU, and Abo M. 2013.** Osmotic stress responses and plant growth controlled by potassium transporters in *Arabidopsis*. *Plant Cell* **25**:609-624.
- Qin Y, Wu W, and Wang Y. 2019.** ZmHAK5 and ZmHAK1 function in K<sup>+</sup> uptake and distribution in maize under low K<sup>+</sup> conditions. *Journal of Integrative Plant Biology* **61**(6):691-705.
- Rigas S, Debrosses G, Haralampidis K, Vicente-Agullo F, and Hatzopoulos P. 2001.** TRH1 encodes a potassium transporter required for tip growth in *Arabidopsis* root hairs. *Plant Cell* **13**(1):139-151.
- Rigas S, Ditengou FA, Ljung K, Daras G, Tietz O, Palme K, and Hatzopoulos P. 2012.** Root gravitropism and root hair development constitute coupled developmental responses regulated by auxin homeostasis in the *Arabidopsis* root apex. *New Phytologist* **197**(4):1130-1141.
- Rubio F, Santa-Maria GE, and Rodriguez-Navarro A. 2000.** Cloning of *Arabidopsis* and barley cDNAs encoding HAK potassium transporters in root and shoot cells. *Physiologia Plantarum* **109**:34-43.

- Sah SK, Reddy KR, and Li J. 2016. Absciscic acid and abiotic stress tolerance in crop plants. *Frontiers in Plant Science* 7:571.
- Schleyer M, and Bakker EP. 1993. Nucleotide sequence and 3'-end deletion studies indicate that the K<sup>(+)</sup>-uptake protein kup from Escherichia coli is composed of a hydrophobic core linked to a large and partially essential hydrophilic C terminus. *Journal of Bacteriology* 175:6925-6931.
- Shen Y, Shen L, Shen Z, Jing W, Ge H, Zhao J, and Zhang W. 2015. The potassium transporter OsHAK21 functions in the maintenance of ion homeostasis and tolerance to salt stress in rice. *Plant Cell and Environment* 38:2766-2779.
- Song J, Mo X, Yang H, Yue L, Song J, and Mo B. 2017. The U-box family genes in Medicago truncatula: Key elements in response to salt, cold, and drought stresses. *PLoS One* 12:e0182402.
- Very AA, Nieves-Cordones M, Daly M, Khan I, Fizames C, and Sentenac H. 2014. Molecular biology of K<sup>+</sup> transport across the plant cell membrane: what do we learn from comparison between plant species? *Journal of Plant Physiology* 171:748-769.
- Vicente-Agullo F, Rigas S, Desbrosses G, Dolan L, and Grabov A. 2004. Potassium carrier TRH1 is required for auxin transport in Arabidopsis roots. *Plant Journal for Cell & Molecular Biology* 40:523-535.
- Wang Y, Tang H, Debarry JD, Tan X, Li J, Wang X, Tae-Ho L, Jin H, Barry M, and Guo H. 2012. MCScanX: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. *Nucleic Acids Research* 40:e49-e49.
- Yang T, Zhang S, Hu Y, Wu F, Hu Q, Chen G, Cai J, Wu T, Moran N, Yu L, and Xu G. 2014. The role of a potassium transporter OsHAK5 in potassium acquisition and transport from roots to shoots in rice at low potassium supply levels. *Plant Physiology* 166:945-959.
- Young ND, Debelle F, Oldroyd GED, Geurts R, Cannon SB, Udvardi MK, Benedito VA, Mayer KFX, Gouzy J, Schoof H, Van de Peer Y, Proost S, Cook DR, Meyers BC, Spannagl M, Cheung F, De Mita S, Krishnakumar V, Gundlach H, Zhou SG, Mudge J, Bharti AK, Murray JD, Naoumkina MA, Rosen B, Silverstein KAT, Tang HB, Rombauts S, Zhao PX, Zhou P, Barbe V, Bardou P, Bechner M, Bellec A, Berger A, Berges H, Bidwell S, Bisseling T, Choisine N, Couloux A, Denny R,

- Deshpande S, Dai XB, Doyle JJ, Dudez AM, Farmer AD, Fouteau S, Franken C, Gibelin C, Gish J, Goldstein S, Gonzalez AJ, Green PJ, Hallab A, Hartog M, Hua A, Humphray SJ, Jeong DH, Jing Y, Jocker A, Kenton SM, Kim DJ, Klee K, Lai HS, Lang CT, Lin SP, Macmil SL, Magdelenat G, Matthews L, McCorrison J, Monaghan EL, Mun JH, Najar FZ, Nicholson C, Noirot C, O'Bleness M, Paule CR, Poulain J, Prion F, Qin BF, Qu CM, Retzel EF, Riddle C, Sallet E, Samain S, Samson N, Sanders I, Saurat O, Scarpelli C, Schiex T, Segurens B, Severin AJ, Sherrier DJ, Shi RH, Sims S, Singer SR, Sinharoy S, Sterck L, Viollet A, Wang BB, Wang KQ, Wang MY, Wang XH, Warfsmann J, Weissenbach J, White DD, White JD, Wiley GB, Wincker P, Xing YB, Yang LM, Yao ZY, Ying F, Zhai JX, Zhou LP, Zuber A, Denarie J, Dixon RA, May GD, Schwartz DC, Rogers J, Quetier F, Town CD, and Roe BA. 2011. The Medicago genome provides insight into the evolution of rhizobial symbioses. *Nature* **480**:520-524.
- Zhang ML, Huang PP, Ji Y, Wang S, Wang SS, Li Z, Guo Y, Ding Z, Wu WH, and Wang Y. 2020. KUP9 maintains root meristem activity by regulating K<sup>+</sup> and auxin homeostasis in response to low K. *EMBO Reports* **21**(6):e50164.
- Zhang Z, Zhang J, Chen Y, Li R, Wang H, and Wei J. 2012. Genome-wide analysis and identification of *HAK* potassium transporter gene family in maize (*Zea mays* L.). *Molecular Biology Reports* **39**:8465-8473.
- Zhao P, Liu Y, Kong W, Ji J, Cai T, and Guo Z. 2021. Genome-wide identification and characterization of calcium-dependent protein kinase (*CDPK*) and *CDPK-related kinase* (*CRK*) gene families in *medicago truncatula*. *International Journal of Molecular Science* **22**(3):1044.
- Zhao S, Zhang M-L, Ma T-L, and Wang Y. 2016. Phosphorylation of ARF2 relieves Its repression of transcription of the K<sup>+</sup> transporter gene *HAK5* in response to low potassium stress. *Plant Cell* **28**:3005-3019.
- Zhao Y, Wang L, Sun X, Bao Y, Liu H, and Zhao Y. 2022. Inhibition of ribosome biogenesis by actinomycin D affects Arabidopsis root development. *Biochem Biophys Res Commun* **588**:61-67.

**Table 1** (on next page)

Characteristics of *MtHAK* genes in *M. truncatula*

Gene name	Gene ID	No. of aa	MW (kDa)	pI	TMS	Subcellular localization
MtHAK1	Medtr2g008820.1	849	94.74	5.66	11	PM
MtHAK2	Medtr2g438150.1	856	95.67	8.26	13	PM
MtHAK3	Medtr2g438160.1	619	69.03	9.28	13	PM
MtHAK4	Medtr3g094090.1	794	89.3	7.24	13	PM
MtHAK5	Medtr4g094660.1	787	88	8.08	13	PM
MtHAK6	Medtr4g099260.1	815	90.92	8.74	11	PM
MtHAK7	Medtr5g034500.1	782	87.25	8.24	13	PM
MtHAK8	Medtr5g070670.1	849	95.05	5.44	12	PM
MtHAK9	Medtr5g071630.1	725	81.46	6.63	12	PM
MtHAK10	Medtr5g071827.1	666	74.26	7.22	10	PM
MtHAK11	Medtr5g071860.1	754	84.24	7.02	11	PM
MtHAK12	Medtr6g007697.1	776	87.1	7.71	13	PM
MtHAK13	Medtr6g033165.1	819	91.56	8.71	12	PM
MtHAK14	Medtr7g108480.1	773	87.24	7.77	12	PM
MtHAK15	Medtr8g022130.1	766	85.45	7.77	12	PM
MtHAK16	Medtr8g063840.1	840	93.11	6.51	12	PM
MtHAK17	Medtr8g063900.1	745	83.49	8.45	11	PM
MtHAK18	Medtr8g088200.1	782	87.17	9.03	10	PM
MtHAK19	Medtr8g099090.1	792	88.52	9.39	12	PM
MtHAK20	Medtr8g107510.1	782	86.83	8.32	12	PM

1 aa, amino acid; MW, molecular weight; pI, isoelectric points; TMS, transmembrane segments;


2 PM, plasma membrane.

3

# Figure 1

Phylogenetic analysis of HAK proteins in *M. truncatula* (red circle), *A. thaliana* (green triangle), and *O. sativa* (blue square).

The tree was constructed using MEGA7.0 software by the Neighbor-Joining method. The numbers next to the branch represent the 1000 bootstrap replicates expressed in percentage.




1

# Figure 2

Phylogenetic tree, gene structure, and conserved motifs of the HAKs in *M. truncatula*.

(A) Phylogenetic tree of the MtHAK proteins. (B) Exon-intron structure distribution. (C) Conserved protein motifs.






1  
 2

# Figure 3

The synteny analysis of MtHAKs displayed between the *M. truncatula* and *Arabidopsis* genomes.

The *M. truncatula* and *Arabidopsis* chromosomes are represented by yellow and green boxes, respectively. Blue lines indicate the collinear relationship of *MtHAKs* between *M. truncatula* and *Arabidopsis*, while green lines indicate the *MtHAK* gene pairs.

1



2

# Figure 4

Analysis of the *cis*-acting regulatory elements in the promoter region of the *MtHAK* genes.

Depending on the functional annotation, the elements were classified into three main categories: phytohormone-responsive, abiotic stress-responsive, and plant growth and development-related. The frequency of these elements in the promoter region was represented by the numbers and the depth of the red color.

1

	Phytohormone responsive									Abiotic stress									Growth and development							
	TGA-element	AuxRR-core	GARE-motif	P-box	ERE	ABRE	TGACG-motif	TGA-element	CGTCA	STRE	ARE	WRE3	MBS	WUN-motif	LTR	DRE-core	TC-rich repeats	DRE1	O2-site	GCN4	CAT-box	AACA_motif	RY-element	CCGTCC-box	MSA-like	HD-Zip 1
MhHAK1				2	2	4		1		1	4			4					1	1						
MhHAK2	1					1		1	1	1			1	1							1	1	1			
MhHAK3						1	4	1	1	1									1	1	1	1				
MhHAK4						4				1	3									1	1			2		
MhHAK5		1		1					1	3			1	2			2			1						
MhHAK6	1				1	1	3	3	3	3	2				1	1	1									1
MhHAK7	3							1	1			1	2	1					1							
MhHAK8	2					1	2	2	2			1	1													
MhHAK9	1					1	1	1	1	1						2	1									
MhHAK10	1									1				1												
MhHAK11	2					2	1	1	1	2	1		1	1												
MhHAK12				1				1	1	1	3				1		1		1							
MhHAK13						3		1	1	1	2	1	2	1					1						1	
MhHAK14	1					1	2	2	2		2		1					1								
MhHAK15	2						1		1	1	1	2	1	1						1						
MhHAK16						4	4	2	1	2	2							1	1		1					
MhHAK17	1	1	1			1		3	1	3	1	3	1	2	1				2				4			
MhHAK18						2	3				1	2			1		1									
MhHAK19	2					3	2	3	3		5			1	1		1				3		2			
MhHAK20																										


2

3

# Figure 5

Expression patterns of the *MtHAK* genes in different developmental tissues.

The microarray data were normalized based on the mean value of each gene in all the analyzed plant organs. The heat map was portrayed by the relative expressions after  $\log_2$  transformed.




# Figure 6

Relative expression of the *MtHAK* genes in response to K<sup>+</sup> deficiency treatment.

Two-week-old seedlings were placed in K<sup>+</sup> deficient conditions for 0, 1, 6, 12, 24, and 48 h. Mean values and standard errors were calculated from three biological replicates. \* indicates the significant difference between K<sup>+</sup> deficiency and control at  $p < 0.05$ .






# Figure 7

Relative expression of the *MtHAK* genes in response to salt stress.

Two-week-old seedlings were treated with 300 mM NaCl for 0, 1, 6, 12, 24, and 48 h. Mean values and standard errors were calculated from three biological replicates. \*indicates the significant difference between the salt-stressed and control at  $p < 0.05$ .



# Figure 8

Relative expression of the *MtHAK* genes in response to drought stress.

Two-week-old seedlings were treated with 18% PEG6000 for 0, 1, 6, 12, 24, and 48 h.

Meanvalues and standard errors were calculated from three biological replicates. \*indicates the significant difference between the drought-stressed and control at  $p < 0.05$ .

