

Objective monitoring of functional recovery after total knee and hip arthroplasty using sensor-derived gait measures

Ramon Boekesteijn^{Corresp., 1, 2}, José Smolders³, Vincent Busch³, Noël Keijsers¹, Alexander Geurts², Katrijn Smulders¹

¹ Department of Research, Sint Maartenskliniek, Nijmegen, The Netherlands

² Department of Rehabilitation, Donders Institute for Brain Cognition and Behaviour, Radboud University Medical Center, Nijmegen, The Netherlands

³ Department of Orthopedic Surgery, Sint Maartenskliniek, Nijmegen, The Netherlands

Corresponding Author: Ramon Boekesteijn
Email address: r.boekesteijn@maartenskliniek.nl

Background: Inertial sensors hold the promise to objectively measure functional recovery after total knee (TKA) and hip arthroplasty (THA), but their value in addition to patient-reported outcome measures (PROMs) has yet to be demonstrated. This study investigated recovery of gait after TKA and THA using inertial sensors, and compared results to recovery of self-reported scores of pain and function. **Methods:** PROMs and gait parameters were assessed before and at two and fifteen months after TKA (n=24) and THA (n=24). Gait parameters were compared with healthy individuals (n=27) of similar age. Gait data were collected using inertial sensors on the feet, lower back, and trunk. Participants walked for two minutes back and forth over a 6m walkway with 180° turns. PROMs were obtained using the Knee Injury and Osteoarthritis Outcome Scores and Hip Disability and Osteoarthritis Outcome Score. **Results:** Gait parameters recovered to the level of healthy controls after both TKA and THA. Early improvements were found in gait-related trunk kinematics, while spatiotemporal gait parameters mainly improved between two and fifteen months after TKA and THA. Compared to the large and early improvements found in of PROMs, these gait parameters showed a different trajectory, with a marked discordance between the outcome of both methods at two months post-operatively. **Conclusion:** Sensor-derived gait parameters were responsive to TKA and THA, showing different recovery trajectories for spatiotemporal gait parameters and gait-related trunk kinematics. Fifteen months after TKA and THA, there were no remaining gait differences with respect to healthy controls. Given the discordance in recovery trajectories between gait parameters and PROMs, sensor-derived gait parameters seem to carry relevant information for evaluation of physical function that is not captured by self-reported scores.

Objective monitoring of functional recovery after total knee and hip arthroplasty using sensor-derived gait measures

Ramon J. Boekesteijn ^{1,2}, José M.H. Smolders ³, Vincent J.J.F. Busch ³, Noël L.W. Keijsers ¹, Alexander C.H. Geurts ², Katrijn Smulders ¹

¹ Department of Research, Sint Maartenskliniek, Nijmegen, The Netherlands.

² Department of Rehabilitation, Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Center, Nijmegen, The Netherlands.

³ Department of Orthopedic Surgery, Sint Maartenskliniek, Nijmegen, The Netherlands.

Corresponding Author:

Ramon J. Boekesteijn ^{1,2}

Hengstdal 3, Nijmegen, 6500 GM, The Netherlands

Email address: r.boekesteijn@maartenskliniek.nl

Abstract

Background: Inertial sensors hold the promise to objectively measure functional recovery after total knee (TKA) and hip arthroplasty (THA), but their value in addition to patient-reported outcome measures (PROMs) has yet to be demonstrated. This study investigated recovery of gait after TKA and THA using inertial sensors, and compared results to recovery of self-reported scores of pain and function.

Methods: PROMs and gait parameters were assessed before and at two and fifteen months after TKA (n=24) and THA (n=24). Gait parameters were compared with healthy individuals (n=27) of similar age. Gait data were collected using inertial sensors on the feet, lower back, and trunk. Participants walked for two minutes back and forth over a 6m walkway with 180° turns. PROMs were obtained using the Knee Injury and Osteoarthritis Outcome Scores and Hip Disability and Osteoarthritis Outcome Score.

Results: Gait parameters recovered to the level of healthy controls after both TKA and THA. Early improvements were found in gait-related trunk kinematics, while spatiotemporal gait parameters mainly improved between two and fifteen months after TKA and THA. Compared to the large and early improvements found in of PROMs, these gait parameters showed a different trajectory, with a marked discordance between the outcome of both methods at two months post-operatively.

Conclusion: Sensor-derived gait parameters were responsive to TKA and THA, showing different recovery trajectories for spatiotemporal gait parameters and gait-related trunk kinematics. Fifteen months after TKA and THA, there were no remaining gait differences with respect to healthy controls. Given the discordance in recovery trajectories between gait parameters and PROMs, sensor-derived gait parameters seem to carry relevant information for evaluation of physical function that is not captured by self-reported scores.

Introduction

Walking is essential for many activities of daily living, and a good walking capacity is key for participation in society. Previous reports have identified walking speed as ‘sixth vital sign’, given its correlation with essential health parameters, including quality of life (1), risk of future hospitalization (2), and mortality (3). In individuals with end-stage osteoarthritis (OA) of the knee and hip, walking capacity is reduced (4), thereby leading to decreased physical functioning and a lower quality of life (5). As final step in the treatment of severe knee and hip OA, total joint arthroplasty can be performed in order to resolve OA-related symptoms (e.g. pain, stiffness, instability) and improve physical functioning.

Although total knee arthroplasty (TKA) and total hip arthroplasty (THA) are very successful and cost-effective procedures (6), a subset of patients is dissatisfied with treatment outcome (7-9). In addition to patients with identified complications, this includes patients who had an uneventful procedure, but did not achieve their expected level of functional recovery (7). Early identification of individuals at-risk of limited functional recovery is crucial in order to enable clinicians to intervene timely, and may help to readjust patient expectations (10). However, it has been challenging to identify these patients. In part, this is due to a lack of outcomes of physical functioning with good psychometric properties (11). Current diagnostics (e.g. radiographs, physical exam, self-reported outcomes) are limited to static or non-weightbearing situations, or are not necessarily reflective of someone’s actual performance during daily life activities (12, 13). Moreover, patient-reported outcomes (PROMs) are inherently subjective, largely influenced by pain, and suffer from early ceiling effects (14). Although PROMs often contain subscales related to limitations in activities of daily life, such as KOOS/HOOS-ADL or WOMAC function score, these outcomes seem to be more reliant on a patients’ own reflections on their capacity rather than their actual performance (13). Hence, there is a need for objective data that can bridge this gap in clinical assessment.

As an alternative to these subjective scores, performance-based tests have been proposed to objectively capture physical function. For example, evaluation of sit-to-stand transfers, walking short distances, and stair negotiation has been endorsed by the OARSI as core-activities for individuals with knee and hip OA (15). While these tests are well-suited to quickly obtain a global picture of a patient’s physical function, they are limited to a single outcome measure, being the time to perform the task or activity, completed distance, or number of repetitions. These tests provide no information about compensations or underlying biomechanics relevant to the performance, and thus may lack important details. Wearable, inertial sensors, are promising tools to instrument performance-based tests in order to obtain more detailed insights into physical functioning. These inertial sensors are easy to use, have been proven to be valid and reliable (16), do not require lengthy procedures or specialized laboratories, and can be used in clinical settings or even remotely in the home environment (17). Not surprisingly, inertial sensors have gained interest

over the past few years to objectively monitor changes in physical function after total knee and hip arthroplasty (18, 19). In particular, the focus has been on studying gait recovery (18, 19), potentially due to the fact that gait parameters are predictive of limitations in other activities of daily living (20) and gait improvements are an important goal for patients after TKA and THA (21). In the same settings, turning could also be evaluated (22), which has been suggested to be even more sensitive to sensorimotor impairments than straight ahead gait (23). However, before such technologies can be clinically adopted, it is important that the derived outcome measures fulfill the following requirements: they must 1) be sensitive to pre-operative impairment, 2) be responsive to interventions aimed at improving mobility, and 3) provide clinically relevant information about physical functioning.

Multiple gait and turning parameters derived from inertial sensors have shown to be sensitive to mobility impairment in end-stage knee and hip OA (22). The next step herein is to evaluate responsiveness of these parameters to unilateral TKA and THA, and to assess whether post-operative function recovers to the level of healthy individuals. While recovery of gait has previously been investigated using inertial sensors at different timepoints after TKA (13, 17, 24-28) and THA (12, 29-31), a comprehensive study is lacking that maps the recovery trajectory – including turning capacity – at multiple timepoints matching routine follow-up after TKA and THA. In addition, there is a lack of clarity whether gait can be assumed to be ‘normal’ one year after joint replacement (32-34). Finally, little is known about how gait recovery compares to self-reported recovery of physical function (e.g. PROMs). Therefore, the aims of this study were threefold: 1) to investigate gait recovery at two and fifteen months after TKA and THA using inertial sensors, 2) to compare gait 15 months after TKA and THA with data from healthy participants, and 3) to compare recovery trajectories between objective gait parameters and self-reported scores physical functioning.

Materials & Methods

Participants

Individuals with end-stage OA scheduled for TKA (n=24) or THA (n=24) at the Sint Maartenskliniek participated in this study. A group of healthy controls (HC; n=27) within the same age range of 50 to 75 years old was recruited from the community for reference purposes. Healthy participants had no pain in the lower extremities, nor were they familiar with a clinical diagnosis of knee or hip OA. All participants had to be able to walk for more than two minutes without the use of any assistive device. Exclusion criteria were: 1) joint replacement within a year following surgery (including revisions), or symptomatic OA in another weight-bearing joint than the joint scheduled for surgery, 2) BMI > 40 kg/m², and 3) any other musculoskeletal or neurological impairment interfering with gait or balance. Participants who received any other joint replacement to the lower extremities, or had a revision surgery within the period of fifteen months follow-up, were labeled as lost to follow-up. In these cases, data that had been collected until the time of the

second surgery was still used for analysis. Written informed consent was obtained from all participants prior to testing. This study was exempt from ethical review by the CMO Arnhem/Nijmegen (2018-4452) as it was not subject to the Medical Research Involving Human Subjects Act (WMO). All study procedures were conducted in accordance with the Declaration of Helsinki.

Power calculation

Sample sizes were based on the smallest difference that we aimed to detect in this study, which was the difference in gait parameters between individuals 15 months after arthroplasty and HC. Effect sizes for this comparison were informed by studies from Senden *et al.* (25) and Kluge *et al.* (27). When using a standardized mean difference for stride length of 1.1, a power of 80%, and a significance level of 0.05, 22 participants were required per group. To account for potential drop-outs, 24 individuals were recruited for each study group.

Surgical procedure

TKA was performed using the medial parapatellar approach. All individuals scheduled for TKA received the Genesis II posterior stabilized knee prosthesis (Smith & Nephew, Memphis, TN). The patella was resurfaced in 58% of the patients. THA was performed using the posterolateral approach. Specific types of hip implants differed among individuals scheduled for THA and are listed in Supplementary file 1. In total, TKA was performed by seven different surgeons in this study, whereas THA was performed by ten different surgeons. All patients followed an enhanced recovery protocol with mobilization on the day of surgery and hospital discharge within two days. All patients were referred to out-of-hospital physical therapy, which was focused on optimizing functionality, mobility, muscle power, coordination, stability, and walking improvement. Although physical therapy protocols were not standardized, patients usually continued physical therapy for 6-12 months, until their functional goals had been reached.

Demographic and clinical assessment

Severity of radiological OA was determined using Kellgren and Lawrence (KL) grades (35) as scored by JS and VB. Baseline anthropometric characteristics (e.g. body mass, height, and BMI) were obtained during the pre-operative screening visit. In addition, PROMs were assessed using the Knee Injury and Osteoarthritis Outcomes Score (KOOS) for TKA (36) and Hip Disability Osteoarthritis Outcome Score (HOOS) (37) for THA patients. More specifically, HOOS and KOOS subscales “Pain” and “Activities of Daily Living (ADL)” were used to represent pain and physical function. PROMs and gait were assessed pre-operatively – on the same day as the pre-operative screening visit – and at two and fifteen months follow-up. Follow-up measurements were initially set to take place at one year, but measurements were delayed with three months due to the COVID-19 pandemic. Timepoints of follow-up were chosen to match routine follow-up after TKA and THA in the Netherlands, and roughly reflect the moments when patients can walk

independently without an assistive device (e.g. 2 months) and when full recovery has been achieved (e.g. 1 year). For HC, gait was investigated at only one occasion.

Gait protocol

Experimental procedures of the gait assessments were similar to the methods described in Boekesteijn *et al.* (22). Four inertial sensors (*Opal V2, APDM Inc., Portland, OR*) were attached to the dorsum of both feet, the waist (sacrolumbar level), and the sternum. Participants walked back and forth along a six meter trajectory making 180° turns for a total duration of two minutes (Figure 1). Gait tests were performed at comfortable, self-selected speed.

Data analysis

Raw inertial data was processed using validated Mobility Lab v2 software (38). Turning steps were separated from straight walking based on the gyroscope data of the lumbar sensor (39). Gait parameters were calculated for each stride during steady-state walking phases, excluding the two steps preceding and following a turn. Parameters were summarized as mean value of all valid strides or turns. Based on non-redundancy and size of the difference between individuals with end-stage knee and hip OA and HC as found previously (22), the following outcomes were extracted (Figure 1): 1) gait speed, 2) stride length, 3) cadence 4), step time asymmetry, 5) stride time variability, 6) peak turning velocity, 7) lumbar sagittal range of motion, 8) lumbar coronal range of motion, and 9) trunk coronal range of motion. Parameters were only evaluated for the TKA or THA group in case they were previously found to be sensitive to mobility impairment in knee or hip OA (22). For this reason, step time asymmetry, lumbar sagittal range of motion, and lumbar coronal range of motion were not evaluated in the TKA group.

Statistical analysis

Recovery trajectories of gait parameters and KOOS/ HOOS scores were visualized on group level by the mean and 95% confidence intervals (CI). Linear mixed models with gait parameters and KOOS or HOOS scores as dependent variable, time as two independent dummy variables (e.g. T2 and T15), and subject ID as random effect factor were constructed to investigate the effect of time on gait and KOOS/HOOS scores for TKA and THA separately. Addition of random slopes was evaluated, but these were not included in the final model for reasons of parsimony, as this did not contribute to a better model fit. Gait parameters of TKA and THA groups were compared with HC at 15 month follow-up using an independent samples t-test or non-parametric Mann-Whitney U test in case data was not normally distributed. Inferences of statistical significance were based on $p < 0.05$. Since multiple outcome parameters were used for the same construct (e.g. gait) we controlled the family-wise error rate using the Hommel procedure (40), by adjusting the p-values for the number of gait parameters involved in each comparison. To assess discrepancies between gait and self-reported scores of physical function, we compared trajectories between gait speed, which was found to be most sensitive to gait impairment in knee and hip OA (22), and KOOS/HOOS-ADL scores. Meaningful improvements were defined as a change in gait speed

>0.10 m/s (41) and a change in KOOS/HOOS ADL score >20 points (42). Data were processed in Python 3.8.3 and statistical analyses were conducted in RStudio 3.6.1 using the lme4 package (version 1.1-26) (43).

Results

Participant characteristics

The study groups did not differ significantly in age, sex, height, or BMI (Table 1). Compared to HC, body mass was significantly higher in individuals scheduled for TKA and THA. All individuals scheduled for TKA or THA had moderate to severe OA (KL grades 3 or 4). In total we had missing data for eleven participants. Three participants had a complication within the study window. For details regarding missing data and complications, see Supplementary file 2.

Recovery of gait after arthroplasty

Two months after surgery, gait speed, stride length, and cadence were not significantly different from baseline, both after TKA and THA (Table 2; Figure 2A-C). Peak turning velocity improved with 19.1 deg/s (95% CI: 6.9, 31.5) in the first two months after THA, but not after TKA (Table 2). There were no changes in step time asymmetry within the first two months after THA (Table 2), nor were there changes in stride time variability after TKA and THA at this timepoint (Table 2). As for kinematics of the trunk, trunk coronal RoM was slightly lower two months after TKA (mean diff: -1.0 deg, 95% CI: -1.6, -0.3) compared to pre-operatively, whereas lumbar sagittal RoM was lower two months after THA (mean diff: -1.9 deg, 95% CI: -3.0, -0.8) (Table 2).

Between two and fifteen months, large improvements in gait speed, cadence, and stride length were observed after both TKA and THA (Table 2; Figure 2A-C). For gait speed, the gain between two and fifteen months was 0.22 m/s (95% CI: 0.15, 0.29) after TKA and 0.14 m/s (95% CI: 0.06, 0.20) after THA. Peak turning velocity did not change significantly (mean diff: 17.4 deg/s, 95% CI: 1.7, 33.0, $p_{\text{adj}} = 0.105$) between two and fifteen months after TKA. There were no significant improvements in turning velocity between two and fifteen months after THA (Table 2). Step time asymmetry did not change between two and fifteen months after THA. There were no changes in stride time variability, or trunk coronal RoM between two and fifteen months after TKA and THA (Table 2). Individuals after THA showed an increase of 1.4 degrees (95% CI: 0.6, 2.1) in lumbar coronal RoM between two and fifteen months. Finally, none of the gait parameters were significantly different from HC at fifteen months after TKA and THA (Table 3; Figure 2A-I).

Changes on PROMs after arthroplasty

Two months after TKA, individuals improved on all KOOS subscales, except for ‘Symptoms’ (Table 4). For all other subscales, self-reported scores showed large improvements (> 20 points) with some individuals already reaching (sub)maximal scores (≥ 90 points) within the first two

months (Figure 2J & 2K). Further improvements were found for all KOOS subscales from two to fifteen months follow-up (Table 4). As for the HOOS, all subscales improved from baseline to two months after THA, as well as from two to fifteen months follow-up, with the largest magnitude of effects taking place in the first two months (Table 4).

Relation between recovery trajectories of gait parameters and PROMs

When comparing recovery trajectories of self-reported scores with gait parameters, substantial differences were observed (Figure 2). Where KOOS and HOOS scores showed large improvements over almost all subscales in the first two months after surgery (Table 4), gait parameters generally improved between 2 and 15 months, with the exception of trunk-related gait parameters. More specifically, discrepancies between HOOS/KOOS-ADL scores and spatiotemporal parameters were present at two months after surgery. For gait speed specifically, there were no significant changes between baseline and two months after TKA and THA, while HOOS/KOOS-ADL improved with 42 points and 21 points, respectively. To illustrate, two months after surgery, 10/23 individuals after TKA reported meaningful improvements in ADL scores, while merely 4/23 showed a meaningful improvement in gait speed. Similarly, after THA, 20/23 individuals reported meaningful improvements in ADL scores at 2 months, with 10/23 individuals showing meaningful improvements in gait speed.

Discussion

This study evaluated the use of inertial sensors to monitor functional recovery after TKA and THA. In concordance with our previous work, that sensor-derived gait parameters show sensitive to knee and hip OA (22), this study showed that these parameters were also responsive to TKA and THA at two and fifteen months after surgery, and recovered to the same level as HC fifteen months after surgery. In addition, discrepancies between recovery trajectories of spatiotemporal gait parameters and HOOS/KOOS scores were observed, particularly at two months post-operatively.

Recovery trajectory of gait after TKA and THA

There were limited improvements in spatiotemporal gait parameters two months after TKA and THA, which is in agreement with previous studies (25, 33). However, the observed faster turning in absence of higher gait speed two months after THA is interesting, and may suggest that turning is more sensitive to short-term improvements in physical function after THA than gait speed. In contrast to these basic spatiotemporal parameters, normalization of trunk movement was found already two months after TKA and THA. Pre-operatively, individuals with knee OA may increase lateral trunk lean as a strategy to reduce knee joint loading and/or pain (44-46), which is no longer required two months after TKA. Increased lumbar RoM in the sagittal plane, in its turn, may serve as pre-operative compensation for individuals with hip OA to overcome pain and hip joint stiffness (47, 48). Taken together, these results suggest that while two months is too early for meaningful

recovery of spatiotemporal gait parameters, pre-operative compensations of the trunk and pelvis already disappear within the first two months after TKA and THA.

Large and clinically relevant improvements were observed on spatiotemporal parameters between two and fifteen months after TKA and THA. This is in agreement with literature investigating gait with inertial sensors one year after TKA (13, 24, 27) and THA (12, 31). Recovery of muscle strength (e.g. quadriceps and hip abductors) – which coincides with this period (49, 50) – may underly these improvements in walking capacity. As for trunk kinematics, both individuals after TKA and THA showed an increase in lumbar coronal RoM from two to fifteen months after surgery, which may relate to the restored ability of the hip abductors to control frontal plane pelvic movement (12, 51). Compensations like lateral trunk lean, which limit pelvic RoM, are then longer required (52). When combining these results with those of gait recovery at two months, it can thus be concluded that a wide range of sensor-derived gait metrics is responsive to TKA and THA, with spatiotemporal parameters and trunk kinematics each showing a distinctive recovery trajectory.

None of the gait parameters were different from HC mean values at fifteen months after TKA and THA. This in contrast with some earlier studies reporting remaining gait differences between HC and individuals one year after TKA (27, 32, 53) or THA (33). Although one year after arthroplasty is generally considered as endpoint of recovery, these differences between studies might be attributed to the longer follow-up time in our study. This seems like a reasonable explanation given that improvements in gait were larger in our study compared to these earlier studies (27, 32, 33). Our findings underscore the success of TKA and THA in improving physical functioning, and indicate that normal spatiotemporal gait parameters and normal trunk kinematics may be achieved 15 months after TKA and THA. Whether other aspects of gait, including lower-extremity kinematics and kinetics, also recover to the level of healthy controls remains to be elucidated. Despite our findings of full recovery after TKA and THA, current literature suggest that more advanced parameters, including lower extremity kinematics and kinetics, may still reveal deficits in gait one year after surgery (32, 33, 53).

Relationship between PROMs and objective gait measures

Objective gait parameters showed a different recovery trajectory than subjective reports of physical function and pain. Scores on the KOOS and HOOS greatly improved within the first two months, while spatiotemporal gait parameters mainly improved between two and fifteen months after surgery. Similar discrepancies between PROMs, gait, and performance-based tests have previously been recognized in the literature (12, 14, 32, 54-56). For example, inverse recovery trajectories (i.e. early improvements in PROMs compared to worsening of performance-based outcomes) have been observed between KOOS/HOOS ADL scores and performance-based outcomes, including the 6 minute walk test, stair climbing test, and timed up and go test, during the first month of recovery after TKA and THA (14, 54-56). For sensor-derived gait parameters specifically, poor agreement with PROM scores has been found after TKA and THA (12, 24). On

a similar note, Fransen *et al.* found that, although perceived walking ability and self-reported physical function improved, there were no improvements in quality or quantity of daily life gait three months after surgery (17). The current study adds that the discordance between gait parameters and self-reported function scores is most prominent at two months after surgery, with the exception of parameters related to trunk motion. The general consensus is that physical function subscales of PROMs assess a different domain than performance-based tests and gait analysis (13). This discrepancy may first be related to a strong relation of physical function subscales with pain (14), as was also apparent from the similarity between the recovery trajectories of HOOS/KOOS Pain and ADL subscales in our study. One potential explanation for this is that improvements in pain directly translate to a more positive reflection on daily life performance, and that patients considered pain as the main limiting factor in their daily life activities. Second, these self-reported scores ask about experienced difficulty during a wide range of activities, rather than how they execute a specific activity, which is inherently different from what these gait parameters measure. Finally, there is evidence that objective parameters of physical function are more sensitive to remaining functional deficits after TKA than PROMs (32), which may attributed to early ceiling effects of PROMs. Since improving mobility – specifically walking – is an important goal of joint replacement (57), these sensor-derived parameters may thus add a relevant dimension to evaluation of physical functioning, although their clinical value still has to be demonstrated.

Limitations and future directions

This study has a number of limitations which merit attention. First, we measured gait recovery in a well-defined cohort of patients with unilateral osteoarthritis without pain complaints in any other joint or previous joint replacement. While this was relevant for the aims of the current study, this limits the generalizability of our findings. Second, in the present study, evaluation of physical function was limited to gait and turning in the present study while other daily life activities, including sit-to-stand transfers and stair climbing, are also relevant for physical functioning after TKA and THA (15). Third, gait parameters in this study were limited to spatiotemporal parameters and gait-related trunk kinematics. Other parameters, such as knee and hip kinematics that can be derived from a different set-up of inertial sensors may provide additional information about gait recovery after TKA and THA, especially in light of remaining gait deficits (33). While the current study touches upon the potential value of objective measurement of physical function, the actual value of clinical implementation of gait tests cannot be derived from our study results. Future studies with larger samples and a more diverse population are required to investigate the applicability of objective gait assessment systems to identify poor-responders. Another valuable direction would be to explore whether such data can be used to adjust patient expectations during clinical visits and to further tailor post-operative care. Finally, there is a need for studies employing inertial sensors for remote monitoring during daily life, which may not only enable more efficient (digital) healthcare pathways in the future, but may also contribute to data with greater ecological validity (58, 59).

Conclusions

This study showed that objective gait measures derived from inertial sensors are responsive to TKA and THA. Not only speed-related parameters, but also turning and trunk motion provide important information about functional status before and at two and fifteen months after joint replacement. There were no remaining gait differences between individuals after TKA or THA and healthy participants at fifteen months. Recovery trajectories of objective gait data were different from those of KOOS and HOOS ADL subscales, with a marked discordance at two months after surgery. Altogether, these results strengthen the premise that sensor-derived gait metrics may provide meaningful information about recovery of physical functioning after TKA and THA that is not captured by self-reported ADL or pain scores.

References

- Schmid A, Duncan Pamela W, Studenski S, Lai Sue M, Richards L, Perera S, et al. Improvements in Speed-Based Gait Classifications Are Meaningful. *Stroke*. 2007;38(7):2096-100.
- Montero-Odasso M, Schapira M, Soriano ER, Varela M, Kaplan R, Camera LA, et al. Gait Velocity as a Single Predictor of Adverse Events in Healthy Seniors Aged 75 Years and Older. *The Journals of Gerontology: Series A*. 2005;60(10):1304-9.
- Hardy SE, Perera S, Roumani YF, Chandler JM, Studenski SA. Improvement in Usual Gait Speed Predicts Better Survival in Older Adults. *Journal of the American Geriatrics Society*. 2007;55(11):1727-34.
- Thomas SG, Pagura SMC, Kennedy D. Physical Activity and its Relationship to Physical Performance in Patients With End Stage Knee Osteoarthritis. *Journal of Orthopaedic & Sports Physical Therapy*. 2003;33(12):745-54.
- Neogi T. The epidemiology and impact of pain in osteoarthritis. *Osteoarthritis and Cartilage*. 2013;21(9):1145-53.
- Ethgen O, Bruyère O, Richy F, Dardennes C, Reginster J-Y. Health-Related Quality of Life in Total Hip and Total Knee Arthroplasty: A Qualitative and Systematic Review of the Literature. *JBJS*. 2004;86(5).
- Gunaratne R, Pratt DN, Banda J, Fick DP, Khan RJK, Robertson BW. Patient Dissatisfaction Following Total Knee Arthroplasty: A Systematic Review of the Literature. *The Journal of Arthroplasty*. 2017;32(12):3854-60.
- Anakwe RE, Jenkins PJ, Moran M. Predicting Dissatisfaction After Total Hip Arthroplasty: A Study of 850 Patients. *The Journal of Arthroplasty*. 2011;26(2):209-13.
- Nilsdotter AK, Toksvig-Larsen S, Roos EM. Knee arthroplasty: are patients' expectations fulfilled? *Acta Orthopaedica*. 2009;80(1):55-61.
- Tolk JJ, Janssen RPA, Haanstra TM, van der Steen MC, Bierma-Zeinstra SMA, Reijman M. The influence of expectation modification in knee arthroplasty on satisfaction of patients: a randomized controlled trial. *The Bone & Joint Journal*. 2021;103-B(4):619-26.
- Hossain FS, Konan S, Patel S, Rodriguez-Merchan EC, Haddad FS. The assessment of outcome after total knee arthroplasty. *The Bone & Joint Journal*. 2015;97-B(1):3-9.

12. Bolink SAAN, Lenguerrand E, Brunton LR, Wylde V, Gooberman-Hill R, Heyligers IC, et al. Assessment of physical function following total hip arthroplasty: Inertial sensor based gait analysis is supplementary to patient-reported outcome measures. *Clinical Biomechanics*. 2016;32:171-9.
13. Fransen BL, Mathijssen NMC, Slot K, de Esch NHH, Verburg H, Temmerman OPP, et al. Gait quality assessed by trunk accelerometry after total knee arthroplasty and its association with patient related outcome measures. *Clinical Biomechanics*. 2019;70:192-6.
14. Stevens-Lapsley JE, Schenkman ML, Dayton MR. Comparison of Self-Reported Knee Injury and Osteoarthritis Outcome Score to Performance Measures in Patients After Total Knee Arthroplasty. *PM&R*. 2011;3(6):541-9.
15. Dobson F, Hinman RS, Roos EM, Abbott JH, Stratford P, Davis AM, et al. OARSI recommended performance-based tests to assess physical function in people diagnosed with hip or knee osteoarthritis. *Osteoarthritis and Cartilage*. 2013;21(8):1042-52.
16. Kobsar D, Charlton JM, Tse CTF, Esculier J-F, Graffos A, Krowchuk NM, et al. Validity and reliability of wearable inertial sensors in healthy adult walking: a systematic review and meta-analysis. *Journal of NeuroEngineering and Rehabilitation*. 2020;17(1):62.
17. Fransen BL, Pijnappels M, Butter IK, Burger BJ, van Dieën JH, Hoozemans MJM. Patients' perceived walking abilities, daily-life gait behavior and gait quality before and 3 months after total knee arthroplasty. *Archives of Orthopaedic and Trauma Surgery*. 2021.
18. Small SR, Bullock GS, Khalid S, Barker K, Trivella M, Price AJ. Current clinical utilisation of wearable motion sensors for the assessment of outcome following knee arthroplasty: a scoping review. *BMJ Open*. 2019;9(12):e033832.
19. Kobsar D, Masood Z, Khan H, Khalil N, Kiwan MY, Ridd S, et al. Wearable Inertial Sensors for Gait Analysis in Adults with Osteoarthritis—A Scoping Review. *Sensors*. 2020;20(24).
20. Potter JM, Evans AL, Duncan G. Gait speed and activities of daily living function in geriatric patients. *Archives of Physical Medicine and Rehabilitation*. 1995;76(11):997-9.
21. Scott CEH, Bugler KE, Clement ND, MacDonald D, Howie CR, Biant LC. Patient expectations of arthroplasty of the hip and knee. *The Journal of Bone and Joint Surgery British volume*. 2012;94-B(7):974-81.
22. Boekesteijn RJ, Smolders JMH, Busch VJJF, Geurts ACH, Smulders K. Independent and sensitive gait parameters for objective evaluation in knee and hip osteoarthritis using wearable sensors. *BMC Musculoskeletal Disorders*. 2021;22(1):242.
23. Mancini M, Schlueter H, El-Gohary M, Mattek N, Duncan C, Kaye J, et al. Continuous Monitoring of Turning Mobility and Its Association to Falls and Cognitive Function: A Pilot Study. *The Journals of Gerontology: Series A*. 2016;71(8):1102-8.
24. Bolink SAAN, Grimm B, Heyligers IC. Patient-reported outcome measures versus inertial performance-based outcome measures: A prospective study in patients undergoing primary total knee arthroplasty. *The Knee*. 2015;22(6):618-23.
25. Senden R, Grimm B, Meijer K, Savelberg H, Heyligers IC. The importance to including objective functional outcomes in the clinical follow up of total knee arthroplasty patients. *The Knee*. 2011;18(5):306-11.
26. Jolles BM, Grzesiak A, Eudier A, Dejnabadi H, Voracek C, Pichonnaz C, et al. A randomised controlled clinical trial and gait analysis of fixed- and mobile-bearing total knee replacements with a five-year follow-up. *The Journal of Bone and Joint Surgery British volume*. 2012;94-B(5):648-55.
27. Kluge F, Hannink J, Pasluosta C, Klucken J, Gaßner H, Gelse K, et al. Pre-operative sensor-based gait parameters predict functional outcome after total knee arthroplasty. *Gait & Posture*. 2018;66:194-200.

28. Youn I-H, Leutzinger T, Youn J-H, Zeni JA, Knarr BA. Self-Reported and Performance-Based Outcome Measures Estimation Using Wearables After Unilateral Total Knee Arthroplasty. *Frontiers in Sports and Active Living*. 2020;2(128).
29. Reininga IHF, Stevens M, Wagenmakers R, Boerboom AL, Groothoff JW, Bulstra SK, et al. Comparison of gait in patients following a computer-navigated minimally invasive anterior approach and a conventional posterolateral approach for total hip arthroplasty: A randomized controlled trial. *Journal of Orthopaedic Research*. 2013;31(2):288-94.
30. Nelms NJ, Birch CE, Halsey DH, Blankstein M, McGinnis RS, Beynon BD. Assessment of Early Gait Recovery After Anterior Approach Compared to Posterior Approach Total Hip Arthroplasty: A Smartphone Accelerometer-Based Study. *The Journal of Arthroplasty*. 2020;35(2):465-70.
31. Wada O, Asai T, Hiyama Y, Nitta S, Mizuno K. Gait Variability in Women With Hip Osteoarthritis Before and After Total Hip Replacement: A Prospective Cohort Study. *American Journal of Physical Medicine & Rehabilitation*. 2019;98(10).
32. Naili JE, Iversen MD, Esbjörnsson A-C, Hedström M, Schwartz MH, Häger CK, et al. Deficits in functional performance and gait one year after total knee arthroplasty despite improved self-reported function. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2017;25(11):3378-86.
33. Bahl JS, Nelson MJ, Taylor M, Solomon LB, Arnold JB, Thewlis D. Biomechanical changes and recovery of gait function after total hip arthroplasty for osteoarthritis: a systematic review and meta-analysis. *Osteoarthritis and Cartilage*. 2018;26(7):847-63.
34. Milner CE. Is gait normal after total knee arthroplasty? Systematic review of the literature. *Journal of Orthopaedic Science*. 2009;14(1):114-20.
35. Kellgren JH, Lawrence JS. Radiological Assessment of Osteo-Arthrosis. *Annals of the Rheumatic Diseases*. 1957;16(4):494.
36. de Groot IB, Favejee MM, Reijman M, Verhaar JAN, Terwee CB. The Dutch version of the knee injury and osteoarthritis outcome score: A validation study. *Health and Quality of Life Outcomes*. 2008;6(1):16.
37. de Groot IB, Reijman M, Terwee CB, Bierma-Zeinstra SMA, Favejee M, Roos EM, et al. Validation of the Dutch version of the Hip disability and Osteoarthritis Outcome Score. *Osteoarthritis and Cartilage*. 2007;15(1):104-9.
38. Morris R, Stuart S, McBarron G, Fino PC, Mancini M, Curtze C. Validity of Mobility Lab (version 2) for gait assessment in young adults, older adults and Parkinson's disease. *Physiological Measurement*. 2019;40(9):095003.
39. El-Gohary M, Pearson S, McNamara J, Mancini M, Horak F, Mellone S, et al. Continuous Monitoring of Turning in Patients with Movement Disability. *Sensors*. 2014;14(1).
40. Hommel G. A stagewise rejective multiple test procedure based on a modified Bonferroni test. *Biometrika*. 1988;75(2):383-6.
41. Bohannon RW, Glenney SS. Minimal clinically important difference for change in comfortable gait speed of adults with pathology: a systematic review. *Journal of Evaluation in Clinical Practice*. 2014;20(4):295-300.
42. Lyman S, Lee Y-Y, McLawhorn AS, Islam W, MacLean CH. What Are the Minimal and Substantial Improvements in the HOOS and KOOS and JR Versions After Total Joint Replacement? *Clinical Orthopaedics and Related Research*. 2018;476(12).
43. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*; Vol 1, Issue 1 (2015). 2015.
44. Mündermann A, Dyrby CO, Andriacchi TP. Secondary gait changes in patients with medial compartment knee osteoarthritis: Increased load at the ankle, knee, and hip during walking. *Arthritis & Rheumatism*. 2005;52(9):2835-44.

45. Hunt MA, Birmingham TB, Bryant D, Jones I, Giffin JR, Jenkyn TR, et al. Lateral trunk lean explains variation in dynamic knee joint load in patients with medial compartment knee osteoarthritis. *Osteoarthritis and Cartilage*. 2008;16(5):591-9.
46. Linley HS, Sled EA, Culham EG, Deluzio KJ. A biomechanical analysis of trunk and pelvis motion during gait in subjects with knee osteoarthritis compared to control subjects. *Clinical Biomechanics*. 2010;25(10):1003-10.
47. Hurwitz DE, Hulet CH, Andriacchi TP, Rosenberg AG, Galante JO. Gait compensations in patients with osteoarthritis of the hip and their relationship to pain and passive hip motion. *Journal of Orthopaedic Research*. 1997;15(4):629-35.
48. Lenaerts G, Mulier M, Spaepen A, Van der Perre G, Jonkers I. Aberrant pelvis and hip kinematics impair hip loading before and after total hip replacement. *Gait & Posture*. 2009;30(3):296-302.
49. Mizner RL, Petterson SC, Snyder-Mackler L. Quadriceps Strength and the Time Course of Functional Recovery After Total Knee Arthroplasty. *Journal of Orthopaedic & Sports Physical Therapy*. 2005;35(7):424-36.
50. Ismailidis P, Kvarda P, Vach W, Cadosch D, Appenzeller-Herzog C, Mündermann A. Abductor Muscle Strength Deficit in Patients After Total Hip Arthroplasty: A Systematic Review and Meta-Analysis. *The Journal of Arthroplasty*. 2021.
51. Reininga IHF, Stevens M, Wagenmakers R, Bulstra SK, Groothoff JW, Zijlstra W. Subjects with hip osteoarthritis show distinctive patterns of trunk movements during gait-a body-fixed-sensor based analysis. *Journal of NeuroEngineering and Rehabilitation*. 2012;9(1):3.
52. Bolink SAAN, Brunton LR, van Laarhoven S, Lipperts M, Heyligers IC, Blom AW, et al. Frontal Plane Pelvic Motion during Gait Captures Hip Osteoarthritis Related Disability. *HIP International*. 2015;25(5):413-9.
53. Outerleys JB, Dunbar MJ, Richardson G, Hubley-Kozey CL, Astephen Wilson JL. Quantifying Achievable Levels of Improvement in Knee Joint Biomechanics During Gait After Total Knee Arthroplasty Relative to Osteoarthritis Severity. *Journal of Applied Biomechanics*. 2021;37(2):130-8.
54. Dayton MR, Judd DL, Hogan CA, Stevens-Lapsley JE. Performance-Based Versus Self-Reported Outcomes Using the Hip Disability and Osteoarthritis Outcome Score After Total Hip Arthroplasty. *American journal of physical medicine & rehabilitation*. 2016;95(2):132-8.
55. Luna IE, Kehlet H, Peterson B, Wede HR, Høevsgaard SJ, Aasvang EK. Early patient-reported outcomes versus objective function after total hip and knee arthroplasty. *The Bone & Joint Journal*. 2017;99-B(9):1167-75.
56. Mizner RL, Petterson SC, Clements KE, Zeni JA, Irrgang JJ, Snyder-Mackler L. Measuring Functional Improvement After Total Knee Arthroplasty Requires Both Performance-Based and Patient-Report Assessments: A Longitudinal Analysis of Outcomes. *The Journal of Arthroplasty*. 2011;26(5):728-37.
57. Lange T, Schmitt J, Kopkow C, Rataj E, Günther K-P, Lützner J. What Do Patients Expect From Total Knee Arthroplasty? A Delphi Consensus Study on Patient Treatment Goals. *The Journal of Arthroplasty*. 2017;32(7):2093-9.e1.
58. Van Ancum JM, van Schooten KS, Jonkman NH, Huijben B, van Lummel RC, Meskers CGM, et al. Gait speed assessed by a 4-m walk test is not representative of daily-life gait speed in community-dwelling adults. *Maturitas*. 2019;121:28-34.
59. Takayanagi N, Sudo M, Yamashiro Y, Lee S, Kobayashi Y, Niki Y, et al. Relationship between Daily and In-laboratory Gait Speed among Healthy Community-dwelling Older Adults. *Scientific Reports*. 2019;9(1):3496.

Table 1(on next page)

Baseline characteristics

Note: TKA = total knee arthroplasty, THA = total hip arthroplasty, HC = healthy controls, BMI = body mass index, KL = Kellgren Lawrence. Data are presented as mean [95% CI].

1

2 **Table 1** : baseline characteristics

	TKA (n=24)	THA (n=24)	HC (n=27)	Main effect	Post-hoc analysis
Age (y)	63 [61, 66]	64 [62, 67]	66 [63, 68]	F(2,72) = 0.81, p = 0.448	
Sex (M:F)	12:12	16:8	13:14	χ^2 (2, N=75) = 2.07, p = 0.355	
Height (m)	1.73 [1.69, 1.77]	1.75 [1.72, 1.79]	1.72 [1.68, 1.75]	F(2,72) = 0.98, p = 0.381	
Body mass (kg)	84.6 [78.6, 90.6]	86.0 [78.1, 94.0]	75.7 [71.5, 80.0]	F(2,72) = 3.66, p = 0.031	TKA vs. HC: t(49) = 2.527 p = 0.015 THA vs. HC: t(49) = 2.428 ; p = 0.019
BMI (kg/m ²)	28.2 [26.6, 29.9]	27.9 [25.6, 30.2]	25.7 [24.5, 26.8]	F(2,72) = 2.91, p = 0.060	
KL score (I:II:III:IV)	0:0:8:16	0:0:6:18	-		

3 *Note:* TKA = total knee arthroplasty, THA = total hip arthroplasty, HC = healthy controls, BMI = body mass index, KL = Kellgren Lawrence. Data are presented as mean [95%
4 CI].

Table 2 (on next page)

Effects of time on gait parameters in the TKA and THA group

Note: TKA = total knee arthroplasty, THA = total hip arthroplasty, RoM = range of motion,

P_{corr} = Hommel adjusted p-value. Data are presented as mean (95% CI).

Table 2: Effects of time on gait parameters in the TKA and THA group.

Gait parameters	TKA (n=24)							THA (n=24)						
	baseline – 2 months				2 months – 15 months			baseline – 2 months				2 months – 15 months		
	Baseline (intercept)	Mean difference (95% CI)	P _{value}	P _{corr}	Mean difference (95% CI)	P _{value}	P _{corr}	Baseline (intercept)	Mean difference (95% CI)	P _{value}	P _{corr}	Mean difference (95% CI)	P _{value}	P _{corr}
Gait speed (m/s)	0.99	-0.04 (-0.10, 0.03)	0.272	0.569	0.22 (0.15, 0.29)	<0.001	<0.001	0.96	0.04 (-0.02, 0.10)	0.245	0.514	0.14 (0.06, 0.20)	<0.001	0.003
Stride Length (m)	1.16	-0.002 (-0.05, 0.05)	0.924	0.924	0.14 (0.09, 0.19)	<0.001	<0.001	1.11	0.05 (0.002, 0.10)	0.049	0.306	0.07 (0.02, 0.13)	0.013	0.076
Cadence (steps/min)	102.8	-3.8 (-6.8, -0.8)	0.016	0.081	10.1 (7.0, 13.2)	<0.001	<0.001	103.0	-1.1 (-3.9, 1.7)	0.453	0.514	6.9 (3.8, 10.1)	<0.001	<0.001
Peak turning velocity (deg/s)	164.0	8.5 (-6.5, 23.7)	0.275	0.759	17.4 (1.7, 33.0)	0.035	0.105	171.2	19.1 (6.9, 31.5)	0.004	0.033	11.1 (-2.5, 24.9)	0.121	0.484
Step time asymmetry (%)	-	-	-	-	-	-	-	4.3	-0.7 (-2.0, 0.6)	0.292	0.514	-1.0 (-2.4, 0.3)	0.155	0.553
Stride time variability (%)	2.3	0.2 (-0.2, 0.6)	0.380	0.569	-0.3 (-0.8, 0.1)	0.117	0.234	2.5	-0.1 (-0.5, 0.3)	0.514	0.514	-0.2 (-0.7, 0.2)	0.277	0.831
Lumbar sagittal RoM (deg)	-	-	-	-	-	-	-	8.1	-1.9 (-3.0, -0.8)	0.001	0.013	0.1 (-1.1, 1.3)	0.870	0.870
Lumbar coronal RoM (deg)	-	-	-	-	-	-	-	5.2	0.4 (-0.3, 1.3)	0.255	0.514	1.4 (0.6, 2.1)	0.001	0.010
Trunk coronal RoM (deg)	7.8	-1.0 (-1.6, -0.3)	0.009	0.049	0.1 (-0.6, 0.9)	0.710	0.710	8.1	-0.6 (-1.3, 0.1)	0.087	0.439	0.1 (-0.6, 0.8)	0.797	0.870

2 Note: TKA = total knee arthroplasty, THA = total hip arthroplasty, RoM = range of motion, P_{corr} = Hommel adjusted p-value.

3 Data are presented as mean (95% CI).

Table 3 (on next page)

Post-operative situation compared to HC

Note: HC = healthy control, TKA = total knee arthroplasty, THA = total hip arthroplasty, RoM = range of motion, P_{corr} = Hommel adjusted p-value. Non-normal distributed data are presented in *italic* and are summarized as median (IQR) with median difference (95% CI). Test statistics represent either the t-value (normal data) or U (non-normal data).

Table 3: Post-operative situation compared to HC

Gait parameters	HC (n=27)	TKA – 15 mo (n = 21)	TKA vs HC				THA – 15 mo (n=18)	THA vs HC			
			Mean diff (95%CI)	Test statistic (df = 46)	P _{TKA-HC}	P _{corr}		Mean diff (95%CI)	Test statistic (df =43)	P _{THA-HC}	P _{corr}
Gait speed (m/s)	1.24 (1.18, 1.31)	1.19 (1.13, 1.24)	-0.06 (-0.15, 0.03)	1.31	0.197	0.569	1.16 (1.10, 1.22)	-0.08 (-0.18, 0.02)	1.70	0.096	0.441
Stride Length (m)	1.32 (1.26, 1.37)	1.30 (1.25, 1.36)	-0.02 (-0.10, 0.07)	0.40	0.691	0.691	1.26 (1.20, 1.31)	-0.06 (-0.15, 0.03)	-1.32	0.194	0.581
Cadence (steps/min)	113 (110, 117)	109 (105, 112)	-4 (-9, 1)	1.66	0.104	0.402	110 (107, 113)	-3 (-8, 2)	-1.13	0.265	0.794
Peak turning velocity (deg/s)	205 (190, 220)	190 (174, 206)	-15 (-38, 8)	1.30	0.201	0.569	207 (190, 223)	1 (-23, 26)	1.40	0.909	0.909
Step time asymmetry (%)	2.7 (1.9, 3.4)	-	-	-	-	-	2.6 (1.9, 3.2)	-0.2 (-1.3, 1.0)	-0.28	0.778	0.909
Stride time variability (%)	1.8 (1.6, 2.0)	2.1 (1.8, 2.4)	0.3 (-0.1, 0.7)	1.68	0.099	0.398	2.1 (1.8, 2.4)	0.3 (-0.03, 0.7)	1.86	0.069	0.397
Lumbar sagittal RoM (deg)	5.3 (4.7, 5.9)	-	-	-	-	-	6.3 (5.7, 6.8)	0.9 (-0.2, 2.1)	1.63	0.111	0.444
Lumbar coronal RoM (deg)	<i>8.1 (6.3, 9.0)</i>	-	-	-	-	-	<i>6.5 (5.3, 7.3)</i>	<i>-1.6 (-2.7, 0.1)</i>	U = 165.5	0.074	0.397
Trunk coronal RoM (deg)	6.6 (5.9, 7.3)	7.1 (6.2, 8.1)	0.5 (-0.7, 1.7)	0.89	0.379	0.691	7.5 (6.6, 8.5)	0.8 (-0.6, 2.1)	1.16	0.254	0.761

2 Note: HC = healthy control, TKA = total knee arthroplasty, THA = total hip arthroplasty, RoM = range of motion, P_{corr} = Hommel adjusted p-value.

3 Non-normal distributed data are presented in *italic* and are summarized as median (IQR) with median difference (95% CI). Test statistics represent either the t-value (normal data)

4 or U (non-normal data).

5

Table 4(on next page)

Patient-reported outcome scores for both groups at each timepoint

Note: OA = osteoarthritis, TKA = total knee arthroplasty, THA = total hip arthroplasty, HC = healthy controls, ADL = activities of daily living.

1 **Table 4:** Patient-reported outcome scores for both groups at each timepoint

PROM scores	TKA (n=24)					THA (n=24)				
	Pre-operative – 2 months			2 months – 15 months		Pre-operative – 2 months			2 months – 15 months	
	Baseline (estimate)	Mean difference (95% CI)	P-value	Mean difference (95%CI)	P-value	Baseline (estimate)	Mean difference (95% CI)	P-value	Mean difference (95%CI)	P-value
HOOS/KOOS										
1) Symptoms	50	5 (-3, 14)	0.210	27 (18, 36)	<0.001	41	37 (30, 44)	<0.001	12 (4, 20)	0.007
2) Pain	42	28 (20, 36)	<0.001	19 (10, 28)	<0.001	39	45 (39, 52)	<0.001	9 (2, 16)	0.017
3) ADL	52	21 (13, 29)	<0.001	18 (10, 26)	<0.001	39	42 (35, 45)	<0.001	12 (4, 20)	0.004
4) Sports/Recreation	16	22 (9, 34)	0.001	30 (17, 43)	<0.001	15	48 (39, 57)	<0.001	13 (3, 22)	0.014
5) Quality of life	26	25 (17, 33)	<0.001	29 (20, 37)	<0.001	24	42 (32, 51)	<0.001	19 (9, 29)	<0.001

2 *Note:* OA = osteoarthritis, TKA = total knee arthroplasty, THA = total hip arthroplasty, HC = healthy controls, ADL = activities of daily living.

3

Figure 1

Overview of the experimental set-up and outcome parameters

Wearable inertial sensors were used to capture gait parameters during a two minute walk test over a six meter walkway with 180 degree turns. The figure is adapted from Boekesteijn *et al.* (19) .




Figure 2

Recovery trajectories of gait parameters and PROMs.

Dots with error bars represent group means with 95% CI, whereas grey areas display HC group means with 95% CI. Individual datapoints are represented as small dots. Please note that dashed lines indicate linear recovery trajectories, which may deviate from the actual situation. *Note:* TKA = total knee arthroplasty, THA = total hip arthroplasty, HC = healthy controls.

