

# What's the effectiveness of stocking actions in small creeks? the role of water discharge behind hatchery trout downstream movement

Stefano Brignone <sup>Corresp., 1</sup>, Vanessa De Santis <sup>1</sup>, Tiziano Putelli <sup>2</sup>, Christophe Molina <sup>2</sup>, Armando Piccinini <sup>3</sup>, Richard A. Carmichael <sup>3</sup>, Pietro Volta <sup>1</sup>

<sup>1</sup> National Research Council CNR-IRSA, Verbania, Italy, Italy

<sup>2</sup> Ufficio della caccia e della pesca, Repubblica e Cantone Ticino, Bellinzona, Canton Ticino, Switzerland

<sup>3</sup> Biomark, Boise, USA

Corresponding Author: Stefano Brignone  
Email address: stefano.brignone@irsa.cnr.it

Fish stocking to enhance freshwater fisheries or to improve the conservation status of endangered fish species is a common practice in many countries. Little is known, however, of the effectiveness of these practices in spite of the high efforts and investments required. The movement of subadult/adult hatchery-released brown trout *Salmo trutta* L. was studied by passive telemetry in a small tributary of Lake Lugano (i.e., Laveggio Creek, Canton Ticino, Switzerland). Hatchery fish, together with some resident wild individuals sampled during electrofishing surveys, were tagged with Passive Integrated Transponders (PIT) tags. Hatchery fish were released upstream and downstream a submersible monitoring antenna, which was anchored to the streambed in a pass-over orientation. The number of hatchery fish detected daily by the antenna (divided between fish released upstream and downstream the antenna) was analyzed in relation to the daily water discharge, to search for similar patterns in their fluctuation over time. Only the movement of fish released upstream the antenna displayed a significant relationship with water discharge, with the highest number of fish detected during periods of high-water flow, occurring after heavy rains. High-water discharge events had a significant role in hatchery trout downstream movement in our study site, likely acting as a driver for the downstream migration to Lake Lugano. Such events contributed to the poor effectiveness of stocking actions in this small tributary, providing further evidence against stocking strategies based on subadult/adult fish.

# What's the effectiveness of stocking actions in small creeks? the role of water discharge behind hatchery trout downstream movement

Stefano Brignone<sup>1\*</sup>, Vanessa De Santis<sup>1</sup>, Tiziano Putelli<sup>2</sup>, Christophe Molina<sup>2</sup>, Armando Piccinini<sup>3</sup>, Richard A. Carmichael<sup>3</sup>, Pietro Volta<sup>1</sup>

<sup>1</sup> CNR-IRSA, Largo Tonolli 50, 28922 Verbania Pallanza, VB, Italy.

<sup>2</sup> Ufficio della caccia e della pesca, Repubblica e Cantone Ticino, Via F. Zorzi 13, 6500 Bellinzona

<sup>3</sup> Biomark, LLC 705 S. 8th St. Boise, ID 83702, USA.

## Corresponding Author:

Stefano Brignone, CNR-IRSA, Largo Tonolli 50, 28922 Verbania Pallanza, VB, Italy. Email: stefano.brignone@irsa.cnr.it

## ABSTRACT

Fish stocking to enhance freshwater fisheries or to improve the conservation status of endangered fish species is a common practice in many countries. Little is known, however, of the effectiveness of these practices in spite of the high efforts and investments required. The movement of subadult/adult hatchery-released brown trout *Salmo trutta* L. was studied by passive telemetry in a small tributary of Lake Lugano (i.e., Laveggio Creek, Canton Ticino, Switzerland). Hatchery fish, together with some resident wild individuals sampled during electrofishing surveys, were tagged with Passive Integrated Transponders (PIT) tags. Hatchery fish were released upstream and downstream a submersible monitoring antenna, which was anchored to the streambed in a pass-over orientation. The number of hatchery fish detected daily by the antenna (divided between fish released upstream and downstream the antenna) was analyzed in relation to the daily water discharge, to search for similar patterns in their fluctuation over time. Only the movement of fish released upstream the antenna displayed a significant relationship with water discharge, with the highest number of fish detected during periods of high-water flow, occurring after heavy rains. High-water discharge events had a significant role in hatchery trout downstream movement in our study

39 site, likely acting as a driver for the downstream migration to  
 40 Lake Lugano. Such events contributed to the poor effectiveness of  
 41 stocking actions in this small tributary, providing further  
 42 evidence against stocking strategies based on subadult/adult fish.

# INTRODUCTION

Brown trout *Salmo trutta* L. is one of the fish species with the highest economic value, both for professional and recreational fisheries (ICES, 2016; Schwinn et al., 2017). Hatchery-reared brown trout have been used for a long time to enhance and/or maintain artificial populations in fresh waters (Einum & Fleming, 2001, Aprahamian et al., 2003; Flowers et al., 2019). However, hatchery fish face several problems compared with wild conspecifics, such as reduced fitness and feeding efficiency, and high mortality (Waples, 1999; Brannon et al., 2004; Araki, Cooper & Blouin, 2007). This is usually due to rearing conditions presenting high densities, unnatural food sources and absence of external stimuli, like predation and variations in water flow (Fernö & Järvi, 1998; Aarestrup et al., 2005; Pedersen, Koed & Malte, 2008).

Millions of hatchery-reared trout are still released into creeks/streams every year (Waples, 1999; Aprahamian et al., 2003; Brockmark & Johnsson, 2010; Polgar et al., 2022) despite their fate and survival being largely unknown after the release into the wild. Furthermore, a few studies have investigated the environmental variables behind the movements of hatchery-reared brown trout following their release into water courses, especially in creeks and little streams (Helfrich & Kendall, 1982; Pedersen, Dieperink & Geertz-Hansen, 2003; Aarestrup et al., 2005).

Fish movement is influenced by a variety of factors. One of the main driving forces for large-scale fish migrations is the search for suitable spawning habitats (Northcote, 1992; Larinier, 2000; Young et al., 2010; Alò et al., 2021). However, fish can move along streams and rivers for several other reasons, such as to seek better feeding grounds or better water conditions, including more suitable thermal conditions (Hughes, 1998; Gowan & Fausch, 2002; Hughes et al., 2003).

Movement is not always a volunteer action, as fish can be displaced downstream because of high-water discharge, and this is especially true for juveniles, less capable to withstand high-water flows compared to adults (Jowett & Richardson, 1989; Hayes, 1995; Nislow et al., 2002).

Contrastingly, periods of high-water flow may also favor upstream movements allowing fish to take advantage of temporary passages created during high-flow conditions and, in general, to overcome in-river barriers more easily (Jonsson, 1991; Jowett & Richardson, 1994; Dedual & Jowett, 1999; Schwartz & Herricks, 2005).

In many salmonid species water discharge variation, combined with other environmental factors such as water temperature and photoperiod, regulates the downstream migration from nursery grounds (Elliott, 1994; Klemetsen et al., 2003; Byrne et al., 2004). Fish move to better feeding areas or to areas with better environmental conditions, such as lakes or seas/oceans (Klemetsen et al., 2003), to maximize fitness (Olsson & Greenberg, 2004).

This migration is part of the smoltification process, which is preceded by a series of morphological, physiological and behavioral changes in juveniles, and starts as they reach a minimum size, depending on the salmonid species (Thorstad et al., 2011).

Brown trout smolts, for instance, initiate their downstream migration a few years after hatching, usually as 2- and 3-year-old individuals. However, hatching can also take place in 1- to 8-year-old fish/individuals (Elliott, 1994; Jonsson et al., 1999; Klemetsen et al., 2003).

Despite the migratory potential of the species, stocking subadult and adult fish is still considered as a useful method to increase biomass of catchable fish both in the short and long term. To test the effectiveness (i.e., the short-term availability of catchable fish) of brown trout stocking with subadult and adult fish in a small creek, we analyzed the movement of hatchery-reared brown trout released in Laveggio Creek (Lake Lugano catchment, Canton Ticino, Southern Switzerland), assessing the role exerted by water discharge in altering their post-release residence time and abundance.

We hypothesize that water discharge plays an important role in fish movement, either as a physical stimulus behind hatchery trout migration towards lake habitats (active movement), or as a physical factor forcing hatchery fish to move downstream, due to their scarce attitude to sustained swimming during high water discharge events (passive movement).

# MATERIALS & METHODS

## Study area

Laveggio is a small creek located in the southern side of the Alps (45°54'N 8°58'E), at the Swiss-Italian border within the Lake Lugano basin. Lake Lugano is a eutrophic lake with a surface area of 49 km<sup>2</sup> (Fig. 1). Laveggio Creek has a spring origin, flowing for 11 km; the streambed width ranges from 4 to 8 meters, with many human-altered and channelized sections, except for some of its uppermost portions (<https://map.geo.admin.ch>). Located in an area with high rainfall (mean daily precipitation: 4.0 ± 10.7 mm, max daily precipitation: 82.5 ± 11.2 mm, last 5 years data), this creek can undergo dramatic changes in its discharge rate, reaching more than 10 m<sup>3</sup> s<sup>-1</sup> (mean daily discharge: 1.1 ± 1.2 m<sup>3</sup> s<sup>-1</sup>).

Water temperature varies significantly during the year, peaking in the summer (~ 27 °C) and reaching its coldest point in the winter (~ 6 °C).

Precipitation and water temperature data were taken from the Environmental Observatory of Switzerland (<https://www.oasi.ti.ch/web/dati/idrologia.html>).

The fish community is made of brown trout *S. trutta*, European river lamprey *Lampetra fluviatilis* (L.), European perch *Perca fluviatilis* L., roach *Rutilus rutilus* (L.), Italian chub *Squalius squalus* (Bonaparte, 1837) and the Italian dace *Telestes muticellus* (Bonaparte, 1837).

On the streambed of Laveggio, around 1.2 km upstream from the confluence with Lake Lugano (Fig. 1), we positioned two high-density polyethylene (HDPE) antennas (3 meters each; Biomark, Boise, Idaho, USA), which were lined up to cover the entire width of the streambed, forming a singular flat-bed PIT monitoring system. The antennas were anchored to the streambed of Laveggio, at the end of a large pool in approx. 30 cm of water, to make sure to detect fish during high-water events.

This system usually registers the passage of PIT tagged fish moving up to 10 m s<sup>-1</sup> and within a range up to 0.9 meters, although the latter can vary according to the physico-chemical properties of waters and to the background noise (e.g., electromagnetic interference from a nearby-electrified railway). Once fish enter

the electromagnetic field generated by the antennas, the implanted PIT tag is charged and can transmit its unique identification code.

The PIT tag antenna system together with its recording station (Biomark Master Controller), was installed and anchored to the streambed at the end of August 2020.

# **Fish sampling**

Fish used in this study came from the fish hatchery managed by the Office of hunting and fishing of Canton Ticino, Bellinzona (Switzerland). After eggs hatched, trout were reared under natural light conditions and fed with commercial pellet.

Stocking with subadult and adult fish in Laveggio Creek is forbidden by law and, thus, the stocking is usually carried out using eggs and fry. However, to study the movement of trout in Laveggio, we were allowed to mark and release 998 subadult/adult hatchery-reared trout during autumn 2020, and 520 during autumn 2021.

PIT tags (Biomark APT12, Advanced Performance Tag, 12 x 2 mm, ISO Standard, 134.2 kHz) were implanted in the intraperitoneal cavity through a Biomark MK25 implanter and pre-loaded single-use needles to minimize fish stress and possible disease transmission and to reduce the healing time. Fish underwent a light eugenol anesthesia during handling and tagging operations, while feeding was stopped 2 days prior to the marking day. All trout remained in the hatchery, in a dedicated tank for at least 10 days after tagging and before release to monitor the healing of the implant incision and to allow a full recovery. Feeding resumed 24 hours after the tagging day. No mortality was observed during the 10 days of recovery time.

Fish released in 2020 were equally divided between fish of age 1<sup>+</sup> (total length range: 15-20 cm) and 2<sup>+</sup> (total length range: 20-30 cm), whereas all fish released in 2021 were of age 1<sup>+</sup>. Hatchery trout were marked and released at the end of October 2020 and 2021 at four different sectors of the creek, one sector close to the confluence with Lake Lugano, downstream of the antenna, and three sectors upstream of the antenna (Fig. 1).

After a few days of data monitoring, some of the fish seemed to pause in the proximity of the antenna; this location was probably used as temporary feeding area and refuge. A delay of 15 seconds between each recording was necessary to avoid memory overload.

During the study period, in order to search for marked trout, we conducted six electrofishing surveys using a Hans-Grassl EL64GII electrofisher, with settings of 7000 W, 300 V. Because of the study area extension, we had to focus on different portions of the creek in each electrofishing survey: between sector 3 and 4 during the first survey (January 2021), near sector 1 and 3 during the second (February 2021), near sector 2 and 3 during the third (first half of April 2021), near the antenna and between the sector 3 and 4 during the fourth (second half of April 2021), between sector 3 and 4 during the fifth (June 2021), and near the antenna during the sixth survey (December 2021). During three surveys (first and second half of April and December 2021), 226 wild fish were captured, measured (total length and total weight) and marked with PIT tags before being released back to their site of capture.

Wild trout movement data was used to look for possible differences between hatchery and wild trout response to high water discharge events, although the number of wild trout marked was not comparable with hatchery conspecifics.

The experiment was conducted with permission of the 'Dipartimento della sanità e della socialità-Esperimenti su animali', authorization N° TI-27-2020 (Supplementary materials).



## Data collection

Fish passages started to be recorded by the antenna from October 29<sup>th</sup>, 2020, and terminated on April 30<sup>th</sup>, 2022, the last date in which hatchery trout were detected by the antenna. The main variable measured was the Number of Fish Detected Daily (hereinafter referred as NFDD) by the antenna, which is the total amount of individuals registered by the antenna during one day. Hatchery fish were divided into two groups based on whether they were released upstream (NFDD upstream) or downstream (NFDD downstream) the antenna, in order to discriminate the role of high-water events.

To assess the relation between NFDD (upstream and downstream) and daily water discharge, hydrological data were retrieved from the Environmental Observatory of Switzerland (<https://www.oasi.ti.ch/web/dati/idrologia.html>). Hydrological data were recorded by a monitoring station positioned close to the PIT tag antenna (Riva San Vitale).

## Statistical analysis

NFDD released upstream and downstream and water discharge fluctuated over time, so they were analyzed implementing a time series analysis approach.

The ARIMA (Autoregressive Integrated Moving Average) model was applied to analyze the structure of each variable. The optimal model was determined by automatic model selection based on the Akaike information criterion (AIC) function, included in the package "forecast" (Hyndman et al., 2020). The residuals of each resulting model were examined to check possible structures within each time series. In case a trend in the residual distribution of the time series is present, an adjustment for the estimated coefficients and standard errors is required to fit the generalized estimating equation (GEE) of generalized linear model, performed using the "geepack" package included in R software (Højsgaard et al., 2016).

Cross-correlation (CC) analysis was used to measure the similarity between time series and, in particular, to look for shifted patterns between water discharge time series and NFDD time series;

272 this was done using the R package “tseries” (Trapletti et al.,  
273 2015) .

# RESULTS

In total, 1518 hatchery trout were PIT-tagged and released in Laveggio Creek. Among these trout, 222 fish were registered by the antenna at least once, which corresponded to only the 14.6 % of the total amount of trout released in the creek. The highest number of fish registered by the antenna in a day was 38, one day after the release date in 2021.

From November 29<sup>th</sup> to December 7<sup>th</sup>, 2020, the antenna stopped recording data because of memory overload, whereas from June 7<sup>th</sup> to July 9<sup>th</sup>, 2021, it stopped working for an electrical malfunction (Fig. 2,3).

During the five electrofishing surveys conducted in the first half of 2021, only nine marked trout were recaptured: six in January, one in February, two in the first half of April and zero during the last two surveys in the second half of April and in June. During the survey of June 2021 23 marked wild trout were recaptured. During the electrofishing survey of December 2021, one month after the second release of hatchery trout in Laveggio Creek, only 4 hatchery trout were recaptured, together with one marked wild trout.

Electrofishing efficiency was supposed to be high, since together with the few hatchery trout found during these surveys, we were able to sample several wild trout along Laveggio Creek, most of them under-size for PIT tag marking and therefore not considered further.

During three electrofishing surveys, 226 wild trout were marked with PIT tags (75 and 139 fish in the first and second part of April 2021, respectively, and 12 in December 2021), with an average length of  $18.7 \pm 4.7$  cm. The number of wild fish detected daily by the antenna is shown in Figure 4.

Six of the wild fish detected by the antenna stationed in the proximity of the antenna for a very long time (two individuals for almost one year), suggesting that these trout particularly appreciated the antenna area which was potentially used as refuge/feeding ground.

Water discharge was on average  $1.0 \pm 1.0$  m<sup>3</sup> s<sup>-1</sup> and exceeded 3.5 m<sup>3</sup> s<sup>-1</sup> for 13 non-consecutive days, usually after short periods of

heavy rains, and water levels remained high for several days after these events (Fig. 2,3).

For each analyzed time series, ARIMA model outputs revealed that residuals were randomly distributed (Supplementary materials). ACF and PACF did not show any relevant peak for the time series of NFDD released downstream (Fig. S2), whereas the ACF and PACF of NFDD released upstream and water discharge time series showed one or two peaks far from the zero, respectively (Fig. S1 and S3 for NFDD released upstream and water discharge series, respectively). Since residuals of the ARIMA model were randomly distributed, with no evident structure, there was no need to adjust the estimated coefficient and standard error, and the generalized estimating equation (GEE) could be fitted.

In the generalized linear regression model obtained through the GEE, only NFDD released upstream displayed a significant relationship with water discharge (Table 1). No significant relationship was found between NFDD released downstream and water discharge (Table 2).

Between NFDD released upstream and water discharge, Cross-correlation (CC) found a correlation at lag 1, suggesting a 1-day shift between the two time series (Fig. 5).

CC analysis highlighted a shifted pattern between NFDD released downstream and water discharge at lag 2, suggesting a 2-days shift between the two time series (Fig. 6). However, since there was no clear relation between these time series, this result could be misleading.

# DISCUSSION

Trout stocking into fresh waters has been a common practice for decades (Brockmark & Johnsson, 2010), but its effectiveness and utility to support local populations and/or local fishing is debated (Halverson, 2008; Araki & Schmid, 2010; Hunt & Jones, 2018; Pinter et al., 2018).

The life stage of fish used for stocking may vary depending on the final aims of the stocking activities (conservation and/or supplementation for recreational fishing), as well as on the hatchery dimensions and production capacity (Cowx, 1994; Letcher & Terrick, 2001). Stocking with eggs (Vibert boxes, cocooning) and with yolk-sac fry is one of the most used procedures for stocking (Kirkland, 2012). However, the use of juveniles and subadult fish is still implemented in many freshwater fisheries and considered a suitable system to artificially increase the productivity of water courses (Baer, Blasel & Diekmann, 2007). Nevertheless, the adverse effects of domestication on fish physiology and behavior are not often considered and fish mortality is usually underestimated (Araki & Schmid, 2010).

In this study we used PIT tag telemetry to monitor the movement of post-release subadult/adult hatchery trout in a small creek, a typical environment subject to stoking practices for fishery enhancement. The number of fish detected by the PIT tag antenna steadily decreased monthly and, only six months after both release dates, no more hatchery trout were detected by the antenna. The steep reduction in hatchery trout abundance after the release in wild ecosystems has been also observed in several other studies (Näslund, 1992; Wiley et al., 1993; Pedersen, Dieperink & Geertz-Hansen, 2003; Pinter et al. 2018), including also water courses of Canton Ticino (Molina, 2019) and Switzerland (Gmünder & Friedl, 2002).

Water discharge significantly influenced the movement of hatchery trout across the antenna in this study. The effects of water discharge strongly depended on the release site: peaks in water discharge were mirrored by peaks in fish passages only for fish released upstream the antenna, with one-day lag after high-water events.

The relationship between fish movement and water discharge has been observed by many authors, in several fish species and environments. For example, Tytler, Thorpe & Shearer (1978) and Thorpe et al. (1981) observed a strong passive displacement in *Salmo salar* L. smolts, which were mainly transported to the ocean after high spring runoffs, which is one of the factors controlling their anadromous behavior (Jonsson & Jonsson, 2009). The downstream movement of smolts can also be a survival strategy, as highlighted by Hansen & Jonsson (1985), where moving into the main current of the river allows smolts to avoid water sloughs and remnant pools. In contrast, Brown, Power & Beltaoa (2001) observed an opposite behavior: many fish take advantage of shelters in the creek margins, to minimize energy expenditure, avoid strong currents, and mitigate the risk of uncontrolled downstream dispersal. It has been observed that brown trout, in particular, move to side channels searching for refuges from peak flows (Bunt et al., 1999). These strategies enable fish to avoid physical injuries and getting trapped in shallow waters, where they would inevitably die as water dries up or because of anoxia (Ortlepp & Mürle, 2003). An alternative tactic used by some fish, described by Brown, Power & Beltaoa (2001), is to take refuge in very large and deep pools during floods: these pools usually have a very low water velocity close to the bottom, which provides a suitable place where fish can endure flood events.

Water discharge variation is a well-known stimulus initiating brown trout smolt descend to lentic habitats, such as alpine lakes (Klemetsen et al., 2003); such habitats provide higher food availability and more favorable environmental conditions compared to nursery grounds (Økland et al., 1993; Olsson et al., 2006). The observed peaks in water discharge followed by peaks in fish moving downstream after a one-day lag suggest that these steep variations in water flow triggered the downstream movement of post-release hatchery trout.

Hatchery trout released in Laveggio Creek were brown trout of age 1+ and 2+, in which the smoltification process, although not assessed in this study, might have played a fundamental role in their downstream movement towards Lake Lugano, even after a few days/weeks from the release (Flowers et al., 2019).

Space and food competition with wild trout could also play an important role in the downstream migration of hatchery trout, since wild fish could induce hatchery trout to search for areas with lower intraspecific competition (low population density, higher food availability), e.g., lake habitats (Olsson et al., 2006).

Another hypothesis behind the hatchery trout downstream movement, may be attributed to a passive transportation due to the high-water events occurred in Laveggio Creek. Laveggio is for a good portion of its lower course a channelized creek, with most of its banks being completely human altered; furthermore, it exhibits few slow and deep pools where fish can take refuge. Therefore, trout may struggle to find any suitable cover on creek margins or in deep pools during high-water events. As stated before, spatial competition with established wild trout could also limit the number of available covers for post-release hatchery trout, through a prior-resident effect (Deverill, Adams & Bean, 1999; Sánchez-González & Nicieza, 2021).

Hatchery trout are more vulnerable to high-current conditions compared to wild trout (Pedersen, Koed & Malte, 2008): hatchery trout usually have reduced fin quality due to abrasion with tank sides and over-density conditions (Drucker & Lauder, 2003; Ojanguren & Brana, 2003). Lack of genetic diversity is also common in hatchery-reared fish, which, coupled with artificial selection, can favor the maintenance of stocks made by sedentary individuals that invest energy intake in growth rather than in movement (Duthie, 1987; Molina, 2019). Hatchery fish are usually bred under low flow rates, which may also reduce their activity levels, reducing activity of fatty acid oxidation enzymes in slow and fast muscle (Johnston & Moon, 1980).

Nevertheless, a good portion of the hatchery fish released upstream the antenna where registered again after the first passage across the antenna following a high-water event, so this passive transportation, if any, was limited to a little percentage of the released fish. Therefore, the results of the present study point mainly to the first hypothesis, for which water discharge acted mainly as a driver for the active downstream migration of the released hatchery trout.

These assumptions on the fate of hatchery trout along Laveggio Creek should take into account the antenna detection range and efficiency, which is a potential source of bias, quite difficult to estimate. In general, the detection efficiency of PIT tag antennas is not constant over time, and can rarely reach 100% (Horton, Dubreuil & Letcher, 2007; Connolly et al., 2008). Several factors can alter PIT tag antenna detection efficiency, such as tag orientation, water properties (e.g., water salinity), tag collision and external electromagnetic interferences such as high-voltage power line (Zydlewski et al., 2001; Aymes & Rives, 2009; Zentner et al., 2021). However, Laveggio is a shallow creek, and the noise sources are limited (according to the noise levels measured by the recording station Biomark Master Controller), thus the detection efficiency during normal flows should be potentially very high. Nevertheless, high-water discharge events (such as the ones occurred during the study period) may have decreased the detection range and efficiency of the PIT tag antenna (Zydlewski et al., 2006; Horton, Dubreuil & Letcher, 2007). Hence, we could not exclude that there may have been other fish that moved over the antenna position, but which were not detected because of the reduced antenna detection range during periods of high-water flow. Yet, the low number of hatchery trout remained in the study area after a few months from the release was also pointed out by the results of five electrofishing surveys performed during the same period, in which only a few marked trout (13 trout, 0.9% of the total amount released into the creek) were detected. This low number could also be related to the low percentage of creek sampled during the electrofishing surveys, compared to the extension of the study area.

Despite our data suggest the importance of high-water discharge events behind the downstream movement of hatchery-reared brown trout (only fish released upstream the antenna were significantly related to water discharge), we cannot exclude the contribution of other factors in the apparent steep decrease of hatchery trout abundance. For instance, harvest by anglers can be very impactful on trout populations, especially in little creeks like Laveggio (Almodóvar & Nicola, 2004; Wallace, 2010; Flowers et al., 2019). Since anglers do not have the ability to detect or report interactions with PIT tagged fish, we were limited in our ability to assess angler exploitation and impact on movement and survival.



However, the minimum size limit for trout harvest is set at 24 cm and at least 70% of the hatchery-reared fish were under this limit, so the role of fishing harvest on our results, if any, was limited.

Another important factor behind the drop in hatchery trout abundance from Laveggio could be linked to avian predation. This predation can be very effective and cause huge drops in trout abundance, and this is particularly relevant for hatchery fish (Matkowski, 1989; Dieperink, Pedersen & Pedersen, 2001). As stated before, hatchery fish are not used to sustained swimming and/or to predator avoidance, which, coupled with their more surface-oriented behaviors, makes them more vulnerable to bird predation than wild conspecifics (Maynard, Flagg & Mahnken, 1995; Dieperink, Pedersen & Pedersen, 2001). The collection of a PIT tag inside the stomach of a dead grey heron *Ardea cinerea* (L.) along the Laveggio Creek banks (Molina pers. comm.) could suggest a hidden effect of waterbirds on hatchery trout.

These factors, coupled with the downstream movement following high-water events, may help to explain the low percentage of recaptured trout during the electrofishing surveys, as well as the low percentage of fish detected by the antenna (0.9 % and 14.6 % of the marked hatchery trout, respectively), suggesting that most of the released trout died or moved downstream undetected by the antenna. Unfortunately, it was not possible to discriminate between these two hypotheses, and this aspect should be considered in future studies.

Wild trout movements seemed to be less influenced by high water events than hatchery trout. Some of the wild fish marked were detected by the antenna for the entire duration of the study period, despite the repeated high-water events particularly occurring in summer and early autumn 2021. However, comparisons between wild and hatchery trout should be taken with caution since data on wild fish were scarce, both for the low number of wild fish marked and for the low number of wild fish detected by the antenna; besides, wild trout were marked in different time period than hatchery trout. For these reasons any further assumptions on the resilience of wild trout and their differences with hatchery trout behavior in Laveggio Creek are limited.

In conclusion, our results suggest that stocking for fishery enhancement in creeks like Laveggio should probably be adjusted and moved toward different methods, since the traditional stocking activities with subadult/adult trout is not effective. We highlighted how fragile the stability of artificial populations made of hatchery trout is in this type of environment, and how easily different biotic and abiotic factors can disrupt this stability.

We cannot exclude the usefulness of these stocking actions on the long-term period, since hatchery trout moved downstream may potentially return back in Laveggio Creek as spawners, enhancing the natural stock of this freshwater system. The PIT tag antenna monitoring system used in this study will be maintained for the next years and it will contribute to shed new light on this possible hidden aspect.

# CONCLUSIONS

This study showed how the effectiveness of stocking practices using subadult and adult hatchery brown trout is limited by downstream movements toward lentic connected systems, eased by high-water discharge events. Peaks in water discharge mirrored peaks in PIT-tagged hatchery trout detections registered by a PIT tag antenna system. The number of fish detected by the antenna decreased monthly until, after only six months from the two release dates, the number of hatchery trout detected was close to zero. The absence of hatchery trout was also confirmed by different electrofishing surveys carried out in the study site. High-water flow events, coupled with other factors, could be the main causes behind the poor effectiveness of stocking actions using subadult/adult fish.

# Acknowledgements

We thank the office of hunting and fishing of Canton Ticino, Switzerland, for the data collection, field work and revisions. In particular, we thank Danilo Foresti (Ufficio della caccia e della pesca, Repubblica e Cantone Ticino) and Diego Dagani (BAFU) for the insightful comments.

# REFERENCES

- Aarestrup K, Jepsen N, Koed A, Pedersen S. 2005. Movement and mortality of stocked brown trout in a stream. *Journal of Fish Biology*, 66(3), 721-728. <https://doi.org/10.1111/j.0022-1112.2005.00634.x>
- Almodóvar A, Nicola GG. 2004. Angling impact on conservation of Spanish stream-dwelling brown trout *Salmo trutta*. *Fisheries Management and Ecology*, 11(3-4), 173-182. <https://doi.org/10.1111/j.1365-2400.2004.00402.x>
- Alò D, Lacy SN, Castillo A, Samaniego HA, Marquet PA. 2021. The macroecology of fish migration. *Global Ecology and Biogeography*, 30(1), 99-116. <https://doi.org/10.1111/geb.13199>
- Aprahamian MW, Smith KM, McGinnity P, McKelvey S, Taylor J. 2003. Restocking of salmonids—opportunities and limitations. *Fisheries Research*, 62(2), 211-227. [https://doi.org/10.1016/S0165-7836\(02\)00163-7](https://doi.org/10.1016/S0165-7836(02)00163-7)
- Araki H, Cooper B, Blouin MS. 2007. Genetic effects of captive breeding cause a rapid, cumulative fitness decline in the wild. *Science*, 318(5847), 100-103. DOI: 10.1126/science.1145621
- Araki H, Schmid C. 2010. Is hatchery stocking a help or harm? Evidence, limitations and future directions in ecological and genetic surveys. *Aquaculture*, 308, S2-S11. <https://doi.org/10.1016/j.aquaculture.2010.05.036>
- Aymes JC, Rives J. 2009. Detection efficiency of multiplexed passive integrated transponder antennas is influenced by environmental conditions and fish swimming behaviour. *Ecology of Freshwater Fish*, 18(4), 507-513. <https://doi.org/10.1111/j.1600-0633.2009.00373.x>
- Baer J, Blaschel K, Diekmann M. 2007. Benefits of repeated stocking with adult, hatchery-reared brown trout, *Salmo trutta*, to recreational fisheries? *Fisheries Management and Ecology*, 14(1), 51-59. <https://doi.org/10.1111/j.1365-2400.2006.00523.x>
- Brannon EL, Amend DF, Cronin MA, Lannan JE, LaPatra S, McNeil W J, Noble RE, Smith CE, Talbot AJ, Wedemeyer GA, Westers H. 2004. The controversy about salmon hatcheries. *Fisheries*, 29(9), 12-31. [https://doi.org/10.1577/1548-8446\(2004\)29\[12:TCASH\]2.0.CO;2](https://doi.org/10.1577/1548-8446(2004)29[12:TCASH]2.0.CO;2)

618 Brockmark S, Johnsson JI. 2010. Reduced hatchery rearing density  
619 increases social dominance, postrelease growth, and survival in  
620 brown trout (*Salmo trutta*). *Canadian Journal of Fisheries and*  
621 *Aquatic Sciences*, 67(2), 288-295. <https://doi.org/10.1139/F09-185>

622 Brown RS, Power G, Beltaoa S. 2001. Winter movements and habitat  
623 use of riverine brown trout, white sucker and common carp in  
624 relation to flooding and ice break-up. *Journal of Fish Biology*,  
625 59(5), 1126-1141. [https://doi.org/10.1111/j.1095-](https://doi.org/10.1111/j.1095-8649.2001.tb00180.x)  
626 [8649.2001.tb00180.x](https://doi.org/10.1111/j.1095-8649.2001.tb00180.x)

627 Bunt CM, Cooke SJ, Katopodis C, McKinley RS. 1999. Movement and  
628 summer habitat of brown trout (*Salmo trutta*) below a pulsed  
629 discharge hydroelectric generating station. *Regulated Rivers:*  
630 *Research & Management: An International Journal Devoted to River*  
631 *Research and Management*, 15(5), 395-403.  
632 [https://doi.org/10.1002/\(SICI\)10991646\(199909/10\)15:5<395::AIDRR](https://doi.org/10.1002/(SICI)10991646(199909/10)15:5<395::AIDRR)  
633 [R556>3.0.CO;2-1](https://doi.org/10.1002/(SICI)10991646(199909/10)15:5<395::AIDRR)

634 Byrne CJ, Poole R, Dillane M, Rogan G, Whelan KF. 2004. Temporal  
635 and environmental influences on the variation in sea trout (*Salmo*  
636 *trutta* L.) smolt migration in the Burrishoole system in the west  
637 of Ireland from 1971 to 2000. *Fisheries Research*, 66(1), 85-94.  
638 [https://doi.org/10.1016/S0165-7836\(03\)00146-2](https://doi.org/10.1016/S0165-7836(03)00146-2)

639 Connolly PJ, Jezorek IG, Martens KD, Prentice EF. 2008. Measuring  
640 the performance of two stationary interrogation systems for  
641 detecting downstream and upstream movement of PIT-tagged  
642 salmonids. *North American Journal of Fisheries Management*, 28(2),  
643 402-417. <https://doi.org/10.1577/M07-008.1>

644 Cowx IG. 1994. Stocking strategies. *Fisheries management and*  
645 *ecology*, 1(1), 15-30. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2400.1970.tb00003.x)  
646 [2400.1970.tb00003.x](https://doi.org/10.1111/j.1365-2400.1970.tb00003.x)

647 Dedual M, Jowett IG. 1999. Movement of rainbow trout (*Oncorhynchus*  
648 *mykiss*) during the spawning migration in the Tongariro River, New  
649 Zealand. *New Zealand Journal of Marine and Freshwater Research*,  
650 33(1), 107-117. <https://doi.org/10.1080/00288330.1999.9516861>

651 Deverill JI, Adams CE, Bean CW. 1999. Prior residence, aggression  
652 and territory acquisition in hatchery-reared and wild brown trout.  
653 *Journal of fish Biology*, 55(4), 868-875.  
654 <https://doi.org/10.1111/j.1095-8649.1999.tb00723.x>

655 Dieperink C, Pedersen S, Pedersen MI. 2001. Estuarine predation on  
656 radiotagged wild and domesticated sea trout (*Salmo trutta* L.)  
657 smolts. *Ecology of Freshwater Fish*, 10(3), 177-183.  
658 <https://doi.org/10.1034/j.1600-0633.2001.100307.x>

659 Drucker EG, Lauder GV. 2003. Function of pectoral fins in rainbow  
660 trout: behavioral repertoire and hydrodynamic forces. *Journal of*  
661 *experimental biology*, 206(5), 813-826.  
662 <https://doi.org/10.1242/jeb.00139>

663 Duthie GG. 1987. Observations of poor swimming performance among  
664 hatchery-reared rainbow trout, *Salmo gairdneri*. *Environmental*  
665 *Biology of Fishes*, 18(4), 309-311.  
666 <https://doi.org/10.1007/BF00004884>

667 Einum S, Fleming IA. 2001. Implications of stocking: ecological  
668 interactions between wild and released salmonids. *Nordic Journal*  
669 *of Freshwater Research*, 75, 56-70.

670 Elliott J. M. 1994. Quantitative ecology and the brown trout.  
671 *Oxford University Press*.

672 Fernö A, Järvi T. 1998. Domestication genetically alters the anti-  
673 predator behaviour of anadromous brown trout (*Salmo trutta*)-a  
674 dummy predator experiment. *Nordic Journal of Freshwater Research*,  
675 95-100. <http://113.160.249.209:8080/xmlui/handle/123456789/4842>

676 Flowers HJ, Kwak TJ, Fischer JR, Cope WG, Rash JM, Besler DA. 2019.  
677 Behavior and survival of stocked trout in southern Appalachian  
678 Mountain streams. *Transactions of the American Fisheries Society*,  
679 148(1), 3-20. <https://doi.org/10.1002/tafs.10113>

680 Gmünder R, Friedl C. 2002. Efficacité des repeuplements piscicoles  
681 effectués en Suisse. *Informations concernant la pêche N° 71, 1st*  
682 *edition*. Office fédéral de l'environnement, des forêts et du  
683 paysage OFEFP, Bern.

684 Gowan C, Fausch KD. 2002. Why do foraging stream salmonids move  
685 during summer? *Environmental Biology of Fishes*, 64(1), 139-153.  
686 <https://doi.org/10.1023/A:1016010723609>

687 Halverson MA. 2008. Stocking trends: a quantitative review of  
688 governmental fish stocking in the United States, 1931 to 2004.  
689 *Fisheries*, 33(2), 69-75. [https://doi.org/10.1577/1548-8446-](https://doi.org/10.1577/1548-8446-33.2.69)  
690 [33.2.69](https://doi.org/10.1577/1548-8446-33.2.69)

691 Hansen LP, Jonsson B. 1985. Downstream migration of hatchery-  
 692 reared smolts of Atlantic salmon (*Salmo salar* L.) in the River  
 693 Imsa, Norway. *Aquaculture*, 45(1-4), 237-248.  
 694 [https://doi.org/10.1016/0044-8486\(85\)90273-X](https://doi.org/10.1016/0044-8486(85)90273-X)

695 Hayes JW. 1995. Spatial and temporal variation in the relative  
 696 density and size of juvenile brown trout in the Kakanui River,  
 697 North Otago, New Zealand. *New Zealand Journal of Marine and*  
 698 *Freshwater Research*, 29(3), 393-407.  
 699 <https://doi.org/10.1080/00288330.1995.9516674>

700 Helfrich LA, Kendall WT. 1982. Movements of hatchery-reared  
 701 rainbow, brook, and brown trout stocked in a Virginia mountain  
 702 stream. *The Progressive Fish-Culturist*, 44(1), 3-7.  
 703 [https://doi.org/10.1577/1548-8659\(1982\)44\[3:MOHRBA\]2.0.CO;2](https://doi.org/10.1577/1548-8659(1982)44[3:MOHRBA]2.0.CO;2)

704 Højsgaard S, Halekoh U, Yan J, Højsgaard MS. 2016. Package  
 705 'geepack'. R package version, 1-2.

706 Horton GE, Dubreuil TL, Letcher BH. 2007. A model for estimating  
 707 passive integrated transponder (PIT) tag antenna efficiencies for  
 708 interval-specific emigration rates. *Transactions of the American*  
 709 *Fisheries Society*, 136(5), 1165-1176. [https://doi.org/10.1577/T06-](https://doi.org/10.1577/T06-053.1)  
 710 [053.1](https://doi.org/10.1577/T06-053.1)

711 Hughes NF. 1998. A model of habitat selection by drift-feeding  
 712 stream salmonids at different scales. *Ecology*, 79(1), 281-294.  
 713 [https://doi.org/10.1890/0012-9658\(1998\)079\[0281:AMOHBS\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1998)079[0281:AMOHBS]2.0.CO;2)

714 Hughes NF, Hayes JW, Shearer KA, Young RG. 2003. Testing a model  
 715 of drift-feeding using three-dimensional videography of wild brown  
 716 trout, *Salmo trutta*, in a New Zealand river. *Canadian Journal of*  
 717 *Fisheries and Aquatic Sciences*, 60(12), 1462-1476.  
 718 <https://doi.org/10.1139/f03-126>

719 Hunt TL, Jones P. 2018. Informing the great fish stocking debate:  
 720 An Australian case study. *Reviews in Fisheries Science &*  
 721 *Aquaculture*, 26(3), 275-308.  
 722 <https://doi.org/10.1080/23308249.2017.1407916>

723 Hyndman RJ, Athanasopoulos G, Bergmeir C, Caceres G, Chhay L,  
 724 O'Hara-Wild M, Petropoulos F, Razbash S, Wang E. 2020. Package  
 725 'forecast'. [Online] [https://cran.](https://cran.r-project.org/web/packages/forecast/forecast.pdf) [r-project.](https://cran.r-project.org/web/packages/forecast/forecast.pdf)  
 726 [org/web/packages/forecast/forecast.pdf](https://cran.r-project.org/web/packages/forecast/forecast.pdf).

ICES. 2016. Report of the Baltic Salmon and Trout Assessment Working Group (WGBAST). 30 March-6 April 2016, Klaipeda, Lithuania. ICES CM 2016/ACOM:09 257 pp.

Johnston IA, Moon TW. 1980. Exercise training in skeletal muscle of brook trout (*Salvelinus fontinalis*). *Journal of Experimental Biology*, 87(1), 177-194. <https://doi.org/10.1242/jeb.87.1.177>

Jonsson N. 1991. Influence of water flow, water temperature and light on fish migration in rivers. *Nordic journal of freshwater research*, 66(1991), 20-35.

Jonsson N, Næsje TF, Jonsson B, Saksgård R, Sandlund OT. 1999. The influence of piscivory on life history traits of brown trout. *Journal of Fish Biology*, 55(6), 1129-1141. <https://doi.org/10.1111/j.1095-8649.1999.tb02064.x>

Jonsson B, Jonsson N. 2009. A review of the likely effects of climate change on anadromous Atlantic salmon *Salmo salar* and brown trout *Salmo trutta*, with particular reference to water temperature and flow. *Journal of fish biology*, 75(10), 2381-2447. <https://doi.org/10.1111/j.1095-8649.2009.02380.x>

Jowett IG, Richardson J. 1989. Effects of a severe flood on instream habitat and trout populations in seven New Zealand rivers. *New Zealand journal of marine and freshwater research*, 23(1), 11-17. <https://doi.org/10.1080/00288330.1989.9516335>

Jowett IG, Richardson J. 1994. Comparison of habitat use by fish in normal and flooded river conditions. *New Zealand Journal of Marine and Freshwater Research*, 28(4), 409-416. <https://doi.org/10.1080/00288330.1994.9516631>

Klemetsen A, Amundsen PA, Dempson JB, Jonsson B, Jonsson N, O'connell MF, Mortensen E. 2003. Atlantic salmon *Salmo salar* L., brown trout *Salmo trutta* L. and Arctic charr *Salvelinus alpinus* (L.): a review of aspects of their life histories. *Ecology of freshwater fish*, 12(1), 1-59. <https://doi.org/10.1034/j.1600-0633.2003.00010.x>

Kirkland D. 2012. A review of factors influencing artificial salmonid incubation success and a spate river-specific incubator design. *Fisheries Management and Ecology*, 19(1), 1-9. <https://doi.org/10.1111/j.1365-2400.2011.00807.x>



763 Larinier M. 2000. Dams and fish migration. World Commission on  
764 Dams. (23 March 2012; [www.unep.org/dams/WCD/report.asp](http://www.unep.org/dams/WCD/report.asp))

765 Letcher BH, Terrick TD. 2001. Effects of developmental stage at  
766 stocking on growth and survival of Atlantic salmon fry. *North*  
767 *American Journal of Fisheries Management*, 21(1), 102-110.  
768 [https://doi.org/10.1577/1548-8675\(2001\)021<0102:EODSAS>2.0.CO;2](https://doi.org/10.1577/1548-8675(2001)021<0102:EODSAS>2.0.CO;2)

769 Matkowski SM. 1989. Differential susceptibility of three species  
770 of stocked trout to bird predation. *North American Journal of*  
771 *Fisheries Management*, 9(2), 184-187.  
772 [https://doi.org/10.1577/15488675\(1989\)009<0184:DSOTSO>2.3.CO;2](https://doi.org/10.1577/15488675(1989)009<0184:DSOTSO>2.3.CO;2)

773 Maynard DJ, Flagg TA, Mahnken CV. 1995. A review of seminatural  
774 culture strategies for enhancing the post release survival of  
775 anadromous salmonids. In *American Fisheries Society Symposium*  
776 (Vol. 15, pp. 307-314).

777 Molina C. 2019. Certificate of Advanced Studies (CAS), Poissons  
778 d'eau douce d'Europe-Ecologie et Gestion. *Caractérisation*  
779 *génétique des truites de rivière du Canton du Tessin*.

780 Näslund I. 1992. Survival and distribution of pond-and  
781 hatchery-reared 0+ brown trout, *Salmo trutta* L., released in a  
782 Swedish stream. *Aquaculture Research*, 23(4), 477-488.  
783 <https://doi.org/10.1111/j.1365-2109.1992.tb00791.x>

784 Nislow KH, Magilligan FJ, Folt CL, Kennedy BP. 2002. Within-basin  
785 variation in the short-term effects of a major flood on stream  
786 fishes and invertebrates. *Journal of Freshwater Ecology*, 17(2),  
787 305-318. <https://doi.org/10.1080/02705060.2002.9663899>

788 Northcote TG. 1992. Migration and residency in stream salmonids-  
789 some ecological considerations and evolutionary consequences.  
790 *Nordic journal of freshwater research*. Drottningholm, 67, 5-17.

791 Ojanguren AF, Brana F. 2003. Effects of size and morphology on  
792 swimming performance in juvenile brown trout (*Salmo trutta* L.).  
793 *Ecology of Freshwater Fish*, 12(4), 241-246.  
794 <https://doi.org/10.1046/j.1600-0633.2003.00016.x>

795 Økland F, Jonsson B, Jensen AJ, Hansen LP. 1993. Is there a  
796 threshold size regulating seaward migration of brown trout and  
797 Atlantic salmon? *Journal of Fish Biology*, 42(4), 541-550.  
798 <https://doi.org/10.1111/j.1095-8649.1993.tb00358.x>

799 Olsson IC, Greenberg LA. 2004. Partial migration in a landlocked  
800 brown trout population. *Journal of Fish Biology*, 65(1), 106-121.  
801 <https://doi.org/10.1111/j.0022-1112.2004.00430.x>

802 Olsson IC, Greenberg LA, Bergman E, Wysujack K. 2006.  
803 Environmentally induced migration: the importance of food. *Ecology*  
804 *letters*, 9(6), 645-651. [https://doi.org/10.1111/j.1461-](https://doi.org/10.1111/j.1461-0248.2006.00909.x)  
805 [0248.2006.00909.x](https://doi.org/10.1111/j.1461-0248.2006.00909.x)

806 Ortlepp J, Mürle U. 2003. Effects of experimental flooding on brown  
807 trout (*Salmo trutta fario* L.): The River Spöl, Swiss National Park.  
808 *Aquatic Sciences*, 65(3), 232-238. [https://doi.org/10.1007/s00027-](https://doi.org/10.1007/s00027-003-0666-5)  
809 [003-0666-5](https://doi.org/10.1007/s00027-003-0666-5)

810 Pedersen SS, Dieperink C, Geertz-Hansen P. 2003. Fate of stocked  
811 trout *Salmo trutta* L. in Danish streams: Survival and exploitation  
812 of stocked and wild trout by anglers. *International Journal of*  
813 *Ecohydrology and Hydrobiology*, 3(1), 39-50.

814 Pedersen LF, Koed A, Malte H. 2008. Swimming performance of wild  
815 and Fl-hatchery-reared Atlantic salmon (*Salmo salar*) and brown  
816 trout (*Salmo trutta*) smolts. *Ecology of Freshwater Fish*, 17(3),  
817 425-431. <https://doi.org/10.1111/j.1600-0633.2008.00293.x>

818 Pinter K, Weiss S, Lautsch E, Unfer G. 2018. Survival and growth  
819 of hatchery and wild brown trout (*Salmo trutta*) parr in three  
820 Austrian headwater streams. *Ecology of Freshwater Fish*, 27(1),  
821 146-157. <https://doi.org/10.1111/eff.12332>

822 Polgar G, Iaia M, Righi T, Volta P. 2022. The Italian Alpine and  
823 Subalpine trouts: Taxonomy, Evolution, and Conservation. *Biology*,  
824 11(4), 576. <https://doi.org/10.3390/biology11040576>

825 R Core Team. 2021. R: A language and environment for statistical  
826 computing. R Foundation for Statistical Computing, Vienna,  
827 Austria. Available at <https://www.R-project.org/>.

828 Sánchez-González JR, Nicieza AG. 2021. Individual differences in  
829 dominance-related traits drive dispersal and settlement in  
830 hatchery-reared juvenile brown trout. *Scientific reports*, 11(1),  
831 1-14. <https://doi.org/10.1038/s41598-021-86613-4>

832 Schwartz JS, Herricks EE. 2005. Fish use of stage-specific fluvial  
833 habitats as refuge patches during a flood in a low-gradient  
834 Illinois stream. *Canadian Journal of Fisheries and Aquatic*  
835 *Sciences*, 62(7), 1540-1552. <https://doi.org/10.1139/f05-060>

836 Schwinn M, Baktoft H, Aarestrup K, Koed A. 2017. A comparison of  
837 the survival and migration of wild and F1-hatchery-reared brown  
838 trout (*Salmo trutta*) smolts traversing an artificial lake.  
839 *Fisheries Research*, 196, 47-55.  
840 <https://doi.org/10.1016/j.fishres.2017.08.011>

841 Thorpe JE, Ross LG, Struthers G, Watts W. 1981. Tracking Atlantic  
842 salmon smolts, *Salmo salar* L., through Loch Voil, Scotland. *Journal*  
843 *of Fish Biology*, 19(5), 519-537. [https://doi.org/10.1111/j.1095-](https://doi.org/10.1111/j.1095-8649.1981.tb03820.x)  
844 [8649.1981.tb03820.x](https://doi.org/10.1111/j.1095-8649.1981.tb03820.x)

845 Thorstad EB, Whoriskey F, Rikardsen AH, Aarestrup K. 2011. Aquatic  
846 nomads: the life and migrations of the Atlantic salmon. *Atlantic*  
847 *salmon ecology*, 1(6), 1-32.  
848 <https://doi.org/10.1002/9781444327755.ch1>

849 Trapletti A, Hornik K, LeBaron B, Hornik MK. 2015. Package  
850 'tseries'.

851 Tytler P, Thorpe JE, Shearer WM. 1978. Ultrasonic tracking of the  
852 movements of Atlantic salmon smolts (*Salmo salar* L) in the  
853 estuaries of two Scottish rivers. *Journal of Fish Biology*, 12(6),  
854 575-586. <https://doi.org/10.1111/j.1095-8649.1978.tb04204.x>

855 Wallace BC. 2010. Trout population and production dynamics in North  
856 Carolina state park streams. Master's thesis. North Carolina State  
857 University, Raleigh.

858 Waples RS. 1999. Dispelling some myths about hatcheries.  
859 *Fisheries*, 24(2), 12-21. [https://doi.org/10.1577/1548-](https://doi.org/10.1577/1548-8446(1999)024<0012:DSMAH>2.0.CO;2)  
860 [8446\(1999\)024<0012:DSMAH>2.0.CO;2](https://doi.org/10.1577/1548-8446(1999)024<0012:DSMAH>2.0.CO;2)

861 Wiley RW, Whaley RA, Satake JB, Fowden M. 1993. An evaluation of  
862 the potential for training trout in hatcheries to increase  
863 poststocking survival in streams. *North American Journal of*  
864 *Fisheries Management*, 13(1), 171-177.  
865 [https://doi.org/10.1577/15488675\(1993\)013<0171:AEOTPF>2.3.CO;2](https://doi.org/10.1577/15488675(1993)013<0171:AEOTPF>2.3.CO;2)

866 Young RG, Hayes JW, Wilkinson J, Hay J. 2010. Movement and  
867 mortality of adult brown trout in the Motupiko River, New Zealand:  
868 effects of water temperature, flow, and flooding. *Transactions of*  
869 *the American Fisheries Society*, 139(1), 137-146.  
870 <https://doi.org/10.1577/T08-148.1>

871 Zentner DL, Wolf SL, Brewer SK, Shoup DE. 2021. A Review of Factors  
 872 Affecting PIT Tag Detection Using Mobile Arrays and Use of Mobile  
 873 Antennas to Detect PIT-Tagged Suckers in a Wadeable Ozark Stream.  
 874 *North American Journal of Fisheries Management*.  
 875 <https://doi.org/10.1002/nafm.10578>


876 Zydlewski GB, Haro A, Whalen KG, McCormick SD. 2001. Performance  
 877 of stationary and portable passive transponder detection systems  
 878 for monitoring of fish movements. *Journal of Fish Biology*, 58(5),  
 879 1471-1475. <https://doi.org/10.1111/j.1095-8649.2001.tb02302.x>

880 Zydlewski GB, Horton G, Dubreuil T, Letcher B, Casey S, Zydlewski  
 881 J. 2006. Remote monitoring of fish in small streams: a unified  
 882 approach using PIT tags. *Fisheries*, 31(10), 492-502.  
 883 [https://doi.org/10.1577/1548-8446\(2006\)31\[492:RMOFIS\]2.0.CO;2](https://doi.org/10.1577/1548-8446(2006)31[492:RMOFIS]2.0.CO;2)

# Figure 1

## Study area


Figure 1: Study area. Black triangles are the four stations where marked hatchery trout were released.



# Figure 2

## NFDD upstream & Water discharge

Figure 2. Trend comparison between Number of fish detected daily (NFDD) released upstream the antenna and daily water discharge measured in Laveggio creek between October 2020 and April 2022. Dashed areas are the periods in which the antenna did not work.






# Figure 3

## NFDD downstream & Water discharge


Figure 3. Trend comparison between Number of fish detected daily (NFDD) released downstream the antenna and daily water discharge measured in Laveggio creek between October 2020 and April 2022. Dashed areas are the periods in which the antenna did not work.



# Figure 4

## NFDD Wild & Water discharge

Figure 4. Trend comparison between Number of wild fish detected daily (NFDD) and daily water discharge measured in Laveggio creek between April 2021 and April 2022. Dashed areas are the periods in which the antenna did not work.






# Figure 5

NFDD upstream vs Water discharge


Figure 5: Cross-Correlation (CC) analysis between NFDD released upstream and water discharge.



# Figure 6

NFDD downstream vs Water discharge

Figure 6: Cross-Correlation (CC) analysis between NFDD released downstream and water discharge.



# **Table 1**(on next page)

Table 1: Results of the generalized linear model analysis using the generalized estimating equation (GEE), performed between NFDD released upstream and water discharge.

<b>Analysis of GEE parameter estimates</b>			
	Estimate	Std. Error	<i>P</i> -value
(Intercept)	0.227	0.105	0.031*
Water discharge	0.33	0.141	0.02*

1

2

## Table 2 (on next page)

Table 2: Results of the generalized linear model analysis using the generalized estimating equation (GEE), performed between NFDD released downstream and water discharge.

<b>Analysis of GEE parameter estimates</b>			
	Estimate	Std. Error	<i>P</i> -value
(Intercept)	0.1961	0.0389	0.00000046*
Water discharge	0.0245	0.0232	0.29

1