

Planting *Cyperus esculentus* augments soil microbial biomass and diversity, but not enzymatic activities

Yulong Duan¹, Wei Ren², Jianhua Zhao³, Chun Luo⁴, Yang Liu⁵

¹ Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou, Gansu, China

² Naiman Desertification Research Station, Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Tongliao, China

³ Agricultural Biotechnology Institute, Jilin Academy of Agricultural Sciences, Changchun, China

⁴ Shanghai Majorbio Bio-pharm Technology Co., Ltd, Shanghai, China

⁵ Gansu Institute of Architectural Design and Research Company, Lanzhou, Gansu, China

Corresponding Author: Yulong Duan

Email address: duanyulong@nieer.ac.cn

The planting of *Cyperus esculentus*, a member of the grass family Cyperaceae which includes nut sedge weeds, is being increasingly promoted in northern China's semi-arid and arid regions. Yet the effects of planting *C. esculentus* upon soil quality and soil microbial characteristics of sandy land remain unclear. This study examined the short-term (1 year) impact of this grass species on soil microbial biomass indices, enzymatic activities, and microbiome characteristics in the Horqin Sandy Land area of China. The results show that planting *C. esculentus* could increase microbial biomass in the form of carbon (MBC), nitrogen (MBN), and phosphorus (MBP), but it negligibly influenced the enzymatic activities of soil β -1,4-glucosidase (BG), cellobiohydrolase (CBH), leucine aminopeptidase (LAP), and β -1,4-N-acetaminoglycosidase (NAG). Over 1 year, we found that planting *C. esculentus* significantly increased the soil bacterial richness and diversity of sandy land, yet also altered community composition of soil bacteria and eukaryotes in way that could promote their homogenization. In this respect, the relative abundances of *Acidobacteria* and *Proteobacteria* significantly decreased and increased, respectively; hence, they may be considered for use as important indicators of soil nutrient-rich conditions. Overall, the results could be explained by greater soil organic carbon (SOC) and total nitrogen (TN), mainly derived from cumulative plant litter input to soils, which then increased the sandy soil's C:N ratio. Future research should focus on exploring the long-term effects of planting *C. esculentus* on soil quality and soil microbial characteristics of sandy lands in China and abroad.

Planting *Cyperus esculentus* augments soil microbial biomass and diversity, but not enzymatic activities

Yulong Duan^{a, b, #, *}, Wei Ren^{c, #}, Jianhua Zhao^d, Chun Luo^d, Yang Liu^e

^a Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou 730000, China.

^b Naiman Desertification Research Station, Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Tongliao 028300, China.

^c Agricultural Biotechnology Institute, Jilin Academy of Agricultural Sciences, Changchun 130033, China.

^d Shanghai Majorbio Bio-pharm Technology Co., Ltd, Shanghai 200120, China.

^e Gansu Institute of Architectural Design and Research Company, Lanzhou 730000, China.

[#] Co-first authors who contributed equally to this work (Y. Duan and W. Ren).

***Corresponding author** Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, 320 Donggang West Road, Lanzhou 730000, China (Y. Duan).

E-mail address: duanyulong@nieer.ac.cn (Y. Duan); renwei@cjaas.com (W. Ren).

Abstract

The planting of *Cyperus esculentus*, a member of the grass family *Cyperaceae* which includes nut sedge weeds, is being increasingly promoted in northern China's semi-arid and arid regions. Yet the effects of planting *C. esculentus* upon soil quality and soil microbial characteristics of sandy land remain unclear. This study examined the short-term (1 year) impact of this grass species on soil microbial biomass indices, enzymatic activities, and microbiome characteristics in the Horqin Sandy Land area of China. The results show that planting *C. esculentus* could increase microbial biomass in the form of carbon (MBC), nitrogen (MBN), and phosphorus (MBP), but it negligibly influenced the enzymatic activities of soil β -1,4-glucosidase (BG), cellobiohydrolase (CBH), leucine aminopeptidase (LAP), and β -1,4-N-acetaminoglycosidase (NAG). Over 1 year, we found that planting *C. esculentus* significantly increased the soil bacterial richness and diversity of sandy land, yet also altered community composition of soil bacteria and eukaryotes in way that could promote their homogenization. In this respect, the relative abundances of *Acidobacteria* and *Proteobacteria* significantly decreased and increased, respectively; hence, they may be considered for use as important indicators of soil nutrient-rich conditions. Overall, the results could be explained by greater soil organic carbon (SOC) and total nitrogen (TN), mainly derived from cumulative plant litter input to soils, which then increased the sandy soil's C:N ratio. Future research should focus on exploring the long-term effects of planting *C. esculentus* on soil quality and soil microbial characteristics of sandy lands in China and abroad.

Keywords: *Cyperus esculentus*, soil microorganisms, soil organic carbon, soil enzymatic activities, semi-arid and arid regions

1. Introduction

Cyperus esculentus, also known as chufa sedge, yellow nut sedge, tiger nut sedge, or earth almond, belongs to the grass family *Cyperaceae*. This plant species is a perennial weed native to the Mediterranean region, but has since spread globally, in part because of its well-developed root system, strong tillering ability, short growing period, large biomass, high oil content, strong resistance to stress, and a wide range of adaptive traits. Nevertheless, the tubers of *C. esculentus* are valued for their health and nutritive benefits (Asare et al., 2020; Oderinde and Tairu, 1988), and in recent years, the distribution of introduced *C. esculentus* has extended considerably in the semi-arid and arid regions of northern China. There, the total cultivated area of *C. esculentus* reached 6886 km² in 2021, mainly in the Inner Mongolia Autonomous Region, Xinjiang Uygur Autonomous Region, Liaoning Province, and Jilin Province, whose coverage is expected to continue expanding (Sha et al., 2022; Zhang and Wu, 2022). In these regions, *C. esculentus* is generally grown on sandy soil or sandy loam to conveniently harvest its valued tuberous roots, given the typically loose texture of sandy land. Surprisingly, however, the effects of planting *C. esculentus* on the soil quality of sandy land and its soil microbial characteristic are still poorly understood.

Soil microbes are indispensable components of terrestrial ecosystems and their biological characteristics, including diversity, community composition, biomass, quantity, and enzymatic activity, are highly sensitive to changing environmental conditions (Hu et al., 2019; Lange et al., 2015; Neilson et al., 2012; Yu et al., 2022). A meta-analysis of 81 available studies from 28 countries indicates that plant crops enhance soil microbial diversity and biomass, and altered community structure compared to no plant crop, and the responses of microbial parameters to plant crop varied with soil and climatic conditions (Muhammad et al., 2021). Actually, such soil microbial biological characteristics can more directly engage in feedback with altered soil quality than can physicochemical properties such as soil organic matter (SOC) and soil nutrient content (Bünemann et al., 2018; Caldwell, 2005). Microbial biomass, namely microbial biomass in terms of carbon (MBC), nitrogen (MBN), and phosphorus (MBP) is the main driving force behind the transformation and cycling of soil organic matter and nutrients (N and P); it is also functions as a crucial reserve pool for soil nutrients to provision plants' growth, and can thus serve as an indicator of soil fertility (Jenkinson and Ladd, 1981; Pan et al., 2018; Powlson et al.,

2001; Sparling et al., 1997). Soil extracellular enzymes are special secondary metabolites secreted by microorganisms to obtain energy and nutrients from soil. Accordingly, changed extracellular enzymes activities may reflect the metabolic status of a soil microbe taxon or community as well as the potential for specific soil biochemical reactions (Dirks et al., 2012; Ma et al., 2022). Notably, in response to sudden changes in environmental conditions, soil extracellular enzymes are often expressed and released, such as β -1,4-glucosidase (BG), β -1,4-N-acetylglucosaminidase (NAG) and leucine aminopeptidase (LAP), acid phosphatase (ACP); these have been used to successfully characterize the cycling dynamics of carbon (C), nitrogen (N), and phosphorus (P) in soil, respectively (Allison and Jastrow, 2006; Brunner et al., 2003; Shao et al., 2020). In particular, cellobiohydrolase (CBH) are a class of key enzymes functioning in cellulose hydrolysis and biomass conversion, which may be used to characterize the decomposition capacity of cellulose material (Haan et al., 2013).

The sandy land of semi-arid and arid ecosystems is generally characterized by low microbial biomass indices, enzymatic activities, and α -diversity, these features usually ascribed to its low soil moisture and poor availability of limiting nutrients. Therefore, we hypothesized that planting *C. esculentus* followed by its growth (cultivation) for 1 year would increase the soil microbial biomass, richness, and diversity. Our study's findings provide timely scientific data to inform future strategies for planting *C. esculentus* on sandy soils in the semi-arid areas of northern China.

2. Materials and methods

2.1 Site description and soil sampling

The two sampling sites were established, one on bare sandy land (BSL) and the other on sandy land in which *C. esculentus* was planted and cultivated for 1 year (CE) (Fig. S1). Both experimental sites were located in Naiman, Tongliao, of Inner Mongolia, in a semi-arid region of China's Horqin Sandy Land (42°55'N, 120°42'E; 348 m a.s.l.). This region has a continental semi-arid monsoon temperate climate, with a mean annual temperature of 6.8°C and a mean annual precipitation of 360 mm, 70% of which occurs from May through September. The dominant native plant species of the sandy land were *Caragana microphylla*, *Setaria viridis* and *Echinops gmelinii*, with a low vegetation cover of 30%. The distance

between these two sites is less than 500 m and they were freshly reclaimed in December 2020. The soil type is degraded sandy chestnut soil and aeolian sandy soil, and the topsoil (depth: 0–20 cm) was taken for nutrient determination before reclamation. The soil organic carbon content was $2.61 \text{ g}\cdot\text{kg}^{-1}$, soil total nitrogen content was $0.20 \text{ g}\cdot\text{kg}^{-1}$, the available phosphorus content was $0.71 \text{ g}\cdot\text{kg}^{-1}$. It is characterized by high sand content, low nutrient content and poor water and fertilizer retention.

In this study, *C. esculentus* were sown by mechanical seeding method in mid-May. To control for the disturbance caused by fertilization and irrigation, both sites received the same fertilization and irrigation application (urea, $675 \text{ kg}\cdot\text{hm}^{-2}$; $(\text{NH}_4)_2\text{HPO}_4$, $300 \text{ kg}\cdot\text{hm}^{-2}$; K_2SO_4 , $375 \text{ kg}\cdot\text{hm}^{-2}$; humid acid, $300 \text{ kg}\cdot\text{hm}^{-2}$; total irrigation volume was $5800 \text{ m}^3\cdot\text{hm}^{-2}$). Of these, the base fertilizer applied before planting (urea, $45 \text{ kg}\cdot\text{hm}^{-2}$; $(\text{NH}_4)_2\text{HPO}_4$, $300 \text{ kg}\cdot\text{hm}^{-2}$; K_2SO_4 , $375 \text{ kg}\cdot\text{hm}^{-2}$; humid acid $300\text{kg}\cdot\text{hm}^{-2}$ and the irrigation amount before sowing was $2025 \text{ m}^3\cdot\text{hm}^{-2}$. After that, all treatments adopted the water-fertilizer integration technology for trickle irrigation and fertilization, and the trickle irrigation belt was laid along the planting direction of *C. esculentus*, with an interval of 30 cm for each trickle irrigation belt. At the initial stage of growth (from May to June), water was dropped 4 times, each time $187.5 \text{ m}^3\cdot\text{hm}^{-2}$, each hectare 33.75 kg urea and 15 kg K_2SO_4 . In the middle irrigation period (from June to mid-August), 12 drops were applied, and in the late irrigation period (from mid-August to mid-September), 4 drops were applied, with the same amount of water and fertilizer per drop as in the initial irrigation period.

A total of 10 plots were set up, evenly split between the *BSL* and *CE* sites (i.e., 5 plots per site), and each plot randomly distributed on the *BSL* and *CE sites* (the distance between the plots is ca. 50 m). An auger (2.5-cm diameter) was used to collect topsoil samples (depth: 0–20 cm) on 28 August 2021, this corresponding to the peak of the growing season at *CE*. Within each plot ($1 \text{ m} \times 1 \text{ m}$), five random locations were sampled and mixed together to generate a single composite soil sample; all such samples were then individually packed in sterilized polyethylene bags and taken to the laboratory in coolers (portable car refrigerators) as quickly as possible. Any surface litter in the soil samples was removed by passing them through a 2-mm square-aperture mesh. Each composite soil sample was divided into two subsamples: one was stored at 4°C to later determine the soil properties, and the other was stored at -80°C until the DNA extractions.

2.2 Measurement of soil microbial biomass carbon, nitrogen, and phosphorus contents

A routine fumigation-extraction methodology was employed to measure the microbial biomass carbon (MBC), nitrogen (MBN), and phosphorus (MBP) (Joergensen, 1996). Briefly, triplicate fresh 50-g soil subsamples were placed separately in 100-mL beakers and incubated in the dark for 7 days at 25°C with a relative humidity (RH) of 70%. The first subsample was set as the control group and the second one fumigated for 24 h with ethanol-free CHCl_3 ; the third one was used to quantify the soil moisture content. The control and fumigated samples were transferred into 250-mL Erlenmeyer flasks, to which 100 mL of 0.5 M K_2SO_4 (for MBC and MBN) or NaHCO_3 (for MBP) was added to each, and the solution shaken for 30 min at 25°C to obtain the soil extracts. These extracts were filtered through 0.45- μm cellulose filters and stored at -20°C until their analysis. The MBC, MBN, and MBP contents were all measured using an Elementar Vario TOC (Elementar, Langenselbold, Germany). The MBC, MBN, and MBP were respectively calculated as the difference between the extractable C, N, and P contents of fumigated and control samples, using fixed conversion factors: 0.45 for both the K_{EC} of C and K_{EN} of N, while the K_{EP} for P was 0.40.

2.3 Measurement of five types of soil enzymatic activity

For soil enzymes β -1,4-glucosidase (BG), cellobiohydrolase (CBH), leucine aminopeptidase (LAP), β -1, 4-N-acetaminoglycosidase (NAG), and acid phosphatase (ACP), their respective activity was quantified using commercial enzyme kits and following the manufacturer's protocol (Solarbio, Beijing, China). Briefly, BG decomposes p-nitrobenzene- β -D-glucopyranoside, yielding p-nitrophenol; CBH decomposes nitrobenzene fibrodiglycoside (PNPC), yielding p-nitrophenol; LAP decomposes L-leucine-p-nitroaniline, yielding p-nitroaniline; NAG decomposes p-nitrobenzene β -N-acetylglucosamine, yielding p-nitrophenol; and ACP decomposes disodium phenyl phosphate hydrate to yield phenol and Na_2HPO_4 . The maximum absorption peak for BG, CBH, or NAG occurs at 400 nm, for LAP at 405 nm, and for ACP at 660 nm; to measure absorbance, a UV-VIS spectrophotometer was used (UV-1800, Mapada Instruments Co., Shanghai, China).

149

150 **2.4 Soil DNA extraction and quantitative PCR**

151 Soil total DNA was extracted from a 0.5-g subsample using the PowerSoil kit (Omega Laboratories
152 Inc., Mogadore, OH, USA) according the manufacturer's instructions. The integrity of DNA was checked
153 by electrophoresis in 1.0% agarose gel, and its purity and concentration measured spectrophotometrically
154 by a NanoDrop ND5000 (Thermo Fisher Scientific Inc., USA). Next, PCR was conducted in a Line Gene
155 9600 Plus Cycler (Thermo Fisher Scientific Inc.) to amplify the hyper-variable 460-bp V3–V4 region of
156 the 16s rRNA, using the primer pair Eub338/806 (338F, 5'-ACTCCTACGGGAGGCAGCAG-3' (Dennis
157 et al., 2013); 806R, 5'-GGACTACHVGGGTWTCTAAT-3'). Likewise, the 300-bp V5–V7 region of 18
158 rRNA was amplified using the primer pair SSU0817/1196 (0817F, 5'-
159 TTAGCATGGAATAATRAATAGGA-3'; 1196R, 5'-TCTGGACCTGGTGAGTTTCC-3') (Rousk et al.,
160 2010).

161 Real-time quantitative PCR (*qPCR*) was carried out with 25 mg of the DNA mixed with 12.5 mL of
162 ChamQ SYBR color *qPCR* Master Mix (2x) (Vazyme Biotech Co., Ltd, Nanjing, China), 0.5 mL of a
163 buffer solution (10 mM) of the primer pair, and 9.5 mL of dd H₂O. Standard and environmental DNA
164 samples were added at 2.0 mL per reaction. Every reaction was run on an ABI7300 unit (Applied
165 Biosystems, USA) with this cycling program: initial denaturation at 95°C for 3 min, followed by 40
166 cycles of 94°C for 30 s, 53°C for 30 s, and 72°C for 45 s, with a final extension at 72°C for 5 min (Niu et
167 al., 2021). Melting curve and gel electrophoresis analyses were used to confirm that the amplified
168 products were of the appropriate size. Bacterial and fungal gene copy numbers were generated using a
169 regression equation for each assay that related the cycle threshold (*CT*) value to a known number of
170 copies in a given standard. All *qPCR* reactions were run in triplicate for each soil sample. The average
171 bacterial PCR efficiency was 101.00%, with an R^2 of 0.9992 for the standard curves; the fungal PCR
172 efficiency was 105.56% with an R^2 of 0.9999 for the standard curves.

173

174 **2.5 High-throughput amplicon sequencing bioinformatics analysis**

175 Each PCR was carried out in triplicate, in a 20-μL reaction volume that contained 4 μL of a 5×

reaction buffer, 4 μ L of dNTPs (2.5 mM), 0.8 μ L of each primer (5 μ M), 1 μ L of template DNA (ca. 10 ng), and 0.4 μ L of Pfu DNA Polymerase (TransStart-FastPfu DNA Polymerase, TransGen Biotech, Beijing, China), topped up with double-distilled H₂O. The PCR program consisted of an initial denaturation phase at 95°C for 3 min; followed by 35 cycles at 94°C for 30 s, annealing at 55°C for 30 s, and an extension at 70°C for 45 s; a final extension was done at 72°C for 10 min. The PCR was performed in an ABI GeneAmp 9700 Cyclor (Thermo Fisher Scientific Inc.).

Unique barcode sequences were added at the 5' end of the forward primer, to separate corresponding reads from the data pool generated in a single sequencing run. The ensuing amplicons were extracted by electrophoresis in 2.0% agarose gel and purified with a Gel Extraction Kit (Axygen Co., Hangzhou, China), according to the manufacturer's instructions, then quantified with a QuantiFluor-ST fluorimeter (Promega, Fitchburg, WI, USA). The purified amplicons were pooled in an equimolar way and paired-end sequenced (2 \times 300 bp) by Majorbio Co. Ltd. (Shanghai, China), according to recognized protocols.

Raw FASTQ files were de-multiplexed and quality-filtered, by using the 'fastp' tool (v.0.20.0) (Chen et al., 2018) according to these criteria: (i) The 300-bp reads were truncated at any site that had an average quality score < 20 over a 50-bp sliding window, discarding any truncated reads shorter than 50-bp; (ii) Extracted matching barcodes were removed, and as well as any two-nucleotide mismatches vis-à-vis the primer, or reads that contained ambiguous characters; (iii) Overlapping sequences > 10 bp in length were assembled according to their overlapping sequence. Reads that could not be assembled were discarded.

The resulting quality-checked sequences were aligned as per the SILVA alignment database (<https://www.arb-silva.de>) (Quast et al., 2013), and clustered into operational taxonomic units (OTUs) by USEARCH software (v. 7.1) (<https://www.drive5.com/usearch/>) (Edgar, 2013). Only those OTUs with at least a 97% similarity level were used to draw the rarefaction curves, and the α -diversity indices—Chao, ACE, Shannon, and Simpson indices (<http://www.mothur.org/wiki/Calculators>), and their performed coverage analysis—were calculated using Mothur software (v1.30.2) (<https://www.mothur.org>) (Schloss et al., 2009). The taxonomic assignment of OTUs at a 97% similarity cut-off was conducted by Mothur, with a 70% confidence interval, in accordance with the SILVA (138) or Unite (8.0) database

(<http://unite.ut.ee/index.php>) for bacteria and eukaryota, respectively.

All sequencing data associated with this study can be found in the NCBI Sequence Read Archive (<https://www.ncbi.nlm.nih.gov/sra>), under project accession no. PRJNA836307.

2.6 Statistical analysis

Significant differences in the means of soil microbial biomass indices, enzyme activity, quantity, and α -diversity between two sites (*BSL* vs. *CE*) were determined by one-way analysis of variance (ANOVA). All analyses were implemented in SPSS statistical software (v. 19.0) (<https://www.ibm.com/analytics/spss-statistics-software>). Unless otherwise noted, a difference between means was considered statistically significant at $P < 0.05$. Principal co-ordinates analysis (PCoA) was used to uncover differences in β -diversity between samples of the two sites (*BSL* vs. *CE*). Mantel tests were used to evaluate the relationships of the bacterial or eukaryotic community composition to measure site-level soil variables. Pearson correlation coefficient (r) were used to indicate a positive correlation ($r > 0$) or a negative correlation ($r < 0$). Both PCoA and Mantel tests were carried out in the R platform (v3.2.1) (www.r-project.org) with its ‘vegan’ package. This package was also used to build the matrices for the pairwise taxonomic distances between the bacterial or eukaryotic communities (Bray–Curtis’s dissimilarity) and environmental variables (Euclidean distance). Pearson correlation analysis was carried out in the R platform (v3.2.1) (www.r-project.org) with its function `cor()`.

3. Results

3.1 Comparative analysis of soil microbial biomass indices and enzymatic activities

The soil biomass microbial indices for the two sites are shown in Fig. 1A. Evidently, the mean values for MBC, MBN, and MBP were one-third to three times greater at *CE* than *BSL*. ($P < 0.05$) (Table S1, Table S3). For the soil MBC:MBN ratio, its mean (\pm SD) was 18.293 ± 6.868 at *BSL*, which was significantly higher than the 8.324 ± 4.700 at *CE* ($P < 0.05$) (Table S1, Table S3). In Fig. 1B are the soil enzymatic activities (BG, CBH, LAP, NAG, and ACP) for the two sites. In this respect, only ACP differed significantly between *BSL* and *CE*, being almost 20% lower in the latter ($P < 0.05$) (Table S2,

Table S3).

3.2 Microbial diversity characterization and community composition of the two sites

Gene copy numbers for total bacteria (16S RNA) and eukaryota (18S RNA) species at the two sampling sites are presented in Fig 2. For bacteria, copy abundance was nearly two times higher in the CE than BSL soil ($P < 0.05$), while the eukaryota were similarly abundant between the two sites ($P < 0.05$). Fig. 3 shows the two α -diversity indexes, community species richness (Chao 1 index) and community diversity (Shannon index), at the two sites. For bacteria, the Chao1 index ranged from 2329.50 to 2677.07 and 2690.12 to 2799.07 at BSL and CE, with means of ca. 2500 and 2700, respectively (Table S4), while the Shannon index ranged from 5.76 to 6.45 and 6.33 to 6.42 respectively at BSL and CE, also with similar means (Table S4). For eukaryota, the Chao1 index ranged from 73.00 to 110.5 and 83.88 to 117.6 at BSL and CE, with means of ca. 90 and 97, respectively (Table S5); corresponding values for the Shannon index were 1.10 to 2.73 and 2.16 to 2.55, with nearly identical means (Table S5). The plot-level ACE and Simpson indices for the diversity of bacteria and eukaryota are summarized in Tables S4 and S5, respectively.

Across all 10 soil samples, a total of 272 464 valid sequences and 4 379 OTUs were obtained. All sequences could be assigned to one of 36 bacterial phyla. Of these, the 19 most dominant bacterial phyla across all samples were *Actinobacteria* (8.01%–12.64%), *Acidobacteria* (5.90%–17.92%), *Proteobacteria* (7.21%–14.38%), *Chloroflexi* (6.55%–12.26%), *Gemmatimonadota* (6.05%–13.09%), *Firmicutes* (6.96%–12.07%), *Bacteroidetes* (7.17%–12.39%), *Myxococcota* (7.46%–13.81%), *Cyanobacteria* (0.69%–32.44%), *Methylomirabilota* (2.84%–18.56%), *Nitrospirae* (5.88%–16.48%), *Bacteria_unclassified* or *norank* (4.46%–15.02%), *Patescibacteria* (3.01%–15.39%), *Planctomycetota* (6.45%–14.72%), *Verrucomicrobiota* (3.77%–14.87%), *Bdellovibrionota* (6.34%–16.28%), *Armatimonadota* (4.52%–16.96%), RCP2-54 (2.35%–22.15%), and *Entotheonellaeota* (4.27%–21.35%), which together accounted for ca. 99.55% of all bacterial sequences (Fig. 4; Table S6). The remaining phyla only comprising < 1% of the total bacterial library

For eukaryota, 350 670 valid sequences and 202 OTUs were obtained. All sequences could be

assigned to one of 22 eukaryota phyla. Of these, the 12 most dominant eukaryota phyla across all samples were *Ascomycota* (59.31%–96.30%), *Basidiomycota* (1.68%–35.14%), *SAR_norank* (0.81%–7.38%), *Mucoromycota* (0.29%–1.71%), *Fungi_norank* (0.01%–9.50%), *Eukaryota_norank* (0.08%–3.04%), *Schizoplasmodiida* (0.03%–1.10%), *Zoopagomycota* (0%–1.01%), *Fungi_unclassified* (0.03%–0.90%), *Eukaryota_unclassified* (0.03%–0.25%), *Amoebozoa_norank* (0%–0.36%), and *Chytridiomycota* (0%–0.46%). Among them, *Ascomycota*, *Basidiomycota*, *Mucoromycota*, *Fungi_norank*, *Zoopagomycota*, *Fungi_unclassified*, and *Chytridiomycota* belonged to the fungi and constituted 88.44%–98.71% of the total eukaryota library.

3.3 Bacterial and eukaryota community composition at the two sites

The PCoA done at the OTU level was used to infer the β -diversity patterning of soil bacterial and eukaryotic community composition (Fig. 5). Bacterial communities of the *CE* samples strongly clustered at the far right of the PC1 axis (explaining 52.96% of the total variation), unlike the *BSL* samples which were well scattered along the entire PC2 axis (accounting 11.32% of the total variation). Likewise, for eukaryota communities, those of the *CE* samples were clumped along the right of PC1 axis (accounting for 34.49% of the total variation), whereas those of the *BSL* samples were interspersed along the length of the PC2 axis (accounting for 11.32 % of the total variation).

The student's t-test bar plots indicated differences between the two sites (*BSL* vs. *CE* below, respectively) in their bacterial and eukaryotic communities at the phylum level (Fig. 6). A total of 13 bacterial phyla, namely *Acidobacteria* (30.06% vs. 16.96%), *Proteobacteria* (15.23% vs. 23.6%), *Chloroflexi* (12.50% vs. 6.21%), *Cyanobacteria* (2.13% vs. 0.20%), *Nitrospirota* (0.48% vs. 0.73%), *Bacteria_unclassified* (0.43% vs. 0.69%), *Planctomycetota* (0.43% vs. 0.29%), *Deinococcota* (0.03% vs. 0.07%), *NB1-j* (0.008% vs. 0.087%), *WS2* (0.064% vs. 0.0194%), *GAL15* (0.0055% vs. 0.048% at *CE*), *SAR324_cladeMarine_group_B* (0.00526% vs. 0.02216%), and *Dadabacteria* (0% vs. 0.023%) differed significantly in their relative abundance between the *BSL* and *CE* sites (Fig. 6; Table S8). For eukaryota, only *Eukaryota_unclassified* (0.075% vs. 0.183%), *Amoebozoa_norank* (0.018% vs. 0.232%), and *Aphelidea* (0% vs. 0.060%), occurred in significantly different proportions between the two sites (Fig.

6; Table S9).

Mantel tests were used to discern the several environmental factors that shaped the soil bacterial and eukaryota community composition across all 10 samples (Fig. 7). For bacteria, SOC ($R = 0.808$, $P = 0.008$) was the most critical environmental factor that individually explained the variation in soil bacterial community composition, followed by soil pH ($R = 0.808$, $P = 0.008$), soil TN ($R = 0.769$, $P = 0.005$), soil EC ($R = 0.710$, $P = 0.006$), soil C:N ratio ($R = 0.605$, $P = 0.004$), and MBN ($R = 0.624$, $P = 0.009$) (Fig. 7; Table S10). The soil eukaryota community composition was most associated with SOC ($R = 0.397$, $P = 0.003$), soil pH ($R = 0.341$, $P = 0.015$), soil TN ($R = 0.298$, $P = 0.013$), MBN ($R = 0.271$, $P = 0.031$), and soil EC ($R = 0.236$, $P = 0.041$) (Fig. 7; Table S10).

4. Discussion

4.1 Effects of planting *C. esculentus* on soil microbial biomass indices and enzymatic activities

Our results showed that the soil MBC, MBN, and MBP values of planted *C. esculentus* cultivated for 1 year were all higher than those of the control site (BSL), and significantly so for MBC and MBN (Fig. 1A; Table S3). This result is probably due to the significantly greater soil nutrient contents, especially for SOC and TN, that arose in the presence of *C. esculentus* over 1 year (Table S3), likely driven by accumulating organic matter derived from litter decomposition and root exudates. Generally, the sandy soil in semi-arid and arid ecosystems are considered oligotrophic environments with poor nutrient status. (Zhao et al., 2016). Although there are some exogenous nutrient inputs, the sandy soil still cannot accumulate nutrients effectively because of its loosened soil texture that is also prone to leaching. Evidently, planting and then cultivating *C. esculentus* for 1 year could improve soil nutrient conditions of sandy land. An earlier study in arid and semi-arid grassland of China showed that the total biomass of soil microorganisms, as well as bacterial or fungal biomass, were significantly positively correlated with the SOC content (Hu et al., 2014). In other work, soil bacterial diversity was also positively correlated with the TN content (Wang et al., 2015). Generally, the main factor responsible was larger concentrations of key soil nutrients, especially organic matter and nitrogen, which was beneficial for the growth of soil microbial populations and microbial biomass (Ralte et al., 2005).

Compared with the control site, the four soil enzymatic activities (BG, CBH, LAP, and NAG) under planting with *C. esculentus* were hardly changed, in that only ACP was significantly decreased ($P < 0.05$) (Fig. 1B; Table S1). Soil enzymes have been considered as indicators of soil quality, indicating soil fertility quality and soil health quality. Nonetheless, distinct enzymes have different molecular compositions and structures, and catalyze different reactions, which are affected by multiple environmental factors, such as soil temperature, moisture, pH, fertilization, vegetation type, plant diversity, and community structure (Fu et al., 2012; Pan et al., 2018). Although the soil nutrient status of the sandy land was upgraded after the planting with *C. esculentus*, the change was not sufficient to cause significant changes in soil enzymatic activities (BG, CBH, LAP, and NAG), because the treatment time was only one year. By comparison, ACP are mainly produced by plant roots and microorganisms and play a key role in the P cycling, an often-limiting element in terrestrial ecosystems, and its activities were commonly positively correlated with soil organic matter, total nitrogen and organic phosphorus contents (Margalef et al., 2021). Therefore, our results are different from conventional studies, and the specific reason still needs to be confirmed through long-term positioning tests.

4.2 Effects of planting *C. esculentus* on soil microbial diversity and community composition

To compare the soil microbial characteristics of bare sandy land (BSL, the control) and sandy land in which *C. esculentus* was cultivated for 1 year (CE), real-time PCR and MiSeq high-throughput sequencing were used in tandem to reveal differences in the soil microbial abundance, α -diversity, and community composition between the two sampling sites.

The *q*PCR and sequencing results detected high bacterial abundance and diversity but a relatively low eukaryotic abundance and diversity at the two sampling sites. For microbial abundance, twice as many bacterial 16S rRNA gene copies were found in soil at CE than the control whereas the number of eukaryota 18S rRNA gene copies was quite similar between sites (Fig. 2). The great variation in bacterial 16S rRNA and eukaryota 18S rRNA gene copies and their site-level disparity are probably attributable to different feedback processes between bacteria and eukaryota and the soil C:N ratio, with a lower C:N ratio likely benefiting more the reproduction of bacteria than fungi (Hu et al., 2001).

Soil bacterial phyla were dominated by *Actinobacteria*, *Acidobacteria*, *Proteobacteria*, and *Chloroflexi*, while eukaryota phyla were dominated by *Ascomycota*, *Basidiomycota*, and SAR_norank (Fig. 4). These phyla are widely distributed in various terrestrial ecosystems, especially in arid and semi-arid regions (Baubin et al., 2019; Crits-Christoph et al., 2013; Sun et al., 2018; Zhang et al., 2019). The PCoA results uncovered stark differences in soil bacterial or eukaryotic community composition of bare sandy land across all its samples (Fig. 5), whereas either composition was very similar among samples where planted *C. esculentus* grew for 1 year. This pattern strongly implied that the short-term presence of *C. esculentus* could markedly promote the homogenization of soil bacterial and eukaryotic communities. Of these, the relative abundances of several bacterial and eukaryotic phyla distinctly changed: for bacteria, compared with the control, the relative abundance of *Acidobacteria* in response to *C. esculentus* planting fell sharply from 30.06% to 16.96%, and that of *Proteobacteria* and *Chloroflexi* increased sharply from 15.23% to 23.6% and from 12.50% to 16.21%, respectively (Fig. 6). Previous studies have shown that *Acidobacteria* are oligotrophic bacteria, being typically more abundant in low-nutrient soils, while *Proteobacteria* are copiotrophic bacteria usually found in nutrient-rich habitats (Fierer et al., 2007; Wang et al., 2017; Zeng et al., 2019). Accordingly, from our results, we may infer that the planted *C. esculentus* probably enriched the soil fertility of bare sandy soil. For eukaryota, compared with the control, the relative abundance of Eukaryota_unclassified in response to *C. esculentus* planting increased from 0.075% to 0.18%; likewise, the relative abundances of Amoebozoa_norank and *Aphelidea* respectively increased from 0.018% to 0.232% and from 0% to 0.060%.

Mantel tests indicated that soil microbial community composition was associated with multiple edaphic drivers whose roles differed marginally between the two microbial groups (Fig. 7; Table S10). For bacteria, SOC was the most fundamental environmental driver shaping the bacterial community composition across all soil samples, followed by the soil pH, TN, EC, C:N ratio, and MBN. Similarly, soil eukaryota community composition was mainly related to the SOC, pH, TN, MBN, and EC (Fig. 7, Table S10). A predominant role for soil pH in shaping bacterial community composition has been widely demonstrated at different scales and in different ecosystems (Jiao and Lu, 2020; Shen et al., 2013; Tian et al., 2018; Yu et al., 2021). Furthermore, soil nutrients' content is also a limiting factor of microbial

activity and can pivotally impact soil microbial community distribution patterns in arid and semi-arid regions (Chen et al., 2015). In our study, SOC strongly altered the community composition of both soil bacteria and eukaryota, which is consistent with other studies' findings (Hu et al., 2001; Tian et al., 2018).

5. Conclusion

In this study, we evaluated the short-term (1 year) effects of planted *C. esculentus* on the soil microbial biomass indices, enzymatic activities, and microbiome characteristics in the Horqin Sandy Land, China. The presence of *C. esculentus* for 1 year is capable of inducing the differential accumulation of SOC and TN, thereby increasing the soil C:N ratio. Furthermore, *C. esculentus* can promote three microbial biomass indices (MBC, MBN, and MBP) but it has very little impact on four enzymes' (BG, CBH, LAP, and NAG) activities. Moreover, planting with *C. esculentus* significantly increases the richness and diversity of soil bacteria rather than eukaryota in sandy land, while reducing and augmenting the relative abundance of *Acidobacteria* and *Proteobacteria*, respectively, suggesting their potential use as bioindicators of soil nutrient-rich conditions. Overall, our findings demonstrate that cultivating *C. esculentus* for just 1 year could improve the nutrient status and microbial characteristics of sandy soils in a semi-arid region.

Author Contributions

Sample collection: YLD.

Data curation: YLD, CL, JHZ.

Funding acquisition: YLD

Methodology: YLD, CL.

Project administration: YLD

Writing-original draft: YLD, WR, and YL.

Writing-review & editing: YLD, WR, and YL.

All authors contributed to the article and approved the submitted version.

391

392 **Funding**

393 This research work was supported by the National Key R&D Program of China (grant no.
394 2019YFC0507602) and the “Light of West China” Program of the Chinese Academy of Sciences
395 (E0290501).

396

397 **Data statement**

398 The following information was supplied regarding data availability: The raw measurements are
399 available as a Supplemental File.

400

401 **Declaration of competing interest**

402 The authors declare no conflicts of interest. The authors Jianhua Zhao and Chun Luo were employed
403 by Shanghai Majorbio Bio-Pharm Technology Co. Ltd. The author Yang Liu is employed by Gansu
404 Institute of Architectural Design and Research Company. The remaining authors confirm the research was
405 conducted in the absence of any commercial or financial relationships that could be construed as a
406 potential conflict of interest. The funders had no role in the design of the study; nor, in the collection,
407 analyses, or interpretation of data, in the writing of the manuscript, or in the decision to publish the results.

408

Figure legends

Fig. 1. Soil microbial biomass indices (MBC, MBN, MBC/MBN ratio, and MBP) (A) and soil enzyme activities (BG, CBH, LAP, NAG, and ACP) (B) in the topsoil (to a depth of 20 cm) at the two sampling sites. *BSL*: bare sandy land; *CE*: sandy land where *Cyperus esculentus* was cultivated for 1 year. Bars are the mean \pm SD, $n = 5$. *MBC*: microbial biomass carbon; *MBN*: microbial biomass nitrogen; *MBP*: microbial biomass phosphorus; *BG*: 1,4-glucosidase; *CBH*: cellobiohydrolase; *LAP*: leucine aminopeptidase; *NAG*: β -1, 4-N-acetaminoglycosidase; *ACP*: acid phosphatase. Significant levels (ANOVA followed by LSD test): * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Fig. 2. Comparison of the 16S rRNA gene (for bacteria) and 18 rRNA gene (for eukaryota) copy numbers from the soils at the two sampling sites as determined by qPCR. *BSL*: bare sandy land; *CE*: sandy land where *Cyperus esculentus* was cultivated for 1 year. Bars are the mean \pm SD, $n = 5$. Significant levels (ANOVA followed by LSD test): * $P < 0.05$.

Fig. 3. The α -diversity (Chao and Shannon indices) of the microbial community between the two sites. *BSL*: bare sandy land; *CE*: sandy land where *Cyperus esculentus* was cultivated for 1 year. Bars are the mean \pm SD, $n = 5$. Significant levels (ANOVA followed by LSD test): * $P < 0.05$.

Fig. 4. Relative abundances of the bacterial and eukaryota community members at the phylum level across all 10 soil samples. *BSL*: bare sandy land; *CE*: sandy land in which *Cyperus esculentus* was cultivated for 1 year.

Fig. 5. Results of the principal co-ordinates analysis (PC1 and PC2) for the soil bacterial and eukaryotic communities at the two sampling sites. *BSL*: bare sandy land; *CE*: sandy land in which *Cyperus esculentus* was cultivated for 1 year.

Fig. 6. Proportions at the phylum level for the two groups and tests for their significant differences. *BSL*: bare sandy land; *CE*: sandy land in which *C. esculentus* was cultivated for 1 year.

Fig. 7. Pairwise comparisons of environmental factors with soil bacterial or eukaryota community composition, for which a color gradient denotes Pearson's correlation coefficient.

SOC: soil organic carbon; *TN*: total nitrogen; *TP*: total phosphorus; *EC*: electric conductivity; *MBC*: microbial biomass carbon; *MBN*: microbial biomass nitrogen; *MBP*: microbial biomass phosphorus. Pearson's $r > 0$

indicates a positive correlation and $r < 0$ indicates a negative correlation. The boxed-edge width corresponds to the distance dependence of Mantel's R statistic, whose statistical significance was based on $n = 9999$ permutations. An inner square's size is proportional to the strength of Mantel's R correlation. The color of the curved lines indicates the correlation strength between bacterial or eukaryota community composition and the environmental factors.

References

- Allison SD, Jastrow JD. 2006. Activities of extracellular enzymes in physically isolated fractions of restored grassland soils. *Soil Biology & Biochemistry* **38**: 3245-3256. DOI 10.1016/j.soilbio.2006.04.011.
- Asare PA, Kpankpari R, Adu MO, Afutu E, Adewumi AS. 2020. Phenotypic characterization of tiger nuts (*Cyperus esculentus*) from major growing areas in Ghana. *The Scientific World Journal* 7232591.
- Baubin C, Farrell AM, Oviék A, Ghazaryan L, Gillor O. 2019. Seasonal and spatial variability in total and active bacterial communities from desert soil. *Pedobiologia*. **74**: 7-14. DOI 10.1016/j.pedobi.2019.02.001.
- Brunner K, Peterbauer CK, Mach RL, Lorito M, Zeilinger S, Kubicek CP. 2003. The NagI N-acetylglucosaminidase of *Trichoderma atroviride* is essential for chitinase induction by chitin and of major relevance to biocontrol. *Current Genetics*. **43**: 289-295. DOI10.1007/s00294-003-0399-y.
- Bünemann E, Bongiorno G, Bai Z, Creamer RE, Brussaard L. 2018. Soil quality - A critical review. *Soil Biology & Biochemistry* **120**: 105-125. DOI 10.1016/j.soilbio.2018.01.030.
- Caldwell B. 2005. Enzyme activities as a component of soil biodiversity: A review. *Pedobiologia*. **49**: 637-644. DOI 10.1016/j.pedobi.2005.06.003.
- Chen DM, Mi J, Chu PF, Cheng JH, Zhang LX, Pan QM, Xie YC, Bai YF. 2015. Patterns and drivers of soil microbial communities along a precipitation gradient on the Mongolian Plateau. *Landscape Ecology* **30**: 1669-1682. DOI 10.1007/s10980-014-9996-z.
- Chen SF, Zhou YQ, Chen YR, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**: 884-890. DOI 10.1093/bioinformatics/bty560.
- Crits-Christoph A, Robinson CK, Barnum T, Fricke WF, Davila AF, Jedynak B, McKay CP Di Ruggiero J. 2013. Colonization patterns of soil microbial communities in the Atacama Desert. *Microbiome* **1**: 28. DOI 10.1186/2049-2618-1-28.
- Dennis KL, Wang Y, Blatner NR, Wang S, Saadalla A, Trudeau E. 2013. Adenomatous polyps are driven by microbe-instigated focal inflammation and are controlled by IL-10-producing T Cells. *Cancer Research* **73**: 5905-5913. DOI 10.1158/0008-5472.CAN-13-1511.
- Dick WA, Cheng L, Wang P. 2000. Soil acid and alkaline phosphatase activity as pH adjustment indicators. *Soil Biology & Biochemistry* **32**: 1915-1919. DOI 10.1016/S0038-0717(00)00166-8.

- 469 **Dirks RC, Singh M, Potter GS, Sobotka LG, Schaefer J. 2012.** Carbon partitioning in soybean (*Glycine max*)
470 leaves by combined ¹¹C and ¹³C labeling. *New Phytologist* **196**: 1109-1121. DOI 10.1111/j.1469-
471 8137.2012.04333. x.
- 472 **Edgar RC. 2013.** UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nature Methods* **10**:
473 996-998. DOI 10.1038/nmeth.2604.
- 474 **Fierer N, Bradford MA, Jackson RB. 2007.** Toward an ecological classification of soil bacteria. *Ecology* **88**:
475 1354-1364. DOI 10.1890/05-1839.
- 476 **Fu QL, Liu C, Ding NF, Lin YC, Guo B, Luo JF, Wang HH. 2012.** Soil microbial communities and enzyme
477 activities in a reclaimed coastal soil chronosequence under rice–barley cropping. *Journal of Soils and*
478 *Sediments* **12**: 1134-1144. DOI 10.1007/s11368-012-0544-7.
- 479 **Haan RD, Kroukamp H, Zyl JHDV, Zyl WHV. 2013.** Cellobiohydrolase secretion by yeast: Current state and
480 prospects for improvement. *Process Biochemistry* **48**: 1-12. DOI 10.1016/j.procbio.2012.11.015.
- 481 **Hu S, Chapin FS, Firestone MK, Field CB, Chiariello NR. 2001.** Nitrogen limitation of microbial decomposition
482 in a grassland under elevated CO₂. *Nature* **409**: 188-191. DOI 10.1038/35051576.
- 483 **Hu YJ, Dan X, Veresoglou SD, Chen FL, Chen YL, Hao ZP, Zhang X. 2014.** Soil organic carbon and soil
484 structure are driving microbial abundance and community composition across the arid and semi-arid grasslands
485 in northern China. *Soil Biology & Biochemistry* **77**: 51-57. DOI 10.1016/j.soilbio.2014.06.014.
- 486 **Hu YG, Zhang ZS, Huang L, Qi Q, Liu LC, Zhao Y, Wang ZR, Zhou HK, Lv XY, Mao ZC, Yang YF, Zhou**
487 **JZ, Kardol P. 2019.** Shifts in soil microbial community functional gene structure across a 61-year desert
488 revegetation chronosequence. *Geoderma* **347**: DOI 126-134. 10.1016/j.geoderma.2019.03.046.
- 489 **Jenkinson DS, Ladd JN. 1981.** Microbial biomass in soil: Measurement and turnover. *Soil biochemistry* **5**.
- 490 **Jiao S, Lu YH. 2020.** Soil pH and temperature regulate assembly processes of abundant and rare bacterial
491 communities in agricultural ecosystems. *Environment Microbiology* **22**: 1052-1065. DOI 10.1111/1462-
492 2920.14815.
- 493 **Joergensen RG. 1996.** The fumigation-extraction method to estimate soil microbial biomass: Calibration of the
494 kEC value. *Soil Biology & Biochemistry* **28**: 25-31. DOI 10.1016/0038-0717(95)00102-6.
- 495 **Lange M, Eisenhauer N, Sierra CA, Bessler H, Engels C, Griffiths RI, Mellado-Vázquez PG, Malik AA, Roy**
496 **J, Scheu S, Steinbeiss S, Thomson BC, Trumbore SE, Gleixner G. 2015.** Plant diversity increases soil
497 microbial activity and soil carbon storage. *Nature Communications* **6**: 6707. DOI 10.1038/ncomms7707.
- 498 **Ma J, Rabin SS, Anthoni P, Bayer AD, Nyawira SS, Olin S. 2022.** Assessing the impacts of agricultural
499 managements on soil carbon stocks, nitrogen loss, and crop production – a modelling study in eastern Africa.
500 *Biogeosciences* **19**: 2145-2169. DOI 10.5194/bg-2021-352.
- 501 **Margalef O, Sardans J, Maspons J, Molowny-Horas R, Fernández-Martínez M, Janssens IA, Richter A, Ciais**
502 **P, Obersteiner M, Peñuelas J. 2021.** The effect of global change on soil phosphatase activity. *Global Change*
503 *Biology* **27**: 5989-6003. DOI 10.1111/gcb.15832.
- 504 **Muhammad I, Wang J, Sainju UM, Zhang SH, Zhao FZ, Khan A. 2021.** Cover cropping enhances soil

- microbial biomass and affects microbial community structure: A meta-analysis. *Geoderma* 381: 114696. DOI 10.1016/j.geoderma.2020.114696.
- Neilson JW, Quade J, Ortiz M, Nelson WM, Legatzki A, Tian F, LaComb F, Betancourt JL, Wing RA, Soderlund CA, Maier RM. 2012.** Life at the hyperarid margin: novel bacterial diversity in arid soils of the Atacama Desert, Chile. *Extremophiles* 16: 553-566. DOI 10.1007/s10482-013-9911-7.
- Niu YY, Duan YL, Li YQ, Wang XY, Chen Y, Wang LL. 2021.** Soil microbial community responses to short-term nitrogen addition in China's Horqin Sandy Land. *PLoS ONE* 16: e0242643. DOI 10.1371/journal.pone.0242643.
- Oderinde RA, Tairu OA. 1988.** Evaluation of the properties of yellow nutsedge (*Cyperus esculentus*) tuber oil. *Food Chemistry* 28(3): 233-237. DOI 10.1016/0308-8146(88)90055-6.
- Pan FJ, Zhang W, Liang YM, Liu SJ, Wang KL. 2018.** Increased associated effects of topography and litter and soil nutrients on soil enzyme activities and microbial biomass along vegetation successions in karst ecosystem, southwestern China. *Environmental Science and Pollution Research* 25: 16979-16990. DOI 10.1007/s11356-018-1673-3.
- Powlson DS, Hirsch PR, Brookes PC. 2001.** The role of soil microorganisms in soil organic matter conservation in the tropics. *Nutrient Cycling in Agroecosystems* 61: 41-51. DOI 10.1023/A:1013338028454.
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner FO. 2013.** The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research* 41: 590-596. DOI 10.1093/nar/gks1219.
- Ralte V, Pandey HN, Barik SK, Tripathi RS, Prabhu SD. 2005.** Changes in microbial biomass and activity in relation to shifting cultivation and horticultural practices in subtropical evergreen forest ecosystem of north-east India. *Acta Oecologica* 28: 163-172. DOI 10.1016/j.actao.2005.04.001.
- Rousk J, Bååth E, Brookes PC, Lauber CL, Lozupone C, Caporaso JG, Knight R, Fierer N. 2010.** Soil bacterial and fungal communities across a pH gradient in an arable soil. *The ISME Journal* 4: 1340-1351. DOI 10.1038/ismej.2010.58.
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF. 2009.** Introducing mothur: opensource, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology* 75: 7537-7541. DOI 10.1128/AEM.01541-09.
- Sha RN, Yu MH, Ding GD, Zhao YY, Sai K, Zhang GX. 2022.** Shelter efficiency of conservation tillage model in *Cyperus esculentus* in sandy farmland. *Journal of Arid Land Resources and Environment* 36: 87-94. DOI 10.13448/j.cnki.jalre.2022.096 (In Chinese).
- Shao TY, Zhao JJ, Liu AH, Long XH, Rengel Z. 2020.** Effects of soil physicochemical properties on microbial communities in different ecological niches in coastal area. *Applied Soil Ecology* 150: 103486. DOI 10.1016/j.apsoil.2019.103486.
- Shen CC, Xiong JB, Zhang HY, Feng YZ, Lin XG, Li XY, Liang WJ, Chu HY. 2013.** Soil pH drives the spatial

- 541 distribution of bacterial communities along elevation on Changbai Mountain. *Soil Biology & Biochemistry* **57**:
542 204-211. DOI 10.1016/j.soilbio.2012.07.013.
- 543 **Sparling GP, Pankhurst C, Doube B, Gupta V. 1997.** Soil microbial biomass, activity and nutrient cycling as
544 indicators of soil health. Biological indicators of soil health. Wallingford; New York (USA): CAB International
545 97-119.
- 546 **Sun Y, Shi YL, Wang H, Zhang T, Yu LY, Sun H, Zhang YQ. 2018.** Diversity of bacteria and the characteristics
547 of actinobacteria community structure in Badain Jaran Desert and Tengger Desert of China. *Frontiers in*
548 *Microbiology* **9**: 1068. DOI 10.3389/fmicb.2018.01068.
- 549 **Tian J, He NP, Hale L, Niu SL, Yu GR, Liu Y, Blagodatskaya E, Kuzyakov Y, Gao Q, Zhou JZ. 2018.** Soil
550 organic matter availability and climate drive latitudinal patterns in bacterial diversity from tropical to cold
551 temperate forests. *Functional Ecology* **32**: 61-70. DOI 10.1111/1365-2435.12952.
- 552 **Wang JC, Song Y, Ma TF, Raza W, Li J, Howland JG, Huang QW, Sheng QR. 2017.** Impacts of inorganic and
553 organic fertilization treatments on bacterial and fungal communities in a paddy soil. *Applied Soil Ecology* **112**:
554 42-50. DOI 10.1016/j.apsoil.2017.01.005.
- 555 **Wang XB, Van Nostrand, JD, Deng Y, Lv XT, Wang C, Zhou JZ, Han XG. 2015.** Scale-dependent effects of
556 climate and geographic distance on bacterial diversity patterns across northern China's grasslands. *FEMS*
557 *Microbiology Ecology* **91**: 12. DOI 10.1093/femsec/fiv133.
- 558 **Yu J, Yin Q, Niu JM, Yan ZJ, Wang H, Wang YQ, Chen DM. 2022.** Consistent effects of vegetation patch type
559 on soil microbial communities across three successional stages in a desert ecosystem. *Land Degradation &*
560 *Development* **33**: 1552-1563. DOI 10.1002/ldr.4194.
- 561 **Yu WJ, Lawrence NC, Sooksa-nguan T, Smith SD, Tenesaca C, Howe AC, Hall SJ. 2021.** Microbial linkages to
562 soil biogeochemical processes in a poorly drained agricultural ecosystem. *Soil Biology and Biochemistry*
563 **156**: 108228. DOI 10.1016/j.soilbio.2021.108228.
- 564 **Zeng QC, An SS, Liu Y, Wang HL, Wang Y. 2019.** Biogeography and the driving factors affecting forest soil
565 bacteria in an arid area. *Science of The Total Environment* **680**: 124-131. DOI 10.1016/j.scitotenv.2019.04.184.
- 566 **Zhang KP, Shi Y, Cui XP, Yue P, Li KH, Liu XJ, Tripathi BM, Chu HY, Lozupone C. 2019.** Salinity is a key
567 determinant for soil microbial communities in a desert ecosystem. *mSystems* **4**: e00225-18. DOI
568 10.1128/mSystems.00225-18.
- 569 **Zhang XT, Wu XQ. 2022.** Cultivation and production potential evaluation of *Cyperus esculentus* L. in marginal
570 land of Inner Mongolia. *Transactions of the Chinese Society of Agricultural Engineering* **38**: 289-295. DOI
571 10.11975/j.issn.1002-6819.2022.02.032. (In Chinese).
- 572 **Zhao CC, Miao Y, Yu CD, Zhu LL, Wang F, Jiang L, Hui DF, Wan SQ. 2016.** Soil microbial community
573 composition and respiration along an experimental precipitation gradient in a semiarid steppe. *Scientific*
574 *Reports* **6**: 24317. DOI 10.1038/srep24317.

Figure 1

Soil microbial biomass indices (MBC, MBN, MBC/MBN ratio, and MBP) (A) and soil enzyme activities (BG, CBH, LAP, NAG, and ACP) (B) in the topsoil (to a depth of 20 cm) at the two sampling sites

BSL : bare sandy land; *CE*: sandy land where *Cyperus esculentus* was cultivated for 1 year.

Bars are the mean \pm SD, $n = 5$. *MBC* : microbial biomass carbon; *MBN*: microbial biomass nitrogen; *MBP*: microbial biomass phosphorus; *BG*: 1,4-glucosidase; *CBH*: cellobiohydrolase; *LAP*: leucine aminopeptidase; *NAG*: β -1, 4-N-acetaminoglycosidase; *ACP*: acid phosphatase.

Significant levels (ANOVA followed by LSD test): * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.





Figure 2

Fig. 2. Comparison of the 16S rRNA gene (for bacteria) and 18 rRNA gene (for eukaryota) copy numbers from the soils at the two sampling sites as determined by qPCR.

BSL: bare sandy land; *CE*: sandy land where *Cyperus esculentus* was cultivated for 1 year.

Bars are the mean \pm SD, $n = 5$. Significant levels (ANOVA followed by LSD test): * $P < 0.05$.




Figure 3

The α -diversity (Chao and Shannon indices) of the microbial community between the two sites.

BSL: bare sandy land; CE: sandy land where *Cyperus esculentus* was cultivated for 1 year.

Bars are the mean \pm SD, $n = 5$. Significant levels (ANOVA followed by LSD test): * $P < 0.05$.





Figure 4

Relative abundances of the bacterial and eukaryota community members at the phylum level across all 10 soil samples.

BSL: bare sandy land; *CE*: sandy land in which *Cyperus esculentus* was cultivated for 1 year.

Bacteria



Eukaryota




Figure 5

Results of the principal co-ordinates analysis (PC1 and PC2) for the soil bacterial and eukaryotic communities at the two sampling sites.

BSL: bare sandy land; *CE*: sandy land in which *Cyperus esculentus* was cultivated for 1 year.





Figure 6

Proportions at the phylum level for the two groups and tests for their significant differences.

BSL: bare sandy land ; CE: sandy land in which *C. esculentus* was cultivated for 1 year.




Figure 7

Pairwise comparisons of environmental factors with soil bacterial or eukaryota community composition, for which a color gradient denotes Pearson's correlation coefficient.

SOC: soil organic carbon; *TN*: total nitrogen; *TP*: total phosphorus; *EC*: electric conductivity; *MBC*: microbial biomass carbon; *MBN*: microbial biomass nitrogen; *MBP*: microbial biomass phosphorus. Pearson's $r > 0$ indicates a positive correlation and $r < 0$ indicates a negative correlation. The boxed-edge width corresponds to the distance dependence of Mantel's R statistic, whose statistical significance was based on $n = 9999$ permutations. An inner square's size is proportional to the strength of Mantel's R correlation. The color of the curved lines indicates the correlation strength between bacterial or eukaryota community composition and the environmental factors.

