

Convex-hull mass estimates of the dodo (*Raphus cucullatus*): application of a CT-based mass estimation technique

Charlotte A Brassey, Thomas O'Mahoney, Andrew C Kitchener, Phillip L Manning, William I Sellers

The external appearance of the dodo (*Raphus cucullatus*) has been a source of considerable intrigue as contemporaneous accounts or depictions are rare. The body mass of the dodo has been particularly contentious, with the flightless pigeon alternatively reconstructed as slim or overweight depending upon the skeletal metric used as the basis for mass prediction. Resolving this dichotomy and obtaining a reliable estimate for mass is essential before future analyses regarding dodo life history, physiology or biomechanics can be conducted. Previous mass estimates of the dodo have relied upon predictive equations based upon hind limb dimensions of extant pigeons. Yet the hind limb proportions of dodo have been found to differ considerably from those of their modern relatives, particularly with regards to midshaft diameter. Therefore, application of predictive equations to unusually robust fossil skeletal elements may bias mass estimates. We present a whole-body computed tomography (CT) -based mass estimation technique for application to the dodo. We generate 3D volumetric renders of the articulated skeletons of 20 species of extant pigeons, and wrap minimum-fit 'convex hulls' around their bony extremities. Convex hull volume is subsequently regressed against mass to generate predictive models based upon whole skeletons. Our best-performing predictive model is characterized by high correlation coefficients and low mean squared error ($a=-2.31$, $b=0.90$, $r^2=0.97$, $MSE=0.0046$). When applied to articulated composite skeletons of the dodo (National Museums Scotland, NMS.Z.1993.13; Natural History Museum, NHM A.9040 and S/1988.50.1), we estimate eviscerated body masses of 8-10.8 kg. When accounting for missing soft tissues, this may equate to live masses of 10.6-14.3 kg. Mass predictions presented here fall towards the lower end of those previously published, and support recent suggestions of a relatively slim dodo. CT-based reconstructions provide a means of objectively estimating mass and body segment properties of extinct species using whole articulated skeletons.

Convex-hull mass estimates of the dodo (*Raphus cucullatus*): application of a CT-based mass estimation technique

Brassey, C.A.^{1*}, O'Mahoney, T.¹, Kitchener, A.C.², Manning, P. L.³, Sellers, W.I.¹

¹Faculty of Life Sciences, University of Manchester, Manchester M13 9PL

²Department of Natural Sciences, National Museums Scotland, Edinburgh EH1 1JF

³Interdisciplinary Centre for Ancient Life, School of Earth, Atmospheric and Environmental Sciences, University of Manchester, Manchester M13 9PL

*corresponding author (charlotte.brassey@manchester.ac.uk)

Abstract

The external appearance of the dodo (*Raphus cucullatus*) has been a source of considerable intrigue as contemporaneous accounts or depictions are rare. The body mass of the dodo has been particularly contentious, with the flightless pigeon alternatively reconstructed as slim or overweight depending upon the skeletal metric used as the basis for mass prediction. Resolving this dichotomy and obtaining a reliable estimate for mass is essential before future analyses regarding dodo life history, physiology or biomechanics can be conducted.

Previous mass estimates of the dodo have relied upon predictive equations based upon hind limb dimensions of extant pigeons. Yet the hind limb proportions of dodo have been found to differ considerably from those of their modern relatives, particularly with regards to midshaft diameter. Therefore, application of predictive equations to unusually robust fossil skeletal elements may bias mass estimates. We present a whole-body computed tomography (CT) -based mass estimation technique for application to the dodo. We generate 3D volumetric renders of the articulated skeletons of 20 species of extant pigeons, and wrap minimum-fit 'convex hulls' around their bony extremities. Convex hull volume is subsequently regressed against mass to generate predictive models based upon whole skeletons.

Our best-performing predictive model is characterized by high correlation coefficients and low mean squared error ($a=-2.31$, $b=0.90$, $r^2=0.97$, $MSE=0.0046$). When applied to articulated composite skeletons of the dodo (National Museums Scotland,

NMS.Z.1993.13; Natural History Museum, NHM A.9040 and S/1988.50.1), we estimate eviscerated body masses of 8-10.8 kg. When accounting for missing soft tissues, this may equate to live masses of 10.6-14.3 kg. Mass predictions presented here fall towards the lower end of those previously published, and support recent suggestions of a relatively slim dodo. CT-based reconstructions provide a means of objectively estimating mass and body segment properties of extinct species using whole articulated skeletons.

Key words: dodo, *Raphus cucullatus*, body mass, computed tomography, convex hulls

Introduction

Body mass (M_b) is a fundamental descriptor of an organism and co-varies with important ecological and physiological traits, such as population density, metabolism and cost-of-transport [1]. Key evolutionary scenarios, such as the origin of avian flight [2] and the extinction of island flightless avian species [3], have been diagnosed on the basis of estimated M_b . Therefore, the reconstruction of body mass in extinct bird species is a subject of considerable interest within the palaeontological and evolutionary biology literature [2–6].

An often-applied technique for estimating the body mass of an extinct vertebrate has been to measure a skeletal dimension from modern species, such as femur circumference [7] or glenoid diameter [8], and apply this as the independent variable in a regression against body mass. However, ‘overdevelopment’ of the pelvic apparatus has been found to be significantly correlated with the flightless condition in extant birds [9]. Therefore, the application of mass prediction equations, based solely on hind limb material of flightless avian taxa, has been questioned in extinct species such as the moa [6].

The dodo (*Raphus cucullatus*) is an iconic representative of island flightlessness and human-induced extinction, and its external appearance has been a source of considerable intrigue due to the scarcity of contemporaneous accounts or depictions [10]. This extinct flightless columbiform was endemic to the island of Mauritius. However, the skeletal anatomy of the dodo is comparatively well known, and its pelvic

morphology has been thoroughly investigated. Hind limb bones of *R. cucullatus* have been found to differ considerably in both their length and width relative to their volant relatives [11, although see 12]. Yet previous attempts to estimate the body mass of the dodo have predominantly relied upon predictive equations derived solely from the hind limb metrics of extant species [13–15].

An alternative approach to mass estimation involves the reconstruction of 3D volumetric models. An early volumetric reconstruction of the dodo was conducted by physically sculpting a scale model of an individual and estimating volume via fluid displacement [12]. Whilst such volumetric techniques are less liable to bias by individual robust/gracile postcranial elements than traditional linear bivariate equations, they do inevitably involve some degree of artistic licence in the sculpting of soft tissue contours and require an estimate for fossil body density to be assigned.

Following advances in 3D imaging technology, the use of digital skeletal models in mass estimation of fossil skeletons has become increasingly popular [16–19]. These studies typically involve the ‘wrapping’ of geometric shapes or lofted smooth surfaces around the skeleton in order to replicate the original soft tissue contour of the animal. Zero-density cavities such as lung and tracheal space may also be modeled [20]. However, similar to physical sculpting with clay, assumptions must still be made regarding body density and the extent of soft tissues beyond the skeleton. Therefore, it is essential that reconstructions are grounded within a quantitative understanding of extant species in order to avoid subjective modeling of soft tissues (both body and plumage).

89 Here we present new mass estimates for the dodo based on an alternative ‘convex hull’
 90 volumetric reconstruction approach [21,22]. The convex hull (*CH*) of a set of points is
 91 defined as the smallest convex polytope that contains all said points, and intuitively can
 92 be thought of as a shrink-wrap fit around an object (see Figure 1). Application of the
 93 convex hulling technique to mass estimation does not involve any subjective
 94 reconstruction of soft tissue anatomy and solely relies upon the underlying skeleton. We
 95 calculate minimum convex hull volumes for a sample of composite articulated dodo
 96 skeletons, and convert these to body mass estimates using a computed tomography
 97 (CT) calibration dataset of 20 species of extant pigeon. To our knowledge, this is the
 98 first time such an extensive CT dataset of extant animals has been used to reconstruct
 99 the body mass of a fossil of an extinct species.

Methods and Materials

The modern dataset consists of 20 columbiform individuals, spanning a wide variety of body sizes from a 70 g fruit dove (*Ptilinopus*) up to the largest extant pigeon, the 2 kg Victoria crowned pigeon (*Goura victoria*). We also cover a broad taxonomic range (including the closest extant relative of the dodo [23], the Nicobar pigeon (*Caloenas nicobarica*)). Frozen carcasses were sourced from National Museums Scotland, Edinburgh, and the University of Manchester (see Table 1). Carcasses were CT scanned at Leahurst Veterinary School, University of Liverpool, in a Toshiba Aquilion PRIME helical scanner at a slice thickness of 0.5mm and a pixel spacing of between 0.24-0.51mm, depending on the maximum size of the specimen. 3D models of the skeletons were generated in Seg3D [24], using an automatic threshold with subsequent manual masking to remove the dense rachises attached to the forelimb.

Models were exported into Geomagic Studio (www.geomagic.com), where each skeleton was divided into functional units (skull, neck, trunk, humerus, radius+ulna, carpometacarpals, femur, tibiotarsus+fibula, tarsometatarsus, feet). The cervical series was further subdivided in order to achieve a tight-fitting hull around the curving neck. Minimum convex hulls were calculated in MATLAB (www.mathworks.com), using the 'convhull' function implementing the Quickhull (qhull) algorithm [25], and total convex hull volume was calculated as the sum of individual segment volumes (see Figure 1). Body mass was measured for each carcass, and the relationship between M_b and convex hull volume (CH_{vol}) was estimated using ordinary least squares (OLS)

regression on \log_{10} transformed data. As the purpose of the regression was to derive a predictive equation, a type-I regression, such as OLS, was deemed most appropriate [26]. Additionally we accounted for the statistical non-independence of phylogenetically-related data points by carrying out phylogenetic generalized least squares (PGLS) regressions, implemented in MATLAB using 'Regression2' software [27]. A composite phylogeny was modified from Johnson et al. [28] and Heupink et al. [29], with all branch lengths equal to 1.

To reconstruct the body masses of articulated dodo skeletons, we generated 3D digital models of these specimens. The Edinburgh dodo (National Museums Scotland, NMS.Z.1993.13) was scanned using a Z+F Imager 5010 LiDAR (Light Detection And Range) scanner and reconstructed in the Z+F LaserControl software. The Natural History Museum (NHM), London specimens (Tring skeleton, S/1988.50.1; South Kensington specimen NHM A9040) were digitized using the photogrammetric technique detailed elsewhere [30,31] and reconstructed in VisualSFM [32]. Despite application of two alternative imaging techniques, previous studies have found the results obtained via photogrammetry and laser scanning to be comparable [30], and convex hull results to be insensitive to point cloud density [22,33]. 3D models of the dodo skeletons were cleaned up in Geomagic and subdivided into functional units. Our only intervention with the dodo models was to mirror the right hand side of the Edinburgh ribcage to account for missing ribs on its left side. Convex hulls were fitted according to the methodology applied to modern pigeons.

Results

Total convex hull volumes for the modern pigeons are reported in Table 1, and segment-specific CH_{vol} values can be found in Supplementary Material S1. Convex hull models are available for download from <http://www.animalsimulation.org>. We found considerable variation between frozen pigeon specimens in the posture of the digits forming the foot i.e. adduction vs. abduction of the digits. This influenced the overall shape, and hence calculated CH_{vol} , of the foot functional units (see Figure 1D). Given repositioning of the skeleton was not possible due to the frozen nature of the carcasses, here we report total CH_{vol} values with and without feet included. External inspection of the carcasses suggested evisceration had been carried out on some specimens. Using CT scans the occurrence of evisceration was confirmed across our modern dataset (see Table 1). Therefore, we report separate predictive models derived from 'eviscerated' carcasses (n=13), 'intact' carcasses (n=7), and a third 'combined' model comprising both eviscerated and intact specimens (n=20).

The results of the OLS regression analyses are presented in Table 2, and phylogenetically corrected (PGLS) regressions are given in Supplementary Material S2 alongside the composite phylogeny used in this analysis. PGLS regressions did not provide a better fit to the data than uncorrected OLS regressions (as determined by Akaike Information Criterion values, AIC) for the 'eviscerated' and 'combined' models (Table 2). However, a PGLS model was found to fit the 'intact' extant pigeon data better than an uncorrected OLS model (Table 2).

Removing CH_{vol} of the feet from the analyses had very little effect on the results of the regression, although mean squared error (MSE) decreased slightly in all models (Table 2) and therefore only regression models minus feet are discussed any further in the text. Figure 2 illustrates a strong positive correlation between M_b and CH_{vol} for the eviscerated specimens within the sample ($a=-2.31$, $b=0.90$, $r^2=0.97$, $MSE=0.0046$). In contrast, the relationship between M_b and CH_{vol} in intact specimens illustrates a weak positive correlation characterized by low correlation coefficients and high mean square error ($a=-1.06$, $b=0.66$, $r^2=0.70$, $MSE=0.0161$). Intact specimens do not plot consistently above the eviscerated pigeon slope (Figure 2) and are instead characterized by a high degree of scatter. When combining the eviscerated and intact specimens into one dataset, M_b and CH_{vol} remain tightly correlated ($a=-2.08$, $b=0.85$, $r^2=0.92$, $MSE=0.0084$).

Total CH_{vol} calculated for the mounted dodo skeletons are reported in Table 3 (see Supplementary Material S3 for segment-specific values) and an example of a photogrammetric model is illustrated in Figure 3. Using the ‘eviscerated’ predictive model, dressed M_b is estimated as 8.0 kg (95% prediction interval (PI) 4.6-13.9 kg), 8.7 kg (95%PI 5.0-15.0 kg) and 10.8 kg (95%PI 6.1-19.0 kg) respectively for the NHM Tring, NHM South Kensington and Edinburgh dodos. Applying the ‘combined’ predictive equation results in wider and therefore more conservative prediction intervals (NHM Tring, 6.7 kg 95%PI 3.5-13.1 kg; NHM South Kensington, 7.3 kg 95%PI 3.7-14.3 kg; Edinburgh, 9.0 kg 95%PI 4.5-17.9 kg).

189 Figure 4 illustrates the distribution of segment-specific convex hull volumes as a
 190 proportion of total CH_{vol} within the models. In extant pigeons trunk CH_{vol} represents on
 191 average 69% of total CH_{vol} . The NHM Tring dodo skeleton has a percentage trunk
 192 volume significantly lower than that of extant pigeons (67%, 1-tailed t -test, $t=3.23$,
 193 $p<0.01$), whilst percentage trunk volume in the NHM South Kensington and Edinburgh
 194 skeletons is significantly higher than extant pigeons (71% and 80%, $t=-2.23$ and -13.0
 195 respectively, $p<0.05$). With the exception of the tarsometatarsii of the NHM South
 196 Kensington skeleton, pelvic convex hull segments of the dodos comprise a significantly
 197 greater proportion of total CH_{vol} than in extant pigeons ($p<0.05$). In contrast, dodo
 198 pectoral convex hull segments contribute proportionally less to total CH_{vol} than in extant
 199 pigeons ($p<0.0001$) (Figure 4).

Discussion

Predictive equation derived from modern CT dataset

To our knowledge the present study represents the first application of a predictive equation derived solely from whole-body CT to the problem of body mass estimation for extinct animals. Previous volumetric mass estimate studies have relied upon articulated skeletons of extant species to derive a calibration equation [6,21]. Yet articulated skeletons are often missing crucial specimen information, such as a recorded body mass. By working with frozen carcasses, body mass is directly measurable and uncertainties associated with mounting and posing of the skeletons can be avoided [22].

Our dataset consists of both 'intact' and 'eviscerated' pigeons as determined by examination of CT scans. Previous analyses of carcass composition have found eviscerated mass to represent 62-66% of live body mass in rock doves [34,35], yet no data exist regarding the possible scaling of internal organ mass across a range of body sizes in the Columbiformes. As can be seen in Figure 2, there is no consistent disparity between intact and eviscerated specimens, and the relationship between M_b and CH_{vol} in intact pigeons is relatively weak ($r^2=70$, $p=0.019$). This correlation improves considerably when accounting for phylogeny (Supplementary Material S2), but remains weaker than the relationship between M_b and CH_{vol} characterizing eviscerated specimens. Live body mass has been shown to vary considerably in wild animals due to hydration, nutrition and gut content [36] and therefore some degree of scatter is to be expected in intact carcasses. Particularly striking is the variability in gizzard contents between similar-sized specimens visible in CT scans (see Figure 5).

This suggests intact pigeon M_b cannot be corrected for the presence of internal organs using a single factor representing average percentage eviscerated mass as a function of live mass (i.e. multiplying by values of 0.62 or 0.66 previously found in the literature). Additionally, attempting to correct intact M_b by substituting intact CH_{vol} into the eviscerated regression model would be highly circular and result in artificially inflated correlation coefficients, if the equation were used in a predictive capacity. Therefore, we apply both the uncorrected OLS eviscerated model and combined (eviscerated and intact) model to bracket the range of likely dodo body masses. Interestingly, the very high correlation coefficient and low mean squared error of the eviscerated equation ($r^2=0.97$, $MSE=0.0046$) suggest that once the variability associated with fluid and gut content is removed, the relationship between the mass of the remaining musculoskeletal system and CH_{vol} is more tightly constrained.

Volumetric body mass estimation applied to the dodo

No reliable records of the body mass of dodo exist prior to its extinction in the 17th Century and subsequent mass estimates have varied considerably. Early accounts of the flightless bird suggested an average mass of 50 lb (22 kg) [37], although such accounts “have a tendency towards exaggeration” [10]. More recently a ‘slim’ dodo (mean 10.2 kg) was proposed on the basis of femoral, tibiotarsal and tarsometatarsal length scaling in modern birds [13]. However hind limb bone length has been shown to correlate poorly with body mass relative to other cross-sectional geometric properties and frequently contains a strong functional signal [38]. Alternatively, a predictive

equation based on femoral and tibiotarsal least circumference in ground-dwelling birds has suggested mass estimates between 9.5-12.3 kg [14].

The application of volumetric mass estimation techniques to the dodo has been rare. A sculpted scale model of a 'slim' dodo based upon mean skeletal measures was created to replicate sketches dating contemporaneously to its survival on Mauritius and resulted in mass estimated of 12.5-16.1 kg [12]. In the same study a 'fat' dodo model based on later 'exaggerated' artworks was predicted to weigh between 21.7-27.8 kg.

Here we estimate mean eviscerated body masses for articulated composite dodo skeletons of between 8.0-10.8 kg. Without further information regarding the effect of within-subject variability in gizzard, crop or gut content or interspecific scaling of viscera mass, any extrapolation to a live mass should be treated with caution. However, with this caveat in mind, a 33% increase in mass to account for missing organs (as quantified in extant *C. livia*) would take our results to 10.6-14.3 kg. This overlaps with the slim sculpted model based on contemporaneous accounts [12]. Including our 95% confidence intervals takes both the NHM Tring and South Kensington skeletons to a maximum of 18.2 kg and 19.9 kg whole body masses, still considerably below the 22kg suggested historically [37]. In contrast, the 95% confidence intervals of the Edinburgh dodo include 22kg once multiplied by 1.33.

Unlike all previous volumetric studies, our convex hulling technique does not require a value for body density to be assigned from the literature. Instead we directly derive the

relationship between M_b and CH_{vol} in order to avoid uncertainty regarding assigning literature values, which have been shown to differ considerably across avian groups and with various methodologies for estimating body density [17]. However, this does implicitly rely upon the predictive equation being applied to a fossil of an extinct species that is closely related to (and can therefore be assumed to share a similar body density to) the modern dataset from which the predictive equation was derived. In this case of estimating dodo mass based on extant pigeons, we believe this assumption can be upheld. Alternatively, CH_{vol} may be multiplied by a given value of carcass density to give a hard lower limit to body mass (as carcass volume cannot be less than convex hull volume). The sole literature value for intact feathered pigeon density is 648kg/m^3 from Hamershock et al. [39], producing hard lower bounds to estimated body mass 5.8kg, 6.3kg, and 7.9kg for the Tring, Kensington and Edinburgh composite skeletons respectively.

We consider the convex hulling technique to be superior to other sculpting-based volumetric methods (such as manual sculpting with clay [12] or digital sculpting with non-uniform rational B-spline (NURBs) curves [19]) for the purpose of mass estimation as soft tissues and hypothesized respiratory systems need not be reconstructed for fossils of extinct species. When values for centre of mass (COM) and segment inertial properties are required for further biomechanical analyses, NURBs may be required in order to achieve a representative mass distribution across the skeleton. In such situations it is essential that soft tissue reconstructions are based on quantitative comparative dissection data from relevant modern species in order to minimize

subjectivity in model creation. However, for the sole purpose of mass estimation, convex hulling should be the preferred technique.

Composite and articulated skeletons

The dodo specimens included in this study are composite skeletons, comprising skeletal material from more than one individual and including sculpted or cast elements. Therefore, our study is limited to estimating the body mass of the hypothetical animal represented by each articulated skeleton, rather than a known individual. Currently there exists only one near-complete dodo skeleton comprising a single individual (the Thirioux dodo), upon which research is currently still continuing [40].

Whilst the use of composite skeletons should clearly be treated with caution when used in biomechanical analyses, their composite nature does not entirely rule out their use, particularly in the case of mass estimation. A recent large-scale macroevolutionary [41] study of body size in a fossil lineage relied upon mass data derived solely from humeral and femoral circumferences of one individual specimen per species. While this approach is often the only one feasible, given the highly fragmentary nature of the fossil record, taking one individual as being representative of an entire species leaves us vulnerable to the possibility of high levels of intraspecific variation.

In contrast, a volumetric reconstruction based on a composite skeleton is likely to reflect a species average by virtue of being a combination of several individuals and will be less skewed by isolated robust or gracile elements. If subsequent biomechanical

analyses are to be carried out (such as FEA on a particular musculoskeletal unit), then it is important that the body mass entered into the analyses is representative of that specific individual. However, for the case of volumetric body mass estimation alone, a representative species mean can be derived from a composite skeleton.

Of more concern is the frequency of missing, deformed or reconstructed material within a fossil mount. Known issues with the dodo mounts included in this study include missing ribs (Edinburgh skeleton), missing carpals (NHM South Kensington skeleton), deformation of the fragile pubis (NHM South Kensington skeleton) or the loss of the most of the ischium, pubis and caudal vertebrae (NHM Tring skeleton). For a given object, the extent of the convex hull fitted to that object is dictated solely by its geometric extremes. In many ways this is advantageous for volumetric fossil reconstructions as damage occurring within the bounds of the convex hull does not affect our volume estimate. However when extremities are missing (such as the caudal tip of the pubis), the shape and volume of the convex hull are strongly affected. This is evident in the low percentage trunk volume of the NHM Tring skeleton (Figure 5) compared to those of extant pigeons and other dodos.

Whilst some evidence of paedomorphosis (retention of juvenile characters as an adult) of pectoral elements and peramorphosis (characters that mature past adulthood) of pelvic elements in the dodo is discernable relative to extant volant pigeons, Figure 5 predominantly illustrates the important contribution of trunk volume to total mass estimates. The Edinburgh skeleton has a proportionally more voluminous trunk than that

338 of extant pigeons and other dodo skeletons, and therefore all other skeletal elements
 339 contribute proportionally less to total CH_{vol} . The more voluminous trunk relative to other
 340 specimens may be attributed to the anterior positioning of the sternum due to
 341 constraints associated with the armature supporting the mount. The opposite is true of
 342 the NHM Tring skeleton, in which damage to the extremities of the pelvic girdle result in
 343 a reduced trunk volume. This highlights the sensitivity of volumetric reconstructions of
 344 fossils of extinct species to trunk morphology, and should be a concern when working
 345 with both composite and complete fossil specimens.

Summary

Here we present the first volumetric reconstruction of fossil body mass based entirely on modern whole-animal CT data. The eviscerated body mass of three articulated composite dodo skeletons is estimated to fall between 8.0-10.8 kg. When accounting for missing organ mass, our mean values still fall towards the lower range of previously published mass estimates. As the availability and cost of CT improves, we believe this non-subjective convex hull approach will become increasingly commonplace. Mass estimation of extinct species from fossils relies upon two key components; a reliable calibration equation derived from extant species, and an accurate reconstruction of the extinct individual from its fossil. We discuss the issues surrounding the use of articulated composite skeletons, and highlight the particular importance of trunk morphology to volume reconstructions. We suggest future efforts should focus on quantifying ribcage and sternal geometry in extant groups in order to bracket the possible trunk shape in fossils of extinct species.

360 **Acknowledgements**

361 We thank Zoller+Fröhlich (Z+F) Limited for their continued support, Jennifer Anné and
 362 Dr Victoria Egerton (University of Manchester), Martin Baker (University of Liverpool),
 363 Malgosia Nowak-Kemp (Oxford Museum of Natural History) and Julian Hume, Judith
 364 White and Sandra Chapman (Natural History Museum, London).

References

1. Schmidt-Nielsen K. *Scaling: Why is animal size so important?* Cambridge: Cambridge University Press; 1984.
2. Turner AH, Pol D, Clarke JA, Erickson GM, Norell MA. A basal dromaeosaurid and size evolution preceding avian flight. *Science* (80-). 2007;317: 1378–1381. doi:10.1126/science.1144066
3. Boyer AG. Extinction patterns in the avifauna of the Hawaiian islands. *Divers Distrib.* 2008;14: 509–517. doi:10.1111/j.1472-4642.2007.00459.x
4. Hone DWE, Dyke GJ, Haden M, Benton MJ. Body size evolution in Mesozoic birds. *J Evol Biol.* 2008;21: 618–624. doi:10.1111/j.1420-9101.2007.01483.x
5. Butler RJ, Goswami A. Body size evolution in Mesozoic birds: Little evidence for Cope's rule. *J Evol Biol.* 2008;21: 1673–1682. doi:10.1111/j.1420-9101.2008.01594.x
6. Brassey CA, Holdaway RN, Packham AG, Anne J, Manning PL, Sellers WI. More Than One Way of Being a Moa: Differences in Leg Bone Robustness Map Divergent Evolutionary Trajectories in Dinornithidae and Emeidae (Dinornithiformes). *PLoS One.* 2013;8: e82668. doi:10.1371/journal.pone.0082668
7. Campione NE, Evans DC. A universal scaling relationship between body mass and proximal limb bone dimensions in quadrupedal terrestrial tetrapods. *BMC Biol.* 2012;10: 60.
8. Field DJ, Lynner C, Brown C, Darroch SAF. Skeletal Correlates for Body Mass Estimation in Modern and Fossil Flying Birds. *PLoS One.* 2013;8: e82000. doi:10.1371/journal.pone.0082000
9. Cubo J, Arthur W. Patterns of correlated character evolution in flightless birds: a phylogenetic approach. *Evol Ecol.* 2000;14: 693–702.
10. Hume JP. The history of the Dodo *Raphus cucullatus* and the penguin of Mauritius. *Hist Biol.* 2006;18: 65–89.
11. Livezey BC. An ecomorphological review of the dodo (*Raphus cucullatus*) and solitaire (*Pezophaps solitaria*), flightless Columbiformes of the Mascarene Islands. *J Zool.* 1993;230: 247–292.
12. Kitchener AC. The external appearance of the dodo, *Raphus cucullatus*. *Arch Nat Hist.* 1993;20: 279–301.

- 398 13. Angst D, Buffetaut E, Abourachid A. The end of the fat dodo? A new mass
399 estimate for *Raphus cucullatus*. *Naturwissenschaften*. 2011;98: 233–236.
- 400 14. Louchart A, Mourer-Chauviré C. The dodo was not so slim : leg dimensions and
401 scaling to body mass. *Naturwissenschaften*. 2011;98: 357–358.
402 doi:10.1007/s00114-011-0771-6
- 403 15. Angst D, Buffetaut E, Abourachid A. In defence of the slim dodo : a reply to
404 Louchart and Mourer-Chauviré. *Naturwissenschaften*. 2011;98: 359–360.
405 doi:10.1007/s00114-011-0772-5
- 406 16. Seebacher F. A new method to calculate allometric length-mass relationships of
407 dinosaurs. *J Vertebr Paleontol*. 2001;21: 51–60.
- 408 17. Hutchinson JR, Ng-Thow-Hing V, Anderson FC. A 3D interactive method for
409 estimating body segmental parameters in animals: Application to the turning and
410 running performance of *Tyrannosaurus rex*. *J Theor Biol*. 2007;246: 660–680.
- 411 18. Gunga HC, Suthau T, Bellmann A, Stoinski S, Friedrich A, Trippel T, et al. A new
412 body mass estimation of *Brachiosaurus brancai* Janensch, 1914 mounted and
413 exhibited at the Museum of Natural History (Berlin, Germany). *Foss Rec*. 2008;11:
414 33–38. doi:10.1002/mmng.200700011
- 415 19. Bates KT, Manning PL, Hodgetts D, Sellers WI. Estimating mass properties of
416 dinosaurs using laser imaging and 3D computer modelling. *PLoS One*. 2009;4:
417 e4532.
- 418 20. Allen V, Paxton H, Hutchinson JR. Variation in Center of Mass Estimates for
419 Extant Sauropsids and its Importance for Reconstructing Inertial Properties of
420 Extinct Archosaurs. *Anat Rec*. 2009;292: 1442–1461.
- 421 21. Sellers WI, Hepworth-Bell J, Falkingham PL, Bates KT, Brassey CA, Egerton VM,
422 et al. Minimum convex hull mass estimations of complete mounted skeletons. *Biol*
423 *Lett*. 2012;8: 842–845.
- 424 22. Brassey CA, Sellers WI. Scaling of convex hull volume to body mass in modern
425 primates, non-primate mammals and birds. *PLoS One*. 2014;9: e91691.
426 doi:10.1371/journal.pone.0091691
- 427 23. Shapiro B, Sibthorpe D, Rambaut A, Austin J, Wragg GM, Bininda-Emonds ORP,
428 et al. Flight of the dodo. *Science*. 2002;295: 1683.
429 doi:10.1126/science.295.5560.1683
- 430 24. CIBC. Seg3D: Volumetric Image Segmentation and Visualization. Scientific
431 Computing and Imaging Institute (SCI). 2014.

- 432 25. Barber CB, Dobkin DP, Huhdanpaa HT. The Quickhull algorithm for convex hulls.
433 ACM Trans Math Softw. 1996;22: 469–483.
- 434 26. Smith RJ. Use and misuse of the reduced major axis for line-fitting. Am J Phys
435 Anthropol. 2009;140: 476–486.
- 436 27. Lavin SR, Karasov WH, Ives AR, Middleton KM, Garland Jr T. Morphometrics of
437 the Avian Small Intestine Compared with That of Nonflying Mammals: A
438 Phylogenetic Approach. Physiol Biochem Zool. 2008;81: 526–550.
- 439 28. Johnson KP, de Kort S, Dinwoodey K, Mateman AC, Ten Cate C, Lessells CM, et
440 al. A molecular phylogeny of the dove genera *Streptopelia* and *Columba*. Auk.
441 2001;118: 874–887. doi:10.1642/0004-8038(2001)118[0874:AMPOTD]2.0.CO;2
- 442 29. Heupink TH, van Grouw H, Lambert DM. The mysterious Spotted Green Pigeon
443 and its relation to the Dodo and its kindred. BMC Evol Biol. 2014;14: 136.
444 doi:10.1186/1471-2148-14-136
- 445 30. Falkingham PL. Acquisition of high resolution three-dimensional models using
446 free, open-source, photogrammetric software. Palaeontol Electron. 2012;15: 1T.
- 447 31. Mallison H, Wings O. Photogrammetry in Paleontology – A Practical Guide. J
448 Paleontol Tech. 2014;12: 1–31.
- 449 32. Wu C. VisualSFM: A visual structure from motion system. [Internet]. 2011.
- 450 33. Brassey C, Gardiner J. An advanced shape-fitting algorithm applied to the
451 vertebrate skeleton: improving volumetric mass estimates. R Soc Open Sci.
- 452 34. Ibrahim T, Bashrat O. Carcass characteristics of pigeons reared in Bauchi
453 Metropolis, north eastern Nigeria. Anim Prod Res Adv. 2009;5.
- 454 35. Omojola AB, Isa MA, Jibir M, S ABTG, Kassim OR, Omotoso AB, et al. Carcass
455 Characteristics and Meat Attributes of Pigeon (*Columbia livia*) as Influenced by
456 Strain and Sex. J Anim Sci Adv. 2012;2: 475–480.
- 457 36. Roth VL. Insular dwarf elephants: a case study in body mass estimation and
458 ecological inferenc. In: Damuth J, MacFadden B, editors. Body size in mammalian
459 paleobiology: Estimates and biological implications. Cambridge: Cambridge
460 University Press; 1990. pp. 151–180.
- 461 37. Herbert T. A relation of some yeares' travaile, begunne Anno 1626, into Afrique
462 and the greater Asia, especially the territories of the Persian Monarchie, and
463 some parts of the Oriental Indies and Isles adjacent. London: W Stansby & J
464 Bloome; 1634.

- 465 38. Brassey CA, Kitchener AC, Withers PJ, Manning PL, Sellers WI. The Role of
466 Cross-Sectional Geometry, Curvature, and Limb Posture in Maintaining Equal
467 Safety Factors: A Computed Tomography Study. *Anat Rec.* 2013;296: 395–413.
- 468 39. Hamershock DM, Seamans TW, Bernhard GE. Determination of body density for
469 twelve bird species. Flight Dynamics Directorate, Wright Laboratory. Flight
470 Dynamics Directorate, Wright Laboratory; 1993.
- 471 40. Randall A, Pepe A, Yamartino K, Kimelblatt A, Claeesens L. How fat was the
472 Dodo? The first mass estimate from digital body reconstruction based on a
473 complete skeleton of *Raphus cucullatus*. Society of Vertebrate Paleontology
474 Annual Meeting 2014, Berlin Poster Abstract. 2014.
- 475 41. Benson RBJ, Campione NE, Carrano MT, Mannion PD, Sullivan C, Upchurch P,
476 et al. Rates of dinosaur body mass evolution indicate 170 million years of
477 sustained ecological innovation on the avian stem lineage. *PLoS Biol.* 2014;12:
478 e1001853. doi:10.1371/journal.pbio.1001853
- 479 42. Rosset A, Spadola L, Ratib O. OsiriX: an open-source software for navigating in
480 multidimensional DICOM images. *J Digit Imaging.* 2004;17: 205–216.
- 481 43. Mascaro J, Litton CM, Hughes RF, Uowolo A, Schnitzer SA. Minimizing Bias in
482 Biomass Allometry : Model Selection and Log-transformation of Data. 2011;43:
483 649–653.

Figure 1. Example of the convex hulling process applied to the CT scanned carcass of a Victoria crowned pigeon (*Goura victoria*) from which the skeleton has been segmented. A and C, skeleton in dorsal and lateral view respectively and; B and D, corresponding convex hulls fitted to the functional units of the skeleton. Note convex hulls fitted to the feet in 1D are strongly influenced by the positioning of the toes (see in text for discussion).





Figure 2. OLS regression results. Body mass (g) against convex hull volume (mm^3). For slope equations see Table 2. Filled circles and solid line, eviscerated carcasses; crosses and dashed line, intact carcasses; dot-dash line, combined sample.



495 Figure 3. A, Photogrammetry model of the Tring dodo skeleton (S/1988.50.1); B,
496 volumetric convex hulls fitted around the skeleton.





Figure 4. The distribution of segment CH_{vol} as a proportion of total CH_{vol} within the convex hulled skeletons of extant pigeons and articulated dodo skeletons. Mean values are illustrated for extant pigeons. Error bars represent 95% confidence intervals of the mean. The underdevelopment of the pectoral girdle (humerus, radius and ulna and carpometacarpals) in dodo relative to extant pigeons is particularly striking.



504 Figure 5. Volumetric renderings of a rock dove (*Columba livia*, A-B) and collared dove
 505 (*Streptopelia decaocto*, C-D) generated from CT scans. A and C illustrate the outer soft
 506 tissue contours of the carcass, while B and D illustrate the position of the gizzard and
 507 associated gizzard contents. There is considerable variation in the quantity and size of
 508 gizzard stones between intact pigeon specimens within the dataset. Renderings were
 509 generated in OsiriX [42]



Species name	Common name	Body mass (g)	Preparation	CH_{vol} (mm ³)	CH_{vol} – feet (mm ³)
<i>Goura victoria</i>	Victoria crowned pigeon	1951	Eviscerated	1487180	1436777
<i>Streptopelia decaocto</i> *	Collared dove	201	Intact	203875	196791
<i>Columba livia</i>	Rock dove	290	Intact	115945	113074
<i>Columba palumbus</i> *	Wood pigeon	305	Intact	337993	328279
<i>Ducula bicolor</i>	Pied imperial pigeon	450	Eviscerated	337377	329220
<i>Petrophassa rufipennis</i>	Chestnut-quilled rock pigeon	314	Eviscerated	303511	286104
<i>Otidiphaps nobilis</i>	Pheasant pigeon	401	Eviscerated	344368	329238
<i>Nesoenas mayeri</i>	Pink pigeon	200	Eviscerated	197171	185981
<i>Ducula</i> sp.	Imperial pigeon	336	Eviscerated	314985	305344
<i>Caloenas nicobarica</i>	Nicobar pigeon	539	Eviscerated	383736	367753
<i>Phaps chalcoptera</i>	Common bronze-winged pigeon	249	Intact	213953	208942
<i>Ducula aenea</i>	Green imperial pigeon	483	Intact	348268	336968
<i>Columba guinea</i>	Speckled pigeon	158	Intact	105156	102041
<i>Zenaida graysoni</i>	Socorro dove	176	Eviscerated	105776	102441

<i>Gallicolumba</i> sp.	Bleeding heart dove	215	Eviscerated	163764	152136
<i>Phapitreron leucotis</i>	White-eared brown dove	107	Eviscerated	69424	67088
<i>Ptilinopus</i> sp.	Fruit dove	71	Eviscerated	47816	46635
<i>Ptilinopus superbus</i>	Superb fruit dove	137	Eviscerated	77882	74691
<i>Treron vernans</i>	Pink-necked green pigeon	167	Eviscerated	104991	101984
<i>Ocyphaps lophotes</i>	Crested pigeon	107	Intact	67451	64011

Table 1. Specimen list of modern pigeons sourced from National Museums Scotland, Edinburgh. CH_{vol} , minimum convex hull volume of the skeleton; $CH_{vol} - \text{feet}$, minimum convex hull volume minus the volume of the feet. * indicates specimens were sourced from the University of Manchester. All other specimens were loaned from National Museums Scotland, Edinburgh.

516

517

518

Model	a	a ($\pm 95\%$)	b	b ($\pm 95\%$)	r^2	p	MSE	AIC _{OLS}	AIC _{PGLS}
Eviscerated	-2.31	-2.90 - -1.72	0.89	0.78-1.00	0.97	<0.001	0.0049	-28.42	-24.66

- minus feet	-2.31	-2.87 - -1.74	0.90	0.79-1.00	0.97	<0.001	0.0046	-29.38	-25.35
Intact	-1.08	-3.69 - -1.53	0.66	0.16-1.16	0.70	0.019	0.0163	-5.29	-9.76
- minus feet	-1.06	-3.62 - -1.50	0.66	0.17-1.15	0.70	0.018	0.0161	-5.41	-10.03
Combined	-2.08	-2.75 - -1.42	0.85	0.72-0.98	0.92	<0.001	0.0086	-34.42	-30.12
- minus feet	-2.08	-2.73 - -1.42	0.85	0.73-0.98	0.92	<0.001	0.0084	-34.94	-30.46

519 Table 2. Ordinary least squares regressions of \log_{10} body mass (g) against \log_{10} convex hull volume (CH_{vol} , mm^3). $\pm 95\%$,
520 95% confidence intervals of the intercept and slope; MSE, mean square error of the regression; AIC, Akaike Information
521 Criterion calculated for Ordinary Least Squares (OLS) and Phylogenetically Generalised Least Squares (PGLS).

522

523

524

Model	Accession number	CH_{vol} (mm^3)	$CH_{vol} - \text{feet}$ (mm^3)	Eviscerated M_b (g) *	95% PI (g)
NHM Tring dodo	S/1988.50.1	8942820	8445134	7980	4653- 13685
NHM Kensington dodo	NHM A.9040	9730367	9283795	8687	5027-15011

Edinburgh dodo	NMS.Z.1993.13	12147000	11787000	10760	6106-18961
----------------	---------------	----------	----------	-------	------------

Table 3. M_b estimated using 'eviscerated' equation minus feet (Table 2) and applying correction factor $e(MSE/2)$ to account for back-transformation of a log-linear model into a power function [43], where MSE is the mean square error reported in Table 2. 95%PI, 95% prediction intervals.

528

529

530

531

